



Comparative Effectiveness Review
Number 231

Care Interventions for People Living With Dementia and Their Caregivers



Comparative Effectiveness Review

Number 231

Care Interventions for People Living With Dementia and Their Caregivers

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
5600 Fishers Lane
Rockville, MD 20857
www.ahrq.gov

Contract No. 290-2015-00008-I

Prepared by:

Minnesota Evidence-based Practice Center
Minneapolis, MN

Investigators:

Mary Butler, Ph.D., M.B.A.
Joseph E. Gaugler, Ph.D.
Kristine M.C. Talley, Ph.D., R.N., G.N.P.-B.C., F.G.S.A.
Hamdi I. Abdi, M.P.H.
Priyanka J. Desai, Ph.D., M.S.P.H.
Susan Duval, Ph.D.
Mary L. Forte, Ph.D., D.C.
Victoria A. Nelson, M.Sc.
Weiwen Ng, M.P.H.
Jeannine M. Ouellette, M.F.A.
Edward Ratner, M.D.
Jayati Saha, Ph.D., M.P.H.
Tetyana Shippee, Ph.D.
Brittin L. Wagner, Ph.D.
Timothy J. Wilt, M.D., M.P.H.
Lobsang Yeshi, M.Sc.

AHRQ Publication No. 20-EHC023

August 2020

Errata December 2020

This report is based on research conducted by the Minnesota Evidence-based Practice Center (EPC) under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. 290-2015-00008-I). The findings and conclusions in this document are those of the authors, who are responsible for its contents; the findings and conclusions do not necessarily represent the views of AHRQ. Therefore, no statement in this report should be construed as an official position of AHRQ or of the U.S. Department of Health and Human Services.

None of the investigators have any affiliations or financial involvement that conflicts with the material presented in this report.

The information in this report is intended to help healthcare decisionmakers—patients and clinicians, health system leaders, and policymakers, among others—make well-informed decisions and thereby improve the quality of healthcare services. This report is not intended to be a substitute for the application of clinical judgment. Anyone who makes decisions concerning the provision of clinical care should consider this report in the same way as any medical reference and in conjunction with all other pertinent information, i.e., in the context of available resources and circumstances presented by individual patients.

This report is made available to the public under the terms of a licensing agreement between the author and the Agency for Healthcare Research and Quality. This report may be used and reprinted without permission except those copyrighted materials that are clearly noted in the report. Further reproduction of those copyrighted materials is prohibited without the express permission of copyright holders.

AHRQ or U.S. Department of Health and Human Services endorsement of any derivative products that may be developed from this report, such as clinical practice guidelines, other quality enhancement tools, or reimbursement or coverage policies, may not be stated or implied.

This report may periodically be assessed for the currency of conclusions. If an assessment is done, the resulting surveillance report describing the methodology and findings will be found on the Effective Health Care Program Web site at www.effectivehealthcare.ahrq.gov. Search on the title of the report.

Suggested citation:

Butler M, Gaugler JE, Talley KMC, Abdi HI, Desai PJ, Duval S, Forte ML, Nelson VA, Ng W, Ouellette JM, Ratner E, Saha J, Shippee T, Wagner BL, Wilt TJ, Yeshi L. Care Interventions for People Living With Dementia and Their Caregivers. Comparative Effectiveness Review No. 231. (Prepared by the Minnesota Evidence-based Practice Center under Contract No. 290-2015-00008-I.) AHRQ Publication No. 20-EHC023. Rockville, MD: Agency for Healthcare Research and Quality; August 2020. Errata December 2020. Posted final reports are located on the Effective Health Care Program [search page](#). DOI: <https://doi.org/10.23970/AHRQEPCCER231>.

Errata

In the original version of the report, the strength of evidence grade for intensive multicomponent interventions (i.e. discrete adaptations of REACH II) on informal caregiver quality of life at 6 months was inconsistently reported. The strength of evidence grade for this finding was erroneously reported as “low” rather than “insufficient” in five instances. These instances have been removed from the Evidence Summary key points and the full report Results and Discussion sections (pp. 77, 78, 81, and 107).

Additional references were added in two instances:

- Page 78: Reference 528 was added to the following sentence: “This result was consistent with a second study that found no difference between groups but improvement over baseline in both intervention and control groups.”
- Page 95: The reference in row 2 of Table 8.16 was corrected to read “616” and not “641.”

Preface

The Agency for Healthcare Research and Quality (AHRQ), through its Evidence-based Practice Centers (EPCs), sponsors the development of systematic reviews to assist public- and private-sector organizations in their efforts to improve the quality of healthcare in the United States. These reviews provide comprehensive, science-based information on common, costly medical conditions, and new healthcare technologies and strategies. The National Institute on Aging (NIA) of the National Institutes of Health (NIH) requested this report from the AHRQ Evidence-based Practice Center (EPC) Program. The report was presented April 15, 2020, at the Health and Medicine Division, National Academies of Sciences, Engineering, and Medicine public meeting on Care Interventions for Individuals With Dementia and Their Caregivers.

The reports and assessments provide organizations with comprehensive, evidence-based information on common medical conditions and new healthcare technologies and strategies. They also identify research gaps in the selected scientific area, identify methodological and scientific weaknesses, suggest research needs, and move the field forward through an unbiased, evidence-based assessment of the available literature. The EPCs systematically review the relevant scientific literature on topics assigned to them by AHRQ and conduct additional analyses when appropriate prior to developing their reports and assessments.

To bring the broadest range of experts into the development of evidence reports and health technology assessments, AHRQ encourages the EPCs to form partnerships and enter into collaborations with other medical and research organizations. The EPCs work with these partner organizations to ensure that the evidence reports and technology assessments they produce will become building blocks for healthcare quality improvement projects throughout the Nation. The reports undergo peer review and public comment prior to their release as a final report.

AHRQ expects that these systematic reviews will be helpful to health plans, providers, purchasers, government programs, and the healthcare system as a whole. Transparency and stakeholder input are essential to the Effective Health Care Program.

If you have comments on this systematic review, they may be sent by mail to the Task Order Officer named below at: Agency for Healthcare Research and Quality, 5600 Fishers Lane, Rockville, MD 20857, or by email to epc@ahrq.hhs.gov.

Gopal Khanna, M.B.A.
Director
Agency for Healthcare Research and Quality

Arlene S. Bierman M.D., M.S.
Director
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Stephanie Chang, M.D., M.P.H.
Director
Evidence-based Practice Center Program
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Kim Wittenberg, M.A.
Task Order Officer
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Acknowledgments

Many people were instrumental to the process that led to this final report. In particular, we wish to thank Melinda Kelley, Elena Fazio, and Courtney Wallin from the NIA for their guidance and editorial skill; Kim Wittenberg from AHRQ for her unflagging support; Ken Tobacman, Mary Walters, Morgaine Butler, and Caroline Au-Yeung for help with abstracting studies for the evidence map; Michelle Brasure for her unparalleled skill in developing search algorithms; Michael Anderson for creating graphics; Cheryl Cole-Hill for helping to collate the draft report; and Carla Kahl for her technical skills that kept all our computers functional.

Key Informants and Technical Expert Panel

The role of the Key Informants was filled by the Health and Medicine Division (HMD) Committee of the National Academies of Sciences, Engineering, and Medicine (NASEM) that will use the report to help develop its own consensus report on which care interventions are supported by sufficient evidence to be widely disseminated and implemented to NASEM and the NIA. (An overview of the NASEM conflict of interest policies can be found at <https://www.nationalacademies.org/about/institutional-policies-and-procedures/conflict-of-interest-policies-and-procedures>) Because the HMD committee would not see the draft Key Questions, the population, intervention, comparator, outcomes, and study timing and setting (PICOTS) to specify the Key Questions, and analytic framework until the KQs were posted for public comment, a panel of content experts from federal agencies acted as proxy Key Informants prior to posting. The proxy Key Informants disclosed any financial conflicts of interest greater than \$10,000 and any other relevant business or professional conflicts of interest.

The list of proxy Key Informants who provided input to this report follows:

*Susan Cooley, Ph.D.
Chief, Dementia Initiatives
U.S. Department of Veterans Affairs
Washington, DC

Valerie Edwards, Ph.D.
Alzheimer's Disease and Healthy Aging Program
Centers for Disease Control and Prevention
Atlanta, GA

Melissa Gerald, Ph.D.
Program Director
Division of Behavioral and Social Research
National Institute on Aging
Bethesda, MD

Margaret Kabat, L.C.S.W.-C.
National Director Caregiver & Family Support
U.S. Department of Veterans Affairs
Washington, DC

Rohini Khillan, M.P.H.
Office of Disability, Aging, and Long Term Care Policy
Office of the Assistant Secretary for Planning and Evaluation
Washington, DC

Greg Link, M.A.
Director, Office of Supportive and Caregiver Services
Administration for Community Living
Washington, DC

*Erin Long, M.S.W.
Aging Services Program Specialist, Dementia Team Lead
Administration for Community Living
Washington, DC

Lisa McGuire, Ph.D.
Lead, Alzheimer's Disease and Healthy Aging Program
Centers for Disease Control and Prevention
Atlanta, GA

*Anthony Pacifico, Ph.D.
Program Manager, Peer Reviewed Alzheimer's Research Program
Congressionally Directed Medical Research Program
Fort Detrick, MD

*This TEP member also provided review of the draft report.

Peer Reviewers

Prior to publication of the final evidence report, EPCs sought input from independent Peer Reviewers without financial conflicts of interest. However, the conclusions and synthesis of the scientific literature presented in this report do not necessarily represent the views of individual reviewers.

Peer Reviewers must disclose any financial conflicts of interest greater than \$10,000 and any other relevant business or professional conflicts of interest. Because of their unique clinical or content expertise, individuals with potential nonfinancial conflicts may be retained. The Task Order Officer and the EPC work to balance, manage, or mitigate any potential nonfinancial conflicts of interest identified.

The list of Peer Reviewers follows:

Richard H. Fortinsky, Ph.D.
Professor
University of Connecticut Center on Aging
University of Connecticut
Farmington, CT

Helen Lamont, Ph.D.
Long-Term Care Policy Analyst
Office of Disability, Aging, and Long Term Care Policy
Office of the Assistant Secretary for Planning and Evaluation
Washington, DC

Katherine Ornstein, Ph.D.
Associate Professor
Icahn School of Medicine at Mount Sinai
New York, NY

Tia Powell, M.D.
Professor
Albert Einstein College of Medicine
Montefiore Medical Center
Bronx, NY

Barbara Resnick, Ph.D., C.R.N.P., F.A.A.N.
Professor
University of Maryland
College Park, MD

Care Interventions for People Living With Dementia and their Caregivers

Structured Abstract

Objective. To understand the evidence base for care interventions for people living with dementia (PLWD) and their caregivers, and to assess the potential for broad dissemination and implementation of that evidence.

Data sources. We searched Ovid Medline, Ovid Embase, Ovid PsycINFO, CINAHL, and the Cochrane Central Register of Controlled Trials (CENTRAL) to identify randomized controlled trials, nonrandomized controlled trials, and quasi-experimental designs published and indexed in bibliographic databases through March 2020.

Review methods. We searched for nondrug interventions targeting PLWD, their informal or formal caregivers, or health systems. Two investigators screened abstracts and full-text articles of identified references for eligibility. Eligible studies included randomized controlled trials and quasi-experimental observational studies enrolling people with Alzheimer’s disease or related dementias or their informal or formal caregivers. We extracted basic study information from all eligible studies. We assessed risk of bias and summarized results for studies not judged to be NIH Stage Model 0 to 2 (pilot or small sample size studies) or to have high risk of bias. We grouped interventions into categories based on intervention target.

Results. We identified 9,217 unique references, of which 627 unique studies with an additional 267 companion articles were eligible. We classified interventions into 37 major categories. With few exceptions, we did not combine data quantitatively due to variability of interventions, comparison groups, outcomes measured, and study timing. Low-strength evidence shows that an intensive multicomponent intervention for informal caregiver support, with education, group discussion, in-home and phone support, and caregiver feedback (i.e., discrete adaptations of REACH II), may improve informal caregiver depression at 6 months. Low-strength evidence also shows that collaborative care models (i.e., Care Ecosystems or discrete adaptations of the ACCESS models) may improve quality of life for PLWD and health system–level markers, including improvements in guideline-based quality indicators and reducing emergency room visits. The literature does not allow for further determination of whether the very small to small average effects in quality of life applied to all enrolled PLWD or if larger effects were concentrated in an unidentified subgroup. For all other interventions and outcomes, we found the evidence insufficient to draw conclusions. Insufficient evidence does not mean that the intervention is determined to be of no value to PLWD or their caregivers. Rather, it means that, due to the uncertainty of the evidence, we cannot draw meaningful conclusions at this time.

Conclusions. Despite hundreds of studies, very little evidence supports widespread dissemination of any general care approaches for PLWD or caregivers. This review demonstrates the need for larger, longer-term, and more-rigorous studies of interventions.

Contents

Evidence Summary	ES-1
Chapter 1. Introduction	1
Background	1
Scope and Key Questions	4
Report Organization.....	10
Chapter 2. Methods	13
Study Selection	13
Data Extraction	14
Assessing Methodological Risk of Bias and NIH Stage of Individual Studies	16
Data Synthesis.....	17
Grading the Strength of Evidence for Major Comparisons and Outcomes	18
Chapter 3. Search Results	20
Chapter 4. Care Interventions for Managing BPSD in PLWD	23
Assisted Therapy.....	23
Multi-Sensory Stimulation/Snoezelen	26
Complementary and Alternative Medicine (CAM) Therapies	28
Bright Light Therapy	30
Psychosocial Therapies for BPSD	31
Multicomponent Interventions for BPSD	32
Conclusion	33
Chapter 5. Care Interventions for PLWD Well-Being	34
Exercise.....	34
Music.....	38
Reminiscence Therapy.....	40
Cognitive Rehabilitation	43
Cognitive Training	46
Cognitive Stimulation Therapy.....	49
Recreation Therapy.....	51
Psychosocial Therapy for PLWD Well-being	53
Creative Expression Therapy.....	54
Multicomponent Interventions for PLWD Well-being.....	55
Assistive Technology.....	57
Electrostimulation	57
Other Interventions for PLWD Well-being	59
Conclusion	60
Chapter 6. Care Interventions for Informal Caregivers	60
Psychosocial Interventions to Support Informal Caregiver Well-being	60
Social Support.....	70
Lifestyle Interventions	73
Respite Care.....	76
Multicomponent Interventions.....	77
Other Interventions for Caregiver Well-being	79
Conclusion	80
Chapter 7. Care Interventions for Formal Caregivers	81

Formal Caregiver Well-being	81
Conclusion	81
Chapter 8. Care Delivery Interventions.....	82
Care Service Provision.....	82
Care Delivery Models and Programs	93
Care Staff Education and Support Needs.....	98
Conclusion	103
Chapter 9. Implementation of Care Interventions	105
Grey Literature Search	105
Chapter 10. Discussion	107
Overview.....	107
Strengths and Weaknesses of the Review.....	107
Future Research	109
Abbreviations and Acronyms	115
References	117

Tables

Table 1.1. PICOTS.....	7
Table 1.2 Glossary of Terms.....	11
Table 2.1. Study inclusion criteria	12
Table 3.1 Identified unique eligible studies bt intervention category, by results chapter	21
Table 4.1. Basic characteristics of literature set: robot-assisted therapy	24
Table 4.2. Summary of findings for PLWD outcomes: robot-assisted therapy.....	25
Table 4.3. Basic characteristics of literature set: live animal- and doll-assisted therapy	25
Table 4.4. Basic characteristics of literature set: multisensory stimulation/Snoezelen	26
Table 4.5. Summary of findings for PLWD outcomes: multisensory stimulation/Snoezelen.....	27
Table 4.6. Basic characteristics of literature set: complementary and alternative medicine	28
Table 4.7. Summary of findings for PLWD outcomes: complementary and alternative medicine	29
Table 4.8. Basic characteristics of literature set: bright light therapy	30
Table 4.9. Basic characteristics of literature set: psychosocial therapies for PLWD	31
Table 4.10 Basic characteristics of literature set: multicomponent interventions for BPSD.....	32
Table 5.1. Basic characteristics of literature set: exercise	35
Table 5.2. Summary of findings for PLWD outcomes: exercise.....	37
Table 5.3. Basic characteristics of literature set: music.....	38
Table 5.4 Summary of findings for PLWD outcomes: music	39
Table 5.5 Summary of findings for caregiver outcomes: music.....	40
Table 5.6 Basic characteristics of literature set: reminiscence therapy	41
Table 5.7 Summary of findings for PLWD outcomes: reminiscence therapy	42
Table 5.8 Summary of findings for caregiver outcomes: reminiscence therapy	43
Table 5.9 Basic characteristics of literature set: cognitive rehabilitation	44
Table 5.10: Summary of findings for PLWD outcomes: cognitive rehabilitation	45
Table 5.11. Basic characteristics of literature set: cognitive training.....	46
Table 5.12 Summary of findings for PLWD outcomes: cognitive training.....	47
Table 5.13 Summary of findings for caregiver outcomes: cognitive training	48
Table 5.14. Basic characteristics of literature set: cognitive stimulation therapy	49

Table 5.15 Summary of findings for PLWD outcomes: cognitive stimulation therapy	50
Table 5.16 Basic characteristics of literature set: recreational therapy	51
Table 5.17 Summary of findings for PLWD outcomes: recreational therapy	52
Table 5.18 Basic characteristics of literature set: psychosocial interventions for PLWD.....	53
Table 5.19 Basic characteristics of literature set: creative expression therapy.....	55
Table 5.20 Basic characteristics of literature set: multicomponent for PLWD well-being.....	55
Table 5.21 Summary of findings for PLWD outcomes: multicomponent for PLWD well-being	56
Table 5.22 Basic characteristics of literature set: assistive home technology	57
Table 5.23 Basic characteristics of literature set:electrostimulation	58
Table 5.24 Basic characteristics of literature set:other interventions for BPSD.....	58
Table 6.1 Basic characteristics of literature set: psychosocial interventions for informal caregivers	61
Table 6.2 Summary of findings for PLWD outcomes: psychosocial interventions for CG/P	64
Table 6.3 Summary of findings for caregiver outcomes: psychosocial interventions for CG/P ..	67
Table 6.4 Basic characteristics of literature set:in-person social support.....	70
Table 6.5 Summary of findings for caregiver outcomes: social support	71
Table 6.6 Basic characteristics of literature set: mindfulness, meditation, and spiritually-focused activities	72
Table 6.7 Summary of findings for caregiver outcomes: mindfulness, meditation, and spiritually-focused activities.....	73
Table 6.8 Basic characteristics of literature set:physical activity and other lifestyle interventions	74
Table 6.9 Basic characteristics of literature set: respite care	75
Table 6.10 Basic characteristics of literature set: multicomponent for CG/P.....	76
Table 6.11 Summary of findings for caregiver outcomes:multicomponent for CG/P.....	77
Table 6.12 Basic characteristics of literature set: other interventions for CG/P.....	78
Table 7.1 Basic characteristics of literature set: formal caregiver well-being.....	81
Table 8.1 Basic characteristics of literature set: consultation services	83
Table 8.2 Summary of findings for PLWD outcomes: consultation services.....	83
Table 8.3 Summary of findings for caregiver outcomes: consultation services	82
Table 8.4 Basic characteristics of literature set: case management	85
Table 8.5 Summary of findings for PLWD outcomes: case management.....	85
Table 8.6 Summary of findings for caregiver outcomes: case management	86
Table 8.7 Basic characteristics of literature set: care protocols for PLWD.....	87
Table 8.8. Basic characteristics of literature set: advance care planning.....	88
Table 8.9. Summary of findings for PLWD outcomes: advance care planning	88
Table 8.10 Summary of findings for caregiver outcomes: advance care planning.....	89
Table 8.11. Basic characteristics of literature set: palliative care.....	89
Table 8.12 Basic characteristics of literature set: other service provision interventions.....	90
Table 8.13 Summary of findings for PLWD outcomes: other service provision interventions....	91
Table 8.14 Summary of findings for caregiver outcomes: other service provision interventions	91
Table 8.15. Basic characteristics of literature set: collaborative care models	93
Table 8.16 Summary of findings for PLWD outcomes: collaborative care models	94
Table 8.17. Summary of findings for caregiver outcomes: collaborative care models	95
Table 8.18. Basic characteristics of literature set: individualized person-centered care (non-U.S.)	97

Table 8.19. Basic characteristics of literature set: formal caregiver staff training	98
Table 8.20 Basic characteristics of literature set: CG/P training	99
Table 8.21 Basic characteristics of literature set: family education and partnering	100
Table 8.22 Basic characteristics of literature set: multitier training	101
Table 8.23 Summary of findings for PLWD outcomes: multitier training.....	101
Table 8.24 Summary of findings for caregiver outcomes: multitier training	102

Figures

Figure 1.1. Framework for care interventions	3
Figure 1.2. Analytic framework.....	10
Figure 2.1. Review extraction and assessment processes	15
Figure 3.1 Literature flow diagram.....	20
Figure 6.1 Collaboration network visualization of informal caregiver psychosocial intervention components.....	61

Appendixes

Appendix A. Methods
Appendix B. Excluded Studies
Appendix C. Chapter 4 Treatment for BPSD
Appendix D. Chapter 5 Interventions for PLWD Well-Being
Appendix E. Chapter 6 and 7 Interventions for Caregiver Well-Being
Appendix F. Chapter 8 Care Delivery Interventions Consultation

Evidence Summary

Main Points

- An intensive multicomponent intervention with education, group discussion, in-home and phone support sessions, and caregiver feedback for informal caregiver support (i.e. discrete adaptations of REACH II), may improve informal caregiver depression at 6 months. (low-strength evidence)
- Collaborative care models (i.e. Care Ecosystems or discrete adaptations of the ACCESS models) may improve people living with dementia (PLWD) quality of life. (low-strength evidence) The literature does not allow for further determination of whether the very small to small average effects applied to all enrolled PLWD or if larger effects were concentrated in an unidentified subgroup.
- Collaborative care models (i.e. discrete adaptations of the ACCESS model) may improve system-level markers, including guideline-based quality indicators and reduction in emergency department visits. (low-strength evidence)
- For all other outcomes and interventions, we found the evidence was insufficient because the uncertainty of the evidence was too high to draw conclusions.
- We found little information to determine whether interventions are equally appropriate for or have been successfully adapted to other race/ethnic cultures within the United States, rural communities/communities with low resources, or specific populations of PLWD, such as people with Down syndrome or complex presentations of dementias.

Background and Purpose

The aging of the U.S. population and the concurrent rise in the number of adults living with dementia underscore the urgent need for a systematic review of the available evidence for care interventions for PLWD and their formal and informal caregivers. The National Institute on Aging commissioned such a review from the Evidence-based Practice Center Program at the Agency for Healthcare Research and Quality. Dementia affects about 5 million U.S. adults 65 years and older (with disparities by race and ethnicity), and that number may grow to almost 14 million by 2060.^{1,2}

The goal is to understand the evidence base for effective care interventions, and to assess the potential for broad dissemination and implementation of that evidence. Subsequently, a National Academies of Sciences, Engineering, and Medicine committee of experts, established at the request of the National Institute on Aging, will use the evidence findings delivered in this report to help develop its own independent recommendations regarding which care interventions are supported by sufficient evidence to be widely disseminated and implemented, as well as to identify research gaps.

Methods

The methods for this systematic review follow the Agency for Healthcare Research and Quality Methods Guide for Effectiveness and Comparative Effectiveness Reviews.⁹ See the review protocol (<https://effectivehealthcare.ahrq.gov/products/care-interventions-pwd/protocol>) and the full report of the review for additional details. We searched Ovid Medline, Ovid Embase, Ovid PsycINFO, CINAHL, and the Cochrane Central Register of Controlled Trials (CENTRAL)

to identify randomized controlled trials, nonrandomized controlled trials, and quasi-experimental designs published and indexed in bibliographic databases through October, 2019.

Results

We identified 9217 unique references, 894 of which were eligible for our review, comprising 627 unique studies with an additional 267 companion publications. We sorted eligible studies into 37 major intervention categories. Approximately 60 percent of the literature emerged from research conducted outside of the United States.

While the literature was highly diverse, we found little information to determine whether interventions are equally appropriate for or have been successfully adapted to other race/ethnic cultures within the United States, with only a handful of studies providing sufficient inclusion of African-American or Hispanic/Latino populations. Interventions for rural communities were found to be even more rare in the literature. Additionally, the many countries in which care interventions were evaluated were almost exclusively high-resource, with very few low-resource countries represented. Finally, few studies gathered the granular detail necessary for a deeper understanding of the applicability of the interventions. Many important groups were rarely studied, including people with Down syndrome, who are living longer and who overall experience higher rates of dementia than the general population, and individuals with complex presentations of dementia.

We found low-strength evidence that one multicomponent intervention for informal caregiver support, REACH II (comprised of education, group discussion, in-home and phone support sessions, and caregiver feedback), may improve caregiver depression.³⁻⁵ Collaborative care models based on the ACCESS or Care Ecosystems models may improve quality of life for PLWD⁶⁻⁹ and health system-level markers, including improvements in guideline-based quality indicators^{7,8} and reducing emergency room visits.⁶ For all other interventions and outcomes, we found the uncertainty of the evidence was too high to draw conclusions. The vast majority of studies had small sample size, were pilots that had not undergone traditional efficacy testing, and/or had high risk of bias, and provided insufficient evidence to draw conclusions.

Limitations

As stated, the goal of the review was to understand the evidence base for effective care interventions for PLWD and their caregivers in order to assess the potential for broad dissemination and implementation of that evidence. All decisions about the review methodology followed from this goal, which has implications for our findings. Therefore, some specific approaches for a particular intervention, or even whole classes of interventions, may not have been captured. Similarly, we may have missed some community services and support approaches such as tool kits, referral services and links, or awareness-raising outreach.

Because we excluded studies with fewer than 10 participants per study arm, we may not have identified some interventions with only very preliminary research. Our approach to risk of bias assessment was generous, relative to how risk of bias is assessed in more targeted systematic review topics. This is in part due to the unusually varied studies included in this review as well as the complexity of the condition and the care approaches.

Furthermore, included studies had methodological problems such as low subject retention, widely varying measures of success, and relatively small size trials that may have lacked the power to detect benefits. These problems stem from well-recognized challenges in researching these populations; therefore, we erred on the side of assessing a body of research as insufficient

rather than ineffective. Additionally, given the progressive nature of dementia and the anticipated increase in care needs over time, it may be difficult to identify improvement in caregiver burden. The goal may in fact be to slow the rate of burden—and studies may be underpowered to detect such a small effect.

Finally, given the breadth of the topic, our systematic review is naturally reductionist in nature. That is, small but true differences may exist between many of the categories we summarized into our outcomes.

Implications and Conclusions

Ultimately, we uncovered very little evidence to support interventions and programs for active, widespread dissemination because evidence was insufficient to draw conclusions about the effects of the vast majority of interventions studied.

Dementia care research has been slow to incorporate key elements of rigorous intervention design. Until relatively recently, many dementia care intervention studies were not held to reporting standards (e.g. the Consolidated Standards of Reporting Trials statement), pre-registration of trials, data safety and monitoring boards, or other standards more common in other areas of clinical science. As a direct result, despite a few positive findings, our global conclusions, largely similar to past reviews,¹⁰ are that the amount of high-quality evidence is insufficient to draw firm conclusions about interventions.

In order for Federal funders and stakeholders to expedite the translational pipeline of idea development to implementation, as they aim to do, critical improvements are needed in dementia care research. Only with such improvements will we be able to draw clearer, less ambiguous conclusions related to efficacy.

References

1. Matthews KA, Xu W, Gaglioti AH, et al. Racial and ethnic estimates of Alzheimer's disease and related dementias in the United States (2015-2060) in adults aged ≥ 65 years. *Alzheimers Dement*. 2019 Jan;15(1):17-24. doi: 10.1016/j.jalz.2018.06.3063. PMID: 30243772.
2. Hebert LE, Weuve J, Scherr PA, et al. Alzheimer disease in the United States (2010-2050) estimated using the 2010 census. *Neurology*. 2013 May 7;80(19):1778-83. doi: 10.1212/WNL.0b013e31828726f5. PMID: 23390181.
3. Belle SH, Burgio L, Burns R, et al. Enhancing the quality of life of dementia caregivers from different ethnic or racial groups: a randomized, controlled trial.[Summary for patients in *Ann Intern Med*. 2006 Nov 21;145(10):I39; PMID: 17116914]. *Annals of Internal Medicine*. 2006 Nov 21;145(10):727-38. PMID: 17116917.
4. Berwig M, Heinrich S, Spahlholz J, et al. Individualized support for informal caregivers of people with dementia - effectiveness of the German adaptation of REACH II. *BMC Geriatrics*. 2017 Dec 12;17(1):286. doi: <https://dx.doi.org/10.1186/s12877-017-0678-y>. PMID: 29233097.
5. Luchsinger JA, Burgio L, Mittelman M, et al. Comparative Effectiveness of 2 Interventions for Hispanic Caregivers of Persons with Dementia. *Journal of the American Geriatrics Society*. 2018 Sep;66(9):1708-15. doi: <https://dx.doi.org/10.1111/jgs.15450>. PMID: 30084133.

6. Possin KL, Merrilees JJ, Dulaney S, et al. Effect of Collaborative Dementia Care via Telephone and Internet on Quality of Life, Caregiver Well-being, and Health Care Use: The Care Ecosystem Randomized Clinical Trial. *JAMA Internal Medicine*. 2019 Sep 30;30:30. doi: <https://dx.doi.org/10.1001/jamainternmed.2019.4101>. PMID: 31566651.
7. Chodosh J, Colaiaco BA, Connor KI, et al. Dementia Care Management in an Underserved Community: The Comparative Effectiveness of Two Different Approaches. *Journal of Aging & Health*. 2015 Aug;27(5):864-93. doi: <https://dx.doi.org/10.1177/0898264315569454>. PMID: 25656074.
8. Vickrey BG, Mittman BS, Connor KI, et al. The effect of a disease management intervention on quality and outcomes of dementia care: a randomized, controlled trial.[Summary for patients in *Ann Intern Med*. 2006 Nov 21;145(10):I31; PMID: 17116913]. *Annals of Internal Medicine*. 2006 Nov 21;145(10):713-26. PMID: 17116916.
9. Thyrian JR, Hertel J, Wucherer D, et al. Effectiveness and Safety of Dementia Care Management in Primary Care: A Randomized Clinical Trial. *JAMA Psychiatry*. 2017 Oct 01;74(10):996-1004. doi: <https://dx.doi.org/10.1001/jamapsychiatry.2017.2124>. PMID: 28746708.
10. Gaugler JE, Jutkowitz E, Gitlin L. Non-Pharmacological Interventions for Persons Living with Alzheimer's Disease: Decadal Review and Recommendations. National Academies of Sciences, Engineering, and Medicine: Washington DC. 2020.

Chapter 1. Introduction

Background

The aging of the U.S. population and the concurrent rise in the number of adults living with dementia underscore the urgent need for a systematic review of the available evidence for care interventions for people living with dementia (PLWD) and their formal and informal caregivers.¹ The National Institute on Aging (NIA) has commissioned such a review from the Evidence-based Practice Center Program at the Agency for Healthcare Research and Quality (AHRQ). The goal is to understand the evidence base for effective care interventions, and to assess the potential for broad dissemination and implementation of that evidence.

Dementia affects about 5 million U.S. adults 65 years and older (with disparities by race and ethnicity), and that number may grow to almost 14 million by 2060.¹⁻³ A further 200,000 individuals under age 65 have some form of early-onset dementia.¹ As a clinical syndrome and a disability, dementia is characterized by an acquired cognitive deficit that interferes with independence in daily activities.⁴ Alzheimer's disease is the most common form of progressive dementia and, grouped with Lewy body, frontotemporal, vascular, and mixed forms, it has been referred to as AD/ADRD (i.e., Alzheimer's disease [AD] and Alzheimer's disease related dementias [ADRD]).⁵ Dementia can lower an individual's quality of life, burden caregivers (even those who find caregiving very rewarding), increase institutionalization, and increase costs to families and society.⁶ Agitation, aggression, and other behavioral disturbances are common, especially late in the disease course.⁷

The significant public health implications of dementia led to the 2011 passage of the National Alzheimer's Project Act, an effort to create a national research strategy to accelerate scientific discovery of curative treatments, preventive approaches, and effective strategies to manage and alleviate the many clinical symptoms of AD/ADRD. The National Alzheimer's Project Act has spurred considerable Federal investment; the National Institutes of Health (NIH) funding for AD/ADRD research has more than tripled since 2015.⁸ (See https://report.nih.gov/categorical_spending.aspx.)

In addition to attracting new investigators, the NIA and other Federal agencies have leveraged funding increases to initiate and/or contribute to several important, complementary, large-scale efforts to improve the design and delivery of care for PLWD. Among these are the first ever National Research Summit on Care, Services, and Supports for Persons with Dementia and Their Caregivers. This 2017 summit generated a number of recommendations to advance the science of dementia care and catalyzed several important actions, including a substantial investment in several Funding Opportunity Announcements, one of which resulted in award of the NIA IMbedded Pragmatic Alzheimer's disease (AD) and AD Related Dementias (AD/ADRD) Clinical Trials (IMPACT) Collaboratory. The IMPACT Collaboratory is designed to provide technical assistance and pilot grant support to "embed" dementia care trials within healthcare systems across the United States, a decadal review of the state of behavioral and social science research (including but not limited to AD/ADRD care intervention research) commissioned by the National Academies of Sciences, Engineering and Medicine (NASEM), as well as other efforts.

While these initiatives are designed for rapid advancement of the science of dementia, this systematic review of the available evidence for care interventions for PLWD and their formal and informal caregivers will provide valuable information about the efficacy or effectiveness of certain types of strategies/interventions. This review also pinpoints areas that require greater

attention when evaluating, disseminating, or implementing certain dementia care and caregiver interventions.

Dementia has no known cure, but both drug and nondrug interventions are available to treat symptoms, support function, and improve quality of life. Nondrug interventions have been recommended as first-line treatments for behavioral and psychological symptoms of dementia (BPSD), but pharmacological treatment options such as antipsychotics are also available.⁹ And although nondrug interventions are generally presumed safe, few trials have reported information on their harms or other unintended consequences. (Drugs and over-the-counter supplements to treat clinical Alzheimer's-type dementia and BPSD are being addressed by a separate AHRQ systematic review; please see <https://effectivehealthcare.ahrq.gov/products/alzheimers-type-dementia/research>.)

Dementia care is costly, and more than 83 percent of community-residing older adults who need it rely on help from family members.¹⁰ In 2017, informal (unpaid) caregivers for PLWD provided an estimated 17 billion hours of care at an economic value of \$232.1 billion, and about two-thirds of informal caregivers are women.¹¹ Caregiving for dementia is multifaceted and can be both rewarding and burdensome, sometimes simultaneously. Many surveys suggest that, for some, caregiving instills confidence, provides lessons on dealing with difficult situations, and increases feelings of closeness to the care recipient.¹² However, evidence also suggests that caregivers have lower self-ratings of physical health, elevated levels of stress hormones, higher rates of chronic disease, and impaired health behaviors. Therefore, many research teams have developed and tested interventions for supporting the health and well-being of informal caregivers. Some examples include social support, therapeutic counseling, skills training, respite, and combined approaches.¹³ Additionally, many frontline paid caregivers, such as home health aides in home-based settings or certified nursing assistants in institutional settings, lack adequate training and support for this difficult work.¹² A recent NASEM report recommended an increase in Federal requirements for training of direct care workers—from 75 hours to 120 hours—along with more focus on knowledge and skills related to caring for PLWD.¹⁴


Care interventions for PLWD encompass a broad range of activities that support, enhance, or otherwise help the care recipient. Likewise, care interventions comprise an array of options that, as noted by the NASEM committee's feedback on the framing and parameters for this review [<https://www.nationalacademies.org/our-work/care-interventions-for-individuals-with-dementia-and-their-caregivers>], “contribute to a person's well-being, happiness, identity, privacy, capacity, autonomy, or authority. They can be supports, services, programs, accommodations, or practices that include behavioral, environmental, technological, and psychological methods or approaches. They may be delivered by healthcare, social services, and other community organizations or caregivers with the intention of having a direct impact on either a person with dementia or their caregiver or both.”

Necessarily, then, interventions addressing care for PLWD and their caregivers can be complicated and multifaceted. Unfortunately, no consensus has been reached on classification systems for types of interventions, and investigators are left to categorize interventions themselves, based on varied criteria. In our own effort to categorize interventions for this review, we identified two basic intervention groups: 1) interventions testing a type of care that is aimed at improving the health and well-being of PLWD and/or their caregivers (e.g., interventions that use music or essential oils to help calm the care recipient, or respite care that provides a break for the caregiver and 2) interventions testing the manner in which care is delivered in order to improve effectiveness, efficiency, and/or accessibility and availability of care (e.g., staff training

for caregivers, coordination of care). This review refers to interventions testing a type of care as care interventions, and interventions testing the manner in which care is delivered as care delivery interventions.

Intervention complexity also stems from the diversity of PLWD (e.g., younger adults with Down syndrome or other genetic risk factors, younger and middle-aged adults with frontotemporal dementia, and older adults with AD, from very early to advanced stages) as well as different caregiver populations (e.g., spousal caregivers, adult child caregivers, paid caregivers). Intervention designs may be straightforward and aimed at supporting a single, well-defined group, such as formal or informal caregivers, or they may be very complex and target several levels of a system simultaneously, from a care system (e.g., healthcare or social services) to family units or caregiver/PLWD dyads to individual formal or informal caregivers. (Figure 1.1) Furthermore, complexity in outcomes may arise because interventions targeting one level of a system, such as PLWD, may benefit other individuals, such as caregivers, or other levels of the system, such as reduced use of healthcare services for an accountable care organization.

Figure 1.1. Framework for care interventions



Abbreviations: CG=caregiver; CR=care recipient
 Source: NASEM, 2016, *Families caring for an aging America*. p. 163. Used with permission.

Given these complexities, our review aims to specify intervention characteristics that link to benefits. Unfortunately, informal and formal caregivers may not always be easily characterized according to the levels outlined in Figure 1.1; paid caregivers may be hired as independent contractors by family of a PLWD, whereas unpaid volunteers may be affiliated with a larger organization. Nonetheless, information regarding relationships between PLWD and caregiver characteristics and outcomes will help clinicians, care providers, and other stakeholders make decisions about the best interventions for their specific circumstances or PLWD.

Assessing whether a care intervention is ready for broad implementation is challenging. For this review, we were guided by the NIH Stage Model for Behavioral Interventions.¹⁵ This model provides a conceptual framework of intervention research development, ranging from basic science research (Stage 0) to new intervention creation (Stage 1), research-setting efficacy (Stage 2), “real-world” community-clinic efficacy (Stage 3), broad community-based effectiveness (Stage 4), to eventually dissemination and implementation research (Stage 5). This model not only describes the stages of behavioral intervention development, but also supports eventual implementation. While the stages are not a direct assessment of implementation readiness, the model suggests that interventions at Stage 3 or higher are more likely to be ready for broad dissemination. Interventions at Stage 4 that use pragmatic study designs move research closer to “real world” conditions and population levels.

Scope and Key Questions

This review examines a large number of nondrug care interventions targeted at PLWD, their informal and formal caregivers, and the larger health systems, including collaborative or integrated care. The intended audience is similarly broad, from PLWD and their families, to care services and support providers, to research organizations and policymakers at national, local, state, tribal, and Federal levels.

Given such a wide range of stakeholders, we have prioritized the readability and usability of our review by striving for plain language and avoiding technical and field-specific jargon as much as possible throughout this report. We acknowledge that the wide readership for these findings makes communication challenging. Stakeholder groups differ in how they define and address terms and concepts related to care interventions for PLWD and their caregivers, and some of these differences may be philosophical. Additionally, individual members of stakeholder groups may differ on these matters. This report cannot resolve the rich and nuanced discussions that would be needed to come to a wide-ranging consensus about which terms to use and where. Therefore, we have opted to use the terms most commonly found within the studies we identified for this review. Our choice reflects our priority to communicate the evidence base in the clearest and most accessible way to the largest number of people. In no way does our choice of terms reflect any particular philosophical position. We recognize that the various perspectives among readers may lead to different interpretations of our report. Our intention is to honor all perspectives and value all audiences.

The review specifically focuses on AD/ADRD, informal and formal caregivers, and the effect of interventions on outcomes for people or systems beyond the intended intervention target. We *did not exclude* any care setting. However, because the purpose is to inform readiness for dissemination, the review *does not include* education interventions conducted in educational settings. Because of the interest in daily caregiving, we also *did not include* workplace-based training programs that targeted professional staff (such as physicians and registered nurses) rather than auxiliary staff.

Key Questions

The Key Questions (KQs) are structured to organize the literature by the intervention target and grouped such that outcomes for PLWD and caregivers were examined regardless of the intervention target. The KQs are further specified by the populations, interventions, comparators, outcomes, timing, and settings (PICOTS) laid out in Table 1.1.

Care Interventions for Behavioral and Psychological Symptoms of Dementia (BPSD) in People Living With Dementia (PLWD):

- KQ1: For people living with dementia (**PLWD**), what are the benefits and harms of care interventions aimed at treating the behavioral and psychological symptoms of dementia (BPSD) in PLWD?
 - KQ1a: What evidence is available on how outcomes differ by **PLWD** characteristics?
 - KQ1b: What evidence is available on how outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ1c: Which intervention characteristics or components are associated with effectiveness?
- KQ2: For informal and/or formal **PLWD Caregivers**, what are the benefits and harms for care interventions aimed at treating BPSD in **PLWD**?
 - KQ2a: What evidence is available on how outcomes differ by **PLWD** characteristics?
 - KQ2b: What evidence is available on how outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ2c: Which intervention characteristics or components are associated with effectiveness?

Care Interventions for Quality of Life, Function, or Non-BPSD Symptoms in PLWD:

- KQ3: For people living with dementia (**PLWD**), what are the benefits and harms for care interventions aimed at improving quality of life, function, or non-BPSD symptoms in **PLWD**?
 - KQ3a: What evidence is available on how outcomes differ by **PLWD** characteristics?
 - KQ3b: What evidence is available on how outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ3c: Which intervention characteristics or components are associated with effectiveness?
- KQ4: For informal and/or formal **PLWD Caregivers**, what are the benefits and harms for care interventions aimed at improving quality of life, function, or non-BPSD symptoms in **PLWD**?
 - KQ4a: What evidence is available on how outcomes differ by **PLWD** characteristics?
 - KQ4b: What evidence is available on how outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ4c: Which intervention characteristics or components are associated with effectiveness?

Care Interventions for Quality of Life and Health Outcomes for Informal and Formal PLWD Caregivers:

- KQ5: For people living with dementia (**PLWD**), what are the benefits and harms for care interventions aimed at supporting the quality of life and health outcomes of the informal **PLWD Caregivers**?
 - KQ5a: What evidence is available on how quality of life and outcomes differ by **PLWD** characteristics?
 - KQ5b: What evidence is available on how quality of life and outcomes differ by informal or formal **PLWD Caregiver** characteristics?

- KQ5c: Which intervention characteristics or components are associated with effectiveness?
- KQ6: For informal and/or formal **PLWD Caregivers**, what are the benefits and harms for care interventions aimed at supporting the quality of life and health outcomes of the informal **PLWD Caregivers**?
 - KQ6a: What evidence is available on how quality of life and outcomes differ by **PLWD** characteristics?
 - KQ6b: What evidence is available on how quality of life and outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ6c: Which intervention characteristics or components are associated with effectiveness?
- KQ7: For people living with dementia (**PLWD**), what are the benefits and harms for care interventions aimed at supporting the quality of life and health outcomes of the formal **PLWD Caregivers**?
 - KQ7a: What evidence is available on how quality of life and outcomes differ by **PLWD** characteristics?
 - KQ7b: What evidence is available on how quality of life and outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ7c: Which intervention characteristics or components are associated with effectiveness?
- KQ8: For informal and/or formal **PLWD Caregivers**, what are the benefits and harms for care interventions aimed at supporting the quality of life and health outcomes of the formal **PLWD Caregivers**?
 - KQ8a: What evidence is available on how quality of life and outcomes differ by **PLWD** characteristics?
 - KQ8b: What evidence is available on how quality of life and outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ8c: Which intervention characteristics or components are associated with effectiveness?

Interventions for How Care Is Delivered:

- KQ9: For people living with dementia (**PLWD**), what are the benefits and harms for care delivery interventions?
 - KQ9a: What evidence is available on how outcomes differ by **PLWD** characteristics?
 - KQ9b: What evidence is available on how outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ9c: Which intervention characteristics or components are associated with effectiveness?
- KQ10: For informal and formal **PLWD Caregivers**, what are the benefits and harms for care delivery interventions?
 - KQ10a: What evidence is available on how outcomes differ by **PLWD** characteristics?
 - KQ10b: What evidence is available on how outcomes differ by informal and/or formal **PLWD Caregiver** characteristics?
 - KQ10c: Which intervention characteristics or components are associated with effectiveness?

Dissemination and Implementation Research:

- Guiding Question 1: What is the state of the empirical evidence on implementation of interventions that have at least low-strength evidence for “real-world” benefits and harms (i.e., NIH Stage Model for Behavioral Intervention Development Stages 3-5)?

Note that in Table 1.1, outcomes are loosely organized to correspond with the levels shown in Figure 1.1. Importantly, we based our final organization of outcomes according to the outcomes examined and measures used in eligible studies, as well as the intentions of the authors.

Table 1.1. PICOTS

Element	PLWD	PLWD Caregiver
Population	<p>PLWD, including individuals with possible or diagnosed AD/ADRD.</p> <p>PLWD Subgroups: Age, sex, sexual orientation/gender identity, race/ethnicity, education, socioeconomic status, prior disability, age at diagnosis, dementia type, dementia severity [e.g. stage of dementia (early stage, moderate, or severe), level of cognitive impairment rate of cognitive decline], family/household characteristics, health insurance, geographic location (e.g. urban, rural), setting type</p>	<p>Informal PLWD Caregivers, such as spouses, family, friends, and volunteers</p> <p>Informal PLWD Caregiver Subgroups, including age, sex, sexual orientation/gender identity, race/ethnicity, family history of dementia, education, socioeconomic status, employment status, relationship with PLWD, living distance from PLWD, dementia care training, general health status, caregiving networks, setting type</p> <p>Formal PLWD Caregivers, such as certified nursing assistants (CNAs), home health aides, auxiliary workers, personal care aides, hospice aides, promotoras or promotores, and community health workers</p> <p>Formal PLWD Caregiver Subgroups, including age, sex, race/ethnicity, education, job position, skill, training, general health status, setting type</p>
Intervention	<p>KQ 1-4. Any nondrug care intervention intended to benefit PLWD except interventions to treat conditions other than dementia, including but not limited to CPAP, and those that use supplements/natural products.</p> <p>(See list of example intervention types in Appendix A.)</p> <p>Guiding Question: Any quality improvement or implementation science study that informs the dissemination or implementation of a care intervention at least low-strength evidence for “real-world” benefits and harms (i.e., NIH Stage Model for Behavioral Intervention Development Stages 3-5)</p>	<p>KQ 5-6. Any care intervention intended to support informal PLWD caregivers’ well-being except interventions to treat health conditions unrelated to providing care to PLWD.</p> <p>KQ 7-8. Any care intervention intended to support formal PLWD caregivers’ well-being except interventions to treat health conditions unrelated to providing care to PLWD.</p> <p>KQ 9-10. Any care delivery intervention to improve how care is delivered IF the training intervention is incorporated as on-going operational procedures into the structure or processes of the organization. Interventions carried out by higher education organizations or professional organizations to provide training toward licensed professionals, and continuing education for degreed health professionals are also excluded.</p> <p>(See list of example intervention types in Appendix A.)</p> <p>Guiding Question: Any quality improvement or implementation science study that informs the dissemination or implementation of a care intervention at least low-strength evidence for</p>

Element	PLWD	PLWD Caregiver
Population	<p>PLWD, including individuals with possible or diagnosed AD/ADRD.</p> <p>PLWD Subgroups: Age, sex, sexual orientation/gender identity, race/ethnicity, education, socioeconomic status, prior disability, age at diagnosis, dementia type, dementia severity [e.g. stage of dementia (early stage, moderate, or severe), level of cognitive impairment rate of cognitive decline], family/household characteristics, health insurance, geographic location (e.g. urban, rural), setting type</p>	<p>Informal PLWD Caregivers, such as spouses, family, friends, and volunteers</p> <p>Informal PLWD Caregiver Subgroups, including age, sex, sexual orientation/gender identity, race/ethnicity, family history of dementia, education, socioeconomic status, employment status, relationship with PLWD, living distance from PLWD, dementia care training, general health status, caregiving networks, setting type</p> <p>Formal PLWD Caregivers, such as certified nursing assistants (CNAs), home health aides, auxiliary workers, personal care aides, hospice aides, promotoras or promotores, and community health workers</p> <p>Formal PLWD Caregiver Subgroups, including age, sex, race/ethnicity, education, job position, skill, training, general health status, setting type</p>
		<p>"real-world" benefits and harms (i.e., NIH Stage Model for Behavioral Intervention Development 3-5)</p>
Comparator	<p>Inactive Comparator: No intervention, usual care, waitlist, attention control</p> <p>Active Comparator: Different intervention</p>	<p>Inactive Comparator: No intervention, usual care, waitlist, attention control</p> <p>Active Comparator: Different intervention</p>
Outcomes (Generally organized to correspond with Figure 1.1 Framework for care interventions)	<p>Quality of life and subjective well-being Burden of care* Satisfaction with care Perceived Support</p> <p>Expenditures/financial burden (informal caregivers)</p> <p>Health-related outcomes: Psychological health (e.g., depression, anxiety) Neuropsychiatric symptoms (including apathy, aggression, and agitation) Function (e.g., ADL, IADL, ability to care for one's self, ability to recreate/socialize) Weight loss Sleep problems Use of restraints Use of anti-psychotics Harm reduction (e.g. driving, firearms)</p> <p>Palliative care/hospice outcomes: Completion of advanced directives Comfort during dying process Concordance with preferred location of death</p> <p>Social/Community level outcomes: Engagement in community activities, Perceived inclusion Safety/perceived safety</p> <p>Utilization of healthcare service outcomes:</p>	<p>Quality of life and subjective well-being Burden of care* Satisfaction with care for PLWD (informal caregivers) Perceived Support</p> <p>Expenditures/financial burden (informal caregivers)</p> <p>Health-related outcomes: Psychological health (e.g., depression, anxiety) Immune function (e.g., inflammation or cortisol) Sleep problems Weight loss due to stress Health behaviors (e.g., exercise, substance use)</p> <p>Caregiving self-efficacy Confidence to manage caregiver tasks</p> <p>Social/Community level outcomes (informal caregivers): Engagement in community activities, Perceived inclusion Safety/perceived safety</p> <p>Turnover and retention (formal caregivers) Utilization of healthcare service (e.g., physician visits, antidepressant or anti-anxiety medication usage) Societal costs including caregiving time/time spent on activities</p> <p>Harms, including isolation, loneliness, perceived stigma, caregiver PTSD</p>

Element	PLWD	PLWD Caregiver
Population	<p>PLWD, including individuals with possible or diagnosed AD/ADRD.</p> <p>PLWD Subgroups: Age, sex, sexual orientation/gender identity, race/ethnicity, education, socioeconomic status, prior disability, age at diagnosis, dementia type, dementia severity [e.g. stage of dementia (early stage, moderate, or severe), level of cognitive impairment rate of cognitive decline], family/household characteristics, health insurance, geographic location (e.g. urban, rural), setting type</p>	<p>Informal PLWD Caregivers, such as spouses, family, friends, and volunteers</p> <p>Informal PLWD Caregiver Subgroups, including age, sex, sexual orientation/gender identity, race/ethnicity, family history of dementia, education, socioeconomic status, employment status, relationship with PLWD, living distance from PLWD, dementia care training, general health status, caregiving networks, setting type</p> <p>Formal PLWD Caregivers, such as certified nursing assistants (CNAs), home health aides, auxiliary workers, personal care aides, hospice aides, promotoras or promotores, and community health workers</p> <p>Formal PLWD Caregiver Subgroups, including age, sex, race/ethnicity, education, job position, skill, training, general health status, setting type</p>
	<p>Admission to nursing home Access to care and services ICU and ED usage Hospital admission and readmission Primary, Specialty, Long-term Care usage</p> <p>Quality of care and services (e.g., overutilization of unnecessary antibiotics, other quality care metrics.)</p> <p>Societal costs, including caregiving time/time spent on activities</p> <p>Harms, including isolation, loneliness, perceived stigma, suicidal ideation or suicide, elder abuse (e.g., physical harm, abuse, neglect, exploitation, family violence)</p>	
Timing	No minimum duration or followup	No minimum duration or followup
Setting	Any setting; no exclusion based on geographic location or setting. Includes home, home health care, adult day care, acute care settings, social service agencies, nursing homes, assisted living, memory care units, hospice, rehabilitation centers/ skilled nursing facilities, long-distance caregiving, and nonplace-based settings	Any setting; no exclusion based on geographic locations or setting. Includes home, home health care, adult day care, acute care settings, social service agencies, nursing homes, assisted living, memory care units, hospice, rehabilitation centers/ skilled nursing facilities, long-distance caregiving, and nonplace-based settings


*We are obligated to use the proper names of measurement tools, such as the Zarit Caregiver Burden Scale, when used in a specific study.

Abbreviations: AD=Alzheimer’s disease; ADL=activities of daily living; ADRD=Alzheimer’s disease–related dementias; CNA=certified nursing assistant; CPAP=continuous positive airway pressure; ED=emergency department; IADL=instrumental activities of daily living; ICU=intensive care unit; KQ=Key Question; NIH=National Institutes of Health; PICOTS=population, intervention, comparator, outcome, timing, setting; PLWD=person/people living with dementia; PTSD=post-traumatic stress disorder

Analytic Framework

Figure 1.2 is a traditional analytic framework, illustrating the relationships between interventions and outcomes. Due to limited space, not all baseline characteristics or outcomes listed in Table 1.1 are specifically listed in Figure 1.2.

Figure 1.2. Analytic framework



Abbreviations: KQ=Key Question; PLWD=people living with dementia; SES=socioeconomic status

Report Organization

This report provides, in Chapter 2, details intended to familiarize readers with the methods used to conduct this systematic review. Chapter 3 presents the overall results of the search for the review's eligible studies. Beginning in Chapter 4, results are organized by the intervention target, then by outcome. Due to the breadth and complexity of the interventions, we also present for each result section a brief description of the intervention as well as the intervention's research context, especially regarding how the eligible studies may or may not represent how that

particular body of research has progressed along the NIH Stage Model. A glossary of terms for the report is provided in Table 1.2.

Results Chapters are structured to present each pair of KQs, keeping the PLWD and caregiver outcomes together for each intervention. And, as discussed earlier, we have categorized interventions as either care interventions (which test the effects of a specific type of care) or care delivery interventions (which test the effects of different manners in which care is delivered). We present the care interventions addressed in KQs 1-8 in Chapters 4 – 7, and the care delivery interventions addressed in KQs 8 – 9 in Chapter 8. Readers interested in considering potential implementation costs and investments (of technologies, support materials, and personnel required for intervention implementation) to suit their unique settings and contexts may find this separation helpful. Chapter 9 provides a brief response to the Guiding Question 1. The report then concludes with the Discussion in Chapter 10.

Table 1.2. Glossary of terms

Term	Description
Analytic set	For the purposes of this review, the analytic set is the set of studies that underwent synthesis. It consists of the studies not judged to be pilots or have a high potential for bias that might have interfered with the ability of the study to answer its research question.
Care delivery intervention	Care delivery interventions aim to improve the manner in which care is delivered, including the scheduling of staff and tasks as well as the ways in which staff and tasks are interdependent. Care delivery interventions can change the tasks that are performed, the set of staff who perform the tasks, or the way the staff work together. While much of this change is implemented through education and training, care delivery interventions differ from the type of education and training targeted at improving already established roles and tasks.
Care intervention	Care interventions contribute to a person's well-being, happiness, identity, privacy, capacity, autonomy, or authority. They can be supports, services, programs, accommodations, or practices that include behavioral, environmental, technological, and psychological methods or approaches. They may be delivered by healthcare, social services, and other community organizations or caregivers with the intention of having a direct impact on either a person with dementia or their caregiver or both.
Eligible study	An eligible study is one that meets the initial study criteria that were defined in advance regarding the type of study that would be included in the systematic or comparative effectiveness review.
Evidence map	An evidence map is the result of a systematic search of a defined topic area that can facilitate evidence-informed decision making or identify gaps in knowledge and future research needs.
Explanatory studies	Explanatory studies aim to test whether an intervention works under optimal situations.
Exploratory study	Exploratory studies are preliminary research designed to clarify the exact nature of the problem to be solved.
Formal caregiver	Formal caregivers are paid caregivers, such as certified nursing assistants (CNAs), home health aides, auxiliary workers, personal care aides, hospice aides, promotoras or promotores, and community health workers.
Heterogeneity	Heterogeneity is a word that signifies diversity in something. A classroom consisting of people from lots of different backgrounds would be considered having the quality of heterogeneity. Likewise, a wide range of study designs in a group of studies would be considered heterogeneous.
Informal caregiver or Caregiver/Care Partner (CG/P)	For the purposes of this review, Informal caregivers are spouses, family, friends, and volunteers providing care to one or more PLWD. Informal caregivers are typically unpaid. While caregiver has been a term commonly used in the literature, some people prefer the term "care partner."
People Living With Dementia (PLWD)	For the purposes of this review, People Living With Dementia (PLWD) is a term/ abbreviation that refers to individuals living with Alzheimer's disease or Alzheimer's disease related dementias.
Person-centered	Person-centered , in this context, is a way of designing interventions with consideration for the needs of the people using health and social services in planning, developing and monitoring care.

Term	Description
Analytic set	For the purposes of this review, the analytic set is the set of studies that underwent synthesis. It consists of the studies not judged to be pilots or have a high potential for bias that might have interfered with the ability of the study to answer its research question.
Pragmatic studies	Pragmatic studies or trials are designed to evaluate the effectiveness of interventions in real-life routine practice conditions.
Preliminary studies	A preliminary study is an initial exploration of issues related to a proposed intervention.
Protocol	A protocol is set of steps or procedures for health systems or units providing the care. They can also specify the tools and tasks that need to be carried out, and they help processes to be understood by staff regardless of staff's tenure or experience.
Pilot study	A pilot study is a small-scale test of the methods and procedures to be used on a larger scale in a future study.
Risk of bias	Risk of bias is the extent to which the design and conduct of a study are likely to have prevented bias in the results.
Small sample	Sample size is a count of the individual people or observations in any statistical setting, such as a scientific experiment or a public opinion survey. Too small a sample yields unreliable results, while an overly large sample requires a significant commitment of time and resources.

Abbreviations: CG/P=caregiver/care partner; CNA=certified nursing assistant; PLWD=person/people living with dementia

Chapter 2. Methods

The methods for this systematic review followed the Agency for Healthcare Research and Quality (AHRQ) Methods Guide for Effectiveness and Comparative Effectiveness Reviews (available at <https://effectivehealthcare.ahrq.gov/topics/ceer-methods-guide/overview>). This systematic review also reports in accordance with the Preferred Items for Reporting in Systematic Reviews and Meta-Analyses (PRISMA),¹⁶ A Measurement Tool to Assess Systematic Reviews (AMSTAR 2),¹⁷ and any relevant extension statements.

The topic of this review was initially developed by the National Institute on Aging (NIA) in consultation with AHRQ. The role of the Key Informants was filled by the NASEM committee, which, as noted earlier, will use the review to help develop its own recommendations regarding which interventions are ready for dissemination and implementation on a broad scale. However, the NASEM committee did not see the draft Key Questions (KQs), PICOTS, and analytic framework until the KQs were posted for public comment; therefore, a panel of content experts from Federal agencies acted as proxy Key Informants, providing input on the KQs to be examined. Federal content experts were drawn from the NIA, the Department of Veterans Affairs, The Department of Defense, the Center for Disease Control and Prevention, the Office of the Assistant Secretary for Planning and Evaluation, and the Administration for Community Living within the Department of Health and Human Services. The NASEM committee also served as Technical Experts, providing high-level content and methodological expertise throughout development of the review protocol. The final protocol is posted on the Effective Health Care website at <https://effectivehealthcare.ahrq.gov/products/care-interventions-pwd/protocol>.

Study Selection

Studies were included in the review based on the PICOTS framework outlined above in Table 1.1 and the study-specific inclusion criteria described in Table 2.1.

Table 2.1. Study inclusion criteria

Category	Criteria for Inclusion
Study Enrollment	Adults with possible or diagnosed AD/ADRD. No age requirement is made, that is, early onset disease that may be experienced by people with Down syndrome or other genetic risk factors are included. Study populations may include adults with mild cognitive impairment (MCI) if 15% or less of total sample, or must report results for dementia population separately.
Study Objective	KQ 1-2: Evaluate benefits and harms of care interventions for BPSD symptoms in PLWD KQ 3-4: Evaluate benefits and harms of care interventions for quality of life, function, or non-BPSD symptoms in PLWD KQ 5-6: Evaluate benefits and harms of care interventions for quality of life and health outcomes of informal caregivers for PLWD KQ 7-8: Evaluate benefits and harms of care interventions for quality of life and health outcomes of formal caregivers for PLWD KQ 9-10: Evaluate benefits and harms of care delivery interventions that address how care is delivered KQ subquestions: Evaluate possible effect modifiers of intervention benefits and harms
Study Design	RCTs, and prospective studies with concurrent comparator arms, and at least 10 participants per arm at study analysis. Interrupted time series with at least 3 measures both pre- and post-intervention (therefore excluding simple controlled before/after studies without comparator arm).

Category	Criteria for Inclusion
Study Enrollment	Adults with possible or diagnosed AD/ADRD. No age requirement is made, that is, early onset disease that may be experienced by people with Down syndrome or other genetic risk factors are included. Study populations may include adults with mild cognitive impairment (MCI) if 15% or less of total sample, or must report results for dementia population separately.
Outcomes	Outcomes listed in Table 1. Actual outcome measures will be defined by study authors. Common measures are provided in Appendix A. We will only include studies with immune function, turnover, or retention of caregivers if the study also includes another PLWD or quality outcomes; that is, we will not include the study if it only examines turnover or retention as an intermediate outcome in isolation.
Publication type	Published in peer-reviewed journals and grey literature with full text available (if sufficient information to assess eligibility and risk of bias are provided). Letters and conference abstracts are excluded due to the inability of such short publications to provide the information needed to fully describe the interventions.
Language of Publication	English only, due to resource limitations

Abbreviations: AD=Alzheimer’s disease; ADRD=Alzheimer’s disease–related dementias; BPSD=behavioral and psychological symptoms of dementia; KQ=Key Question; MCI=mild cognitive impairment; PLWD=person/people living with dementia; RCT=randomized controlled trial

The following discussion about the review search processes is organized by type of research question—first the KQs, then the guiding question.

For the KQs, we searched Ovid Medline®, Ovid® Embase, Ovid PsycInfo®, CINAHL, and the Cochrane Central Register of Controlled Trials (CENTRAL) to identify studies published and indexed in bibliographic databases. The search algorithm included relevant controlled vocabulary and natural language terms for the concepts of Alzheimer’s disease (AD) and related dementias (ADRD) (Appendix A).

We reviewed bibliographic database search results for studies relevant to our PICOTS framework and study-specific criteria. Search results were downloaded to EndNote. Two reviewers independently reviewed titles and abstracts to identify studies meeting PICOTS framework and inclusion/exclusion criteria. Two reviewers independently performed full-text screening to determine if inclusion criteria were met. Differences in screening decisions were resolved by consultation between reviewers, and, if necessary, consultation with a third reviewer. We documented the inclusion and exclusion status of citations that underwent full-text screening. Throughout the screening process, team members met regularly to discuss training material and issues as they arose to ensure consistent application of inclusion criteria.

We conducted limited additional searching of grey literature (research or other written material produced outside of traditional academic publishing) to identify relevant completed and ongoing studies that met the study design inclusion criteria. Grey literature search results were used to identify studies, outcomes, and analyses not reported in the peer-reviewed published literature to assess publication and reporting bias and inform future research needs. We also tracked published protocols for studies that have not published results in the public domain.

For the guiding question, we conducted forward citation searching of studies with low to moderate strength of evidence for companion articles describing implementation processes.

Lastly, to provide resources for care interventions which may not have been empirically studied using study designs that met inclusion criteria, we searched websites of relevant governmental agencies, professional associations, and AD or ADRD nongovernmental groups for curated lists of known interventions. An example list of organizations is provided in Appendix A.

We will update searches while the draft report is under public/peer review.


Data Extraction

Studies that met inclusion criteria were distributed to EPC reviewers for data extraction. Data extraction used a two-stage process: (1) we first used an **evidence map** table for basic data extraction, and (2) we then created a series of **analytic set** tables including comprehensive evidence and assessment tables for those studies that went on to further analysis. Figure 2.1 provides a graphic illustration of the flow of studies through the review processes. Data fields for both the evidence map and analytic set included author, year of publication, population of interest (including a granular checklist of PLWD and caregiver characteristics), intervention, comparison, setting, outcomes cited, intervention duration, and study followup.

As mentioned in Chapter 1, we were guided by the National Institutes of Health (NIH) Stage Model for Behavioral Interventions.¹⁵ This model provides a conceptual framework of intervention research development, ranging from basic science research (Stage 0) to new intervention creation (Stage 1), research-setting efficacy (Stage 2), “real-world” community-clinic efficacy (Stage 3), broad community-based effectiveness (Stage 4), to eventually dissemination and implementation research (Stage 5). This model not only describes the stages of behavioral intervention development, but also supports eventual implementation. While the stages are not a direct assessment of implementation readiness, the model suggests that interventions at Stage 3 or higher are more likely to be ready for broad dissemination. Interventions at Stage 4 that use pragmatic study designs move research closer to “real world” conditions and population levels.

For studies that appeared to be at NIH Stage 0 to 2 (pilot, feasibility, and small sample size studies), extraction was complete at the evidence map stage; *these studies did not advance to further extraction or outcome assessment*. For NIH Stage 3 (efficacy or explanatory studies) and NIH Stage 4 studies (effectiveness, or pragmatic studies), if a study was assessed as high risk of bias or over threshold risk of bias (see section below on risk of bias assessment for further details), *these studies did not advance to further extraction or outcome assessment*, however, we do present details of the risk of bias assessment.

Figure 2.1. Review assessment and extraction processes



Abbreviations: NIH=National Institutes of Health; RoB= risk of bias

The remaining studies that were assessed as having low to moderate risk of bias and appearing to be NIH Stage 3 to 5 comprise the analytic set. We extracted additional data fields including subject inclusion criteria, more detailed PLWD and caregiver characteristics, intervention and comparison characteristics, descriptions and results of included outcomes and harms, risk of bias elements, elements to distinguish NIH Stage 3 or 4 (where appropriate), and study funding source. Intervention characteristics included theory base, components and activities, timing, frequency, duration, use of technology, training, delivery approach (prescriptive or manualized vs. tailored), other delivery modalities, and use of cultural adaptations or modifications. We noted the point on the disease continuum (i.e. stage of dementia) for which the intervention was intended and methods for targeting the interventions to PLWD and/or caregivers and their identified goals and priorities.

We extracted relevant data into Microsoft Excel. Evidence map tables were verified for accuracy by a second EPC reviewer. For the analytic set, one reviewer extracted data to evidence and outcomes tables, and a second reviewer reviewed and verified the data for accuracy. Given the number of included studies, we did not contact study authors for missing data.

Assessing Methodological Risk of Bias and NIH Stage of Individual Studies

Based on AHRQ guidance,¹⁸ two EPC reviewers independently assessed risk of bias for all eligible studies. Reviewers consulted to reconcile discrepancies in overall risk of bias. Overall risk of bias assessments for each study were classified as low, moderate, or high based on the collective risk of bias inherent in each domain and the level of confidence that the results were believable given the study's limitations. However, the approach differed based on the KQ and

study NIH Stage mode; these differences are detailed below. We began with an initial sorting into NIH Stages 0 to 2 versus NIH Stages 3 to 5 by simple examination of the study aims.

For KQ 1-8: For studies of individual care interventions, we used a modified Cochrane risk of bias tool to assess them as high, medium, or low for each of the following domains: (1) selection bias (adequacy of randomization method [RCTs], accounting for imbalance in prognostic variables [observational studies]); (2) attrition bias (differentiated by mortality versus loss to followup); (3) detection bias (outcome measurement quality, outcome assessor masking); (4) performance bias (intention to treat or test analysis, adjustment for potential confounding variables, participant masking to treatment assignment); (5) reporting bias (selective reporting of outcomes). (Appendix A) While we were not expressly looking for studies identified as quality improvement interventions, we recognize that complex care delivery interventions use multicomponent approaches similar to quality improvement interventions. Therefore, for these complex interventions, risk of bias included domains similar to those outlined in a risk of bias tool for quality improvement, e.g., fidelity to the program.¹⁹

For KQs 9–10: We anticipated that care delivery studies would generally fall in the range of NIH Stage 3 to 4 effectiveness trials, with the possibility that a few were carried out as quality improvements and thus Stage 5. Along with categorizing studies by NIH Stage Model, we also broadly labeled study designs as explanatory or pragmatic. **Explanatory** studies test whether an intervention works under optimal conditions, similar to Stage 3, while **pragmatic** studies evaluate effectiveness of interventions in real-life practice conditions, similar to Stage 4.²⁰ Since study designs exist on a continuum, rather than as discrete categories, we included a “**balanced**” category for study designs that appeared poised between explanatory and pragmatic. Because both the higher NIH stages and pragmatic trials are explicitly designed to balance, or trade off, internal and external validity, we approached risk of bias assessment as a threshold requirement rather than a continuum for suspected pragmatic design studies. We targeted studies self-identifying as pragmatic and studies using advanced study designs such as cluster or stepped wedge designs. We assessed whether such studies were below the threshold of high risk of bias based on selection bias, level of attrition, and fidelity to the intervention. If a study was over threshold risk of bias, we abstracted it into the evidence map with no further action. If a study was determined to be below the threshold, we then assessed it for NIH stage. To assess NIH stage, we used a modified PRECIS-2 tool,²¹ initially developed to help interventionists design pragmatic trials. Because explanatory and pragmatic classifications fall along a continuum rather than being discrete categories,²⁰ and because reporting details for fine distinctions are often lacking in publications, we used the three categories outlined above (explanatory, balanced, and pragmatic) rather than the 5-point scale of the PRECIS-2 tool. Appendix A provides the modified tool. (Advanced study designs such as cluster trials were considerably less prevalent in KQ 1 – 8 and often readily identified as explanatory based on the stated purpose of the study and the reason for using a cluster design. We therefore found it essentially unnecessary to use this staged assessment process for KQ 1 – 8.)

Data Synthesis

We summarized results in evidence maps or analytic set tables and synthesized evidence for each unique population, comparison, and outcome or harm. *Evidence maps* provide a quick synthesis of what the identified literature has studied. *Analytic set tables* gather a more comprehensive set of data, allowing the reviewer to attempt to go further and answer “what did the included studies find.” For this review, we organized analytic set tables by intervention

targets, interventions, comparators, and PLWD, caregiver, or other system-level outcomes. We reported descriptive information regarding numbers of studies reporting more granular PLWD or caregiver characteristics.

Because we could not identify a consensus taxonomy of interventions to apply to the literature, we categorized interventions empirically by intervention and comparator pairs. Because splitting into very narrow categories can make drawing conclusions impossible (due to few studies of often questionable risk of bias), we sought to balance two competing concerns: (1) a need to group broadly conceptual ideas together as much as possible in order to have sufficient studies informing the synthesis, and (2) avoiding excessive heterogeneity in the studies grouped together because that makes interpretation difficult. We used the explanatory or pragmatic classifications along with the NIH Stage Model to inform our qualitative synthesis of the **intervention’s research context** (i.e., a brief representation of the current state of the research and its development over time, including a summary and description of the eligible studies). The intervention research context sections function as the main results reporting for the evidence maps, and as contextual information for outcomes reported for the analytic sets.

For the KQs, we assessed the effects of outcomes using clinically important differences if well-established, but for many outcomes this was not the case. Because of the very wide range of outcomes of interest across the panel of potential interventions, we did not list specific priority outcomes beyond those noted in Table 1.1. For any individual study, we examined no more than five to seven outcomes per PLWD or caregiver population, prioritizing person-centered outcomes, (e.g., quality of life, function, and harms), over intermediate outcomes (e.g., laboratory test values, subscales of outcome measurement tools). Our rationale for this decision is that excessive reporting of outcomes generally happens with the latter type of outcome.

When pooling outcomes across studies was possible, we used random effects models. For continuous outcomes, we calculated weighted mean differences and/or standardized mean differences with the corresponding 95 percent confidence intervals. We assessed the clinical and methodological heterogeneity and variation in effect size to determine appropriateness of pooling data. We assessed statistical heterogeneity with Cochran’s Q test and measure magnitude with I² statistic.

Grading the Strength of Evidence for Major Comparisons and Outcomes

The overall strength of evidence for select outcomes for KQs 1 – 10 were evaluated based on five required domains: (1) study limitations (risk of bias); (2) consistency (similarity of effect direction and size); (3) directness (single, direct link between intervention and outcome); (4) precision (degree of certainty around an estimate); and (5) reporting bias.²² An outcome with an overall rating of “high strength of evidence” implies that the included contributing studies were randomized controlled trial studies with both a low risk of bias, and with consistent, direct, and precise domains. We assessed strength of evidence for key final health outcomes measured with validated scales.

Based on study design and risk of bias, we rated study limitations as low, medium, or high. Consistency was rated as consistent, inconsistent, or unknown/not applicable (e.g., single study) based on whether intervention effects were similar in direction and magnitude, and statistical significance of all studies. Directness was rated as either direct or indirect based on the need for indirect comparisons when inference requires observations across studies (i.e., more than one step was needed to reach the conclusion). Precision was rated as precise or imprecise based on

the degree of certainty surrounding each effect estimate or qualitative finding. An imprecise estimate is one for which the confidence interval is wide enough to include clinically distinct conclusions. If we had found any outcome to have at least moderate or high strength of evidence, we would have evaluated reporting bias by the potential for publication bias, selective outcome reporting bias, and selective analysis reporting bias. We would have done this by comparing reported results with those mentioned in the methods section and an assessment of the grey literature to assess potentially unpublished studies. However, no findings rose to this level. Other factors considered in assessing strength of evidence included weighting by strength of study design to address broad dissemination (thus pragmatic trials hold stronger weight), dose-response relationship, the presence of confounders, and strength of association.

Based on these factors, we rated the overall strength of evidence for each outcome as:

High: Very confident that estimate of effect lies close to true effect. Few or no deficiencies in body of evidence, findings are believed to be stable.

Moderate: Moderately confident that estimate of effect lies close to true effect. Some deficiencies in body of evidence; findings likely to be stable, but some doubt.

Low: Limited confidence that estimate of effect lies close to true effect; major or numerous deficiencies in body of evidence. Additional evidence necessary before concluding that findings are stable or that estimate of effect is close to true effect.


Insufficient: No evidence, unable to estimate an effect, or no confidence in estimate of effect. Available evidence or lack of evidence precludes judgment.

Notably, an assessment of insufficient evidence does not mean that the intervention is ineffective. Rather, it means that due to the uncertainty of the evidence, we could not draw meaningful conclusions about its effectiveness at this time.

Chapter 3. Search Results

Our search identified 9,217 publications for screening. Based on inclusion criteria, we identified 595 unique eligible studies discussed in 850 publications. We list studies excluded at full text screening, by exclusion category, in Appendix B. See Figure 3.1 for details of the screening process.

Figure 3.1. Literature flow diagram



Of the eligible 627 unique studies, we categorized 409 studies as pilot or small sample studies which thus appeared to be National Institutes of Health (NIH) Stage Model Stage 0 – 2 (i.e., for newer interventions that would not be appropriate to address readiness for implementation). A further 218 were assessed as high risk of bias, most commonly due to issues with selection bias, attrition, or intervention fidelity. This left 100 low to moderate risk of bias studies for the analytic set. (Table 3.1) Non-U.S. studies were a large portion of the eligible studies. Overall, approximately 67 percent of the literature emerged from research conducted outside of the United States. This was heavily weighted by care interventions for treating or managing behavioral or psychological symptoms of dementia for people living with dementia (PLWD) (Chapter 4) or PLWD well-being (Chapter 5).

Unfortunately, many important groups warrant further inclusion in research, including people with Down syndrome, who are living longer and who overall experience higher rates of dementia than the general population. Individuals with complex presentations of dementias, e.g., early-onset, amnesic forms, also need further consideration. We also found little information to determine whether interventions are equally appropriate for or have been successfully adapted to other race/ethnic cultures within the United States, with only a handful of studies providing

sufficient inclusion of African-American or Hispanic/Latino populations. Interventions for rural communities were found to be even more rare in the literature. Additionally, the many countries in which these care interventions were evaluated were almost exclusively high-resource, with very few low-resource countries represented. Finally, few studies gathered the granular detail necessary for a deeper understanding of the applicability of the interventions. For example, the minority of studies that measured PLWD disability used medical approaches, like the Charlson score, rather than functional approaches that note disability prior to dementia onset.

Based on the structure of the Key Questions and the eligible studies identified in the search process, we developed categories and assigned each study (or comparison within a multi-arm study) to one of 37 major intervention categories. A few interventions exhibited characteristics that could be ascribed to more than one category. Further, some care interventions are variably defined and described within the literature, making categorization more challenging. We attempted to keep clearly defined interventions together and classified less clearly defined interventions into more general categories such as psychosocial therapies or multicomponent interventions. If an intervention’s characteristics were balanced between a care intervention and a care delivery intervention, we tended to classify the intervention into the care intervention category and present it in Chapters 4 – 7.

Table 3.1. Identified unique eligible studies by intervention category, by results chapter

Location	Intervention	Total Unique Studies	# Analytic Set	# Evidence map	# non-U.S.
Chapter 4 Managing PLWD BPSD	Assisted Therapy	16	1	15	15
	Multisensory Stimulation/Snoezelen	9	2	7	6
	Complementary and Alternative Medicine (CAM) Therapies	21	5	16	19
	Bright Light Therapy	9	0	9	4
	Psychosocial Therapies for BPSD	6	0	6	1
	Multicomponent Interventions for BPSD	9	0	9	5
	Chapter 4 TOTAL		70	8	62
Chapter 5 PLWD Wellbeing	Exercise	53	10	43	48
	Music	35	5	30	26
	Reminiscence Therapy	25	4	21	22
	Cognitive Rehabilitation	23	3	20	19
	Cognitive Training	18	5	13	15
	Cognitive Stimulation Therapy	12	2	10	11
	Recreation Therapy	14	3	11	11
	Psychosocial Interventions for PLWD well-being	7	0	7	4
	Creative Expression Therapy	5	0	5	2
	Multicomponent Interventions	24	3	21	14
	Assistive Technology	4	0	4	2
	Electrostimulation	14	0	11	12
	Other Interventions for PLWD well-being	7	0	7	4
Chapter 5 TOTAL		241	35	207	190
Chapter 6 Informal Caregivers	Psychosocial Interventions for Informal Caregiver Wellbeing	122	29	93	74
	Social Support	13	2	11	10
	Lifestyle Interventions	19	1	18	10
	Respite Care	3	0	3	0
	Multicomponent for Informal Caregivers	22	7	15	13

Location	Intervention	Total Unique Studies	# Analytic Set	# Evidence map	# non-U.S.
	Other Interventions for Informal Caregiver Wellbeing	6	0	6	4
	Chapter 6 TOTAL	185	39	146	111
Chapter 7 Formal Caregivers	Formal Caregiver Wellbeing	3	0	3	3
	Chapter 7 TOTAL	3	0	3	3
Chapter 8 Care Delivery	Care Service Provision	50	6	44	29
	Consultation Services	5	1	4	4
	Case Management	9	3	6	6
	Care Protocols for PLWD	17	0	17	11
	Advance Care Planning	9	1	8	3
	Palliative Care	5	0	5	3
	Other Service Provision Interventions	5	1	4	2
	Care Delivery Models or Programs	31	12	18	18
	Care Delivery Staff Education and Support Needs	46	1	45	27
	Caregiver Staff Training	22	0	22	18
	Informal Caregiver Staff Training	12	0	12	6
	Family Education and Partnering	5	0	5	1
	Mutitier Training	7	1	6	2
	Chapter 8 TOTAL	127	19	107	74

Abbreviations: BPSD=behavioral and psychological symptoms of dementia; CAM=complementary and alternative medicine; PLWD=person/people living with dementia

Chapter 4. Care Interventions for Managing BPSD in PLWD

This chapter includes care interventions that aimed to address behavioral or psychological symptoms of dementia (BPSD) in people living with dementia (PLWD). Studies in this chapter enrolled PLWD currently identified as experiencing BPSD, the term most commonly used in this literature set. In contrast, Chapter 5 includes studies of interventions aimed at improving the general well-being of PLWD (e.g., exercise, music) and tested with PLWD regardless of presence of BPSD. Care delivery interventions specifically designed to address BPSD in PLWD are presented in Chapter 8.

For each intervention, we present Key Points followed by results in three general sections: Intervention Description, Eligible Studies, and Intervention Research Context. For interventions with no eligible studies assessed as low to medium risk of bias, we present the studies from the evidence map with a brief discussion of what has been examined and the research context. For interventions for which low- to medium-risk-of-bias studies were available for an analytic set, we present Outcomes sections by PLWD, by caregiver, and by variation in outcomes when available. Because differences in outcome measures and intervention complexity prohibited combining outcomes for a statistical meta-analysis, we present summary findings as brief statements of how many studies reported statistically significant benefit or no difference between the intervention and the comparator. Detailed information on all eligible studies can be found in Appendix C.

Assisted Therapy

Assisted therapy interventions aim to reduce BPSD in PLWD with the assistance of an animal or an object that represents a living being. Eligible studies examined therapy with robots, dogs, and dolls. While the use of such interventions does not depend on the setting, all studies examined the use of assisted therapy in nursing homes or other long-term care settings.

Robot-Assisted Therapy

Key Point

- Evidence was insufficient to draw conclusions about the effect of robot-assisted therapy on PLWD and their caregivers.

Intervention Description

Robot-assisted therapy studies primarily focused on the use of PARO, an autonomous, robotic baby harp seal.

Eligible Studies

Table 4.1 summarizes the characteristics of the literature set. We identified eight unique studies from 13 publications that examined robot-assisted therapy for the treatment of dementia.²³⁻³³ Four studies were pilots or small sample studies^{30, 31, 33, 34} and another three were assessed as high risk of bias;^{27-29, 32} therefore, these six studies were excluded from the analytic set.^{27-29, 32} We present information on all pilot studies and high risk of bias studies in the evidence map in Appendix C.

The remaining study (n=415) was a three-arm cluster randomized trial comparing PARO versus usual care and PARO with the robotic features deactivated in PLWD in Australian nursing homes.²³⁻²⁶ The study was assessed as medium risk of bias and categorized as explanatory, or Stage 3 of the National Institutes of Health Stage Model. It enrolled individuals living in long-term care facilities with a documented dementia diagnosis (all types and severities). Appendix C provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 4.1. Basic characteristics of literature set: robot-assisted therapy

Characteristic	Information
Total Studies	8 studies
Non-U.S. studies	7 studies
Evidence map studies	2 pilot studies 2 small sample studies 3 high risk of bias studies
Analytic set studies	1 cluster randomized controlled trial
Risk of bias of analytic set	1 medium
Number of PLWD in analytic set	415
Dementia type/definition	All dementia types and severities
Caregiver type (number)	Not reported

Abbreviations: PLWD=person/people living with dementia

Intervention Research Context

The literature on robot-assisted therapy generally follows the NIH Stage Model, and consists primarily of pilots. With the exception of one U.S. study,³² the studies were conducted in Australia,^{31, 34} New Zealand,³⁰ the Netherlands,²⁷ Norway,²⁸ and Spain.³³ We identified two pilots comparing PARO versus usual care or an attention control.^{30, 31, 34} We also identified a pilot comparing PARO versus live animal therapy, a human-like robot, and usual care.³³ The PARO study included in our analytic set appears to be one of the first larger cluster RCTs of sufficient methodological rigor that compares PARO versus usual care.²³⁻²⁶ The study also appears to be one of the first larger cluster RCTs to compare PARO versus PARO with robotic features deactivated. While the study also included outcomes related to motor activity and sleep patterns, these outcomes were exploratory and data were available for less than half the study population.²⁵

PLWD Outcomes

Evidence was insufficient to draw conclusions about either PARO versus usual care (n=257) or PARO versus PARO with robotic features deactivated (n=278).²³⁻²⁶ Table 4.2 summarizes the primary findings. Results for reduction in agitation were mixed. PLWD showed more visual and verbal engagement with PARO than with PARO with robotic features deactivated. The two groups did not differ for positive behavioral engagement or social engagement. No harms were assessed.²³⁻²⁶ An assessment of insufficient evidence does not mean that the intervention is determined to be of no value. Rather, it means that due to the uncertainty of the evidence we could not draw meaningful conclusions at this time.

Table 4.2. Summary of findings for PLWD outcomes: robot-assisted therapy

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Agitation measures PARO vs. usual care	1 cluster RCT ²⁶ (n=257) 10 weeks	All dementia severities and types in Australia	1 found benefit 1 found no difference	Insufficient
Agitation measures PARO vs. deactivated PARO	1 cluster RCT ²⁶ (n=278) 10 weeks	All dementia severities and types in Australia	0 found benefit 2 found no difference	Insufficient
Engagement measures PARO vs. deactivated PARO	1 cluster RCT ²⁶ (n=278) 10 weeks	All dementia severities and types in Australia	2 found benefit 2 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: n=number; RCT=randomized controlled trials

Caregiver Outcomes

The analytic study set did not report outcomes related to informal or formal caregivers.

Variation in Outcomes

In the PARO group, lower levels of agitation at baseline were associated with greater behavioral positive engagement and visual engagement at 10 weeks.²⁶ No similar analysis was conducted for the PARO with robotic features deactivated group. The analytic study set did not report variation in outcomes by caregiver or intervention characteristics.

Live Animal- and Doll-Assisted Therapy

Key Point

- Studies of live animal- and doll-assisted therapy were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Live animal-assisted therapy involves treatment sessions that include animals with a focus on specific goals (e.g., mental, emotional and/or social). Doll-assisted therapy involves offering dolls to PLWD in order to improve their comfort, engagement, and quality of life.

Eligible Studies

We identified five unique studies from five publications that examined animal-assisted therapy (with dogs) in the treatment of dementia. (Table 4.3) Two were pilot or small sample studies and the remaining three studies were assessed as high risk of bias.³⁵⁻³⁹ Three unique pilot studies examined doll-assisted therapy.⁴⁰⁻⁴² We present information on all pilot studies and high risk of bias studies in the evidence map in Appendix C.

Table 4.3. Basic characteristics of literature set: live animal- and doll-assisted therapy

Characteristic	Information
Total Studies	8 studies
Non-U.S. studies	8 studies

Characteristic	Information
Evidence map studies	5 pilot studies 3 high risk of bias
Analytic set studies	0 studies

Intervention Research Context

One study compared the use of doll-therapy versus an active control of hand warmers, which would mimic the sensory characteristics of holding a doll, in PLWD residing in an Italian nursing home.⁴² A second compared doll-therapy to teaching PLWD to use gestures to improve communication in Italian nursing homes.⁴⁰ An third study compared doll-therapy to usual care in Australian nursing homes.⁴¹ One study compared individual animal therapy versus usual care in PLWD in German nursing homes,³⁵ while another used group animal therapy in Italian adult daycare centers.³⁶ Two others compared group animal therapy versus usual care in PLWD in nursing homes in Norway,^{37, 38} while the third compared group animal therapy versus group therapy without an animal in Australian nursing homes.³⁹

Multisensory Stimulation/Snoezelen

Key Point

- Evidence was insufficient to draw conclusions about the effect of multisensory stimulation (MSS) on PLWD and their caregivers.

Intervention Description

MSS is intended to have both relaxing and activating effects that promote calm engagement for PLWD in nursing homes through an experiential process that includes light, sound, scents, and music, usually with the accompaniment of an aide or therapist. The “Snoezelen rooms” often used for this intervention were developed in the Netherlands in the 1970s, and are designed to deliver stimuli to various senses, including through different tactile materials and floors that may be adjusted to stimulate the sense of balance.

Eligible Studies

Table 4.4 summarizes the characteristics of the literature set. We identified nine unique studies from 11 publications.⁴³⁻⁵⁰ Two studies (N = 256) were assessed as medium risk of bias and classified as explanatory, or Stage 3 of the NIH Stage Model.^{44, 45} One was a two-arm randomized trial comparing Snoezelen versus activity sessions, which were treated as an active control.⁴⁴ The other was a three-arm randomized trial comparing Sonas, a different type of MSS intervention, to reading sessions and to usual care.⁴⁵ Appendix C provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.⁴⁶⁻⁵² Pilot studies, small sample studies, or studies assessed as high risk of bias were excluded from the analytic set. We present information on these studies in the evidence map in Appendix C.

Table 4.4. Basic characteristics of literature set: multisensory stimulation/Snoezelen

Characteristic	Information
Total studies	9 studies
Non-U.S. studies	6 studies

Characteristic	Information
Evidence map studies	2 pilot studies 4 small sample studies 1 high risk of bias studies
Analytic set studies	2 randomized controlled trials
Risk of bias of analytic set	2 medium
Number of PLWD in analytic set	256
Dementia type/definition	Diagnoses of Alzheimer's, vascular, or mixed dementia
Caregiver type (number)	Not reported

Abbreviation: PLWD=people living with dementia

Intervention Research Context

This literature set did not clearly follow progression along the NIH Stage Model. Earlier studies were either pilots⁵⁰ or full studies that reported a pilot component^{43, 44} However, later studies did not appear to progress to pragmatic trials. The studies generally used active controls, such as activity or reminiscence sessions.

PLWD Outcomes

Evidence was insufficient to draw conclusions about the effect of MSS on BPSD compared with an active control.⁴⁴ There was no significant mean change on several measures of behavioral problems in the study. Table 4.5 summarizes these findings. Similarly, evidence was insufficient to draw conclusions about the effect of MSS on social function versus both an attention control and usual care.⁴⁵ There was a statistically significant benefit for MSS versus both comparators on improvements in PLWD communication.

Table 4.5. Summary of findings for PLWD outcomes: multisensory stimulation/Snoezelen

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Neuropsychiatric symptoms MSS vs. attention control	1 RCT ⁴⁴ (n=136) 4 weeks, 8 weeks	Persons with Alzheimer's, vascular, or mixed dementia in Norway	0 found benefit 1 found no difference	Insufficient
Social function MSS vs. attention control	1 RCT ⁴⁵ (n = 120) 24 weeks	Persons with Alzheimer's, vascular, or mixed dementia in Norway	1 found benefit 0 found no difference	Insufficient
Social function MSS vs. usual care	1 RCT ⁴⁵ (n = 120) 24 weeks	Persons with Alzheimer's, vascular, or mixed dementia in Norway	1 found benefit 0 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: MSS=multisensory stimulation; n=number; RCT=randomized controlled trial

Caregiver Outcomes

No studies in the analytic set reported caregiver outcomes.

Variation in Outcomes

In PLWD with severe cognitive impairment (Mini-Mental State Examination (MMSE) scores 0 to 9), MSS appeared slightly more effective than the activity control at improving behavior at 4 weeks (mean difference -1.0 points, $p < 0.05$).⁴⁴ For PLWD with moderate cognitive impairment,

activity sessions appeared to improve the behavior more than MSS (mean difference 0.8 points, p-value not significant).⁴⁴ However, this trend was not seen with other outcomes.

In PLWD with severe cognitive impairment (MMSE scores 0 to 10), MSS was reported more effective than the active control group at improving HCS scores at 12 weeks, but not at 24 weeks. In addition, MSS did not show a significant advantage in improving HCS scores when compared with usual care.⁴⁵

Complementary and Alternative Medicine (CAM) Therapies

Key Point

- Evidence was insufficient to draw conclusions about the effects of aromatherapy or foot massage on PLWD and their caregivers.

Intervention Description

CAM is a set of therapies that are intended to achieve health effects, but are not part of conventional medical practice. While this category has the potential to be very broad, only a few interventions were examined in the eligible literature, mainly aromatherapy, massage, acupuncture, and healing touch. Interventions that involve physical movement, such as yoga or tai chi/taiji, are presented with the section on exercise in Chapter 5.

Eligible Studies

Table 4.6 summarizes the characteristics of the literature set. We identified 21 unique studies from 23 publications that examined CAM for the treatment of dementia.⁵³⁻⁷¹ Five studies were assessed as low to moderate risk of bias and included in the analytic set.^{57, 67, 72-74} Four studies (n=278) evaluated the effects of aromatherapy in the form of lavender and lemon-balm oils versus either sunflower oil or usual care in the United Kingdom,^{72, 73} Hong Kong,⁷⁴ or Taiwan.⁶⁷ The fifth study (n=55) assessed the effects of foot massage for PLWD versus attention control.⁵⁷ No study in the analytic set was conducted in the United States. Appendix C provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes. The evidence map in Appendix C provides information on all other studies.^{53-56, 58-66, 68, 71, 75}

Table 4.6. Basic characteristics of literature set: complementary and alternative medicine

Characteristic	Information
Total studies	21 studies
Non-U.S. studies	19 studies
Evidence map studies	6 pilot studies 8 small samples 2 high risk of bias studies
Analytic set studies	5 randomized controlled trials
Risk of bias of analytic set	4 medium (aromatherapy) 1 low (foot massage)
Number of PLWD in analytic set	278 (aromatherapy) 5 (foot massage)
Dementia type/definition	All dementia types and severity
Caregiver type (number)	Not reported

Abbreviation: PLWD=people living with dementia

Intervention Research Context

As evaluated against the NIH Stage Model, this literature is overall preliminary and based on small sample studies that have not developed beyond a basic explanatory stage, or Stage 3. Evidence map studies examined various CAM techniques such as healing touch and body talk cortices,⁶⁰ acupressure/acupuncture,^{61, 64} mixed aroma exposure,⁶² back and leg massage with moisturizing cream,⁷⁰ aromatherapy,^{53, 54, 58, 66, 67, 71, 76} aromatherapy and hand massage,⁵⁵ aromatherapy massage,⁵⁶ aromatherapy massage plus acupressure,^{59, 67} and therapeutic touch.^{65, 68} Evidence map trials are from Hong Kong,^{59, 64} China,⁶¹ Japan^{53, 62} Taiwan,⁶⁷ Canada,^{65, 68} Australia,^{55, 58} Spain,⁷⁰ United States,^{54, 58-60} the United Kingdom,⁵⁶ and Israel.⁶⁶

PLWD Outcomes

Evidence was insufficient for all outcomes. The most commonly investigated outcomes were agitation and neuropsychiatric symptoms. Table 4.7 summarizes the number of studies investigating each outcome and the number of studies that found a benefit and no difference between PLWD and comparison groups.

Agitation was investigated by three of the four included studies. One found that lavender oil versus sunflower oil resulted in short-term improvements (3 weeks).⁷⁴ Another compared lavender plus orange oil versus usual care and found no improvement,⁶⁷ while two other studies found mixed results, with lemon balm oil versus sunflower oil reducing agitation at 4 weeks,⁷² but not 12 weeks.⁷³

Depression was investigated in one study that compared lavender plus orange oil versus usual care, and reported improved depression scores with the intervention at 9 weeks.⁶⁷

Neuropsychiatric symptoms were investigated in two studies comparing lavender oil and lemon balm oil versus sunflower oil. Lavender oil seemed to improve neuropsychiatric symptoms at 3 weeks,⁷⁴ but not at 12 weeks.⁷³

Quality of life and activities of daily living were examined in one study that evaluated lemon balm oil versus sunflower oil and reported that quality of life seemed to improve over 12 weeks, but activities of daily living did not.⁷³

One study examined foot massage versus attention control.⁵⁷ Both increased agitation in PLWD, but the increase was greater in the comparator group than in the intervention group (p=0.03).⁵⁷

Table 4.7. Summary of findings for PLWD outcomes: complementary and alternative medicine

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Finding	Strength of Evidence*
Agitation Lavender vs sunflower oil	1 RCT ⁷⁴ (n=70) 3 weeks	PLWD with significant agitation in Hong Kong	1 found benefit 0 found no difference	Insufficient
NPI Lavender vs sunflower oil	1 RCT ⁷⁴ (n=70) 3 weeks	PLWD with significant agitation in Hong Kong	1 found benefit 0 found no difference	Insufficient
Agitation Lavender and orange oil vs. Usual care	1 RCT ⁶⁷ (n=59) 9 weeks	PLWD with mild to severe dementia in Taiwan	0 found benefit 1 found no difference	Insufficient
Depression Lavender and	1 RCT ⁶⁷ (n=59) 9 weeks	PLWD with mild to severe dementia in Taiwan	1 found benefit 0 found no difference	Insufficient

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Finding	Strength of Evidence*
orange oil vs. usual care				
Agitation Lemon Balm oil vs sunflower oil	2 RCT ^{72, 73} (n=149) 4 - 12 weeks	PLWD with agitation in the United Kingdom	1 of 2 found benefit 1 of 2 found no difference	Insufficient
NPI Lemon Balm oil vs sunflower oil	1 RCT ⁷³ (n=77) 12 weeks	PLWD with agitation in the United Kingdom	0 found benefit 1 found no difference	Insufficient
QoL Lemon Balm oil vs sunflower oil	1 RCT ⁷³ (n=77) 12 weeks	PLWD with agitation in the United Kingdom	1 found benefit 0 found no difference	Insufficient
Caregiver Burden Lemon Balm oil vs sunflower oil	1 RCT ⁷³ (n=77) 12 weeks	PLWD with agitation in the United Kingdom	0 found benefit 1 found no difference	Insufficient
Agitation Foot massage vs attention control	1 RCT ⁵⁷ (n=55) 3 weeks	Moderate to late stage PLWD in Australia with history of agitation	1 found benefit for control group	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: n=number; NPI=Neuropsychiatric Inventory; PLWD=people living with dementia; QoL=quality of life; RCT=randomized controlled trial

Caregiver Outcomes

The analytic study set did not report outcomes related to informal or formal caregivers.

Variation in Outcomes

The analytic set did not report variations in outcomes by PLWD, caregiver, or intervention characteristics.

Bright Light Therapy

Key Point

- Studies on bright light therapy were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Bright light therapy interventions focus on technology to influence the circadian rhythm of PLWD in order to improve a range of psychiatric symptoms and behavioral disturbances including sleep disruption, agitation, or depression.

Eligible Studies

We identified nine unique studies from 12 publications that examined the use of bright light therapy interventions for individuals with dementia.⁷⁷⁻⁸⁵ (Table 4.8) Excluded from the analytic set are five pilot or small sample studies and four studies assessed as high risk of bias. We present information on all studies in the evidence map in Appendix C.

Table 4.8. Basic characteristics of literature set: bright light therapy

Characteristics	Information
Total unique studies	9 studies
Non-U.S. studies	4 studies
Evidence map studies	1 pilot 4 small sample studies 4 high risk of bias studies
Analytic set studies	0

Intervention Research Context

This literature set is preliminary. Studies examined bright light therapy on PLWD^{77-79, 81-83, 86} tailored lighting,⁸⁵ or dawn-dusk simulation, a “naturalistic” form of light therapy.⁸⁰ Publications date from 1998, but U.S.-based studies continued until 2019, while the last non-U.S. study was published in 2009. Non-U.S. settings included Japan,⁸² Switzerland,⁸⁰ and the Netherlands.⁸⁶ Nursing home or long-term care facilities were the most common setting with the exception of one study that took place in the geriatric unit of a psychiatric hospital.⁸¹

Psychosocial Therapies for BPSD

Key Point

- Studies on psychosocial therapies were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Psychosocial therapies for PLWD include a diverse set of interventions, ranging from cognitive behavioral training for anxiety to individualized social activities to improve sleep to validation group therapy.

Eligible Studies

We identified six unique studies from seven publications that examined the use of psychosocial interventions for individuals with dementia.⁸⁷⁻⁹² (Table 4.9) Excluded from the analytic set are four pilot or small sample studies and two studies assessed as high risk of bias.⁸⁷⁻⁹² We present information on all studies in the evidence map in Appendix C.

Table 4.9. Basic characteristics of literature set: psychosocial therapies for PLWD

Characteristics	Information
Total unique studies	6 studies
Non-U.S. studies	1 study
Evidence map studies	1 pilot 3 small sample studies 2 high risk of bias studies
Analytic set studies	0

Abbreviation: PLWD=person/people living with dementia

Intervention Research Context

This literature set is preliminary. One study examined problem adaptation therapy for depression and suicidal ideation in U.S. PLWD with early stage dementia.⁹¹ One study examined the effect of individualized social activity on disruptive behaviors,⁹⁰ while another examined on sleep patterns in PLWD in U.S. nursing homes.⁹² Two studies examined the use of cognitive

behavioral training to control anxiety in PLWD living in the community with mild to moderate dementia in the United Kingdom⁸⁸ and United States.⁸⁹ An early 1997 publication examined validation therapy in PLWD in U.S. nursing homes.⁸⁷

Multicomponent Interventions for BPSD

Key Point

- Studies of various unique multicomponent interventions were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

This section encompasses a small and diverse set of interventions combining several components targeted at improving BPSD.

Eligible Studies

We identified nine unique studies from 10 publications that examined the use of various multicomponent interventions among PLWD.^{90, 93-100} (Table 4.10) All the studies were either small; sample studies or assessed as high risk of bias; we describe them in the evidence map in Appendix C.

Table 4.10. Basic characteristics of literature set: multicomponent interventions for BPSD

Characteristics	Information
Total unique studies	9 studies
Non-U.S. studies	5 studies
Evidence map studies	1 pilot 3 small sample studies 5 high risk of bias
Analytic set studies	0

Abbreviation: BPSD=behavioral and psychological symptoms of dementia

Intervention Research Context

This research is generally preliminary, consisting of a small set of distinct and mostly unrelated studies. One study used question-asking, reading, reminiscence, and cognitive behavioral techniques, as well as environmental supports and individualized behavioral activity for depression in PLWD in U.S. nursing homes.⁹⁴ One small study examined an intervention in U.S. nursing homes to balance periods of high and low arousal in PLWD throughout the day.⁹⁶ An earlier 1998 study used a somewhat similar stimulation-retreat model.⁹⁷ One study conducted in Taiwan long-term care facilities examined a combined acupressure and Montessori-based activity to address anxiety in PLWD.⁹⁸ One community-based study in Hong Kong examined self-management support plus therapeutic exercise for knee osteoarthritis for PLWD.⁹³ One study in a German nursing home modified a cognitive engagement program for people with autism and combined it with music therapy for PLWD with moderate dementia.⁹⁵

One set of two studies on emotion-oriented care were conducted in the Netherlands.^{99, 100} Emotion-oriented care is a combination of validation therapy and sensory stimulation (discussed briefly in the sections above) and reminiscence therapy, which we present in Chapter 5. Although publications did not clarify whether authors of the separate studies collaborated on study development and purpose, the study by Schrijnemaeker and colleagues tested an

intervention with more elements aimed at broader intervention implementation.⁹⁹ Reported results between the two studies were mixed.

The study by Beck and colleagues examined several groups of PLWD in U.S. nursing homes.⁹⁰ It is also included in this chapter's section on psychosocial interventions and Chapter 5's section on cognitive rehabilitation. We present it here because one study group used a combination of ADL-focused cognitive rehabilitation plus the psychosocial engagement activity.⁹⁰

Conclusion

We found 63 unique studies from 76 publications that coalesced into six categories of care interventions aimed at managing BPSD in PLWD. Applying the framework for care interventions from the NASEM Families Caring for an Aging America 2016 report (Figure 1.1 in Chapter 1), the vast majority of the care interventions were delivered at the individual level. Almost 90 percent of the studies were pilots or small sample studies, Stage 0 to 2 according to the NIH Stage Model, or assessed as high risk of bias. Of the three care interventions that had low to moderate risk of bias studies—robot-assisted therapy, multisensory stimulation/Snoezelen, and CAM—we found the uncertainty of the evidence was too high to draw conclusions. However, our being unable to draw a conclusion does not mean that the intervention has no effect. Research on interventions to change behavioral and psychological outcomes is challenging, and many factors can influence the outcomes. Future research may reduce uncertainty enough to allow for conclusions about the effect of these interventions to be made with greater confidence.

Chapter 5. Care Interventions for PLWD Well-Being

This chapter includes care interventions intended to improve the quality of life and well-being of people living with dementia (PLWD). Interventions use a wide range of approaches from physical to cognitive to environmental, both alone and in combination. Most studies did not base study enrollment on the absence or presence of behavioral or psychological symptoms of dementia (BPSD).

We organized this chapter by the most commonly studied interventions first, followed by whole person, multisensory stimulation, and various “other” intervention categories, including assistive technology. For each intervention, we present Key Points followed by results in three general sections: Intervention Description, Eligible Studies, and Intervention Research Context. For interventions with no studies assessed as low to medium risk of bias, we present the studies from the evidence map with a brief discussion of what has been examined and the research context. For interventions for which low- to medium-risk-of-bias studies were available for an analytic set, we present Outcomes sections by PLWD, by caregiver, and by variation in outcomes when available. Because we were generally unable to pool outcomes for any given intervention and comparison group, we synthesized the information qualitatively; therefore, we present summary findings as brief statements of how many studies reported statistically significant benefits or no difference between the intervention and the comparator. We present detailed information on all eligible studies in Appendix D.

Exercise

Key Point

- For both community-dwelling PLWD and PLWD living in residential care facilities, evidence was insufficient to draw conclusions about the effects of exercise interventions on PLWD and their informal caregivers or care partners (CG/P).

Intervention Description

Exercise interventions are used to improve daily function, aerobic conditioning, strength, gait and balance. While exercise can be done in a wide range of activities and intensity, most exercise interventions compared moderate- to high-intensity aerobic plus strength training exercise with usual care or seated group activities. The type and duration of exercise varied widely. Aerobic training usually involved walking or stationary cycling. Strength training involved repetitive functional maneuvers that relied on body weight, or standard weight training via gym or therapy equipment. Specific balance training was uncommon and involved functional maneuvers conducted with a therapist.¹⁰¹ More often, improvements in balance and gait speed were anticipated outcomes from general aerobic and lower extremity strengthening interventions. Exercise programs commonly included a build-up phase. Settings varied from unidentified areas of nursing homes to gyms, adult daycare, outpatient rehabilitation, homes, or church halls. Most exercises were supervised and conducted in small groups. Those supervising the interventions had varied training and experience. Only two studies also involved incorporating CG/P in exercise training activities.^{102, 103}

Eligible Studies

Below, we provide a summary of exercise intervention studies for adults with dementia (Table 5.1). We identified 53 unique studies from 69 eligible publications that reported the effects of exercise interventions on quality of life and functional outcomes in PLWD. Of these, eight randomized trials (RCT) with low or medium risk of bias were included in the analytic set.^{101, 102, 104-111} (Table 5.1). Information on all pilot, small sample, and high risk of bias studies is provided as part of the evidence map in Appendix D.^{75, 103, 112-150} Appendix D provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes of the analytic set.

Table 5.1. Basic characteristics of literature set: exercise

Characteristic	Information
Total studies	53 studies
Non-U.S. studies	48 studies
Evidence map studies	13 pilot 21 small sample studies 9 high risk of bias
Analytic set studies	7 randomized controlled trials 3 cluster randomized controlled trials
Risk of bias of analytic set	2 low, 8 medium
Number of PLWD in analytic set	1,706
Dementia type/definition	Mostly mild to moderate Alzheimer's disease, although the range included mild to severe and type included vascular and mixed dementias
Caregiver type (number)	NA

Abbreviations: NA=not available; PLWD=people living with dementia

Intervention Research Context

Exercise was the most commonly examined care intervention for PLWD. Most studies assessed the effects of exercise on functional and cognitive outcomes in adults with mild or moderate dementia. Enrolled adults typically had few mobility restrictions, could follow directions, were healthy enough to exercise moderately, and were dependent in at least one activity of daily living (ADL). Few studies included adults with severe dementia, and when included, their outcomes were not separately reported.

Overall, this literature is broad but dominated by pilot and small sample studies, reflecting that preliminary investigations for PLWD are more common. All eligible trials were published within the last two decades, and most (85%) within the past 10 years. The higher quality, analytic set studies were all published within the past decade, yet small sample and pilot studies are scattered throughout the publication dates.

Outcome domains included quality of life, function, neuropsychiatric symptoms, intervention adherence, adverse events, health or social service use, costs, and cognition. Daily functioning, measured with various ADL scales, was the most common outcome reported. The interpretability and clinical utility of reported outcomes varied considerably. Assessed outcomes were highly heterogeneous, even within domains and residential sites. Most outcomes were proxy-reported, except quality of life. Adverse events were proxy reported or chart-identified.

PLWD Outcomes

Evidence was insufficient to address any PLWD outcome. Table 5.2 summarizes the primary findings from 10 exercise intervention studies. We could not pool data for any outcome due to the wide variety of outcome measures, outcomes timing, and idiosyncratic reporting that

rendered some outcomes uninterpretable or noncomparable. Given these and other study limitations, the exercise intervention literature was insufficient for all intervention-outcome comparisons. An assessment of insufficient evidence does not mean that the intervention is determined to be of no value. Rather, it means that due to the uncertainty of the evidence we could not draw meaningful conclusions at this time.

Three exercise trials (n=864) assessed quality of life.^{104, 105, 107} Two trials (n=694) tested moderate to high intensity aerobic conditioning plus strength training exercises in community settings compared with usual care.^{104, 105} One trial compared high-intensity functional exercise with seated group activities in 18 nursing homes (n=170).¹⁰⁷ Intervention duration was 3¹⁰⁷ to 4 months,^{104, 105} and final outcomes were reported from 4 to 12 months. Studies used four quality of life measures, and one study reported dual quality of life measures.¹⁰⁴ Self-reported quality of life did not differ significantly between group aerobic with strength training exercises versus usual care in community-dwelling PLWD with mild to moderate dementia. Similarly, no meaningful between-group differences were seen for quality of life with high-intensity functional exercise compared with seated group activities in nursing home residents with mild to moderate dementia.¹⁰⁷

Eight randomized trials (n=1,518) assessed daily functioning using nine different outcome measures.^{101, 102, 104-107, 110, 111} Five of these were conducted in nursing facilities^{101, 106, 107, 110, 111} Six studies found no benefit for any outcome from exercise versus an inactive comparator, while two found some benefit on activities of daily living. The use of multiple daily functioning measures over varied followup time frames precluded pooling of data.

Balance was reported in two nursing home-based RCTs that examined high-intensity functional exercise versus seated group activities, and both reported balance outcomes with the Berg Balance Scale^{101, 107} in adults with mild to moderate dementia. One RCT (n=186) reported a post-intervention benefit of exercise on balance at 4 months, but improvement was not sustained at 7 months;¹⁰¹ the other trial found no difference at 3 or 6 months.¹⁰⁷ Other mobility-related outcomes were varied, incompletely reported, or shown only as percent change from baseline.^{102, 107, 108} Worse baseline motor function was associated with greater functional gains in one RCT.¹⁰⁸

Although neuropsychiatric symptoms were not required for study enrollment, half of analytic set RCTs reported at least one neuropsychiatric outcome.^{105, 107, 109, 111, 151} Group Tai Chi improved neuropsychiatric symptoms and depression in one RCT (n=80), while all other exercise interventions had no significant impact on the neuropsychiatric symptoms, depression, or anxiety.

Exercise-related adverse events, when reported, were most often musculoskeletal, such as muscle soreness. Serious adverse events were uncommon, variably reported, questionably related to the intervention, and similar between groups.^{101, 102, 105, 107, 108, 111, 151}

Other outcomes included exercise intervention adherence, attendance, or compliance, which ranged from 65 percent to 94 percent in studies that reported it.

Estimated costs of healthcare services in adults enrolled in a community-based group exercise intervention were statistically significantly lower than those of the usual care group at 24 months,¹⁰² but home exercise and usual care costs did not differ.

Table 5.2. Summary of findings for PLWD outcomes: exercise

Outcome Comparison	# Studies (# NH) n Analyzed Outcome Timing	Population	Finding	Strength of Evidence*
Quality of Life Group exercise vs usual care or seated group activity	3 RCTs. ^{104, 105, 107} (1 NH) n=864 3-12 months	Mild to Moderate	0 of 3 found benefit 3 of 3 found no difference	Insufficient
Daily function Group exercise vs usual care or home exercise	3 RCTs ^{102, 104, 105} (0 NH) n=904 4-12 months	1 Mild 1 Mild-Moderate 1 Mild-Severe	1 of 3 found benefit for group exercise vs UC, and in-home exercise vs UC; 2 of 3 found no difference	Insufficient
Daily function Group exercise vs seated group activity	3 RCTs ^{101, 106, 107} (3 NH) n=474 3-7 months	2 Mild-Moderate 1 Mild-Severe	1 of 3 found benefit for combined group exercise vs social activity, and walking vs social activity; 2 of 3 found no difference	Insufficient
Daily function Group Tai Chi vs usual care	1 RCT ¹¹¹ (1 NH) n=80 10 months	Mild	0 found benefit 1 found no difference	Insufficient
Daily function: Eating help needed. Hand exercise program vs usual care	1 RCT ¹¹⁰ (1 NH) n=60 6 months	Mild to Moderate	1 found benefit 0 found no difference	Insufficient
Daily function: Autonomous eating time. Hand exercise program vs usual care	1 RCT ¹¹⁰ (1 NH) n=60 6 months	Mild to Moderate	1 found benefit 0 found no difference	Insufficient
Balance HIFE (strength + balance) vs. seated group activity	2 RCT ^{101, 107} (2 NH) n=356 3-7 months	Mild to moderate	1 of 2 found early benefit 1 of 2 found no difference	Insufficient
Neuropsychiatric symptoms Group exercise vs usual care (2) or seated group activity (1)	3 RCTs. ^{104, 105, 107} (1 NH) n=864 3-12 months	Mild to Moderate	1 of 3 found benefit (non-NH) 2 of 3 found no difference (1 NH)	Insufficient
Neuropsychiatric symptoms: Group Tai Chi vs usual care	1 RCT ¹¹¹ (1 NH) n=80 10 months	Mild	1 found benefit	Insufficient
Depression Group exercise vs usual care (1) or seated group activity (2)	3 RCTs. ^{101, 104, 107} (2 NH) n=436 3-6 months	Mild to Moderate	0 of 3 found benefit 3 of 3 found no difference	Insufficient
Anxiety Group hand movement program vs. attention control	1 RCT ¹⁰⁷ (NH) n=66 3 months	Moderate	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: HIFE= high-intensity functional exercise (repetitive strength, balance, and mobility moves that mimic daily activities, such as rising from a chair); n=number; NH= nursing home; PLWD=person/people living with dementia; RCT=randomized controlled trial; UC=usual care

Caregiver Outcomes

Caregiver quality of life and burden did not differ over 6 and 12 months in the one RCT that measured it.¹⁵¹

Variation in Outcomes

Two of eight RCTs reported subgroup outcomes for adults with Alzheimer’s disease (AD) versus other dementias.^{101, 102} Adults with AD showed higher healthcare costs in all groups (group exercise in adult daycare versus home exercise versus usual care), but a similar cost pattern per intervention group to those with other dementias.¹⁰² Compared with adults with AD, adults with non-AD dementia showed greater balance improvements with high-intensity functional exercise versus seated group activity.¹⁰¹ Men were more likely to enroll in¹⁰¹ and adhere to¹⁵¹ exercise interventions than women. Session attendance was higher for exercise than for control group activities.

Music

Key Point

- Evidence was insufficient to draw conclusions about the effects of music interventions for PLWD or CG/P.

Intervention Description

Music interventions are intended to be calming or to provide pleasure or cognitive and sensory stimulation. They can be delivered as individual or group activities, often with music choices based on familiarity or preferences of PLWD. PLWD may be encouraged to move with the music (ranging from small hand movements to dancing) or listen passively (e.g., background music at mealtimes).

Eligible Studies

Table 5.3 summarizes the effect of music interventions for PLWD and caregivers. We identified 35 unique studies from 38 publications of music interventions for PLWD and caregivers. Five studies, all RCTs, were assessed as medium risk of bias and included in the analytic set.¹⁵²⁻¹⁵⁹ Four used group music as an intervention compared with usual care.¹⁵³⁻¹⁵⁶ One compared music-with-movement with music listening combined with social activity.¹⁵² The analytic set studies were conducted in either Hong Kong or Taiwan. We describe the 10 pilot, 13 small sample, and seven high risk of bias studies in the evidence map in Appendix D.^{153, 155, 160-184} Appendix D also includes an evidence table, summary of risk of bias assessments, and strength of evidence.

Table 5.3. Basic characteristics of literature set: music

Characteristic	Information
Total unique studies	35 studies
Non-U.S. studies	26 studies
Total Evidence map studies	10 pilot studies 13 small sample studies 7 high risk of bias studies
Analytic set studies	5 randomized controlled trials
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	404

Characteristic	Information
Dementia type/definition	All types of dementia with all stages of severity
Caregiver type (number)	1 randomized controlled trial with 59 family, 30 nurses

Abbreviation: PLWD=people living with dementia

Intervention Research Context

Using the National Institutes of Health (NIH) Stage Model, about 65 percent of the studies were categorized as preliminary pilot or small sample studies and the rest were categorized as explanatory, or Stage 3. The analytic set consisted of basic explanatory designs; the largest study enrolled 165 PLWD. We did not see the evolution of research on any particular intervention over time; instead, each study tested a different music intervention approach. Interventions in the evidence map were similar to those in the analytic set. Interventions were administered in groups^{115, 162, 168, 170, 172, 174, 176, 178, 180-182, 185-188} or individually^{160, 165-167, 175, 177, 183, 189} and included watching live music^{164, 169, 190} or listening to familiar songs from recordings,^{182, 183} playing with musical instruments,^{95, 173} and singing along or dancing to music.^{171, 174} Only one study used background music as an intervention.¹⁶¹ Non-U.S settings include: Australia,^{164, 165, 172} France,^{167, 182} Germany,¹⁸³ Hong Kong,¹⁸⁴ Iceland,¹⁷⁸ Italy,^{160, 174, 191, 192} Japan,¹⁷³ the Netherlands,¹⁸⁷ Norway,¹⁷⁵ Singapore,¹⁶⁸ South Korea,¹⁶² Taiwan,^{161, 176, 188} Turkey,¹⁷⁷ and the United Kingdom.¹⁶⁹

PLWD Outcomes

Evidence was insufficient to draw conclusions about music therapy for agitation, anxiety, depression, mood, and quality of life. Table 5.4 provides a summary of findings.

One study (n=84) found the intervention improved quality of life in PLWD.¹⁵⁵

Three studies examined agitation in PLWD; two compared (n=155) group music versus usual care over a period of 6 weeks,^{154, 156} and one (n=165) compared music-with-movement versus music listening combined with social activity for 6 weeks.¹⁵² Results were inconsistent. While one study reported a significant effect of group music on agitation,¹⁵⁴ the other two reported no statistical difference between intervention and control groups.

One study (n=55) found a group music intervention decreased anxiety compared with usual care.¹⁵⁶ One study reported that a group music intervention significantly decreased depression after 6 weeks.¹⁵⁸ A three-armed study (n=84) found music improved mood.¹⁵⁵

Table 5.4. Summary of findings for PLWD outcomes: music

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Quality of Life Group music vs. usual care	1 RCT ¹⁵⁵ (n=84) 10 weeks	Dementia (all types) with mild-moderate	1 found benefit 0 found no difference	Insufficient
Agitation Group music vs. Standard/usual care	3 RCT ^{152, 154, 156} (n=320) 6 weeks	All dementia types with mild to severe severity	1 of 3 found benefit 2 of 3 found no difference	Insufficient
Agitation Music with movement vs. music listening & social activity	1 RCT ¹⁵² (n=165) 6 weeks	Any type of dementia with moderate severity	0 found benefit 1 found no difference	Insufficient

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Anxiety Group music vs. usual care	1 RCT ¹⁵⁶ (n=55) 6 weeks	Any type of dementia with moderate severity	0 found benefit 1 found no difference	Insufficient
Depression Group music vs. usual care	1 RCT ¹⁵⁶ (n=55) 6 weeks	Any type of dementia with moderate severity	0 found benefit 1 found no difference	Insufficient
Mood Group music vs. usual care	1 RCT ¹⁵⁵ (n=84) 10 weeks	Dementia (all types) with mild-moderate	1 found benefit 0 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Evidence was insufficient for the effect of music on stress and caregiver burden for family caregivers. One study reported psychological well-being outcomes (stress and burden) for family caregivers and found benefit at 9 months.¹⁵⁵ Table 5.5 summarizes the primary findings. No significant reduction in stress was reported.

Table 5.5. Summary of findings for caregiver outcomes: music

Outcome Comparison	Studies/ Design (n=analyzed) Timing	Population	Findings	Strength of Evidence*
Stress & burden Group music vs. usual care	1 RCT ¹⁵⁵ (n=84) 9 months	Informal caregivers	One benefit	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: n=number; RCT=randomized controlled trial

Variation in Outcomes

A single study reported that both singing and listening were more effective in improving mood in PLWD with mild dementia and AD, while listening-only interventions were more beneficial for PLWD with moderate non-AD dementia (e.g., vascular and mixed dementias, frontotemporal dementia).¹⁵⁵

Reminiscence Therapy

Key Point

- Evidence was insufficient to draw conclusions about the effect of reminiscence therapy versus usual care for PLWD and CG/P.

Intervention Description

Reminiscence therapy for PLWD involves a discussion of past life events and experiences with the goal of improving memory and well-being. These discussions can be held in groups,

individually, or as a combination of both. Sessions may incorporate sentimental objects or photographs and include family members/caregivers.

Eligible Studies

We identified 25 unique studies from 27 publications examining reminiscence therapy for PLWD.^{47, 193-217} Table 5.6 summarizes the characteristics of the literature set. Four studies were included in the analytic set.¹⁹³⁻¹⁹⁸ All studies in the analytic set were RCTs comparing group reminiscence therapy to usual care for PLWD. One study enrolled PLWD-CG/P dyads.^{197, 198} Three studies examined the use of reminiscence therapy for PLWD with mild to moderate dementia.^{193, 194, 197, 198} One study examined the use of reminiscence therapy for PLWD with mild to severe dementia.^{195, 196} Appendix D provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Eleven studies were pilots or had fewer than 50 participants and were not included as part of the analytic study set.^{47, 199-207, 218} Ten additional studies were assessed as high risk of bias.^{208-217, 219} We present information on all pilot studies and high risk of bias in the evidence map in Appendix D.

Table 5.6. Basic characteristics of literature set: reminiscence therapy

Characteristic	Information
Total Studies	25 studies
Non-U.S. studies	22 studies
Evidence map studies	7 pilot studies 4 small sample size 10 high risk of bias
Analytic set studies	4 randomized controlled trials
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	904
Dementia type/definition	Predominantly mild to moderate dementia
Caregiver type (number)	Informal caregivers (n=904)

Abbreviations: n=number; PLWD=people living with dementia

Intervention Research Context

This literature set comprises mostly pilot studies and small RCTs. Included studies were published between 1999 and 2019. Only three studies were conducted in the United States, and none of these were in the analytic set.^{203, 205, 208} Eleven studies, including the four in the analytic set, examined group reminiscence therapy.^{193-199, 209, 210, 212, 213, 217, 218} Ten studies examined individual reminiscence therapy, including life-story approaches.^{47, 201, 205-207, 211, 214-216} Four studies examined reminiscence therapy using technology such as audio recordings and web-based videos.^{200, 202, 203, 208} We observed no clear trend in the publication of pilot studies or larger RCTs. Most of the studies that examined technology in reminiscence therapy were pilots, and we identified a mix of both pilots and small to large RCTs for group and individual reminiscence published across a two-decade span. The continuation of pilot studies might be partially attributed to study quality, as we found that a large proportion of nonpilot studies had high risk of bias. Potential detection and performance bias were significant concerns in nonpilot studies rated as high risk of bias.

PLWD Outcomes

Evidence was insufficient to draw conclusions about the effects of reminiscence therapy for PLWD. Four studies in the analytic study set reported outcomes for PLWD.¹⁹³⁻¹⁹⁸ Two studies

enrolled PLWD with mild to moderate AD,^{193, 194} one study enrolled PLWD with mild to moderate dementia of any type,^{197, 198} and one study enrolled PLWD with dementia of any type or severity.^{195, 196} Studies measured a range of outcomes for PLWD. We were able to quantitatively analyze and pool PLWD outcomes in four domains: activities of daily living, BPSD, depression, and quality of life. However, we assessed the evidence as insufficient due to the variety of measures used within domains, lack of minimally important differences for measures, and variation in analytic methods across studies.

Table 5.7 summarizes the primary outcomes for PLWD. Three studies with a total of 802 PLWD reported measures of activities of daily living.^{193, 194, 197, 198} At 3 months, groups did not differ significantly across all three studies. One study also reported outcomes at 6 months, finding no difference between groups.¹⁹⁴ Two studies with a total of 416 PLWD reported measures of BPSD.^{193, 194} Reminiscence therapy and usual care groups did not differ at 3 months.^{193, 194} Four studies with a total of 1,006 PLWD reported measures of depression.¹⁹³⁻¹⁹⁸ Individual study findings were mixed; however the pooled effect size showed no difference between groups at 2 to 3 months. Two studies with a total of 814 PLWD reported measures quality of life. Both found no difference between groups at 3 months.^{193, 197, 198}

Two studies reported additional PLWD outcomes. One (n=488) reported a measure of anxiety and a PLWD assessment of the PLWD-caregiver relationship.^{197, 198} The second (n=326) reported a measure of apathy and a measure of dependency.¹⁹³ Groups did not differ at 3 months for either measure.

Table 5.7. Summary of findings for PLWD outcomes: reminiscence therapy

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Activities of Daily Living Reminiscence vs usual care	3 RCTs ^{193, 194, 197, 198} (n=802) 3 months	Mild to moderate dementia	0 of 3 found benefit 3 of 3 found no difference	Insufficient
BPSD Reminiscence vs usual care	2 RCTs ^{193, 194} (n=416) 3 months	Mild to moderate AD	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Depression Reminiscence vs usual care	4 RCTs ¹⁹³⁻¹⁹⁸ (n=1,006) 2 to 3 months	All dementia types and severities	0 of 4 found benefit 4 of 4 found no difference	Insufficient
Quality of Life Reminiscence vs usual care	2 RCTs ^{193, 197, 198} (n=814) 3 months	Mild to moderate dementia	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Anxiety Reminiscence vs usual care	1 RCT ^{197, 198} (n=488) 3 months	Mild to moderate dementia	0 found benefit 1 found no difference	Insufficient
PLWD-Caregiver Reminiscence vs usual care Relationship	1 RCT ^{197, 198} (n=488) 3 months	Mild to moderate dementia	0 found benefit 1 found no difference	Insufficient
Apathy Reminiscence vs usual care	1 RCT ¹⁹³ (n=326) 3 months	Mild to moderate AD	0 found benefit 1 found no difference	Insufficient
Dependency Reminiscence vs usual care	1 RCT ¹⁹³ (n=326) 3 months	Mild to moderate AD	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: AD=Alzheimer’s disease; BPSD=behavioral and psychological symptoms of dementia; n=number; PLWD=people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Evidence was insufficient to draw conclusions about the effects of PLWD reminiscence therapy for CG/P. Three studies in the analytic set reported a range of outcomes for CG/P of PLWD.^{193, 194, 197, 198} Overlap between outcome measures was insufficient to assess the potential benefit of PLWD reminiscence therapy for CG/P. Table 5.8 summarizes the primary outcomes for CG/P.

Two studies enrolled PLWD with mild to moderate AD.^{193, 194} One study (n=326) measured caregiver burden, finding no difference between the reminiscence therapy group and the usual care group at 3 months.¹⁹³ Another study (n=90) measured CG/P distress, with no difference between groups at 3 months or 6 months.¹⁹⁴

One study (n=488) enrolled PLWD with mild to moderate dementia of any type. The study reported measures of mental health, quality of life, stress, anxiety, and depression in CG/Ps.^{197, 198} The study also measured CG/Ps' ratings of their relationship with the PLWD. At 3 months, groups did not differ in any of these measures.^{197, 198}

Table 5.8. Summary of findings for caregiver outcomes: reminiscence therapy

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Caregiver Burden Reminiscence vs usual care	1 RCT ¹⁹³ (n=326) 3 months	Informal caregivers	1 found no benefit.	Insufficient
Caregiver Distress Reminiscence vs usual care	1 RCT ¹⁹⁴ (n=90) 3 months, 6 months	Informal caregivers	1 found no benefit.	Insufficient
Mental Health Reminiscence vs usual care	1 RCT ^{197, 198} (n=488) 3 months	Informal caregivers	1 found no benefit.	Insufficient
Quality of Life Reminiscence vs usual care	1 RCT ^{197, 198} (n=488) 3 months	Informal caregivers	1 found no benefit.	Insufficient
Stress Reminiscence vs usual care	1 RCT ^{197, 198} (n=488) 3 months	Informal caregivers	1 found no benefit.	Insufficient
Anxiety Reminiscence vs usual care	1 RCT ^{197, 198} (n=488) 3 months	Informal caregivers	1 found no benefit.	Insufficient
Depression Reminiscence vs usual care	1 RCT ^{197, 198} (n=488) 3 months	Informal caregivers	1 found no benefit.	Insufficient
PLWD-Caregiver Relationship Reminiscence vs usual care	1 RCT ^{197, 198} (n=488) 3 months	Informal caregivers	1 found no benefit.	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: n=number; PLWD=people living with dementia; RCT=randomized controlled trial

Variation in Outcomes

The analytic study set did not report variation in outcomes by PLWD, caregiver, or intervention characteristics.

Cognitive Rehabilitation

Key Point

- Evidence was insufficient to draw conclusions about the effects of cognitive rehabilitation for PLWD and their CG/P.

Intervention Description

Cognitive rehabilitation, originally developed for people experiencing cognitive impairment from brain injury, was later adapted for use among PLWD with the goal of enabling them to maintain as much independent function as possible in areas they care most about.^{220, 221} We classified studies as cognitive rehabilitation when interventions focused on cognitive activities required for daily living (such as memory and executive function), and/or were aimed at recovery²²² or “reactivation of retained daily skills.”²²³ Therefore, studies in this section range from cognitive process activities in PLWD with mild dementia, to specific, task-based guidance to complete single or multiple ADLs in adults with more advanced dementia.

Eligible Studies

We identified 23 unique studies from 28 publications that examined various cognitive rehabilitation interventions for adults with dementia (Table 5.9).^{90, 193, 221, 222, 224-243} Three randomized trials were assessed as low or medium risk of bias and are included in the analytic set.^{193, 224, 244} The remaining 23 studies were pilot, small sample trials, or assessed as high risk of bias.^{90, 154, 221, 222, 225-230, 232-234, 239-245} We present information on all eligible studies in the evidence map in Appendix D.

Table 5.9. Basic characteristics of literature set: cognitive rehabilitation

Characteristics	Information
Total unique studies	23 studies
Non-U.S. studies	19 studies
Evidence map studies	2 pilot studies 15 small samples studies 3 high risk of bias studies
Analytic set studies	3 studies
Risk of bias of analytic set	Low/medium
Number of PLWD in analytic set	890
Dementia type/definition	Mild to moderate dementia
Caregiver type (number)	Informal caregivers (n=474)

Abbreviations: n=number; PLWD=people living with dementia

Intervention Research Context

The cognitive rehabilitation literature largely comprises unique interventions that have been tested only in small sample studies, thus the literature is largely preliminary. Trials included small group or individually tailored activities that engaged a variety of cognitive functions to improve the completion of daily activities in PLWD. Interventions were diverse, and details about both cognitive exercises and task guidance were often lacking, especially for individual rehabilitation of basic functional tasks.

Most included studies compared cognitive rehabilitation to other active cognitive approaches or usual care. The longest study lasted 2 years,¹⁹³ although most ranged from 3 to 6 months. Some studies reported both task-based outcomes and measures of positive affect surrounding task accomplishment, such as satisfaction, quality of life, or reduction in neuropsychiatric

symptoms. Studies were community,^{193, 220, 221, 224, 225, 228, 230, 232} hospital,²³⁹ or nursing home-based^{90, 229, 231, 235, 240 227, 233, 241} interventions, and two studies included a mix of nursing home and community residents.^{222, 228} Most studies (83%) took place outside of the United States.

PLWD Outcomes

Evidence was insufficient to draw conclusions about the effects of cognitive rehabilitation to improve quality of life, function, neuropsychiatric symptoms, mood, health service use, institutionalization, and harms for adults with dementia. We could not combine studies because the three trials tested very different interventions and assessed different outcomes at varying time points. The three analytic set trials (n=890) were conducted in noninstitutionalized adults with mild to moderate dementia.^{193, 224, 244} Table 5.10 summarizes the primary findings.

The largest, medium risk of bias RCT of 653 community-dwelling adults found no benefit of individual cognitive rehabilitation versus group usual care (n=311 for these study arms) on function, ADLs, behavioral issues, quality of life, depression, or caregiver burden over 24 months.¹⁹³ However, authors reported greater 24-month clinical benefits, including slower functional declines, improved survival, and marginal benefits on behavior, caregiver burden and resource use.¹⁹³

A medium risk of bias RCT of 475 community-dwelling adults found a statistically but not clinically significant benefit of individual cognitive rehabilitation compared with usual care on function assessed as self-rated goal attainment, at three and nine months.²⁴⁴ The mean Mini Mental State Exam scores of enrolled adults was 24 (range 18 to 30), which marginally met criteria for mild dementia. There was no benefit of cognitive rehabilitation on quality of life, depression, or anxiety, compared with usual care.²⁴⁴

One RCT found no benefit of errorless learning with feed-forward (continuous verbal) training over trial and error learning on ADLs in adults with AD and mixed dementia who were living at home.²²⁴

Table 5.10. Summary of findings for PLWD outcomes: cognitive rehabilitation

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Quality of Life Individual cognitive rehabilitation vs usual care	1 RCT ¹⁹³ (n=311) 24 months	Mild to moderate dementia	0 found benefit 1 found no difference	Insufficient
Quality of Life Individual cognitive rehabilitation vs usual care	1 RCT ²⁴⁴ (n=475) 9 months	Mild dementia	0 found benefit 1 found no difference	Insufficient
Daily function Individual cognitive rehabilitation vs usual care	1 RCT ¹⁹³ (n=311) 24 months	Mild to moderate dementia	0 found benefit 1 found no difference	Insufficient
Daily function Errorless Learning (feed-forward instructions) vs Trial & Error Learning	1 RCT ²²⁴ (n=161) 6 months	Mild to moderate dementia	0 found benefit 1 found no difference	Insufficient
Self-rated goal attainment Individual cognitive rehabilitation vs usual care	1 RCT ²⁴⁴ (n=475) 9 months	Mild dementia	1 found benefit 0 found no difference	Insufficient
Dependency Individual cognitive rehabilitation vs usual care	1 RCT ¹⁹³ (n=311) 24 months	Mild to moderate dementia	1 found benefit 0 found no difference	Insufficient
Neuropsychiatric symptoms Individual cognitive rehabilitation vs usual care	1 RCT ¹⁹³ (n=311) 24 months	Mild to moderate dementia	0 found benefit 1 found no difference	Insufficient
Neuropsychiatric symptoms Errorless Learning (feed-forward instructions) vs Trial & Error Learning	1 RCT ²²⁴ (n=161) 6 months	Mild to moderate dementia	0 found benefit 1 found no difference	Insufficient
Depression Individual cognitive rehabilitation vs usual care	1 RCT ²⁴⁴ (n=475) 9 months	Mild dementia	0 found benefit 1 found no difference	Insufficient
Anxiety Individual cognitive rehabilitation vs usual care	1 RCT ²⁴⁴ (n=475) 9 months	Mild dementia	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Cognitive rehabilitation offered no significant benefit over reminiscence on CG/P burden or apathy in the one trial that measured it.¹⁹³ Cognitive rehabilitation offered no significant benefit over usual care on CG/P stress, quality of life, or health-related quality of life over 9 months.²⁴⁴

Variation in Outcomes

No analytic set studies reported variations in outcomes by PLWD, caregiver, or intervention characteristics.

Cognitive Training

Key Point

- Evidence was insufficient to draw conclusions about the effect of cognitive training interventions for PLWD and their CG/P.

Intervention Description

Cognitive Training interventions involve repetitive or progressive drill-type exercises aimed at improving various domains of cognitive function (e.g., memory or executive functions).¹⁹³ The cognitive training interventions included in eligible studies incorporated some form of cognitive drills, whether paper-pencil tests or computer-based.

Eligible Studies

We identified 18 unique studies from 18 publications that examined cognitive training interventions among PLWD (Table 5.11).^{189, 191, 193, 246-260} Five studies were assessed as medium risk of bias and were included in the analytic set.^{189, 193, 246-249} Appendix D provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes. Ten studies had small sample sizes,^{191, 250-252, 254-257, 259-261} and two were assessed as high risk of bias.^{253, 258} We present information on all pilot studies and high risk of bias studies as part of the evidence map in Appendix D.

Table 5.11. Basic characteristics of literature set: cognitive training

Characteristics	Information
Total unique studies	18 studies
Non-U.S. studies	15 studies
Evidence map studies	3 pilot 9 small sample studies 1 high risk of bias
Analytic set studies	5 randomized controlled trials
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	748
Dementia type/definition	All types and severities
Caregiver type (number)	Informal (n=324), Formal (n=NR)

Abbreviations: n=number; NR=not reported; PLWD=people living with dementia

Intervention Research Context

The majority of eligible studies were small sample size trials involving fewer than 50 total participants each. Studies were published between 2001 and 2019, and were mostly conducted outside the United States. Intervention mode ranged from paper-and-pencil tasks to software or

web-based cognitive training. Cognitive training tasks targeted a range of functions that included memory, language, and executive function. Interventions were delivered in community-based (e.g., adult day care centers) or institutional settings (e.g., nursing homes, veterans homes). Three of the studies included in the analytic set took place at nursing homes, dementia special care units, or assisted-living residences,^{189, 246, 249} while two took place at adult day care or memory clinics.^{193, 247, 248}

PLWD Outcomes

Evidence was insufficient to draw conclusions about the effects of cognitive therapy for PLWD. Five studies reported outcomes for PLWD. Four studies compared cognitive training interventions to usual care.^{189, 193, 247-249} One study compared cognitive training with an attention control.²⁴⁶ Table 5.12 summarizes the primary outcomes for PLWD. Two studies enrolling a total of 471 PLWD reported measures of quality of life and found no difference between groups at 3 months.^{193, 247, 248} Two studies enrolling a total of 404 PLWD reported measures of depression found no difference between groups from 3 to 6 months.^{193, 246}

Two studies enrolling a total of 197 PLWD with excessive eating, or hyperphagia, reported outcome measures of hyperphagic behavior.^{189, 249} Both studies found that hyperphagia behaviors improved in the cognitive training group versus usual care at 6 weeks.^{189, 249} One study measured hyperphagia behavior up to 6 months post-intervention (30 weeks total), finding that this benefit was sustained in the cognitive training group.²⁴⁹ Both studies reported a variety of additional measures related to excessive eating. One study (n=97) reported food intake, finding a reduction in average food intake for the cognitive training group versus usual care at 6 weeks.¹⁸⁹ The other study (n=100) reported measures of short meal frequency, pica behavior, and change in eating habits from post-intervention (6 weeks) to 6-months post-intervention (30 weeks).²⁴⁹ Short-meal frequency improved in the cognitive training group versus the usual care group at all measured time points.²⁴⁹ Pica behavior improved in the cognitive training group immediately post-intervention as well as at 1 and 3 months post-intervention, but the benefit was not sustained 6 months post-intervention.²⁴⁹ Groups did not differ for change in eating habits.²⁴⁹

One study (n=324) also reported measures of BPSD, function/ADLs, apathy, and dependence, and found no difference between groups at 3 months.¹⁹³ Another study (n=80) reported a measure of anxiety and found no difference between groups at 3 and 6 months.²⁴⁶ Appendix D provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 5.12. Summary of findings for PLWD outcomes: cognitive training

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Quality of Life Cognitive training vs usual care	2 RCTs ^{193, 247, 248} (n=471) 3 months	Mild to moderate AD or other dementia	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Depression Cognitive training vs usual care	2 RCTs ^{193, 246} (n=404) 3 to 6 months	Early-stage to Moderate AD	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Hyperphagia Behavior Cognitive training vs usual care	2 RCTs ^{189, 249} (n=197) 6 to 30 weeks	Dementia with hyperphagia	2 of 2 found benefit 0 of 2 found no difference	Insufficient

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Food Intake Cognitive training vs usual care	1 RCT ¹⁸⁹ (n=97) 6 weeks	Dementia with hyperphagia	1 found benefit 0 found no difference	Insufficient
Short meal frequency Cognitive training vs usual care	1 RCT ²⁴⁹ (n=100) 6 to 30 weeks	Dementia with hyperphagia	1 found benefit at all time points 0 found no difference	Insufficient
Pica behavior Cognitive training vs usual care	1 RCT ²⁴⁹ (n=100) 6 to 30 weeks	Dementia with hyperphagia	1 found benefit favoring intervention post-intervention, but not sustained 6-months post-intervention 0 found no difference	Insufficient
Change in eating habits Cognitive training vs usual care	1 RCT ²⁴⁹ (n=100) 6 to 30 weeks	Dementia with hyperphagia	0 found benefit 0 found no difference	Insufficient
BPSD Cognitive training vs attention control	1 RCT (n=324) 3 months	Mild to moderate AD	0 found benefit 1 found no difference	Insufficient
Function/ADLs Cognitive training vs attention control	1 RCT ¹⁹³ (n=324) 3 months	Mild to moderate AD	0 found benefit 1 found no difference	Insufficient
Apathy Cognitive training vs attention control	1 RCT ¹⁹³ (n=324) 3 months	Mild to moderate AD	0 found benefit 1 found no difference	Insufficient
Dependence Cognitive training vs attention control	1 RCT ¹⁹³ (n=324) 3 months	Mild to moderate AD	0 found benefit 1 found no difference	Insufficient
Anxiety Cognitive training vs attention control	1 RCT ²⁴⁶ (n=80) 3 and 6 months	Early-stage AD	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: AD=Alzheimer's disease; ADL=activities of daily living; BPSD=behavioral and psychological symptoms of dementia; n=number; PLWD=people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Evidence was insufficient to draw conclusions about the effects of cognitive therapy for informal and formal caregivers of PLWD. Two studies reported outcomes for caregivers and compared cognitive training interventions compared with usual care.^{193, 249} Table 5.13 summarizes primary outcomes for caregivers. One study (n=324) reported a measure of caregiver distress for CG/P, and found no difference between groups at 3 months.¹⁹³ The second study reported a measure of formal caregiver distress specific to PLWD hyperphagic behavior, and found benefit for formal caregivers of the cognitive training group versus the usual care group post-intervention (6 weeks).²⁴⁹ However, the benefit was not sustained at 6 months.²⁴⁹

Table 5.13. Summary of findings for caregiver outcomes: cognitive training

Outcome Comparator	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Caregiver Distress Cognitive training vs usual care	1 RCT ¹⁹³ (n=324) 3 months	Informal caregivers	0 found benefit 1 found no difference	Insufficient
Caregiver Distress, PLWD hyperphagic behavior Cognitive training vs usual care	1 RCT ²⁴⁹ (n=NR) 6 to 30 weeks	Formal caregivers	1 found benefit favoring intervention group immediately post-intervention (6 weeks); however, difference was not sustained over the 6-month period post-intervention.	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: n=number; NR=not reported; PLWD=people living with dementia; RCT=randomized controlled trial

Variation in Outcomes

Studies in the analytic set did not examine variation in outcomes by PLWD, caregiver, or intervention characteristics.

Cognitive Stimulation Therapy

Key Point

- Evidence was insufficient to draw conclusions about the effect of cognitive stimulation therapy (CST) for PLWD and their CG/P.

Intervention Description

Cognitive stimulation is an intervention for people with mild to moderate dementia that offers engagement in various activities and discussions (usually in a group) aimed at general enhancement of cognitive and social functioning. Activities and material are presented in a game-like manner rather than as schooling.

Eligible Studies

Table 5.14 summarizes the characteristics of the literature set. We identified 12 unique studies from 15 publications.^{44, 212, 262-272} One of these was a pilot study, four had small samples, and two were assessed as high risk of bias; these seven studies were not part of the analytic set but information about them is provided as part of the evidence map in Appendix D. One study (n=201) was assessed as low risk of bias,^{262, 264} and the other (n=236) was assessed as medium risk of bias.²⁶³ The first compared CST versus usual care.^{262, 264} The second compared CST maintenance therapy versus withdrawal of CST (in both cases after an initial period of CST).²⁶³ Neither study reported caregiver outcomes. Appendix D provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 5.14. Basic characteristics of literature set: cognitive stimulation therapy

Characteristic	Information
Total studies	12 studies
Non-U.S. studies	11 studies

Characteristic	Information
Evidence map studies	1 pilot study 5 small sample studies 4 high risk of bias studies
Analytic set studies	2 randomized controlled trials
Risk of bias of analytic set	1 low, 1 medium
Number of PLWD in analytic set	437
Dementia type/definition	Mild to moderate severity
Caregiver type (number)	Not reported

Abbreviation: PLWD=people living with dementia

Intervention Research Context

The literature set remains mostly preliminary, using the NIH Stage Model, with pilot or small sample studies. One study reported a pilot component,²⁶⁴ but the other did not. One study in the literature set described itself as a pilot study to adapt CST to the Hong Kong context.²⁷³ Three studies had an active control,^{265, 268, 272} and another had both an active control and a usual care arm.²¹² The remaining studies used usual care or some variation thereof as the control.

PLWD Outcomes

Evidence was insufficient to determine the effect of CST on quality of life, function, neuropsychiatric symptoms, and mood. Table 5.15 summarizes the primary outcomes for PLWD. Only one study addressed the effect of CST versus usual care on quality of life, cognition, neuropsychiatric symptoms, anxiety, and depression (n = 201 subjects analyzed).⁴⁴ The other study took place after an initial session of CST, and examined the effect of CST maintenance therapy versus withdrawal of CST on quality of life, cognition, and neuropsychiatric symptoms.²⁶³

Table 5.15. Summary of findings for PLWD outcomes: cognitive stimulation therapy

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Quality of life CST vs. usual care	1 RCT ⁴⁴ (n=201) 8 weeks	PLWD	1 found benefit 0 found no difference	Insufficient
Function CST vs. usual care	1 RCT ⁴⁴ (n=201) 8 weeks	PLWD	0 found benefit 1 found no difference	Insufficient
Neuropsychiatric symptoms CST vs. usual care	1 RCT ⁴⁴ (n=201) 8 weeks	PLWD	0 found benefit 1 found no difference	Insufficient
Anxiety CST vs. usual care	1 RCT ⁴⁴ (n=201) 8 weeks	PLWD	0 found benefit 1 found no difference	Insufficient
Depression CST vs. usual care	1 RCT ⁴⁴ (n=201) 8 weeks	PLWD	0 found benefit 1 found no difference	Insufficient
Quality of life CST maintenance therapy after initial CST vs. withdrawal of CST after initial CST	1 RCT ²⁶³ (n=236) 12 weeks 24 weeks	PLWD	1 found benefit 0 found no difference	Insufficient

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Function CST maintenance therapy after initial CST vs. withdrawal of CST after initial CST	1 RCT ²⁶³ (n=236) 12 weeks 24 weeks	PLWD	1 found benefit 0 found no difference	Insufficient
Neuropsychiatric symptoms CST maintenance therapy after initial CST vs. withdrawal of CST after initial CST	1 RCT ²⁶³ (n=236) 12 weeks 24 weeks	PLWD	1 found benefit 0 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: CST=cognitive stimulation therapy; n=number; PLWD=people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

No studies reported caregiver outcomes.

Variation in Outcomes

One study reported that women appeared to benefit more than men for quality of life, neuropsychiatric symptoms, and function. The study did not report the magnitude of the differences, nor did it report p-values.²⁶⁴ Neither study reported variation in outcomes by caregiver or intervention characteristics.

Recreation Therapy

Key Point

- Evidence was insufficient to draw conclusions about recreation therapy interventions for PLWD or their CG/Ps.

Intervention Description

Recreation therapy interventions may involve art, games, music, or exercise. They are usually structured and facilitated, and may be conducted in groups or individually. Most of these interventions aim to involve participants actively in the chosen activity, rather than letting them participate passively (e.g. listen to music or read).

Eligible Studies

Table 5.16 summarizes the effect of recreation therapy for PLWD. We identified 14 unique studies from 14 publications that examined the effect of recreation therapy for PLWD. Three studies were assessed as low or medium risk of bias and included in the analytic set. All studies were randomized controlled trials (RCTs; n = 552),²⁷⁴⁻²⁷⁶ and all used usual care as the comparator. The studies were conducted in the United States, Australia, and Italy.. All pilot, small sample, and high-risk-or-bias studies are described as part of the evidence map in Appendix D.²⁷⁶⁻²⁸⁷ An evidence table, summary of risk of bias assessments, and strength of evidence is provided in Appendix D.

Table 5.16. Basic characteristics of literature set: recreational therapy

Characteristic	Information
Total unique studies	14 studies
Non-U.S. studies	11 studies
Total Evidence map studies	1 pilot study 5 small sample studies 5 high risk of bias studies
Analytic set studies	3 randomized controlled trial
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	552
Dementia type/definition	Type of dementia not reported, severity not reported
Caregiver type (number)	Not reported

Abbreviation: PLWD=people living with dementia

Intervention Research Context

The literature for recreational therapy interventions is still quite preliminary. Using the NIH Stage Model, large majority of studies were assessed as pilot and small sample studies. The largest study enrolled 389 PLWD. All but two studies were conducted outside of the United States. We did not see evolution of a particular intervention over time; instead, each study tested a different recreational therapy approach. Two studies involved activity sessions conducted one—on-one with an aide, a wheelchair tandem bicycle ride,²⁷⁷ and individualized activities such as singing or working with clay.²⁸⁸ The remainder involved larger group activities. The studies involved a diverse set of activities, and some studies involved more than one type of activity. One involved some form of group sporting activity.^{277, 279} Four involved different types of games: GO,²⁸¹ chess,²⁸⁶ dhakonan,²⁸⁷ or party games.²⁸⁵ One based activities on Chinese folk art, music, and games.²⁸⁰ Three involved art in some form, such as singing in a choir or visits to art museum plus art-making.^{278, 280, 283} Of the evidence map studies, only four used usual care as a comparator.^{277, 279-281}

PLWD Outcomes

Evidence was insufficient to draw conclusions about PLWD outcomes, including neuropsychological symptoms, depression, agitation, social function, physical function, quality of life, and antipsychotic medication use. Table 5.17 provides a summary of findings. For the most part, each outcome was only covered in one study.

Two studies (n=552) compared neuropsychiatric symptoms using the NPI, with one finding a benefit and the other showing no difference. The studies also reported markedly different baseline levels of neuropsychiatric symptoms.^{274, 275} Two other studies (n=517)^{274, 276} compared agitation; neither found a difference.

At both 13 and 26 weeks, the humor therapy study (n=389) reported quality of life, depression and agitation symptoms, and social function, and found no difference between groups at either time point.²⁷⁴

The indoor garden study (n=163) reported a small but statistically insignificant benefit for physical function. However, the study reported a statistically significant reduction in antipsychotic use at 24 weeks.²⁷⁵

Table 5.17. Summary of findings for PLWD outcomes: recreational therapy

Outcome Comparison	# Studies/Design (n=analyzed) Timing	Population	Findings	Strength of Evidence*
Neuropsychiatric symptoms Recreational therapy vs. usual care	2 RCTs ^{274, 275} (n=552) 24-26 weeks	Dementia type and severity not reported	1 of 2 found benefit 1 of 2 found no difference	Insufficient
Depression Recreational therapy vs. usual care	1 RCT ²⁷⁴ (n = 389) 13 and 26 weeks	Dementia type and severity not reported	1 found benefit 1 found no difference	Insufficient
Agitation Recreational therapy vs. usual care	2 RCTs ^{274, 276} (n = 517) Up to 13 weeks	Dementia type and severity not reported	0 found benefit 2 no difference	Insufficient
Function, social Recreational therapy vs. usual care	1 RCT ²⁷⁴ (n = 389) 13 and 26 weeks	Dementia type and severity not reported	0 found benefit 1 found no difference	Insufficient
Function, physical Recreational therapy vs. usual care	1 RCT ²⁷⁵ (n = 163) 24 weeks	Dementia type and severity not reported	1 found benefit 0 found no difference	Insufficient
Quality of life Recreational therapy vs. usual care	1 RCT ²⁷⁴ (n = 389) 13 and 26 weeks	Dementia type and severity not reported	0 found benefit 1 found no difference	Insufficient
Antipsychotic use Recreational therapy vs. usual care	1 RCT ²⁷⁵ (n = 163) 24 weeks	Dementia type and severity not reported	1 found benefit 0 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

No studies in the analytic set reported caregiver outcomes.

Variation in Outcomes

No studies in the analytic set reported any variation in outcomes.

Psychosocial Interventions for PLWD Well-Being

Key Point

- Studies of psychosocial interventions to improve PLWD well-being were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Psychosocial interventions include components such as group counseling, use of an assigned nursing home resident buddy, or approaches to improve effective communication in order to help psychosocial and behavioral aspects by improving communication, memory, and interpersonal skills among PLWD.

Eligible Studies

We identified seven unique studies from eight publications that examined various psychosocial session interventions among PLWD.²⁸⁹⁻²⁹⁶ (Table 5.18) All studies were either

pilots or small sample studies, and we provide information about them in the evidence map in Appendix D.

Table 5.18. Basic characteristics of literature set: psychosocial interventions for PLWD

Characteristics	Information
Total unique studies	7 studies
Non-U.S. studies	4 studies
Evidence map studies	1 pilot study 4 small sample studies 2 high risk of bias studies
Analytic set studies	0

Abbreviation: PLWD=person/people living with dementia

Intervention Research Context

The research is largely preliminary with small sample or pilot trials. Psychosocial session topics included psychodynamic interpersonal therapy for community-dwelling PLWD in the United Kingdom,²⁸⁹ using noncognitively impaired nursing home residents as “buddies” for PLWD in the United States,²⁹⁰ a self-management group intervention for PLWD in the early stages of dementia in the United Kingdom,^{291, 292} therapeutic conversation counseling sessions for community-dwelling PLWD in the United States,²⁹³ and peer support group sessions for community-dwelling PLWD in Hong Kong.²⁹⁴ Lastly, two studies examined the use of reality orientation therapy among PLWD in the United States in 1997²⁹⁵ and Italy in 2005.²⁹⁶

Creative Expression Therapy

Key Point

- Studies to assess the effect of creative expression therapy were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Creative expression programs are emerging interventions targeted at improving quality of care and life for PLWD in long-term care facilities. Creative expression interventions, such as storytelling or theatrical improvisation, do not require PLWD to access or use memories. They instead focus on creativity in the moment.

Eligible Studies

We identified five unique studies from five publications that examined creative expression interventions for PLWD and formal caregivers. (Table 5.19) Three studies were assessed as high risk of bias and two were pilot studies; all five were therefore excluded from the analytic set. We present information on all studies as part of the evidence map in Appendix D.

Table 5.19. Basic characteristics of literature set: creative expression therapy

Characteristics	Information
Total unique studies	5 studies
Non-U.S. studies	2 studies
Evidence map studies	2 pilot studies 3 high risk of bias studies
Analytic set studies	0

Intervention Research Context

Four interventions examined “TimeSlips (TS),” a group storytelling method,²⁹⁷⁻³⁰⁰ while one study investigated the “Veder Method” which integrates theater improvisation techniques with elements from other communication methods.³⁰¹ These interventions are relatively new, and the research is emerging (published between 2009 and 2019). Three studies were conducted in the United States,^{297, 298, 300} and the other two were conducted in China²⁹⁹ and the Netherlands.³⁰¹ All five studies focused mainly on the outcomes of behavior, mood, and quality of life for PLWD, although one study examined the caregiver outcomes of job satisfaction, attitude towards residents, and burnout.²⁹⁷

Multicomponent Interventions for PLWD Well-Being

Key Point

- Evidence is insufficient to draw conclusion about the effects of multicomponent care interventions in PLWD or their caregivers.

About the Interventions

The multicomponent studies in this set are largely unrelated. Earlier in this chapter we reported on another multicomponent intervention, CST. Since CST had a relatively distinct literature set, we chose to report it in a separate section.

Multicomponent interventions aimed at improving PLWD include a combination of components used simultaneously to support cognitive function, quality of life, and other health outcomes for PLWD. Intervention components may include cognitive and/or motor stimulation, physical activity using daily living activities, strategies adapted from cognitive training and neurorehabilitation, reminiscence therapy, and reality orientation, and exposure to bright light. Multicomponent interventions are intended to improve PLWD outcomes such as functional ability (to engage in activities of daily living) and depression. Multicomponent may or may not use special personnel such as nurses, aides, and therapists, and are delivered in varied settings including the PLWD’s home, day care centers, and nursing homes.

Eligible Studies

Table 5.20 summarizes the characteristics of the literature set. We identified 24 unique studies from 39 publications. Three studies were assessed as low to medium risk of bias and were included in the analytic set. Multicomponent interventions targeting quality of life, function, and/or non-BPSD symptoms in PLWD included three unique studies across four publications. One study examined an intervention in Germany including motor stimulation, activities of daily living, and cognitive stimulation components, delivered by two therapists, one aide, and nurses when necessary.^{302, 303} One study examined walking, light therapy, and combination exercise, light therapy, and education in the United States.³⁰⁴ One Japanese study examined reminiscence and reality orientation therapy care methods, in addition to a routine day-care service.³⁰⁵ Excluded from the analytic set are 21 pilot, small sample, and high risk of bias studies and are described as part of the evidence map in Appendix D.^{44, 94, 114, 116, 303, 306-324} Appendix D provides an evidence table, summary of risk of bias assessments, and strength of evidence for the analytic set.

Table 5.20. Basic characteristics of literature set: multicomponent for PLWD well-being

Characteristics	Information
Total unique studies	24 studies
Non-U.S. studies	14 studies
Evidence map studies	3 pilot studies 7 small sample studies 11 high risk of bias
Analytic set studies	3 randomized controlled trials
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	333
Dementia type/definition	Varied across studies
Caregiver type (number)	NA

Abbreviations: NA=not available; PLWD=people living with dementia

Intervention Research Context

Evidence map studies published between 2004 and 2020 tended to be largely preliminary or early Stage 3 of the NIH Stage Model. Studies in the analytic set were likewise explanatory in nature and as a whole, they did not exhibit much progression along the NIH Stage Model. Study duration ranged from 6 weeks to 18 months, targeted highly varied PLWD populations, and focused primarily on PLWD with mild and moderate dementia residing in various community-based residences and nursing homes. Two of the trials targeting PLWD with mild to moderate dementia focused on an intervention for motor stimulation, activities of daily living, and cognitive stimulation.^{302, 325} These trials represented the most pragmatic approach to multicomponent interventions for PLWD health outcomes.

PLWD Outcomes

Evidence was insufficient for drawing conclusions about the effects of multicomponent interventions for PLWD well-being. The findings are summarized in Table 5.21. One study reported small benefit for PLWD quality of life (social behavior), cognitive function, and instrumental activity of daily living for PLWD with mild to moderate dementia living in nursing homes.^{302, 303} One study reported a small reduction of sleep disturbances.³⁰⁴ and the other reported small improvements in overall cognitive and social function.³⁰⁵

Table 5.21 Summary of findings for PLWD outcomes: multicomponent for PLWD well-being

Outcome Comparison	# Studies/Design (n=analyzed) Timing	Population	Findings	Strength of Evidence*
Instrumental Activities of Daily Living Multicomponent: motor stimulation, activities of daily living, and cognitive stimulation components contact control	1 RCT 302, 303 n=98 6 months	ADRD	1 found benefit 0 found no difference	Insufficient
Sleep Walking, light, combination treatment versus contact control	1 RCT 304 n=66 6 months	People with AD and their in-home caregivers.	0 found benefit 1 found no difference.	Insufficient
Cognitive and Social Function Reminiscence and reality orientation care methods, routine day-care service versus usual care	1 RCT 305 n=60 10 weeks	Community-dwelling older PLWD.	1 found benefit 0 found no difference	Insufficient

*Insufficient ratings due to few studies and imprecision in the findings.

Abbreviations: AD=Alzheimer’s disease; ADRD=Alzheimer’s disease–related dementias; n=number; PLWD=people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Where studies may have included caregiver outcomes, these lacked sufficient or clear reporting for caregiver outcomes.

Variation in Outcomes

Studies did not report variation in PLWD outcomes by PLWD characteristics, caregiver characteristics, or setting.

Assistive Technology

We classified as assistive technology interventions include technology that controls house functions such as lights and/or systems that alert caregivers when a PLWD is at risk. Additionally, these interventions evaluated caregiver outcomes such as job satisfaction, relieving worry, and sleep quality.

Key Point

- Studies of assistive technology interventions were described in the evidence map but not considered for analysis due to limitations in study designs.

Eligible Studies

We identified four unique studies from four publications that examined the use of assistive technology for PLWD and informal/formal caregivers. (Table 5.22) All included studies were pilot studies. Information on all studies is provided as part of the evidence map in Appendix D.^{320, 326-329}

Table 5.22. Basic characteristics of literature set: assistive home technology

Characteristics	Information
Total unique studies	4 studies
Non-U.S. studies	2 studies
Evidence map studies	4 pilot studies
Analytic set studies	0

Intervention Research Context

Assistive technology interventions identified are relatively novel and thus these studies were preliminary in design, spanning nine years from 2010 to 2019. Intervention topics include the effectiveness of implementation of assistive technology in group homes in Amsterdam on the quality of life of PLWD and on the job satisfaction of caregivers.³²⁶ The effectiveness of home-based technologies coupled with teleassistance service was also assessed in order to prevent indoor falls of individuals with dementia in France.³²⁸ Additionally, a home monitoring system designed to track the movements of PLWD was assessed to see if it would relieve worry and improve sleep in caregivers in the United States.³³⁰ A second U.S. study examined wearable technology to alert caregivers of activities, generating reports of activity patterns over time.³²⁹

Electrostimulation

Key Point

- Studies of electrostimulation were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Electrostimulation interventions encompass a variety of techniques using either electric current or electromagnetic pulses of different intensity in various of parts of brain to improve cognitive and behavioral functioning for PLWD.

Eligible Studies

We identified 14 unique studies from 15 publications that examined electrostimulation among PLWD. (Table 5.23) All studies were either pilots or had small sample sizes.^{50, 320, 326-328, 331-343} We present information on all studies as part of the evidence map in Appendix D.

Table 5.23. Basic characteristics of literature set: electrostimulation

Characteristics	Information
Total unique studies	14 studies
Non-U.S. studies	12 studies
Evidence map studies	5 pilot 9 small sample studies
Analytic set studies	0

Intervention Research Context

The research is early in development and based on small sample sizes, thus preliminary. Nursing home or other long-term care facilities were the most common settings. Non-U.S. locations included Brazil, Canada, China, Egypt, Japan, Italy, The Netherlands, and Spain. Electrostimulation topics are highly technical neuroscience research. Please refer to the evidence map for more details.³³¹⁻³⁴⁵

Other Interventions for PLWD Well-Being

Key Point

- No studies of other care interventions for PLWD well-being advanced from the evidence map to further analysis.

Intervention Description

This section encompasses a wide range of interventions. These included passive exercise (movement therapy or finger movement performed by a formal caregiver on PLWD), suicide prevention programs to pain assessments to verbal cues, and handfeeding techniques to help improve various (physical, mental, cognitive and behavioral) aspects of functioning.

Eligible Studies

We identified seven unique studies from eight publications that examined the various interventions among PLWD.^{133, 302, 325, 346-352} (Table 5.24) All the studies were either pilot or small sample studies. We present information on all studies in the evidence map in Appendix D.

Table 5.24. Basic characteristics of literature set: other interventions for BPSD

Characteristics	Information
Total unique studies	7 studies
Non-U.S. studies	4 studies
Evidence map studies	4 pilot 1 small sample studies 1 high risk of bias
Analytic set studies	0

Abbreviation: BPSD=behavioral and psychological symptoms of dementia

Intervention Research Context

This research is preliminary. Publications included a suicide prevention program tested in South Korea for older adults with early-stage dementia,³⁵¹ pain assessment in Norwegian nursing homes,³⁵² passive finger movement exercise to improve grip strength in Chinese PLWD in residential facilities,¹³³ various handfeeding techniques in PLWD with advanced dementia in the United States,³⁴⁶ and passive movement therapy for severe paratonia, or muscle stiffness, in late-stage dementia.³⁴⁹ One study from 1997 examined directed verbal prompts and positive reinforcements for eating independence in PLWD in U.S. nursing homes.³⁴⁷

Conclusion

We found 223 unique studies that we grouped into 13 categories of care interventions aimed at improving PLWD quality of life and well-being. The large majority of studies, 85 percent, were conducted in non-U.S. settings. Eighty-five percent of the studies were pilots or small sample studies, Stage 0 to 2 according to the NIH Stage Model, or assessed as high risk of bias. Of those interventions with an analytic set of low to moderate risk of bias studies, we found the uncertainty of the evidence was too high to draw conclusions. Challenges with clear definitions of classes of interventions, and variability both in outcomes and how they were measured, hampered our ability to combine studies. Loss of PLWD participants after study enrollment was a frequent cause of increased risk of bias. However, evidence that is insufficient does not mean that none of the individual interventions described are potentially useful for individual PLWD or their caregivers. It simply means the uncertainty of the evidence is too high for us to draw conclusions, at present.

Chapter 6. Care Interventions for Informal Caregivers

This chapter includes care interventions for informal caregivers, with the goal of improving their quality of life and well-being. While the large majority of studies use the term “informal caregiver,” the term care partner is also occasionally present; we use the term caregiver/partner, or CG/P. The interventions include psychosocial interventions, social support, lifestyle interventions, respite care, multicomponent interventions, and other interventions that did not fall into these previous categories. We also present in this chapter interventions that target dyads of people living with dementia (PLWD) and their CG/P.

For each intervention, we present Key Points followed by results in three general sections: Intervention Description, Eligible Studies, and Intervention Research Context. For interventions with no studies assessed as low to medium risk of bias, we present the studies from the evidence map with a brief discussion of what has been examined and research context. For interventions for which low- to medium-risk-of-bias studies were available for an analytic set, we present Outcomes sections by PLWD, by caregiver, and by variation in outcomes when available.

We present a qualitative synthesis of the findings, because differences in outcome measures and intervention complexity prohibited the pooling of outcomes for a statistical analysis, such as a meta-analysis. We present summary findings as brief statements of how many studies reported statistically significant benefits or no difference between the intervention and comparator. Effect sizes and the data to calculate them were rarely reported in this literature. In studies where an effect size was reported, we present the study’s originally reported effect size where it may be helpful for interpreting the study’s findings. We present detailed information on eligible studies in Appendix E.

Psychosocial Interventions To Support Informal Caregiver Well-Being

Key Point

- For both PLWD and CG/Ps, evidence is insufficient to draw conclusions about the effects of psychosocial interventions targeting CG/P.
- For CG/P, we found no studies that assessed harms for psychosocial interventions targeting CG/P well-being.

Intervention Descriptions

As an umbrella term, we use “psychosocial interventions” to capture a wide range of psychoeducational and psychotherapeutic interventions that seek to improve the well-being of caregivers. Newly formulated conceptual definitions for psychoeducational and psychotherapeutic caregiver interventions do exist. However, overlap in intervention components made it impossible for us to use these newer definitions to categorize our eligible interventions.³⁵³ Figure 6.1 provides a visual representation of the overlap of intervention components drawn from the analytic set (for which results are provided below). No clear pattern distinguished interventions as either psychoeducational or psychotherapeutic.

Figure 6.1. Collaboration network visualization of CG/P psychosocial intervention components




Figure 6.1 Legend for network, starting at the top and moving clockwise	
Engage PLWD in Daily/Social Activities	
Relationship Building	
Increasing Rewarding Activities	
Counseling	
Network Analysis	
Activity Analysis	
CBT	
Control of Activation	
Cognitive Reappraisal/ restructuring	
Dementia Education	
Training - Managing BPSD	
Training – Caregiving Skill	
Support	
Reminiscence	
Enhance Self-efficacy	
Stress Management	
Self-Care	
Problem Solving	
Communication Skills	
Environmental Modification	
Community Resources	
Goal Setting	
Coping Skills	
Positive Emotion Regulation	
Crisis Management	

Abbreviations: BPSD=behavioral and psychological symptoms of dementia; CBT=Cognitive Behavioral Therapy; CG/P=caregiver/care partner; PLWD=person/people living with dementia

We also could not categorize interventions based on duration, dose, delivery elements, or the interventionists. In almost all of the studies, interventionists were highly trained healthcare professionals or graduate level trainees, such as psychologists, social workers, registered nurses, occupational therapists, and licensed therapists. Interventions lasted anywhere from 6 weeks to 10 months, and participants spent anywhere from 5.5 to 32 hours receiving education, skills training, or counseling. Interventions were delivered to groups, individuals, and caregiver/ PLWD dyads. Most sessions occurred in person, a few by phone, and one online. We summarize intervention components, interventionists, duration, dose, and delivery format in Appendix E.

Eligible Studies

Table 6.1 summarizes the characteristics of the literature set on psychosocial interventions for CG/P. We identified 122 unique studies from 161 publications. Excluded from the analytic set are 46 pilot and small sample studies and 47 studies assessed as high risk of bias. The analytic dataset includes 29 studies with medium to low risk of bias, and represents 5054 caregiver/ PLWD dyads.³⁵⁴⁻³⁸¹ The majority of PLWD had mild to moderate dementia; however, studies infrequently reported dementia severity. We provide information on all pilot studies and high risk of bias studies as part of the evidence map in Appendix E.³⁸²⁻⁴⁷⁶ Appendix E also provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 6.1. Basic characteristics of literature set: psychosocial interventions for CG/P

Characteristic	Information
Total unique studies	122 studies
Non-U.S. studies	74 studies (17 analytic set non-U.S.)

Characteristic	Information
Evidence map studies	28 pilot studies 18 small sample studies 47 high risk of bias studies
Analytic set studies	27 randomized controlled trials 1 cluster randomized controlled trial
Risk of bias of analytic set	29 medium/low
Number of PLWD in analytic set	5054
Dementia type	Predominantly mild to moderate dementia, commonly not defined
Caregiver type (number)	Informal caregivers (4952)

Abbreviations: CG/P=caregiver/care partner; PLWD=people living with dementia

Intervention Research Context

Using the National Institutes of Health (NIH) Stage Model, about 40 percent of the studies were categorized as pilot studies and the rest were categorized as explanatory, or Stage 3. Most of the randomized controlled trials had high risk of bias. All of the 28 medium to low risk of bias studies in the analytic dataset were categorized as explanatory studies using the NIH Stage Model. None were categorized as a pragmatic trial, or Stage 4. All but two of the studies occurred in high-resource countries including the United States, the United Kingdom, France, Hong Kong, Finland, Spain, Germany, Denmark, Norway, and the Netherlands. These interventions tended to be costly and resource intensive. Two smaller studies occurred in Peru and Iran, low-resource countries. The Peruvian intervention was a modification of the 10/66 Helping Carer’s to Care program from India that was designed for diverse low to middle income countries with limited health and social care resources. The Iranian intervention promoted resilience training. Few studies investigated diverse caregivers (three investigated Latino caregivers, and one investigated African American caregivers). About half of the studies used attention control groups of generic psychoeducational programs, reminiscence, or support. A single comparative effectiveness study compared two cognitive behavioral therapy programs, one conducted by phone and one in person.

Modifications of the REACH Palo Alto Coping with Caregiving program were investigated in several countries including the United States,^{371, 372} the United Kingdom,³⁶³ Spain,^{375 364} and Hong Kong.³⁵⁴ Over time, the Coping with Caregiving group program³⁷¹ was adapted for individual sessions occurring in the caregiver’s home or by phone and for shorter sessions.

A research program of occupational therapy interventions were also investigated in the United States,^{359, 373} Germany,²²⁴ and the Netherlands.³⁷⁴ The occupational therapy interventions added components of environmental adaptations and compensatory actions for the PLWD’s daily activities to common psychoeducational activities.

PLWD Outcomes

Only 16 of the 29 medium to low risk of bias studies reported outcomes related to PLWD. We synthesized results qualitatively because differences in outcome reporting, outcome measures, time to outcomes, and comparison groups prohibited quantitative pooling. The most commonly investigated outcomes were function, neuropsychiatric symptoms, and quality of life, and the least commonly investigated were depression, healthcare use, social support, harms, and institutionalization. Table 6.2 summarizes the number of studies investigating each outcome and the number of studies that found a statistically significant benefit or no difference in outcomes between PLWD and controls.

Evidence was insufficient to draw conclusions about the effects of psychosocial interventions for any PLWD outcome. The inability to combine outcomes from multiple studies made it

impossible to show precision in any outcome. Several outcomes were only investigated by a single study, making the consistency of findings unknown. Many PLWD outcomes were deemed indirect because the researchers relied on proxy respondents. Although a few comparisons had reasonably large numbers of participants (500 to 900+) and several studies contributing, the studies often failed to report effect sizes or the data needed to calculate them. The uncertainty regarding a true “no difference between groups” finding versus an “inability to show a difference” finding remained too high.

Depression was investigated by four studies using usual care and one study using attention control groups, with only one study reporting benefit for the psychosocial intervention in the short term and one study reporting benefit in the long term. One RCT reported improvements in PLWD depression at 6 weeks for the psychosocial treatment group.³⁷⁴ Three RCTs reported no group differences in depression scores over 12 to 52 weeks between psychosocial groups and usual care,^{224, 355, 367} with one study reporting a Cohen’s d effect size of 0.³⁶⁷ One RCT reported improved depression scores over 1 year for their psychosocial treatment group versus attention control.³⁶⁹

Function was investigated by three studies using usual care and five using attention control groups, with mixed benefit for the psychosocial interventions. Two^{373, 374} found short-term improvements in function for the psychosocial treatment group versus usual care, but one did not.³⁵⁵ Three studies found no differences in long-term function between psychosocial intervention groups and usual care comparisons.^{224, 356, 367} Outcomes were mixed for psychosocial groups versus attention control groups; one study found improved functioning for the treatment group,³⁵⁹ but a second found no difference.³⁶⁹

Neuropsychiatric symptoms were investigated by three studies using usual care and five studies with attention control groups. For studies using usual care comparisons, only one found improved neuropsychiatric symptoms for the psychosocial group,³⁷⁷ while three found no difference over 3 to 6 months.^{355, 361, 373} For studies using an attention control group, two found short-term benefits in neuropsychiatric symptoms for their psychosocial treatment groups,^{359, 360} but three found no long-term differences in neuropsychiatric symptoms.^{356, 369, 370}

Quality of life was investigated by six studies using usual care and two using attention control groups. For studies using usual care comparisons, only one found improved PLWD quality of life at 6 weeks,³⁷⁷ while five found no difference in quality of life over 6 to 24 months.^{224, 356, 362, 363, 367} For studies using an attention control group, one study found benefits in quality of life for their psychosocial treatment group,³⁶⁹ but another³⁵⁹ found no difference.

Social support was investigated by one study using usual care and another using an attention control. Both found no difference in PLWD social support over 6 months.^{367, 368}

Unmet needs was investigated by one study using education and information support over 6 months. No difference was found between groups.⁴⁷⁷

Healthcare use was investigated by four studies using usual care and one using an attention control group. One English study found that participants in the psychosocial group spent 436 fewer pounds sterling over 2 years ($p=0.035$) on healthcare services than their usual care group.³⁶² However, four other studies found no difference in healthcare use over 4 to 24 months.^{363, 375, 378}

Harms were investigated by one study using a usual care control group. It found zero adverse events for participants in both the psychosocial treatment and control groups.³⁷⁸

Nursing home placement was investigated by two studies using usual care and two using attention control groups. One of the two studies using a usual care control group found a benefit

for the psychosocial intervention,³⁷⁶ and the other found no difference.³⁵⁵ The studies using an attention control group found no difference between groups.^{368, 370}

Table 6.2. Summary of findings for PLWD outcomes: psychosocial interventions for CG/P

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Depression psychosocial vs Usual care	4 RCTs ^{224, 355, 367, 374} (n=592) 6-52 weeks	CG/P	1 of 4 found benefit 3 of 4 found no difference	Insufficient
Depression Psychosocial vs Attention control	1 RCT ³⁶⁹ n=330 52 weeks	CG/P	1 found benefit 0 found no difference	Insufficient
Function Psychosocial vs Usual care	3 RCTs ^{355, 373, 374} (n=545) 6-12 weeks	CG/P	2 of 3 found benefit 1 of 3 found no difference	Insufficient
Function Psychosocial vs Usual care	3 RCTs ^{224, 356, 367} (n=540) 6-12 months	CG/P	0 of 3 found benefit 3 of 3 found no difference	Insufficient
Function Psychosocial vs Attention control	2 RCTs ^{359, 369} (n=490) 4-12 months	CG/P	1 of 2 found benefit 1 of 2 found no difference	Insufficient
Neuropsychiatric symptoms Psychosocial vs Usual care	4 RCTs ^{355, 361, 373, 457} (n=740) 3-6 months	CG/P	1 of 4 found benefit 3 of 4 found no difference	Insufficient
Neuropsychiatric symptoms Psychosocial vs Attention control	2 RCTs ^{359, 360} (n=227) 4-6 months	CG/P	2 of 2 found benefit 0 of 2 found no difference	Insufficient
Neuropsychiatric symptoms Psychosocial vs Attention control	3 RCTs ^{356, 369, 370} (n=916) 12-18 months	CG/P	0 of 3 found benefit 3 of 3 found no difference	Insufficient
Quality of life Psychosocial vs Usual care	1 RCT ⁴⁵⁷ (n=135) 6 weeks	CG/P	1 found benefit 0 found no difference	Insufficient
Quality of life Psychosocial vs Usual care	5 RCTs ^{224, 356, 362, 363, 367} (n=936) 6-24 months	CG/P	0 of 5 found benefit 5 of 5 found no difference	Insufficient
Quality of life Psychosocial vs Attention control	2 RCTs ^{359, 369} (n=490) 4-12 months	CG/P	1 of 2 found benefit 1 of 2 found no difference	Insufficient
Social support Psychosocial vs Usual care	1 RCT ³⁶⁷ (n=108) 6 months	CG/P	0 found benefit 1 found no difference	Insufficient
Social support Psychosocial vs Attention control	1 RCT ³⁶⁸ (n=250) 6 months	CG/P	0 found benefit 1 found no difference	Insufficient
Healthcare usage Psychosocial vs Usual care	4 RCTs ^{224, 362, 363, 375} (n=652) 4-24 months	CG/P	1 of 4 found benefit 3 of 4 found no difference	Insufficient
Healthcare usage Psychosocial vs Attention control	1 RCT ³⁶⁸ (n=250) 6 months	CG/P	0 found benefit 1 found no difference	Insufficient

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Harms – adverse events Psychosocial vs Usual care	1 RCT ²²⁴ (n=141) 52 weeks	CG/P	0 found benefit 1 found no difference	Insufficient
Institutionalization Psychosocial vs Usual care	2 RCTs ^{355, 376} (n=288) 6-12 months	CG/P	1 of 2 found benefit 1 of 2 found no difference	Insufficient
Institutionalization Psychosocial vs Attention control	2 RCT ^{368, 370} (n=545) 6-18 months	CG/P	0 of 2 found benefit 2 of 2 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: CG/P=caregiver/care partner; n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

All 29 medium to low risk of bias studies reported caregiver outcomes related to the caregivers’ psychological health, quality of life, social support, healthcare use, and their perception of caregiving burden, bother or distress, confidence, knowledge, and skills. We synthesized results qualitatively because differences in outcome measures, time to outcomes, and comparison groups prohibited quantitative pooling. The most commonly investigated outcomes were depression, quality of life, caregiver burden, and caregiver bother or distress, and the least common were coping, caregiving knowledge, social support, healthcare use, and relationship with PLWD. No studies looked for caregiver harms associated with the psychosocial intervention. Table 6.3 summarizes the number of studies investigating each outcome and the number of studies that found a statistically significant benefit or no difference in outcomes for caregivers enrolled in the psychosocial intervention group and those enrolled in the comparison group.

Evidence was insufficient for all caregiver outcomes. The insufficient ratings were attributed to several limitations across the bodies of evidence. The inability to quantitatively combine outcomes from multiple studies made it impossible to show precision for any outcome. Several outcomes were only investigated by a single study, making the consistency of findings unknown. Even though a few comparisons had reasonably large numbers of PLWD (500 to 900+) combined from several studies, those studies rarely reported effect sizes or the data to calculate them. The uncertainty regarding a true no difference between groups versus an inability to show a difference remained too high.

Anxiety was investigated by two studies using usual care and two using attention control groups, and only one study reported benefit for the psychosocial intervention. For studies using usual care comparisons, just one reported short-term improvements at 6 weeks for the psychosocial group with a Cohen’s d effect size of -0.32 (-.63, -.02).³⁶⁶ However, another study found no group difference in anxiety over the long term.³⁶³ Both of the two studies using attention control comparisons found no group differences in anxiety over 6 to 12 months.^{356, 360}

Depression was investigated by 10 studies using usual care and 10 others using attention control groups; most found no statistically significant difference between the psychosocial and comparison groups. For studies using usual care comparisons, two found short-term benefits for the psychosocial groups,^{366, 377} while three found no short-term group differences in depression.^{358, 366, 367} Two studies found long-term benefits for the psychosocial group,^{355, 379}

while three found no group differences in depression over 1 to 2 years.^{224, 356, 363} For studies using attention control groups, six found short-term benefits for the psychosocial groups,^{354, 357, 360, 368, 372, 380} with one reporting a moderate Cohen's d effect size of -0.66 (-1.04, -0.74).³⁵⁴ However, one study reported no short-term group differences in depression.³⁵⁹ We calculated the standardized mean difference (SMD) and 95 percent confidence intervals for each of these studies and found a nonsignificant overall SMD of 0.207 (-0.117, 0.531) (p=0.15). See forest plots in Appendix E. Depression in the long-term studies was investigated by three studies that all reported no group differences in depression at 12 or 18 months.^{356, 369, 370}

Stress was investigated by three studies using usual care and one using attention control groups, with only one study reporting benefit for the psychosocial intervention. Three studies using usual care comparisons found no group differences in caregiver stress in the short or long term,^{355, 366, 367} with one study reporting a nonsignificant Cohen's d effect size of -0.20 (-0.50, 0.11).³⁶⁶ A single study using an attention control comparison found improvements in stress for the psychosocial group.³⁷²

Caregiving burden was investigated by six studies using usual care and four using attention control groups with no apparent pattern of benefit for the psychosocial interventions. For studies using usual care comparisons, three found no short-term group differences,^{364, 366, 377} with one reporting a nonsignificant Cohen's d effect size of -0.16 (-0.46, 0.14).³⁶⁶ Results for long-term caregiver burden were mixed, with two studies finding benefit for the psychosocial intervention^{361, 376} and three finding no group differences.^{224, 358, 375} For studies using attention control groups, results were mixed, with three studies finding benefit for the psychosocial group,^{354, 357, 380} and two finding no group differences in caregiving burden.^{359, 368}

Caregiving bother/distress was investigated by four studies using usual care and five using attention control groups, with most studies finding no group differences. For studies using usual care comparisons, only one study found benefit for the psychosocial group,³⁷⁷ while three studies found no group differences in caregiver bother.^{361, 366, 373} For studies using attention control comparisons, results were mixed, with three studies finding benefit for the psychosocial interventions,^{359, 360, 372} but two finding no group differences for caregiving burden.^{356, 368}

Caregiving knowledge was investigated by a single study using a usual care comparison. It found that the psychosocial group had greater improvements in their caregiving knowledge than the usual care comparison group at 6 months.³⁵⁸

Caregiving confidence was investigated by three studies using usual care and three using attention control groups, and more studies found no group differences than benefit for the psychosocial intervention. For studies using a usual care comparison, only one study found a benefit for the psychosocial group,³⁷⁷ while two found no group differences at 12 to 24 weeks.^{367, 373} For studies using attention controls, two studies found a benefit for the psychosocial interventions^{354, 360} one of which reported moderate Cohen's d effect size of 0.86 (1.24, 0.46),³⁵⁴ while another study found no group differences.³⁶⁸

Caregiving skill was investigated by five studies using usual care and two using attention control groups, with six of the seven reporting no long-term group differences, and a single study reporting short-term benefit for the psychosocial intervention. For studies using usual care comparisons, one found short-term benefit at 6 weeks for the psychosocial group,³⁷⁴ one found no short-term benefit,⁴⁵⁷ and three found no difference in caregiving skill long term.^{224, 358, 362} An additional two studies using attention control comparisons also found no long-term group differences in caregiving skill.^{368, 370}

Coping was investigated by a single study using a usual care comparison. This study reported higher coping skills in the psychosocial group at 6 months.³⁵⁸

Quality of life was investigated by 11 studies using usual care and three using attention control groups. Eight studies found no group differences while six found benefit for the psychosocial group. For studies using usual comparison groups, four found short-term benefits for the psychosocial groups,^{366, 374, 376, 381} and three found no group differences.^{361, 364, 367} In the long term, only one study found a benefit for the psychosocial group,³⁷⁵ while three did not.^{224, 362, 363} For studies using attention control comparisons, a single study found short-term benefits for the psychosocial group;³⁵⁷ however, two found no group differences long term.^{356, 368}

The caregiver/PLWD relationship was investigated by three studies using attention control groups. One found a benefit for the psychosocial group and reported a moderate Cohen's d effect size of 0.44 (0.82, 0.07),³⁵⁴ while two found no group differences.^{356, 368}

Social support was investigated by two studies using usual care comparisons and one using attention control groups. The two studies using usual care comparisons found no group differences in social support.^{367, 376} A single study using an attention control group found a benefit for the psychosocial group at 6 months.³⁶⁸

Healthcare use was investigated by two studies using usual care and two using attention control groups. Only one study using an attention control group found a benefit for the psychosocial group,³⁶⁸ while all the others found no group differences.^{362, 365, 375}

No studies reported outcomes on caregiver harms for psychosocial interventions.

Table 6.3. Summary of findings for caregiver outcomes: psychosocial interventions for CG/P

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Anxiety Psychosocial vs Attention control	2 RCT ^{356, 360} n=358 24-52 weeks	CG/P	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Anxiety Psychosocial vs Usual care	1 RCT ³⁶⁶ n=170 6 weeks	CG/P	1 found benefit 0 found no difference	Insufficient
Anxiety Psychosocial vs Usual care	1 RCT ³⁶³ n=260 2 years	CG/P	0 found benefit 1 found no difference	Insufficient
Burden of care Psychosocial vs Attention control	5 RCT ^{354, 357, 359, 368, 380} n=725 8-24 weeks	CG/P	3 of 5 found benefit 2 of 5 found no difference	Insufficient
Burden of care Psychosocial vs Usual care	3 RCT ^{364, 366, 457} n=680 6-16 weeks	CG/P	0 of 3 found benefit 3 of 3 found no difference	Insufficient
Burden of care Psychosocial vs Usual care	5 RCT ^{224, 358, 361, 375, 376} n=561 24-52 weeks	CG/P	2 of 5 found benefit 3 of 5 found no difference	Insufficient
Depressive symptoms Psychosocial vs Attention control	7 RCT ^{354, 357, 359, 360, 368, 372, 380} n=976 8-24 weeks	CG/P	6 of 7 found benefit 1 of 7 found no difference	Insufficient
Depressive symptoms Psychosocial vs Attention control	3 RCT ^{356, 369, 370} n=916 52-72 weeks	CG/P	0 of 3 found benefit 3 of 3 found no difference	Insufficient

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Depressive symptoms Psychosocial vs Usual care	5 RCT ^{358, 366, 367, 374, 457} n=852 6-24 weeks	CG/P	2 of 5 found benefit 3 of 5 found no difference	Insufficient
Depressive symptoms Psychosocial vs Usual care	5 RCT ^{224, 355, 356, 363} n=1073 52-104 weeks	CG/P	2 of 5 found benefit 3 of 5 found no difference	Insufficient
Caregiving bother/distress/affect Psychosocial vs Attention control	5 RCT ^{356, 359, 360, 368, 372} n=952 4-12 months	CG/P	3 of 5 found benefit 2 of 5 found no difference	Insufficient
Caregiving bother/distress/affect Psychosocial vs Usual care	4 RCT ^{361, 366, 373, 457} n=702 6-24 weeks	CG/P	1 of 4 found benefit 3 of 4 found no difference	Insufficient
Healthcare usage Psychosocial vs Attention control	2 RCT ^{365, 368} n=359 3-6 months	CG/P	1 of 2 found benefit 1 of 2 found no difference	Insufficient
Healthcare usage Psychosocial vs Usual care	2 RCT ^{362, 375} n=251 4-24 months	CG/P	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Caregiving knowledge Psychosocial vs Usual care	1 RCT ³⁵⁸ n=167 24 weeks	CG/P	1 found benefit 0 found no difference	Insufficient
Quality of life Psychosocial vs Attention control	1 RCT ³⁵⁷ n=132 8 weeks	CG/P	1 found benefit 0 found no difference	Insufficient
Quality of life Psychosocial vs Attention control	2 RCT ^{356, 368} n=541 24-52 weeks	CG/P	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Quality of life Psychosocial vs Usual care	7 RCT ^{361, 364, 366, 367, 374, 376, 381} n=843 6-24 weeks	CG/P	4 of 7 found benefit 3 of 7 found no difference	Insufficient
Quality of life Psychosocial vs Usual care	4 RCT ^{224, 362, 363, 375} n=652 9 months to 2 years	CG/P	1 of 4 found benefit 3 of 4 found no difference	Insufficient
Relationship with person with dementia Psychosocial vs Attention control	4 RCT ^{354, 356, 368, 477} n=652 20-52 weeks	CG/P	1 of 4 found benefit 3 of 4 found no difference	Insufficient
Caregiving confidence Psychosocial vs Attention control	3 RCT ^{354, 360, 368} n=428 20-24 weeks	CG/P	2 of 3 found benefit 1 of 3 found no difference	Insufficient
Caregiving confidence Psychosocial vs Usual care	3 RCT ^{367, 373, 457} n=582 12-24 weeks	CG/P	1 of 3 found benefit 2 of 3 found no difference	Insufficient
Caregiving skill Psychosocial vs Attention control	2 RCT ^{368, 370} n=545 6-18 months	CG/P	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Caregiving skill Psychosocial vs Usual care	2 RCT ^{374, 457} n=407 6-16 weeks	CG/P	1 of 2 found benefit 1 of 2 found no difference	Insufficient

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Caregiving skill Psychosocial vs Usual care	3 RCT ^{224, 358, 362} n=444 24-52 weeks	CG/P	0 of 3 found benefit 3 of 3 found no difference	Insufficient
Stress Psychosocial vs Attention	1 RCT ³⁷² n=184 6 months	CG/P	1 found benefit 0 found no difference	Insufficient
Stress Psychosocial vs Usual care	3 RCT ^{355, 366, 367} n=486 6-52 weeks	CG/P	0 of 3 found benefit 3 of 3 found no difference	Insufficient
Social support Psychosocial vs Attention control	1 RCT ³⁶⁸ n=250 6 months	CG/P	1 found benefit 0 found no difference	Insufficient
Social support Psychosocial vs Usual care	2 RCT ^{367, 376} n=188 12-24 weeks	CG/P	0 of 2 found benefit 2 of 2 found no difference	Insufficient
Caregiving coping Psychosocial vs Usual care	1 RCT ³⁵⁸ n=167 24 weeks	CG/P	1 found benefit 0 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: CG/P=caregiver/care partner; n=number; RCT=randomized controlled trial

Variation in Outcomes

By PLWD Characteristics

Only one study identified how PLWD characteristics may influence outcomes. A subgroup analysis for the Farran study (2004) found that data from 143 caregivers of family members with baseline agitated behaviors indicated more reduction of emotional distress with the skill-building intervention than with an information- and support-oriented comparison over an 18-month period. These researchers suggested their findings indicate that dementia caregivers exposed to agitated behaviors can benefit from psychosocial interventions, particularly those aimed at building behavioral management skills.³⁷⁰

By Caregiver Characteristics

Only a few studies identified how caregiver characteristics may influence outcomes. Gallagher-Thompson and colleagues (2003) found no differences between Hispanic and nonHispanic white female caregivers (n=213) in caregiver coping, depression, social support, and caregiver burden 3 months after completing the Coping with Caregiving intervention.³⁷¹

Gallagher-Thompson and colleagues in 2008 investigated differences in caregiver coping, depression, stress, support, burden, and bother between Hispanic and nonHispanic white female caregivers (n=184).³⁷² The only differences they found were that nonHispanic white female caregivers reported higher bother managing neuropsychiatric symptoms at baseline (p=0.03) and post intervention (p=0.010) than Hispanic female caregivers. The two groups did not differ in depressive symptoms or perceived stress before or 6 months after completing a psychosocial intervention.

Gitlin and colleagues (2001) reported that intervention spouses reported reduced upset (p=.049), women reported enhanced self-efficacy in managing PLWD behaviors (p=.038), and women (p=.049) and racial/ethnic minorities (p=.037) reported enhanced self-efficacy in managing PLWD functional dependency.³⁷³

A moderator analysis of the Gallagher-Thompson (2003) intervention³⁷¹ found that caregiver self-efficacy predicted differential outcomes in a randomized trial comparing a cognitive behavior psychoeducational intervention versus an enhanced support group. The four key outcomes were depression, anxiety, social support, and coping. The findings showed that low baseline self-efficacy scores better predicted positive response to treatment in the psychosocial intervention than in the enhanced support group intervention. This study supports the use of self-efficacy as a screening tool for appropriate caregiver intervention assignment.⁴⁷⁸

Otero and colleagues (2015) reported that caregivers younger than 65 years and with higher emotional distress at baseline were more likely than those receiving usual care to improve their depressive symptoms after completing five weeks of a group-based cognitive behavioral program focused on problem solving.³⁷⁹

By Intervention Characteristics

No studies investigated whether intervention characteristics modified the intervention effects.

Social Support

Key Point

- Evidence was insufficient to draw conclusions about the effect of in-person social support on PLWD and their CG/P.
- Phone-based social support for CG/P were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

We defined social support interventions as those targeted to provide information, resources, and a form of social interaction to caregivers of PLWD. Intervention delivery modes could involve talking to peers or other supports in-person or over the phone, or automated support programs that were phone or web-based. Phone and web-based platforms provided some form of social interaction either through automated conversations, voicemail, or chat groups.

Eligible Studies

We identified 13 unique studies from 16 publications (Table 6.4).⁴⁷⁹⁻⁴⁹³ Two studies were assessed as medium risk of bias for the analytic set.⁴⁷⁹⁻⁴⁸¹ We provide information on all pilot studies and high risk of bias studies as part of the evidence map in Appendix E.

The two studies in the analytic set were categorized as being in the explanatory stage, or Stage 3 of the NIH Stage Model. One (n=78) was a randomized trial comparing in-person, peer-led mutual support groups for CG/P with usual care.⁴⁸¹ Support groups met bi-weekly over 24 months. The other (n=100) was a randomized trial comparing an automated phone support system for CG/P with usual care.^{479, 480} The phone support system included an activity-respite module that engaged PLWD in an 18-minute automated conversation, weekly automated conversations to check stress levels, a personal voice mailbox, and a phone support group “bulletin board” (which functioned like an internet chat group). The support system was available for 22 hours a day over 12 months.^{479, 480} Appendix E provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 6.4. Basic characteristics of literature set: in-person social support

Characteristic	Information
Total Studies	13 studies
Non-U.S. studies	10 studies
Evidence map studies	4 pilot studies 7 high risk of bias studies
Analytic set studies	2 randomized controlled trials
Risk of bias of analytic set	2 medium
Number of PLWD in analytic set	178
Dementia type/definition	All dementia types and severities (primarily AD and cerebrovascular) Probable AD with functional impairments and AD-related disturbing behavior
Caregiver type (number)	Informal caregivers (n=178)

Abbreviations: AD=Alzheimer’s disease; n=number; PLWD=people living with dementia

Intervention Research Context

The eligible literature on social support services for caregivers generally comprises pilot or explanatory research. Studies were published between 1990 and 2019 and tended to be pilot, small sample, or high risk of bias studies. One intervention did show studies progressing along the NIH Stage Model.^{489, 490}

In-person social support approaches included a befriending intervention where volunteers were recruited and trained to befriend CG/Ps.^{481, 482, 485, 487} One study examined social support groups that engaged CG/P and PLWD dyads separately and together.⁴⁸⁵ The automated social support for CG/P comprised phone-based automated support and web-based automated support. We identified only one study that examined the use of phone-based automated support, which was included in our analytic study set.^{479, 480} The lack of additional studies on phone-based automated support may be due to a shift towards web-based platforms as technology has evolved. Four web-based approaches were piloted in Europe.^{486, 488, 489} One piloted a tool allowing caregivers to assess their needs for care and support and providing tailored advice as well as links to local support organizations. Two piloted platforms providing information, relaxation exercises, a social networking platform, and allowing caregivers to assess their health and reach out to clinical sites. An additional pilot examining the use of a website to provide stress reduction, support, and information for caregivers was conducted in the United States.⁴⁹⁴ The pilots suggested some benefit to web-based social support tools. However, a later small study was assessed as high risk of bias.⁴⁹⁰ One study examined email contacts with a specialist dementia nurse with online videos and e-bulletins.⁴⁹³

PLWD Outcomes

No reportable PLWD outcomes were available from the analytic set.

Caregiver Outcomes

Evidence was insufficient to draw conclusions about in-person social support versus usual care for CG/P. Table 6.5 summarizes the primary outcomes for CG/P. At 28 weeks, findings showed statistically significant differences in Neuropsychiatric Inventory-Caregiver Distress Scale scores between the peer support and usual care groups, with the peer support group reporting lower levels of distress.⁴⁸¹ The peer support group also had statistically significant improvements in quality of life at 28 weeks compared with the usual care group. Groups did not differ in the use of mental health services. No harms were reported.

Evidence was also insufficient to draw conclusions about automated social support versus usual care for CG/P. At 6 months, findings showed no difference between the automated phone-based support group and usual care in depressive symptoms.^{479, 480} Additionally, no difference was found between groups in anxiety.^{479, 480} No study assessed harms.

Table 6.5. Summary of findings for caregiver outcomes: social support

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Quality of Life In person social support vs. Usual care	1 RCT ⁴⁸¹ (n=78) 28 weeks	CG/P, respite centers in China	1 found benefit 0 found no difference	Insufficient
Caregiver Distress In person social support vs. Usual care	1 RCT ⁴⁸¹ (n=78) 28 weeks	CG/P, respite centers in China	1 found benefit 0 found no difference	Insufficient
Depression Automated social support vs. Usual care	1 RCT ⁴⁷⁹ (n=100) 6 months	CG/P	0 found benefit 1 found no difference	Insufficient
Anxiety Automated social support vs. Usual care	1 RCT ⁴⁷⁹ (n=100) 6 months	CG/P	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: CG/P=caregiver/care partner; n=number; RCT=randomized controlled trial

Variation in Outcomes

No studies reported variation in outcomes by PLWD, caregiver, or intervention characteristics.

Lifestyle Interventions

Lifestyle interventions include a range of interventions targeted at improving the health and well-being of participants, including physical activity, leisure activities, visual arts, and mindfulness, meditation, or spiritually focused activities.

Mindfulness, Meditation, or Spiritually Focused Activities

Key Point

- Evidence was insufficient to draw conclusions about the effect of mindfulness-based stress reduction on PLWD and their CG/P.

Eligible Studies

We identified 10 unique studies from 10 publications examining mindfulness, meditation, or spiritually focused activities. Table 6.6 summarizes the characteristics of the literature set. Only one study was included in the analytic study set.⁴⁹⁵ The study was assessed as medium risk of bias and categorized as explanatory, or Stage 3 of the NIH stage model. Three studies were pilot studies and four studies had fewer than 50 participants.⁴⁹⁶⁻⁵⁰² Two studies were assessed as high risk of bias due to potential performance bias.^{503, 504} We provide information on all pilot studies and high risk of bias studies in the evidence map in Appendix E. Appendix E also provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 6.6. Basic characteristics of literature set: mindfulness, meditation, and spiritually focused activities

Characteristic	Information
Total Studies	10 studies
Non-U.S. studies	5 studies
Evidence map studies	3 pilot studies 4 small sample size studies 2 high risk of bias studies
Analytic set studies	1 randomized controlled trial
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	31
Dementia type/definition	Progressive dementia diagnosis
Caregiver type (number)	Informal caregivers (31)

Abbreviation: PLWD=people living with dementia

Intervention Research Context

The literature set follows the NIH Stage Model, with most research consisting of pilots and small RCTs. Studies were published between 2010 and 2019, reflecting the relatively recent interest in these types of interventions for CG/P of PLWD. These studies generally used group sessions as the intervention delivery mode, and they focused on therapy or education using mindfulness, religion, or meditation as a main feature of the intervention. Interventions lasted from 5 to 12 weeks. Five studies specifically examined the effect of mindfulness therapy or mindfulness-based stress reduction for CG/P.^{495, 496, 498, 501, 502} Two studies examined spiritually based therapy or education.^{503, 504} Three studies examined meditation.^{497, 499, 500}

PLWD Outcomes

The analytic study set did not report outcomes related to PLWD.

Caregiver Outcomes

One study was included in the analytic set and reported outcomes for CG/P.⁴⁹⁵ The study (n=78) randomized CG/P to either mindfulness-based stress reduction or education and support. Evidence was insufficient to draw conclusions about the impact of mindfulness-based stress reduction versus education and support. In both arms, participants attended in-person group sessions over 8 weeks. The education and support group also received support calls by phone. The study reported outcomes for caregiver stress, burden, depression, anxiety, mental health, physical health, and social support at 2 months (immediately post-intervention) and 6 months.⁴⁹⁵

Table 6.7 summarizes the outcomes for CG/P. Groups did not differ significantly at either time point for measures of caregiver burden, social support, or physical health.⁴⁹⁵ Participants in the mindfulness group showed statistically significant improvements in mental health versus the education and support group at 2 and 6 months.⁴⁹⁵ Participants in the mindfulness group also showed a reduction in anxiety at 2 and 6 months versus the education and support group. At 2 months, participants in the mindfulness group showed a statistically significant reduction in stress versus those in the education and support group. This difference was not sustained at 6 months.⁴⁹⁵ However, this was also true for depression, where the mindfulness group showed a statistically significant improvement in depressive symptoms versus the education and support group at 2 months, but not 6 months.⁴⁹⁵

Table 6.7. Summary of findings for caregiver outcomes: mindfulness, meditation, and spiritually focused activities

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Caregiver Stress Mindfulness vs Active Control	1 RCT (n=78) ⁴⁹⁵ 2 months, 6 months	CG/P	1 found benefit 0 found no difference	Insufficient
Caregiver Burden Mindfulness vs Active Control	1 RCT (n=78) ⁴⁹⁵ 2 months, 6 months	CG/P	0 found benefit 1 found no difference	Insufficient
Depression Mindfulness vs Active Control	1 RCT (n=78) ⁴⁹⁵ 2 months, 6 months	CG/P	1 found benefit at 2 months but no difference at 6	Insufficient
Anxiety Mindfulness vs Active Control	1 RCT (n=78) ⁴⁹⁵ 2 months, 6 months	CG/P	1 found benefit at 2 months but no difference at 6	Insufficient
Mental Health Mindfulness vs Active Control	1 RCT (n=78) ⁴⁹⁵ 2 months, 6 months	CG/P	1 found benefit 0 found no difference	Insufficient
Physical Health Mindfulness vs Active Control	1 RCT (n=78) ⁴⁹⁵ 2 months, 6 months	CG/P	0 found benefit 1 found no difference	Insufficient
Social Support Mindfulness vs Active Control	1 RCT (n=78) ⁴⁹⁵ 2 months, 6 months	CG/P	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: CG/P=caregiver/care partner; n=number; RCT=randomized controlled trial

Variation in Outcomes

The analytic study set did not report variation in outcomes by PLWD, caregiver, or intervention characteristics.

Physical Activity and Other Lifestyle Interventions

Key Point

- No physical activity or other lifestyle interventions advanced from the evidence map for further analysis.

Intervention Description

Physical activity interventions may involve caregivers alone or caregiver/PLWD dyads, and often involve an interventionist encouraging (in person or via phone or through written materials) physical activity, helping with goal setting, and/or supervising the implementation of assigned physical activity.

Eligible Studies

We identified five unique studies from six publications on physical activity interventions⁵⁰⁵⁻⁵⁰⁹ and four unique studies from four publications focused on other various lifestyle interventions for CG/P.⁵¹⁰⁻⁵¹³ Table 6.8 summarizes the characteristics of the literature set. No studies were eligible for the analytic set. Four studies were assessed as high risk of bias due to potential performance bias.^{505-507, 509} The remaining study was a pilot.⁵⁰⁸ Appendix E presents information on all studies as part of the evidence map.

Table 6.8. Basic characteristics of literature set: physical activity and other lifestyle interventions

Characteristic	Information
Total Studies	9 studies
Non-U.S. studies	5 studies
Evidence map studies	2 pilot study 7 high risk of bias studies
Analytic set studies	0 studies

Intervention Research Context

Studies were published between 2002 and 2018. Generally, this literature did not progress along the NIH Stage Model, as we identified only two pilot studies. One study enrolled CG/P and PLWD dyads to jointly participate in physical activity.⁵⁰⁹ Of the five studies that examined physical activity, three focused on phone-based physical activity interventions.⁵⁰⁵⁻⁵⁰⁷ Phone contact was used to encourage physical activity, set goals, and/or supervise the implementation of assigned physical activity. The remaining two physical activity studies assigned CG/P a physical activity regimen without the use of phone support.^{508, 509} Of four studies that examined leisure interventions, two focused on leisure activities or leisure education for CG/P,^{510, 513} while two others enrolled CG/P and PLWD dyads.^{511, 512} Of the dyad studies, one study compared a structured visual arts education program (with visual arts projects of increasing difficulty) for CG/P and PLWD versus painting and discussion about art.⁵¹¹ The second study compared providing assistance to female spouse caregivers and PLWD while on vacation versus a waitlist control.⁵¹²

Respite Care

Key Point

- No respite care interventions advanced from the evidence map for further analysis.

Intervention Description

Respite care interventions provide temporary breaks to caregivers of PLWD. Interventions include in-home care for PLWD, adult day care programs, and institutional respite services. These interventions provide breaks for a few hours a day or allow individuals to take a full break from caregiving for a short-term period of time.

Eligible Studies

We identified three unique studies from four publications (Table 6.9).⁵¹⁴⁻⁵¹⁷ All three were assessed as high risk of bias due to potential attrition bias, and were not included in the analytic set. Two studies used a quasi-experimental study design.⁵¹⁵⁻⁵¹⁷ The remaining study was an RCT.⁵¹⁴ We present information on all high risk of bias studies as part of the evidence map in Appendix E.

Table 6.9. Basic characteristics of literature set: respite care

Characteristic	Information
Total Studies	3 studies
Non-U.S. studies	1 study
Evidence map studies	3 high risk of bias
Analytic set studies	0 studies

Intervention Research Context

The studies were published between 1989 and 2019 and compared a range of respite care activities for CG/P versus usual care. One study examined in-home 24-hour respite care for 5 to 14 days,^{515, 516} one study examined adult day care for PLWD (twice or more per week),⁵¹⁷ and one study examined a program offering in-home day care versus institutional respite services used at the discretion of CG/P.⁵¹⁴

Multicomponent Interventions

Key Point

- Intensive multicomponent intervention with education, group discussion, in-home and phone support sessions, and caregiver feedback for CG/P support (i.e. discrete adaptations of REACH II), improved CG/P depression at 6 months.(low-strength evidence)
- Evidence was insufficient to draw conclusions about the effect of other forms of multicomponent interventions on PLWD and their CG/P.

Intervention Description

Multicomponent interventions include more than one care technique or delivery method to foster support, expertise, information, or skills for caregivers in order to improve caregiver quality of life and health outcomes. Care techniques and delivery methods found in multicomponent interventions include therapy and support such as counseling (in groups and/or one-on-one, phone-based and/or in person), training (in person with key personnel, one-on-one and/or paired, group, or classroom style, and/or via written materials), supportive feedback, goal-setting, and planning (in any setting, with family members, in groups, and in person or remote/via phone), as well as relaxation and physical exercises. Offered as structured, programmatic approaches to strengthen communication, develop resources and skills, and/or create or maintain relationships for caregivers and PLWD, multicomponent interventions may benefit informal (family) caregivers.

Eligible Studies

Multicomponent interventions targeting quality of life and other outcomes for informal and formal caregivers included seven studies of three different multicomponent interventions across 22 publications on multicomponent interventions (Table 6.10).^{383, 518-540} All seven multicomponent interventions in the analytic set targeted outcomes for caregivers by offering multiple intervention components within a structured, programmatic approach for community-dwelling caregivers living with PLWD. We provide information on all pilot studies and high risk of bias studies as part of the evidence map in Appendix E. Appendix E also provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 6.10. Basic characteristics of literature set: multicomponent for CG/P

Characteristic	Information
Total unique studies	22 studies
Non-U.S. studies	13 studies

Characteristic	Information
Total Evidence map studies	4 pilot studies 2 small sample studies 9 high risk of bias studies
Analytic set studies	6 randomized controlled trials 1 pragmatic trial
Risk of bias of analytic set	7 medium/low
Number of PLWD in analytic set	1688
Dementia type/definition	All types of dementia with all stages of severity
Caregiver type (number)	Informal caregivers (1688)

Abbreviation: PLWD=people living with dementia

Intervention Research Context

Studies published from 1993 to 2017 tended to be small or high risk of bias, and largely based on pilot or explanatory research. More trials of structured multicomponent interventions, focusing on how treatments correspond to improvement in outcomes for caregivers (via intervention for caregiver and PLWD dyads) appeared in six of the studies.

The literature in the evidence map and the analytic set contains a considerable amount of preliminary work. However, we observed more development along the NIH Stage Model in this set than in most other intervention categories. This literature set demonstrates growth over time toward the development of both pragmatic trials as well as dissemination/implementation research. Specific examples include the REACH II intervention^{519, 520} and the RDAD dual-component intervention.^{535, 537}

PLWD Outcomes

One study reported benefits for PLWD physical role functioning and depression.⁵⁴¹ Evidence was insufficient to draw conclusions about PLWD outcomes.

Caregiver Outcomes

Table 6.11 provides a summary of findings. Structured multicomponent interventions presented low-strength evidence for improving caregiver depression.^{519, 520, 528}

Three studies tested an intensive multicomponent intervention aimed at improving outcomes for community-dwelling informal (family) caregivers. The intervention incorporated education, group discussion, in-home and phone support sessions, and caregiver feedback, four components, for CG/P support across a range of characteristics.^{519, 520, 528} This intervention was developed to manage behavioral problems, reduce functional dependence, and prevent functional impairment. Low-strength evidence showed reduced depression for CG/P. One U.S.-based study found moderate effect size for depression at 6 months.⁴⁸² This result was consistent with a second study that found no difference between groups but improvement over baseline in both intervention and control groups.⁵²⁸ For outcomes including caregiver quality of life, stress, burden, caregiver support, and nursing home placement, we found too much variation in outcomes definition, measurement, and reporting within⁴⁸² and across studies^{482, 483, 491} to draw conclusions. An assessment of insufficient evidence does not mean that the intervention is determined to be of no value. Rather, it means that due to the uncertainty of the evidence we could not draw meaningful conclusions at this time.

Evidence was insufficient from three studies to assess one intervention that used a structured, three-component approach comprising in-person counseling sessions, family counseling sessions

involving family members whom caregivers invited, and phone counseling sessions for caregivers on-demand across three countries. Evidence was also insufficient from two studies for one intervention examining combined exercise and CG/P support.

Table 6.11. Summary of findings for caregiver outcomes: multicomponent for CG/P

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Health Combined exercise & support vs. monthly phone calls & mailed bulletins	1 RCT ^{535, 541} (n=153) 12 weeks	Community-dwelling PLWD and Informal (family) caregivers	1 found benefit 0 found no difference	Insufficient
Depression Counseling sessions (in-person, family, and phone) for caregivers vs. usual care	1 RCT ^{528, 531, 533} (n=371) 4-6 months	Spouse caregivers living at home with PLWD	1 found benefit 0 found no difference	Insufficient
Caregiver stress perception Counseling sessions (in-person, family, and phone) for caregivers vs. usual care	1 RCT ^{528, 531, 533} (n=158) 4-6 months	Spouse caregivers living at home with PLWD	1 found benefit 0 found no difference	Insufficient
Depression Education, group discussion, in-home and phone support sessions, and caregiver feedback vs usual care**	3 RCT ^{519, 520, 528} (n=895) 6 months	Family caregivers (living with or sharing cooking) with PLWD	2 of 3 found benefit 1 of 3 found no difference	Low discrete adaptations of REACH II
Caregiver stress, burden Education, group discussion, in-home and phone support sessions, and caregiver feedback vs usual care**	3 RCT ^{519, 520, 528} (n=895) 6 months	Family caregivers (living with or sharing cooking) with PLWD	1 of 3 found benefit 2 of 3 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

**Luchsinger et al. compares REACH II to New York University Caregiver interventions and finds no difference between groups but improvement over baseline in both groups.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial; REACH II=Resources for Enhancing Alzheimer’s Caregiver Health II

Variation in Outcome

Studies reported variation in benefit across racial and ethnic groups studied.^{519, 528} In one study, spouse caregivers in the intervention experienced significantly greater improvement in quality of life than those in the control group for all included racial and ethnic groups (Hispanic or Latino, white, and black or African-American).⁵¹⁹ Another study reported benefits for both a four-component intervention and a three-component intervention⁵³³ for Hispanic caregivers.⁵²⁸

Other Interventions for Caregiver Well-Being

Key Point

- Studies of several other types of interventions were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

This set of diverse interventions included approaches such as support meetings, decision aids, virtual reality experience to increase empathy, and cranial electrical stimulation of the CG/P.

Eligible Studies

We identified six unique studies from seven publications, each examining a unique intervention.⁵⁴²⁻⁵⁴⁷ We provide characteristics of the literature set in Table 6.12, and information on all studies in the evidence map in Appendix E.

Table 6.12. Basic characteristics of literature set: other interventions for CG/P

Characteristic	Information
Total Studies	6 studies
Non-U.S. studies	4 studies
Evidence map studies	3 pilot studies 3 high risk of bias studies
Analytic set studies	0 studies

Intervention Research Context

The literature set was highly varied, comprising unrelated studies that represented earlier stages of the NIH Stage Model. Two studies were conducted in the United States. One study examined support meetings for CG/P to assist in conversations with PLWD about driving cessation.⁵⁴⁶ The other study examined cranial electrical stimulation for CG/P to improve sleep.⁵⁴⁴ Four studies were conducted in non-U.S. settings. One study examined an aid for CG/P in Australia to make decisions about community services including respite care.⁵⁴⁷ Another study examined an aid for CG/P in the United Kingdom to make decisions about place of residence for PLWD.⁵⁴³ One study in the Netherlands examined virtual reality devices to give CG/P a simulated experience of having dementia.⁵⁴² The last study, conducted in the United Kingdom, examined the use of a caregiver-held record folder that included sections for caregiver/family and professional feedback as well as a caregiver diary.⁵⁴⁵

Conclusion

We identified many studies with a large number of participants that investigated the efficacy of care interventions to improve the well-being of CG/P. Applying the framework for care interventions from the NASEM Families Caring for an Aging America 2016 report (Figure 1.1 in Chapter 1), this category included interventions delivered at two levels: the individual level (i.e., psychosocial programs, lifestyle interventions, and multicomponent interventions) and the social or community level (i.e., social support and respite care). The vast majority of studies investigated psychosocial interventions delivered at the individual level. Fewer investigated multicomponent and lifestyle interventions delivered at the individual level. Very few investigated social support and respite care delivered at the social or community level.

Using the NIH Stage Model as a guide, about one-third of studies were pilot studies, and almost all of the others were explanatory, or Stage 3. Only a few multicomponent interventions were Stage 4 pragmatic trials. Of the explanatory studies, two thirds were rated as high risk of bias and excluded from the analytic set. Ultimately, just over 20 percent of the literature was eligible for the analytic set.

Included studies investigated many types of PLWD and caregiver outcomes. Only psychosocial studies examined PLWD outcomes; the most common were function, neuropsychiatric symptoms, and proxy-rated quality of life. Several important PLWD outcomes, such as depression, healthcare use, social support, and nursing home admission, were infrequently investigated. All of the studies investigated outcomes for the CG/P; the most common were depression, quality of life, caregiving burden, and caregiving bother and distress.

Several important CG/P outcomes, such as coping, caregiving knowledge, social support, and healthcare use, were infrequently investigated. Other important outcomes, such as those related to resilience and the positive aspects of caregiving, and social health outcomes, such as social isolation, were also missing from the literature.

While the literature set was large, the overwhelming majority of the evidence was insufficient to draw conclusions about whether or not these interventions work. We found, however, that one intensive multicomponent intervention (i.e. discrete adaptations of REACH II) did have low-strength evidence for improving depression among CG/P. Of all included interventions to improve caregiver well-being, multicomponent interventions used the most targeted components. Possibly, these comprehensive interventions addressed at least one critical need (across a wide range of individual caregiver needs), thus improving outcomes. Multicomponent interventions were structured to provide the same components to all participants. Very few studies were delivered at the social and community level. Only 12 studies investigated social support, and just two of these had low to medium risk of bias. The three studies that examined respite care were all assessed as high risk of bias.

About half of the studies were conducted outside of the United States, and all but one of these occurred in high-resource countries. The vast majority of studies conducted in the United States enrolled white urban or suburban dwelling caregivers. Only a few studies investigated Hispanic and African American caregivers and caregivers living in rural and underserved urban areas.

Our assessment of the evidence as insufficient should not be interpreted to mean that we concluded these interventions do not work. Rather, it means that the findings were too uncertain to draw conclusions about their combined effect. Research on interventions to change behavioral and psychological outcomes is challenging, and many factors can influence the outcomes. The insufficient ratings were driven mainly by the inconsistency and imprecision of study findings. For almost every outcome investigated, findings from several trials were inconsistent; some studies found improvement for the intervention group, and some found no difference in outcomes between the intervention and comparison groups. For studies that found no difference, often the comparison groups scores improved more than the intervention group (although not enough to be statistically significant). The findings were also generally imprecise. Our ability to combine results from several studies was hampered by the heterogeneity of the intervention components, duration and intensity of treatment, and comparison groups. Our ability to combine results from several studies statistically was generally not possible, because studies rarely reported the data necessary to calculate effect sizes.

Chapter 7. Care Interventions for Formal Caregivers

This chapter includes care interventions with elements designed to address the well-being of formal caregivers in the workplace. We present Key Points followed by results in three general sections: Intervention Description, Eligible Studies, and Intervention Research Context. Since no studies were assessed as low to medium risk of bias, we present the studies from the evidence map with a brief discussion of what has been examined, and research context. We present detailed information on all eligible studies in Appendix E.

Formal Caregiver Well-Being

Key Point

- Studies of interventions to improve formal caregiver well-being were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Interventions categorized as targeting formal caregiver well-being include components such as peer support, stress management, and relaxation techniques.

Eligible Studies

We identified three unique studies from five publications that examined interventions to improve the well-being of formal caregiving staff in long-term care facilities.⁵⁴⁸⁻⁵⁵² (Table 7.1) All were pilot studies. We provide information on all studies in the evidence map in Appendix E.

Table 7.1. Basic characteristics of literature set: formal caregiver well-being

Characteristics	Information
Total unique studies	3 studies
Non-U.S. studies	3 studies
Evidence map studies	3 pilot studies
Analytic set studies	0

Intervention Research Context

The research is very preliminary, with small pilot studies conducted in non-U.S. settings from 2003 to 2015. Similarly to research conducted to support informal caregivers, the Canadian 2003 study tested the premise that education and training to improve skills will reduce formal caregiver stress (in this case as measured by burnout).⁵⁵¹ Later studies examined more direct interventions such as peer support in Australia^{550, 552} and stress management and relaxation techniques to reduce job-related stress and burnout in Portugal.^{548, 549}

Conclusion

The literature for formal caregivers is preliminary. Possibly, our search algorithm (designed to maximize the ability to find therapies for PLWD) limited our ability to locate related literature. We chose not to include the workforce literature, since this review is focused on the well-being of both PLWD and caregiver. We anticipated that this association would be more prominent in the informal caregiver literature, but not well-studied in the workforce literature.

Chapter 8. Care Delivery Interventions

This chapter includes care delivery interventions intended to improve how care is delivered. These interventions target the organization of care, including the scheduling of staff and tasks as well as the ways in which staff and tasks are interdependent. While care delivery interventions may be implemented through education and training, they differ from most education and training in that they do not aim to improve already established roles and tasks. Instead, they aim to *change* the tasks that are performed, the set of staff who perform the tasks, or the way the staff work together. This chapter is organized into three categories: care service provision, care delivery models or programs, and care staff education and support needs.

For each intervention, we present Key Points followed by results in three general sections: Intervention Description, Eligible Studies, and Intervention Research Context. For interventions with no studies assessed as low to medium risk of bias, we present the studies from the evidence map with a brief discussion of what has been examined as well as the research context. For interventions for which low to medium risk of bias studies were available for an analytic set, we present Outcomes sections by people living with dementia (PLWD), by caregiver, and by variation in outcomes when available. We use the term caregiver/partner, or CG/P, for informal caregiver outcomes. Because we generally could not pool outcomes for any given intervention and comparison group, we synthesized the information qualitatively; therefore, we present summary findings as brief statements of how many studies reported statistically significant benefits or no statistically significant difference between the intervention and the comparator. We present detailed information on all eligible studies in Appendix F.

Care Service Provision

Care service provision refers to the act of providing care for the PLWD. In general business terms, specialized service provision usually requires qualified staff members to be regularly available to perform the services offered. The qualifications needed for staff will depend on the care being provided or supported.

Consultation Services

Key Point

- Evidence was insufficient to draw conclusions about the effect of consultation services on PLWD and their caregivers.

Intervention Description

Consultation services include individualized and multidisciplinary approaches targeting PLWD as well as their caregivers. Most of the consultants interacted with CG/P not only to improve well-being among PLWD, but also to improve the health of CG/P by improving their overall knowledge for managing dementia and reducing their burden and depression. While this category of interventions has the potential to be very broad, we identified only a few forms examined in the eligible literature, namely a dementia outreach service, individualized consultations, multidisciplinary care consultations, and a needs assessment for older adults.

Eligible Studies

Table 8.1 summarizes the characteristics of the literature set. We identified five unique studies that examined the use of consultation interventions for improving quality of life and reducing hospitalizations and depression among PLWD, and improving efficiency of response among CG/P.⁵⁵³⁻⁵⁵⁷ Four studies were assessed as high risk of bias.^{553, 555-557} Only one study was included in the analytic set.⁵⁵⁴ We present information on the high risk of bias studies in the evidence map in Appendix E.

The included study (n=84) was a randomized controlled trial targeted at family (informal) caregivers. It compared consultation services versus an attention control.⁵⁵⁴ The study was assessed as medium risk of bias and categorized as explanatory, or Stage 3 of the National Institutes of Health (NIH) Stage Model. Appendix C Tables provide evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 8.1. Basic characteristics of literature set: consultation services

Characteristics	Information
Total unique studies	5 studies
Non-U.S. studies	4 studies
Evidence map studies	4 high risk of bias studies
Analytic set studies	1 randomized controlled trial
Risk of bias of analytic set	1 medium
Number of PLWD in analytic set	84
Dementia type/definition	All dementia types and severities
Caregiver type (number)	Informal caregivers (84)

Abbreviation: PLWD=people living with dementia

Intervention Research Context

The literature on consultation services comprises mostly high risk of bias studies in non-U.S. locations. We identified a dementia outreach study led by nurse practitioners in Australia; the intervention aimed to assist staff at residential care facilities to better manage behavioral and psychological symptoms of dementia (BPSD).⁵⁵³ We also identified two other Australian studies that used multidisciplinary consultation approaches to help nursing home staff improve care in order to reduce PLWD depression⁵⁵⁵ and BPSD.⁵⁵⁶ An additional study conducted in the United Kingdom examined a liaison-mediated intervention to reduce unmet needs and improve quality of life among community-dwelling PLWD.⁵⁵⁷

PLWD Outcomes

Evidence was insufficient to draw conclusions about nursing home admission among PLWD who received individualized consultation versus attention control interventions (n=84)⁵⁵⁴ Table 8.2 summarizes the findings.

Table 8.2. Summary of findings for PLWD outcomes: consultation services

Outcomes Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Nursing home admission Consultation vs Usual Care	1 RCT ⁵⁵⁴ (n=84) 12 months	PLWD	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Evidence was insufficient to draw conclusions about individualized consultation versus attention control (n=84).⁵⁵⁴ Table 8.3 summarizes the findings. The study reported no statistically significant improvement among caregivers in physical health symptoms or self-efficacy for managing dementia, and no reduction in caregiver depression or care burden.

Table 8.3. Summary of findings for caregiver outcomes: consultation services

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Physical health Consultation vs Usual Care	1 RCT ⁵⁵⁴ (n=84) 12 months	Primary family caregivers	0 found benefit 1 found no difference	Insufficient
Self-efficacy score Consultation vs Usual Care	1 RCT ⁵⁵⁴ (n=84) 12 months	Primary family caregivers	0 found benefit 1 found no difference	Insufficient
Depression Consultation vs Usual Care	1 RCT ⁵⁵⁴ (n=84) 12 months	Primary family caregivers	0 found benefit 1 found no difference	Insufficient
Burden Consultation vs Usual Care	1 RCT ⁵⁵⁴ (n=84) 12 months	Primary family caregivers	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial

Variation in Outcomes

Studies in the analytic set did not examine variation in outcomes by PLWD, caregiver, or intervention characteristics.

Case Management

Key Point

- Evidence was insufficient to draw conclusions about the effect of case management for PLWD and their caregivers.

Intervention Description

Case management is a service to support CG/P and their PLWD care recipient. Case managers help coordinate health and social services to support PLWD and their caregivers. Case managers commonly work with CG/P while the PLWD is still living in the community, although case management can also be engaged in nursing homes or other long-term care facilities. We categorized an intervention as case management if the study did not report the case or care manager as being embedded in a team-based care approach.

Eligible Studies

Table 8.4 summarizes the characteristics of the literature set. We identified nine unique studies from 12 publications that examined the use of case management for improving health outcomes for PLWD and their caregivers. Three studies, all non-U.S. settings, were assessed as

low to medium risk of bias and included in the analytic set.⁵⁵⁸⁻⁵⁶¹ Excluded from the analytic set were high risk of bias studies,⁵⁶²⁻⁵⁶⁷ which we describe in the evidence map in Appendix F. Evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes for the analytic set are also in Appendix F.

Table 8.4. Basic characteristics of literature set: case management

Characteristics	Information
Total unique studies	9 studies
Non-U.S. studies	6 studies
Evidence map studies	1 pilot/demonstration study 5 high risk of bias studies
Analytic set studies	3 (non-U.S.) studies
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	294
Dementia type/definition	Alzheimer's disease or dementia unspecified
Caregiver type (number)	Informal caregivers (294)

Abbreviation: PLWD=people living with dementia

Intervention Research Context

Eligible studies on case management were published between 1993 and 2016, although the analytic set was published between 2001 and 2011. The majority were non-U.S. settings, including Australia,⁵⁶² Hong Kong,^{558, 559, 561} Finland,⁵⁶⁰ and the Netherlands.⁵⁶⁴ One study compared case management versus consultation services in a nursing home.⁵⁶² The remaining studies were community-based. No particular research program in this literature set showed evolution over time, although the majority of studies were explanatory, or Stage 3 of the NIH Stage Model.

PLWD Outcomes

Evidence was insufficient to assess the effect of case management on PLWD outcomes. Table 8.5 provides the summary findings. All studies reported outcomes for PLWD.⁵⁵⁸⁻⁵⁶¹ Studies reported mixed results for rate of institutionalization⁵⁵⁸⁻⁵⁶⁰ and change in dementia symptoms.^{558, 559, 561} Groups showed no statistically significant difference for quality of life or depression.⁵⁶¹

Table 8.5. Summary of findings for PLWD outcomes: case management

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Institutionalization Case management vs usual care	2 RCTs ^{558, 560} (n=192) 18-24 months	PLWD dementia unspecified	1 of 2 found benefit 1 of 2 found no difference	Insufficient
NPI Case management vs usual care	2 RCT ^{558, 561} (n=194) 12-18 months	PLWD dementia unspecified	1 of 2 found benefit 1 of 2 found no difference	Insufficient
Quality of Life Case management vs usual care	1 RCT ⁵⁶¹ (n=102) 12 months	PLWD dementia unspecified	0 found benefit 1 found no difference	Insufficient
Depression Case management vs usual care	1 RCT ⁵⁶¹ (n=102) 12 months	PLWD dementia unspecified	0 found benefit 1 found no benefit	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: n=number; NPI=Neuropsychiatric Inventory; PLWD=person/people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Evidence was insufficient to assess the effect of case management on CG/P outcomes. Table 8.6 summarizes the primary outcomes for CG/P. Two studies reported outcomes for CG/P.^{558, 559, 561} The studies found mixed results for quality of life and caregiving burden. One study reported no statistically significant difference between groups for caregiver health status.⁵⁶¹

Table 8.6. Summary of findings for caregiver outcomes: case management

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Quality of Life Case management vs usual care	2 RCT ^{558, 561} (n=194) 12-18 months	CG/P	1 found benefit 1 found no difference	Insufficient
Caregiver Burden Case management vs usual care	2 RCT ^{558, 561} (n=194) 12-18 months	CG/P	1 found benefit 1 found no difference	Insufficient
Caregiver health status Case management vs usual care	1 RCT (n=102) 12 months	CG/P	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: CG/P=caregiver/care partner; n=number; RCT=randomized controlled trial

Variation in Outcomes

Studies in the analytic set did not examine variation in outcomes by PLWD, caregiver, or intervention characteristics.

Care Protocols for PLWD

Key Point

- Studies of care protocols for PLWD were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Interventions in this section targeted the health system level and aimed to improve quality of care through the implementation of protocols. Generally speaking, protocols are rules and procedures for providing care in a health system or in units within an organization. They also specify the tools and tasks to be carried out, and can therefore help formal caregiving staff, regardless of their length of service or experience, understand the processes. Protocols can be based on published guidelines or other sources for what are considered best practices. Staff training is an essential process for implementing protocols. Care protocols can be wide ranging, from decision trees to help formal caregivers systematize care decisions to protocols to improve nutrition care. Many protocols addressed agitation in nursing home settings.

Eligible Studies

We identified 17 unique studies from 25 publications that examined the impact of training and delivery of evidence-based guidelines/protocols targeted toward PLWD. (Table 8.7) We provide information on all studies in the evidence map in Appendix Tables F.⁵⁶⁸⁻⁵⁸³

Table 8.7. Basic characteristics of literature set: care protocols for PLWD

Characteristics	Information
Total unique studies	17 studies
Non-U.S. studies	11 studies
Evidence map studies	6 pilot or feasibility studies 11 high risk of bias
Analytic set studies	0

Intervention Research Context

Eligible studies on case management were published between 2006 and 2019. The majority were non-U.S. settings, including Australia,⁵⁷⁵ Spain,⁵⁸⁰ Germany,⁵⁷⁹ Norway,⁵⁸⁴ Taiwan,²⁶¹ and the Netherlands.^{568, 581, 582} Using the NIH Stage Model as a framework for assessment, this literature set comprised Stage 3 explanatory studies. A few studies built directly upon a prior publication, showing some evolution of the research over time. With the exception of the one nutrition protocol,⁵⁸⁰ the non-U.S. studies were published in the last 6 years and focused on preventing or addressing BPSD in nursing home settings. While also focused on nursing home settings, U.S.-based studies tended to be older publications, published between 2006 and 2012. However, one U.S.-based study tested a decision algorithm to help care managers create care plans to reduce burden and depression among CG/P in the community.⁵⁷⁶

Advance Care Planning

Key Point

- Evidence was insufficient to draw conclusions about the effect of decision support tools for advance care planning on PLWD and their caregivers.

Intervention Description

Advance care planning is a process that broadly involves individuals making healthcare decisions for themselves or for others about future healthcare needs. Within this section, studies focused on family members/caregivers participating in advance care planning as decisionmakers for PLWD unable to make their own decisions. Advance care planning interventions can include a range of tools such as decision support tools or treatment plans, and they usually include some form of discussion with healthcare professionals. Studies in our analytic set focused on the use of decision support tools for advance care planning.

Eligible Studies

We identified nine unique studies from 12 publications.⁵⁸⁵⁻⁵⁹⁴ Table 8.8 summarizes the characteristics of the literature set. Three studies were pilots^{587, 589, 593, 595} and an additional three were assessed as high risk of bias, therefore these six studies are not part of the analytic set.^{588, 590-592} We provide information about them in the evidence map in Appendix F.

One study examining the use of decision support tools in advance care planning was assessed as medium risk of bias and is the analytic set.^{585, 586, 596-598} This study enrolled PLWD with

advanced dementia and their surrogates, and it examined a brief video decision support tool focused on advance directives, goals of care, and burdensome treatments.^{585, 586} Appendix F provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 8.8. Basic characteristics of literature set: advance care planning

Characteristic	Information
Total Studies	9 studies
Non-U.S. studies	3 studies
Evidence map studies	5 pilot studies 3 high risk of bias studies
Analytic set studies	1 cluster randomized controlled trial
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	402
Dementia type/definition	Advanced dementia (type not specified)
Caregiver type (number)	Informal caregivers (402)

Abbreviation: PLWD=people living with dementia

Intervention Research Context

Eligible studies on advance care planning were published between 2011 and 2019 and were primarily conducted in the United States. One study took place in an adult-day care setting among African American family caregivers.⁵⁸⁷ The remaining studies took place in nursing homes. The fairly recent publication dates of this literature set and the prevalence of pilot studies may reflect increased interest over the last decade in person-centered care and decision support tools. Four of eight eligible studies examined the use of decision support tools.^{585, 586, 589-591} The other four studies examined advance care planning conversations between health professionals and CG/P or decision-makers incorporating education, treatment plans, and other tools.^{587, 588, 592, 593, 595}

PLWD Outcomes

Evidence was insufficient to assess the effect of advance care planning decision support tools on PLWD outcomes. Table 8.9 summarizes the primary outcomes for PLWD. One study (n=402) reported hospitalization rates at 12 months, finding them similar between the decision support and usual care groups.⁵⁸⁶

Table 8.9. Summary of findings for PLWD outcomes: advance care planning

Outcome Comparator	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Hospitalizations Advance care planning decision support tool vs usual care	1 cluster RCT ⁵⁸⁶ (n=402) 12 months	Advanced dementia	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Evidence was insufficient to assess the effect of advance care planning decision support tools on CG/P outcomes. Table 8.10 summarizes the primary outcomes for CG/P. One study (n=402) reported outcomes for whether CG/P completed an advance directive for no hospitalization on

behalf of PLWD, participated in a goals of care discussion, or stated a preference for comfort care (over basic or intensive care). Groups did not differ for any outcome at 6 or 12 months.⁵⁸⁶

Table 8.10. Summary of findings for caregiver outcomes: advance care planning

Outcome	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Documented Do Not Hospitalize Directive ACP decision support tool vs usual care	1 cluster RCT ⁵⁸⁶ (n=402) 6 months, 12 months	Advanced dementia	1 found no benefit.	Insufficient
Goal of care discussions ACP decision support tool vs usual care	1 cluster RCT ⁵⁸⁶ (n=402) 6 months, 12 months	Advanced dementia	1 found no benefit.	Insufficient
Proxies Preferring Comfort Care ACP decision support tool vs usual care	1 cluster RCT ⁵⁸⁶ (n=402) 6 months, 12 months	Advanced dementia	1 found no benefit.	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: ACP=advance care planning; n=number; RCT=randomized controlled trial

Variation in Outcomes

The analytic set study did not examine variation in outcomes by PLWD, caregiver, or intervention characteristics.

Palliative Care

Key Point

- Studies of palliative care for PLWD were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Palliative care interventions aim improve symptom relief, distress, hospitalization burden, and comfort with death among PLWD and their caregivers.

Eligible Studies

We identified five unique studies from six publications that examined the use of palliative care interventions among PLWD.⁵⁹⁹⁻⁶⁰⁴ (Table 8.11) All studies were either pilots or assessed as high risk of bias. We present information on all studies in the evidence map in Appendix F.

Table 8.11. Basic characteristics of literature set: palliative care

Characteristics	Information
Total unique studies	5 studies
Non-U.S. studies	3 studies
Evidence map studies	1 pilot study 4 high risk of bias studies
Analytic set studies	0

Intervention Research Context

Eligible studies on palliative care were published between 2000 and 2019. The three non-U.S. studies were set in nursing homes.^{599, 600, 602, 604} One examined case conferencing in Australia,^{599, 600} one examined a multicomponent intervention that included training, communication, routine palliative care tasks, and a nurse facilitator in Canada,⁶⁰⁴ and one compared general versus personalized feedback for end-of-life care in the Netherlands.⁶⁰² The U.S.-based studies examined palliative care triggered by acute hospital stays for PLWD.^{601, 603}

Other Service Provision Interventions

Key Point

- Evidence was insufficient to draw conclusions about the effect of other service provision interventions on PLWD or their caregivers.

Intervention Description

This set of diverse interventions included approaches such as decision aids, technology to assist CG/P, and a tool-kit to assist CG/P in improving the safety of the home for PLWD.

Eligible Studies

We identified five unique studies from seven publications.^{596-598, 605-608} Table 8.12 summarizes the characteristics of the literature set. Two studies were a pilot^{605, 608} and two were assessed as high risk of bias;^{606, 607} these studies were therefore excluded from the analytic set. We provide information about them in the evidence map in Appendix F.

One study was assessed as medium risk of bias and was classified as an NIH Stage 3 explanatory study. This study enrolled PLWD with advanced dementia and feeding issues, along with their surrogates, to compare a print decision aid for feeding options versus usual care.⁵⁹⁶⁻⁵⁹⁸ Appendix F provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 8.12. Basic characteristics of literature set: other service provision interventions

Characteristic	Information
Total Studies	5 studies
Non-U.S. studies	2 studies
Evidence map studies	2 pilot studies 2 high risk of bias studies
Analytic set studies	1 cluster randomized controlled trial
Risk of bias of analytic set	Medium
Number of PLWD in analytic set	256
Dementia type/definition	Advanced dementia (type not specified)
Caregiver type (number)	Informal caregivers (n=256)

Abbreviations: n=number; PLWD=people living with dementia

Intervention Research Context

All studies were unrelated to each other and U.S.-based. Two studies examined video-based telehealth technology to support community-based CG/P.^{605, 607} Another study examined a self-directed educational program and home safety tool-kit to improve home safety for PLWD.⁶⁰⁶ Another examined a decision aid to help CG/P make decisions about feeding care for their

PLWD with advanced dementia and swallowing difficulties.⁵⁹⁶ A fourth study examined including PLWD living in nursing homes in care planning meetings.⁶⁰⁸

PLWD Outcomes

Evidence was insufficient to address PLWD outcomes for a decision aid for PLWD feeding. Table 8.13 provides a summary of the findings. At 3 months, the decision aid study (n=256) found a statistically significant benefit with the intervention in the number of PLWD receiving a specialized dysphagia diet.⁵⁹⁶ However, groups did not differ for other types of feeding interventions (e.g., specialized utensils, specialized staff assistance, high-calorie diet).⁵⁹⁶

Table 8.13. Summary of findings for PLWD outcomes: other service provision interventions

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Assisted Feeding Intervention-Specialized Dysphagia Diet Decision aid vs usual care	1 cluster RCT ⁵⁹⁶ (n=256) 3 months	Advance dementia with feeding issues	1 found benefit 0 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial

Caregiver Outcomes

Evidence was insufficient to address CG/P outcomes for a decision aid for PLWD feeding. Table 8.14 provides a summary of the findings. One study (n=256) reported outcomes for CG/P decisional conflict, satisfaction with decisions, decisional regret, and feeding discussions with nursing home staff. At 3 months, CG/P who reviewed the decision support tool reported less decisional conflict than the usual care group.⁵⁹⁶ Informal caregivers who reviewed the decision support tool also reported more conversations about feeding issues with doctors, nurse practitioners, and physician's assistants than usual care.⁵⁹⁶ However, the groups did not differ in number of conversations with other nursing home staff.⁵⁹⁶ Nor did groups differ in measures of decision satisfaction and regret.⁵⁹⁶

Table 8.14. Summary of findings for caregiver outcomes: other service provision interventions

Outcome Comparison	# Studies/Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Decisional conflict Decision aid vs usual care	1 cluster RCT ⁵⁹⁶ (n=256) 3 months	Advance dementia with feeding issues	1 found benefit 0 found no difference	Insufficient
Satisfaction with Decisions Decision aid vs usual care	1 cluster RCT ⁵⁹⁶ (n=256) 3 months	Advance dementia with feeding issues	0 found benefit 1 found no difference	Insufficient
Decisional Regret Decision aid vs usual care	1 cluster RCT ⁵⁹⁶ (n=256) 3 months	Advance dementia with feeding issues	0 found benefit 1 found no difference	Insufficient
Feeding Discussions Decision aid vs usual care	1 cluster RCT ⁵⁹⁶ (n=256) 3 months	Advance dementia with feeding issues	1 found benefit for discussions with doctors, PA, or NP. No benefit for other nursing home staff	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: n=number; NP=nurse practitioner; PA=physician assistant; RCT=randomized controlled trial

Care Delivery Models and Programs

We use the term “model” in a generic manner to refer to developed conceptual approaches. We use the term “program” for more clearly identified sets of components and/or protocols. To further distinguish between interventions presented in Chapters 4 – 7 and the care delivery interventions in this chapter, we use the term intervention in a slightly different manner than in previous chapters. Here, we reserve the term intervention only for discretely identified interventions that, if supported by research, could be incorporated into an evidence-based program or model. Person-centered or individualized programs are interventions that are evidence-based, manualized or algorithmic, and ideally built from studies of individual components that may be incorporated into an individualized plan or approach. This differs from the many tailored interventions presented in Chapters 4 – 7, which rely on care staff to use their knowledge and experience to adapt the intervention to the individual receiving care.

Collaborative Care and Care Coordination Models

Key Points

- Collaborative care models (i.e. Care Ecosystems or discrete adaptations of the ACCESS models) may improve PLWD quality of life. (low-strength evidence) This improvement may be very small to small, or it may be larger but concentrated in some not yet identified subgroup of people.
- Collaborative care models (i.e. discrete adaptations of the ACCESS model) may improve system-level markers, including guideline-based quality indicators and reduction in emergency department visits. (low-strength evidence)
- Evidence was insufficient to draw conclusions about all other outcomes for both PLWD and CG/P.

Intervention Description

Collaborative care models use multidisciplinary teams to integrate medical and psychosocial approaches to healthcare for PLWD. Care coordination functions are usually assigned to specific care coordinators. Team members may be co-located or spread across locations with a designated hub. Since care coordination is central to collaborative care models, we include care coordination studies in this section. Collaborative care is organized at the healthcare system level and may be provided through services either in the community or in residential facilities. As such, these approaches try to leverage local care and support resources. Most are aimed at providing CG/P support, along with coordinating care for PLWD. Often, studies required care coordinators or navigators to have a minimum of one contact per month with CG/P, or with PLWD directly. However, contact could be more frequent if necessary for caregivers to accomplish the care plans. Contact took place by phone or internet except in one German study that used in-home visits. Studies ran from 6 months to almost 2 years, but most outcomes were measured at 12 months.

Eligible Studies

We identified 13 unique studies from 32 publications that examined collaborative care. (Table 8.15) We provide information on all pilot and high risk of bias studies as part of the

evidence map in Appendix F.⁶⁰⁹⁻⁶¹⁴ The analytic set includes seven low to medium bias of bias studies and represents care for 2,597 PLWD.⁶¹⁵⁻⁶⁴⁰ Given the pragmatic nature of most of the studies, information on dementia type and severity is less available. Appendix F provides evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 8.15. Basic characteristics of literature set: collaborative care models

Characteristics	Information
Total unique studies	13 studies
Non-U.S. studies	4 studies
Evidence map studies	4 pilot studies 2 high risk of bias studies
Analytic set studies	5 pragmatic, 2 explanatory studies
Risk of bias of analytic set	6 medium
Number of PLWD in analytic set	2,641
Dementia type/definition	Generally dementia unspecified, usually more mild to moderate
Caregiver type (number)	Informal caregivers (2,405)

Abbreviation: PLWD=people living with dementia

Intervention Research Context

This literature falls mainly into Stage 4 of the NIH Stage Model, or mostly pragmatic trial designs, although we did identify a few pilot studies. The pragmatic studies either cited previous evidence-based research or related pilot studies, but we identified no literature showing any collaborative care approach as progressing through all stages of the model. The pragmatic trials appeared along a spectrum of balanced explanatory/pragmatic to fully pragmatic. Most studies were conducted in high-resource locations, including the United States, Germany, Singapore, and the Netherlands. Few studies investigated diverse caregivers. However, one study enrolled 50 percent urban African-American caregivers,⁶¹⁶ and one study used the ACCESS protocol for Latino/immigrant populations in an underserved urban setting.⁶¹⁷ Two high risk of bias studies of multidisciplinary teams, reported in the evidence map, were conducted in nursing homes. One was a relatively large pragmatic trial of 793 PLWD, but only 40 percent of participants received the intervention, leaving it difficult to understand what drove the lack of difference between the intervention and the control groups.⁶¹³

PLWD Outcomes

Six of the seven medium to low risk of bias studies reported outcomes related to PLWD. We synthesized results qualitatively because differences in outcome measures, time to outcomes, and analytic methods prohibited quantitative pooling. The most commonly investigated outcome was quality of life. We also include quality indicators derived from guideline recommendations as PLWD-related outcomes. Table 8.16 summarizes the number of studies investigating each outcome and the number of studies that found a statistically significant benefit or no difference in outcomes for PLWD in the intervention group versus the comparison group.

Quality of life improvement was supported by low-strength evidence from four studies using usual care controls. One large pragmatic study found statistically significant but very small to small benefit using the QoL-AD.⁶¹⁸ Another large pragmatic trial found benefit greater than the established minimally important difference for the health utility index, which indicates the improvement may have been large enough for the study participants to notice the change.⁶²⁰ Two other trials found no difference for QoL-AD⁶¹⁹ and the health utility index.⁶¹⁷ However, when

assessing the strength of evidence, we gave less weight to the findings from these studies due to the smaller sample sizes and other concerns with the statistical analyses they used.

Quality indicators were supported by low-strength evidence drawn from two studies. These indicators were based on adherence to 19 to 23 dementia guideline recommendations in four domains: assessment, treatment, education and support, and safety. Both studies found benefit of about the same magnitude across the quality indicators.^{617, 620}

Low-strength evidence also showed a decrease in rate of emergency department visits over a 12-month period.⁶¹⁸ Five PLWD needed to participate in collaborative care to achieve a decrease of one emergency room visit.

Evidence was insufficient for the remaining outcomes. Some outcomes were only investigated by a single small study, making the consistency in findings unknown. Also, the uncertainty regarding a true “no difference between groups” finding versus an “inability to show a difference” finding remained too high. Two studies investigated neuropsychiatric symptoms, but the outcome was assessed as high risk of bias in one, which was therefore not included. One explanatory study found benefit for the neuropsychiatric symptom inventory.⁶¹⁶ Groups did not differ significantly for daily activities of living,^{616, 619} depression as measured by the CSDD,⁶¹⁶ or hospitalization or ambulance.⁶¹⁸ Nursing home placement had mixed results.^{616, 619}

Table 8.16. Summary of findings for PLWD outcomes: collaborative care models

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Findings	Strength of Evidence*
Quality of Life Collaborative care vs usual care	4 Pragmatic trials ⁶¹⁷⁻⁶²⁰ (n=1,746) 6-18 months	2 of 4 found benefit 2 of 4 found no difference	Low for benefit (Weighted to larger pragmatic trials)
Neuropsychiatric symptoms Collaborative care vs usual care	1 Explanatory cluster trial ⁶¹⁶ (n=152) 12 months	1 found benefit 0 found no difference	Insufficient
Function Collaborative care vs usual care	1 Pragmatic trial ⁶¹⁹ 1 Explanatory cluster trial ⁶¹⁶ (n=560) 6-12 months	0 found benefit 2 found no difference	Insufficient
Depression Collaborative care vs usual care	1 Explanatory cluster trial ⁶¹⁶ (n=152) 12 months	0 found benefit 1 found no difference	Insufficient
Quality indicators Collaborative care vs usual care	2 Pragmatic trials ^{617, 620} (n=559) 6-18 months	2 of 2 found benefit 0 of 2 found no difference	Low for benefit (Weighted to larger pragmatic trials)
Emergency room visits Collaborative care vs usual care	1 Pragmatic trial ⁶¹⁸ (n=780) 12 months	1 found benefit 0 found no difference (no difference for hospitalization or ambulance use)	Low for benefit (Weighted to larger pragmatic trials)
Nursing home placement Collaborative care vs usual care	1 Pragmatic trial ⁶¹⁹ 1 Explanatory cluster trial ⁶¹⁶ 1 RCT ⁶⁴⁰ (n=794) 6-18 months	0 found benefit 2 found no difference 1 found benefit at 1.6 years but not at 2 years	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: n=number; RCT=randomized controlled trial

Caregiver Outcomes

Six of the seven medium- to low-risk-of-bias studies reported CG/P outcomes. Again, we synthesized results qualitatively because differences in outcome measures, time to outcomes, and analytic methods prohibited quantitative pooling. The most commonly investigated outcomes were caregiver burden and depression. Table 8.17 summarizes the number of studies investigating each outcome and the number of studies that found a statistically significant benefit or no difference in outcomes for CG/P.

Evidence was insufficient to draw conclusions about collaborative care versus usual care for CG/P. The inability to combine outcomes from multiple studies made it impossible to show precision in any outcome. Several outcomes were only investigated by a single study, making the consistency of findings unknown. The uncertainty regarding a true “no difference between groups” finding versus an “inability to show a difference” finding remained too high. One study found no statistical difference between groups for quality of life as measured by the EuroQual-5D.⁶²⁰ Caregiver burden was variably measured with mixed findings. One large pragmatic trial found very small benefit,⁶¹⁸ while the other study reported mixed findings.^{617, 619, 620} Caregiver depression was slightly reduced in one large study,⁶¹⁸ but two other studies found no difference.⁶¹⁵⁻⁶¹⁷ One trial reported no difference between groups for caregiver self-efficacy.⁶¹⁸ Another trial found mixed results in quality measures.⁶¹⁵

Table 8.17. Summary of findings for caregiver outcomes: collaborative care models

Outcome Comparison	# Studies/Design (n analyzed) Timing	Findings	Strength of Evidence*
Quality of Life Collaborative care vs usual care	1 Pragmatic trial ⁶²⁰ (n=408) 18 months	0 found benefit 1 found no difference	Insufficient
Caregiver burden Collaborative care vs usual care	4 Pragmatic trials ^{615, 617-620} (n=1,719) 6-18 months	2 of 4 found benefit 2 of 4 found no difference (also no difference in several related measures such as role captivity or physical health strain)	Insufficient
Depression Collaborative care vs usual care	3 Pragmatic trials ^{615, 617, 618} 1 Explanatory cluster trial ⁶¹⁶ (n=1,570) 6-18 months	1 of 4 found benefit 3 of 4 found no difference	Insufficient
Self-efficacy Collaborative care vs usual care	1 Pragmatic trial ⁶¹⁸ (n=780) 12 months	0 found benefit 1 found no difference	Insufficient
Quality measures Collaborative care vs usual care	1 Pragmatic trial ⁶¹⁵ (n=486) 12 months	1 found benefit in unmet needs but no difference in use of support services or informal helpers	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviation: n=number

Variation in Outcomes

By PLWD Characteristics

One pragmatic study, the ACCESS study, enrolled primarily white urban caregivers and found variation in outcomes by caregiver education. Those who had not graduated from high school showed greater improvement than college graduates in the quality indicators (44.4 vs 29.5

for the assessment dimension, 36.9 vs. 15.7 for the treatment dimension, and 52.7 vs. 40.9 for the safety dimension [$P < .001$ for all three]).⁶³²

The ACCESS protocol was also later tested with a more intensive in-person protocol in an urban Latino population with a large immigrant population. This study also found improvement in quality indicators, further supporting the ACCESS trial. However, they were unable to confirm that more intensive in-person delivery further improved scores. Since the adapted model was also tested with a different population, we could not separate the potential effects of the new model from those of the new population.

By Caregiver Characteristics

The VA Partners in Dementia Care program reported that benefits were larger among CG/P of PLWD with higher baseline cognitive impairment (unmet needs: $B=-0.97$, $p<0.001$), caregivers with higher baseline conflict with caregiving (role captivity: $B=-0.23$, $p=0.02$), and caregivers of PLWD with higher baseline behavior problems (physical health strain: $B=-0.09$, $p=0.06$; number informal helpers: $B=0.61$, $p=0.005$).⁶¹⁵

By Intervention Characteristics

The ACCESS study also reported increased benefit with increased contact between CG/P and providers. For every additional monthly encounter between a health organization care manager and a caregiver, the mean percentage of quality care indicators rose between 10 and 16 percentage points across four quality domains.⁶³¹

Nursing Home–Based Interdisciplinary Individualized Care/Person-Centered Care and Dementia Care Mapping

Intervention Description

Similar to Collaborative Care models for community-dwelling PLWD and their CG/P, models and programs to improve care have also been used in nursing home settings. Collaborative Care models incorporate working with CG/P, which means care is personalized to the informal caregivers' PLWD. Programs for nursing home settings focus more on providing individualized care through training formal caregivers to foster PLWD personhood and attending to unmet needs through individualized care plans. These programs support person-centered care structurally through training, embedding caregivers in care teams, and/or providing tools to support systematic observation of factors that enhance person-centered care.

This form of individualized care for nursing homes has been explored predominantly by non-U.S. countries. Only one recent small pilot⁶⁴² and one study from 1997 were based in the United States; these studies examined interdisciplinary team approaches to provide individualized care.⁶⁴³ The remaining locations in which these programs were investigated include the United Kingdom,⁶⁴⁴⁻⁶⁴⁷ Australia,⁶⁴⁸⁻⁶⁵⁰ Norway,^{641, 651} Germany,⁶⁵² and the Netherlands.⁶⁵³ Because the programs have been tested in non-U.S. long-term care systems, we could not determine their adaptability to U.S. healthcare and support systems. Interpreting the usefulness of the findings would be challenging. For this reason, we did not further analyze these studies, but we briefly describe them in the Intervention Research Context subsection without performing further analysis.

Eligible Studies

We identified 16 unique studies from 22 publications that examined individualized person-centered care in nursing homes. (Table 8.18) We provide information on all pilot and high risk of bias studies as part of the evidence map in Appendix F.^{608, 642, 644, 645, 648, 649, 651, 652, 654, 655} Five studies were rated as low to medium risk of bias.^{641, 643, 647, 650, 653, 656-659} These studies enrolled 1,722 PLWD with generally moderate to severe dementia. Appendix F provides evidence tables and summary risk of bias assessments.

Table 8.18. Basic characteristics of literature set: individualized person-centered care (non-U.S.)

Characteristics	Information
Total unique studies	16 studies
Non-U.S. studies	13 studies
Evidence map studies	4 pilot studies high risk of bias studies
Analytic set studies	1 pragmatic, 2 balanced, and 2 explanatory studies
Risk of bias of analytic set	5 medium

Intervention Research Context

With the exception of one study from 1996, this literature is relatively recent, with the eight studies published within the last 5 years. The predominant research design involved assigning care facilities to treatment or control study arms. Using the NIH Stage Model, these studies were assessed as Stage 4 and more pragmatic in study design. Problems with ensuring the intervention was actually delivered to the PLWD residents was a common cause of high risk of bias. Study authors were frequently very transparent regarding reporting fidelity to the interventions.

The largest set of literature examined person-centered care based on Kitwood’s framework, generally coupled with Dementia Care Mapping™ (DCM™).^{644, 653} DCM™ is an intervention that promotes care practice development. It involves training formal caregivers to apply a systematic approach to observing factors associated with PLWD well-being, and then share the information with staff to support care planning. DCM™ is a trade-marked tool that has been used in several countries including Europe, the United Kingdom, and Australia. Over 60 percent of published trials of DCM™ are in the evidence map as pilots or high risk of bias studies, and results were mixed.^{649, 651, 652, 654} Recently, two large pragmatic trials both found no benefit for DCM™, but these trials also reported challenges with ensuring that the intervention was fully and widely implemented.^{644, 653}

A few studies in the analytic set used a different structural approach and supported person-centered care within interdisciplinary teams.^{641, 647} These larger studies, published in 2018, stemmed from prior work presented in the evidence map, and showed indications that the research was developing along a distinct line of inquiry. These studies noted small improvement in PLWD quality of life and reduction in agitation over 8 weeks to 9 months. However, these program approaches will require further investigation into their sustainability.

Care Staff Education and Support Needs

As noted above, specialized service provision requires qualified staff members to be regularly available to perform the services offered to PLWD or their CG/P. Interventions addressing staff education and support needs include approaches targeted at helping informal and formal caregivers work together to support PLWD well-being.

Caregiver Staff Training

We considered training programs as being at the pilot stage if the training was not embedded in the organizational structure, such that the knowledge or skills developed by the training will sustain across staff turnover.

Key Point

- Studies of training interventions to improve formal caregiver staff knowledge and skills were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Formal staff training is intended to improve staff knowledge and facilitate skill-building.

Eligible Studies

We identified 22 unique studies from 23 publications that examined the use of training interventions for formal caregiver staff.^{214, 660-680} (Table 8.19) No studies were assessed as low or moderate risk of bias, while two were assessed as high risk of bias. Remaining studies were all pilot or small sample studies. We provide information on all studies in the evidence map in Appendix F.

Table 8.19. Basic characteristics of literature set: formal caregiver staff training

Characteristics	Information
Total unique studies	22 studies
Non-U.S. studies	12 studies
Evidence map studies	12 pilot studies 3 small sample studies 4 training pilot studies (without evidence of sustained training) 3 high risk of bias studies
Analytic set studies	0

Intervention Research Context

Training topics in the evidence map included general education about dementia and dementia care,^{668, 669, 678, 680} communication techniques,^{660, 663, 671, 672, 679} bathing techniques,^{667, 676} identifying and addressing pain,^{666, 680} morning care,⁶⁷⁷ integrating physical activity into daily routines,⁶⁶⁵ sleep,⁶⁷³ and feeding skills.⁶⁶¹ Several used activities or taught skills targeted at improving behavioral issues,^{664, 674} including apathy.⁶⁷⁰ One intervention examined training to identify signs of awareness in PLWD with severe dementia.⁶⁶² While nursing homes or other long-term care facilities were the most common settings, one study examined education on dementia for acute care hospital staff,⁶⁷⁵ and another study examined education for community-based care staff.⁶⁶³ One examined an online training portal on dementia care available to both formal and CG/P.⁶⁶⁸ Non-U.S. locations included Australia, Canada, France, Germany, Taiwan, the United Kingdom, and the European Union.

The research is largely preliminary across a varied set of training interventions. Most interventions stopped at the pilot stage or did not embed the training intervention into the care organization in order that new skills and knowledge would be sustained over time even with staff turnover. Research activity grew between 1999 and 2017, with new unrelated pilot studies scattered throughout the whole period and across locations. We found, however, one small indication of research progressing through the NIH Stage Model. One training intervention

(Bathing Without a Battle) published both a small sample initial study⁶⁷⁶ and a later larger multisite study that enrolled PLWD with a broader range of dementia severity.⁶⁶⁷

Informal Caregiver Staff Training

Key Point

- Studies of training interventions to improve CG/Ps’ knowledge and skills were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Training interventions for CG/P differ from interventions in Chapter 6 because they do not incorporate any psychoeducation or therapy to support CG/P’ well-being. They consist, instead, of dementia education and training focused on skill-building with an emphasis on role-training.

Eligible Studies

We identified 12 unique studies from 12 publications that examined training interventions to improve CG/P skills.^{668, 681-691} (Table 8.20) Three pilot studies and five small sample studies were excluded from the analytic study set. Four additional studies were assessed as high risk of bias. We provide information on all pilot studies and high risk of bias studies as part of the evidence map in Appendix F.

Table 8.20. Basic characteristics of literature set: CG/P training

Characteristics	Information
Total unique studies	12 studies
Non-U.S. studies	6 studies
Evidence map studies	3 pilot studies 5 small sample studies 4 high risk of bias studies
Analytic set studies	0

Abbreviation: CG/P=caregiver/care partner

Intervention Research Context

The research is largely preliminary and includes studies examining a varied set of training interventions. Most interventions in the evidence map stopped at the small sample or pilot stage. Research spanned over 20 years, from 1994 to 2015. Studies conducted in the United States were older, from 1994 to 2007, and mostly focused on the role of caregiving, building general dementia caregiver skills, or skills for behavioral management.^{681, 684-686, 689, 691} Non-U.S.-based studies, conversely, were published from 2001 to 2015 and were balanced across general care^{683, 690} and communication-specific skills.^{682, 687, 688} Studies also examined portable training materials in the United States⁶⁸⁴ and internet-accessible training materials in the European Union.⁶⁶⁸

Family Education and Partnering

Key Point

- Studies of training interventions to improve formal caregiver staff knowledge and skills were described in the evidence map but not considered for analysis due to limitations in study designs.

Intervention Description

Family involvement interventions provide education, training, and other supports to establishing a collaborative relationship between family members and formal caregivers of PLWD in nursing homes. These interventions seek to improve PLWD well-being through: 1) improving the partnership between the staff and family members to identify unmet needs among PLWD, and 2) helping family members structure visits to avoid triggering behavioral symptoms in PLWD. Training in communication techniques is prominent.

Eligible Studies

We identified five unique studies from five publications that examined family education and partnership interventions.^{76, 692-695} (Table 8.21) Three studies were pilots and two were assessed as high risk of bias, thus none were included in the analytic set. We provide information on all studies as part of the evidence map in Appendix F.

Table 8.21. Basic characteristics of literature set: family education and partnering

Characteristics	Information
Total unique studies	5 studies
Non-U.S. studies	1 study (Australia)
Evidence map studies	3 pilot studies 2 high risk of bias studies
Analytic set studies	0

Intervention Research Context

The research is preliminary, comprising pilot or explanatory studies with design or conduct challenges. Studies conducted in the United States were published from 1999 to 2007. Family Involvement in Care was examined in several studies,⁶⁹²⁻⁶⁹⁴ including a 2011 study in Australia.⁷⁶ The 1999 study was a foundation for the Family Involvement in Care program.⁶⁹⁴ The Partners in Caregiving adaptation for Special Care Units in the 2007 publication added concurrent staff and family training, thereby broadening the intervention focus beyond primarily family members.⁶⁹⁵ PLWD outcomes were not prominent in this literature set; outcomes primarily applied to family members and staff.

Multitier Training

Multitier training interventions involve formal caregivers being trained to provide training to CG/P for specific care needs of the PLWD.

Key Point

- Evidence was insufficient to draw conclusions about the effect of multi-tier training on PLWD and their caregiver.

Intervention Description

This literature set comprises mostly practice guideline-based interventions encompassing multidisciplinary care interventions. These interventions aimed to improve communication between PLWD and their caregivers, and used nutrition and activity planning to help improve various aspects (psychosocial, physical, and behavioral) related to better functioning among PLWD and their caregivers.

Eligible Studies

Table 8.22 summarizes the characteristics of the literature set. We identified seven unique studies from nine publications that examined the use of training plus delivery of evidence-based guidelines/protocols for caregivers, with the goal in improving their own health outcomes as well as the health outcomes of PLWD.⁶⁹⁶⁻⁷⁰² One study was assessed as high risk of bias⁶⁹⁷ and five were pilots.^{696, 698, 700-702} Only one study was included in the analytic set.⁶⁹⁹ We provide information on the high risk of bias studies in the evidence map in Appendix F.

The included study (n=95) was a randomized controlled trial targeted toward family (informal) caregivers. The trial compared STAR-C (interactive sessions between community consultants and caregivers) with routine medical care.⁶⁹⁹ It was assessed as medium risk of bias and categorized as explanatory. PLWD were eligible if they were community dwellers living with their caregivers and had a probable diagnosis of AD (with moderate cognitive impairment). Family caregivers were also enrolled in the study. Appendix F provide evidence tables, summary risk of bias assessments, and strength of evidence for key comparisons and outcomes.

Table 8.22. Basic characteristics of literature set: multitier training

Characteristics	Information
Total unique studies	7 studies
Non-U.S. studies	2 studies
Evidence map studies	5 pilot studies 1 high risk of bias studies
Analytic set studies	1 randomized controlled trial
Risk of bias of analytic set	1 medium
Number of PLWD in analytic set	95
Dementia type/definition	Moderate cognitive impaired probable Alzheimer's disease
Caregiver type (number)	Informal caregivers (95)

Abbreviation: PLWD=people living with dementia

Intervention Research Context

The literature on multitier training interventions comprises mostly pilot and high risk of bias studies. Four studies were conducted in United States^{697, 699, 701, 702} while the others occurred in the United Kingdom^{696, 700} and Finland.⁶⁹⁸ We identified a manual-based intervention targeted at improving caregiver's understanding of sleep and dementia.⁶⁹⁶ We also identified a study that provided caregivers with instruction on how to recognize pain among PLWD.⁶⁹⁷ Also evaluated was an intervention that used tailored nutritional guidance on the basis of the food diaries to prevent weight gain among PLWD.⁶⁹⁸

PLWD Outcomes

Evidence was insufficient to draw conclusions on quality of life and memory or behavioral problems of PLWD using the STAR-C intervention versus routine medical care (n=95).⁶⁹⁹ Table 8.23 summarizes the primary findings.

Table 8.23. Summary of findings for PLWD outcomes: multitier training

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
RMB-PC STAR-C vs RMC	1 RCT ⁶⁹⁹ (n=95) 2 months	Probable Alzheimer's disease	0 found benefit 1 found no difference	Insufficient

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
Quality of life STAR-C vs RMC	1 RCT ⁶⁹⁹ (n=95) 2 months	Probable Alzheimer's disease	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: n=number; PLWD=person/people living with dementia; RCT=randomized controlled trial; RMB-PC=Revised Memory and Behavior Problem Checklist; RMC=routine medical care

Caregiver Outcomes

Evidence was insufficient to draw conclusions about using the STAR-C intervention versus RMC (n=95) training plus delivery of evidence-based guidelines/protocols targeted toward caregivers. (Table 8.24)

Table 8.24. Summary of findings for caregiver outcomes: multitier training

Outcome Comparison	# Studies/ Design (n analyzed) Timing	Population	Findings	Strength of Evidence*
CES-D STAR-C vs RMC	1 RCT (n=95) 2 months	Primary family caregivers	1 found benefit 0 found no difference	Insufficient
HDRS STAR-C vs RMC	1 RCT (n=95) 2 months	Primary family caregivers	0 found benefit 1 found no difference	Insufficient
Caregiver burden STAR-C vs RMC	1 RCT (n=95) 2 months	Primary family caregivers	1 found benefit 0 found no difference	Insufficient
Caregiver reaction STAR-C vs RMC	1 RCT (n=95) 2 months	Primary family caregivers	1 found benefit 0 found no difference	Insufficient
Caregiver sleep questionnaire STAR-C vs RMC	1 RCT (n=95) 2 months	Primary family caregivers	0 found benefit 1 found no difference	Insufficient

*Insufficient ratings due to study limitations and imprecision in the findings.

Abbreviations: CES-D=Center for Epidemiological Studies Depression Inventory; HDRS=Hamilton Depression Rating scale; n=number; RCT=randomized controlled trial; RMC=routine medical care; STAR-C=staff training in assisted living residences-caregivers

Conclusion

We found 123 unique studies that investigated 11 care delivery interventions to improve how care is delivered. We grouped the 11 intervention categories into three main themes of care service delivery, care delivery models and programs, and care staff education and support needs. These care delivery interventions conform well to the framework for care interventions from the NASEM Families Caring for an Aging America 2016 report (Figure 1.1 in Chapter 1), delivering the interventions at the system level.

We found collaborative care models (i.e. discrete adaptations of the ACCESS⁶²⁰ or Care Ecosystem⁶¹⁸) may improve PLWD quality of life. It is difficult to estimate an effect size from a qualitative synthesis of study results. The largest, rigorously designed study found a statistically significant improvement in quality of life, but the effect size was very small. Based on reported data, we could not determine whether the average effect was broadly distributed across the full study population, but too small of a benefit to be noticeable by PLWD, or if the benefits were largely concentrated in some not yet identified subgroup of PLWD for whom the improvement would be noticeable. The other pragmatic trial also found improvement in health-related quality of life, with the average change being greater than what is considered a minimally detectable difference. These two studies may represent the range of possible effect sizes.

We also found collaborative care models may improve system-level markers, including guideline-based quality indicators and reduction in emergency department visits. Cost was not presented as an outcome. Cost data for the interventions were provided, but a cost-effectiveness analysis is outside the scope of this review.

For the remainder of the PLWD and CG/P outcomes for collaborative care, and for all other care delivery interventions, we found the uncertainty of the evidence was too high to draw conclusions. However, our being unable to draw a conclusion does not mean that the intervention has no effect. Research on interventions to change behavioral and psychological outcomes is challenging, and many factors can influence the outcomes. Future research may reduce uncertainty enough to allow for conclusions about the effect of these interventions, including potential benefits.

Chapter 9. Implementation of Care Interventions

This chapter addresses Guiding Question 1 on providing context for implementation and dissemination of care interventions. We present the results of the grey literature search conducted to provide resources for care interventions which may not have been empirically studied using study designs required by the review inclusion criteria. Because so few interventions were assessed to have at least low-strength evidence to support our review findings, we were concerned that providing implementation and dissemination information based only on those studies would leave readers with an unbalanced or skewed view of implementation and dissemination. Therefore, we do not report on this evidence here.

Grey Literature Search

We searched 15 different sources from the grey literature (i.e., research or other written material produced outside of traditional academic publishing) during April, 2019 to identify repositories of care interventions and criteria for evaluating and/or categorizing care interventions for people living with dementia (PLWD) and their caregivers. Most of these sources provided educational materials, webinars, and information on research and policy. However, three housed lists or libraries of interventions: the Benjamin Rose Institute on Aging, Family Caregiver Alliance, and Rosalynn Carter Institute for Caregiving. Subsequently, the Benjamin Rose Institute on Aging, in partnership with the Family Caregiver Alliance and the Gerontological Society of America, has formally launched their online resource for caregiving for dementia, the Best Practice Caregiving resource (see <https://www.benrose.org/best-practice-caregiving>). The Rosalynn Carter Institute has since discontinued their online resource.

The Best Practice Caregiving resource, a free online searchable database, provides information about programs for informal caregivers. The standard of evidence used by this resource differs substantially from that of this review. To be listed in the Best Practice database, a program needed to have at least one published statistically significant benefit for a caregiver outcome from a study of any empirical research design, plus have been implemented in at least one organization's regular service portfolio. The published research needed to be U.S.-based and have at least 50 percent of the informal caregivers providing care to PLWD.

The Family Caregiver Alliance provides a list of interventions (formerly their "Innovations Clearinghouse") for family caregivers of people with chronic disabling conditions. Housed under an umbrella of "program development," this list groups interventions into three categories: evidence-based practices, emerging practices, and model programs. The list is partially searchable, and the options under the program development umbrella can be narrowed by caregiver role or specific topics. Again, the standard of evidence used by this resource differs substantially from that of this review. The website does not provide uniform criteria to evaluate implementation readiness of level of evidence. Practices categorized as evidence-based are published in the literature after 1990, and must provide credible evidence for improved caregiver outcomes. Additionally, included publications must provide adequate information on methodology to allow for replication (see <https://www.caregiver.org/evidence-based-practices>). Emerging practices must use innovative methods and/or focus on underserved populations in diverse settings and populations (see <https://www.caregiver.org/emerging-practices>). Model programs are selected based on expert input, and must have been proven effective, replicated, or adapted, and provide training materials (see <https://www.caregiver.org/model-programs>). Both

emerging practices and model programs must be from a credible source and remain available for the foreseeable future.

A fourth source, the Administration for Community Living (ACL), through its National Alzheimer's and Dementia Resource Center (NADRC) (see <https://nadrc.acl.gov/node/140>) provides a report compendium listing dementia specific evidence-based and evidence-informed interventions that have been implemented through its grant programs. The interventions meet the ACL criteria and have been implemented by Alzheimer's Disease Supportive Services Program (ADSSP), Alzheimer's Disease Initiative Specialized Supportive Services (ADI-SSS) and Alzheimer's Disease Program Initiative (ADPI) grantees from 2007 to 2018.

Chapter 10. Discussion

Overview

This systematic review sought to assess the evidence base for effective care interventions for people living with dementia (PLWD) and their caregivers. Our findings were intended to support the task of identifying which approaches are ready for wider dissemination and implementation. In a society experiencing unprecedented population longevity, this is a crucial task. To accomplish it, we tried to identify interventions and programs supported by evidence that met a minimum threshold of quality. We identified 595 unique eligible studies discussed in 850 publications, in which we found a remarkably diverse set of interventions. We used the National Institutes of Health (NIH) Stage Model framework to classify the interventions into pilot, explanatory, and “real world” pragmatic trials. This classification method allowed us to focus on the studies within this literature set that were best designed to look for real-world effects. We also removed from consideration studies with the potential to bias the outcomes due to concerns with study design or conduct.

Ultimately, we uncovered no moderate- or high-strength evidence to support care interventions and programs for active, widespread dissemination. We found low-strength evidence that collaborative care models (i.e. Care Ecosystems or discrete adaptations of the ACCESS models), may improve quality of life for PLWD and health system-level markers, including improvements in guideline-based quality indicators and reduction of emergency room visits, but the evidence was insufficient for informal caregiver outcomes. We also found low-strength evidence that an intensive multicomponent intervention, REACH II or discrete adaptations, improved informal caregiver depression at 6 months.

For all other interventions and outcomes, we found the evidence was insufficient. This does not mean that none of the individual interventions described are potentially useful for individual PLWD, their caregivers, or healthcare systems. Rather, it means that current available evidence cannot yet provide clear answers about which interventions offer consistent benefits. Therefore, the uncertainty of the evidence is too high for us to draw conclusions, at present. Further, when the evidence overall does not find a difference between groups, uncertainty is even higher about whether the lack of difference is truly because the interventions being compared did not differ in effect, or because the studies were designed to detect *differences* rather than *no difference*.

The lack of sufficient evidence to support widespread dissemination of all other interventions analyzed in this review leaves PLWD, caregivers, programs that support PLWD and caregivers, funders, and policymakers without clear answers. These groups, when deciding whether to disseminate or implement these interventions, will continue to depend on subjective observations, low-quality evidence, economics, and local and institutional policies. For individual PLWD and caregivers, trial and error with interventions, either one at a time or in combination, will likely continue as the norm.

Strengths and Weaknesses of the Review

We determined the methods for this review in order to best answer the question of readiness for broad dissemination. Decision making through this lens has implications for our findings.

We also adopted a review scope based on concerns about whether results from a literature search would give a biased view of interventions to address symptoms, safety, or quality of life, all of which overlap with frailty in older adults. For example, if falls risk and prevention differs

greatly between older adults with and without dementia, then research that specifically targets dementia would be useful. Without a big difference in outcomes, study populations likely remained mixed and these studies were screened out. Therefore, some specific approaches for a particular intervention, or even whole classes of interventions, may not have been captured. Similarly, we may have missed some community services and support approaches such as tool kits, referral services and links, or awareness-raising outreach. The case management literature revealed some of these studies, but search terms are diffuse and may have resulted in some studies being overlooked.

Our approach not to advance pilot, small sample, and high risk of bias studies to full analysis resulted in a very high-level assessment of the state of the science. It is possible that in many instances, the inclusion of the preliminary literature may have provided enough data for quantitative pooling for specific outcomes. Systematic reviews of specific interventions can investigate and report very fine details. This review was not intended for that purpose. Our use of the NIH Stage Model as a framework to focus on studies that would best support broad dissemination precluded reporting deeply on pilot studies. Additionally, current guidance for Evidence-based Practice Center (EPC) program methods supports not looking to studies assessed as high risk of bias to fortify what is already at best low-strength evidence.⁷⁰³ Lastly, small-study bias in reporting large effect sizes presents an important issue in a literature set where the majority of research is preliminary.⁷⁰⁴

Because we excluded studies with fewer than 10 participants per study arm, we may not have identified some interventions with very preliminary research supporting them. We accepted this limitation because exact precision for the research context findings was not feasible due to the wide range of care approaches and large literature set. Likewise, our decision not to include single-arm pre/post or evaluation studies limited our ability to address practices supported through evaluation studies.

Our approach to risk of bias assessment was generous, compared with how risk of bias is assessed in more targeted systematic review topics. In part, we based this decision on the unusually varied studies included in this review as well as the complexity of dementia and its associated care approaches. We allowed attrition to reach relatively high levels before assigning high risk of bias. Likewise, we treated fidelity generously, giving credit based on relatively brief mention. Concurrent treatments for specific interventions aimed at PLWD, especially for behavioral and psychiatric symptoms of dementia, were given a secondary position in the assessment. Unfortunately, studies rarely presented such information in a way that would allow for a sound evaluation of the implication for bias.

We also used the truncated risk of bias approach for studies assessed as pragmatic. The use of the PRagmatic Explanatory Continuum Indicator Summary-2 (PRECIS-2) tool in systematic reviews is preliminary for classifying and understanding the relative pragmatism in research design. We conservatively used a threshold approach, classifying studies as mostly explanatory, pragmatic, or balanced. We further labeled studies that were mostly pragmatic as good, fair, or marginal based on the analytic techniques. All of this represents modifications to EPC systematic review methods as we expand the boundaries of the topics systematic reviews are being asked to address. We cannot say with certainty how the practices used here affect the findings, regardless of whether we used more or less conservative methods for any particular finding. Nonetheless, as systematic reviews attempt to answer increasingly complex questions, we must find novel ways to answer them. The experiences from this review process may help inform future efforts.

Finally, given the already expansive breadth of this topic, our systematic review and meta-analysis is naturally reductionist in nature. That is, small but true differences may exist between many of the interventions within the various intervention categories we created for summarizing outcomes.

Future Research

The questions of “what works” in dementia care and how to deliver that care greatly interests researchers, funders, care providers, healthcare systems, and PLWD and their families. The intensive investment in dementia funding at the Federal level reflects the increasing public health importance not only of finding a way to prevent and treat dementia, but also of developing and eventually disseminating optimal dementia care and caregiver programs. While our review offers no firm conclusions, our findings provide valuable insights for the further development and improvement of dementia care science.

Current practice regarding how PLWD are diagnosed, treated, and supported throughout the disease trajectory are underpinned by concurrent and sometimes overlapping streams of research on dementia diagnosis and medical treatment, geriatric and chronic disease models of care, and dementia care support.⁷⁰⁵ Together, these streams of research have informed best practice recommendations for dementia care,^{706, 707} including the central idea that support for needs of family caregivers should be incorporated into any care plan.⁷⁰⁷

Methodological Rigor

Dementia care research has been slow to incorporate key elements of rigorous intervention design. Until relatively recently, many dementia care and caregiver intervention studies were not held to pre-registration of trials, data safety and monitoring boards, or other standards more common in other areas of clinical science including reporting standards required by journals (e.g., the Consolidated Standards of Reporting Trials [CONSORT] statement). As a direct result, despite a few positive findings, we reached the global conclusion that the evidence is insufficient to say with certainty that interventions were beneficial or not.

Federal funding requirements have instigated change in favor of oversight and reporting mechanisms that will likely lead to more transparent and reproducible research. However, many of these improvements in rigor did not occur until late 2015 and 2016, which coincided with increased investment in research on dementia care interventions. Therefore, many of the trials initially subject to more stringent data monitoring oversight and reporting are just now ending, and their findings have yet to be captured in reviews such as this one. This new, more rigorous research base will hopefully propel future comprehensive reviews to draw conclusions beyond insufficient evidence. Indeed, we did note an improvement in rigor from 2016 to 2017 in the published eligible studies.

In order for Federal funders and stakeholders to fulfill their goal of expediting the translational pipeline of idea development to implementation, critical improvements must be made in dementia care and caregiver research. Only with such improvements will we be able to draw clearer, less ambiguous conclusions related to efficacy.

Populations

During the topic refinement period for this review, we received many requests to ensure that certain groups that experience dementia were included as research participants in the examined literature. However, the published research rarely included many important populations. For

example, people with Down syndrome aging into dementia were absent, despite our later *ad hoc* literature searches to assure no relevant studies were missed. People with frontotemporal dementia were rarely included, and the few identified studies were limited to pilot or small sample studies. Few studies addressed racial or ethnic differences, and those that did were limited to major race/ethnic categories of Black/African American or Hispanic/Latinx. No study specifically studied LGBTQ populations. Likewise, culturally sensitive or culturally adapted interventions were rare. Some identified non-U.S. based research may help inform future intervention adaptations for PLWD or caregivers with immigrant or related racial/ethnic heritages.

Lastly, we identified very few studies of interventions specifically designed for low-resource areas (including rural and/or tribal communities) beyond pilot or small sample studies, which could not reach any level of certainty of the finding. All of these populations represent areas for future research.

Outcomes

Ultimately, care interventions aim to support quality of life and well-being and prevent harm for PLWD and caregivers, while enabling both to continue in their roles within their families and society. Unfortunately, quality of life was often not measured, and rarely as the outcome of primary interest. Further, PLWD and caregivers exist in relationships with one another, and this literature offers little to help us understand how they change in concert within their dyadic relationship. We attempted to address this issue by placing PLWD and caregiver outcomes adjacent to one another within the results section for each intervention; however, no clear patterns emerged. However, research would be improved by better measures for psychosocial outcomes in PLWD, and better methods of measurement.

Harms were rarely assessed. Although studies frequently measured caregiver burden, they far more often sought to observe reduction in burden than to check for increased burden. Other harms, such as elder abuse, were completely absent.

Additionally, the progressive nature of dementia and the anticipated increase in care needs complicates this research. Studies may be challenged to identify improvement in caregiver burden scales, because the goal of the intervention may in fact be to slow the rate of burden. Studies may be underpowered to detect such a small effect, or small benefits may be overwhelmed by the larger context, including social and financial implications, of caring for a person with a progressive condition.

Lastly, even within intervention categories, outcomes were variably measured and reported. More consistency in the outcomes measured would make it easier to assess bodies of evidence for specific interventions, or to understand how outcomes may differ by setting.

Interventions

The wide range of identified interventions, and the relationships between PLWD and caregivers, highlight the importance of understanding potential intervention mechanisms. We found low-strength evidence that multicomponent interventions may improve select outcomes, but the underlying question of what drives the benefit (i.e., the specific set of components, the mere presence of a multicomponent approach, or both) remains unanswered.

One anomaly of note in our findings is that while evidence was insufficient for all individual interventions, low-strength evidence showed that multicomponent interventions (i.e. discrete adaptations of REACH II) or collaborative care models (i.e. Care Ecosystems or discrete adaptations of the ACCESS model) could improve some outcomes. On the surface, this finding

appears contradictory to our global finding of insufficient evidence for all individual interventions. After all, theoretically, multicomponent interventions incorporate individual intervention components that have been shown to work. This discrepancy in our findings suggests two possibilities to explore with further research. One is that adequate support structures at the health system level, such as those provided by the REACH II⁵¹⁹ or Care Ecosystems interventions,⁶¹⁸ are crucial factors in addressing the needs of PLWD and their caregivers. Another possibility is that the particular set of interventions may matter less than whether PLWD and caregivers feel supported adequately at the health-system level. Perhaps such structural supports, although not specifically examined or measured in this literature, add significant value to the interventions applied within them.

Lack of consensus about taxonomies to classify interventions hampers the work of assessing this evidence base. We found a lack of precision in how individual interventions were described within specific intervention classifications, and this imprecision inhibits understanding. Many research publications used vague and inconsistent terminology for what constituted a specific intervention, especially given the broad range of baseline dementia severity. For example, we found lack of clarity in the differentiation between cognitive rehabilitation (aimed at restoring daily activity function) and cognitive training activities (cognitive drills that lack direct relationship to completion of daily activities). Authors often used both terms to describe intervention components within a single article. Clearly defining the classes of cognitive rehabilitation interventions would improve comparability across studies, as would specifying which interventions are suitable for PLWD with varying degrees of dementia-related impairment. This is made evident by the extent of misclassification across various systematic reviews and meta-analysis of the effectiveness of dementia caregiving interventions¹³ For example, in a synthesis of reviews, Gaugler and colleagues found that among eight meta-analyses and systematic reviews that considered psychoeducation/skills building, 45 individual interventions were inconsistently classified. One review classified them as psychoeducational/skills building interventions, another placed them in an entirely different category.¹³ Although taxonomies exist to improve the reporting of elements of dementia care and caregiver support interventions,⁷⁰⁸ studies often neglect to use them. Improved reporting of dementia care interventions through the use of taxonomy strategies, either in outcome evaluations or protocol reports, would enhance the ability of reviews such as this one to better classify interventions and thus improve inferences of efficacy/effectiveness.

Complex Interventions for Complex Systems

Most importantly, the care approaches examined in this review represent complex interventions nested within complex systems. The framework for care interventions from the NASEM Families Caring for an Aging America 2016 report displayed in Figure 1.1 in the Introduction sought to display this idea graphically. Complex systems, by their nature, always encompass some level of uncertainty; indeed, such irreducible uncertainty is a defining element of complex systems. In this literature, the multiple levels of uncertainty are difficult or impossible to overcome. Therefore, we must emphasize again that low-strength evidence is already a difficult bar to reach. Insufficient evidence, places where the evidence is very uncertain, means we could not, with integrity, say that a care approach is beneficial or not—which, as we have noted, is different from saying it does not work.

For example, even when a care approach focuses on a “lower” complex system level—such as the simple addition of aromatherapy to help a person feel calmer—the challenge is still steep to design a study that rules out all competing influences on that person’s sense of calmness. And

if the effect is small, or moderate but for only a very select group of people, that effect becomes difficult to see in the data, and uncertainty in the findings remains high.

A further problem with researching complex interventions is that complex systems tend to be self-preserving.⁷⁰⁹ Traditional research rests on an assumption that the way the overall process functions does not change as a result of being intervened upon. Yet, a defining feature of a complex system is its ability to adapt to change, even if it takes great effort to propel it from one state of homeostasis toward another, different one. And, when the system does arrive at a new state, it may not be the desired one, but instead an unanticipated adaptation. As a simple example, an intervention may teach staff how to do a new task intended to bring about a desired outcome. But the staff may look for ways to preserve the familiarity of their previous methods, and therefore devise work-arounds. Or they may feel the need to preserve efficiency overall, and make trade-offs in other areas of work. These modifications may multiply across the system as people adapt to the new approaches or requirements.

Many care approaches may actually be aimed, at least in part, at shifting the care culture. As the apex of complex systems, culture is highly diffuse. Yet, culture tends to shift through specific activities that gain traction over time. We see clear examples of this in Veteran-centered care in the United States, or person-centered care in non-U.S. settings. In both examples, training focuses on a relationship-based approach that reframes how caregivers perceive PLWD. Specific activities and tasks are considered secondary, and perhaps even a natural outcome, of this shift in perception and relationship.

Certainly, research is lacking in regard to how components in complex interventions interact to influence key outcomes. The prevailing approach in the dementia care literature is to develop an intervention with multiple components, and deliver it to determine efficacy on caregiver or PLWD outcomes; whether a single or select number of components are essential to an effect (if it exists) is generally unanswered. This has important and adverse implications for the field as a whole, and reduces the dissemination and implementation potential of dementia care interventions. This problem escalates as interventions increase in complexity due to the time, cost, and training requirements to deliver them successfully. In other domains of the intervention literature, approaches such as the Multiphase Optimization Strategy (MOST) have been applied to increase understanding of how singular intervention components interact to influence key outcomes prior to efficacy testing.⁷¹⁰ Such approaches may expedite the timeline from intervention development to potential implementation by crafting and evaluating interventions that are distilled to their essential components. Additionally, these approaches may offer greater insight into not just *whether* an intervention works, but *why*. It would also be helpful if researchers were to conduct and publish process evaluations of dementia care interventions that better described the mechanisms of benefit. Relatedly, basing dementia care and caregiver interventions on theories or conceptual models to test such mechanisms is an essential component of the NIH Stage Model. These theories and conceptual models should inform the design and evaluation of future dementia care interventions.

Intervention Fidelity

Whether interventions are simple or complex, problems with fidelity are significant for this research. Many studies reported no differences between groups. If an intervention showed benefit, delivery of a sufficient dose is assumed. Conversely, if no difference could be demonstrated, the reader is left with uncertainty about whether the dose was large enough. We used a liberal approach to fidelity while assessing risk of bias. Had we imposed a firm restriction on some form of fidelity measure, much or most of the literature would have been excluded.

Even so, problems with fidelity often contributed to high risk of bias. Unfortunately, fidelity assessment approaches lack consensus regarding both the components and how to measure them.

Further, we note that this research treats the concept of fidelity to intervention in training and formal caregiving differently than fidelity to intervention by informal caregivers. Informal caregivers have arguably the best incentives to help PLWD, which would motivate fidelity. However, many factors may inhibit fidelity even when desired. Although companion publications may use mixed-methods research to probe informal caregivers' views on an intervention, the question of whether interventions are delivered as designed and trained is rarely tracked for informal caregivers.

Implementation

Overall, the evidence we reviewed suggests that to consider questions related to dissemination and implementation at the outset of intervention design would result in more dementia care and caregiver interventions becoming ready for rapid implementation in real-world settings. For example, incorporating measures or indicators of implementation (e.g., appropriateness, feasibility, acceptability, cost) alongside clinical outcomes of efficacy/effectiveness would expedite the timeline from dementia care intervention development and evaluation to dissemination and implementation. That, in turn, would help the interventions that demonstrate efficacy to reach and benefit those who need and desire them.⁷¹¹

Threats to scalability in dementia care include reliance on interventions that require extensive training and fidelity monitoring; too great a need to rely on the original developers of the intervention; a requirement for highly trained and skilled professionals to deliver the intervention; highly complex, intense, and costly interventions; lack of implementation manuals; and lack of payment mechanisms to ensure sustainability.⁷¹²

Further, implementation will also differ by setting. Future attempts to transfer or modify interventions to assisted living facilities could be helped by a better understanding of implementation factors. Assisted living facilities were among the least studied settings in the included literature.

Another important question pertains to the evidentiary standard that should apply to the complex interventions so overwhelmingly represented in this literature set. EPC guidance provides, as does the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) working group, clear principles and guidelines to apply to medical treatment or therapies that come with benefits weighed against potential harms. Complex interventions applied to complex whole-person and community systems in varied and often complex settings can be exceedingly difficult to implement with fidelity and potential for replication and dissemination. In addition, the study of these interventions is challenged by the appropriateness of basic assumptions that underlie research designs and statistical tools.

One approach to many of these issues is better use of community-based research methods. Involving PLWD and caregivers in planning, implementing, and disseminating research is increasingly recognized as valuable for addressing the populations for which interventions are intended. Applying this approach to PLWD and their caregivers was recognized at a Health and Human Services Summit in 2017.⁷¹³

Broader Research Context

Another concern is how informal caregivers are perceived, and the way in which this perception informs research designs. While caregiving for PLWD presents challenges, burdens, and risks to the health of caregivers, it is not a pathological condition. Interventions aimed at

mitigating burdens for informal caregivers can also recognize and build on the rewards of caregiving and the bonds it nurtures between caregivers/partners and care recipients.

Additional questions to consider pertain to how dementia care science might be conducted more efficiently and effectively. Perhaps it is possible to create an environment that encourages experimenting with care solutions while maximizing the ability to learn from those efforts. One avenue for growth may lie in more collaborative, open science with collective impact approaches to its development. This represents a different form of “big science,” in which resources such as Benjamin Rose Institute on Aging’s (BRI) Best Practice Caregiving resource (see <https://www.benrose.org/best-practice-caregiving>) could be an important asset, allowing us another way to close feedback loops and gain knowledge from real-world applications. Innovation requires something more than status quo behavior, as does the research needed to push toward paradigm growth or adoption. Innovation in research involves maximizing discovery while minimizing the risk of locking in solutions that are only *currently* relevant or *partially* accurate. Perhaps we can find easier ways for researchers and care systems to test new things efficiently, without overburdening coordination and/or oversight. Many researchers are already thinking deeply about these kinds of questions, including the Center for Open Science (see <https://cos.io/> or <http://www.unesco.org/new/en/communication-and-information/portals-and-platforms/goap/>) or MetaScience (see <https://www.metascience2019.org/>) initiatives.

Much research has limited impact because it is conducted in isolation, among small groups, and with variable timing for study endpoints to measure outcomes. However, demonstration projects have been conducted to examine specific research questions being crowdsourced by the research community, with peer review processes built in prior to and during analysis, rather than through an extensive research and publication process.⁷¹⁴ This type of collaborative research can prioritize outcome measurement tools and timing in a way that leads to science that others can measure and replicate.

Many aspects of care interventions for PLWD and their caregivers need more thorough exploration. We hesitated to give an exhaustive list for fear of overwhelming the readers. We were instead guided by peer and public comments on the draft version of this report to specifically mention the following areas: functional and health status limitations, access to care and intervention services and supports as well as accessibility, transportation, culture, racial/ethnic, and related factors. Public commentators to this report who are living with dementia noted an urgent need for more research on interventions that support personhood, purpose and meaning, social and peer supports, proactive approaches to living with a chronic, progressive illness, and lifestyle and spirituality interventions.

Importantly, we do note a gap in the literature for interventions that address the early stages of dementia and being an informal caregiver, just at the time of diagnosis, when problems associated with disconnection, denial, and misinformation may be most urgent.

In the end, high-level discussions of how future research might be structured should not distract us from the primary need to provide research that is relevant to all of the populations that matter. With the exception of a handful of studies, the current research is silent for many social groups, whether by race/ethnicity, citizen status, geographic locations, or dementia types, to name some of the larger categories. Much crucial work remains to be done.

Abbreviations and Acronyms

ACCESS	Alzheimer's Disease Coordinated Care for San Diego Seniors
ACP	Advance Care Planning
AD	Alzheimer's Disease
ADL	Activities of Daily Living
ADRD	Alzheimer's Disease and Related Dementias
AHRQ	Agency for Healthcare Research and Quality
AMSTAR	Assessment of Multiple Systematic Reviews
BPSD	Behavioral and Psychological Symptoms of Dementia
CAM	Complementary and Alternative Medicine
CONSORT	Consolidated Standards of Reporting Trials
CST	Cognitive Stimulation Therapy
DCM™	Dementia Care Mapping™
EPC	Evidence-based Practice Center
HHS	Health and Human Services
HMD	Health and Medicine Division
IMPACT	IMbedded Pragmatic Alzheimer's disease and Related Dementias Clinical Trials
KQ	Key Question
MMSE	Mini Mental State Examination
MOST	Multiphase Optimization Strategy
MSS	Multisensory Stimulation
N	Number
NASEM	National Academies of Sciences, Engineering, and Medicine
NIA	National Institute on Aging
NIH	National Institutes of Health
NPI	Neuropsychiatric Inventory
PICOTS	Population, Intervention, Comparison, Outcomes and Timing
PRECIS-2	PRagmatic EXplanatory Continuum Indicator Summary-2 tool
PRISMA	Preferred Reporting Items for Systematic Review and Meta-Analysis
PLWD	People Living With Dementia
QoL	Quality of Life
QOL-AD	Quality of Life-Alzheimer's Disease
RCT	Randomized Controlled Trial
RDAD	Reducing Disability in Alzheimer's Disease
REACH II	Resources for Enhancing Alzheimer's Caregivers Health
RMC	Routine Medical Care
SMD	Standardized Mean Difference

STAR-C	Social learning theory and principles of behavior analysis of caregivers
UK	United Kingdom
US	United States

References

1. Alzheimer's Association. 2018 Alzheimer's disease facts and figures. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association*. 2018;14:367–429.
2. Matthews KA, Xu W, Gaglioti AH, et al. Racial and ethnic estimates of Alzheimer's disease and related dementias in the United States (2015-2060) in adults aged ≥ 65 years. *Alzheimers Dement*. 2019 Jan;15(1):17-24. doi: 10.1016/j.jalz.2018.06.3063. PMID: 30243772.
3. Hebert LE, Weuve J, Scherr PA, et al. Alzheimer disease in the United States (2010-2050) estimated using the 2010 census. *Neurology*. 2013 May 7;80(19):1778-83. doi: 10.1212/WNL.0b013e31828726f5. PMID: 23390181.
4. McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia. The journal of the Alzheimer's Association*. 2011 May;7(3):263-9. doi: 10.1016/j.jalz.2011.03.005. PMID: 21514250.
5. Office of the Assistant Secretary for Planning and Evaluation. National Alzheimer's Project Act. <https://aspe.hhs.gov/national-alzheimers-project-act> <https://aspe.hhs.gov/national-alzheimers-project-act>. Accessed on February 22 2019.
6. Hurd MD, Martorell P, Delayande A, et al. Monetary costs of dementia in the United States. *New England Journal of Medicine*. 2013 Apr 4;368(14):1326-34. doi: 10.1056/NEJMs1204629. PMID: 23550670.
7. Zhao QF, Tan L, Wang HF, et al. The prevalence of neuropsychiatric symptoms in Alzheimer's disease: Systematic review and meta-analysis. *Journal of Affective Disorders*. 2016;190:264-71. doi: 10.1016/j.jad.2015.09.069. PMID: 26540080.
8. National Alzheimer's Project Act. 124 STAT. 4100 111th Congress. United States of America; 2011.
9. Reus VI, Fochtmann LJ, Eyler AE, et al. The American Psychiatric Association Practice Guideline on the Use of Antipsychotics to Treat Agitation or Psychosis in Patients With Dementia. *American Journal of Psychiatry*. 2016 May 1;173(5):543-6. doi: 10.1176/appi.ajp.2015.173501. PMID: 27133416.
10. Corbett A, Stevens J, Aarsland D, et al. Systematic review of services providing information and/or advice to people with dementia and/or their caregivers. *International Journal of Geriatric Psychiatry*. 2012 June 27;27(6):628-36. doi: 10.1002/gps.2762. PMID: 22038644.
11. Centers for Disease Control and Prevention. Caregiving for Person with Alzheimer's Disease or a related Dementia. <https://www.cdc.gov/aging/caregiving/alzheimer.htm>. Accessed on Feb 22.
12. Committee on Family Caregiving for Older Adults. Families Caring for an Aging America. In: Shulz R, Eden J, eds. Washington (DC): National Academies Press (US); 2016.
13. Gaugler JE, Jutkowitz E, Shippee TP, et al. Consistency of dementia caregiver intervention classification: an evidence-based synthesis. *International psychogeriatrics / IPA*. 2017;29(1):19-30. doi: 10.1017/S1041610216001514. PMID: 27671663.

14. Warshaw GA, Bragg EJ. Preparing The Health Care Workforce To Care For Adults With Alzheimer's Disease And Related Dementias. *Health Affairs*. 2014;33(4):633-41. doi: 10.1377/hlthaff.2013.1232. PMID: 24711325.
15. Onken LS, Carroll KM, Shoham V, et al. Reenvisioning clinical science: Unifying the discipline to improve the public health. *Clinical Psychological Science*. 2014;2(1):22-34. doi: 10.1177/2167702613497932. PMID: 25821658.
16. Moher D, Altman DG, Liberati A, et al. PRISMA statement. *Epidemiology*. 2011 Jan;22(1):128; author reply doi: <http://dx.doi.org/10.1097/EDE.0b013e3181fe7825>. PMID: 21150360.
17. Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358. doi: 10.1136/bmj.j4008.
18. Viswanathan M, Ansari M, Berkman N, et al. Assessing the Risk of Bias of Individual Studies in Systematic Reviews of Health Care Interventions AHRQ. 2012.
19. Hempel S, Shekelle PG, Liu JL, et al. Development of the quality improvement minimum quality criteria set (QI-MOCS): a tool for critical appraisal of quality improvement intervention publications. *BMJ Quality and Safety*. 2015;24:796-804.
20. Patsopoulos N. A pragmatic view on pragmatic trials. *Dialogues Clin Neurosci*. 2011;13(2):217-24. PMID: 21842619.
21. Loudon K, Treweek S, Sullivan F, et al. The PRECIS-2 tool: designing trials that are fit for purpose. *Bmj*. 2015 May 8;350:h2147. doi: 10.1136/bmj.h2147. PMID: 25956159.
22. Berkman ND, Lohr K, Ansari M, et al. Grading the strength of a body of evidence when assessing health care interventions for the effective health care program of the Agency for Healthcare Research and Quality: An update. Rockville, MD: Agency for Healthcare Research and Quality; 2013. (vol. AHRQ Publication No. 13(14)-EHC 130-EF).
23. Jones C, Moyle W, Murfield J, et al. Does Cognitive Impairment and Agitation in Dementia Influence Intervention Effectiveness? Findings From a Cluster-RCT With the Therapeutic Robot, PARO. *Journal of the American Medical Directors Association*. 2018 Apr 13;13:13. doi: <https://dx.doi.org/10.1016/j.jamda.2018.02.014>. PMID: 29656838.
24. Mervin MC, Moyle W, Jones C, et al. The Cost-Effectiveness of Using PARO, a Therapeutic Robotic Seal, to Reduce Agitation and Medication Use in Dementia: Findings from a Cluster-Randomized Controlled Trial. *Journal of the American Medical Directors Association*. 2018 Jan 09;09:09. doi: <https://dx.doi.org/10.1016/j.jamda.2017.10.008>. PMID: 29325922.
25. Moyle W, Jones C, Murfield J, et al. Effect of a robotic seal on the motor activity and sleep patterns of older people with dementia, as measured by wearable technology: A cluster-randomised controlled trial. *Maturitas*. 2018 01 Apr;110:10-7. doi: <http://dx.doi.org/10.1016/j.maturitas.2018.01.007>. PMID: 621060108.
26. Moyle W, Jones CJ, Murfield JE, et al. Use of a Robotic Seal as a Therapeutic Tool to Improve Dementia Symptoms: A Cluster-Randomized Controlled Trial. *Journal of the American Medical Directors Association*. 2017 Sep 01;18(9):766-73. doi: <https://dx.doi.org/10.1016/j.jamda.2017.03.018>. PMID: 28780395.

27. Bemelmans R, Gelderblom GJ, Jonker P, et al. Effectiveness of Robot Paro in Intramural Psychogeriatric Care: A Multicenter Quasi-Experimental Study. *Journal of the American Medical Directors Association*. 2015 Nov 01;16(11):946-50. doi: <https://dx.doi.org/10.1016/j.jamda.2015.05.007>. PMID: 26115817.
28. Joranson N, Pedersen I, Rokstad AM, et al. Effects on Symptoms of Agitation and Depression in Persons With Dementia Participating in Robot-Assisted Activity: A Cluster-Randomized Controlled Trial. *Journal of the American Medical Directors Association*. 2015 Oct 01;16(10):867-73. doi: <https://dx.doi.org/10.1016/j.jamda.2015.05.002>. PMID: 26096582.
29. Joranson N, Pedersen I, Rokstad AM, et al. Change in quality of life in older people with dementia participating in Paro-activity: a cluster-randomized controlled trial.[Erratum appears in *J Adv Nurs*. 2017 Jun;73(6):1531; PMID: 28485533]. *Journal of Advanced Nursing*. 2016 Dec;72(12):3020-33. doi: <https://dx.doi.org/10.1111/jan.13076>. PMID: 27434512.
30. Liang A, Piroth I, Robinson H, et al. A Pilot Randomized Trial of a Companion Robot for People With Dementia Living in the Community. *Journal of the American Medical Directors Association*. 2017 Oct 01;18(10):871-8. doi: <https://dx.doi.org/10.1016/j.jamda.2017.05.019>. PMID: 28668664.
31. Moyle W, Cooke M, Beattie E, et al. Exploring the effect of companion robots on emotional expression in older adults with dementia: a pilot randomized controlled trial. *Journal of Gerontological Nursing*. 2013 May;39(5):46-53. doi: <https://dx.doi.org/10.3928/00989134-20130313-03>. PMID: 23506125.
32. Petersen S, Houston S, Qin H, et al. The Utilization of Robotic Pets in Dementia Care. *Journal of Alzheimer's Disease*. 2017;55(2):569-74. PMID: 27716673.
33. Valenti SM, Aguera-Ortiz L, Olazarán RJ, et al. Social robots in advanced dementia. *Frontiers in aging neuroscience*. 2015;7(Jun). doi: 10.3389/fnagi.2015.00133. PMID: CN-01103263.
34. Pu L, Moyle W, Jones C, et al. The Effect of Using PARO for People Living With Dementia and Chronic Pain: A Pilot Randomized Controlled Trial. *Journal of the American Medical Directors Association*. 2020 Feb 28;28:28. doi: <https://dx.doi.org/10.1016/j.jamda.2020.01.014>. PMID: 32122797.
35. Majic T, Gutzmann H, Heinz A, et al. Animal-assisted therapy and agitation and depression in nursing home residents with dementia: a matched case-control trial. *American Journal of Geriatric Psychiatry*. 2013 Nov;21(11):1052-9. doi: <https://dx.doi.org/10.1016/j.jagp.2013.03.004>. PMID: 23831177.
36. Menna LF, Santaniello A, Gerardi F, et al. Efficacy of animal-assisted therapy adapted to reality orientation therapy: measurement of salivary cortisol. *Psychogeriatrics: The Official Journal of the Japanese Psychogeriatric Society*. 2019 Sep;19(5):510-2. doi: <https://dx.doi.org/10.1111/psyg.12418>. PMID: 30740833.
37. Olsen C, Pedersen I, Bergland A, et al. Effect of animal-assisted activity on balance and quality of life in home-dwelling persons with dementia. *Geriatric Nursing*. 2016 Jul-Aug;37(4):284-91. doi: <https://dx.doi.org/10.1016/j.gerinurse.2016.04.002>. PMID: 27155968.
38. Olsen C, Pedersen I, Bergland A, et al. Effect of animal-assisted interventions on depression, agitation and quality of life in nursing home residents suffering from cognitive impairment or dementia: a cluster randomized controlled trial. *International Journal of Geriatric Psychiatry*. 2016 Dec;31(12):1312-21. doi: <https://dx.doi.org/10.1002/gps.4436>. PMID: 26807956.

39. Travers C, Perkins J, Rand J, et al. An evaluation of dog-assisted therapy for residents of aged care facilities with dementia. *Anthrozoos*. 2013;26(2):213-25. PMID: CN-00871982.
40. Balzotti A, Filograsso M, Altamura C, et al. Comparison of the efficacy of gesture-verbal treatment and doll therapy for managing neuropsychiatric symptoms in older patients with dementia. *International Journal of Geriatric Psychiatry*. 2019 09;34(9):1308-15. doi: <https://dx.doi.org/10.1002/gps.4961>. PMID: 30136743.
41. Moyle W, Murfield J, Jones C, et al. Can lifelike baby dolls reduce symptoms of anxiety, agitation, or aggression for people with dementia in long-term care? Findings from a pilot randomised controlled trial. *Aging & Mental Health*. 2019;23(10):1442-50. doi: 10.1080/13607863.2018.1498447. PMID: 138051539. Language: English. Entry Date: 20190817. Revision Date: 20190823. Publication Type: Article.
42. Cantarella A, Borella E, Faggian S, et al. Using dolls for therapeutic purposes: A study on nursing home residents with severe dementia. *International Journal of Geriatric Psychiatry*. 2018 July;33(7):915-25. doi: <http://dx.doi.org/10.1002/gps.4872>. PMID: 622559552.
43. Baker R, Bell S, Baker E, et al. A randomized controlled trial of the effects of multi-sensory stimulation (MSS) for people with dementia. *British Journal of Clinical Psychology*. 2001 Mar;40(Pt 1):81-96. PMID: 11317951.
44. Baker R, Holloway J, Holtkamp CCM, et al. Effects of multi-sensory stimulation for people with dementia. *Journal of Advanced Nursing*. 2003;43(5):465-77. doi: 10.1046/j.1365-2648.2003.02744.x. PMID: 106688835. Language: English. Entry Date: 20050425. Revision Date: 20150819. Publication Type: Journal Article.
45. Strom BS, Engedal K, Benth JS, et al. Effect of the Sonas Programme on Communication in People with Dementia: A Randomized Controlled Trial. *Dementia and Geriatric Cognitive Disorders Extra*. 2017 Jan-Apr;7(1):122-35. doi: <https://dx.doi.org/10.1159/000468147>. PMID: 28553314.
46. van Weert JC, van Dulmen AM, Spreeuwenberg PM, et al. The effects of the implementation of snoezelen on the quality of working life in psychogeriatric care. *International Psychogeriatrics*. 2005 Sep;17(3):407-27. PMID: 16252374.
47. Baillon S, Van Diepen E, Prettyman R, et al. A comparison of the effects of Snoezelen and reminiscence therapy on the agitated behaviour of patients with dementia. *International Journal of Geriatric Psychiatry*. 2004 Nov;19(11):1047-52. PMID: 15481068.
48. Collier L, McPherson K, Ellis-Hill C, et al. Multisensory stimulation to improve functional performance in moderate to severe dementia—interim results. *American Journal of Alzheimer's Disease & Other Dementias*. 2010;25(8):698-703. doi: 10.1177/1533317510387582. PMID: 104960420. Language: English. Entry Date: 20101214. Revision Date: 20150820. Publication Type: Journal Article.
49. Staal JA, Sacks A, Matheis R, et al. The effects of Snoezelen (multi-sensory behavior therapy) and psychiatric care on agitation, apathy, and activities of daily living in dementia patients on a short term geriatric psychiatric inpatient unit. *International Journal of Psychiatry in Medicine*. 2007;37(4):357-70. PMID: 18441625.
50. Van Diepen E, Baillon SF, Redman J, et al. A pilot study of the physiological and behavioural effects of snoezelen in dementia. *British Journal of Occupational Therapy*. 2002 February;65(2):61-6. PMID: 35396447.

51. Baker R, Dowling Z, Wareing LA, et al. Snoezelen: its Long-Term and Short-Term Effects on Older People with Dementia. *British Journal of Occupational Therapy*. 1997;60(5):213-8. PMID: 104158278. Language: English. Entry Date: 20131114. Revision Date: 20150820. Publication Type: Journal Article.
52. Maseda A, Cibeira N, Lorenzo-Lopez L, et al. Multisensory Stimulation and Individualized Music Sessions on Older Adults with Severe Dementia: Effects on Mood, Behavior, and Biomedical Parameters. *Journal of Alzheimer's Disease*. 2018;63(4):1415-25. doi: <https://dx.doi.org/10.3233/JAD-180109>. PMID: 29843244.
53. Kouzuki M, Kitao S, Kaju T, et al. Evaluation of the effect of aroma oil as a bath salt on cognitive function. *Psychogeriatrics : the official journal of the Japanese Psychogeriatric Society*. 2019;02. doi: <http://dx.doi.org/10.1111/psyg.12481>. PMID: 629513763.
54. Moorman Li R, Gilbert B, Orman A, et al. Evaluating the effects of diffused lavender in an adult day care center for patients with dementia in an effort to decrease behavioral issues: a pilot study. *Journal of Drug Assessment*. 2017;6(1):1-5. doi: <https://dx.doi.org/10.1080/21556660.2016.1278545>. PMID: 28265482.
55. Chieh-Yu F, Moyle W, Cooke M. A randomised controlled trial of the use of aromatherapy and hand massage to reduce disruptive behaviour in people with dementia. *BMC Complementary & Alternative Medicine*. 2013;13(1):165-73. doi: 10.1186/1472-6882-13-165. PMID: 104229104. Language: English. Entry Date: 20130205. Revision Date: 20150711. Publication Type: Journal Article.
56. Smallwood J, Brown R, Coulter F, et al. Aromatherapy and behaviour disturbances in dementia: a randomized controlled trial. *International Journal of Geriatric Psychiatry*. 2001 Oct;16(10):1010-3. PMID: 11607948.
57. Moyle W, Cooke ML, Beattie E, et al. Foot massage versus quiet presence on agitation and mood in people with dementia: a randomised controlled trial. *International Journal of Nursing Studies*. 2014 Jun;51(6):856-64. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2013.10.019>. PMID: 24216598.
58. Fu CY, Moyle W, Cooke M. A randomised controlled trial of the use of aromatherapy and hand massage to reduce disruptive behaviour in people with dementia. *BMC Complementary & Alternative Medicine*. 2013 Jul 10;13:165. doi: <https://dx.doi.org/10.1186/1472-6882-13-165>. PMID: 23837414.
59. Fung JKK, Tsang HW. Management of behavioural and psychological symptoms of dementia by an aromamassage with acupressure treatment protocol: A randomised clinical trial. *Journal of Clinical Nursing*. 2018 May;27(9-10):1812-25. doi: <https://dx.doi.org/10.1111/jocn.14101>. PMID: 28986942.
60. Lu DF, Hart LK, Lutgendorf SK, et al. Slowing progression of early stages of AD with alternative therapies: a feasibility study. *Geriatric Nursing*. 2013 Nov-Dec;34(6):457-64. doi: <https://dx.doi.org/10.1016/j.gerinurse.2013.07.003>. PMID: 23972540.
61. Jia Y, Zhang X, Yu J, et al. Acupuncture for patients with mild to moderate Alzheimer's disease: a randomized controlled trial. *BMC Complementary & Alternative Medicine*. 2017 Dec 29;17(1):556. doi: <https://dx.doi.org/10.1186/s12906-017-2064-x>. PMID: 29284465.
62. Jimbo D, Kimura Y, Taniguchi M, et al. Effect of aromatherapy on patients with Alzheimer's disease. *Psychogeriatrics*. 2009;9(4):173-9. doi: 10.1111/j.1479-8301.2009.00299.x. PMID: CN-00750884.

63. Rodriguez-Mansilla J, Gonzalez-Lopez-Arza MV, Varela-Donoso E, et al. Ear therapy and massage therapy in the elderly with dementia: a pilot study. *Journal of Traditional Chinese Medicine*. 2013 Aug;33(4):461-7. PMID: 24187866.
64. Kwan RYC, Leung MCP, Lai CKY. A Randomized Controlled Trial Examining the Effect of Acupressure on Agitation and Salivary Cortisol in Nursing Home Residents with Dementia. *Dementia & Geriatric Cognitive Disorders*. 2017;44(1-2):92-104. doi: <https://dx.doi.org/10.1159/000478739>. PMID: 28768251.
65. Woods DL, Craven RF, Whitney J. The effect of therapeutic touch on behavioral symptoms of persons with dementia. *Alternative Therapies in Health & Medicine*. 2005 Jan-Feb;11(1):66-74. PMID: 15712768.
66. Zalomonson S, Freud T, Punchik B, et al. The results of a cross-over placebo-controlled study of the effect of lavender oil on behavioral and psychological symptoms of dementia. *Rejuvenation Research*. 2018 Oct 17;17:17. doi: <https://dx.doi.org/10.1089/rej.2018.2123>. PMID: 30328781.
67. Yang YP, Wang CJ, Wang JJ. Effect of Aromatherapy Massage on Agitation and Depressive Mood in Individuals With Dementia. *Journal of Gerontological Nursing*. 2016 Sep 01;42(9):38-46. doi: <https://dx.doi.org/10.3928/00989134-20160615-03>. PMID: 27319407.
68. Hawranik P, Johnston P, Deatrich J. Therapeutic touch and agitation in individuals with Alzheimer's disease. *Western Journal of Nursing Research*. 2008 Jun;30(4):417-34. doi: <https://dx.doi.org/10.1177/0193945907305126>. PMID: 18272750.
69. Kwan RY, Leung MC, Lai CK. Acupressure for agitation in nursing home residents with dementia: study protocol for a randomized controlled trial. *Trials* [Electronic Resource]. 2014 Oct 27;15:410. doi: <https://dx.doi.org/10.1186/1745-6215-15-410>. PMID: 25346026.
70. Rodriguez-Mansilla J, Gonzalez Lopez-Arza MV, Varela-Donoso E, et al. The effects of ear acupressure, massage therapy and no therapy on symptoms of dementia: a randomized controlled trial. *Clinical Rehabilitation*. 2015 Jul;29(7):683-93. doi: <https://dx.doi.org/10.1177/0269215514554240>. PMID: 25322869.
71. Takahashi Y, Shindo S, Kanbayashi T, et al. Examination of the influence of cedar fragrance on cognitive function and behavioral and psychological symptoms of dementia in Alzheimer type dementia. *Neuropsychopharmacology reports*. 2020. PMID: CN-02084429.
72. Ballard CG, O'Brien JT, Reichelt K, et al. Aromatherapy as a safe and effective treatment for the management of agitation in severe dementia: the results of a double-blind, placebo-controlled trial with Melissa. *Journal of Clinical Psychiatry*. 2002 Jul;63(7):553-8. PMID: 12143909.
73. Burns A, Perry E, Holmes C, et al. A double-blind placebo-controlled randomized trial of Melissa officinalis oil and donepezil for the treatment of agitation in Alzheimer's disease. *Dementia & Geriatric Cognitive Disorders*. 2011;31(2):158-64. doi: <https://dx.doi.org/10.1159/000324438>. PMID: 21335973.
74. Lin PW, Chan WC, Ng BF, et al. Efficacy of aromatherapy (*Lavandula angustifolia*) as an intervention for agitated behaviours in Chinese older persons with dementia: a cross-over randomized trial. *International Journal of Geriatric Psychiatry*. 2007 May;22(5):405-10. PMID: 17342790.

75. Yang SY, Shan CL, Qing H, et al. The Effects of Aerobic Exercise on Cognitive Function of Alzheimer's Disease Patients. *CNS & Neurological Disorders Drug Targets*. 2015;14(10):1292-7. PMID: 26556080.
76. Bramble M, Moyle W, Shum D. A quasi-experimental design trial exploring the effect of a partnership intervention on family and staff well-being in long-term dementia care. *Aging & Mental Health*. 2011 Nov;15(8):995-1007. doi: <https://dx.doi.org/10.1080/13607863.2011.583625>. PMID: 21702706.
77. Ancoli-Israel S, Martin JL, Gehrman P, et al. Effect of light on agitation in institutionalized patients with severe Alzheimer disease. *American Journal of Geriatric Psychiatry*. 2003 Mar-Apr;11(2):194-203. PMID: 12611749.
78. Burns A, Allen H, Tomenson B, et al. Bright light therapy for agitation in dementia: a randomized controlled trial. *International Psychogeriatrics*. 2009 Aug;21(4):711-21. doi: <https://dx.doi.org/10.1017/S1041610209008886>. PMID: 19323872.
79. Dowling GA, Graf CL, Hubbard EM, et al. Light treatment for neuropsychiatric behaviors in Alzheimer's disease. *Western Journal of Nursing Research*. 2007 Dec;29(8):961-75. PMID: 17596638.
80. Fontana Gasio P, Krauchi K, Cajochen C, et al. Dawn-dusk simulation light therapy of disturbed circadian rest-activity cycles in demented elderly. *Experimental Gerontology*. 2003 Jan-Feb;38(1-2):207-16. PMID: 12543279.
81. Hickman SE, Barrick AL, Williams CS, et al. The effect of ambient bright light therapy on depressive symptoms in persons with dementia. *Journal of the American Geriatrics Society*. 2007 Nov;55(11):1817-24. PMID: 17944896.
82. Mishima K, Hishikawa Y, Okawa M. Randomized, dim light controlled, crossover test of morning bright light therapy for rest-activity rhythm disorders in patients with vascular dementia and dementia of Alzheimer's type. *Chronobiology International*. 1998 Nov;15(6):647-54. PMID: 9844752.
83. Onega LL, Pierce TW, Epperly L. Effect of Bright Light Exposure on Depression and Agitation in Older Adults with Dementia. *Issues in Mental Health Nursing*. 2016 Sep;37(9):660-7. PMID: 27254531.
84. Riemersma-van der Lek RF, Swaab DF, Twisk J, et al. Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities: a randomized controlled trial. *JAMA: Journal of the American Medical Association*. 2008;299(22):2642-55. doi: 10.1001/jama.299.22.2642. PMID: 105752996. Language: English. Entry Date: 20080627. Revision Date: 20161112. Publication Type: journal article.
85. Figueiro MG, Plitnick B, Roohan C, et al. Effects of a Tailored Lighting Intervention on Sleep Quality, Rest-Activity, Mood, and Behavior in Older Adults With Alzheimer Disease and Related Dementias: A Randomized Clinical Trial. *Journal of Clinical Sleep Medicine*. 2019 Dec 15;15(12):1757-67. doi: <https://dx.doi.org/10.5664/jcsm.8078>. PMID: 31855161.
86. Riemersma-van der Lek RF, Swaab DF, Twisk J, et al. Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities: a randomized controlled trial. *JAMA*. 2008 Jun 11;299(22):2642-55. doi: <https://dx.doi.org/10.1001/jama.299.22.2642>. PMID: 18544724.

87. Toseland RW, Diehl M, Freeman K, et al. The impact of validation group therapy on nursing home residents with dementia. *Journal of Applied Gerontology*. 1997 March;16(1):31-50. doi: <http://dx.doi.org/10.1177/073346489701600102>. PMID: 27097884.
88. Spector A, Charlesworth G, King M, et al. Cognitive-behavioural therapy for anxiety in dementia: pilot randomised controlled trial. *British Journal of Psychiatry*. 2015 Jun;206(6):509-16. doi: <https://dx.doi.org/10.1192/bjp.bp.113.140087>. PMID: 25698766.
89. Stanley MA, Calleo J, Bush AL, et al. The peaceful mind program: a pilot test of a cognitive-behavioral therapy-based intervention for anxious patients with dementia. *American Journal of Geriatric Psychiatry*. 2013 Jul;21(7):696-708. doi: <https://dx.doi.org/10.1016/j.jagp.2013.01.007>. PMID: 23567399.
90. Beck CK, Vogelpohl TS, Rasin JH, et al. Effects of behavioral interventions on disruptive behavior and affect in demented nursing home residents. *Nursing Research*. 2002 Jul-Aug;51(4):219-28. PMID: 12131234.
91. Kiosses DN, Rosenberg PB, McGovern A, et al. Depression and Suicidal Ideation During Two Psychosocial Treatments in Older Adults with Major Depression and Dementia. *Journal of Alzheimer's Disease*. 2015;48(2):453-62. doi: <https://dx.doi.org/10.3233/JAD-150200>. PMID: 26402009.
92. Richards KC, Beck C, O'Sullivan PS, et al. Effect of individualized social activity on sleep in nursing home residents with dementia. *Journal of the American Geriatrics Society*. 2005 Sep;53(9):1510-7. PMID: 16137280.
93. Law W, Kwok TCY. Impacts of a multicomponent intervention programme on neuropsychiatric symptoms in people with dementia and psychological health of caregivers: A feasibility pilot study. *International Journal of Geriatric Psychiatry*. 2019 Aug 07;07:07. doi: <https://dx.doi.org/10.1002/gps.5191>. PMID: 31390090.
94. Bailey EM, Stevens AB, LaRocca MA, et al. A Randomized Controlled Trial of a Therapeutic Intervention for Nursing Home Residents With Dementia and Depressive Symptoms. *Journal of Applied Gerontology*. 2017 Jul;36(7):895-908. doi: <https://dx.doi.org/10.1177/0733464815627956>. PMID: 26912731.
95. Fischer-Terworth C, Probst P. Evaluation of a TEACCH-and music therapy-based psychological intervention in mild to moderate dementia. In *Geropsych: the journal of gerontopsychology and geriatric psychiatry*
96. Kovach CR, Taneli Y, Dohearty P, et al. Effect of the BACE intervention on agitation of people with dementia. *Gerontologist*. 2004 Dec;44(6):797-806. PMID: 15611216.
97. Lawton MP, Van Haitsma K, Klapper J, et al. A stimulation-retreat special care unit for elders with dementing illness. *International Psychogeriatrics*. 1998 Dec;10(4):379-95. PMID: 9924833.
98. Lin LC, Yang MH, Kao CC, et al. Using acupuncture and Montessori-based activities to decrease agitation for residents with dementia: a cross-over trial. *Journal of the American Geriatrics Society*. 2009 Jun;57(6):1022-9. PMID: 19507295.
99. Schrijnemaekers V, van Rossum E, Candel M, et al. Effects of emotion-oriented care on elderly people with cognitive impairment and behavioral problems. *International Journal of Geriatric Psychiatry*. 2002 Oct;17(10):926-37. PMID: 12325052.

100. Finnema E, Droes RM, Ettema T, et al. The effect of integrated emotion-oriented care versus usual care on elderly persons with dementia in the nursing home and on nursing assistants: a randomized clinical trial. *International Journal of Geriatric Psychiatry*. 2005 Apr;20(4):330-43. PMID: 15799079.
101. Toots A, Littbrand H, Lindelof N, et al. Effects of a High-Intensity Functional Exercise Program on Dependence in Activities of Daily Living and Balance in Older Adults with Dementia. *Journal of the American Geriatrics Society*. 2016 Jan;64(1):55-64. doi: <https://dx.doi.org/10.1111/jgs.13880>. PMID: 26782852.
102. Pitkala KH, Poysti MM, Laakkonen ML, et al. Effects of the Finnish Alzheimer disease exercise trial (FINALEX): a randomized controlled trial. *JAMA Internal Medicine*. 2013 May 27;173(10):894-901. doi: <https://dx.doi.org/10.1001/jamainternmed.2013.359>. PMID: 23589097.
103. Nyman SR, Ingram W, Sanders J, et al. Randomised Controlled Trial Of The Effect Of Tai Chi On Postural Balance Of People With Dementia. *Clinical Interventions In Aging*. 2019;14:2017-29. doi: <https://dx.doi.org/10.2147/CIA.S228931>. PMID: 31819385.
104. Lamb SE, Sheehan B, Atherton N, et al. Dementia And Physical Activity (DAPA) trial of moderate to high intensity exercise training for people with dementia: randomised controlled trial. *BMJ*. 2018;361:k1675. PMID: 29769247.
105. Hoffmann K, Sobol NA, Frederiksen KS, et al. Moderate-to-High Intensity Physical Exercise in Patients with Alzheimer's Disease: A Randomized Controlled Trial. *Journal of Alzheimer's Disease*. 2016;50(2):443-53. doi: 10.3233/JAD-150817. PMID: 112439259. Language: English. Entry Date: 20161122. Revision Date: 20180627. Publication Type: journal article.
106. Bossers WJ, van der Woude LH, Boersma F, et al. Comparison of Effect of Two Exercise Programs on Activities of Daily Living in Individuals with Dementia: A 9-Week Randomized, Controlled Trial. *Journal of the American Geriatrics Society*. 2016 Jun;64(6):1258-66. doi: <https://dx.doi.org/10.1111/jgs.14160>. PMID: 27321604.
107. Telenius EW, Engedal K, Bergland A. Long-term effects of a 12 weeks high-intensity functional exercise program on physical function and mental health in nursing home residents with dementia: a single blinded randomized controlled trial. *BMC Geriatrics*. 2015 Dec 03;15:158. doi: <https://dx.doi.org/10.1186/s12877-015-0151-8>. PMID: 26630910.
108. Hauer K, Schwenk M, Zieschang T, et al. Physical training improves motor performance in people with dementia: a randomized controlled trial. *Journal of the American Geriatrics Society*. 2012 Jan;60(1):8-15. doi: <https://dx.doi.org/10.1111/j.1532-5415.2011.03778.x>. PMID: 22211512.
109. Eggermont LH, Knol DL, Hol EM, et al. Hand motor activity, cognition, mood, and the rest-activity rhythm in dementia: a clustered RCT. *Behavioural Brain Research*. 2009 Jan 23;196(2):271-8. doi: <https://dx.doi.org/10.1016/j.bbr.2008.09.012>. PMID: 18926856.
110. Chen LL, Li H, Chen XH, et al. Effects of Hand Exercise on Eating Action in Patients With Alzheimer's Disease. *American Journal of Alzheimer's Disease & Other Dementias*. 2019 02;34(1):57-62. doi: <https://dx.doi.org/10.1177/1533317518803722>. PMID: 30301358.
111. Huang N, Li W, Rong X, et al. Effects of a Modified Tai Chi Program on Older People with Mild Dementia: A Randomized Controlled Trial. *Journal of Alzheimer's Disease*. 2019;72(3):947-56. doi: <https://dx.doi.org/10.3233/JAD-190487>. PMID: 31743998.

112. Cancela JM, Ayan C, Varela S, et al. Effects of a long-term aerobic exercise intervention on institutionalized patients with dementia. *Journal of Science & Medicine in Sport*. 2016 Apr;19(4):293-8. doi: <https://dx.doi.org/10.1016/j.jsams.2015.05.007>. PMID: 26087884.
113. Chen KM, Kuo CC, Chang YH, et al. Resistance Band Exercises Reduce Depression and Behavioral Problems of Wheelchair-Bound Older Adults with Dementia: A Cluster-Randomized Controlled Trial. *Journal of the American Geriatrics Society*. 2017 Feb;65(2):356-63. doi: <https://dx.doi.org/10.1111/jgs.14526>. PMID: 27879982.
114. Henskens M, Nauta IM, Drost KT, et al. The effects of movement stimulation on activities of daily living performance and quality of life in nursing home residents with dementia: a randomized controlled trial. *Clinical Interventions In Aging*. 2018;13:805-17. doi: <https://dx.doi.org/10.2147/CIA.S160031>. PMID: 29750023.
115. Ho RTH, Fong TCT, Chan WC, et al. Psychophysiological effects of Dance Movement Therapy and physical exercise on older adults with mild dementia: A randomized controlled trial. *Journals of Gerontology Series B Psychological Sciences & Social Sciences*. 2018 Nov 28;28:28. doi: <https://dx.doi.org/10.1093/geronb/gby145>. PMID: 30496547.
116. Masayuki S, Jun-ichi O, Tomoko T, et al. Physical Exercise with Music Maintains Activities of Daily Living in Patients with Dementia: Mihama-Kiho Project Part 21. *Journal of Alzheimer's Disease*. 2017;57(1):85-96. doi: 10.3233/JAD-161217. PMID: 121747018. Language: English. Entry Date: 20180302. Revision Date: 20180619. Publication Type: journal article.
117. Roach KE, Tappen RM, Kirk-Sanchez N, et al. A randomized controlled trial of an activity specific exercise program for individuals with Alzheimer disease in long-term care settings. *Journal of Geriatric Physical Therapy*. 2011 Apr-Jun;34(2):50-6. doi: <https://dx.doi.org/10.1519/JPT.0b013e31820aab9c>. PMID: 21937893.
118. Rolland Y, Pillard F, Klapouszczak A, et al. Exercise program for nursing home residents with Alzheimer's disease: a 1-year randomized, controlled trial. *Journal of the American Geriatrics Society*. 2007 Feb;55(2):158-65. PMID: 17302650.
119. Bossers WJ, Scherder EJ, Boersma F, et al. Feasibility of a combined aerobic and strength training program and its effects on cognitive and physical function in institutionalized dementia patients. A pilot study. *PLoS ONE [Electronic Resource]*. 2014;9(5):e97577. doi: <https://dx.doi.org/10.1371/journal.pone.0097577>. PMID: 24844772.
120. Bossers WJ, van der Woude LH, Boersma F, et al. A 9-Week Aerobic and Strength Training Program Improves Cognitive and Motor Function in Patients with Dementia: A Randomized, Controlled Trial. *American Journal of Geriatric Psychiatry*. 2015 Nov;23(11):1106-16. doi: <https://dx.doi.org/10.1016/j.jagp.2014.12.191>. PMID: 25648055.
121. Burge E, Berchtold A, Maupetit C, et al. Does physical exercise improve adl capacities in people over 65 years with moderate or severe dementia hospitalized in an acute psychiatric setting? A multisite randomized clinical trial. *International psychogeriatrics*. 2016. PMID: CN-01308751.
122. de Souto Barreto P, Cesari M, Denormandie P, et al. Exercise or Social Intervention for Nursing Home Residents with Dementia: A Pilot Randomized, Controlled Trial. *Journal of the American Geriatrics Society*. 2017 Sep;65(9):E123-E9. doi: <https://dx.doi.org/10.1111/jgs.14947>. PMID: 28542742.

123. Holthoff VA, Marschner K, Scharf M, et al. Effects of physical activity training in patients with Alzheimer's dementia: results of a pilot RCT study. *PLoS ONE* [Electronic Resource]. 2015;10(4):e0121478. doi: <https://dx.doi.org/10.1371/journal.pone.0121478>. PMID: 25884637.
124. Morris JK, Vidoni ED, Johnson DK, et al. Aerobic exercise for Alzheimer's disease: A randomized controlled pilot trial. *PLoS ONE* [Electronic Resource]. 2017;12(2):e0170547. doi: <https://dx.doi.org/10.1371/journal.pone.0170547>. PMID: 28187125.
125. Padala KP, Padala PR, Lensing SY, et al. Home-Based Exercise Program Improves Balance and Fear of Falling in Community-Dwelling Older Adults with Mild Alzheimer's Disease: A Pilot Study. *Journal of Alzheimer's Disease*. 2017;59(2):565-74. doi: <https://dx.doi.org/10.3233/JAD-170120>. PMID: 28655135.
126. Stevens J, Killeen M. A randomised controlled trial testing the impact of exercise on cognitive symptoms and disability of residents with dementia. *Contemporary Nurse*. 2006 Feb-Mar;21(1):32-40. PMID: 16594879.
127. Suttanon P, Hill KD, Said CM, et al. Feasibility, safety and preliminary evidence of the effectiveness of a home-based exercise programme for older people with Alzheimer's disease: a pilot randomized controlled trial. *Clinical Rehabilitation*. 2013 May;27(5):427-38. doi: <https://dx.doi.org/10.1177/0269215512460877>. PMID: 23117349.
128. Yu F, Thomas W, Nelson NW, et al. Impact of 6-month aerobic exercise on Alzheimer's symptoms. *Journal of Applied Gerontology*. 2015 Jun;34(4):484-500. doi: <https://dx.doi.org/10.1177/0733464813512895>. PMID: 24652914.
129. Canonici P, Andrade LP, Gobbi S, et al. Functional dependence and caregiver burden in Alzheimer's disease: a controlled trial on the benefits of motor intervention. *Psychogeriatrics*. 2012;12(3):186-92. doi: 10.1111/j.1479-8301.2012.00407.x. PMID: CN-00863385.
130. Dawson N, Judge KS, Gerhart H. Improved Functional Performance in Individuals With Dementia After a Moderate-Intensity Home-Based Exercise Program: A Randomized Controlled Trial. *Journal of Geriatric Physical Therapy*. 2019 Jan/Mar;42(1):18-27. doi: <https://dx.doi.org/10.1519/JPT.000000000000000128>. PMID: 28252473.
131. Henwood T, Neville C, Baguley C, et al. Aquatic exercise for residential aged care adults with dementia: benefits and barriers to participation. *International psychogeriatrics*. 2017;29(9):1439-49. doi: 10.1017/S104161021700028X. PMID: CN-01403352.
132. Kwak YS, Um SY, Son TG, et al. Effect of regular exercise on senile dementia patients. *International Journal of Sports Medicine*. 2008 Jun;29(6):471-4. PMID: 18050054.
133. Liu B, Chen X, Li Y, et al. Effect of passive finger exercises on grip strength and the ability to perform activities of daily living for older people with dementia: a 12-week randomized controlled trial. *Clinical Interventions In Aging*. 2018;13:2169-77. doi: <https://dx.doi.org/10.2147/CIA.S174756>. PMID: 30464426.
134. Nascimento C, Teixeira C, Gobbi L, et al. A controlled clinical trial on the effects of exercise on neuropsychiatric disorders and instrumental activities in women with Alzheimer's disease. *Revista brasileira de fisioterapia (sao carlos (sao paulo, brazil))*. 2012;16(3):197-204. PMID: CN-00863837.

135. Steinberg M, Leoutsakos JM, Podewils LJ, et al. Evaluation of a home-based exercise program in the treatment of Alzheimer's disease: the Maximizing Independence in Dementia (MIND) study. *International Journal of Geriatric Psychiatry*. 2009 Jul;24(7):680-5. doi: <https://dx.doi.org/10.1002/gps.2175>. PMID: 19089875.
136. Stella F, Canonici A, Gobbi S, et al. Attenuation of neuropsychiatric symptoms and caregiver burden in Alzheimer's disease by motor intervention: a controlled trial. *Clinics (sao paulo, brazil)*. 2011;66(8):1353-60. PMID: CN-00860745.
137. Toulotte C, Fabre C, Dangremont B, et al. Effects of physical training on the physical capacity of frail, demented patients with a history of falling: a randomised controlled trial. *Age & Ageing*. 2003 Jan;32(1):67-73. PMID: 12540351.
138. Van de Winckel A, Feys H, De Weerd W, et al. Cognitive and behavioural effects of music-based exercises in patients with dementia. *Clinical Rehabilitation*. 2004 May;18(3):253-60. PMID: 15137556.
139. Venturelli M, Scarsini R, Schena F. Six-month walking program changes cognitive and ADL performance in patients with Alzheimer. *American Journal of Alzheimer's Disease & Other Dementias*. 2011 Aug;26(5):381-8. doi: <https://dx.doi.org/10.1177/1533317511418956>. PMID: 21852281.
140. Vreugdenhil A, Cannell J, Davies A, et al. A community-based exercise programme to improve functional ability in people with Alzheimer's disease: a randomized controlled trial. *Scandinavian Journal of Caring Sciences*. 2012 Mar;26(1):12-9. doi: <https://dx.doi.org/10.1111/j.1471-6712.2011.00895.x>. PMID: 21564154.
141. Brett L, Stapley P, Meedya S, et al. Effect of physical exercise on physical performance and fall incidents of individuals living with dementia in nursing homes: a randomized controlled trial. *Physiotherapy Theory & Practice*. 2019 Mar 26:1-14. doi: <https://dx.doi.org/10.1080/09593985.2019.1594470>. PMID: 30912690.
142. Fan J, Chen K. Using silver yoga exercises to promote physical and mental health of elders with dementia in long-term care facilities. In *International psychogeriatrics / IPA*
143. Karssemeijer EGA, Bossers WJR, Aaronson JA, et al. Exergaming as a Physical Exercise Strategy Reduces Frailty in People With Dementia: A Randomized Controlled Trial. *Journal of the American Medical Directors Association*. 2019 Aug 10;10:10. doi: <https://dx.doi.org/10.1016/j.jamda.2019.06.026>. PMID: 31409559.
144. Lee SM, Joung J, Shin SH. Effects of Fumanet exercise on Korean older adults with mild dementia. *Japan Journal of Nursing Science: JJNS*. 2019 Jul 25:e12286. doi: <https://dx.doi.org/10.1111/jjns.12286>. PMID: 31347294.
145. Taylor ME, Close JCT, Lord SR, et al. Pilot feasibility study of a home-based fall prevention exercise program (StandingTall) delivered through a tablet computer (iPad) in older people with dementia. *Australasian journal on ageing*. 2019;19. doi: <http://dx.doi.org/10.1111/ajag.12717>. PMID: 629382297.
146. Enette L, Vogel T, Merle S, et al. Effect of 9 weeks continuous vs. interval aerobic training on plasma BDNF levels, aerobic fitness, cognitive capacity and quality of life among seniors with mild to moderate Alzheimer's disease: a randomized controlled trial. *European Reviews of Aging & Physical Activity*. 2020;17:2. doi: <https://dx.doi.org/10.1186/s11556-019-0234-1>. PMID: 31921371.

147. Fleiner T, Dauth H, Zijlstra W, et al. A Structured Physical Exercise Program Reduces Professional Caregiver's Burden Caused by Neuropsychiatric Symptoms in Acute Dementia Care: Randomized Controlled Trial Results. *Journal of Alzheimer's Disease*. 2020;74(2):429-33. doi: <https://dx.doi.org/10.3233/JAD-191102>. PMID: 32039852.
148. Liu IT, Lee WJ, Lin SY, et al. Therapeutic Effects of Exercise Training on Elderly Patients With Dementia: A Randomized Controlled Trial. *Archives of Physical Medicine & Rehabilitation*. 2020 Feb 19;19:19. doi: <https://dx.doi.org/10.1016/j.apmr.2020.01.012>. PMID: 32084347.
149. Sanders LMJ, Hortobagyi T, Karssemeijer EGA, et al. Effects of low- and high-intensity physical exercise on physical and cognitive function in older persons with dementia: a randomized controlled trial. *Alzheimer's Research & Therapy*. 2020 Mar 19;12(1):28. doi: <https://dx.doi.org/10.1186/s13195-020-00597-3>. PMID: 32192537.
150. Todri J, Lena O, Martinez Gil JL. A single blind randomized controlled trial of global postural re-education: Cognitive effects on Alzheimer disease patients. *European Journal of Psychiatry*. 2019 April - June;33(2):83-90.
151. Lamb SE, Mistry D, Alleyne S, et al. Aerobic and strength training exercise programme for cognitive impairment in people with mild to moderate dementia: The DAPA RCT. *Health Technology Assessment*. 2018 May;22(28):1-201. doi: <http://dx.doi.org/10.3310/hta22280>. PMID: 622601831.
152. Cheung DSK, Lai CKY, Wong FKY, et al. The effects of the music-with-movement intervention on the cognitive functions of people with moderate dementia: a randomized controlled trial. *Aging & Mental Health*. 2018 Mar;22(3):306-15. doi: <https://dx.doi.org/10.1080/13607863.2016.1251571>. PMID: 27819483.
153. Kwak J, Anderson K, O'Connell Valuch K. Findings From a Prospective Randomized Controlled Trial of an Individualized Music Listening Program for Persons With Dementia. *Journal of Applied Gerontology*. 2018 Jun 01;733464818778991. doi: <https://dx.doi.org/10.1177/0733464818778991>. PMID: 29871544.
154. Lin Y, Chu H, Yang CY, et al. Effectiveness of group music intervention against agitated behavior in elderly persons with dementia. *International Journal of Geriatric Psychiatry*. 2011 Jul;26(7):670-8. doi: <https://dx.doi.org/10.1002/gps.2580>. PMID: 20672256.
155. Sarkamo T, Tervaniemi M, Laitinen S, et al. Cognitive, emotional, and social benefits of regular musical activities in early dementia: randomized controlled study. *Gerontologist*. 2014 Aug;54(4):634-50. doi: <https://dx.doi.org/10.1093/geront/gnt100>. PMID: 24009169.
156. Sung HC, Lee WL, Li TL, et al. A group music intervention using percussion instruments with familiar music to reduce anxiety and agitation of institutionalized older adults with dementia. *International Journal of Geriatric Psychiatry*. 2012 Jun;27(6):621-7. doi: <https://dx.doi.org/10.1002/gps.2761>. PMID: 21823174.
157. Cheung DSK, Lai CKY, Wong FKY, et al. Is music-with-movement intervention better than music listening and social activities in alleviating agitation of people with moderate dementia? A randomized controlled trial. *Dementia*. 2018 Sep 20;1471301218800195. doi: <https://dx.doi.org/10.1177/1471301218800195>. PMID: 30235949.
158. Chu H, Yang CY, Lin Y, et al. The impact of group music therapy on depression and cognition in elderly persons with dementia: a randomized controlled study. *Biological Research for Nursing*. 2014 Apr;16(2):209-17. doi: <https://dx.doi.org/10.1177/1099800413485410>. PMID: 23639952.

159. Sarkamo T, Laitinen S, Numminen A, et al. Clinical and Demographic Factors Associated with the Cognitive and Emotional Efficacy of Regular Musical Activities in Dementia. *Journal of Alzheimer's Disease*. 2016;49(3):767-81. doi: <https://dx.doi.org/10.3233/JAD-150453>. PMID: 26519435.
160. Ceccato E, Vigato G, Bonetto C, et al. STAM protocol in dementia: a multicenter, single-blind, randomized, and controlled trial. *American Journal of Alzheimer's Disease & Other Dementias*. 2012 Aug;27(5):301-10. doi: <https://dx.doi.org/10.1177/1533317512452038>. PMID: 22815078.
161. Chang F, Huang H, Lin K, et al. The effect of a music programme during lunchtime on the problem behaviour of the older residents with dementia at an institution in Taiwan. *Journal of clinical nursing*. 2010;19(7-8):939-48. doi: 10.1111/j.1365-2702.2009.02801.x. PMID: CN-00752730.
162. Choi A, Lee M, Cheong K, et al. Effects of group music intervention on behavioral and psychological symptoms in patients with dementia: a pilot-controlled trial. *International journal of neuroscience*. 2009;119(4):471-81. doi: 10.1080/00207450802328136. PMID: CN-00688678.
163. Giovagnoli AR, Manfredi V, Schifano L, et al. Combining drug and music therapy in patients with moderate Alzheimer's disease: a randomized study. *Neurological Sciences*. 2018 Mar 17;17:17. doi: <https://dx.doi.org/10.1007/s10072-018-3316-3>. PMID: 29550981.
164. Cooke ML, Moyle W, Shum DH, et al. A randomized controlled trial exploring the effect of music on agitated behaviours and anxiety in older people with dementia. *Aging & Mental Health*. 2010 Nov;14(8):905-16. doi: <https://dx.doi.org/10.1080/13607861003713190>. PMID: 20635236.
165. Gerdner LA. Effects of individualized versus classical "relaxation" music on the frequency of agitation in elderly persons with Alzheimer's disease and related disorders. *International Psychogeriatrics*. 2000 Mar;12(1):49-65. PMID: 10798453.
166. Groene RW. Effectiveness of music therapy 1:1 intervention with individuals having Senile Dementia of the Alzheimer's Type. *Journal of Music Therapy*. 1993 Fall;30(3):138-57. doi: <http://dx.doi.org/10.1093/jmt/30.3.138>. PMID: 1994-10861-001.
167. Guetin S, Portet F, Picot MC, et al. Effect of music therapy on anxiety and depression in patients with Alzheimer's type dementia: randomised, controlled study. *Dementia & Geriatric Cognitive Disorders*. 2009;28(1):36-46. doi: <https://dx.doi.org/10.1159/000229024>. PMID: 19628939.
168. Han P, Kwan M, Chen D, et al. A controlled naturalistic study on a weekly music therapy and activity program on disruptive and depressive behaviors in dementia. *Dementia and geriatric cognitive disorders*. 2010;30(6):540-6. PMID: CN-00856782.
169. Holmes C, Knights A, Dean C, et al. Keep music live: music and the alleviation of apathy in dementia subjects. *International Psychogeriatrics*. 2006 Dec;18(4):623-30. PMID: 16805928.
170. Lord TR, Garner JE. Effects of music on Alzheimer patients. *Perceptual & Motor Skills*. 1993 Apr;76(2):451-5. PMID: 8483655.
171. McHugh L, Gardstrom S, Hiller J, et al. The effect of pre-meal, vocal re-creative music therapy on nutritional intake of residents with Alzheimer's disease and related dementias: a pilot study. *Music ther perspect*. 2012;30(1):32-42. PMID: CN-01140174.
172. Ledger A, Baker F. An investigation of long-term effects of group music therapy on agitation levels of people with Alzheimer's disease. *Aging & mental health*. 2007;11(3):330-8. PMID: CN-00738375.

173. Sakamoto M, Ando H, Tsutou A. Comparing the effects of different individualized music interventions for elderly individuals with severe dementia. *International Psychogeriatrics*. 2013 May;25(5):775-84. doi: <https://dx.doi.org/10.1017/S1041610212002256>. PMID: 23298693.
174. Raglio A, Oasi O, Gianotti M, et al. Effects of music therapy on psychological symptoms and heart rate variability in patients with dementia. A pilot study. *Current Aging Science*. 2010 Dec;3(3):242-6. PMID: 20735342.
175. Ridder HM, Stige B, Qvale LG, et al. Individual music therapy for agitation in dementia: an exploratory randomized controlled trial. *Aging & Mental Health*. 2013;17(6):667-78. doi: <https://dx.doi.org/10.1080/13607863.2013.790926>. PMID: 23621805.
176. Sung HC, Chang SM, Lee WL, et al. The effects of group music with movement intervention on agitated behaviours of institutionalized elders with dementia in Taiwan. *Complementary Therapies in Medicine*. 2006 Jun;14(2):113-9. PMID: 16765849.
177. Ugur HG, Orak OS, Aktas YY, et al. Effects of music therapy on the care burden of in-home caregivers and physiological parameters of their in-home dementia patients: a randomized controlled trial. *Complementary medicine research*. 2019;26(1):22-30. PMID: CN-01911571.
178. Svansdottir HB, Snaedal J. Music therapy in moderate and severe dementia of Alzheimer's type: a case-control study. *International Psychogeriatrics*. 2006 Dec;18(4):613-21. PMID: 16618375.
179. Ihara ES, Tompkins CJ, Inoue M, et al. Results from a person-centered music intervention for individuals living with dementia. *Geriatrics & gerontology international*. 2019 Jan;19(1):30-4. doi: <https://dx.doi.org/10.1111/ggi.13563>. PMID: 30460747.
180. Narme P, Clement S, Ehrle N, et al. Efficacy of musical interventions in dementia: evidence from a randomized controlled trial. *Journal of Alzheimer's Disease*. 2014;38(2):359-69. doi: <https://dx.doi.org/10.3233/JAD-130893>. PMID: 23969994.
181. Park J, Tolea MI, Sherman D, et al. Feasibility of Conducting Nonpharmacological Interventions to Manage Dementia Symptoms in Community-Dwelling Older Adults: A Cluster Randomized Controlled Trial. *American Journal of Alzheimer's Disease & Other Dementias*. 2019 Sep 18:1533317519872635. doi: <https://dx.doi.org/10.1177/1533317519872635>. PMID: 31533443.
182. Pongan E, Tillmann B, Leveque Y, et al. Can Musical or Painting Interventions Improve Chronic Pain, Mood, Quality of Life, and Cognition in Patients with Mild Alzheimer's Disease? Evidence from a Randomized Controlled Trial. *Journal of Alzheimer's Disease*. 2017;60(2):663-77. doi: <https://dx.doi.org/10.3233/JAD-170410>. PMID: 28922159.
183. Weise L, Töpfer NF, Deux J, et al. Feasibility and effects of individualized recorded music for people with dementia: A pilot RCT study. *Nordic Journal of Music Therapy*. 2020;29(1):39-56. doi: 10.1080/08098131.2019.1661507. PMID: 141134381. Language: English. Entry Date: 20200117. Revision Date: 20200117. Publication Type: Article.
184. Ho RT, Fong TC, Sing CY, et al. Managing behavioral and psychological symptoms in Chinese elderly with dementia via group-based music intervention: A cluster randomized controlled trial. *Dementia*. 2018 Jan 01:1471301218760023. doi: <https://dx.doi.org/10.1177/1471301218760023>. PMID: 29468887.

185. Raglio A, Bellelli G, Traficante D, et al. Efficacy of music therapy in the treatment of behavioral and psychiatric symptoms of dementia. *Alzheimer Disease & Associated Disorders*. 2008 Apr-Jun;22(2):158-62. doi: <https://dx.doi.org/10.1097/WAD.0b013e3181630b6f>. PMID: 18525288.
186. Raglio A, Bellelli G, Traficante D, et al. Addendum to 'Efficacy of music therapy treatment based on cycles of sessions: A randomised controlled trial' (Raglio et al., 2010). *Aging & Mental Health*. 2012 Mar;16(2):265-7. doi: <http://dx.doi.org/10.1080/13607863.2011.630376>. PMID: 2012-00529-017.
187. Vink AC, Zuidersma M, Boersma F, et al. The effect of music therapy compared with general recreational activities in reducing agitation in people with dementia: a randomised controlled trial. *International Journal of Geriatric Psychiatry*. 2013 Oct;28(10):1031-8. doi: <https://dx.doi.org/10.1002/gps.3924>. PMID: 23280604.
188. Wang SC, Yu CL, Chang SH. Effect of music care on depression and behavioral problems in elderly people with dementia in Taiwan: a quasi-experimental, longitudinal study. *Aging & Mental Health*. 2017 02;21(2):156-62. doi: <https://dx.doi.org/10.1080/13607863.2015.1093602>. PMID: 26443002.
189. Hsu CN, Lin LC, Wu SC. The effects of spaced retrieval training in improving hyperphagia of people living with dementia in residential settings. *Journal of Clinical Nursing*. 2017 Oct;26(19-20):3224-31. doi: <https://dx.doi.org/10.1111/jocn.13671>. PMID: 27878873.
190. Cooke M, Moyle W, Shum D, et al. A randomized controlled trial exploring the effect of music on quality of life and depression in older people with dementia. *Journal of Health Psychology*. 2010 Jul;15(5):765-76. doi: <https://dx.doi.org/10.1177/1359105310368188>. PMID: 20603300.
191. Giovagnoli AR, Manfredi V, Parente A, et al. Cognitive training in Alzheimer's disease: a controlled randomized study. *Neurological Sciences*. 2017 Aug;38(8):1485-93. doi: <https://dx.doi.org/10.1007/s10072-017-3003-9>. PMID: 28577267.
192. Raglio A, Bellelli G, Traficante D, et al. Efficacy of music therapy treatment based on cycles of sessions: a randomised controlled trial. *Aging & Mental Health*. 2010 Nov;14(8):900-4. doi: <https://dx.doi.org/10.1080/13607861003713158>. PMID: 21069596.
193. Amieva H, Robert PH, Grandoulier AS, et al. Group and individual cognitive therapies in Alzheimer's disease: the ETNA3 randomized trial. *International Psychogeriatrics*. 2016 May;28(5):707-17. doi: <https://dx.doi.org/10.1017/S1041610215001830>. PMID: 26572551.
194. Li M, Lyu JH, Zhang Y, et al. Efficacy of Group Reminiscence Therapy on Cognition, Depression, Neuropsychiatric Symptoms, and Activities of Daily Living for Patients With Alzheimer Disease. *Journal of Geriatric Psychiatry & Neurology*. 2019 Oct 23;891988719882099. doi: <https://dx.doi.org/10.1177/0891988719882099>. PMID: 31645180.
195. Wang J, Yen M, OuYang W. Group reminiscence intervention in Taiwanese elders with dementia. In *Archives of gerontology and geriatrics*
196. Wang JJ. Group reminiscence therapy for cognitive and affective function of demented elderly in Taiwan. *International Journal of Geriatric Psychiatry*. 2007 Dec;22(12):1235-40. PMID: 17503545.
197. Woods RT, Bruce E, Edwards RT, et al. Reminiscence groups for people with dementia and their family carers: pragmatic eight-centre randomised trial of joint reminiscence and maintenance versus usual treatment: a protocol. *Trials [Electronic Resource]*. 2009 Jul 30;10:64. doi: <https://dx.doi.org/10.1186/1745-6215-10-64>. PMID: 19642992.

198. Woods RT, Orrell M, Bruce E, et al. REMCARE: Pragmatic Multi-Centre Randomised Trial of Reminiscence Groups for People with Dementia and their Family Carers: Effectiveness and Economic Analysis. PLoS ONE [Electronic Resource]. 2016;11(4):e0152843. doi: <https://dx.doi.org/10.1371/journal.pone.0152843>. PMID: 27093052.
199. Gonzalez J, Mayordomo T, Torres M, et al. Reminiscence and dementia: a therapeutic intervention. *International Psychogeriatrics*. 2015 Oct;27(10):1731-7. doi: <https://dx.doi.org/10.1017/S1041610215000344>. PMID: 25765779.
200. Laird EA, Ryan A, McCauley C, et al. Using Mobile Technology to Provide Personalized Reminiscence for People Living With Dementia and Their Carers: Appraisal of Outcomes From a Quasi-Experimental Study. *JMIR Mental Health*. 2018 Sep 11;5(3):e57. doi: <https://dx.doi.org/10.2196/mental.9684>. PMID: 30206053.
201. Lalanne J, Gallarda T, Piolino P. "The Castle of Remembrance": New insights from a cognitive training programme for autobiographical memory in Alzheimer's disease. *Neuropsychological Rehabilitation*. 2015;25(2):254-82. doi: <https://dx.doi.org/10.1080/09602011.2014.949276>. PMID: 25122521.
202. Manav AI, Simsek N. The effect of reminiscence therapy with internet-based videos on cognitive status and apathy of older people with mild dementia. *Journal of geriatric psychiatry and neurology*. 2019;32(2):104-13. PMID: CN-01707450.
203. Martinez N, Stutzman SE, Olson DM. Electronic interventions aimed at increasing self-worth in mild dementia may not be feasible. *Journal of the American Association of Nurse Practitioners*. 2019 Aug 22;22:22. doi: <https://dx.doi.org/10.1097/JXX.0000000000000280>. PMID: 31453826.
204. Nakamae T, Yotsumoto K, Tatsumi E, et al. Effects of productive activities with reminiscence in occupational therapy for people with dementia: A pilot randomized controlled study. *Hong Kong Journal of Occupational Therapy*. 2014 June;24(1):13-9. doi: <http://dx.doi.org/10.1016/j.hkjot.2014.01.003>. PMID: 53104158.
205. Politis AM, Vozzella S, Mayer LS, et al. A randomized, controlled, clinical trial of activity therapy for apathy in patients with dementia residing in long-term care. *International Journal of Geriatric Psychiatry*. 2004 Nov;19(11):1087-94. PMID: 15481065.
206. Subramaniam P, Woods B, Whitaker C. Life review and life story books for people with mild to moderate dementia: a randomised controlled trial. *Aging & Mental Health*. 2014;18(3):363-75. doi: <https://dx.doi.org/10.1080/13607863.2013.837144>. PMID: 24063317.
207. Van Bogaert P, Van Grinsven R, Tolson D, et al. Effects of SolCos model-based individual reminiscence on older adults with mild to moderate dementia due to Alzheimer disease: a pilot study. *Journal of the American Medical Directors Association*. 2013 Jul;14(7):528.e9-13. doi: <https://dx.doi.org/10.1016/j.jamda.2013.01.020>. PMID: 23583001.
208. Camberg L, Woods P, Ooi WL, et al. Evaluation of Simulated Presence: a personalized approach to enhance well-being in persons with Alzheimer's disease. *Journal of the American Geriatrics Society*. 1999 Apr;47(4):446-52. PMID: 10203120.
209. Duru Asiret G, Kapucu S. The Effect of Reminiscence Therapy on Cognition, Depression, and Activities of Daily Living for Patients With Alzheimer Disease. *Journal of Geriatric Psychiatry & Neurology*. 2016 Jan;29(1):31-7. doi: <https://dx.doi.org/10.1177/0891988715598233>. PMID: 26251112.

210. Hsieh C-J, Chang C, Su S-F, et al. Reminiscence group therapy on depression and apathy in nursing home residents with mild-to-moderate dementia. In *Journal of experimental and clinical medicine*
211. Lai CK, Chi I, Kayser-Jones J. A randomized controlled trial of a specific reminiscence approach to promote the well-being of nursing home residents with dementia. *International Psychogeriatrics*. 2004 Mar;16(1):33-49. PMID: 15190995.
212. Lin H-C, Yang Y-P, Cheng W-Y, et al. Distinctive effects between cognitive stimulation and reminiscence therapy on cognitive function and quality of life for different types of behavioural problems in dementia. In *Scandinavian journal of caring sciences*
213. Lok N, Bademli K, Selcuk-Tosun A. The effect of reminiscence therapy on cognitive functions, depression, and quality of life in Alzheimer patients: Randomized controlled trial. *International Journal of Geriatric Psychiatry*. 2019 Jan;34(1):47-53. doi: <https://dx.doi.org/10.1002/gps.4980>. PMID: 30246408.
214. O'Shea E, Devane D, Cooney A, et al. The impact of reminiscence on the quality of life of residents with dementia in long-stay care. *International Journal of Geriatric Psychiatry*. 2014 Oct;29(10):1062-70. doi: <https://dx.doi.org/10.1002/gps.4099>. PMID: 24633858.
215. Serrani Azcurra DJ. A reminiscence program intervention to improve the quality of life of long-term care residents with Alzheimer's disease: a randomized controlled trial. *Revista Brasileira de Psiquiatria*. 2012 Dec;34(4):422-33. PMID: 23429813.
216. Van Bogaert P, Tolson D, Eerlingen R, et al. SolCos model-based individual reminiscence for older adults with mild to moderate dementia in nursing homes: a randomized controlled intervention study. *Journal of Psychiatric & Mental Health Nursing*. 2016 Nov;23(9-10):568-75. doi: <https://dx.doi.org/10.1111/jpm.12336>. PMID: 27511740.
217. Wu LF, Koo M. Randomized controlled trial of a six-week spiritual reminiscence intervention on hope, life satisfaction, and spiritual well-being in elderly with mild and moderate dementia. *International Journal of Geriatric Psychiatry*. 2016 Feb;31(2):120-7. doi: <https://dx.doi.org/10.1002/gps.4300>. PMID: 25965388.
218. Ching-Teng Y, Ya-Ping Y, Chia-Ju L, et al. Effect of group reminiscence therapy on depression and perceived meaning of life of veterans diagnosed with dementia at veteran homes. *Social Work in Health Care*. 2020;59(2):75-90. doi: 10.1080/00981389.2019.1710320. PMID: 141877576. Language: English. Entry Date: 20200227. Revision Date: 20200319. Publication Type: Article.
219. Inel Manav A, Simsek N. The Effect of Reminiscence Therapy With Internet-Based Videos on Cognitive Status and Apathy of Older People With Mild Dementia. *Journal of Geriatric Psychiatry & Neurology*. 2019 Mar;32(2):104-13. doi: <https://dx.doi.org/10.1177/0891988718819864>. PMID: 30612511.
220. Seyun K. Cognitive rehabilitation for elderly people with early-stage Alzheimer's disease. *Journal of Physical Therapy Science*. 2015;27(2):543-6. doi: 10.1589/jpts.27.543. PMID: 103780951. Language: English. Entry Date: 20150327. Revision Date: 20150820. Publication Type: Journal Article.

221. Clare L, Linden DE, Woods RT, et al. Goal-oriented cognitive rehabilitation for people with early-stage Alzheimer disease: a single-blind randomized controlled trial of clinical efficacy. *American Journal of Geriatric Psychiatry*. 2010 Oct;18(10):928-39. doi: <https://dx.doi.org/10.1097/JGP.0b013e3181d5792a>. PMID: 20808145.
222. Thivierge S, Jean L, Simard M. A randomized cross-over controlled study on cognitive rehabilitation of instrumental activities of daily living in Alzheimer disease. *American Journal of Geriatric Psychiatry*. 2014 Nov;22(11):1188-99. doi: <https://dx.doi.org/10.1016/j.jagp.2013.03.008>. PMID: 23871120.
223. Rogers JC, Holm MB, Burgio LD, et al. Improving morning care routines of nursing home residents with dementia. *Journal of the American Geriatrics Society*. 1999;1049-57. PMID: 107127490. Language: English. Entry Date: 20000801. Revision Date: 20150711. Publication Type: Journal Article.
224. Voigt-Radloff S, de Werd MM, Leonhart R, et al. Structured relearning of activities of daily living in dementia: the randomized controlled REDALI-DEM trial on errorless learning. *Alzheimer's Research & Therapy*. 2017 Mar 23;9(1):22. doi: <https://dx.doi.org/10.1186/s13195-017-0247-9>. PMID: 28335810.
225. Kumar P, Tiwari SC, Goel A, et al. Novel occupational therapy interventions may improve quality of life in older adults with dementia. *International archives of medicine*. 2014;7:26. doi: <https://dx.doi.org/10.1186/1755-7682-7-26>. PMID: 24982692.
226. Quayhagen MP, Quayhagen M, Corbeil RR, et al. A dyadic remediation program for care recipients with dementia. *Nursing Research*. 1995 May-Jun;44(3):153-9. PMID: 7761291.
227. Lin LC, Huang YJ, Su SG, et al. Using spaced retrieval and Montessori-based activities in improving eating ability for residents with dementia. *International Journal of Geriatric Psychiatry*. 2010 Oct;25(10):953-9. doi: <https://dx.doi.org/10.1002/gps.2433>. PMID: 20054841.
228. Lam LC, Lui VW, Luk DN, et al. Effectiveness of an individualized functional training program on affective disturbances and functional skills in mild and moderate dementia--a randomized control trial. *International Journal of Geriatric Psychiatry*. 2010 Feb;25(2):133-41. doi: <https://dx.doi.org/10.1002/gps.2309>. PMID: 19582757.
229. van der Ploeg ES, Eppingstall B, Camp CJ, et al. A randomized crossover trial to study the effect of personalized, one-to-one interaction using Montessori-based activities on agitation, affect, and engagement in nursing home residents with Dementia. *International Psychogeriatrics*. 2013 Apr;25(4):565-75. doi: <https://dx.doi.org/10.1017/S1041610212002128>. PMID: 23237211.
230. Loewenstein DA, Acevedo A, Czaja SJ, et al. Cognitive rehabilitation of mildly impaired Alzheimer disease patients on cholinesterase inhibitors. *American Journal of Geriatric Psychiatry*. 2004 Jul-Aug;12(4):395-402. PMID: 15249277.
231. Lin LC, Huang YJ, Watson R, et al. Using a Montessori method to increase eating ability for institutionalised residents with dementia: a crossover design. *Journal of Clinical Nursing*. 2011 Nov;20(21-22):3092-101. doi: <https://dx.doi.org/10.1111/j.1365-2702.2011.03858.x>. PMID: 21981704.
232. Silva AR, Pinho MS, Macedo L, et al. It is not only memory: effects of sensecam on improving well-being in patients with mild alzheimer disease. *International Psychogeriatrics*. 2017 May;29(5):741-54. doi: <https://dx.doi.org/10.1017/S104161021600243X>. PMID: 28124633.

233. Tanaka S, Honda S, Nakano H, et al. Comparison between group and personal rehabilitation for dementia in a geriatric health service facility: single-blinded randomized controlled study. *Psychogeriatrics: The Official Journal of the Japanese Psychogeriatric Society*. 2017 May;17(3):177-85. doi: <https://dx.doi.org/10.1111/psyg.12212>. PMID: 27612310.
234. Kim S. Cognitive rehabilitation for elderly people with early-stage Alzheimer's disease. *Journal of Physical Therapy Science*. 2015 Feb;27(2):543-6. doi: <https://dx.doi.org/10.1589/jpts.27.543>. PMID: 25729212.
235. Tappen RM. The effect of skill training on functional abilities of nursing home residents with dementia. *Research in Nursing & Health*. 1994 Jun;17(3):159-65. PMID: 8184127.
236. Silva AR, Pinho MS, Macedo L, et al. The cognitive effects of wearable cameras in mild Alzheimer disease-An experimental study. *Current Alzheimer Research*. 2017;14(12):1270-82. doi: <http://dx.doi.org/10.2174/1567205014666170531083015>. PMID: 2018-01500-004.
237. Brunelle-Hamann L, Thivierge S, Simard M. Impact of a cognitive rehabilitation intervention on neuropsychiatric symptoms in mild to moderate Alzheimer's disease. *Neuropsychological Rehabilitation*. 2015;25(5):677-707. doi: <https://dx.doi.org/10.1080/09602011.2014.964731>. PMID: 25312605.
238. Voigt-Radloff S, Leonhart R, Rikkert MO, et al. Study protocol of the multi-site randomised controlled REDALI-DEM trial--the effects of structured relearning methods on daily living task performance of persons with dementia. *BMC Geriatrics*. 2011 Aug 18;11:44. doi: <https://dx.doi.org/10.1186/1471-2318-11-44>. PMID: 21851594.
239. Chen Z, Yu H. Effect of intervention guided by montessori method on improving feeding capacity of patients with dementia. *International Journal of Clinical and Experimental Medicine*. 2020;13(2):1148-55.
240. Mbakile-Mahlanza L, van der Ploeg ES, Busija L, et al. A cluster-randomized crossover trial of Montessori activities delivered by family carers to nursing home residents with behavioral and psychological symptoms of dementia. *International Psychogeriatrics*. 2020 Mar;32(3):347-58. doi: <https://dx.doi.org/10.1017/S1041610219001819>. PMID: 31762434.
241. Wu HS, Lin LC, Wu SC, et al. The effectiveness of spaced retrieval combined with Montessori-based activities in improving the eating ability of residents with dementia. *Journal of Advanced Nursing*. 2014 Aug;70(8):1891-901. doi: <https://dx.doi.org/10.1111/jan.12352>. PMID: 24444172.
242. Kurz A, Thone-Otto A, Cramer B, et al. CORDIAL: cognitive rehabilitation and cognitive-behavioral treatment for early dementia in Alzheimer disease: a multicenter, randomized, controlled trial. *Alzheimer Disease & Associated Disorders*. 2012 Jul-Sep;26(3):246-53. PMID: 21986341.
243. Tsuchiya K, Yamaguchi T, Fujita T, et al. A Quasi-Randomized Controlled Trial of Brain-Activating Rehabilitation in an Acute Hospital. *American Journal of Alzheimer's Disease & Other Dementias*. 2016 Dec;31(8):612-7. PMID: 27303064.
244. Clare L, Kudlicka A, Oyebode JR, et al. Individual goal-oriented cognitive rehabilitation to improve everyday functioning for people with early-stage dementia: A multicentre randomised controlled trial (the GREAT trial). *International Journal of Geriatric Psychiatry*. 2019 May;34(5):709-21. doi: <http://dx.doi.org/10.1002/gps.5076>. PMID: 626609676.
245. Yuen I, Kwok T. Effectiveness of DementiaAbility Methods: The Montessori Way on agitation in long-term care home residents with dementia in Hong Kong. *International Journal of Geriatric Psychiatry*. 2019;34(9):1352-8. doi: <http://dx.doi.org/10.1002/gps.5063>. PMID: 626609449.

246. Cavallo M, Hunter EM, van der Hiele K, et al. Computerized Structured Cognitive Training in Patients Affected by Early-Stage Alzheimer's Disease is Feasible and Effective: A Randomized Controlled Study. *Archives of Clinical Neuropsychology*. 2016 Sep 06;06:06. PMID: 27600448.
247. Kallio EL, Ohman H, Carlson S, et al. Feasibility and baseline findings of a Finnish cognitive training (FINCOG) intervention in a randomised controlled trial among community-dwelling persons with dementia. *European Geriatric Medicine*. 2017 July;8(3):245-9. doi: <http://dx.doi.org/10.1016/j.eurger.2017.04.008>. PMID: 616108233.
248. Kallio EL, Ohman H, Hietanen M, et al. Effects of Cognitive Training on Cognition and Quality of Life of Older Persons with Dementia. *Journal of the American Geriatrics Society*. 2018 Apr;66(4):664-70. doi: <https://dx.doi.org/10.1111/jgs.15196>. PMID: 29345724.
249. Kao CC, Lin LC, Wu SC, et al. Effectiveness of different memory training programs on improving hyperphagic behaviors of residents with dementia: a longitudinal single-blind study. *Clinical Interventions In Aging*. 2016;11:707-20. doi: <https://dx.doi.org/10.2147/CIA.S102027>. PMID: 27307717.
250. Bergamaschi S, Arcara G, Calza A, et al. One-year repeated cycles of cognitive training (CT) for Alzheimer's disease. *Aging-Clinical & Experimental Research*. 2013 Aug;25(4):421-6. doi: <https://dx.doi.org/10.1007/s40520-013-0065-2>. PMID: 23784727.
251. Davis RN, Massman PJ, Doody RS. Cognitive intervention in Alzheimer disease: a randomized placebo-controlled study. *Alzheimer Disease & Associated Disorders*. 2001 Jan-Mar;15(1):1-9. PMID: 11236819.
252. De Luca R, Bramanti A, De Cola MC, et al. Cognitive training for patients with dementia living in a sicilian nursing home: a novel web-based approach. *Neurological Sciences*. 2016 01 Oct;37(10):1685-91. doi: <http://dx.doi.org/10.1007/s10072-016-2659-x>. PMID: 611200073.
253. Giuli C, Papa R, Lattanzio F, et al. The Effects of Cognitive Training for Elderly: Results from My Mind Project. *Rejuvenation Research*. 2016 Dec;19(6):485-94. PMID: 26952713.
254. Kawashima R, Hiller D, Sereda S, et al. SAIDO Learning as a Cognitive Intervention for Dementia Care: a Preliminary Study. In *Journal of the American Medical Directors Association*.
255. Mapelli D, Di Rosa E, Nocita R, et al. Cognitive stimulation in patients with dementia: randomized controlled trial. *Dementia and Geriatric Cognitive Disorders Extra*. 2013;3(1):263-71. doi: <https://dx.doi.org/10.1159/000353457>. PMID: 24052800.
256. McGilton KS, Rivera TM, Dawson P. Can we help persons with dementia find their way in a new environment?.[Erratum appears in *Aging Ment Health*. 2003 Nov;7(6):following author index]. *Aging & Mental Health*. 2003 Sep;7(5):363-71. PMID: 12959805.
257. Tarraga L, Boada M, Modinos G, et al. A randomised pilot study to assess the efficacy of an interactive, multimedia tool of cognitive stimulation in Alzheimer's disease. *Journal of Neurology, Neurosurgery & Psychiatry*. 2006 Oct;77(10):1116-21. PMID: 16820420.
258. Tseng CF, Lee SH, Hsieh TC, et al. Smart Restored by Learning Exercise Alleviates the Deterioration of Cognitive Function in Older Adults with Dementia-A Quasi-Experimental Research. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2019 04 09;16(7):09. doi: <https://dx.doi.org/10.3390/ijerph16071270>. PMID: 30970666.

259. Venturelli M, Sollima A, Ce E, et al. Effectiveness of Exercise- and Cognitive-Based Treatments on Salivary Cortisol Levels and Sundowning Syndrome Symptoms in Patients with Alzheimer's Disease. *Journal of Alzheimer's Disease*. 2016;53(4):1631-40. doi: <http://dx.doi.org/10.3233/JAD-160392>. PMID: 611692625.
260. Zarit SH, Zarit JM, Reever KE. Memory training for severe memory loss: Effects on senile dementia patients and their families. *Gerontologist*. 1982;22(4):373-7. PMID: 13238424.
261. Chen HM, Tsai LJ, Chao SY, et al. Study on the Effects of Individualized Learning Therapy on Cognitive Function and Behavioral and Psychological Symptoms of Dementia in the Institutionalized Older Adults. *Journal of Nursing Research*. 2016 Dec;24(4):300-10. PMID: 27846102.
262. Knapp M, Thorgrimsen L, Patel A, et al. Cognitive stimulation therapy for people with dementia: cost-effectiveness analysis. *British Journal of Psychiatry*. 2006 Jun;188:574-80. PMID: 16738349.
263. Orrell M, Aguirre E, Spector A, et al. Maintenance cognitive stimulation therapy for dementia: single-blind, multicentre, pragmatic randomised controlled trial. *British Journal of Psychiatry*. 2014 Jun;204(6):454-61. doi: <https://dx.doi.org/10.1192/bjp.bp.113.137414>. PMID: 24676963.
264. Spector A, Thorgrimsen L, Woods B, et al. Efficacy of an evidence-based cognitive stimulation therapy programme for people with dementia: randomised controlled trial. *British Journal of Psychiatry*. 2003 Sep;183:248-54. PMID: 12948999.
265. Capotosto E, Belacchi C, Gardini S, et al. Cognitive stimulation therapy in the Italian context: Its efficacy in cognitive and non-cognitive measures in older adults with dementia. *International Journal of Geriatric Psychiatry*. 2017 Mar;32(3):331-40. doi: <http://dx.doi.org/10.1002/gps.4521>. PMID: 2017-05902-002.
266. Cove J, Jacobi N, Donovan H, et al. Effectiveness of weekly cognitive stimulation therapy for people with dementia and the additional impact of enhancing cognitive stimulation therapy with a carer training program. *Clinical Interventions In Aging*. 2014;9:2143-50. doi: <https://dx.doi.org/10.2147/CIA.S66232>. PMID: 25525349.
267. Matsuda O. Cognitive stimulation therapy for Alzheimer's disease: the effect of cognitive stimulation therapy on the progression of mild Alzheimer's disease in patients treated with donepezil. *International psychogeriatrics*. 2007;19(2):241-52. doi: 10.1017/S1041610206004194. PMID: CN-00640778.
268. Niu YX, Tan JP, Guan JQ, et al. Cognitive stimulation therapy in the treatment of neuropsychiatric symptoms in Alzheimer's disease: a randomized controlled trial. *Clinical Rehabilitation*. 2010 Dec;24(12):1102-11. doi: <https://dx.doi.org/10.1177/0269215510376004>. PMID: 20713437.
269. Orgeta V, Leung P, Yates L, et al. Individual cognitive stimulation therapy for dementia: a clinical effectiveness and cost-effectiveness pragmatic, multicentre, randomised controlled trial. *Health Technology Assessment (Winchester, England)*. 2015 Aug;19(64):1-108. doi: <https://dx.doi.org/10.3310/hta19640>. PMID: 26292178.

270. Orrell M, Yates L, Leung P, et al. The impact of individual Cognitive Stimulation Therapy (iCST) on cognition, quality of life, caregiver health, and family relationships in dementia: A randomised controlled trial. *PLoS Medicine / Public Library of Science*. 2017 Mar;14(3):e1002269. doi: <https://dx.doi.org/10.1371/journal.pmed.1002269>. PMID: 28350796.
271. Lok N, Buldukoglu K, Barcin E. Effects of the cognitive stimulation therapy based on Roy's adaptation model on Alzheimer's patients' cognitive functions, coping-adaptation skills, and quality of life: A randomized controlled trial. *Perspectives in Psychiatric Care*. 2020 Jan 12;12:12. doi: <https://dx.doi.org/10.1111/ppc.12472>. PMID: 31930518.
272. Cheung DSK, Li B, Lai DWL, et al. Cognitive Stimulating Play Intervention for Dementia: A Feasibility Randomized Controlled Trial. *American Journal of Alzheimer's Disease & Other Dementias*. 2019 02;34(1):63-71. doi: <https://dx.doi.org/10.1177/1533317518808036>. PMID: 30370782.
273. Wong G, Yek O, Zhang A, et al. Cultural adaptation of cognitive stimulation therapy (CST) for Chinese people with dementia: multicentre pilot study. In *International journal of geriatric psychiatry*
274. Low LF, Brodaty H, Goodenough B, et al. The Sydney Multisite Intervention of LaughterBosses and ElderClowns (SMILE) study: cluster randomised trial of humour therapy in nursing homes. *BMJ Open*. 2013 Jan 11;3(1):11. doi: <https://dx.doi.org/10.1136/bmjopen-2012-002072>. PMID: 23315520.
275. Pedrinolla A, Tamburin S, Brasioli A, et al. An Indoor Therapeutic Garden for Behavioral Symptoms in Alzheimer's Disease: A Randomized Controlled Trial. *Journal of Alzheimer's Disease*. 2019;71(3):813-23. doi: <https://dx.doi.org/10.3233/JAD-190394>. PMID: 31424399.
276. Kolanowski A, Litaker M, Buettner L, et al. A randomized clinical trial of theory-based activities for the behavioral symptoms of dementia in nursing home residents. *Journal of the American Geriatrics Society*. 2011 Jun;59(6):1032-41. doi: <https://dx.doi.org/10.1111/j.1532-5415.2011.03449.x>. PMID: 21649633.
277. Buettner LL, Fitzsimmons S. AD-venture program: therapeutic biking for the treatment of depression in long-term care residents with dementia. *American Journal of Alzheimer's Disease & Other Dementias*. 2002 Mar-Apr;17(2):121-7. PMID: 11954670.
278. Hattori H, Hattori C, Hokao C, et al. Controlled study on the cognitive and psychological effect of coloring and drawing in mild Alzheimer's disease patients. *Geriatrics & gerontology international*. 2011 Oct;11(4):431-7. doi: <https://dx.doi.org/10.1111/j.1447-0594.2011.00698.x>. PMID: 21518170.
279. Hopman-Rock M, Staats PGM, Tak ECPM, et al. The effects of a psychomotor activation programme for use in groups of cognitively impaired people in homes for the elderly. *International Journal of Geriatric Psychiatry*. 1999 August;14(8):633-42. doi: <http://dx.doi.org/10.1002/%28SICI%291099-1166%28199908%2914:8%3C633::AID-GPS996%3E3.0.CO;2-W>. PMID: 29403282.
280. Li D, Li X. The effect of folk recreation program in improving symptoms: a study of Chinese elder dementia patients. *International journal of geriatric psychiatry*. 2017;32(8):901-8. doi: 10.1002/gps.4543. PMID: CN-01628262.
281. Lin Q, Cao Y, Gao J. The impacts of a gO-game (Chinese chess) intervention on alzheimer disease in a Northeast Chinese population. *Frontiers in Aging Neuroscience*. 2015;7 (AUG) (no pagination)(163). doi: <http://dx.doi.org/10.3389/fnagi.2015.00163>. PMID: 607203463.

282. Pongan E, Delphin-Combe F, Krolak-Salmon P, et al. Immediate Benefit of Art on Pain and Well-Being in Community-Dwelling Patients with Mild Alzheimer's. *American Journal of Alzheimer's Disease & Other Dementias*. 2019 Jul 09;1533317519859202. doi: <https://dx.doi.org/10.1177/1533317519859202>. PMID: 31288544.
283. Schall A, Tesky VA, Adams AK, et al. Art museum-based intervention to promote emotional well-being and improve quality of life in people with dementia: The ARTEMIS project. *Dementia*. 2017 Jan 01;1471301217730451. doi: <https://dx.doi.org/10.1177/1471301217730451>. PMID: 28914089.
284. Shalek M, Richeson NE, Buettner LL. Air mat therapy for treatment of agitated wandering: an evidence-based recreational therapy intervention. *American Journal of Recreation Therapy*. 2004 2004 Spring;3(2):18-26. PMID: 106658647. Language: English. Entry Date: 20041105. Revision Date: 20150818. Publication Type: Journal Article.
285. Tse MMY, Lau JL, Kwan R, et al. Effects of play activities program for nursing home residents with dementia on pain and psychological well-being: Cluster randomized controlled trial. *Geriatrics & gerontology international*. 2018 Oct;18(10):1485-90. doi: <https://dx.doi.org/10.1111/ggi.13509>. PMID: 30255637.
286. Laksono WAW, Haryanto J, Bahiyah K. The impact of chess games towards comfortableness of cognitive mind on elderly. *Indian Journal of Public Health Research and Development*. 2019 September;10(9):873-7.
287. Lathifah MN, Haryanto J, Fauziningtyas R. The influence of dhakonan games to impede dementia in elders. *Indian Journal of Public Health Research and Development*. 2019 September;10(9):1440-4.
288. Van Haitsma KS, Curyto K, Abbott KM, et al. A randomized controlled trial for an individualized positive psychosocial intervention for the affective and behavioral symptoms of dementia in nursing home residents. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences*. 2015 Jan;70(1):35-45. doi: <https://dx.doi.org/10.1093/geronb/gbt102>. PMID: 24304555.
289. Burns A, Guthrie E, Marino-Francis F, et al. Brief psychotherapy in Alzheimer's disease: randomised controlled trial. *British Journal of Psychiatry*. 2005 Aug;187:143-7. PMID: 16055825.
290. Cohen CI, Hyland K, Devlin M. An evaluation of the use of the natural helping network model to enhance the well-being of nursing home residents. *Gerontologist*. 1999;39(4):426-33. PMID: 107227653. Language: English. Entry Date: 19991201. Revision Date: 20150711. Publication Type: Journal Article.
291. Quinn C, Anderson D, Toms G, et al. Self-management in early-stage dementia: a pilot randomised controlled trial of the efficacy and cost-effectiveness of a self-management group intervention (the SMART study). *Trials [Electronic Resource]*. 2014 Mar 08;15:74. doi: <https://dx.doi.org/10.1186/1745-6215-15-74>. PMID: 24606601.
292. Quinn C, Toms G, Jones C, et al. A pilot randomized controlled trial of a self-management group intervention for people with early-stage dementia (The SMART study). *International Psychogeriatrics*. 2016 May;28(5):787-800. doi: <https://dx.doi.org/10.1017/S1041610215002094>. PMID: 26674087.
293. Tappen RM, Williams CL. Therapeutic conversation to improve mood in nursing home residents with Alzheimer's disease. *Research in Gerontological Nursing*. 2009 Oct;2(4):267-75. doi: <https://dx.doi.org/10.3928/19404921-20090428-02>. PMID: 20077983.

294. Young DK, Kwok TC, Ng PY. A single blind randomized control trial on support groups for Chinese persons with mild dementia. *Clinical Interventions In Aging*. 2014;9:2105-12. doi: <https://dx.doi.org/10.2147/CIA.S68687>. PMID: 25587218.
295. Wallis GG, Baldwin M, Higginbotham P. Reality orientation therapy--a controlled trial. *British Journal of Medical Psychology*. 1983 Sep;56(Pt 3):271-7. PMID: 6354248.
296. Onder G, Zanetti O, Giacobini E, et al. Reality orientation therapy combined with cholinesterase inhibitors in Alzheimer's disease: randomised controlled trial. *British Journal of Psychiatry*. 2005 Nov;187:450-5. PMID: 16260821.
297. Fritsch T, Kwak J, Grant S, et al. Impact of TimeSlips, a Creative Expression Intervention Program, on Nursing Home Residents with Dementia and their Caregivers. *Gerontologist*. 2009;49(1):117-27. doi: 10.1093/geront/gnp008. PMID: CN-00755001.
298. Houser WS, George DR, Chinchilli VM. Impact of TimeSlips creative expression program on behavioral symptoms and psychotropic medication use in persons with dementia in long-term care: a cluster-randomized pilot study. *American Journal of Geriatric Psychiatry*. 2014 Apr;22(4):337-40. doi: <https://dx.doi.org/10.1016/j.jagp.2012.12.005>. PMID: 23602304.
299. Lin R, Chen HY, Li H, et al. Effects of creative expression therapy on Chinese elderly patients with dementia: an exploratory randomized controlled trial. *Neuropsychiatric Disease & Treatment*. 2019;15:2171-80. doi: <https://dx.doi.org/10.2147/NDT.S200045>. PMID: 31440055.
300. Phillips L, Reid-Arndt S, Pak Y. Effects of a creative expression intervention on emotions, communication, and quality of life in persons with dementia. *Nursing research*. 2010;59(6):417-25. doi: 10.1097/NNR.0b013e3181faff52. PMID: CN-00770848.
301. van Dijk AM, van Weert JC, Droes RM. Does theatre improve the quality of life of people with dementia? *International Psychogeriatrics*. 2012 Mar;24(3):367-81. doi: <https://dx.doi.org/10.1017/S1041610211001992>. PMID: 22040605.
302. Luttenberger K, Donath C, Uter W, et al. Effects of multimodal nondrug therapy on dementia symptoms and need for care in nursing home residents with degenerative dementia: a randomized-controlled study with 6-month follow-up. *Journal of the American Geriatrics Society*. 2012 May;60(5):830-40. doi: <https://dx.doi.org/10.1111/j.1532-5415.2012.03938.x>. PMID: 22468985.
303. Graessel E, Stemmer R, Eichenseer B, et al. Non-pharmacological, multicomponent group therapy in patients with degenerative dementia: a 12-month randomized, controlled trial. *BMC Medicine*. 2011 Dec 01;9:129. doi: <https://dx.doi.org/10.1186/1741-7015-9-129>. PMID: 22133165.
304. McCurry SM, Pike KC, Vitiello MV, et al. Increasing walking and bright light exposure to improve sleep in community-dwelling persons with Alzheimer's disease: results of a randomized, controlled trial. *Journal of the American Geriatrics Society*. 2011 Aug;59(8):1393-402. doi: <https://dx.doi.org/10.1111/j.1532-5415.2011.03519.x>. PMID: 21797835.
305. Tadaka E, Kanagawa K. Randomized controlled trial of a group care program for community-dwelling elderly people with dementia. *Japan Journal of Nursing Science*. 2004 Aug;1(1):19-25. doi: <http://dx.doi.org/10.1111/j.1742-7924.2004.00006.x>. PMID: 2007-06056-002.
306. Brooker DJ, Argyle E, Scally AJ, et al. The enriched opportunities programme for people with dementia: a cluster-randomised controlled trial in 10 extra care housing schemes. *Aging & Mental Health*. 2011 Nov;15(8):1008-17. doi: <https://dx.doi.org/10.1080/13607863.2011.583628>. PMID: 21702705.

307. Burgener SC, Marsh-Yant S, Nega KK. A Combined, Multimodal Intervention for Individuals with Dementia. *Research in Gerontological Nursing*. 2011;4(1):64-75. doi: 10.3928/19404921-20100504-01. PMID: 104897556. Language: English. Entry Date: 20110622. Revision Date: 20150711. Publication Type: Journal Article.
308. Burgener SC, Yang Y, Gilbert R, et al. The effects of a multimodal intervention on outcomes of persons with early-stage dementia. *American Journal of Alzheimer's Disease & Other Dementias*. 2008 Aug-Sep;23(4):382-94. doi: <https://dx.doi.org/10.1177/153331750817527>. PMID: 18453642.
309. Chapman SB, Weiner MF, Rackley A, et al. Effects of cognitive-communication stimulation for Alzheimer's disease patients treated with donepezil. *Journal of Speech Language & Hearing Research*. 2004 Oct;47(5):1149-63. PMID: 15603468.
310. Chen YL, Pei YC. Musical dual-task training in patients with mild-to-moderate dementia: a randomized controlled trial. *Neuropsychiatric Disease & Treatment*. 2018;14:1381-93. doi: <https://dx.doi.org/10.2147/NDT.S159174>. PMID: 29881275.
311. Ferrero-Arias J, Goni-Imizcoz M, Gonzalez-Bernal J, et al. The efficacy of nonpharmacological treatment for dementia-related apathy. *Alzheimer Disease & Associated Disorders*. 2011 Jul-Sep;25(3):213-9. doi: <https://dx.doi.org/10.1097/WAD.0b013e3182087dbc>. PMID: 21346517.
312. Kampragkou C, Iakovidis P, Kampragkou E, et al. Effects of a 12-week aerobic exercise program combined with music therapy and memory exercises on cognitive and functional ability in people with middle type of alzheimer's disease. 2017;4(5):262-8. doi: 10.15621/ijphy/2017/v4i5/159420. PMID: CN-01425536.
313. Kim H-J, Yang Y, Oh J-G, et al. Effectiveness of a community-based multidomain cognitive intervention program in patients with Alzheimer's disease. *Geriatrics & gerontology international*. 2016;16(2):191-9. doi: 10.1111/ggi.12453. PMID: CN-01132849.
314. Okamura H, Otani M, Shimoyama N, et al. Combined Exercise and Cognitive Training System for Dementia Patients: A Randomized Controlled Trial. *Dementia & Geriatric Cognitive Disorders*. 2018;45(5-6):318-25. doi: <https://dx.doi.org/10.1159/000490613>. PMID: 30036871.
315. Tappen RM, Roach KE, Applegate EB, et al. Effect of a combined walking and conversation intervention on functional mobility of nursing home residents with Alzheimer disease. *Alzheimer Disease & Associated Disorders*. 2000 Oct-Dec;14(4):196-201. PMID: 11186596.
316. Wesson J, Clemson L, Brodaty H, et al. A feasibility study and pilot randomised trial of a tailored prevention program to reduce falls in older people with mild dementia. *BMC Geriatrics*. 2013 Sep 03;13:89. doi: <https://dx.doi.org/10.1186/1471-2318-13-89>. PMID: 24004682.
317. Yamagami T, Takayama Y, Maki Y, et al. A randomized controlled trial of brain-activating rehabilitation for elderly participants with dementia in residential care homes. *Dementia and Geriatric Cognitive Disorders Extra*. 2012 Jan;2(1):372-80. doi: <https://dx.doi.org/10.1159/000342614>. PMID: 23300492.
318. Young DK, Ng PY, Kwok T, et al. The effects of an expanded cognitive stimulation therapy model on the improvement of cognitive ability of elderly with mild stage Dementia living in a community - a randomized waitlist controlled trial. *Aging & Mental Health*. 2019 Jul;23(7):855-62. doi: <https://dx.doi.org/10.1080/13607863.2018.1471586>. PMID: 29781725.

319. Femia EE, Zarit SH, Stephens MA, et al. Impact of adult day services on behavioral and psychological symptoms of dementia. *Gerontologist*. 2007 Dec;47(6):775-88. PMID: 18192631.
320. Fernandez-Calvo B, Contador I, Ramos F, et al. Effect of unawareness on rehabilitation outcome in a randomised controlled trial of multicomponent intervention for patients with mild Alzheimer's disease. *Neuropsychological Rehabilitation*. 2015;25(3):448-77. doi: <https://dx.doi.org/10.1080/09602011.2014.948461>. PMID: 25121567.
321. Middelstadt J, Folkerts AK, Blawath S, et al. Cognitive Stimulation for People with Dementia in Long-Term Care Facilities: Baseline Cognitive Level Predicts Cognitive Gains, Moderated by Depression. *Journal of Alzheimer's Disease*. 2016 08 04;54(1):253-68. doi: <https://dx.doi.org/10.3233/JAD-160181>. PMID: 27497474.
322. Rokstad AMM, Engedal K, Kirkevold O, et al. The impact of attending day care designed for home-dwelling people with dementia on nursing home admission: a 24-month controlled study. *BMC Health Services Research*. 2018 Nov 16;18(1):864. doi: <https://dx.doi.org/10.1186/s12913-018-3686-5>. PMID: 30445937.
323. Wimo A, Mattsson B, Adolfsson R, et al. Dementia day care and its effects on symptoms and institutionalization--a controlled Swedish study. *Scandinavian Journal of Primary Health Care*. 1993 Jun;11(2):117-23. PMID: 8356361.
324. Sado M, Funaki K, Ninomiya A, et al. Does the Combination of the Cognitive Interventions Improve the Function of Daily Living and Save the Long-Term Care Cost? A Pilot Study of Effectiveness and Cost Saving Analysis of "Learning Therapy" for People with Dementia. *Journal of Alzheimer's Disease*. 2020 Feb 27;27:27. doi: <https://dx.doi.org/10.3233/JAD-190886>. PMID: 32116248.
325. Luttenberger K, Hofner B, Graessel E. Are the effects of a non-drug multimodal activation therapy of dementia sustainable? Follow-up study 10 months after completion of a randomised controlled trial. *BMC Neurology*. 2012 Dec 05;12:151. doi: <https://dx.doi.org/10.1186/1471-2377-12-151>. PMID: 23217188.
326. Lauriks SM, Meiland FJMP, Oste JPM, et al. Effects of Assistive Home Technology on quality of life and falls of people with dementia and job satisfaction of caregivers; results from a pilot randomized controlled trial. *Assistive Technology*. 2018 Dec 28:1-8. doi: <https://dx.doi.org/10.1080/10400435.2018.1531952>. PMID: 30592439.
327. Rowe M, Kelly A, Horne C, et al. Reducing dangerous nighttime events in persons with dementia by using a nighttime monitoring system. In *Alzheimer's & dementia*
328. Tchalla AE, Lachal F, Cardinaud N, et al. Preventing and managing indoor falls with home-based technologies in mild and moderate Alzheimer's disease patients: Pilot study in a community dwelling. *Dementia and Geriatric Cognitive Disorders*. 2013 September;36(3-4):251-61. doi: <http://dx.doi.org/10.1159/000351863>. PMID: 369900271.
329. Gaugler JE, Zmora R, Mitchell LL, et al. Six-Month Effectiveness of Remote Activity Monitoring for Persons Living With Dementia and Their Family Caregivers: An Experimental Mixed Methods Study. *Gerontologist*. 2019 01 09;59(1):78-89. doi: <https://dx.doi.org/10.1093/geront/gny078>. PMID: 29982413.
330. Rowe MA, Kairalla JA, McCrae CS. Sleep in dementia caregivers and the effect of a nighttime monitoring system. *Journal of Nursing Scholarship*. 2010 Sep 01;42(3):338-47. doi: <https://dx.doi.org/10.1111/j.1547-5069.2010.01337.x>. PMID: 20738745.

331. Ahmed MA, Darwish ES, Khedr EM, et al. Effects of low versus high frequencies of repetitive transcranial magnetic stimulation on cognitive function and cortical excitability in Alzheimer's dementia. *Journal of Neurology*. 2012 Jan;259(1):83-92. doi: <https://dx.doi.org/10.1007/s00415-011-6128-4>. PMID: 21671144.
332. Cotelli M, Manenti R, Brambilla M, et al. Anodal tDCS during face-name associations memory training in Alzheimer's patients. *Frontiers in aging neuroscience*. 2014;6:38. doi: <https://dx.doi.org/10.3389/fnagi.2014.00038>. PMID: 24678298.
333. Hozumi S, Hori H, Okawa M, et al. Favorable effect of transcranial electrostimulation on behavior disorders in elderly patients with dementia: a double-blind study. *International Journal of Neuroscience*. 1996 Nov;88(1-2):1-10. PMID: 9003961.
334. Khedr EM, Salama RH, Abdel Hameed M, et al. Therapeutic Role of Transcranial Direct Current Stimulation in Alzheimer Disease Patients: Double-Blind, Placebo-Controlled Clinical Trial. *Neurorehabilitation & Neural Repair*. 2019 05;33(5):384-94. doi: <https://dx.doi.org/10.1177/1545968319840285>. PMID: 30940012.
335. Lozano AM, Fosdick L, Chakravarty MM, et al. A Phase II Study of Fornix Deep Brain Stimulation in Mild Alzheimer's Disease. *Journal of Alzheimer's Disease*. 2016 09 06;54(2):777-87. doi: <https://dx.doi.org/10.3233/JAD-160017>. PMID: 27567810.
336. Olazaran J, Gonzalez B, Lopez-Alvarez J, et al. Motor effects of REAC in advanced Alzheimer's disease: results from a pilot trial. *Journal of Alzheimer's Disease*. 2013;36(2):297-302. doi: <https://dx.doi.org/10.3233/JAD-130077>. PMID: 23603397.
337. Olazaran J, Gonzalez B, Osa-Ruiz E, et al. Motor effects of radio electric asymmetric conveyer in Alzheimer's disease: results from a cross-over trial. *Journal of Alzheimer's Disease*. 2014;42(1):325-32. doi: <https://dx.doi.org/10.3233/JAD-140417>. PMID: 24898637.
338. Scherder EJ, van Tol MJ, Swaab DF. High-frequency cranial electrostimulation (CES) in patients with probable Alzheimer's disease. *American Journal of Physical Medicine & Rehabilitation*. 2006 Jul;85(7):614-8. PMID: 16788393.
339. Suemoto CK, Apolinario D, Nakamura-Palacios EM, et al. Effects of a non-focal plasticity protocol on apathy in moderate Alzheimer's disease: a randomized, double-blind, sham-controlled trial. *Brain Stimulation*. 2014 Mar-Apr;7(2):308-13. doi: <https://dx.doi.org/10.1016/j.brs.2013.10.003>. PMID: 24262299.
340. Wu Y, Xu W, Liu X, et al. Adjunctive treatment with high frequency repetitive transcranial magnetic stimulation for the behavioral and psychological symptoms of patients with Alzheimer's disease: a randomized, double-blind, sham-controlled study. *Shanghai Jingshen Yixue*. 2015 Oct;27(5):280-8. doi: <https://dx.doi.org/10.11919/j.issn.1002-0829.215107>. PMID: 26977125.
341. Elder GJ, Colloby SJ, Firbank MJ, et al. Consecutive sessions of transcranial direct current stimulation do not remediate visual hallucinations in Lewy body dementia: a randomised controlled trial. *Alzheimer's Research & Therapy*. 2019 01 18;11(1):9. doi: <https://dx.doi.org/10.1186/s13195-018-0465-9>. PMID: 30658705.
342. Sabbagh M, Sadowsky C, Tousi B, et al. Effects of a combined transcranial magnetic stimulation (TMS) and cognitive training intervention in patients with Alzheimer's disease. *Alzheimer's & Dementia*. 2019 Dec 23;23:23. doi: <https://dx.doi.org/10.1016/j.jalz.2019.08.197>. PMID: 31879235.

343. Zhang F, Qin Y, Xie L, et al. High-frequency repetitive transcranial magnetic stimulation combined with cognitive training improves cognitive function and cortical metabolic ratios in Alzheimer's disease. *Journal of Neural Transmission*. 2019 01 Aug;126(8):1081-94.
344. van Dijk KR, Scheltens P, Luijpen MW, et al. Peripheral electrical stimulation in Alzheimer's disease. *Dementia & Geriatric Cognitive Disorders*. 2005;19(5-6):361-8. PMID: 15802911.
345. Leoutsakos JS, Yan H, Anderson WS, et al. Deep Brain Stimulation Targeting the Fornix for Mild Alzheimer Dementia (the ADvance Trial): A Two Year Follow-up Including Results of Delayed Activation. *Journal of Alzheimer's Disease*. 2018;64(2):597-606. doi: <https://dx.doi.org/10.3233/JAD-180121>. PMID: 29914028.
346. Batchelor-Murphy M, McConnell E, Amella E, et al. Experimental Comparison of Efficacy for Three Handfeeding Techniques in Dementia. *Journal of the American Geriatrics Society*. 2017;65(4):E89-e94. doi: 10.1111/jgs.14728. PMID: CN-01415879.
347. Coyne ML, Hoskins L. Improving eating behaviors in dementia using behavioral strategies. *Clinical Nursing Research*. 1997 Aug;6(3):275-90. PMID: 9281930.
348. Henskens M, Nauta IM, Scherder EJA, et al. Implementation and effects of Movement-oriented Restorative Care in a nursing home - a quasi-experimental study. *BMC Geriatrics*. 2017;17:1-11. doi: 10.1186/s12877-017-0642-x. PMID: 125844550. Language: English. Entry Date: 20180724. Revision Date: 20181018. Publication Type: journal article.
349. Hobbelen JH, Tan FE, Verhey FR, et al. Passive movement therapy in severe paratonia: a multicenter randomized clinical trial. *International Psychogeriatrics*. 2012 May;24(5):834-44. doi: <https://dx.doi.org/10.1017/S1041610211002468>. PMID: 22185768.
350. Hobbelen JS, Verhey FR, Bor JH, et al. Passive movement therapy in patients with moderate to severe paratonia; study protocol of a randomised clinical trial (ISRCTN43069940). *BMC Geriatrics*. 2007 Dec 19;7:30. PMID: 18093298.
351. Kim JP, Yang J. Effectiveness of a community-based program for suicide prevention among elders with early-stage dementia: A controlled observational study. *Geriatric Nursing*. 2017 Mar - Apr;38(2):97-105. doi: <https://dx.doi.org/10.1016/j.gerinurse.2016.08.002>. PMID: 27594544.
352. Rostad HM, Utne I, Grov EK, et al. The impact of a pain assessment intervention on pain score and analgesic use in older nursing home residents with severe dementia: A cluster randomised controlled trial. *International Journal of Nursing Studies*. 2018 Aug;84:52-60. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2018.04.017>. PMID: 29763832.
353. Cheng ST, Li KK, Losada A, et al. The effectiveness of nonpharmacological interventions for informal dementia caregivers: an updated systematic review and meta-analysis. *Psychol Aging*. 2020;35(1):55-77. doi: 10.1037/pag0000401. PMID: 31985249.
354. Au A, Yip HM, Lai S, et al. Telephone-based behavioral activation intervention for dementia family caregivers: Outcomes and mediation effect of a randomized controlled trial. *Patient Education & Counseling*. 2019 Nov;102(11):2049-59. doi: <https://dx.doi.org/10.1016/j.pec.2019.06.009>. PMID: 31279613.
355. Bjorge H, Kvaal K, Ulstein I. The effect of psychosocial support on caregivers' perceived criticism and emotional over-involvement of persons with dementia: an assessor-blinded randomized controlled trial. *BMC Health Services Research*. 2019 Oct 24;19(1):744. doi: <https://dx.doi.org/10.1186/s12913-019-4551-x>. PMID: 31651321.

356. Charlesworth G, Burnell K, Crellin N, et al. Peer support and reminiscence therapy for people with dementia and their family carers: a factorial pragmatic randomised trial. *Journal of Neurology, Neurosurgery & Psychiatry*. 2016 Nov;87(11):1218-28. doi: <https://dx.doi.org/10.1136/jnnp-2016-313736>. PMID: 27521377.
357. Cheng ST, Fung HH, Chan WC, et al. Short-Term Effects of a Gain-Focused Reappraisal Intervention for Dementia Caregivers: A Double-Blind Cluster-Randomized Controlled Trial. *American Journal of Geriatric Psychiatry*. 2016 Sep;24(9):740-50. doi: <https://dx.doi.org/10.1016/j.jagp.2016.04.012>. PMID: 27401052.
358. de Rotrou J, Cantegreil I, Faucounau V, et al. Do patients diagnosed with Alzheimer's disease benefit from a psycho-educational programme for family caregivers? A randomised controlled study. *International Journal of Geriatric Psychiatry*. 2011 Aug;26(8):833-42. doi: <https://dx.doi.org/10.1002/gps.2611>. PMID: 20922772.
359. Gitlin LN, Arthur P, Piersol C, et al. Targeting Behavioral Symptoms and Functional Decline in Dementia: A Randomized Clinical Trial. *Journal of the American Geriatrics Society*. 2018 Feb;66(2):339-45. doi: <https://dx.doi.org/10.1111/jgs.15194>. PMID: 29192967.
360. Gonyea JG, Lopez LM, Velasquez EH. The Effectiveness of a Culturally Sensitive Cognitive Behavioral Group Intervention for Latino Alzheimer's Caregivers. *Gerontologist*. 2016 Apr;56(2):292-302. doi: <https://dx.doi.org/10.1093/geront/gnu045>. PMID: 24855313.
361. Guerra M, Ferri CP, Fonseca M, et al. Helping carers to care: the 10/66 dementia research group's randomized control trial of a caregiver intervention in Peru. *Revista Brasileira de Psiquiatria*. 2011 Mar;33(1):47-54. PMID: 20602013.
362. Laakkonen ML, Kautiainen H, Holttä E, et al. Effects of Self-Management Groups for People with Dementia and Their Spouses--Randomized Controlled Trial. *Journal of the American Geriatrics Society*. 2016 Apr;64(4):752-60. doi: <https://dx.doi.org/10.1111/jgs.14055>. PMID: 27060101.
363. Livingston G, Barber J, Rapaport P, et al. START (STRAtegies for RelaTives) study: a pragmatic randomised controlled trial to determine the clinical effectiveness and cost-effectiveness of a manual-based coping strategy programme in promoting the mental health of carers of people with dementia. *Health Technology Assessment (Winchester, England)*. 2014 Oct;18(61):1-242. doi: <https://dx.doi.org/10.3310/hta18610>. PMID: 25300037.
364. Martin-Carrasco M, Dominguez-Panchon AI, Gonzalez-Fraile E, et al. Effectiveness of a psychoeducational intervention group program in the reduction of the burden experienced by caregivers of patients with dementia: the EDUCA-II randomized trial. *Alzheimer Disease & Associated Disorders*. 2014 Jan-Mar;28(1):79-87. doi: <https://dx.doi.org/10.1097/WAD.00000000000003>. PMID: 24113563.
365. Meng H, Marino VR, Conner KO, et al. Effects of in-person and telephone-based cognitive behavioral therapies on health services use and expenditures among African-American dementia caregivers with depressive symptoms. *Ethnicity & Health*. 2019 Mar 19:1-14. doi: <https://dx.doi.org/10.1080/13557858.2019.1590536>. PMID: 30884961.
366. Moskowitz JT, Cheung EO, Snowberg KE, et al. Randomized controlled trial of a facilitated online positive emotion regulation intervention for dementia caregivers. *Health Psychology*. 2019 May;38(5):391-402. doi: <https://dx.doi.org/10.1037/hea0000680>. PMID: 31045422.

367. Nordheim J, Hausler A, Yasar S, et al. Psychosocial Intervention in Couples Coping with Dementia Led by a Psychotherapist and a Social Worker: The DYADEM Trial. *Journal of Alzheimer's Disease*. 2019;68(2):745-55. doi: <https://dx.doi.org/10.3233/JAD-180812>. PMID: 30775982.
368. Tremont G, Davis JD, Papandonatos GD, et al. Psychosocial telephone intervention for dementia caregivers: A randomized, controlled trial. *Alzheimer's & Dementia*. 2015 May;11(5):541-8. doi: <https://dx.doi.org/10.1016/j.jalz.2014.05.1752>. PMID: 25074341.
369. Waldorff FB, Buss DV, Eckermann A, et al. Efficacy of psychosocial intervention in patients with mild Alzheimer's disease: the multicentre, rater blinded, randomised Danish Alzheimer Intervention Study (DAISY). *BMJ*. 2012 Jul 17;345:e4693. doi: <https://dx.doi.org/10.1136/bmj.e4693>. PMID: 22807076.
370. Farran CJ, Gilley DW, McCann JJ, et al. Psychosocial interventions to reduce depressive symptoms of dementia caregivers: A randomized clinical trial comparing two approaches. *Journal of Mental Health and Aging*. 2004 Winter;10(4):337-50. PMID: 41552352.
371. Gallagher-Thompson D, Coon DW, Solano N, et al. Change in indices of distress among Latino and Anglo female caregivers of elderly relatives with dementia: site-specific results from the REACH national collaborative study. *Gerontologist*. 2003 Aug;43(4):580-91. PMID: 12937336.
372. Gallagher-Thompson D, Gray HL, Dupart T, et al. Effectiveness of Cognitive/Behavioral Small Group Intervention for Reduction of Depression and Stress in Non-Hispanic White and Hispanic/Latino Women Dementia Family Caregivers: Outcomes and Mediators of Change. *Journal of Rational-Emotive & Cognitive Behavior Therapy*. 2008 Dec 01;26(4):286-303. PMID: 25067886.
373. Gitlin LN, Corcoran M, Winter L, et al. A randomized, controlled trial of a home environmental intervention: effect on efficacy and upset in caregivers and on daily function of persons with dementia. *Gerontologist*. 2001 Feb;41(1):4-14. PMID: 11220813.
374. Graff MJ, Vernooij-Dassen MJ, Thijssen M, et al. Community based occupational therapy for patients with dementia and their care givers: randomised controlled trial. *BMJ*. 2006 Dec 09;333(7580):1196. PMID: 17114212.
375. Martin-Carrasco M, Martin MF, Valero CP, et al. Effectiveness of a psychoeducational intervention program in the reduction of caregiver burden in Alzheimer's disease patients' caregivers. *International Journal of Geriatric Psychiatry*. 2009 May;24(5):489-99. doi: <https://dx.doi.org/10.1002/gps.2142>. PMID: 18949763.
376. Wang LQ, Chien WT. Randomised controlled trial of a family-led mutual support programme for people with dementia. *Journal of Clinical Nursing*. 2011 Aug;20(15-16):2362-6. doi: <https://dx.doi.org/10.1111/j.1365-2702.2011.03746.x>. PMID: 21752121.
377. Gitlin LN, Winter L, Dennis MP, et al. Targeting and managing behavioral symptoms in individuals with dementia: a randomized trial of a nonpharmacological intervention. *Journal of the American Geriatrics Society*. 2010 Aug;58(8):1465-74. doi: <https://dx.doi.org/10.1111/j.1532-5415.2010.02971.x>. PMID: 20662955.
378. Voigt-Radloff S, Graff M, Leonhart R, et al. A multicentre RCT on community occupational therapy in Alzheimer's disease: 10 sessions are not better than one consultation. *BMJ Open*. 2011;1. doi:10.1136/bmjopen-2011-000096.
379. Otero P, Smit F, Cuijpers P, et al. Long-term efficacy of indicated prevention of depression in non-professional caregivers: randomized controlled trial. *Psychol Med*. 2015;45(7):1401-12. doi: 10.1017/S0033291714002505. PMID: 25331992.

380. Au A, Lai DWL, Biggs S, et al. Perspective-taking interventions for intergenerational caregivers of alzheimer's diseases: a randomized controlled trial. *Research on social work practice*. 2020;30(3):306-19. PMID: CN-02084449.
381. Ghaffari F, Rostami M, Fotokian Z, et al. Effectiveness of resilience education in the mental health of family caregivers of elderly patients with Alzheimer's disease. *Iranian Journal of Psychiatry and Behavioral Sciences*. 2019;13 (3) (no pagination)(e69507).
382. Chang BL. Cognitive-behavioral intervention for homebound caregivers of persons with dementia. *Nursing Research*. 1999 May-Jun;48(3):173-82. PMID: 10337848.
383. Dias A, Dewey ME, D'Souza J, et al. The effectiveness of a home care program for supporting caregivers of persons with dementia in developing countries: a randomised controlled trial from Goa, India. *PLoS ONE [Electronic Resource]*. 2008 Jun 04;3(6):e2333. doi: <https://dx.doi.org/10.1371/journal.pone.0002333>. PMID: 18523642.
384. Marriott A, Donaldson C, Tarrier N, et al. Effectiveness of cognitive-behavioural family intervention in reducing the burden of care in carers of patients with Alzheimer's disease. *British Journal of Psychiatry*. 2000 Jun;176:557-62. PMID: 10974962.
385. Spijker A, Wollersheim H, Teerenstra S, et al. Systematic care for caregivers of patients with dementia: a multicenter, cluster-randomized, controlled trial. *American Journal of Geriatric Psychiatry*. 2011 Jun;19(6):521-31. doi: <https://dx.doi.org/10.1097/JGP.0b013e3182110599>. PMID: 21358385.
386. Stolley JM, Reed D, Buckwalter KC. Caregiving appraisal and interventions based on the progressively lowered stress threshold model. *American Journal of Alzheimer's Disease & Other Dementias*. 2002 Mar-Apr;17(2):110-20. PMID: 11954669.
387. Wawrziczny E, Larochette C, Papo D, et al. A Customized Intervention for Dementia Caregivers: A Quasi-Experimental Design. *Journal of Aging & Health*. 2019 Aug;31(7):1172-95. doi: <https://dx.doi.org/10.1177/0898264318770056>. PMID: 29665714.
388. Au A, Gallagher-Thompson D, Wong MK, et al. Behavioral activation for dementia caregivers: scheduling pleasant events and enhancing communications. *Clinical Interventions In Aging*. 2015;10:611-9. doi: <https://dx.doi.org/10.2147/CIA.S72348>. PMID: 25848237.
389. Chen HM, Huang MF, Yeh YC, et al. Effectiveness of coping strategies intervention on caregiver burden among caregivers of elderly patients with dementia. *Psychogeriatrics: The Official Journal of the Japanese Psychogeriatric Society*. 2015 Mar;15(1):20-5. doi: <https://dx.doi.org/10.1111/psyg.12071>. PMID: 25515800.
390. Cristancho-Lacroix V, Wrobel J, Cantegreil-Kallen I, et al. A web-based psychoeducational program for informal caregivers of patients with Alzheimer's disease: a pilot randomized controlled trial. *Journal of Medical Internet Research*. 2015 May 12;17(5):e117. doi: <https://dx.doi.org/10.2196/jmir.3717>. PMID: 25967983.
391. Gallagher-Thompson D, Wang PC, Liu W, et al. Effectiveness of a psychoeducational skill training DVD program to reduce stress in Chinese American dementia caregivers: results of a preliminary study. *Aging & mental health*. 2010 Apr;14(3):263-73. doi: <http://dx.doi.org/10.1080/13607860903420989>. PMID: 359062574.
392. Gant JR, Steffen AM, Lauderdale SA. Comparative outcomes of two distance-based interventions for male caregivers of family members with dementia. *American Journal of Alzheimer's Disease & Other Dementias*. 2007 Apr-May;22(2):120-8. PMID: 17545139.

393. Gonzalez EW, Polansky M, Lippa CF, et al. Enhancing resourcefulness to improve outcomes in family caregivers and persons with Alzheimer's disease: a pilot randomized trial. *International journal of Alzheimer's disease*. 2014;2014:323478. doi: <https://dx.doi.org/10.1155/2014/323478>. PMID: 25328754.
394. Hebert R, Leclerc G, Bravo G, et al. Efficacy of a support group programme for care-givers of demented patients in the community: A randomized controlled trial. *Archives of Gerontology and Geriatrics*. 1994;18(1):1-14. doi: [http://dx.doi.org/10.1016/0167-4943\(94\)90043-4](http://dx.doi.org/10.1016/0167-4943(94)90043-4). PMID: 24087060.
395. Kales HC, Gitlin LN, Stanislawski B, et al. Effect of the WeCareAdvisor™ on family caregiver outcomes in dementia: a pilot randomized controlled trial. *BMC Geriatrics*. 2018 05 10;18(1):113. doi: <https://dx.doi.org/10.1186/s12877-018-0801-8>. PMID: 29747583.
396. Lok N, Bademli K. Pilot testing of the "First You Should Get Stronger" program among caregivers of older adults with dementia. *Archives of Gerontology & Geriatrics*. 2017 Jan - Feb;68:84-9. doi: <https://dx.doi.org/10.1016/j.archger.2016.09.006>. PMID: 27689315.
397. Martin-Cook K, Remakel-Davis B, Svetlik D, et al. Caregiver attribution and resentment in dementia care. *American Journal of Alzheimer's Disease & Other Dementias*. 2003 Nov-Dec;18(6):366-74. PMID: 14682086.
398. McCurry SM, Logsdon RG, Vitiello MV, et al. Successful behavioral treatment for reported sleep problems in elderly caregivers of dementia patients: a controlled study. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences*. 1998 Mar;53(2):P122-9. PMID: 9520929.
399. Nunez-Naveira L, Alonso-Bua B, de Labra C, et al. UnderstAID, an ICT Platform to Help Informal Caregivers of People with Dementia: A Pilot Randomized Controlled Study. *BioMed Research International*. 2016;2016:5726465. doi: <https://dx.doi.org/10.1155/2016/5726465>. PMID: 28116300.
400. Steffen AM. Anger management for dementia caregivers: A preliminary study using video and telephone interventions. *Behavior Therapy*. 2000 Spr;31(2):281-99. doi: [http://dx.doi.org/10.1016/S0005-7894\(98\)0016-7](http://dx.doi.org/10.1016/S0005-7894(98)0016-7). PMID: 2000-16043-005.
401. Ta Park VM, Ton V, Tiet QQ, et al. Promising results from a pilot study to reduce distress in Vietnamese American dementia and memory loss caregivers. *Alzheimer's & Dementia : Translational Research & Clinical Interventions*. 2019;5:319-27. doi: <https://dx.doi.org/10.1016/j.trci.2019.05.006>. PMID: 31384663.
402. Zauszniewski JA, Lekhak N, Burant CJ, et al. Preliminary Evidence for Effectiveness of Resourcefulness Training in Women Dementia Caregivers. *Journal of Family Medicine*. 2016;3(5). PMID: 27500286.
403. Aboulafia-Brakha T, Suchecki D, Gouveia-Paulino F, et al. Cognitive-behavioural group therapy improves a psychophysiological marker of stress in caregivers of patients with Alzheimer's disease. *Aging & Mental Health*. 2014;18(6):801-8. doi: <https://dx.doi.org/10.1080/13607863.2014.880406>. PMID: 24499394.
404. Akkerman RL, Ostwald SK. Reducing anxiety in Alzheimer's disease family caregivers: the effectiveness of a nine-week cognitive-behavioral intervention. *American Journal of Alzheimer's Disease & Other Dementias*. 2004 Mar-Apr;19(2):117-23. PMID: 15106393.
405. Au A, Li S, Lee K, et al. The Coping with Caregiving Group Program for Chinese caregivers of patients with Alzheimer's disease in Hong Kong. In *Patient education and counseling*

406. Brodaty H, Gresham M, Luscombe G. The Prince Henry Hospital dementia caregivers' training programme. *International Journal of Geriatric Psychiatry*. 1997 Feb;12(2):183-92. PMID: 9097211.
407. Davis JD, Tremont G, Bishop DS, et al. A telephone-delivered psychosocial intervention improves dementia caregiver adjustment following nursing home placement. *International Journal of Geriatric Psychiatry*. 2011 Apr;26(4):380-7. doi: <https://dx.doi.org/10.1002/gps.2537>. PMID: 20842759.
408. Dowling GA, Merrilees J, Mastick J, et al. Life enhancing activities for family caregivers of people with frontotemporal dementia. *Alzheimer Disease & Associated Disorders*. 2014 Apr-Jun;28(2):175-81. doi: <https://dx.doi.org/10.1097/WAD.0b013e3182a6b905>. PMID: 24113564.
409. Garand L, Buckwalter KC, Lubaroff D, et al. A pilot study of immune and mood outcomes of a community-based intervention for dementia caregivers: the PLST intervention. *Archives of Psychiatric Nursing*. 2002 Aug;16(4):156-67. PMID: 12143075.
410. Gaugler JE, Reese M, Sauld J. A Pilot Evaluation of Psychosocial Support for Family Caregivers of Relatives with Dementia in Long-Term Care: The Residential Care Transition Module. *Research in Gerontological Nursing*. 2015 Jul-Aug;8(4):161-72. doi: <https://dx.doi.org/10.3928/19404921-20150304-01>. PMID: 25751083.
411. Gendron C, Poitras L, Dastoor DP, et al. Cognitive-behavioral group intervention for spousal caregivers: Findings and clinical considerations. *Clinical Gerontologist*. 1996;17(1):3-19. doi: http://dx.doi.org/10.1300/J018v17n01_02. PMID: 26250550.
412. Glueckauf R, Sharma D, Davis W, et al. Telephone-based cognitive-behavioral intervention for distressed rural dementia caregivers: initial findings. In *Clinical gerontologist*
413. Huang H-L, Shyu Y-IL, Chen M-C, et al. A pilot study on a home-based caregiver training program for improving caregiver self-efficacy and decreasing the behavioral problems of elders with dementia in Taiwan. *International Journal of Geriatric Psychiatry*. 2003;18(4):337-45. doi: 10.1002/gps.835. PMID: 104731281. Language: English. Entry Date: 20110610. Revision Date: 20150711. Publication Type: Journal Article.
414. Kamkhagi D, Costa A, Kusminsky S, et al. Benefits of psychodynamic group therapy on depression, burden and quality of life of family caregivers to Alzheimer's disease patients. *Revista de psiquiatria clinica*. 2015;42(6):157-60. doi: 10.1590/0101-60830000000067. PMID: CN-01162210.
415. Meichsner F, Theurer C, Wilz G. Acceptance and treatment effects of an internet-delivered cognitive-behavioral intervention for family caregivers of people with dementia: A randomized-controlled trial. *Journal of Clinical Psychology*. 2018 Dec 31;31:31. doi: <https://dx.doi.org/10.1002/jclp.22739>. PMID: 30597537.
416. Moniz-Cook E, Agar S, Gibson G, et al. A preliminary study of the effects of early intervention with people with dementia and their families in a memory clinic. *Aging & Mental Health*. 1998;2(3):199-211. PMID: 107163779. Language: English. Entry Date: 20070101. Revision Date: 20150711. Publication Type: Journal Article.
417. Nobili A, Riva E, Tettamanti M, et al. The effect of a structured intervention on caregivers of patients with dementia and problem behaviors: a randomized controlled pilot study. *Alzheimer Disease & Associated Disorders*. 2004 Apr-Jun;18(2):75-82. PMID: 15249851.

418. Orgeta V, Tuijt R, Leung P, et al. Behavioral Activation for Promoting Well-Being in Mild Dementia: Feasibility and Outcomes of a Pilot Randomized Controlled Trial. *Journal of Alzheimer's Disease*. 2019 Oct 08;08:08. doi: <https://dx.doi.org/10.3233/JAD-190696>. PMID: 31609689.
419. Paun O, Farran CJ, Fogg L, et al. A chronic grief intervention for dementia family caregivers in long-term care. *Western Journal of Nursing Research*. 2015 Jan;37(1):6-27. doi: <https://dx.doi.org/10.1177/0193945914521040>. PMID: 24510968.
420. de Oliveira AM, Radanovic M, Homem de Mello PC, et al. An intervention to reduce neuropsychiatric symptoms and caregiver burden in dementia: Preliminary results from a randomized trial of the tailored activity program-outpatient version. *International Journal of Geriatric Psychiatry*. 2018 Jul 23;23:23. doi: <https://dx.doi.org/10.1002/gps.4958>. PMID: 30035341.
421. Gitlin LN, Winter L, Burke J, et al. Tailored activities to manage neuropsychiatric behaviors in persons with dementia and reduce caregiver burden: a randomized pilot study. *American Journal of Geriatric Psychiatry*. 2008 Mar;16(3):229-39. doi: <https://dx.doi.org/10.1097/JGP.0b013e318160da72>. PMID: 18310553.
422. Kwok T, Wong B, Ip I, et al. Telephone-delivered psychoeducational intervention for Hong Kong Chinese dementia caregivers: a single-blinded randomized controlled trial. *Clinical Interventions In Aging*. 2013;8:1191-7. doi: <https://dx.doi.org/10.2147/CIA.S48264>. PMID: 24072965.
423. Meichsner F, Theurer C, Wilz G. Acceptance and treatment effects of an internet-delivered cognitive-behavioral intervention for family caregivers of people with dementia: A randomized-controlled trial. *Journal of Clinical Psychology*. 2019 Apr;75(4):594-613. doi: <https://dx.doi.org/10.1002/jclp.22739>. PMID: 30597537.
424. Novelli M, Machado SCB, Lima GB, et al. Effects of the Tailored Activity Program in Brazil (TAP-BR) for Persons With Dementia: A Randomized Pilot Trial. *Alzheimer Disease & Associated Disorders*. 2018 Apr 25;25:25. doi: <https://dx.doi.org/10.1097/WAD.0000000000000256>. PMID: 29698251.
425. O'Connor CM, Clemson L, Brodaty H, et al. The tailored activity program (TAP) to address behavioral disturbances in frontotemporal dementia: a feasibility and pilot study. *Disability & Rehabilitation*. 2017 Oct 15:1-12. doi: <https://dx.doi.org/10.1080/09638288.2017.1387614>. PMID: 29034719.
426. Sepe-Monti M, Vanacore N, Bartorelli L, et al. The Savvy Caregiver Program: A Probe Multicenter Randomized Controlled Pilot Trial in Caregivers of Patients Affected by Alzheimer's Disease. *Journal of Alzheimer's Disease*. 2016 10 04;54(3):1235-46. PMID: 27567824.
427. Supiano KP, Haynes LB, Pond V. The Process of Change in Complicated Grief Group Therapy for Bereaved Dementia Caregivers: An Evaluation Using the Meaning of Loss Codebook. *Journal of Gerontological Social Work*. 2017 Feb-Mar;60(2):155-69. doi: <https://dx.doi.org/10.1080/01634372.2016.1274930>. PMID: 28051926.
428. Teri L, Logsdon RG, Uomoto J, et al. Behavioral treatment of depression in dementia patients: a controlled clinical trial. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences*. 1997 Jul;52(4):P159-66. PMID: 9224439.

429. Tremont G, Davis JD, Bishop DS, et al. Telephone-delivered psychosocial intervention reduces burden in dementia caregivers. *Dementia: The International Journal of Social Research and Practice*. 2008 Nov;7(4):503-20. doi: <http://dx.doi.org/10.1177/1471301208096632>. PMID: 2008-16304-006.
430. van Knippenberg RJM, de Vugt ME, Ponds RW, et al. An Experience Sampling Method Intervention for Dementia Caregivers: Results of a Randomized Controlled Trial. *American Journal of Geriatric Psychiatry*. 2018 Dec;26(12):1231-43. doi: <https://dx.doi.org/10.1016/j.jagp.2018.06.004>. PMID: 30126766.
431. Yoo R, Yeom J, Kim GH, et al. A Multicenter, Randomized Clinical Trial to Assess the Efficacy of a Therapeutic Intervention Program for Caregivers of People with Dementia. *Journal of Clinical Neurology*. 2019 Apr;15(2):235-42. doi: <https://dx.doi.org/10.3988/jcn.2019.15.2.235>. PMID: 30938110.
432. Boots LM, de Vugt ME, Kempen GI, et al. Effectiveness of a Blended Care Self-Management Program for Caregivers of People With Early-Stage Dementia (Partner in Balance): Randomized Controlled Trial. *Journal of Medical Internet Research*. 2018 Jul 13;20(7):e10017. doi: <https://dx.doi.org/10.2196/10017>. PMID: 30006327.
433. Huang HL, Shyu YI, Chen MC, et al. A pilot study on a home-based caregiver training program for improving caregiver self-efficacy and decreasing the behavioral problems of elders with dementia in Taiwan. *International Journal of Geriatric Psychiatry*. 2003 Apr;18(4):337-45. PMID: 12673611.
434. Spalding-Wilson KN, Guzman-Velez E, Angelica J, et al. A novel two-day intervention reduces stress in caregivers of persons with dementia. *Alzheimer's and Dementia: Translational Research and Clinical Interventions*. 2018 01 Jan;4:450-60. doi: <http://dx.doi.org/10.1016/j.trci.2018.08.004>. PMID: 2001115918.
435. Whitlatch CJ, Heid AR, Femia EE, et al. The Support, Health, Activities, Resources, and Education program for early stage dementia: Results from a randomized controlled trial. *Dementia*. 2019 08;18(6):2122-39. doi: <https://dx.doi.org/10.1177/1471301217743033>. PMID: 29171296.
436. Wilz G, Reder M, Meichsner F, et al. The Tele.TAnDem Intervention: Telephone-based CBT for Family Caregivers of People With Dementia. *Gerontologist*. 2018 03 19;58(2):e118-e29. doi: <https://dx.doi.org/10.1093/geront/gnx183>. PMID: 29190357.
437. Au A. Developing Volunteer-Assisted Behavioral Activation Teleprograms to Meet the Needs of Chinese Dementia Caregivers. *Clinical gerontologist*. 2015;38(3):190-202. PMID: CN-01071414.
438. Blom MM, Zarit SH, Groot Zwaafink RB, et al. Effectiveness of an Internet intervention for family caregivers of people with dementia: results of a randomized controlled trial. *PLoS ONE [Electronic Resource]*. 2015;10(2):e0116622. doi: <https://dx.doi.org/10.1371/journal.pone.0116622>. PMID: 25679228.
439. Callahan CM, Boustani MA, Schmid AA, et al. Targeting Functional Decline in Alzheimer Disease: A Randomized Trial. *Annals of Internal Medicine*. 2017 Feb 07;166(3):164-71. doi: <https://dx.doi.org/10.7326/M16-0830>. PMID: 27893087.
440. Chiu M, Pauley T, Wesson V, et al. Evaluation of a problem-solving (PS) techniques-based intervention for informal carers of patients with dementia receiving in-home care. *International psychogeriatrics*. 2015;27(6):937-48. doi: 10.1017/S1041610214002798. PMID: CN-01102329.

441. Taati F, Bahramnezhad F, Seyedfatemi N, et al. The effect of participation in support groups on depression, anxiety and stress in family caregivers of people with Alzheimer's: randomized clinical trial. *International journal of medical research & health sciences*. 2016;5(7):256-62. PMID: CN-01308282.
442. Whitlatch CJ, Heid AR, Femia EE, et al. The Support, Health, Activities, Resources, and Education program for early stage dementia: Results from a randomized controlled trial. *Dementia*. 2017 Jan 01:1471301217743033. doi: <https://dx.doi.org/10.1177/1471301217743033>. PMID: 29171296.
443. Arango-Lasprilla JC, Panyavin I, Merchan EJH, et al. Evaluation of a group cognitive-behavioral dementia caregiver intervention in Latin America. *American Journal of Alzheimer's Disease and Other Dementias*. 2014 Sep;29(6):548-55. doi: <http://dx.doi.org/10.1177/1533317514523668>. PMID: 2014-38564-010.
444. Bruvik FK, Allore HG, Ranhoff AH, et al. The effect of psychosocial support intervention on depression in patients with dementia and their family caregivers: an assessor-blinded randomized controlled trial. *Dementia and Geriatric Cognitive Disorders Extra*. 2013;3(1):386-97. doi: <https://dx.doi.org/10.1159/000355912>. PMID: 24348500.
445. Gallagher-Thompson D, Tzuang M, Hinton L, et al. Effectiveness of a fotonovela for reducing depression and stress in Latino dementia family caregivers. *Alzheimer Disease & Associated Disorders*. 2015 Apr-Jun;29(2):146-53. doi: <https://dx.doi.org/10.1097/WAD.000000000000077>. PMID: 25590939.
446. Huang HL, Kuo LM, Chen YS, et al. A home-based training program improves caregivers' skills and dementia patients' aggressive behaviors: a randomized controlled trial. *American Journal of Geriatric Psychiatry*. 2013 Nov;21(11):1060-70. doi: <https://dx.doi.org/10.1016/j.jagp.2012.09.009>. PMID: 23933422.
447. Losada A, Marquez-Gonzalez M, Romero-Moreno R, et al. Cognitive-behavioral therapy (CBT) versus acceptance and commitment therapy (ACT) for dementia family caregivers with significant depressive symptoms: Results of a randomized clinical trial. *Journal of Consulting & Clinical Psychology*. 2015 Aug;83(4):760-72. doi: <https://dx.doi.org/10.1037/ccp0000028>. PMID: 26075381.
448. Otero P, Smit F, Cuijpers P, et al. Differential response to depression prevention among a sample of informal caregivers: Moderator analysis of longer-term follow-up trial data. *Psychiatry Research*. 2015 Dec 15;230(2):271-8. doi: <https://dx.doi.org/10.1016/j.psychres.2015.09.005>. PMID: 26456895.
449. Passoni S, Moroni L, Toraldo A, et al. Cognitive behavioral group intervention for Alzheimer caregivers. *Alzheimer Disease & Associated Disorders*. 2014;28(3):275-82. doi: 10.1097/WAD.0000000000000033. PMID: 107799151. Language: English. Entry Date: 20150508. Revision Date: 20150712. Publication Type: Journal Article.
450. Chu H, Yang CY, Liao YH, et al. The effects of a support group on dementia caregivers' burden and depression. *Journal of aging and health*. 2011;23(2):228-41. PMID: CN-00858523.

451. Joling KJ, van Marwijk HW, Smit F, et al. Does a family meetings intervention prevent depression and anxiety in family caregivers of dementia patients? A randomized trial. *PLoS ONE* [Electronic Resource]. 2012;7(1):e30936. doi: <https://dx.doi.org/10.1371/journal.pone.0030936>. PMID: 22303473.
452. Judge KS, Yarry SJ, Looman WJ, et al. Improved Strain and Psychosocial Outcomes for Caregivers of Individuals with Dementia: Findings from Project ANSWERS. *Gerontologist*. 2013 Apr;53(2):280-92. doi: <https://dx.doi.org/10.1093/geront/gns076>. PMID: 22899427.
453. Kajiyama B, Thompson LW, Eto-Iwase T, et al. Exploring the effectiveness of an internet-based program for reducing caregiver distress using the iCare Stress Management e-Training Program.[Erratum appears in *Aging Ment Health*. 2013;17(5):c1]. *Aging & Mental Health*. 2013;17(5):544-54. doi: <https://dx.doi.org/10.1080/13607863.2013.775641>. PMID: 23461355.
454. Kuo LM, Huang HL, Huang HL, et al. A home-based training program improves Taiwanese family caregivers' quality of life and decreases their risk for depression: a randomized controlled trial. *International Journal of Geriatric Psychiatry*. 2013 May;28(5):504-13. doi: <https://dx.doi.org/10.1002/gps.3853>. PMID: 22778053.
455. Moore RC, Chattillion EA, Ceglowski J, et al. A randomized clinical trial of Behavioral Activation (BA) therapy for improving psychological and physical health in dementia caregivers: results of the Pleasant Events Program (PEP). *Behaviour Research & Therapy*. 2013 Oct;51(10):623-32. doi: <https://dx.doi.org/10.1016/j.brat.2013.07.005>. PMID: 23916631.
456. Gavrilova SI, Ferri CP, Mikhaylova N, et al. Helping carers to care--the 10/66 dementia research group's randomized control trial of a caregiver intervention in Russia. *International Journal of Geriatric Psychiatry*. 2009 Apr;24(4):347-54. doi: <https://dx.doi.org/10.1002/gps.2126>. PMID: 18814197.
457. Gitlin LN, Winter L, Dennis MP, et al. A biobehavioral home-based intervention and the well-being of patients with dementia and their caregivers: the COPE randomized trial. *JAMA*. 2010 Sep 01;304(9):983-91. doi: <https://dx.doi.org/10.1001/jama.2010.1253>. PMID: 20810376.
458. Kurz A, Wagenpfeil S, Hallauer J, et al. Evaluation of a brief educational program for dementia carers: the AENEAS study. *International Journal of Geriatric Psychiatry*. 2010 Aug;25(8):861-9. doi: <https://dx.doi.org/10.1002/gps.2428>. PMID: 19946869.
459. Losada A, Marquez-Gonzalez M, Romero-Moreno R. Mechanisms of action of a psychological intervention for dementia caregivers: effects of behavioral activation and modification of dysfunctional thoughts. *International Journal of Geriatric Psychiatry*. 2011 Nov;26(11):1119-27. doi: <https://dx.doi.org/10.1002/gps.2648>. PMID: 21061414.
460. Marquez-Gonzalez M, Losada A, Izal M, et al. Modification of dysfunctional thoughts about caregiving in dementia family caregivers: description and outcomes of an intervention programme. *Aging & Mental Health*. 2007 Nov;11(6):616-25. PMID: 18074249.
461. Williams V, Bishop-Fitzpatrick L, Lane J, et al. Video-Based Coping Skills to Reduce Health Risk and Improve Psychological and Physical Well-Being in Alzheimer's Disease Family Caregivers. *Psychosomatic medicine*. 2010;72(9):897-904. doi: 10.1097/PSY.0b013e3181fc2d09. PMID: CN-00892462.

462. Wilz G, Schinkothe D, Soellner R. Goal attainment and treatment compliance in a cognitive-behavioral telephone intervention for family caregivers of persons with dementia. *GeroPsych: The Journal of Gerontopsychology and Geriatric Psychiatry*. 2011 01 Sep;24(3):115-25. doi: <http://dx.doi.org/10.1024/1662-9647/a000043>. PMID: 362501128.
463. Beauchamp N, Irvine AB, Seeley J, et al. Worksite-based internet multimedia program for family caregivers of persons with dementia. *Gerontologist*. 2005 Dec;45(6):793-801. PMID: 16326661.
464. Burgio L, Stevens A, Guy D, et al. Impact of two psychosocial interventions on white and African American family caregivers of individuals with dementia. *Gerontologist*. 2003 Aug;43(4):568-79. PMID: 12937335.
465. Burns R, Nichols LO, Martindale-Adams J, et al. Primary care interventions for dementia caregivers: 2-year outcomes from the REACH study. *Gerontologist*. 2003 Aug;43(4):547-55. PMID: 12937333.
466. Coon DW, Thompson L, Steffen A, et al. Anger and depression management: psychoeducational skill training interventions for women caregivers of a relative with dementia. *Gerontologist*. 2003 Oct;43(5):678-89. PMID: 14570964.
467. Gitlin LN, Winter L, Corcoran M, et al. Effects of the home environmental skill-building program on the caregiver-care recipient dyad: 6-month outcomes from the Philadelphia REACH Initiative. *Gerontologist*. 2003 Aug;43(4):532-46. PMID: 12937332.
468. Gonyea JG, O'Connor MK, Boyle PA. Project CARE: a randomized controlled trial of a behavioral intervention group for Alzheimer's disease caregivers. *Gerontologist*. 2006 Dec;46(6):827-32. PMID: 17169938.
469. Ulstein ID, Sandvik L, Wyller TB, et al. A one-year randomized controlled psychosocial intervention study among family carers of dementia patients-- effects on patients and carers. *Dementia & Geriatric Cognitive Disorders*. 2007;24(6):469-75. PMID: 17986818.
470. Buckwalter KC, Gerdner L, Kohout F, et al. A nursing intervention to decrease depression in family caregivers of persons with dementia. *Archives of Psychiatric Nursing*. 1999 Apr;13(2):80-8. PMID: 10222636.
471. Fung W, Chien W. The effectiveness of a mutual support group for family caregivers of a relative with dementia. *Archives of Psychiatric Nursing*. 2002;16(3):134-44. PMID: 106975399. Language: English. Entry Date: 20021108. Revision Date: 20150819. Publication Type: Journal Article.
472. Hebert R, Levesque L, Vezina J, et al. Efficacy of a psychoeducative group program for caregivers of demented persons living at home: a randomized controlled trial. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences*. 2003 Jan;58(1):S58-67. PMID: 12496309.
473. Ostwald SK, Hepburn KW, Caron W, et al. Reducing caregiver burden: a randomized psychoeducational intervention for caregivers of persons with dementia. *Gerontologist*. 1999 Jun;39(3):299-309. PMID: 10396888.
474. Wright LK, Litaker M, Laraia MT, et al. Continuum of care for Alzheimer's disease: a nurse education and counseling program. *Issues in Mental Health Nursing*. 2001 Apr-May;22(3):231-52. PMID: 11885210.
475. Terracciano A, Artese A, Yeh J, et al. Effectiveness of Powerful Tools for Caregivers on Caregiver Burden and on Care Recipient Behavioral and Psychological Symptoms of Dementia: A Randomized Controlled Trial. *Journal of the American Medical Directors Association*. 2019 Dec 19;19:19. doi: <https://dx.doi.org/10.1016/j.jamda.2019.11.011>. PMID: 31866419.

476. Uyar F, Ozmen D, Mavioglu H, et al. Assessment of the impact of dementia care and support program in both patient and caregiver outcomes: An intervention study. *Turk Geriatri Dergisi*. 2019;22(4):482-93.
477. Kunik ME, Stanley MA, Shrestha S, et al. Aggression Prevention Training for Individuals With Dementia and Their Caregivers: A Randomized Controlled Trial. *American Journal of Geriatric Psychiatry*. 2020 Feb 03;03:03. doi: <https://dx.doi.org/10.1016/j.jagp.2020.01.190>. PMID: 32115311.
478. Rabinowitz YG, Mausbach BT, Coon DW, et al. The moderating effect of self-efficacy on intervention response in women family caregivers of older adults with dementia. *American Journal of Geriatric Psychiatry*. 2006 Aug;14(8):642-9. PMID: 16861368.
479. Mahoney DF, Tarlow BJ, Jones RN. Effects of an automated telephone support system on caregiver burden and anxiety: findings from the REACH for TLC intervention study. *Gerontologist*. 2003 Aug;43(4):556-67. PMID: 12937334.
480. Mahoney DM, Tarlow B, Jones RN, et al. Factors affecting the use of a telephone-based intervention for caregivers of people with Alzheimer's disease. *Journal of Telemedicine & Telecare*. 2001;7(3):139-48. PMID: 11346473.
481. Wang LQ, Chien WT, Lee IY. An experimental study on the effectiveness of a mutual support group for family caregivers of a relative with dementia in mainland China. *Contemporary Nurse*. 2012 Feb;40(2):210-24. doi: <https://dx.doi.org/10.5172/conu.2012.40.2.210>. PMID: 22554214.
482. Charlesworth G, Shepstone L, Wilson E, et al. Befriending carers of people with dementia: randomised controlled trial. *BMJ*. 2008 Jun 07;336(7656):1295-7. doi: <https://dx.doi.org/10.1136/bmj.39549.548831.AE>. PMID: 18505757.
483. Charlesworth G, Shepstone L, Wilson E, et al. Does befriending by trained lay workers improve psychological well-being and quality of life for carers of people with dementia, and at what cost? A randomised controlled trial. *Health Technology Assessment (Winchester, England)*. 2008 Mar;12(4):iii, v-ix, 1-78. PMID: 18284895.
484. Goodman C. Evaluation of a model self-help telephone program: impact on natural networks. *Social Work*. 1990 Nov;35(6):556-62. PMID: 2284602.
485. Logsdon RG, Pike KC, McCurry SM, et al. Early-stage memory loss support groups: outcomes from a randomized controlled clinical trial. *Journals of Gerontology Series B: Psychological Sciences & Social Sciences*. 2010;65B(6):691-7. doi: [geronb/gbq054](https://doi.org/10.1093/geronb/gbq054). PMID: 104963362. Language: English. Entry Date: 20110128. Revision Date: 20150820. Publication Type: Journal Article.
486. Metcalfe A, Jones B, Mayer J, et al. Online information and support for carers of people with young-onset dementia: A multi-site randomised controlled pilot study. *International Journal of Geriatric Psychiatry*. 2019 01 Oct;34(10):1455-64. doi: <http://dx.doi.org/10.1002/gps.5154>. PMID: 2002845656.
487. Pillemer K, Suiitor JJ. Peer support for Alzheimer's caregivers: Is it enough to make a difference? *Research on Aging*. 2002;24(2):171-92. PMID: 34163761.
488. Torkamani M, McDonald L, Saez Aguayo I, et al. A randomized controlled pilot study to evaluate a technology platform for the assisted living of people with dementia and their carers. *Journal of Alzheimer's Disease*. 2014;41(2):515-23. doi: <https://dx.doi.org/10.3233/JAD-132156>. PMID: 24643137.

489. van der Roest HG, Meiland FJ, Jonker C, et al. User evaluation of the DEMentia-specific Digital Interactive Social Chart (DEM-DISC). A pilot study among informal carers on its impact, user friendliness and, usefulness. *Aging & Mental Health*. 2010 May;14(4):461-70. doi: <https://dx.doi.org/10.1080/13607860903311741>. PMID: 20455122.
490. Van Mierlo LD, Meiland FJ, Van de Ven PM, et al. Evaluation of DEM-DISC, customized e-advice on health and social support services for informal carers and case managers of people with dementia; a cluster randomized trial. *International Psychogeriatrics*. 2015 Aug;27(8):1365-78. doi: <https://dx.doi.org/10.1017/S1041610215000423>. PMID: 25872457.
491. Wilson E, Thalanany M, Shepstone L, et al. Befriending carers of people with dementia: a cost utility analysis. *International Journal of Geriatric Psychiatry*. 2009 Jun;24(6):610-23. doi: <https://dx.doi.org/10.1002/gps.2164>. PMID: 19101921.
492. Winter L, Gitlin LN. Evaluation of a telephone-based support group intervention for female caregivers of community-dwelling individuals with dementia. *American Journal of Alzheimer's Disease & Other Dementias*. 2006 Dec-2007 Jan;21(6):391-7. PMID: 17267370.
493. Huis In Het Veld JG, Willemse BM, van Asch IF, et al. Online Self-Management Support for Family Caregivers Dealing With Behavior Changes in Relatives With Dementia (Part 2): Randomized Controlled Trial. *Journal of Medical Internet Research*. 2020 Feb 25;22(2):e13001. doi: <https://dx.doi.org/10.2196/13001>. PMID: 32130142.
494. Gustafson DH, Cody OJ, Chih MY, et al. Pilot Test of a Computer-Based System to Help Family Caregivers of Dementia Patients. *Journal of Alzheimer's Disease*. 2019;70(2):539-50. doi: <http://dx.doi.org/10.3233/JAD-190052>. PMID: 628746285.
495. Whitebird RR, Kreitzer M, Crain AL, et al. Mindfulness-based stress reduction for family caregivers: a randomized controlled trial. *Gerontologist*. 2013 Aug;53(4):676-86. doi: <https://dx.doi.org/10.1093/geront/gns126>. PMID: 23070934.
496. Brown KW, Coogle CL, Wegelin J. A pilot randomized controlled trial of mindfulness-based stress reduction for caregivers of family members with dementia. *Aging & Mental Health*. 2016 Nov;20(11):1157-66. PMID: 26211415.
497. Danucalov MA, Kozasa EH, Afonso RF, et al. Yoga and compassion meditation program improve quality of life and self-compassion in family caregivers of Alzheimer's disease patients: A randomized controlled trial. *Geriatrics & gerontology international*. 2017 Jan;17(1):85-91. doi: <https://dx.doi.org/10.1111/ggi.12675>. PMID: 26685923.
498. Kor PPK, Liu JYW, Chien WT. Effects of a modified mindfulness-based cognitive therapy for family caregivers of people with dementia: A pilot randomized controlled trial. *International Journal of Nursing Studies*. 2019 Oct;98:107-17. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2019.02.020>. PMID: 30922609.
499. Lavretsky H, Epel ES, Siddarth P, et al. A pilot study of yogic meditation for family dementia caregivers with depressive symptoms: effects on mental health, cognition, and telomerase activity. *International Journal of Geriatric Psychiatry*. 2013 Jan;28(1):57-65. doi: <https://dx.doi.org/10.1002/gps.3790>. PMID: 22407663.
500. Leach MJ, Francis A, Ziaian T. Transcendental Meditation for the improvement of health and wellbeing in community-dwelling dementia caregivers [TRANSCENDENT]: a randomised wait-list controlled trial. *BMC Complementary & Alternative Medicine*. 2015 May 08;15:145. doi: <https://dx.doi.org/10.1186/s12906-015-0666-8>. PMID: 25952550.

501. Oken BS, Fonareva I, Haas M, et al. Pilot controlled trial of mindfulness meditation and education for dementia caregivers. *Journal of Alternative & Complementary Medicine*. 2010 Oct;16(10):1031-8. doi: <https://dx.doi.org/10.1089/acm.2009.0733>. PMID: 20929380.
502. Waelde LC, Meyer H, Thompson JM, et al. Randomized Controlled Trial of Inner Resources Meditation for Family Dementia Caregivers. *Journal of Clinical Psychology*. 2017 Dec;73(12):1629-41. doi: <https://dx.doi.org/10.1002/jclp.22470>. PMID: 28263398.
503. Mahdavi B, Fallahi-Khoshknab M, Mohammadi F, et al. Effects of spiritual group therapy on caregiver strain in home caregivers of the elderly with Alzheimer's disease. *Archives of psychiatric nursing*. 2016. PMID: CN-01308784.
504. Salamizadeh AMS, Mirzaei TP, Ravari AP. The Impact of Spiritual Care Education on the Self-Efficacy of the Family Caregivers of Elderly People with Alzheimer's Disease. *International Journal of Community Based Nursing & Midwifery*. 2017 Jul;5(3):231-8. PMID: 28670585.
505. Castro CM, Wilcox S, O'Sullivan P, et al. An exercise program for women who are caring for relatives with dementia. *Psychosomatic Medicine*. 2002 May-Jun;64(3):458-68. PMID: 12021419.
506. Connell CM, Janevic MR. Effects of a Telephone-Based Exercise Intervention for Dementia Caregiving Wives: A Randomized Controlled Trial. *Journal of Applied Gerontology*. 2009 Apr;28(2):171-94. PMID: 21709757.
507. Farran CJ, Etkin CD, Eisenstein A, et al. Effect of Moderate to Vigorous Physical Activity Intervention on Improving Dementia Family Caregiver Physical Function: A Randomized Controlled Trial. *Journal of Alzheimer's Disease & Parkinsonism*. 2016 Aug;6(4). doi: <https://dx.doi.org/10.4172/2161-0460.1000253>. PMID: 28752016.
508. Hirano A, Suzuki Y, Kuzuya M, et al. Influence of regular exercise on subjective sense of burden and physical symptoms in community-dwelling caregivers of dementia patients: a randomized controlled trial. *Archives of Gerontology & Geriatrics*. 2011 Sep-Oct;53(2):e158-63. doi: <https://dx.doi.org/10.1016/j.archger.2010.08.004>. PMID: 20850878.
509. Lowery D, Cerga-Pashoja A, Iliffe S, et al. The effect of exercise on behavioural and psychological symptoms of dementia: the EVIDEM-E randomised controlled clinical trial. *International Journal of Geriatric Psychiatry*. 2014 Aug;29(8):819-27. doi: <https://dx.doi.org/10.1002/gps.4062>. PMID: 24338799.
510. Hirano A, Umegaki H, Suzuki Y, et al. Effects of leisure activities at home on perceived care burden and the endocrine system of caregivers of dementia patients: a randomized controlled study. *International Psychogeriatrics*. 2016 Feb;28(2):261-8. doi: <https://dx.doi.org/10.1017/S1041610215001295>. PMID: 26289496.
511. Richards AG, Tietyen AC, Jicha GA, et al. Visual Arts Education Improves self-esteem for persons with dementia and reduces caregiver burden: A randomized controlled trial. *Dementia*. 2018 Jan 01:1471301218769071. doi: <https://dx.doi.org/10.1177/1471301218769071>. PMID: 29653492.
512. Wilz G, Fink-Heitz M. Assisted vacations for men with dementia and their caregiving spouses: evaluation of health-related effects. *Gerontologist*. 2008 Feb;48(1):115-20. PMID: 18381838.
513. Carbonneau H, Caron CD, Desrosiers J. Effects of an adapted leisure education program as a means of support for caregivers of people with dementia. *Archives of Gerontology & Geriatrics*. 2011 Jul-Aug;53(1):31-9. doi: <https://dx.doi.org/10.1016/j.archger.2010.06.009>. PMID: 20598758.

514. Lawton MP, Brody EM, Saperstein A, et al. Respite services for caregivers: research findings for service planning. *Home Health Care Services Quarterly*. 1989;10(1-2):5-32. PMID: 10318337.
515. Vandepitte S, Putman K, Van Den Noortgate N, et al. Effectiveness of an in-home respite care program to support informal dementia caregivers: A comparative study. *International Journal of Geriatric Psychiatry*. 2019 Oct;34(10):1534-44. doi: <https://dx.doi.org/10.1002/gps.5164>. PMID: 31243801.
516. Vandepitte S, Van Den Noortgate N, Putman K, et al. Effectiveness and cost-effectiveness of an in-home respite care program in supporting informal caregivers of people with dementia: design of a comparative study. *BMC Geriatrics*. 2016 12 02;16(1):207. PMID: 27912740.
517. Zarit SH, Stephens MA, Townsend A, et al. Stress reduction for family caregivers: effects of adult day care use. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences*. 1998 Sep;53(5):S267-77. PMID: 9750575.
518. Baglio F, Griffanti L, Saibene FL, et al. Multistimulation group therapy in Alzheimer's disease promotes changes in brain functioning. *Neurorehabilitation & Neural Repair*. 2015 Jan;29(1):13-24. doi: <https://dx.doi.org/10.1177/1545968314532833>. PMID: 24788581.
519. Belle SH, Burgio L, Burns R, et al. Enhancing the quality of life of dementia caregivers from different ethnic or racial groups: a randomized, controlled trial.[Summary for patients in *Ann Intern Med*. 2006 Nov 21;145(10):I39; PMID: 17116914]. *Annals of Internal Medicine*. 2006 Nov 21;145(10):727-38. PMID: 17116917.
520. Berwig M, Heinrich S, Spahlholz J, et al. Individualized support for informal caregivers of people with dementia - effectiveness of the German adaptation of REACH II. *BMC Geriatrics*. 2017 Dec 12;17(1):286. doi: <https://dx.doi.org/10.1186/s12877-017-0678-y>. PMID: 29233097.
521. Brijoux T, Kricheldorf C, M HL, et al. Supporting Families Living With Dementia in Rural Areas. *Deutsches Arzteblatt International*. 2016 Oct 14;113(41):681-7. doi: <https://dx.doi.org/10.3238/arztebl.2016.0681>. PMID: 27839534.
522. Czaja SJ, Loewenstein D, Schulz R, et al. A videophone psychosocial intervention for dementia caregivers. *American Journal of Geriatric Psychiatry*. 2013 Nov;21(11):1071-81. doi: <https://dx.doi.org/10.1016/j.jagp.2013.02.019>. PMID: 23831174.
523. Ducharme F, Levesque L, Giroux F, et al. Follow-up of an intervention program for caregivers of a relative with dementia living in a long-term care setting: Are there any persistent and delayed effects? *Aging & Mental Health*. 2005 Sep;9(5):461-9. PMID: 16024406.
524. Eisdorfer C, Czaja SJ, Loewenstein DA, et al. The effect of a family therapy and technology-based intervention on caregiver depression. *Gerontologist*. 2003 Aug;43(4):521-31. PMID: 12937331.
525. Finkel S, Czaja SJ, Schulz R, et al. E-care: a telecommunications technology intervention for family caregivers of dementia patients. *American Journal of Geriatric Psychiatry*. 2007 May;15(5):443-8. PMID: 17463195.
526. Gaugler JE, Reese M, Mittelman MS. Effects of the NYU caregiver intervention-adult child on residential care placement. *Gerontologist*. 2013 Dec;53(6):985-97. doi: <https://dx.doi.org/10.1093/geront/gns193>. PMID: 23339050.
527. Koivisto AM, Hallikainen I, Valimaki T, et al. Early psychosocial intervention does not delay institutionalization in persons with mild Alzheimer disease and has impact on neither disease progression nor caregivers' well-being: ALSOVA 3-year follow-up. *International Journal of Geriatric Psychiatry*. 2016 Mar;31(3):273-83. doi: <https://dx.doi.org/10.1002/gps.4321>. PMID: 26177825.

528. Luchsinger JA, Burgio L, Mittelman M, et al. Comparative Effectiveness of 2 Interventions for Hispanic Caregivers of Persons with Dementia. *Journal of the American Geriatrics Society*. 2018 Sep;66(9):1708-15. doi: <https://dx.doi.org/10.1111/jgs.15450>. PMID: 30084133.
529. Marshall A, Spreadbury J, Cheston R, et al. A pilot randomised controlled trial to compare changes in quality of life for participants with early diagnosis dementia who attend a "Living Well with Dementia" group compared to waiting-list control. *Aging & mental health*. 2015;19(6):526-35. PMID: CN-01071410.
530. Martindale-Adams J, Nichols LO, Burns R, et al. A trial of dementia caregiver telephone support. *Canadian Journal of Nursing Research*. 2013 Dec;45(4):30-48. PMID: 24617278.
531. Mittelman MS, Brodaty H, Wallen AS, et al. A three-country randomized controlled trial of a psychosocial intervention for caregivers combined with pharmacological treatment for patients with Alzheimer disease: effects on caregiver depression. *American Journal of Geriatric Psychiatry*. 2008;16(11):893-904. doi: 10.1097/JGP.0b013e3181898095. PMID: 105627901. Language: English. Entry Date: 20090130. Revision Date: 20161207. Publication Type: journal article.
532. Mittelman MS, Ferris SH, Steinberg G, et al. An intervention that delays institutionalization of Alzheimer's disease patients: treatment of spouse-caregivers. *Gerontologist*. 1993 Dec;33(6):730-40. PMID: 8314099.
533. Mittelman MS, Roth DL, Haley WE, et al. Effects of a caregiver intervention on negative caregiver appraisals of behavior problems in patients with Alzheimer's disease: results of a randomized trial. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences*. 2004 Jan;59(1):P27-34. PMID: 14722336.
534. Mohide EA, Pringle DM, Streiner DL, et al. A randomized trial of family caregiver support in the home management of dementia. *Journal of the American Geriatrics Society*. 1990 Apr;38(4):446-54. PMID: 2184186.
535. Prick AE, de Lange J, Twisk J, et al. The effects of a multi-component dyadic intervention on the psychological distress of family caregivers providing care to people with dementia: a randomized controlled trial. *International Psychogeriatrics*. 2015 Dec;27(12):2031-44. doi: <https://dx.doi.org/10.1017/S104161021500071X>. PMID: 26004290.
536. Signe A, Elmståhl S. Psychosocial intervention for family caregivers of people with dementia reduces caregiver's burden: development and effect after 6 and 12 months. *Scandinavian journal of caring sciences*. 2008;22(1):98-109. doi: 10.1111/j.1471-6712.2007.00498.x. PMID: CN-00630033.
537. Teri L, Gibbons LE, McCurry SM, et al. Exercise plus behavioral management in patients with Alzheimer disease: a randomized controlled trial. *JAMA*. 2003 Oct 15;290(15):2015-22. PMID: 14559955.
538. Zhang SY, Wu F, Tang DL, et al. Pilot testing the caregiver self-management intervention for caregivers of relatives with dementia. *Geriatric Nursing*. 2019 Aug 26;26:26. doi: <https://dx.doi.org/10.1016/j.gerinurse.2019.08.006>. PMID: 31466807.
539. Correction: northern Manhattan Hispanic Caregiver Intervention Effectiveness Study: protocol of a pragmatic randomised trial comparing the effectiveness of two established interventions for informal caregivers of persons with dementia (*BMJ Open* (2016) 6 (e014082) DOI: 10.1136/bmjopen-2016-014082). In *BMJ open*

540. Mazurek J, Szczesniak D, Lion KM, et al. Does the Meeting Centres Support Programme reduce unmet care needs of community-dwelling older people with dementia? A controlled, 6-month follow-up Polish study. *Clinical Interventions In Aging*. 2019;14:113-22. doi: <https://dx.doi.org/10.2147/CIA.S185683>. PMID: 30666097.
541. Teri L, Gibbons LE, McCurry SM, et al. Exercise plus behavioral management in patients with Alzheimer disease: a randomized controlled trial. *JAMA : the journal of the American Medical Association*. 2003;290(15):2015-22. doi: 10.1001/jama.290.15.2015.
542. Jutten LH, Mark RE, Sitskoorn MM. Can the Mixed Virtual Reality Simulator Into Dementia Enhance Empathy and Understanding and Decrease Burden in Informal Dementia Caregivers? *Dementia and Geriatric Cognitive Disorders Extra*. 2018 Sep-Dec;8(3):453-66. doi: <https://dx.doi.org/10.1159/000494660>. PMID: 30631337.
543. Lord K, Livingston G, Cooper C. A feasibility randomised controlled trial of the DECIDE intervention: dementia carers making informed decisions. *Bjpsych open*. 2017;3(1):12-4. PMID: CN-01415827.
544. Rose KM, Taylor AG, Bourguignon C. Effects of cranial electrical stimulation on sleep disturbances, depressive symptoms, and caregiving appraisal in spousal caregivers of persons with Alzheimer's disease. *Applied Nursing Research*. 2009 May;22(2):119-25. doi: <https://dx.doi.org/10.1016/j.apnr.2007.06.001>. PMID: 19427574.
545. Simpson R, Wakefield P, Spiers N, et al. Carer-held records for dementia: a controlled trial. *International Psychogeriatrics*. 2006 Jun;18(2):259-68. PMID: 16734920.
546. Stern RA, D'Ambrosio LA, Mohyde M, et al. At the crossroads: development and evaluation of a dementia caregiver group intervention to assist in driving cessation. *Gerontology & Geriatrics Education*. 2008;29(4):363-82. doi: <https://dx.doi.org/10.1080/02701960802497936>. PMID: 19064472.
547. Stirling C, Leggett S, Lloyd B, et al. Decision aids for respite service choices by carers of people with dementia: development and pilot RCT. *BMC Medical Informatics & Decision Making*. 2012 Mar 19;12:21. doi: <https://dx.doi.org/10.1186/1472-6947-12-21>. PMID: 22429384.
548. Barbosa A, Nolan M, Sousa L, et al. Supporting direct care workers in dementia care: effects of a psychoeducational intervention. *American Journal of Alzheimer's Disease & Other Dementias*. 2015 Mar;30(2):130-8. doi: <https://dx.doi.org/10.1177/1533317514550331>. PMID: 25237132.
549. Barbosa A, Nolan M, Sousa L, et al. Effects of a Psychoeducational Intervention for Direct Care Workers Caring for People With Dementia: Results From a 6-Month Follow-Up Study. *American Journal of Alzheimer's Disease & Other Dementias*. 2016 Mar;31(2):144-55. doi: <https://dx.doi.org/10.1177/1533317515603500>. PMID: 26399485.
550. Davison T, McCabe M, Visser S, et al. Controlled trial of dementia training with a peer support group for aged care staff. *International journal of geriatric psychiatry*. 2007;22(9):868-73. doi: 10.1002/gps.1754. PMID: CN-00627010.
551. Mackenzie CS, Peragine G. Measuring and enhancing self-efficacy among professional caregivers of individuals with dementia. *American Journal of Alzheimer's Disease & Other Dementias*. 2003 Sep-Oct;18(5):291-9. PMID: 14569646.

552. Visser SM, McCabe MP, Hudgson C, et al. Managing behavioural symptoms of dementia: effectiveness of staff education and peer support. *Aging & Mental Health*. 2008 Jan;12(1):47-55. doi: <https://dx.doi.org/10.1080/13607860701366012>. PMID: 18297478.
553. Borbasi S, Emmanuel E, Farrelly B, et al. Report of an evaluation of a nurse-led dementia outreach service for people with the behavioural and psychological symptoms of dementia living in residential aged care facilities. *Perspectives in public health*. 2011;131(3):124-30. PMID: CN-00850682.
554. Fortinsky RH, Kulldorff M, Kleppinger A, et al. Dementia care consultation for family caregivers: collaborative model linking an Alzheimer's association chapter with primary care physicians. *Aging & Mental Health*. 2009 Mar;13(2):162-70. doi: <https://dx.doi.org/10.1080/13607860902746160>. PMID: 19347683.
555. McSweeney K, Jeffreys A, Griffith J, et al. Specialist mental health consultation for depression in Australian aged care residents with dementia: a cluster randomized trial. *International Journal of Geriatric Psychiatry*. 2012 Nov;27(11):1163-71. doi: <https://dx.doi.org/10.1002/gps.3762>. PMID: 22344753.
556. Opie J, Doyle C, O'Connor DW. Challenging behaviours in nursing home residents with dementia: a randomized controlled trial of multidisciplinary interventions. *International Journal of Geriatric Psychiatry*. 2002 Jan;17(1):6-13. PMID: 11802224.
557. Orrell M, Hancock G, Hoe J, et al. A cluster randomised controlled trial to reduce the unmet needs of people with dementia living in residential care. *International Journal of Geriatric Psychiatry*. 2007 Nov;22(11):1127-34. PMID: 17394129.
558. Chien WT, Lee IY. Randomized controlled trial of a dementia care programme for families of home-resided older people with dementia. *Journal of Advanced Nursing*. 2011 Apr;67(4):774-87. doi: <https://dx.doi.org/10.1111/j.1365-2648.2010.05537.x>. PMID: 21198803.
559. Chien WT, Lee YM. A disease management program for families of persons in Hong Kong with dementia. *Psychiatric Services*. 2008 Apr;59(4):433-6. doi: <https://dx.doi.org/10.1176/ps.2008.59.4.433>. PMID: 18378844.
560. Eloniemi-Sulkava U, Notkola IL, Hentinen M, et al. Effects of supporting community-living demented patients and their caregivers: a randomized trial. *Journal of the American Geriatrics Society*. 2001 Oct;49(10):1282-7. PMID: 11890485.
561. Lam LC, Lee JS, Chung JC, et al. A randomized controlled trial to examine the effectiveness of case management model for community dwelling older persons with mild dementia in Hong Kong. *International Journal of Geriatric Psychiatry*. 2010 Apr;25(4):395-402. doi: <https://dx.doi.org/10.1002/gps.2352>. PMID: 19606455.
562. Brodaty H, Draper BM, Millar J, et al. Randomized controlled trial of different models of care for nursing home residents with dementia complicated by depression or psychosis. *Journal of Clinical Psychiatry*. 2003 Jan;64(1):63-72. PMID: 12590626.
563. Challis D, von Abendorff R, Brown P, et al. Care management, dementia care and specialist mental health services: an evaluation. *International Journal of Geriatric Psychiatry*. 2002 Apr;17(4):315-25. PMID: 11994884.
564. MacNeil VJ, Bosmans J, Ven P, et al. Community-dwelling patients with dementia and their informal caregivers with and without case management: 2-year outcomes of a pragmatic trial. In *Journal of the American Medical Directors Association*

565. Newcomer R, Spitalny M, Fox P, et al. Effects of the Medicare Alzheimer's Disease Demonstration on the use of community-based services. *Health Services Research*. 1999 Aug;34(3):645-67. PMID: 10445896.
566. Weinberger M, Gold DT, Divine GW, et al. Social service interventions for caregivers of patients with dementia: impact on health care utilization and expenditures. *Journal of the American Geriatrics Society*. 1993 Feb;41(2):153-6. PMID: 8426038.
567. Xiao LD, De Bellis A, Kyriazopoulos H, et al. The Effect of a Personalized Dementia Care Intervention for Caregivers From Australian Minority Groups. *American Journal of Alzheimer's Disease & Other Dementias*. 2016 Feb;31(1):57-67. doi: <https://dx.doi.org/10.1177/1533317515578256>. PMID: 25805891.
568. Appelhof B, Bakker C, de Vugt ME, et al. Effects of a Multidisciplinary Intervention on the Presence of Neuropsychiatric Symptoms and Psychotropic Drug Use in Nursing Home Residents With Young-Onset Dementia: Behavior and Evolution of Young-Onset Dementia Part 2 (BEYOND-II) Study. *American Journal of Geriatric Psychiatry*. 2019 Jan 04;04:04. doi: <https://dx.doi.org/10.1016/j.jagp.2018.12.032>. PMID: 30799167.
569. Chen YH, Lin LC. Ability of the Pain Recognition and Treatment (PRT) Protocol to Reduce Expressions of Pain among Institutionalized Residents with Dementia: A Cluster Randomized Controlled Trial. *Pain Management Nursing*. 2016 Feb;17(1):14-24. doi: <https://dx.doi.org/10.1016/j.pmn.2015.08.003>. PMID: 26584896.
570. Cohen-Mansfield J, Libin A, Marx MS. Nonpharmacological treatment of agitation: a controlled trial of systematic individualized intervention. *Journals of Gerontology Series A-Biological Sciences & Medical Sciences*. 2007 Aug;62(8):908-16. PMID: 17702884.
571. Cohen-Mansfield J, Thein K, Marx MS, et al. Efficacy of nonpharmacologic interventions for agitation in advanced dementia: a randomized, placebo-controlled trial. *Journal of Clinical Psychiatry*. 2012 Sep;73(9):1255-61. doi: <https://dx.doi.org/10.4088/JCP.12m07918>. PMID: 23059151.
572. Kovach C, Simpson M, Joosse L, et al. Comparison of the effectiveness of two protocols for treating nursing home residents with advanced dementia. In *Research in gerontological nursing*
573. Kovach CR, Hekel B, Rababa M. Feasibility Testing of a Protocol to Stop Ineffective Drug and Nondrug Treatments. *Western Journal of Nursing Research*. 2015 Nov;37(11):1404-22. doi: <https://dx.doi.org/10.1177/0193945915599070>. PMID: 26250849.
574. Kovach CR, Logan BR, Noonan PE, et al. Effects of the Serial Trial Intervention on discomfort and behavior of nursing home residents with dementia. *American Journal of Alzheimer's Disease & Other Dementias*. 2006 Jun-Jul;21(3):147-55. PMID: 16869334.
575. McCabe MP, Bird M, Davison TE, et al. An RCT to evaluate the utility of a clinical protocol for staff in the management of behavioral and psychological symptoms of dementia in residential aged-care settings. *Aging & Mental Health*. 2015;19(9):799-807. doi: <https://dx.doi.org/10.1080/13607863.2014.967659>. PMID: 25319535.
576. Montgomery RJ, Kwak J, Kosloski K, et al. Effects of the TCARE intervention on caregiver burden and depressive symptoms: preliminary findings from a randomized controlled study. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences*. 2011 Sep;66(5):640-7. doi: <https://dx.doi.org/10.1093/geronb/gbr088>. PMID: 21840840.

577. Nakanishi M, Endo K, Hirooka K, et al. Psychosocial behaviour management programme for home-dwelling people with dementia: A cluster-randomized controlled trial. *International Journal of Geriatric Psychiatry*. 2018 March;33(3):495-503. doi: <http://dx.doi.org/10.1002/gps.4784>. PMID: 620640680 Pubmed 28857263.
578. Pieper MJ, Francke AL, van der Steen JT, et al. Effects of a Stepwise Multidisciplinary Intervention for Challenging Behavior in Advanced Dementia: A Cluster Randomized Controlled Trial. [Erratum appears in *J Am Geriatr Soc*. 2016 Jun;64(6):1383; PMID: 27321637]. *Journal of the American Geriatrics Society*. 2016 Feb;64(2):261-9. doi: <https://dx.doi.org/10.1111/jgs.13868>. PMID: 26804064.
579. Rapp MA, Mell T, Majic T, et al. Agitation in nursing home residents with dementia (VIDEANT trial): effects of a cluster-randomized, controlled, guideline implementation trial. *Journal of the American Medical Directors Association*. 2013 Sep;14(9):690-5. doi: <https://dx.doi.org/10.1016/j.jamda.2013.05.017>. PMID: 23827658.
580. Salva A, Andrieu S, Fernandez E, et al. Health and nutrition promotion program for patients with dementia (NutriAlz): Cluster randomized trial. *Journal of Nutrition, Health & Aging*. 2011;15(10):822-30. doi: 10.1007/s12603-011-0363-3. PMID: 104625164. Language: English. Entry Date: 20120530. Revision Date: 20150818. Publication Type: Journal Article.
581. Verkaik R, Francke AL, van Meijel B, et al. The effects of a nursing guideline on depression in psychogeriatric nursing home residents with dementia. *International Journal of Geriatric Psychiatry*. 2011 Jul;26(7):723-32. doi: <https://dx.doi.org/10.1002/gps.2586>. PMID: 21495077.
582. Zwijsen SA, Smalbrugge M, Eefsting JA, et al. Coming to grips with challenging behavior: a cluster randomized controlled trial on the effects of a multidisciplinary care program for challenging behavior in dementia. *Journal of the American Medical Directors Association*. 2014 Jul;15(7):531.e1-.e10. doi: <https://dx.doi.org/10.1016/j.jamda.2014.04.007>. PMID: 24878214.
583. Livingston G, Barber J, Marston L, et al. Clinical and cost-effectiveness of the Managing Agitation and Raising Quality of Life (MARQUE) intervention for agitation in people with dementia in care homes: a single-blind, cluster-randomised controlled trial. *The Lancet. Psychiatry*. 2019 04;6(4):293-304. doi: [https://dx.doi.org/10.1016/S2215-0366\(19\)30045-8](https://dx.doi.org/10.1016/S2215-0366(19)30045-8). PMID: 30872010.
584. Husebo BS, Ballard C, Aarsland D, et al. The Effect of a Multicomponent Intervention on Quality of Life in Residents of Nursing Homes: A Randomized Controlled Trial (COSMOS). *Journal of the American Medical Directors Association*. 2019. doi: <http://dx.doi.org/10.1016/j.jamda.2018.11.006>. PMID: 2001439020.
585. Cohen SM, Volandes AE, Shaffer ML, et al. Concordance Between Proxy Level of Care Preference and Advance Directives Among Nursing Home Residents With Advanced Dementia: A Cluster Randomized Clinical Trial. *Journal of Pain & Symptom Management*. 2019 Jan;57(1):37-46.e1. doi: <https://dx.doi.org/10.1016/j.jpainsymman.2018.09.018>. PMID: 30273717.
586. Mitchell SL, Shaffer ML, Cohen S, et al. An Advance Care Planning Video Decision Support Tool for Nursing Home Residents With Advanced Dementia: A Cluster Randomized Clinical Trial. *JAMA Internal Medicine*. 2018 Jul 01;178(7):961-9. doi: <https://dx.doi.org/10.1001/jamainternmed.2018.1506>. PMID: 29868778.

587. Bonner GJ, Wang E, Wilkie DJ, et al. Advance care treatment plan (ACT-Plan) for African American family caregivers: a pilot study. *Dementia*. 2014 Jan;13(1):79-95. doi: <https://dx.doi.org/10.1177/1471301212449408>. PMID: 24381040.
588. Brazil K, Carter G, Cardwell C, et al. Effectiveness of advance care planning with family carers in dementia nursing homes: A paired cluster randomized controlled trial. *Palliative Medicine*. 2018 Mar;32(3):603-12. doi: <https://dx.doi.org/10.1177/0269216317722413>. PMID: 28786323.
589. Einterz SF, Gilliam R, Lin FC, et al. Development and testing of a decision aid on goals of care for advanced dementia. *Journal of the American Medical Directors Association*. 2014 Apr;15(4):251-5. doi: <https://dx.doi.org/10.1016/j.jamda.2013.11.020>. PMID: 24508326.
590. Hanson LC, Song MK, Zimmerman S, et al. Fidelity to a behavioral intervention to improve goals of care decisions for nursing home residents with advanced dementia. *Clinical Trials*. 2016 Dec;13(6):599-604. PMID: 27271683.
591. Hanson LC, Zimmerman S, Song MK, et al. Effect of the Goals of Care Intervention for Advanced Dementia: A Randomized Clinical Trial. *JAMA Internal Medicine*. 2017 Jan 01;177(1):24-31. doi: <https://dx.doi.org/10.1001/jamainternmed.2016.7031>. PMID: 27893884.
592. Reinhardt JP, Chichin E, Posner L, et al. Vital conversations with family in the nursing home: preparation for end-stage dementia care. *Journal Of Social Work In End-Of-Life & Palliative Care*. 2014;10(2):112-26. doi: <https://dx.doi.org/10.1080/15524256.2014.906371>. PMID: 24835382.
593. Sampson EL, Jones L, Thune-Boyle IC, et al. Palliative assessment and advance care planning in severe dementia: an exploratory randomized controlled trial of a complex intervention. *Palliative Medicine*. 2011 Apr;25(3):197-209. doi: <https://dx.doi.org/10.1177/0269216310391691>. PMID: 21228087.
594. Song MK, Ward SE, Hepburn K, et al. Can Persons with Dementia Meaningfully Participate in Advance Care Planning Discussions? A Mixed-Methods Study of SPIRIT. *Journal of Palliative Medicine*. 2019 Nov;22(11):1410-6. doi: <https://dx.doi.org/10.1089/jpm.2019.0088>. PMID: 31373868.
595. Goossens B, Sevenants A, Declercq A, et al. 'We DECide optimized' - training nursing home staff in shared decision-making skills for advance care planning conversations in dementia care: protocol of a pretest-posttest cluster randomized trial. *BMC Geriatrics*. 2019 02 04;19(1):33. doi: <https://dx.doi.org/10.1186/s12877-019-1044-z>. PMID: 30717700.
596. Hanson LC, Carey TS, Caprio AJ, et al. Improving decision-making for feeding options in advanced dementia: a randomized, controlled trial. *Journal of the American Geriatrics Society*. 2011 Nov;59(11):2009-16. doi: <https://dx.doi.org/10.1111/j.1532-5415.2011.03629.x>. PMID: 22091750.
597. Hanson LC, Gilliam R, Lee TJ. Successful clinical trial research in nursing homes: the Improving Decision-Making Study. *Clinical Trials*. 2010 Dec;7(6):735-43. doi: <https://dx.doi.org/10.1177/1740774510380241>. PMID: 20729251.
598. Snyder EA, Caprio AJ, Wessell K, et al. Impact of a decision aid on surrogate decision-makers' perceptions of feeding options for patients with dementia. *Journal of the American Medical Directors Association*. 2013 Feb;14(2):114-8. doi: <https://dx.doi.org/10.1016/j.jamda.2012.10.011>. PMID: 23273855.
599. Agar M, Lockett T, Luscombe G, et al. Effects of facilitated family case conferencing for advanced dementia: A cluster randomised clinical trial. *PLoS ONE [Electronic Resource]*. 2017;12(8):e0181020. doi: <https://dx.doi.org/10.1371/journal.pone.0181020>. PMID: 28786995.

600. Agar M, Beattie E, Luckett T, et al. Pragmatic cluster randomised controlled trial of facilitated family case conferencing compared with usual care for improving end of life care and outcomes in nursing home residents with advanced dementia and their families: the IDEAL study protocol. *BMC Palliative Care*. 2015 Nov 21;14:63. doi: <https://dx.doi.org/10.1186/s12904-015-0061-8>. PMID: 26589957.
601. Ahronheim JC, Morrison RS, Morris J, et al. Palliative care in advanced dementia: a randomized controlled trial and descriptive analysis. *Journal of Palliative Medicine*. 2000;3(3):265-73. PMID: 15859668.
602. Boogaard JA, de Vet HC, van Soest-Poortvliet MC, et al. Effects of two feedback interventions on end-of-life outcomes in nursing home residents with dementia: A cluster-randomized controlled three-armed trial. *Palliative Medicine*. 2018 Mar;32(3):693-702. doi: <https://dx.doi.org/10.1177/0269216317750071>. PMID: 29343173.
603. Hanson LC, Kistler CE, Lavin K, et al. Triggered Palliative Care for Late-Stage Dementia: A Pilot Randomized Trial. *Journal of Pain & Symptom Management*. 2019 Jan;57(1):10-9. doi: <https://dx.doi.org/10.1016/j.jpainsymman.2018.10.494>. PMID: 30342242.
604. Verreault R, Arcand M, Misson L, et al. Quasi-experimental evaluation of a multifaceted intervention to improve quality of end-of-life care and quality of dying for patients with advanced dementia in long-term care institutions. *Palliative Medicine*. 2018 03;32(3):613-21. doi: <https://dx.doi.org/10.1177/0269216317719588>. PMID: 28731379.
605. Williams KN, Perkhounkova Y, Shaw CA, et al. Supporting Family Caregivers With Technology for Dementia Home Care: A Randomized Controlled Trial. *Innovation in Aging*. 2019 Jul;3(3):igz037. doi: <https://dx.doi.org/10.1093/geroni/igz037>. PMID: 31660443.
606. Horvath KJ, Trudeau SA, Rudolph JL, et al. Clinical trial of a home safety toolkit for Alzheimer's disease. *International journal of Alzheimer's disease*. 2013;2013:913606. doi: <https://dx.doi.org/10.1155/2013/913606>. PMID: 24195007.
607. Laver K, Liu E, Clemson L, et al. Does Telehealth Delivery of a Dyadic Dementia Care Program Provide a Noninferior Alternative to Face-To-Face Delivery of the Same Program? A Randomized, Controlled Trial. *American Journal of Geriatric Psychiatry*. 2020 Mar 02;02:02. doi: <https://dx.doi.org/10.1016/j.jagp.2020.02.009>. PMID: 32234275.
608. Villars H, Dupuy C, Soler P, et al. A follow-up intervention in severely demented patients after discharge from a special Alzheimer acute care unit: impact on early emergency room re-hospitalization rate. *International Journal of Geriatric Psychiatry*. 2013 Nov;28(11):1131-40. doi: <https://dx.doi.org/10.1002/gps.3932>. PMID: 23348897.
609. Mavandadi S, Wright EM, Graydon MM, et al. A randomized pilot trial of a telephone-based collaborative care management program for caregivers of individuals with dementia. *Psychological Services*. 2017 Feb;14(1):102-11. doi: <https://dx.doi.org/10.1037/ser0000118>. PMID: 28134558.
610. Chapman DG, Toseland RW. Effectiveness of advanced illness care teams for nursing home residents with dementia. *Social Work*. 2007 Oct;52(4):321-9. PMID: 18232242.
611. Samus QM, Johnston D, Black BS, et al. A multidimensional home-based care coordination intervention for elders with memory disorders: the maximizing independence at home (MIND) pilot randomized trial. *American Journal of Geriatric Psychiatry*. 2014 Apr;22(4):398-414. doi: <https://dx.doi.org/10.1016/j.jagp.2013.12.175>. PMID: 24502822.

612. Saxena N, George PP, Teo KW, et al. Evaluation of an integrated primary care-led dementia shared care program in Singapore: An effectiveness and cost-effectiveness study. *Geriatrics & gerontology international*. 2018 Mar;18(3):479-86. doi: <https://dx.doi.org/10.1111/ggi.13196>. PMID: 29193721.
613. Leontjevas R, Gerritsen DL, Smalbrugge M, et al. A structural multidisciplinary approach to depression management in nursing-home residents: a multicentre, stepped-wedge cluster-randomised trial. *Lancet*. 2013 Jun 29;381(9885):2255-64. doi: [https://dx.doi.org/10.1016/S0140-6736\(13\)60590-5](https://dx.doi.org/10.1016/S0140-6736(13)60590-5). PMID: 23643110.
614. Chen C, Huang Y, Liu C, et al. Effects of an Interdisciplinary Care Team on the Management of Alzheimer's Disease in China. *Journal of gerontological nursing*. 2019 01 May;45(5):39-45. doi: <http://dx.doi.org/10.3928/00989134-20190318-01>. PMID: 627579646.
615. Bass D, Judge K, Snow A, et al. Caregiver outcomes of partners in dementia care: effect of a care coordination program for veterans with dementia and their family members and friends. *Journal of the american geriatrics society*. 2013;61(8):1377-86. doi: 10.1111/jgs.12362. PMID: CN-00875084.
616. Callahan CM, Boustani MA, Unverzagt FW, et al. Effectiveness of collaborative care for older adults with Alzheimer disease in primary care: a randomized controlled trial. *JAMA*. 2006 May 10;295(18):2148-57. PMID: 16684985.
617. Chodosh J, Colaiaco BA, Connor KI, et al. Dementia Care Management in an Underserved Community: The Comparative Effectiveness of Two Different Approaches. *Journal of Aging & Health*. 2015 Aug;27(5):864-93. doi: <https://dx.doi.org/10.1177/0898264315569454>. PMID: 25656074.
618. Possin KL, Merrilees JJ, Dulaney S, et al. Effect of Collaborative Dementia Care via Telephone and Internet on Quality of Life, Caregiver Well-being, and Health Care Use: The Care Ecosystem Randomized Clinical Trial. *JAMA Internal Medicine*. 2019 Sep 30;30:30. doi: <https://dx.doi.org/10.1001/jamainternmed.2019.4101>. PMID: 31566651.
619. Thyrian JR, Hertel J, Wucherer D, et al. Effectiveness and Safety of Dementia Care Management in Primary Care: A Randomized Clinical Trial. *JAMA Psychiatry*. 2017 Oct 01;74(10):996-1004. doi: <https://dx.doi.org/10.1001/jamapsychiatry.2017.2124>. PMID: 28746708.
620. Vickrey BG, Mittman BS, Connor KI, et al. The effect of a disease management intervention on quality and outcomes of dementia care: a randomized, controlled trial.[Summary for patients in *Ann Intern Med*. 2006 Nov 21;145(10):I31; PMID: 17116913]. *Annals of Internal Medicine*. 2006 Nov 21;145(10):713-26. PMID: 17116916.
621. Eichler T, Thyrian JR, Dreier A, et al. Dementia care management: going new ways in ambulant dementia care within a GP-based randomized controlled intervention trial. *International Psychogeriatrics*. 2014 Feb;26(2):247-56. doi: <https://dx.doi.org/10.1017/S1041610213001786>. PMID: 24152974.
622. Eichler T, Thyrian JR, Fredrich D, et al. The benefits of implementing a computerized intervention-management-system (IMS) on delivering integrated dementia care in the primary care setting. *International Psychogeriatrics*. 2014 Aug;26(8):1377-85. doi: <https://dx.doi.org/10.1017/S1041610214000830>. PMID: 24811145.

623. Fis T, Thyrian JR, Wucherer D, et al. Medication management for people with dementia in primary care: description of implementation in the DelpHi study. *BMC Geriatrics*. 2013 Nov 13;13:121. doi: <https://dx.doi.org/10.1186/1471-2318-13-121>. PMID: 24225205.
624. Thyrian JR, Winter P, Eichler T, et al. Relatives' burden of caring for people screened positive for dementia in primary care : Results of the DelpHi study. *Zeitschrift fur Gerontologie und Geriatrie*. 2017 Jan;50(1):4-13. doi: <https://dx.doi.org/10.1007/s00391-016-1119-9>. PMID: 27534949.
625. Thyrian JR, Fis T, Dreier A, et al. Life- and person-centred help in Mecklenburg-Western Pomerania, Germany (DelpHi): study protocol for a randomised controlled trial. *Trials* [Electronic Resource]. 2012 May 10;13:56. doi: <https://dx.doi.org/10.1186/1745-6215-13-56>. PMID: 22575023.
626. Zwingmann I, Hoffmann W, Michalowsky B, et al. Supporting family dementia caregivers: testing the efficacy of dementia care management on multifaceted caregivers' burden. *Aging & Mental Health*. 2017 Nov 20:1-8. doi: <https://dx.doi.org/10.1080/13607863.2017.1399341>. PMID: 29156941.
627. Michalowsky B, Xie F, Eichler T, et al. Cost-effectiveness of a collaborative dementia care management-Results of a cluster-randomized controlled trial. *Alzheimer's & Dementia*. 2019 Oct;15(10):1296-308. doi: <https://dx.doi.org/10.1016/j.jalz.2019.05.008>. PMID: 31409541.
628. Bass DM, Judge KS, Maslow K, et al. Impact of the care coordination program "partners in Dementia Care" on veterans' hospital admissions and emergency department visits. *Alzheimer's and Dementia: Translational Research and Clinical Interventions*. 2015 14 Oct;1(1):13-22. doi: <http://dx.doi.org/10.1016/j.trci.2015.03.003>. PMID: 606362402.
629. Morgan RO, Bass DM, Judge KS, et al. A break-even analysis for dementia care collaboration: Partners in Dementia Care. *JGIM: Journal of General Internal Medicine*. 2015;30(6):804-9. doi: 10.1007/s11606-015-3205-x. PMID: 109738468. Language: English. Entry Date: 20150923. Revision Date: 20160604. Publication Type: journal article.
630. Duru OK, Ettner SL, Vassar SD, et al. Cost evaluation of a coordinated care management intervention for dementia. *American Journal of Managed Care*. 2009 Aug;15(8):521-8. PMID: 19670955.
631. Chodosh J, Pearson ML, Connor KI, et al. A dementia care management intervention: which components improve quality? *American Journal of Managed Care*. 2012 Feb;18(2):85-94. PMID: 22435836.
632. Brown AF, Vassar SD, Connor KI, et al. Collaborative care management reduces disparities in dementia care quality for caregivers with less education. *Journal of the American Geriatrics Society*. 2013 Feb;61(2):243-51. doi: <https://dx.doi.org/10.1111/jgs.12079>. PMID: 23320655.
633. Bass DM, Clark PA, Looman WJ, et al. The Cleveland Alzheimer's managed care demonstration: outcomes after 12 months of implementation. *Gerontologist*. 2003 Feb;43(1):73-85. PMID: 12604748.
634. Clark PA, Bass DM, Looman WJ, et al. Outcomes for patients with dementia from the Cleveland Alzheimer's Managed Care Demonstration. *Aging & Mental Health*. 2004 Jan;8(1):40-51. PMID: 14690867.
635. Bernstein A, Harrison KL, Dulaney S, et al. The Role of Care Navigators Working with People with Dementia and Their Caregivers. *Journal of Alzheimer's Disease*. 2019;71(1):45-55. doi: <https://dx.doi.org/10.3233/JAD-180957>. PMID: 31322558.

636. Possin KL, Merrilees J, Bonasera SJ, et al. Development of an adaptive, personalized, and scalable dementia care program: Early findings from the Care Ecosystem. *PLoS Medicine / Public Library of Science*. 2017 03;14(3):e1002260. doi: <https://dx.doi.org/10.1371/journal.pmed.1002260>. PMID: 28323819.
637. Rosa TD, Possin KL, Bernstein A, et al. Variations in Costs of a Collaborative Care Model for Dementia. *Journal of the American Geriatrics Society*. 2019 Dec;67(12):2628-33. doi: <https://dx.doi.org/10.1111/jgs.16076>. PMID: 31317539.
638. Guterman EL, Allen IE, Josephson SA, et al. Association Between Caregiver Depression and Emergency Department Use Among Patients With Dementia. *JAMA Neurology*. 2019 Jul 08;08:08. doi: <https://dx.doi.org/10.1001/jamaneurol.2019.1820>. PMID: 31282955.
639. Merrilees JJ, Bernstein A, Dulaney S, et al. The Care Ecosystem: Promoting self-efficacy among dementia family caregivers. *Dementia*. 2018 Nov 29;1471301218814121. doi: <https://dx.doi.org/10.1177/1471301218814121>. PMID: 30497302.
640. Eloniemi-Sulkava U, Saarenheimo M, Laakkonen ML, et al. Family care as collaboration: effectiveness of a multicomponent support program for elderly couples with dementia. Randomized controlled intervention study. *Journal of the American Geriatrics Society*. 2009 Dec;57(12):2200-8. doi: <https://dx.doi.org/10.1111/j.1532-5415.2009.02564.x>. PMID: 20121986.
641. Lichtwarck B, Selbaek G, Kirkevold O, et al. Targeted Interdisciplinary Model for Evaluation and Treatment of Neuropsychiatric Symptoms: A Cluster Randomized Controlled Trial. *American Journal of Geriatric Psychiatry*. 2018 January;26(1):25-38. doi: <http://dx.doi.org/10.1016/j.jagp.2017.05.015>. PMID: 617024611.
642. Li J, Grandner M, Chang Y, et al. Person-centered dementia care and sleep in assisted living residents with dementia: a pilot study. *Behavioral sleep medicine*. 2017(2):97-113. doi: 10.1080/15402002.2015.1104686. PMID: CN-01380345.
643. Rovner BW, Steele CD, Shmuelly Y, et al. A randomized trial of dementia care in nursing homes. *Journal of the American Geriatrics Society*. 1996 Jan;44(1):7-13. PMID: 8537594.
644. Surr CA, Griffiths AW, Kelley R, et al. The Implementation of Dementia Care Mapping in a Randomized Controlled Trial in Long-Term Care: Results of a Process Evaluation. *American Journal of Alzheimer's Disease & Other Dementias*. 2019 Sep;34(6):390-8. doi: <https://dx.doi.org/10.1177/1533317519845725>. PMID: 31056923.
645. Ballard C, Orrell M, YongZhong S, et al. Impact of Antipsychotic Review and Nonpharmacological Intervention on Antipsychotic Use, Neuropsychiatric Symptoms, and Mortality in People With Dementia Living in Nursing Homes: A Factorial Cluster-Randomized Controlled Trial by the Well-Being and Health for People With Dementia (WHELPD) Program. *American Journal of Psychiatry*. 2016 Mar 01;173(3):252-62. doi: <https://dx.doi.org/10.1176/appi.ajp.2015.15010130>. PMID: 26585409.
646. Fossey J, Ballard C, Juszcak E, et al. Effect of enhanced psychosocial care on antipsychotic use in nursing home residents with severe dementia: cluster randomised trial. *BMJ (Clinical research ed.)*. 2006 1 Apr;332(7544):756-61. PMID: 43509632.

647. Ballard C, Corbett A, Orrell M, et al. Impact of person-centred care training and person-centred activities on quality of life, agitation, and antipsychotic use in people with dementia living in nursing homes: A cluster-randomised controlled trial. *PLoS Medicine / Public Library of Science*. 2018 02;15(2):e1002500. doi: <https://dx.doi.org/10.1371/journal.pmed.1002500>. PMID: 29408901.
648. Moyle W, Venturato L, Cooke M, et al. Evaluating the capabilities model of dementia care: A non-randomized controlled trial exploring resident quality of life and care staff attitudes and experiences. *International Psychogeriatrics*. 2016 01 Jul;28(7):1091-100. doi: <http://dx.doi.org/10.1017/S1041610216000296>. PMID: 608901615.
649. Chenoweth L, Forbes I, Fleming R, et al. PerCEN: a cluster randomized controlled trial of person-centered residential care and environment for people with dementia. *International Psychogeriatrics*. 2014 Jul;26(7):1147-60. doi: <https://dx.doi.org/10.1017/S1041610214000398>. PMID: 24666667.
650. Chenoweth L, King MT, Jeon YH, et al. Caring for Aged Dementia Care Resident Study (CADRES) of person-centred care, dementia-care mapping, and usual care in dementia: a cluster-randomised trial.[Erratum appears in *Lancet Neurol*. 2009 May;8(5):419]. *Lancet Neurology*. 2009 Apr;8(4):317-25. doi: [https://dx.doi.org/10.1016/S1474-4422\(09\)70045-6](https://dx.doi.org/10.1016/S1474-4422(09)70045-6). PMID: 19282246.
651. Rokstad AM, Rosvik J, Kirkevold O, et al. The effect of person-centred dementia care to prevent agitation and other neuropsychiatric symptoms and enhance quality of life in nursing home patients: a 10-month randomized controlled trial. *Dementia & Geriatric Cognitive Disorders*. 2013;36(5-6):340-53. doi: <https://dx.doi.org/10.1159/000354366>. PMID: 24022375.
652. Dichter M, Quasdorf T, Schwab C, et al. Dementia care mapping: effects on residents' quality of life and challenging behavior in German nursing homes. A quasi-experimental trial. *International psychogeriatrics*. 2015;27(11):1875-92. doi: 10.1017/S1041610215000927. PMID: CN-01215190.
653. van de Ven G, Draskovic I, Adang EM, et al. Effects of dementia-care mapping on residents and staff of care homes: a pragmatic cluster-randomised controlled trial. *PLoS ONE [Electronic Resource]*. 2013;8(7):e67325. doi: <https://dx.doi.org/10.1371/journal.pone.0067325>. PMID: 23844003.
654. Fossey J, Ballard C, Juszcak E, et al. Effect of enhanced psychosocial care on antipsychotic use in nursing home residents with severe dementia: cluster randomised trial.[Erratum appears in *BMJ*. 2006 Apr 1;332(7544):61]. *BMJ*. 2006 Apr 01;332(7544):756-61. PMID: 16543297.
655. Froggatt K, Best A, Bunn F, et al. A group intervention to improve quality of life for people with advanced dementia living in care homes: The namaste feasibility cluster RCT. *Health Technology Assessment*. 2020 January;24(6):vii-139.
656. Lichtwarck B, Selbaek G, Kirkevold O, et al. TIME - Targeted interdisciplinary model for evaluation and treatment of neuropsychiatric symptoms: protocol for an effectiveness-implementation cluster randomized hybrid trial. *BMC Psychiatry*. 2016 Jul 12;16:233. doi: <https://dx.doi.org/10.1186/s12888-016-0944-0>. PMID: 27406242.
657. Lichtwarck B, Myhre J, Selbaek G, et al. TIME to reduce agitation in persons with dementia in nursing homes. A process evaluation of a complex intervention. *BMC Health Services Research*. 2019 May 31;19(1):349. doi: <https://dx.doi.org/10.1186/s12913-019-4168-0>. PMID: 31151437.

658. Fossey J, Garrod L, Tolbol Froiland C, et al. What influences the sustainability of an effective psychosocial intervention for people with dementia living in care homes? A 9 to 12-month follow-up of the perceptions of staff in care homes involved in the WHELD randomised controlled trial. *International Journal of Geriatric Psychiatry*. 2019 May;34(5):674-82. doi: <http://dx.doi.org/10.1002/gps.5066>.
659. Romeo R, Zala D, Knapp M, et al. Improving the quality of life of care home residents with dementia: Cost-effectiveness of an optimized intervention for residents with clinically significant agitation in dementia. *Alzheimer's & Dementia*. 2019 Feb;15(2):282-91. doi: <https://dx.doi.org/10.1016/j.jalz.2018.08.010>. PMID: 30470592.
660. Bourgeois MS, Dijkstra K, Burgio L, et al. Memory aids as an augmentative and alternative communication strategy for nursing home residents with dementia. *AAC: Augmentative and Alternative Communication*. 2001 Sep;17(3):196-210. doi: <http://dx.doi.org/10.1080/714043383>. PMID: 2001-11280-006.
661. Chang CC, Lin LC. Effects of a feeding skills training programme on nursing assistants and dementia patients. *Journal of Clinical Nursing*. 2005 Nov;14(10):1185-92. PMID: 16238764.
662. Clare L, Whitaker R, Woods RT, et al. AwareCare: a pilot randomized controlled trial of an awareness-based staff training intervention to improve quality of life for residents with severe dementia in long-term care settings. *International Psychogeriatrics*. 2013 Jan;25(1):128-39. doi: <https://dx.doi.org/10.1017/S1041610212001226>. PMID: 22840185.
663. Conway ER, Chenery HJ. Evaluating the MESSAGE Communication Strategies in Dementia training for use with community-based aged care staff working with people with dementia: a controlled pretest-post-test study. *Journal of Clinical Nursing*. 2016 Apr;25(7-8):1145-55. doi: <https://dx.doi.org/10.1111/jocn.13134>. PMID: 26821868.
664. Deudon A, Maubourguet N, Gervais X, et al. Non-pharmacological management of behavioural symptoms in nursing homes. *International Journal of Geriatric Psychiatry*. 2009 Dec;24(12):1386-95. doi: <https://dx.doi.org/10.1002/gps.2275>. PMID: 19370714.
665. Galik E, Resnick B, Hammersla M, et al. Optimizing function and physical activity among nursing home residents with dementia: testing the impact of function-focused care. *Gerontologist*. 2014 Dec;54(6):930-43. doi: <https://dx.doi.org/10.1093/geront/gnt108>. PMID: 24092822.
666. Ghandehari OO, Hadjistavropoulos T, Williams J, et al. A controlled investigation of continuing pain education for long-term care staff. *Pain Research & Management*. 2013;18(1):11-8. PMID: 107988179. Language: English. Entry Date: 20130426. Revision Date: 20161117. Publication Type: journal article.
667. Gozalo P, Prakash S, Qato DM, et al. Effect of the bathing without a battle training intervention on bathing-associated physical and verbal outcomes in nursing home residents with dementia: a randomized crossover diffusion study. *Journal of the American Geriatrics Society*. 2014 May;62(5):797-804. doi: <https://dx.doi.org/10.1111/jgs.12777>. PMID: 24697702.

668. Hattink B, Meiland F, van der Roest H, et al. Web-Based STAR E-Learning Course Increases Empathy and Understanding in Dementia Caregivers: Results from a Randomized Controlled Trial in the Netherlands and the United Kingdom. *Journal of Medical Internet Research*. 2015 Oct 30;17(10):e241. doi: <https://dx.doi.org/10.2196/jmir.4025>. PMID: 26519106.
669. Kuske B, Luck T, Hanns S, et al. Training in dementia care: a cluster-randomized controlled trial of a training program for nursing home staff in Germany. *International Psychogeriatrics*. 2009 Apr;21(2):295-308. doi: <https://dx.doi.org/10.1017/S1041610208008387>. PMID: 19193252.
670. Leone E, Deudon A, Bauchet M, et al. Management of apathy in nursing homes using a teaching program for care staff: the STIM-EHPAD study. *International Journal of Geriatric Psychiatry*. 2013 Apr;28(4):383-92. doi: <https://dx.doi.org/10.1002/gps.3836>. PMID: 22700526.
671. Magai C, Cohen CI, Gomberg D. Impact of training dementia caregivers in sensitivity to nonverbal emotion signals. *International Psychogeriatrics*. 2002 Mar;14(1):25-38. PMID: 12094904.
672. McCallion P, Toseland RW, Lacey D, et al. Educating nursing assistants to communicate more effectively with nursing home residents with dementia. *Gerontologist*. 1999 Oct;39(5):546-58. PMID: 10568079.
673. McCurry SM, LaFazia DM, Pike KC, et al. Development and evaluation of a sleep education program for older adults with dementia living in adult family homes. *American Journal of Geriatric Psychiatry*. 2012 Jun;20(6):494-504. doi: <https://dx.doi.org/10.1097/JGP.0b013e318248ae79>. PMID: 22367233.
674. Proctor R, Burns A, Powell HS, et al. Behavioural management in nursing and residential homes: a randomised controlled trial. *Lancet*. 1999 Jul 03;354(9172):26-9. PMID: 10406361.
675. Schindel ML, Gillies L, Coker E, et al. An Education Intervention to Enhance Staff Self-Efficacy to Provide Dementia Care in an Acute Care Hospital in Canada: a Nonrandomized Controlled Study. In *American journal of alzheimer's disease and other dementias*
676. Sloane PD, Hoeffler B, Mitchell CM, et al. Effect of person-centered showering and the towel bath on bathing-associated aggression, agitation, and discomfort in nursing home residents with dementia: a randomized, controlled trial. *Journal of the American Geriatrics Society*. 2004 Nov;52(11):1795-804. PMID: 15507054.
677. Wells DL, Dawson P, Sidani S, et al. Effects of an abilities-focused program of morning care on residents who have dementia and on caregivers. *Journal of the American Geriatrics Society*. 2000 Apr;48(4):442-9. PMID: 10798473.
678. Wenborn J, Challis D, Head J, et al. Providing activity for people with dementia in care homes: a cluster randomised controlled trial. *International Journal of Geriatric Psychiatry*. 2013 Dec;28(12):1296-304. doi: <https://dx.doi.org/10.1002/gps.3960>. PMID: 23637069.
679. Williams KN, Perkhounkova Y, Herman R, et al. A Communication Intervention to Reduce Resistiveness in Dementia Care: A Cluster Randomized Controlled Trial. *Gerontologist*. 2017 Aug 01;57(4):707-18. doi: <https://dx.doi.org/10.1093/geront/gnw047>. PMID: 27048705.
680. Zimmerman S, Mitchell CM, Reed D, et al. Outcomes of a dementia care training program for staff in nursing homes and residential care/assisted living settings. *Alzheimer's Care Today*. 2010;11(2):83-99. PMID: 105194596. Language: English. Entry Date: 20100625. Revision Date: 20150711. Publication Type: Journal Article.
681. Burgener SC, Bakas T, Murray C, et al. Effective caregiving approaches for patients with Alzheimer's disease. *Geriatric Nursing*. 1998 May-Jun;19(3):121-6. PMID: 9708136.

682. Done DJ, Thomas JA. Training in communication skills for informal carers of people suffering from dementia: a cluster randomized clinical trial comparing a therapist led workshop and a booklet. *International Journal of Geriatric Psychiatry*. 2001 Aug;16(8):816-21. PMID: 11536349.
683. Gormley N, Lyons D, Howard R. Behavioural management of aggression in dementia: a randomized controlled trial. *Age & Ageing*. 2001 Mar;30(2):141-5. PMID: 11395344.
684. Hepburn K, Lewis M, Tornatore J, et al. The Savvy Caregiver program: the demonstrated effectiveness of a transportable dementia caregiver psychoeducation program. *Journal of Gerontological Nursing*. 2007 Mar;33(3):30-6. PMID: 17378189.
685. Hepburn KW, Lewis M, Narayan S, et al. Partners in caregiving: A psychoeducation program affecting dementia family caregivers' distress and caregiving outlook. *Clinical Gerontologist*. 2005;29(1):53-69. doi: http://dx.doi.org/10.1300/J018v29n01_05. PMID: 43539927.
686. Hepburn KW, Tornatore J, Center B, et al. Dementia family caregiver training: affecting beliefs about caregiving and caregiver outcomes. *Journal of the American Geriatrics Society*. 2001 Apr;49(4):450-7. PMID: 11347790.
687. Klodnicka Kouri K, Ducharme FC, Giroux F. A psycho-educational intervention focused on communication for caregivers of a family member in the early stage of Alzheimer's disease: Results of an experimental study. *Dementia: The International Journal of Social Research and Practice*. 2011 Aug;10(3):435-53. doi: <http://dx.doi.org/10.1177/1471301211408124>. PMID: 2011-16142-011.
688. Liddle J, Smith-Conway ER, Baker R, et al. Memory and communication support strategies in dementia: effect of a training program for informal caregivers. *International Psychogeriatrics*. 2012 Dec;24(12):1927-42. doi: <https://dx.doi.org/10.1017/S1041610212001366>. PMID: 23092595.
689. Martin-Cook K, Davis BA, Hynan LS, et al. A randomized, controlled study of an Alzheimer's caregiver skills training program. *American Journal of Alzheimer's Disease & Other Dementias*. 2005 Jul-Aug;20(4):204-10. PMID: 16136843.
690. Neely A, Vikstrom S, Josephsson S. Collaborative memory intervention in dementia: caregiver participation matters. In *Neuropsychological rehabilitation*
691. Robinson K, Yates K. Effects of two caregiver-training programs on burden and attitude toward help. *Archives of Psychiatric Nursing*. 1994 Oct;8(5):312-9. PMID: 7993133.
692. Jablonski RA, Reed D, Maas ML. Care intervention for older adults with Alzheimer's disease and related dementias: effect of family involvement on cognitive and functional outcomes in nursing homes. *Journal of Gerontological Nursing*. 2005 Jun;31(6):38-48. PMID: 16138529.
693. Maas ML, Reed D, Park M, et al. Outcomes of family involvement in care intervention for caregivers of individuals with dementia. *Nursing Research*. 2004 Mar-Apr;53(2):76-86. PMID: 15084992.
694. McCallion P, Toseland RW, Freeman K. An evaluation of a family visit education program. *Journal of the American Geriatrics Society*. 1999 Feb;47(2):203-14. PMID: 9988292.
695. Robison J, Curry L, Gruman C, et al. Partners in caregiving in a special care environment: cooperative communication between staff and families on dementia units. *Gerontologist*. 2007 Aug;47(4):504-15. PMID: 17766671.

696. Kinnunen KM, Rapaport P, Webster L, et al. A manual-based intervention for carers of people with dementia and sleep disturbances: An acceptability and feasibility RCT. *Health Technology Assessment*. 2018 December;22(71):I-408. doi: <http://dx.doi.org/10.3310/hta22710>. PMID: 626134933.
697. Kunik ME, Snow AL, Wilson N, et al. Teaching Caregivers of Persons with Dementia to Address Pain. *American Journal of Geriatric Psychiatry*. 2017 Feb;25(2):144-54. doi: <https://dx.doi.org/10.1016/j.jagp.2016.04.009>. PMID: 27743840.
698. Suominen MH, Puranen TM, Jyvakorpi SK, et al. Nutritional Guidance Improves Nutrient Intake and Quality of Life, and May Prevent Falls in Aged Persons with Alzheimer Disease Living with a Spouse (NuAD Trial). *Journal of Nutrition, Health & Aging*. 2015 Nov;19(9):901-7. doi: <https://dx.doi.org/10.1007/s12603-015-0558-0>. PMID: 26482691.
699. Teri L, McCurry SM, Logsdon R, et al. Training community consultants to help family members improve dementia care: a randomized controlled trial. *Gerontologist*. 2005 Dec;45(6):802-11. PMID: 16326662.
700. Livingston G, Barber JA, Kinnunen KM, et al. DREAMS-START (Dementia RElAted Manual for Sleep; STRAtegies for RelaTives) for people with dementia and sleep disturbances: a single-blind feasibility and acceptability randomized controlled trial. *International Psychogeriatrics*. 2019 02;31(2):251-65. doi: <https://dx.doi.org/10.1017/S1041610218000753>. PMID: 30221615.
701. McCurry SM, Gibbons LE, Logsdon RG, et al. Training caregivers to change the sleep hygiene practices of patients with dementia: the NITE-AD project. *Journal of the American Geriatrics Society*. 2003 Oct;51(10):1455-60. PMID: 14511168.
702. McCurry SM, Gibbons LE, Logsdon RG, et al. Nighttime insomnia treatment and education for Alzheimer's disease: a randomized, controlled trial. *Journal of the American Geriatrics Society*. 2005 May;53(5):793-802. PMID: 15877554.
703. Morton S, Murad M, O'Connor E, et al. Quantitative Synthesis—An Update. *Methods Guide for Comparative Effectiveness Reviews*. (Prepared by the Scientific Resource Center under Contract No. 290-2012-0004-C). AHRQ Publication No. 18-EHC007-EF. Rockville, MD: Agency for Healthcare Research and Quality; 2018.
704. Mills EJ, Ayers D, Chou R, et al. Are current standards of reporting quality for clinical trials sufficient in addressing important sources of bias? *Contemporary Clinical Trials*. 2015;45:2-7. doi: 10.1016/j.cct.2015.07.019. PMID: 26232560.
705. Callahan CM, Sachs GA, LaMantia MA, et al. Redesigning systems of care for older adults with Alzheimer's disease. 2014;33(4):626-32.
706. Schultz SK, Llorente MD, Sanders AE, et al. Quality improvement in dementia care: Dementia Management Quality Measurement Set 2018 Implementation Update. 2020;94(5):210-6.
707. Odenheimer G, Borson S, Sanders AE, et al. Quality improvement in neurology: dementia management quality measures. 2013;81(17):1545-9.
708. Czaja SI, Schulz R, Lee CC, et al. A methodology for describing and decomposing complex psychosocial and behavioral interventions. *Psychology & Aging*. 2003;18:385-95. doi: 10.1037/0882-7974.18.3.385 PMID: 14518802.
709. Jackson JW, Arah OA. Making causal inference more social and (social) epidemiology more causal. *American Journal of Epidemiology*. 2019 October 1, 2019;kwz199. doi: 10.1093/aje/kwz199.

710. Collins LM. Optimization of behavioral, biobehavioral, and biomedical interventions: the multiphase optimization strategy (MOST). New York: Springer; 2018.
711. Gitlin LNB, R.; Jutkowitz, E.; Gustavson, A. M.; Sefcik, J. S.; Hodgson, N. A.; Keouth, S.,; Gaugler, J. E. Critical factors for successful dissemination and implementation of interventions designed to improve care for people living with dementia and care partners in health care systems. *Journal of the American Geriatrics Society*. under review.
712. Gitlin LNH, N.; Choi, S.; Marx, K. Interventions to address functional decline in persons with dementia: Closing the gap between what a person “does do” and what they “can do.” In: Jefferson PZB, ed *Neuropsychology of Alzheimer’s Disease and Other Dementias*. New York, NY: Oxford University Press; in press.
713. Office of Assistant Secretary of Health and Human Services. Including People with Dementia and Their Caregivers as Co-Researchers in Studies of Dementia Care and Services. 2018. <https://aspe.hhs.gov/basic-report/including-people-with-dementia-and-their-caregivers-as-co-Researchers-in-studies-of-dementia-care-and-services>. Accessed on March 1, 2020.
714. Silberzahn R, Uhlmann EL, Martin DP, et al. Many analysts, one data set: making transparent how variations in analytic choices affect results. *Advances in Methods and Practices in Psychological Science*. 2018;1(3):337-56. doi: 10.1177/2515245917747646.

Appendixes

Appendix A. Methods

- I. Example Interventions
- II. Common Outcome Measures
- III. Search Strategy
- IV. Risk of Bias – Traditional Studies
- V. Risk of Bias – Cluster Trials
- VI. PRECIS-2 Tool for Care Interventions

Appendix B. Excluded Studies

Appendix C. Chapter 4 Treatment for BPSD

- I. Assisted Therapy
- II. Multi-Sensory Stimulation/Snoezelen
- III. Complementary and Alternative Medicine (CAM) Therapy
- IV. Bright Light Therapy
- V. Psychosocial for BPSD
- VI. Multicomponent Interventions

Appendix D. Chapter 5 Interventions for PWD wellbeing

- I. Exercise
- II. Music Intervention
- III. Reminiscence Therapy
- IV. Cognitive Rehabilitation
- V. Cognitive Training
- VI. Cognitive Stimulation Therapy
- VII. Recreation Therapy
- VIII. Psychosocial for PWD Well-Being
- IX. Creative-Expression Therapy
- X. Multicomponent for PWD Well-Being
- XI. Assistive Technology
- XII. Electrostimulation
- XIII. Other Interventions for PWD Well-Being

Appendix E. Chapter 6 and 7 Interventions for Caregiver Well-Being

- I. Psychosocial Interventions to Support Caregiver's Well-Being
- II. Social Support
- III. Lifestyle Interventions
- IV. Respite Care
- V. Multicomponent for Informal Caregiver Well-Being
- VI. Other Interventions for Caregiver Well-Being
- VII. Interventions for Formal Caregiver Well-Being

Appendix F. Chapter 8 Care delivery interventions

- I. Care Service Provision
 - a. Consultation
 - b. Case Management
 - c. Care Protocols for PWD
 - d. Advance Care Planning
 - e. Palliative Care
 - f. Other Service Provision Interventions

- II. Care Delivery Models and Programs - Collaborative Care
- III. Care Staff Education and Support Needs
 - a. Caregiver Staff Training
 - b. Informal Caregiver Staff Training
 - c. Family Education and Partnering
 - d. Multitier Training

Appendix A. Methods

I. Example Interventions

Essentially, interventions are automatically included unless *specifically* stated as excluded. Note that the list is *not* divided by KQs 1-10. Some interventions may be aimed at both PWD and PWD Caregivers; some may be aimed at one or the other. The list is not intended to be exhaustive, and is a simple categorization based on what may be a more likely classification. The actual distinction between whether an intervention is examining **what** care is delivered or **how** to deliver care would be determined by the study purpose.

- Memory evaluation
- Driving evaluation or encouraging driving cessation
- Meaningful activities
- Advance care planning
- Behavior management
- ADL support
- Home modifications
- Wandering and fall risk management
- Palliative care
- Caregiver support and support groups
- Sensory-based interventions
- Changing the physical environment/environmental modification across settings (e.g., in hospitals, in people's homes)
- Mindfulness training
- Interventions focused on the development of Dementia Friendly Training (e.g., training of police officers in local communities)
- Wandering and Wayfinding
- Reminiscence Therapy
- Prompts and Multicomponent Interventions
- Engagement Interventions
- Exercise Interventions
- Psychoeducational
- Art therapy
- Dance movement therapy
- Music therapy
- Cognitive behavior therapy
- Counseling/care management (including emotionally focused couples therapy)
- General support
- Respite
- Training of PWD
- Psychosocial interventions/studies
- Caregiver support groups
- Therapeutic counseling

- Support interventions, including involving informal caregiver social network to support the primary caregiver
- Cognitive reframing (changing caregivers' maladaptive behaviors or beliefs)
- Web-based multimedia intervention
- Caregiver-therapist e-mail support
- Educational and peer-support website
- Bereavement support
- Improving acute care systems
- Skill training, including for CNAs, home health aides, and/or informal caregivers
- Training for CNAs, home health aides, and/or informal caregivers
- Improving care transitions
- Care coordination
- Multicomponent interventions

II. Common Outcome Measures

Table A-1. Common outcome measures

Test Name	Domain	Data Source	Reference
BEHAVE-AD	General behavior scales & global BPSD		Reisberg et al. 1987
Neuropsychiatric Inventory (NPI)	General behavior scales & global BPSD	informant	Cummings et al 1994
Cohen-Mansfield Agitation Inventory (CMAI)	Agitation/ aggression	informant	Cohen-Mansfield, 1986
Cornell Scale	Depression	patient or informant	Alexopoulos et al. 1988
Patient Health Questionnaire (PHQ-9)	Depression	patient	Spitzer et al., 1999
Geriatric Depression Scale (GDS) 30-item	Depression	patient	Yesavage et al. 1983
Geriatric Depression Scale (GDS) 15-item	Depression	patient	
Montgomery Asberg Depression Rating Scale (MADRS)	Depression		Montgomery & Asberg, 1979
Hamilton Depression Rating Scale (HDRS)	Depression	patient	Hamilton, 1960
Beck Anxiety Inventory (BAI)	Anxiety	patient	
Brief Psychiatric Rating Scale (BPRS)	Mood; Psychosis	clinician administered interview	Overall 1962; Beller 1984
Schedule for Affective Disorders and Schizophrenia (SADS)	Mood; Psychosis	clinician administered interview	Endicott 1978
Schedule for Affective Disorders and Schizophrenia -Lifetime version (SADS-L)	Mood; Psychosis	clinician administered interview	Endicott 1978
Schedule for Affective Disorders and Schizophrenia -Change version (SADS-C)	Mood; Psychosis	clinician administered interview	Endicott 1978
Behavioral Syndromes Scale for Dementia (BSSD)	General behavior scales & global BPSD	informant	Devanand 1992
Barthel index	ADLs	informant	Mahoney and Barthel, 1965
Bristol Activities of Daily Living Scale (BADLS)			Bucks et al. 1996
Direct Assessment of Functional Status	ADLs + IADLs	performance-based	Loewenstein, Amigo, & Duara, 1989
Disability Assessment for Dementia (DAD) Scale		informant	
Functional Activities Questionnaire (FAQ)		informant	Pfeffer et al 1982
Functional Independence Measure (FIM)	ADLs + (social, cogn, etc)	informant	Keith et al. 1987
Health Assessment Questionnaire (HAQ)			
Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE)		informant	Jorm and Jacomb, 1989
Instrumental Activities of Daily Living scale	IADLs		Lawton and Brody, 1969
Katz Index of Independence in ADLs	ADLs		Katz et al. 1963
Modified Health Assessment Questionnaire (MHAQ)			
Older Americans Resources and Services (OARS)	ADLs + IADLs	self-report	George & Fillenbaum, 1985

Test Name	Domain	Data Source	Reference
Physical Self-Maintenance Scale (PSMS)	ADLs		Lawton and Brody, 1969
Minimum Data Set (MDS)-ADL Self Performance Scale	ADLs		
Progressive Deterioration Scale (PDS)	ADLs + IADLs	informant	DeJong 1989
AD-related Quality of Life scale (QoL-AD)		patient or informant	Logsdon et al. 1999
DEMQOL		patient	Smith et al. 2007
DEMQOL		informant	Smith et al. 2007
EuroQol measure		patient or informant	EuroQol Group, 1990
Short Form-36 (SF-36)		patient	Ware & Sherbourne, 1992
General Health Questionnaire (GHQ)	Global Distress		Goldberg & Williams 1988
Zarit Burden Interview	Caregiver Burden		Zarit et al. 1980
Neuropsychiatric Inventory – Distress Scale	Caregiver Distress		Cummings et al 1994
Revised Memory and Behavior Problem Checklist (RMBPC)		informant	Terie et al 1992

III. Search Strategy

Database: Ovid MEDLINE(R) Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily, Ovid MEDLINE and Versions(R) <1946

Search Strategy:

- 1 exp Alzheimer Disease/
- 2 Dementia/
- 3 (dementia or alzheimer*).ti.
- 4 1 or 2 or 3
- 5 limit 4 to "therapy (best balance of sensitivity and specificity)"
- 6 limit 5 to english language
- 7 limit 6 to (addresses or autobiography or bibliography or biography or case reports or clinical conference or comment or comparative study or congresses or consensus development conference or consensus development conference, nih or dataset or dictionary or directory or editorial or evaluation studies or "expression of concern" or festschrift or government publications or guideline or historical article or interactive tutorial or interview or lectures or legal cases or legislation or letter or news or newspaper article or observational study or patient education handout or periodical index or personal narratives or portraits or "review" or "scientific integrity review" or validation studies or video-audio media or webcasts)
- 8 limit 7 to (adaptive clinical trial or clinical study or clinical trial, all or clinical trial, phase i or clinical trial, phase ii or clinical trial, phase iii or clinical trial, phase iv or clinical trial or controlled clinical trial or randomized controlled trial)
- 9 6 not 7
- 10 8 or 9
- 11 limit 10 to ("all child (0 to 18 years)")
- 12 limit 11 to ("middle age (45 to 64 years)" or "middle aged (45 plus years)" or "all aged (65 and over)" or "aged (80 and over)")
- 13 10 not 11
- 14 12 or 13

Database: Embase Classic+Embase <1947

Search Strategy:

- 1 exp *Alzheimer disease/
- 2 *dementia/
- 3 (alzheimer* or dementia*).ti.)
- 4 1 or 2 or 3
- 5 limit 4 to english language
- 6 limit 5 to "therapy (best balance of sensitivity and specificity)"
- 7 limit 6 to "reviews (best balance of sensitivity and specificity)"
- 8 6 not 7
- 9 limit 8 to (embryo <first trimester> or infant <to one year> or child <unspecified age> or preschool child <1 to 6 years> or school child <7 to 12 years> or adolescent <13 to 17 years>)
- 10 limit 9 to (adult <18 to 64 years> or aged <65+ years>)
- 11 8 not 9

- 12 10 or 11
- 13 limit 12 to (book or book series or conference proceeding or trade journal)
- 14 12 not 13
- 15 limit 14 to conference abstracts
- 16 14 not 15
- 17 limit 16 to (abstract report or books or "book review" or chapter or conference abstract or "conference review" or editorial or letter or note or patent or reports or "review" or short survey or tombstone)
- 18 16 not 17
- 19 limit 18 to (amphibia or ape or bird or cat or cattle or chicken or dog or "ducks and geese" or fish or "frogs and toads" or goat or guinea pig or "hamsters and gerbils" or horse or monkey or mouse or "pigeons and doves" or "rabbits and hares" or rat or reptile or sheep or swine)
- 20 18 not 19

Database: PsycINFO <1806

Search Strategy:

-
- 1 exp *ALZHEIMER'S DISEASE/
 - 2 *dementia/
 - 3 (dementia* or alzheimer*).ti.
 - 4 1 or 2 or 3 (64340)
 - 5 limit 4 to "therapy (best balance of sensitivity and specificity)"
 - 6 limit 5 to (childhood <birth to 12 years> or adolescence <13 to 17 years>)
 - 7 limit 6 to adulthood <18+ years>
 - 8 5 not 6
 - 9 7 or 8
 - 10 limit 9 to animal
 - 11 9 not 10
 - 12 limit 11 to (100 childhood <birth to age 12 yrs> or 120 neonatal <birth to age 1 mo> or 140 infancy <2 to 23 mo> or 160 preschool age <age 2 to 5 yrs> or 180 school age <age 6 to 12 yrs> or 200 adolescence <age 13 to 17 yrs> or 320 young adulthood <age 18 to 29 yrs> or 340 thirties <age 30 to 39 yrs>)
 - 13 limit 12 to (360 middle age <age 40 to 64 yrs> or "380 aged <age 65 yrs and older>" or "390 very old <age 85 yrs and older>")
 - 14 11 not 12
 - 15 13 or 14
 - 16 limit 15 to (abstract collection or bibliography or chapter or clarification or "column/opinion" or "comment/reply" or dissertation or editorial or encyclopedia entry or interview or letter or obituary or poetry or publication information or review-book or review-media or review-software & other or reviews)
 - 17 15 not 16
 - 18 limit 17 to ("0200 book" or "0240 authored book" or "0280 edited book" or "0300 encyclopedia" or "0400 dissertation abstract")
 - 19 17 not 18
 - 20 limit 19 to english language
 - 21 limit 20 to "therapy (maximizes specificity)"

IV. Risk of Bias Assessment Guide – Traditional Studies


INSTRUCTIONS: Review the methods of each trial and assess each risk of bias component as described in these instructions. You may need to have separate assessments for different outcomes (i.e. different measures; different time points may have different attrition rates).

1) ATTRITION Bias

Table A-2. Attrition bias

Description/Guiding Questions	Notes
<ul style="list-style-type: none"> ▪ Systematic differences in the loss of participants from the study and how they were accounted for in the results (e.g., incomplete follow-up, differential attrition). Those who drop out of the study or who are lost to follow-up may be systematically different from those who remain in the study. Attrition bias can potentially change the collective (group) characteristics of the relevant groups and their observed outcomes in ways that affect study results by confounding and spurious associations. <p>Reasons for incomplete/missing data adequately explained? Do the author’s attempt to address attrition in the analysis?</p>	<ul style="list-style-type: none"> ▪ Attrition assessment is dependent on overall study duration (see flowchart) Report attrition rate in spreadsheet. If a study reports outcomes at multiple intervals (e.g., 6 months, 12 months, 18 months) assess attrition at the first relevant time point and the last time-point separately, you do not need to do every time point. Analysis should be done with appropriate method (i.e. sensitivity analysis with various scenarios); last value forward would only be appropriate for interventions that are supposed to improve the outcomes (i.e. memory training that intends to improve memory).

Figure A-1. Attrition bias assessment guidance




2) SELECTION BIAS

Table A-3. Selection bias

Description/Guiding Questions	Notes
<ul style="list-style-type: none"> Systematic differences between baseline characteristics of the groups that arise from self-selection of treatments, physician-directed selection of treatments, or association of treatment assignments with demographic, clinical, or social characteristics. <p>Did method of randomization create biased allocation to interventions (inadequate randomization)?</p>	<ul style="list-style-type: none"> “Good” Randomization: Detailed methodology would include providing method of randomization such as use of a random numbers table, or computer random number generator. Limited methodology would be the study saying simply saying they randomized in the methods or provided limited detail such as randomizing by a 2:1 ratio. “Poor”/No Randomization: Randomized based on week of the month of birthday or a non-randomized clinical trial, observational study.

Figure A-2. Selection bias assessment guidance



3) ATTRITION AND SELECTION BIAS OVERALL

Assess joint selection and attrition bias. If either selection or attrition bias is high, the risk of bias is HIGH.

Table A-4. Attrition and selection bias overall

Attrition Bias	Low	Low	Medium	Low	Medium	Medium	High
Selection Bias	Low	Medium	Low	High	Medium	High	
Action	Assess other biases	Assess other biases	Assess other biases	STOP*	Assess other biases	STOP*	STOP*

*Send to evidence map

*****PRIOR TO ASSESSING OTHER BIASES, CHECK PUBLICATION TO MAKE SURE OUTCOMES ARE ABSTRACTABLE. IF OUTCOMES ARE NOT ABSTRACTABLE (e.g., DATA IS PRESENTED AS GRAPHS ONLY), STOP ASSESSMENT AND CHECK WITH TEAM TO CONFIRM THAT PUBLICATION BELONGS IN EVIDENCE MAP*****


4) OTHER BIASES

A. DETECTION BIAS

Table A-5. Detection bias

Description/Guiding Questions	Notes
<ul style="list-style-type: none">▪ <i>Systematic differences in outcomes assessment among groups being compared, including systematic misclassification of the exposure or intervention, covariates, or outcomes because of variable definitions and timings, diagnostic thresholds, recall from memory, inadequate assessor blinding, and faulty measurement techniques. Erroneous statistical analysis might also affect the validity of effect estimates.</i> <p>Were the outcome assessors blinded to the intervention (“outcome assessor blinded”)?</p> <p>Was the timing of the outcome assessment similar in all groups (“comparable timing outcomes assessment”)?</p> <p>Was the scale used to measure outcomes validated, reliable?</p> <p>Were outcomes measured in clinically meaningful ways?</p>	<ul style="list-style-type: none">▪ X

Figure A-3. Detection bias assessment guidance




B. PERFORMANCE BIAS

Table A-6. Performance bias

Description/Guiding Questions	Notes
Systematic differences in the care provided to participants and protocol deviation. Examples include contamination of the control group with the exposure or intervention, problems with fidelity to the intervention, unbalanced provision of additional interventions or co-interventions, difference in co-interventions, and inadequate blinding of providers and participants.	<ul style="list-style-type: none"><li data-bbox="829 373 1424 867">▪ Intention-to-Treat (ITT): Includes every subject according to randomized treatment assignment. Ignores noncompliance, protocol deviations, withdrawal, and anything that happens after randomization. Concurrent Intervention: Study participants are receiving another intervention (i.e., treatment) that is not part of the intervention being tested. Example: Participants are randomized to a physical activity intervention (or no intervention), but are also dieting.

Figure A-4. Performance Bias Assessment Guidance



C. REPORTING BIAS

Table A-7. Reporting bias

Description/Guiding Questions	Notes
<ul style="list-style-type: none"> ▪ Systematic differences between reported and unreported findings (e.g., differential reporting of outcomes or harms, incomplete reporting of study findings, potential for bias in reporting through source of funding). <p>Was a select group of outcomes reported?</p>	<ul style="list-style-type: none"> ▪ Compare results to methods section and/ or protocol. Check if some results are reported in a different publication.

REPORTING BIAS ASSESSMENT GUIDANCE

Table A-8. Reporting bias assessment guidance

Domain	Options	Rating
All outcomes reported	Yes	Low
	No	Medium
	Not Reported	Medium

*****CHECK OVERALL ROB RATING BEFORE ASSESSING FIDELITY TO INTERVENTION. IF THE COMBINATION OF PREVIOUS DOMAINS INDICATES HIGH RISK OF BIAS, YOU DO NOT NEED TO ASSESS FIDELITY*****

D. FIDELITY TO INTERVENTION

Table A-9. Fidelity to intervention


Description/Guiding Questions	Notes
<ul style="list-style-type: none"> ▪ We anticipate that care delivery studies will generally fall in the range of NIH Stage 3 to 4, with the possibility that one or a few may be carried out as quality improvement and thus Stage 5. Since the Stage Model is explicitly designed to balance, or trade off, internal and external validity, we will approach risk of bias assessment as a threshold requirement rather than a continuum. <p>Look for reporting on intervention compliance, any data reported on consistency of intervention use, or any mechanisms used to ensure compliance (e.g., reminders, guides, manuals).</p>	<ul style="list-style-type: none"> ▪ Information may appear in methods, results, or discussion sections.

FIDELITY TO INTERVENTION ASSESSMENT GUIDANCE

Table A-10. Fidelity to intervention assessment guidance

Domain	Options	Rating
Fidelity to intervention	Yes (at least 70%)	Low
	Yes-adaptation planned/ replicable	Medium
	No-adaptation not planned	High
	Unclear/Not Reported	Medium

Figure A-5. Overall risk of bias assessment guidance




V. Risk of Bias Assessment Guide – Cluster Trials

1) ATTRITION Bias

Table A-11. Attrition bias

Description/Guiding Questions	Notes
<ul style="list-style-type: none"> ▪ Systematic differences in the loss of participants from the study and how they were accounted for in the results (e.g., incomplete follow-up, differential attrition). Those who drop out of the study or who are lost to follow-up may be systematically different from those who remain in the study. Attrition bias can potentially change the collective (group) characteristics of the relevant groups and their observed outcomes in ways that affect study results by confounding and spurious associations. <p>Reasons for incomplete/missing data adequately explained? Do the author’s attempt to address attrition in the analysis?</p>	<ul style="list-style-type: none"> ▪ Attrition assessment is dependent on overall study duration (see flowchart) Report attrition rate in spreadsheet. If a study reports outcomes at multiple intervals (e.g., 6 months, 12 months, 18 months) assess attrition at the first relevant time point and the last time-point separately, you do not need to do every time point. Analysis should be done with appropriate method (i.e. sensitivity analysis with various scenarios); last value forward would only be appropriate for interventions that are supposed to improve the outcomes (i.e. memory training that intends to improve memory).

Figure A-6. Attrition bias assessment guidance




3) SELECTION BIAS

Table A-12. Selection bias

Description/Guiding Questions	Notes
<ul style="list-style-type: none"> ▪ Systematic differences between baseline characteristics of the groups that arise from self-selection of treatments, physician-directed selection of treatments, or association of treatment assignments with demographic, clinical, or social characteristics. <p>Did method of randomization create biased allocation to interventions (inadequate randomization)?</p>	<ul style="list-style-type: none"> ▪ “Good” Randomization: Random numbers table, computer random number generator “Poor” Randomization: Randomized based on week of the month of birthday No Randomization: Non-randomized clinical trial, observational study

Figure A-7. Selection bias assessment guidance



3) ATTRITION AND SELECTION BIAS OVERALL

Assess joint selection and attrition bias. If either selection or attrition bias is high, the risk of bias is HIGH.

Table A-13. Attrition and selection bias overall

Attrition Bias	Low	Low	Medium	Low	Medium	Medium	High
Selection Bias	Low	Medium	Low	High	Medium	High	
Action	Assess other biases	Assess other biases	Assess other biases	STOP*	Assess other biases	STOP*	STOP*

*Send to evidence map

*****PRIOR TO ASSESSING OTHER BIASES, CHECK PUBLICATION TO MAKE SURE OUTCOMES ARE ABSTRACTABLE. IF OUTCOMES ARE NOT ABSTRACTABLE (e.g., DATA IS PRESENTED AS GRAPHS ONLY), STOP ASSESSMENT AND CHECK WITH TEAM TO CONFIRM THAT PUBLICATION BELONGS IN EVIDENCE MAP*****

4) FIDELITY TO INTERVENTION

Table A-14. Fidelity to intervention


Description/Guiding Questions	Notes
<p>We anticipate that care delivery studies will generally fall in the range of NIH Stage 3 to 4, with the possibility that one or a few may be carried out as quality improvement and thus Stage 5. Since the Stage Model is explicitly designed to balance, or trade off, internal and external validity, we will approach risk of bias assessment as a threshold requirement rather than a continuum.</p>	<ul style="list-style-type: none"> ▪ X

FIDELITY TO INTERVENTION ASSESSMENT GUIDANCE

Table A-15. Fidelity to intervention assessment guidance

Domain	Options	Rating
Fidelity to intervention	Yes (at least 70%)	Low
	Yes-adaptation planned/ replicable	Medium
	No-adaptation not planned	High
	No/Not Reported	NR

Figure A-8. Overall risk of bias assessment guidance



VI. PRECIS-2 – Modified for Care Delivery Intervention Literature

Table A-16. PRECIS-2—modified for care delivery intervention literature

Domain	Score (1-5)	Rationale for Score	Modified Prompts, With Examples
Eligibility criteria			<p>To what extent are trial participants similar to PWD who would receive the intervention as part of usual care? [5= identical to usual care; 1=many exclusions (highly selected sample, uncommon tests used, exclude noncompliant or non-responders, etc.)]</p> <p><u>Example considerations:</u> PWD: Other comorbidities allowed? Health or behavior restrictions? Mobility or language restrictions? Dementia severity range? Small percentage of eligible chose to participate. Insurance restrictions? Participant had to opt in? Caregiver: Level of mobility/health/cognition necessary? How much time/work loss required?</p>
Recruitment path			<p>How much extra effort is required to recruit participants over usual care? [5=pragmatic, usual care (appt. or clinic); 1=targeted invitation letters, public media announcements, incentives]</p> <p><u>Example possible scores:</u> 5: Invited during routine clinic visit 4: Invitation letter/call from doctor 3: Identified PWD via diagnosis/billing code(s)→sent letter 2: Incentive(s) for participation 1: Worker hired to find participants (clinic, health plan)</p>
Setting			<p>How different is the trial setting from usual care for PWD? [5=identical to usual care; 1=single center, special trial or academic center, etc.]</p> <p><u>Example considerations (if setting not part of study question):</u> Urban only, or likely available in rural settings? Multiple settings included (private group practice, academic, HMO) Components: training for PWD/Caregiver on-site, but implemented at home via case manager?</p>
Intervention organization			<p>How different are intervention resources, provider expertise, and care organization from those available in usual care? How easy to implement without major changes (new staff, funding, policy)?</p> <p><u>Example possible scores:</u> 4: Multicomponent + requires community partners 3: Multicomponent + requires new software 2-4: Requires new staff and funding (some) 1-2: Requires new or proprietary software (1+ sites), policy change, major new staff and funding</p>
Flexibility of intervention: delivery			<p>How different is flexibility of intervention delivery from usual care for PWD?</p> <p><u>Example possible scores:</u></p>

Domain	Score (1-5)	Rationale for Score	Modified Prompts, With Examples
			5: Suggested services obtained based on ability to pay 4-5: Care manager calls/care coordination per care manager/participant discretion 2-3: short training required of PWD/Caregiver (at clinic) 1: lengthy/intensive training required of PWD/Caregiver, or at academic center 2-3: Scheduled calls from case manager (1-2 if frequent; 1 if frequent + case manager calls when needed)
Flexibility of intervention: adherence			How different is the flexibility of intervention adherence requirement from usual care? How rigorous are measures to increase adherence? (Note: rate adherence studies too) <u>Example possible scores:</u> 5: usual encouragement; 1-2: prompts/measures to improve adherence
Follow-up			How different is trial follow-up or measurement intensity from usual care? Does trial follow-up (frequency, intensity, content) result in care that differs from usual care? <u>Example possible scores</u> 5: measurement from usual follow up. 3-4: in home assessment every 6 months by case manager 1: extensive data collection, longer/more frequent clinic visits, event(s) triggered visits
Primary outcome			To what extent is the primary outcome relevant to participants? [5=obviously important; 1=intermediate or physiologic outcome, requires expert assessment, outcome timing/measure differs from usual care] <u>Example possible scores:</u> 5: important to PWD and routinely assessed in usual care 4-5: important to PWD and longer term 3-4: composite primary outcome, some elements unimportant to PWD 2-3: important to PWD but measured earlier than usual care/short-term 1-2: assessment expertise differs from usual care; surrogate, intermediate outcomes.
Primary analysis			To what extent are all data included in the analysis of the primary outcome? <u>Example possible scores:</u> 4-5: ITT or modified ITT 1-2: exclude PWD with low intervention adherence (when adherence ≠ an outcome) 1-2: post hoc-derived subgroup analysis; secondary endpoints 1-2: data merged from > 1 study 1: compliant completer analysis
Applicability			Population: PWD and/or PWD Caregivers: -narrow or broadly generalizable for PWD?
Qualifier(s)			Setting/implementation: -urban setting, practices with linked electronic health records -health plan level with trained case managers -modest vs. intensive electronic health record data extraction required?

Domain	Score (1-5)	Rationale for Score	Modified Prompts, With Examples
			<ul style="list-style-type: none"> -needs proprietary software -costs not reported but startup likely intensive -costs not reported but likely feasible addition to usual care -not likely feasible in US health system -not likely feasible in (some) rural areas

Appendix B. Excluded Studies

Could not locate

1. Cameron H, du Toit S, Richard G, et al. Using lemon balm oil to reduce aggression and agitation in dementia: results of a pilot study. *Journal of Dementia Care*. 2011;19(5):36-8. PMID: 108258536. Language: English. Entry Date: 20111104. Revision Date: 20150712. Publication Type: Journal Article.
2. Essence A. A double blind, placebo controlled trial of aromatherapy using melissa/lavender compared to aricept for the treatment of significant agitation in people with severe dementia.
3. Lpez-Almela As, Gmez-Conesa A. Intervention in dementias by multisensory stimulation (snoezelen). In *Fisioterapia*
4. Lukaszewska B, Tomaszewski W, Mirski A, et al. An evaluation of social isolation of people with dementia of the alzheimer type in the process of neurorehabilitation. In *Psychogeriatra polska*
5. Lyu J, Gao T, Li M, et al. The effect of music therapy on memory, language and psychological symptoms of patients with mild Alzheimer's disease. In *Chinese journal of neurology*
6. Reuster T, Jurjanz L, Schutzwahl M, et al. A randomized controlled trial on occupational therapy for patients with dementia and their caregivers (ERGODEM). In *Zeitschrift fur gerontopsychologie & -psychiatrie*
7. Yang D-L, Li X, Su N, et al. Family medical intervention model of senile dementia with behavioral and psychological symptoms. In *Journal of shanghai jiaotong university (medical science)*

Not English language

8. Correction: northern Manhattan Hispanic Caregiver Intervention Effectiveness Study: protocol of a pragmatic randomised trial comparing the effectiveness of two established interventions for informal caregivers of persons with dementia (*BMJ Open* (2016) 6 (e014082) DOI: 10.1136/bmjopen-2016-014082). In *BMJ Open*
9. Ban S-h, Jun S-S. Spaced Retrieval Effects in Older Adults with Mild Alzheimer's Disease. *Korean Journal of Adult Nursing*. 2012;24(4):398-405. PMID: 104400737. Language: English. Entry Date: 20130125. Revision Date: 20180305. Publication Type: Journal Article.
10. Bang M, Kim O. Effect of the Family Resilience Reinforcement Program for Family Caregivers of the Elderly with Dementia. In *Korean j adult nurs*
11. Chae K. Effects of Laughing and Music Therapy on Depression and Activities of the Autonomic Nervous System in the Elderly with Dementia. In *J korean biol nurs sci*
12. Chu J, Bao Y, Zou C. Effect of long-time retention of scalp needle on the abilities of cognition, daily living activity and P300 in vascular dementia patients. In *Zhen CI yan jiu = acupuncture research*
13. Delphin-Combe F, Rouch I, Martin-Gaujard G, et al. Effect of a non-pharmacological intervention, Voix d'Or®, on behavior disturbances in Alzheimer disease and associated disorders. In *Geriatry ET psychologie neuropsychiatrie du vieillissement*

14. Dijk A, Weert J, Droes R. Theatre as communication method in psychogeriatric care: effects on behaviour, mood and quality of life of people with dementia]. [*Dutch. In Tijdschrift voor gerontologie en geriatrie*
15. Etxeberria AI, Garcia SA, Iglesias DSA, et al. Effects of training in emotional regulation strategies on the well-being of carers of Alzheimer patients]. [*Spanish. In Revista espanola de geriatry y gerontologia*
16. Fernandez-Calvo B, Rodriguez-Perez R, Contador I, et al. Efficacy of cognitive training programs based on new software technologies in patients with Alzheimer-type dementia. In *Psicothema*
17. Fischer-Terworth C, Probst P. Effects of a psychological group intervention on neuropsychiatric symptoms and communication in Alzheimer's dementia. In *Z Gerontol Geriatr*
18. Fleiner T, Dauth H, Zijlstra W, et al. Effects of a Days-structured Training Program on mental and Behavioral Symptoms in Dementia Patients (BPSD) - Results of a randomized controlled Study. In *Z Gerontol Geriatr*
19. Garzon-Maldonado FJ, Gutierrez-Bedmar M, Serrano-Castro V, et al. An assesment of telephone assistance systems for caregivers of patients with Alzheimer's disease. *Neurologia*. 2017 Nov - Dec;32(9):595-601. doi: <https://dx.doi.org/10.1016/j.nrl.2016.03.009>. PMID: 27293022.
20. Geschke K, Scheurich A, Schermuly I, et al. Effectivity of early psychosocial counselling for family caregivers in general practioner based dementia care. In *Deutsche medizinische wochenschrift (1946)*
21. Gok Ugur H, Orak OS, Yaman Aktas Y, et al. Effects of Music Therapy on the Care Burden of In-Home Caregivers and Physiological Parameters of Their In-Home Dementia Patients: A Randomized Controlled Trial. *Complementary Med*. 2018 Nov 30;30:30. doi: <https://dx.doi.org/10.1159/000490348>. PMID: 30497077.
22. Goudour A, Samson S, Bakchine S, et al. Semantic memory training in Alzheimer's disease. In *Geriatry ET psychologie neuropsychiatrie du vieillissement*
23. Graff M, Vernooy-Dassen M, Thijssen M, et al. Home based ergotherapy in patients with dementia. In *Nederlands tijdschrift ergotherapie*
24. Grassel E, Behrmdt E, Straubmeier M. Resource-sustaining therapy in dementia: the MAKS study. In *Public health forum*
25. Haberstroh J, Neumeyer K, Schmitz B, et al. Development and evaluation of a training program for nursing home professionals to improve communication in dementia care. In *Z Gerontol Geriatr*
26. Hong S. Effects of Multi-Component Exercise and Retraining after Detraining on Functional Fitness and Cognitive Function in Elder People with Mild Dementia. In *Korean j health promot*
27. Hori M, Kubota M, Ando K, et al. The effect of videophone communication (with skype and webcam)for elderly patients with dementia and their caregivers. In *Gan to kagaku ryoho Cancer & chemotherapy*
28. Knight S, Singh I. Profile of inpatient falls in patients with dementia: a prospective comparative study between

- 100% single rooms and traditional multibedded wards. In *Journal of clinical gerontology and geriatrics*
29. Li X, Sun Y, Yan Y. Observation on effect of application of sleep restriction therapy for Alzheimer's disease patients with sleep disorders. In *Chinese nursing research*
30. Lopez N, Veliz A, Soto-Anari M, et al. Effects of a combined program of physical activity and cognitive training in Chilean patients with mild Alzheimer. In *Neurologia argentina*
31. Lukas A, Kilian R, Hay B, et al. Maintenance of health and relief for caregivers of elderly with dementia by using "initial case management". Experiences from the lighthouse project on dementia. ulm, ultdem-study. *Z Gerontol Geriatr*. 2012 June;45(4):298-309. doi: <http://dx.doi.org/10.1007/s00391-012-0337-z>. PMID: 51981666.
32. Menezes A, Aguiar A, Alves E, et al. Effectiveness of a cognitive-motorphysiotherapeutical therapy intervention in institutionalized older adults with mild cognitive impairment and mild dementia. In *Ciencia & saude coletiva*
33. Morgenstern U, Ketelhut K, Rösler D. Concentration enhancement for retaining daily living competence in dementia. In *Z Gerontol Geriatr*
34. Muller B, Staudinger B. Evaluation of an inpatient multimodal treatment concept for people with dementia. In *Pflegewissenschaft*
35. Negovanska V, Hergueta T, Guichart-Gomez E, et al. Beneficial effect of a cognitive behavioral and multidisciplinary program in Alzheimer Disease on spouse caregiver anxiety: french study ELMMA. In *Revue neurologique*
36. Pickel S, Grässel E, Luttenberger K. Efficacy of an occupational group therapy in degenerative dementias: a controlled study in the nursing home setting. In *Psychiatrische praxis*
37. Pitteri F, Soulas T, Essertaise A, et al. Contribution of relaxation to a psychoeducational intervention program for family carers of persons with Alzheimer's disease. In *Geriatric ET psychologie neuropsychiatrie du vieillissement*
38. Quibel C, Bonin M, Bonnet M, et al. Evaluation of the animal-assisted therapy in Alzheimer's disease. In *Soins Gerontologie*
39. Ravari A, Mirzaei T, Salamizadeh A, et al. Effect of the spiritual care training on anxiety reduction in home caregivers of the elderly with alzheimer disease. In *Koomesh*
40. Rexhausen E. Psychosocial care assistants caring for residents with dementia according to section 87b SGB XI. Emotional and cognitive effects. In *Pflege zeitschrift*
41. SeungHyun C, Young-ae Y. The Effect of Occupational Therapy Based Multimodal Cognitive Rehabilitation Therapy on Cognitive Function in Elderly People With Mild Dementia: a Randomized Controlled Trial.
42. Song J. Effects of a robot pet-assisted program for elderly people with dementia. In *Journal of korean academy of nursing*
43. Steinert T, Bohnet U, Flammer E, et al. Effects of a training of power and balance on the use of mechanical restraint among in-patients with dementia. In *Psychiatrische praxis*
44. Stemmer R, Grassel E, Schmid M. Individual activation of dementia sufferers in a home setting. A randomized controlled study. *Z Gerontol Geriatr*. 2019;52(3):256-63. PMID: CN-01954578.
45. Tannen A. Effectiveness of a case management program for the elderly in early stage dementia and their caregivers. In *Pflege zeitschrift*
46. Todri J, Lena O, Martinez Gil JL. A single blind randomized controlled trial of global postural re-education: Cognitive effects on Alzheimer disease patients. *European Journal of Psychiatry*. 2019. doi: <http://dx.doi.org/10.1016/j.ejpsy.2019.01.001>. PMID: 2001581733.
47. Wan C, Hu Z, Sun Y. Application of acupoint massage in nursing care of community dementia patients. In *Chinese nursing research*
48. Wang Y, Chiu C, Chen Y, et al. The efficacy of multi-sensory environment therapy on psychological, behavioral symptoms among elderly patients with dementia]. [Chinese. In *Hu li za zhi [journal of nursing]*
49. Zhang HZLHCHKLJ. Clinically multi-central randomized controlled study on scalp electroacupuncture for treatment of vascular dementia. In *Zhongguo zhen jiu [Chinese acupuncture & moxibustion]*
50. Zhou Y, Han H, Jia J. Correlation analysis on changes between cognitive ability and brain fMRI after acupoint thread embedding in Alzheimer's disease patients. In *Zhongguo zhong xi yi jie he za zhi zhongguo zhongxiyi jiehe zazhi = chinese journal of integrated traditional and western medicine*

Ineligible Intervention

51. Aasmul I, Husebo BS, Flo E. Staff Distress Improves by Treating Pain in Nursing Home Patients With Dementia: Results From a Cluster-Randomized Controlled Trial. *J Pain Symptom Manage*. 2016 Dec;52(6):795-805. doi: <https://dx.doi.org/10.1016/j.jpainsymman.2016.07.004>. PMID: 27524403.
52. Adrait A, Perrot X, Nguyen MF, et al. Do Hearing Aids Influence Behavioral and Psychological Symptoms of Dementia and Quality of Life in Hearing Impaired Alzheimer's Disease Patients and Their Caregivers? *J Alzheimers Dis*. 2017;58(1):109-21. doi: <https://dx.doi.org/10.3233/JAD-160792>. PMID: 28269769.
53. Allen V, Methven L, Gosney M. The influence of nutritional supplement drinks on providing adequate calorie and protein intake in older adults with dementia. In *J Nutr Health Aging*
54. Atri A, Shaughnessy LWLJGJH. Long-term course and effectiveness of combination therapy in Alzheimer disease. In *Alzheimer disease and associated disorders*
55. Ayalon L, Ancoli-Israel S, Stepnowsky C, et al. Adherence to continuous positive airway pressure treatment in patients with Alzheimer's disease and obstructive sleep apnea. *Am J Geriatr Psychiatry*. 2006 Feb;14(2):176-80. PMID: 16473983.
56. Beer LE, Hutchinson SR, Skala-Cordes KK. Communicating with patients who have advanced dementia: training nurse aide students. *Gerontol Geriatr*

- Educ. 2012;33(4):402-20. doi: <https://dx.doi.org/10.1080/02701960.2012.702165>. PMID: 23095223.
57. Bellantonio S, Kenny AM, Fortinsky RH, et al. Efficacy of a geriatrics team intervention for residents in dementia-specific assisted living facilities: effect on unanticipated transitions. *J Am Geriatr Soc*. 2008 Mar;56(3):523-8. doi: <https://dx.doi.org/10.1111/j.1532-5415.2007.01591.x>. PMID: 18179497.
58. Chan A, Paskavitz J, Remington R, et al. Efficacy of a vitamin/nutriceutical formulation for early-stage Alzheimer's disease: a 1-year, open-label pilot study with an 16-month caregiver extension. *Am J Alzheimers Dis Other Demen*. 2008 Dec-2009 Jan;23(6):571-85. doi: <https://dx.doi.org/10.1177/1533317508325093>. PMID: 19047474.
59. Chodosh J, Berry E, Lee M, et al. Effect of a dementia care management intervention on primary care provider knowledge, attitudes, and perceptions of quality of care. *J Am Geriatr Soc*. 2006 Feb;54(2):311-7. PMID: 43419419.
60. Cooke JR, Ayalon L, Palmer BW, et al. Sustained use of CPAP slows deterioration of cognition, sleep, and mood in patients with Alzheimer's disease and obstructive sleep apnea: a preliminary study. *J Clin Sleep Med*. 2009 Aug 15;5(4):305-9. PMID: 19968005.
61. Culo S, Mulsant BH, Rosen J, et al. Treating neuropsychiatric symptoms in dementia with Lewy bodies: a randomized controlled-trial. *Alzheimer Dis Assoc Disord*. 2010 Oct-Dec;24(4):360-4. doi: <https://dx.doi.org/10.1097/WAD.0b013e3181e6a4d7>. PMID: 20625270.
62. de Boer B, Hamers JP, Zwakhalen SM, et al. Green Care Farms as Innovative Nursing Homes, Promoting Activities and Social Interaction for People With Dementia. *J Am Med Dir Assoc*. 2017 01;18(1):40-6. doi: <https://dx.doi.org/10.1016/j.jamda.2016.10.013>. PMID: 28012503.
63. Donath C, Grassel E, Grossfeld-Schmitz M, et al. Effects of general practitioner training and family support services on the care of home-dwelling dementia patients--results of a controlled cluster-randomized study. *BMC Health Serv Res*. 2010 Nov 18;10:314. doi: <https://dx.doi.org/10.1186/1472-6963-10-314>. PMID: 21087474.
64. Dopp CM, Graff MJ, Teerenstra S, et al. A new combined strategy to implement a community occupational therapy intervention: designing a cluster randomized controlled trial. *BMC geriatr*. 2011 Mar 30;11:13. doi: <https://dx.doi.org/10.1186/1471-2318-11-13>. PMID: 21450063.
65. Dopp CM, Graff MJ, Teerenstra S, et al. Effectiveness of a multifaceted implementation strategy on physicians' referral behavior to an evidence-based psychosocial intervention in dementia: a cluster randomized controlled trial. *BMC Fam Pract*. 2013 May 30;14:70. doi: <https://dx.doi.org/10.1186/1471-2296-14-70>. PMID: 23718565.
66. Dopp CM, Graff MJ, Teerenstra S, et al. Effectiveness of a training package for implementing a community-based occupational therapy program in dementia: a cluster randomized controlled trial. *Clin Rehabil*. 2015 Oct;29(10):974-86. doi: <https://dx.doi.org/10.1177/0269215514564699>. PMID: 25547113.
67. Fabris F, Molaschi M, Aimonino N, et al. Home care for demented subjects: new models of care and home-care allowance. *Arch Gerontol Geriatr Suppl*. 2004(9):155-62. PMID: 15207409.
68. Gil Gregorio P, Ramirez Diaz SP, Ribera Casado JM, et al. Dementia and Nutrition. Intervention study in institutionalized patients with Alzheimer disease. *J Nutr Health Aging*. 2003;7(5):304-8. PMID: 12917745.
69. Gustavsson A, Cattelin F, Jonsson L. Costs of care in a mild-to-moderate Alzheimer clinical trial sample: key resources and their determinants. *Alzheimer's Dement*. 2011 Jul;7(4):466-73. doi: <https://dx.doi.org/10.1016/j.jalz.2010.06.002>. PMID: 21784355.
70. Hinchliffe AC, Katona C, Livingston G. The assessment and management of behavioural manifestations of dementia: a review and results of a controlled trial. *International Journal of Psychiatry in Clinical Practice*. 1997;1(3):157-68. doi: <https://dx.doi.org/10.3109/13651509709024722>. PMID: 24940831.
71. Hinchliffe AC, Hyman IL, Blizard B, et al. Behavioural complications of dementia - Can they be treated? *Int J Geriatr Psychiatry*. 1995;10(10):839-47. doi: <http://dx.doi.org/10.1002/gps.930101005>. PMID: 25311941.
72. Holle D, Roes M, Buscher I, et al. Process evaluation of the implementation of dementia-specific case conferences in nursing homes (FallDem): study protocol for a randomized controlled trial. *Trials*. 2014 Dec 11;15:485. doi: <https://dx.doi.org/10.1186/1745-6215-15-485>. PMID: 25496425.
73. Huizing AR, Hamers JP, Gulpers MJ, et al. Short-term effects of an educational intervention on physical restraint use: a cluster randomized trial. *BMC geriatr*. 2006 Oct 26;6:17. PMID: 17067376.
74. Iliffe S, Wilcock J, Griffin M, et al. Evidence-based interventions in dementia: A pragmatic cluster-randomised trial of an educational intervention to promote earlier recognition and response to dementia in primary care (EVIDEM-ED). *Trials*. 2010 Feb 10;11:13. doi: <https://dx.doi.org/10.1186/1745-6215-11-13>. PMID: 20146803.
75. Jordan S, Gabe-Walters ME, Watkins A, et al. Nurse-Led Medicines' Monitoring for Patients with Dementia in Care Homes: A Pragmatic Cohort Stepped Wedge Cluster Randomised Trial. *PLoS ONE*. 2015;10(10):e0140203. doi: <https://dx.doi.org/10.1371/journal.pone.0140203>. PMID: 26461064.
76. Judge TG, Urquhart A. Naftidrofuryl--a double blind cross-over study in the elderly. *Curr Med Res Opin*. 1972;1(3):166-72. PMID: 4589711.
77. Kok J, Berg I, Blankevoort G, et al. Rest-activity rhythms in small scale homelike care and traditional care for residents with dementia. In *BMC geriatr*
78. Kotynia-English R, McGowan H, Almeida OP. A randomized trial of early psychiatric intervention in residential care: impact on health outcomes. *Int Psychogeriatr*. 2005 Sep;17(3):475-85. PMID: 16252379.

79. Lea E, Marlow A, Bramble M, et al. Learning opportunities in a residential aged care facility: the role of supported placements for first-year nursing students. *J Nurs Educ*. 2014 Jul 01;53(7):410-4. doi: <https://dx.doi.org/10.3928/01484834-20140620-02>. PMID: 24972402.
80. Lee K, Cho S-J, Kim B, et al. Caregiver Preference and Treatment Compliance in Patients with Mild-to-Moderate Alzheimer's Disease in South Korea: RECAP Study Results. In *Adv Ther*
81. Livingston G, Baio G, Sommerlad A, et al. Effectiveness of an intervention to facilitate prompt referral to memory clinics in the United Kingdom: Cluster randomised controlled trial. *PLoS Medicine*. 2017 March;14 (3) (no pagination)(e1002252). doi: <http://dx.doi.org/10.1371/journal.pmed.1002252>. PMID: 615141552.
82. Logiudice D, Waltrowicz W, Brown K, et al. Do memory clinics improve the quality of life of carers? A randomized pilot trial. *Int J Geriatr Psychiatry*. 1999 Aug;14(8):626-32. PMID: 10489653.
83. Maguire A, Rosato M, O'Reilly D. Mental health and morbidity of caregivers and co-residents of individuals with dementia: a quasi-experimental design. *Int J Geriatr Psychiatry*. 2017 Oct;32(10):1104-13. doi: <https://dx.doi.org/10.1002/gps.4573>. PMID: 27594205.
84. Meeuwse EJ, Melis RJ, Van Der Aa GC, et al. Effectiveness of dementia follow-up care by memory clinics or general practitioners: randomised controlled trial. *Bmj*. 2012 May 15;344:e3086. doi: <https://dx.doi.org/10.1136/bmj.e3086>. PMID: 22589500.
85. Menn P, Holle R, Kunz S, et al. Dementia care in the general practice setting: a cluster randomized trial on the effectiveness and cost impact of three management strategies. *Value Health*. 2012 Sep-Oct;15(6):851-9. doi: <https://dx.doi.org/10.1016/j.jval.2012.06.007>. PMID: 22999135.
86. Nourhashemi F, Andrieu S, Gillette-Guyonnet S, et al. Effectiveness of a specific care plan in patients with Alzheimer's disease: cluster randomised trial (PLASA study). *Bmj*. 2010 Jun 03;340:c2466. doi: <https://dx.doi.org/10.1136/bmj.c2466>. PMID: 20522656.
87. Pai M-C, Aref H, Bassil N, et al. Real-world evaluation of compliance and preference in Alzheimer's disease treatment. In *Clin Interv Aging*
88. Peng X, Dong K. Clinical observation on acupuncture combined with Yizhi Jiannao granules for treatment of Alzheimer's disease. In *Zhongguo zhen jiu [Chinese acupuncture & moxibustion]*
89. Pivi GA, da Silva RV, Juliano Y, et al. A prospective study of nutrition education and oral nutritional supplementation in patients with Alzheimer's disease. *Nutr J*. 2011 Sep 26;10:98. doi: <https://dx.doi.org/10.1186/1475-2891-10-98>. PMID: 21943331.
90. Pond D, Mate K, Stocks N, et al. Effectiveness of a peer-mediated educational intervention in improving general practitioner diagnostic assessment and management of dementia: A cluster randomised controlled trial. *BMJ Open*. 2018 01 Aug;8 (8) (no pagination)(e021125). doi: <http://dx.doi.org/10.1136/bmjopen-2017-021125>. PMID: 623793422.
91. Shikiar R, Shakespeare A, Sagnier PP, et al. The impact of metrifonate therapy on caregivers of patients with Alzheimer's disease: Results from the MALT clinical trial. *J Am Geriatr Soc*. 2000 March;48(3):268-74. PMID: 30164209.
92. Silverberg GD, Levinthal E, Sullivan EV, et al. Assessment of low-flow CSF drainage as a treatment for AD: results of a randomized pilot study. *Neurology*. 2002 Oct 22;59(8):1139-45. PMID: 12391340.
93. Silverberg GD, Mayo M, Saul T, et al. Continuous CSF drainage in AD: results of a double-blind, randomized, placebo-controlled study. *Neurology*. 2008 Jul 15;71(3):202-9. doi: <https://dx.doi.org/10.1212/01.wnl.0000316197.04157.6f>. PMID: 18525029.
94. Smeets CH, Smalbrugge M, Gerritsen DL, et al. Improving psychotropic drug prescription in nursing home patients with dementia: design of a cluster randomized controlled trial. *BMC Psychiatry*. 2013 Nov 01;13:280. doi: <https://dx.doi.org/10.1186/1471-244X-13-280>. PMID: 24180295.
95. Sobów T, Kłoszewska I, Flirski M, et al. Predictors of long-term treatment effect of rivastigmine in Alzheimer's disease: a role for beta-amyloid plasma levels? In *Neurologia i neurochirurgia polska*
96. Steen J, Ribbe M. Pneumonia mortality risk in patients with dementia: nursing home physicians' use and evaluation of a prognostic score. In *Tijdschrift voor gerontologie en geriatric*
97. Swanson EA, Maas ML, Buckwalter KC. Alzheimer's residents' cognitive and functional measures: special and traditional care unit comparison. *Clin Nurs Res*. 1994 Feb;3(1):27-41. PMID: 8167575.
98. Testad I, Aasland AM, Aarsland D. The effect of staff training on the use of restraint in dementia: a single-blind randomised controlled trial. *Int J Geriatr Psychiatry*. 2005 Jun;20(6):587-90. PMID: 15920716.
99. Testad I, Ballard C, Bronnick K, et al. The effect of staff training on agitation and use of restraint in nursing home residents with dementia: a single-blind, randomized controlled trial. *J Clin Psychiatry*. 2010 Jan;71(1):80-6. doi: <https://dx.doi.org/10.4088/JCP.09m05486oli>. PMID: 20129008.
100. Testad I, Mekki TE, Forland O, et al. Modeling and evaluating evidence-based continuing education program in nursing home dementia care (MEDCED)--training of care home staff to reduce use of restraint in care home residents with dementia. A cluster randomized controlled trial. *Int J Geriatr Psychiatry*. 2016 Jan;31(1):24-32. doi: <https://dx.doi.org/10.1002/gps.4285>. PMID: 25845462.
101. Tian JZ, Shi J, Ni JN, et al. Sequential Therapy Based on Evolvement of Patterns: A New Model for Treatment of Alzheimer's Disease. *Chin J Integr Med*. 2019 Aug;25(8):565-73. doi: <https://dx.doi.org/10.1007/s11655-019-3066-y>. PMID: 31069693.
102. van Uffelen JG, Chin APMJ, Hopman-Rock M, et al. The effect of walking and vitamin B supplementation on quality of life in community-dwelling adults with mild cognitive impairment: a randomized, controlled trial. *Qual Life Res*. 2007 Sep;16(7):1137-46. PMID: 17616840.
103. Viola S, Viola P, Buongarzone M, et al. New brain reperfusion rehabilitation therapy improves cognitive

impairment in mild Alzheimer's disease: a prospective, controlled, open-label 12-month study with nirs correlates. In *Aging clinical and experimental research*

104. Vollmar H, Grassel E, Lauterberg J, et al. Multimodal training of general practitioners--evaluation and knowledge increase within the framework of the dementia management initiative in general medicine (IDA)]. [German. In *Zeitschrift für ärztliche Fortbildung und Qualitätssicherung*

105. Vollmar H, Mayer H, Ostermann T, et al. Knowledge transfer for the management of dementia: a cluster-randomised trial of blended learning in general practice. In *Implement Sci*

106. Vollmar HC, Butzlaff ME, Lefering R, et al. Knowledge translation on dementia: a cluster randomized trial to compare a blended learning approach with a "classical" advanced training in GP quality circles. *BMC Health Serv Res*. 2007 Jun 22;7:92. PMID: 17587452.

107. Waldorff FB, Siersma V, Nielsen B, et al. The effect of reminder letters on the uptake of an e-learning programme on dementia: a randomized trial in general practice. *Fam Pract*. 2009 Dec;26(6):466-71. doi: <https://dx.doi.org/10.1093/fampra/cmp072>. PMID: 19858123.

108. Wang F, Xiao LD, Wang K, et al. Evaluation of a WeChat-based dementia-specific training program for nurses in primary care settings: A randomized controlled trial. *Appl Nurs Res*. 2017 Dec;38:51-9. doi: <https://dx.doi.org/10.1016/j.apnr.2017.09.008>. PMID: 29241520.

109. Weiner MF, Tractenberg RE, Sano M, et al. No long-term effect of behavioral treatment on psychotropic drug use for agitation in Alzheimer's disease patients. *J Geriatr Psychiatry Neurol*. 2002;15(2):95-8. PMID: 12083600.

110. Wimo A, Winblad B, Shah SN, et al. Impact of donepezil treatment for Alzheimer's disease on caregiver time. *Curr Med Res Opin*. 2004 Aug;20(8):1221-5. PMID: 15324524.

111. Winblad B, Kawata AK, Beusterien KM, et al. Caregiver preference for rivastigmine patch relative to capsules for treatment of probable Alzheimer's disease. *Int J Geriatr Psychiatry*. 2007 May;22(5):485-91. PMID: 17407176.

112. Wolfs CA, Kessels A, Dirksen CD, et al. Integrated multidisciplinary diagnostic approach for dementia care: randomised controlled trial. *Br J Psychiatry*. 2008 Apr;192(4):300-5. doi: <https://dx.doi.org/10.1192/bjp.bp.107.035204>. PMID: 18378994.

No Eligible Outcomes

113. Ancoli-Israel S, Gehrman P, Martin JL, et al. Increased light exposure consolidates sleep and strengthens circadian rhythms in severe Alzheimer's disease patients. *Behav Sleep Med*. 2003;1(1):22-36. PMID: 15600135.

114. Ancoli-Israel S, Martin JL, Kripke DF, et al. Effect of light treatment on sleep and circadian rhythms in demented nursing home patients. *J Am Geriatr Soc*. 2002 Feb;50(2):282-9. PMID: 12028210.

115. Andrade L, Gobbi L, Coelho F, et al. Benefits of multimodal exercise intervention for postural control and

frontal cognitive functions in individuals with Alzheimer's disease: a controlled trial. In *J Am Geriatr Soc*

116. Aukner C, Eide HD, Iversen PO. Nutritional status among older residents with dementia in open versus special care units in municipal nursing homes: an observational study. *BMC geriatr*. 2013 Mar 14;13:26. doi: <https://dx.doi.org/10.1186/1471-2318-13-26>. PMID: 23496975.

117. Black DS, Cole SW, Irwin MR, et al. Yogic meditation reverses NF-kappaB and IRF-related transcriptome dynamics in leukocytes of family dementia caregivers in a randomized controlled trial. *Psychoneuroendocrinology*. 2013 Mar;38(3):348-55. doi: <https://dx.doi.org/10.1016/j.psyneuen.2012.06.011>. PMID: 22795617.

118. Bossers WJR, van der Woude LHV, Boersma F, et al. A 9-Week Aerobic and Strength Training Program Improves Cognitive and Motor Function in Patients with Dementia: A Randomized, Controlled Trial. *Am J Geriatr Psychiatry*. 2015;23(11):1106-16. doi: [10.1016/j.jagp.2014.12.191](https://dx.doi.org/10.1016/j.jagp.2014.12.191). PMID: 111672087. Language: English. Entry Date: 20160911. Revision Date: 20180629. Publication Type: journal article.

119. Brennan PF. Computer networks promote caregiving collaboration: the ComputerLink Project. *Proc Annu Symp Comput Appl Med Care*. 1992:156-60. PMID: 1482859.

120. Camargo CH, Justus FF, Retzlaff G. The effectiveness of reality orientation in the treatment of Alzheimer's disease. *Am J Alzheimers Dis Other Demen*. 2015 Aug;30(5):527-32. doi: <https://dx.doi.org/10.1177/1533317514568004>. PMID: 25588408.

121. Cavallo M, Angilletta C. Long-Lasting Neuropsychological Effects of a Computerized Cognitive Training in Patients Affected by Early Stage Alzheimer's Disease: Are They Stable Over Time? *J Appl Gerontol*. 2018 Jan 01:733464817750276. doi: <https://dx.doi.org/10.1177/0733464817750276>. PMID: 29295652.

122. Cheng ST, Chan AC, Yu EC. An exploratory study of the effect of mahjong on the cognitive functioning of persons with dementia. *Int J Geriatr Psychiatry*. 2006 Jul;21(7):611-7. PMID: 16779765.

123. Cheng ST, Chow PK, Song YQ, et al. Mental and physical activities delay cognitive decline in older persons with dementia. *Am J Geriatr Psychiatry*. 2014 Jan;22(1):63-74. doi: <https://dx.doi.org/10.1016/j.jagp.2013.01.060>. PMID: 23582750.

124. Cheng ST, Chow PK, Song YQ, et al. Can leisure activities slow dementia progression in nursing home residents? A cluster-randomized controlled trial. *Int Psychogeriatr*. 2014 Apr;26(4):637-43. doi: <https://dx.doi.org/10.1017/S1041610213002524>. PMID: 24411480.

125. Cheung DSK, Li B, Lai DWL, et al. Cognitive Stimulating Play Intervention for Dementia: A Feasibility Randomized Controlled Trial. *Am J Alzheimers Dis Other Demen*. 2019 Feb;34(1):63-71. doi: <https://dx.doi.org/10.1177/1533317518808036>. PMID: 30370782.

126. Chiverton P, Caine ED. Education to assist spouses in coping with Alzheimer's disease: a controlled trial. *J Am Geriatr Soc*. 1989;593-8. PMID: 107540167. Language: English. Entry Date: 19891201. Revision Date: 20150712. Publication Type: Journal Article.
127. Christofolletti G, Oliani M, Gobbi S, et al. A controlled clinical trial on the effects of motor intervention on balance and cognition in institutionalized elderly patients with dementia. In *Clin Rehabil*
128. Coelho FGdM, Andrade LP, Pedroso RV, et al. Multimodal exercise intervention improves frontal cognitive functions and gait in Alzheimer's disease: A controlled trial. *Geriatr Gerontol Int*. 2013;13(1):198-203. doi: 10.1111/j.1447-0594.2012.00887.x. PMID: 104404885. Language: English. Entry Date: 20130110. Revision Date: 20150711. Publication Type: Journal Article.
129. Cotelli M, Adenzato M, Cantoni V, et al. Enhancing theory of mind in behavioural variant frontotemporal dementia with transcranial direct current stimulation. *Cogn Affect Behav Neurosci*. 2018 Dec;18(6):1065-75. doi: <https://dx.doi.org/10.3758/s13415-018-0622-4>. PMID: 29995274.
130. Cotelli M, Manenti R, Cappa S, et al. Transcranial magnetic stimulation improves naming in Alzheimer disease patients at different stages of cognitive decline. In *Eur J Neurol*
131. Dijkstra K, Bourgeois M, Burgio L, et al. Effects of a communication intervention on the discourse of nursing home residents with dementia and their nursing assistants. *Journal of Medical Speech-Language Pathology*. 2002;10(2):143-57. PMID: 106787750. Language: English. Entry Date: 20031212. Revision Date: 20150711. Publication Type: Journal Article.
132. Eggermont LH, Swaab DF, Hol EM, et al. Walking the line: a randomised trial on the effects of a short term walking programme on cognition in dementia. *J Neurol Neurosurg Psychiatry*. 2009 Jul;80(7):802-4. doi: <https://dx.doi.org/10.1136/jnnp.2008.158444>. PMID: 19531688.
133. Eggermont LH, Swaab DF, Hol EM, et al. Walking the line: a randomised trial on the effects of a short term walking programme on cognition in dementia. *J Neurol Neurosurg Psychiatry*. 2009;80(7):802-4. doi: 10.1136/jnnp.2008.158444. PMID: 105369423. Language: English. Entry Date: 20090814. Revision Date: 20171118. Publication Type: journal article.
134. Eggermont LHP, Blankevoort CG, Scherder EJA. Walking and night-time restlessness in mild-to-moderate dementia: a randomized controlled trial. *Age Ageing*. 2010;39(6):746-9. PMID: 105008895. Language: English. Entry Date: 20101027. Revision Date: 20150711. Publication Type: Journal Article.
135. Elder GJ, Colloby SJ, Firbank MJ, et al. Consecutive sessions of transcranial direct current stimulation do not remediate visual hallucinations in Lewy body dementia: a randomised controlled trial. *Alzheimers Res Ther*. 2019 Jan 18;11(1):9. doi: <https://dx.doi.org/10.1186/s13195-018-0465-9>. PMID: 30658705.
136. Epstein C, Auclair U, Mittelman M. Couples Counseling in Alzheimer's Disease: First Observations of a Novel Intervention Study. *Clinical Gerontologist: The Journal of Aging and Mental Health*. 2007;30(2):21-35. doi: http://dx.doi.org/10.1300/J018v30n02_03. PMID: 2007-01997-003.
137. Eritz H, Hadjistavropoulos T, Williams J, et al. A life history intervention for individuals with dementia: a randomised controlled trial examining nursing staff empathy, perceived patient personhood and aggressive behaviours. In *Ageing and society*
138. Finnema E, de Lange J, Dröes R, et al. The quality of nursing home care: do the opinions of family members change after implementation of emotion-oriented care? *J Adv Nurs*. 2001;35(5):728-40. doi: 10.1046/j.1365-2648.2001.01905.x. PMID: 107069044. Language: English. Entry Date: 20050425. Revision Date: 20151008. Publication Type: Journal Article.
139. Frederiksen KS, Larsen CT, Hasselbalch SG, et al. A 16-Week Aerobic Exercise Intervention Does Not Affect Hippocampal Volume and Cortical Thickness in Mild to Moderate Alzheimer's Disease. *Front Aging Neurosci*. 2018;10:293. doi: <https://dx.doi.org/10.3389/fnagi.2018.00293>. PMID: 30319397.
140. Gallagher-Thompson D, Singer LS, Depp C, et al. Effective recruitment strategies for Latino and Caucasian dementia family caregivers in intervention research. *Am J Geriatr Psychiatry*. 2004 Sep-Oct;12(5):484-90. PMID: 15353386.
141. Gao H, Yan L, Liu B, et al. Clinical study on treatment of senile vascular dementia by acupuncture. *J Tradit Chin Med*. 2001 Jun;21(2):103-9. PMID: 11498896.
142. Garland K, Beer E, Eppingstall B, et al. A comparison of two treatments of agitated behavior in nursing home residents with dementia: simulated family presence and preferred music. *Am J Geriatr Psychiatry*. 2007 Jun;15(6):514-21. PMID: 17293386.
143. Garrido S, Stevens CJ, Chang E, et al. Musical Features and Affective Responses to Personalized Playlists in People With Probable Dementia. *American Journal of Alzheimer's Disease and other Dementias*. 2018. doi: <http://dx.doi.org/10.1177/1533317518808011>. PMID: 625754432.
144. Gaugler JE, Zmora R, Mitchell LL, et al. Six-Month Effectiveness of Remote Activity Monitoring for Persons Living With Dementia and Their Family Caregivers: An Experimental Mixed Methods Study. *The Gerontologist*. 2019 09 Jan;59(1):78-89. doi: <http://dx.doi.org/10.1093/geront/gny078>. PMID: 625875732.
145. Graf A, Wallner C, Schubert V, et al. The effects of light therapy on mini-mental state examination scores in demented patients. *Biol Psychiatry*. 2001 Nov 01;50(9):725-7. PMID: 11704081.
146. Hahm SC, Suh HR, Cho HY. The effect of transcutaneous electrical nerve stimulation on pain, muscle strength, balance, and gait in individuals with dementia: A double blind, pilot randomized controlled trial. *European Journal of Integrative Medicine*. 2019 August;29 (no pagination)(100932). doi: <http://dx.doi.org/10.1016/j.eujim.2019.100932>. PMID: 2002272126.
147. Harris M, Richards KC, Grando VT. The effects of slow-stroke back massage on minutes of nighttime sleep in

- persons with dementia and sleep disturbances in the nursing home: a pilot study. *J Holist Nurs*. 2012 Dec;30(4):255-63. doi: <https://dx.doi.org/10.1177/0898010112455948>. PMID: 23007716.
148. Heesterbeek M, van der Zee EA, van Heuvelen MJG. Feasibility of Three Novel Forms of Passive Exercise in a Multisensory Environment in Vulnerable Institutionalized Older Adults with Dementia. *J Alzheimers Dis*. 2019;70(3):681-90. doi: <https://dx.doi.org/10.3233/JAD-190309>. PMID: 31256137.
149. Hoefler B, Talerico KA, Rasin J, et al. Assisting cognitively impaired nursing home residents with bathing: effects of two bathing interventions on caregiving. *Gerontologist*. 2006 Aug;46(4):524-32. PMID: 16921006.
150. Innes KE, Selfe TK, Brundage K, et al. Effects of Meditation and Music-Listening on Blood Biomarkers of Cellular Aging and Alzheimer's Disease in Adults with Subjective Cognitive Decline: An Exploratory Randomized Clinical Trial. *J Alzheimers Dis*. 2018;66(3):947-70. doi: <https://dx.doi.org/10.3233/JAD-180164>. PMID: 30320574.
151. Ismail S, Christopher G, Dodd E, et al. Psychological and Mnemonic Benefits of Nostalgia for People with Dementia. *J Alzheimers Dis*. 2018;65(4):1327-44. doi: <https://dx.doi.org/10.3233/JAD-180075>. PMID: 30149444.
152. Jelcic N, Agostini M, Meneghello F, et al. Feasibility and efficacy of cognitive telerehabilitation in early Alzheimer's disease: a pilot study. *Clin Interv Aging*. 2014;9:1605-11. doi: <https://dx.doi.org/10.2147/CIA.S68145>. PMID: 25284993.
153. Jelcic N, Cagnin A, Meneghello F, et al. Effects of lexical-semantic treatment on memory in early Alzheimer disease: an observer-blinded randomized controlled trial. *Neurorehabil Neural Repair*. 2012 Oct;26(8):949-56. doi: <https://dx.doi.org/10.1177/1545968312440146>. PMID: 22460609.
154. Johnson D, Niedens M, Wilson J, et al. Treatment outcomes of a crisis intervention program for dementia with severe psychiatric complications: the Kansas bridge project. In *Gerontologist*
155. Karssemeijer EGA, Aaronson JA, Bossers WJR, et al. The quest for synergy between physical exercise and cognitive stimulation via exergaming in people with dementia: a randomized controlled trial. *Alzheimers Res Ther*. 2019 Jan 05;11(1):3. doi: <https://dx.doi.org/10.1186/s13195-018-0454-z>. PMID: 30611286.
156. Kawashima R. Mental exercises for cognitive function: clinical evidence. *J Prev Med Pub Health*. 2013 Jan;46 Suppl 1:S22-7. doi: <https://dx.doi.org/10.3961/jpmph.2013.46.S.S22>. PMID: 23412645.
157. Kemoun G, Thibaud M, Roumagne N, et al. Effects of a physical training programme on cognitive function and walking efficiency in elderly persons with dementia. *Dement Geriatr Cogn Disord*. 2010;29(2):109-14. doi: <https://dx.doi.org/10.1159/000272435>. PMID: 105169833. Language: English. Entry Date: 20100604. Revision Date: 20150711. Publication Type: Journal Article.
158. Khedr E, Foly EGN, Abo E-FN, et al. Transcranial direct current stimulation and Alzheimer's disease. In *Clin Neurophysiol*
159. Kidd LI, Zauszniewski JA, Morris DL. Benefits of a poetry writing intervention for family caregivers of elders with dementia. *Issues Ment Health Nurs*. 2011;32(9):598-604. doi: <https://dx.doi.org/10.3109/01612840.2011.576801>. PMID: 21859411.
160. Kim MJ, Han CW, Min KY, et al. Physical Exercise with Multicomponent Cognitive Intervention for Older Adults with Alzheimer's Disease: A 6-Month Randomized Controlled Trial. *Dement Geriatr Cogn Dis Extra*. 2016 May-Aug;6(2):222-32. doi: <https://dx.doi.org/10.1159/000446508>. PMID: 27403134.
161. Kuhn J, Hardenacke K, Lenartz D, et al. Deep brain stimulation of the nucleus basalis of Meynert in Alzheimer's dementia. *Mol Psychiatry*. 2015 12 Mar;20(3):353-60. doi: <http://dx.doi.org/10.1038/mp.2014.32>. PMID: 53127239.
162. Laffan A, Metzler-Baddeley C, Walker I, et al. Making errorless learning more active: self-generation in an error free learning context is superior to standard errorless learning of face-name associations in people with Alzheimer's disease. In *Neuropsychol*
163. Lee M, Cho B, Min G, et al. Effects of therapeutic recreation on the brain quotient in the elderly dementia patients. In *J Phys Ther Sci*
164. Lemke NC, Werner C, Wiloth S, et al. Transferability and Sustainability of Motor-Cognitive Dual-Task Training in Patients with Dementia: A Randomized Controlled Trial. *Gerontology*. 2019;65(1):68-83. doi: <https://dx.doi.org/10.1159/000490852>. PMID: 30041173.
165. Mortimer JA, Ding D, Borenstein AR, et al. Changes in brain volume and cognition in a randomized trial of exercise and social interaction in a community-based sample of non-demented Chinese elders. *J Alzheimers Dis*. 2012;30(4):757-66. doi: <https://dx.doi.org/10.3233/JAD-2012-120079>. PMID: 22451320.
166. Moyle W, Cooke ML, Beattie E, et al. Foot Massage and Physiological Stress in People with Dementia: A Randomized Controlled Trial. *J Altern Complement Med*. 2014;20(4):305-11. doi: [10.1089/acm.2013.0177](https://doi.org/10.1089/acm.2013.0177). PMID: 103928995. Language: English. Entry Date: 20140418. Revision Date: 20171010. Publication Type: Journal Article.
167. Nousia A, Siokas V, Aretouli E, et al. Beneficial Effect of Multidomain Cognitive Training on the Neuropsychological Performance of Patients with Early-Stage Alzheimer's Disease. *Neural Plast*. 2018;2018:2845176. doi: <https://dx.doi.org/10.1155/2018/2845176>. PMID: 30123243.
168. Pahlavanzadeh S, Heidari FG, Maghsudi J, et al. The effects of family education program on the caregiver burden of families of elderly with dementia disorders. *Iran J Nurs Midwifery Res*. 2010;15(3):102-8. PMID: 21589771.
169. Pedrinolla A, Venturelli M, Fonte C, et al. Exercise Training on Locomotion in Patients with Alzheimer's Disease: A Feasibility Study. *J Alzheimers Dis*. 2018;61(4):1599-609. doi: <https://dx.doi.org/10.3233/JAD-170625>. PMID: 29376858.
170. Perren S, Schmid R, Wettstein A. Caregivers' adaptation to change: the impact of increasing impairment

- of persons suffering from dementia on their caregivers' subjective well-being. *Aging Ment Health*. 2006 Sep;10(5):539-48. PMID: 16938689.
171. Pomeroy VM, Warren CM, Honeycombe C, et al. Mobility and dementia: is physiotherapy treatment during respite care effective? *Int J Geriatr Psychiatry*. 1999 May;14(5):389-97. PMID: 10389044.
172. Pot AM, Willemse BM, Horjus S. A pilot study on the use of tracking technology: feasibility, acceptability, and benefits for people in early stages of dementia and their informal caregivers. *Aging Ment Health*. 2012;16(1):127-34. doi: <https://dx.doi.org/10.1080/13607863.2011.596810>. PMID: 21780960.
173. Puranen T, Pitkala K, Suominen M. Tailored nutritional guidance for home-dwelling AD families: The feasibility of and elements promoting positive changes in diet (NuAD-trial). *J Nutr Health Aging*. 2015;19(4):454-9. doi: 10.1007/s12603-014-0567-4. PMID: 103787980. Language: English. Entry Date: 20150414. Revision Date: 20160331. Publication Type: Journal Article.
174. Rosen J, Mulsant BH, Kollar M, et al. Mental health training for nursing home staff using computer-based interactive video: a 6-month randomized trial. *J Am Med Dir Assoc*. 2002;3(5):291-6. PMID: 106804693. Language: English. Entry Date: 20030207. Revision Date: 20150711. Publication Type: Journal Article.
175. Rosvik J, Engedal K, Kirkevold O. Factors to make the VIPS practice model more effective in the treatment of neuropsychiatric symptoms in nursing home residents with dementia. *Dement Geriatr Cogn Disord*. 2014;37(5-6):335-46. doi: <https://dx.doi.org/10.1159/000357773>. PMID: 24503517.
176. Shelton P, Schraeder C, Dworak D, et al. Caregivers' utilization of health services: results from the Medicare Alzheimer's Disease Demonstration, Illinois site. *J Am Geriatr Soc*. 2001 Dec;49(12):1600-5. PMID: 11843991.
177. Spector A, Orrell M, Woods B. Cognitive Stimulation Therapy (CST): effects on different areas of cognitive function for people with dementia. *Int J Geriatr Psychiatry*. 2010 Dec;25(12):1253-8. doi: <https://dx.doi.org/10.1002/gps.2464>. PMID: 20069533.
178. Stocking CB, Hougham GW, Danner DD, et al. Empirical assessment of a research advance directive for persons with dementia and their proxies. *J Am Geriatr Soc*. 2007 Oct;55(10):1609-12. PMID: 17714461.
179. Swanson EA, Maas ML, Buckwalter KC. Catastrophic reactions and other behaviors of Alzheimer's residents: Special unit compared with traditional units. *Arch Psychiatr Nurs*. 1993 Oct;7(5):292-9. doi: <http://dx.doi.org/10.1016/0883-9417%2893%2990007-J>. PMID: 1994-11146-001.
180. Thyrian JR, Eichler T, Michalowsky B, et al. Community-Dwelling People Screened Positive for Dementia in Primary Care: A Comprehensive, Multivariate Descriptive Analysis Using Data from the Delphi-Study. *J Alzheimers Dis*. 2016 03 30;52(2):609-17. doi: <https://dx.doi.org/10.3233/JAD-151076>. PMID: 27031481.
181. Trebbastoni A, Imbriano L, Podda L, et al. Cognitive Training in Patients with Alzheimer's Disease: Findings of a 12-month Randomized Controlled Trial. *Curr Alzheimer Res*. 2018 Mar 14;15(5):452-61. doi: <https://dx.doi.org/10.2174/1567205014666171113105044>. PMID: 29141548.
182. Tsantali E, Economidis D, Rigopoulou S. Testing the benefits of cognitive training vs. cognitive stimulation in mild Alzheimer's disease: A randomised controlled trial. *Brain Impairment*. 2017 01 Sep;18(2):188-96. doi: <http://dx.doi.org/10.1017/BrImp.2017.6>. PMID: 615666653.
183. van Weert JC, Janssen BM, van Dulmen AM, et al. Nursing assistants' behaviour during morning care: effects of the implementation of snoezelen, integrated in 24-hour dementia care. *J Adv Nurs*. 2006 Mar;53(6):656-68. PMID: 16553674.
184. Werner C, Wiloth S, Lemke NC, et al. People with Dementia Can Learn Compensatory Movement Maneuvers for the Sit-to-Stand Task: A Randomized Controlled Trial. *J Alzheimers Dis*. 2017;60(1):107-20. doi: <https://dx.doi.org/10.3233/JAD-170258>. PMID: 28759967.
185. Yaguez L, Shaw KN, Morris R, et al. The effects on cognitive functions of a movement-based intervention in patients with Alzheimer's type dementia: a pilot study. *Int J Geriatr Psychiatry*. 2011 Feb;26(2):173-81. doi: <https://dx.doi.org/10.1002/gps.2510>. PMID: 20878665.
186. Young DK. Multicomponent intervention combining a cognitive stimulation group and tai chi to reduce cognitive decline among community-dwelling older adults with probable dementia: A multi-center, randomized controlled trial. *Dementia*. 2018 Nov 28;1471301218814637. doi: <https://dx.doi.org/10.1177/1471301218814637>. PMID: 30486656.
187. Zhang D, Zhou Z, Wu W, et al. Influence of electroencephalograph bionic electrical stimulation on neuronal activities in patients with Alzheimer's disease: A functional magnetic resonance imaging study. *Radiology of Infectious Diseases*. 2018 March;5(1):26-32. doi: <http://dx.doi.org/10.1016/j.jrid.2018.02.005>. PMID: 620922742.
188. Zhao J, Li Z, Cong Y, et al. Repetitive transcranial magnetic stimulation improves cognitive function of Alzheimer's disease patients. *Oncotarget*. 2017 May 16;8(20):33864-71. doi: <https://dx.doi.org/10.18632/oncotarget.13060>. PMID: 27823981.

Ineligible Population

189. Allen VJ, Methven L, Gosney M. Impact of serving method on the consumption of nutritional supplement drinks: randomized trial in older adults with cognitive impairment. *J Adv Nurs*. 2014;70(6):1323-33. doi: 10.1111/jan.12293. PMID: 103928802. Language: English. Entry Date: 20140418. Revision Date: 20150819. Publication Type: Journal Article.
190. Alves J, Alves-Costa F, Magalhaes R, et al. Cognitive stimulation for Portuguese older adults with cognitive impairment: a randomized controlled trial of efficacy, comparative duration, feasibility, and experiential relevance. *Am J Alzheimers Dis Other Demen*. 2014 Sep;29(6):503-12. doi: <https://dx.doi.org/10.1177/1533317514522541>. PMID: 24526760.

191. Bakker TJ, Duivenvoorden HJ, van der Lee J, et al. Prognostic factors for a favourable long-term outcome from an integrative psychotherapeutic nursing home programme. *Dement Geriatr Cogn Disord*. 2011;32(5):318-31. doi: <https://dx.doi.org/10.1159/000334969>. PMID: 22286593.
192. Bakker TJ, Duivenvoorden HJ, van der Lee J, et al. Integrative psychotherapeutic nursing home program to reduce multiple psychiatric symptoms of cognitively impaired patients and caregiver burden: randomized controlled trial. *Am J Geriatr Psychiatry*. 2011;19(6):507-20. doi: [10.1097/JGP.0b013e3181eafdc6](https://doi.org/10.1097/JGP.0b013e3181eafdc6). PMID: 108240694. Language: English. Entry Date: 20110930. Revision Date: 20171129. Publication Type: journal article.
193. Ball K, Berch DB, Helmers KF, et al. Effects of cognitive training interventions with older adults: a randomized controlled trial. *JAMA: Journal of the American Medical Association*. 2002;288(18):2271-81. PMID: 106804253. Corporate Author: Advanced Cognitive Training for Independent and Vital Elderly Study Group. Language: English. Entry Date: 20030207. Revision Date: 20161125. Publication Type: journal article.
194. Barban F, Annicchiario R, Pantelopoulos S, et al. Protecting cognition from aging and Alzheimer's disease: a computerized cognitive training combined with reminiscence therapy. *Int J Geriatr Psychiatry*. 2016 Apr;31(4):340-8. doi: <https://dx.doi.org/10.1002/gps.4328>. PMID: 26205305.
195. Bateman C, Anderson K, Bird M, et al. Volunteers improving person-centred dementia and delirium care in a rural Australian hospital. *Rural Remote Health*. 2016 Apr-Jun;16(2):3667. PMID: 27303861.
196. Bautmans I, Demarteau J, Cruys B, et al. Dysphagia in elderly nursing home residents with severe cognitive impairment can be attenuated by cervical spine mobilization. *Journal of Rehabilitation Medicine (Stiftelsen Rehabiliteringsinformation)*. 2008;40(9):755-60. PMID: 105703943. Language: English. Entry Date: 20081205. Revision Date: 20150819. Publication Type: Journal Article.
197. Beer C, Horner B, Flicker L, et al. A cluster-randomised trial of staff education to improve the quality of life of people with dementia living in residential care: the DIRECT study. *PLoS ONE*. 2011;6(11):e28155. doi: <https://dx.doi.org/10.1371/journal.pone.0028155>. PMID: 22140531.
198. Beer C, Lowry R, Horner B, et al. Development and evaluation of an educational intervention for general practitioners and staff caring for people with dementia living in residential facilities. *Int Psychogeriatr*. 2011 Mar;23(2):221-9. doi: <https://dx.doi.org/10.1017/S104161021000195X>. PMID: 20950507.
199. Bier N, Grenier S, Brodeur C, et al. Measuring the impact of cognitive and psychosocial interventions in persons with mild cognitive impairment with a randomized single-blind controlled trial: rationale and design of the MEMO+ study. *Int Psychogeriatr*. 2015 Mar;27(3):511-25. doi: <https://dx.doi.org/10.1017/S1041610214001902>. PMID: 25268968.
200. Bird M, Jones RH, Korten A, et al. A controlled trial of a predominantly psychosocial approach to BPSD: treating causality. *Int Psychogeriatr*. 2007;19(5):874-91. PMID: 105828624. Language: English. Entry Date: 20080307. Revision Date: 20161117. Publication Type: journal article.
201. Blair A, Anderson K, Bateman C. The "Golden Angels": effects of trained volunteers on specialising and readmission rates for people with dementia and delirium in rural hospitals. *Int Psychogeriatr*. 2018 Nov;30(11):1707-16. doi: <https://dx.doi.org/10.1017/S1041610218000911>. PMID: 30099973.
202. Brown LJE, Blake D, Berry K, et al. A pilot and feasibility study of the effectiveness of care mapping on person-centred care in neurorehabilitation settings. *Neuropsychol*. 2019 08 Apr;1-18. doi: <http://dx.doi.org/10.1080/09602011.2019.1598446>. PMID: 627288558.
203. Buettner LL, Fitzsimmons S, Atav S, et al. Cognitive stimulation for apathy in probable early-stage Alzheimer's. *J Aging Res*. 2011 Apr 28;2011:480890. doi: <https://dx.doi.org/10.4061/2011/480890>. PMID: 21584241.
204. Burgio LD, Allen-Burge R, Roth DL, et al. Come talk with me: improving communication between nursing assistants and nursing home residents during care routines. *Gerontologist*. 2001;41(4):449-60. PMID: 107066936. Language: English. Entry Date: 20011109. Revision Date: 20150711. Publication Type: Journal Article.
205. Buschert VC, Friese U, Teipel SJ, et al. Effects of a newly developed cognitive intervention in amnesic mild cognitive impairment and mild Alzheimer's disease: a pilot study. *J Alzheimers Dis*. 2011;25(4):679-94. doi: <https://dx.doi.org/10.3233/JAD-2011-100999>. PMID: 21483095.
206. Catic A, Mattison M, Bakaev I, et al. ECHO-AGE: an Innovative Model of Geriatric Care for Long-Term Care Residents With Dementia and Behavioral Issues. In *J Am Med Dir Assoc*
207. Chiu wu CH, Perng SJ, Shi CK, et al. Advance Care Planning and Advance Directives: A Multimedia Education Program in Community-Dwelling Older Adults. *Journal of Applied Gerontology*. 2019. doi: <http://dx.doi.org/10.1177/0733464819831596>.
208. Conradsson M, Littbrand H, Lindelöf N, et al. Effects of a high-intensity functional exercise programme on depressive symptoms and psychological well-being among older people living in residential care facilities: a cluster-randomized controlled trial. *Aging Ment Health*. 2010;14(5):565-76. doi: [10.1080/13607860903483078](https://doi.org/10.1080/13607860903483078). PMID: 105049096. Language: English. Entry Date: 20100827. Revision Date: 20150711. Publication Type: Journal Article.
209. Corbett A, Owen A, Hampshire A, et al. The Effect of an Online Cognitive Training Package in Healthy Older Adults: An Online Randomized Controlled Trial. *J Am Med Dir Assoc*. 2015 Nov 01;16(11):990-7. doi: <https://dx.doi.org/10.1016/j.jamda.2015.06.014>. PMID: 26543007.
210. Cox KL, Cyarto EV, Ellis KA, et al. A Randomized Controlled Trial of Adherence to a 24-Month Home-Based Physical Activity Program and the Health Benefits for Older Adults at Risk of Alzheimer's Disease: The AIBL Active-Study. *J Alzheimers Dis*. 2018 Oct 26;26:26. doi: <https://dx.doi.org/10.3233/JAD-180521>. PMID: 30372680.

211. Dimitriou TD, Verykoui E, Papatriantafyllou J, et al. Non-pharmacological interventions for agitation/aggressive behaviour in patients with dementia: a randomized controlled crossover trial. *Funct Neurol*. 2018 Jul;33(3):143-7. PMID: 30457967.
212. Folkerts AK, Dorn ME, Roheger M, et al. Cognitive Stimulation for Individuals with Parkinson's Disease Dementia Living in Long-Term Care: Preliminary Data from a Randomized Crossover Pilot Study. *Parkinson's dis*. 2018;2018:8104673. doi: <https://dx.doi.org/10.1155/2018/8104673>. PMID: 30631420.
213. Gaitan A, Garolera M, Cerulla N, et al. Efficacy of an adjunctive computer-based cognitive training program in amnesic mild cognitive impairment and Alzheimer's disease: a single-blind, randomized clinical trial. *Int J Geriatr Psychiatry*. 2013 Jan;28(1):91-9. doi: <https://dx.doi.org/10.1002/gps.3794>. PMID: 22473855.
214. Garand L, Morse JQ, ChiaRebecca L, et al. Problem-solving therapy reduces subjective burden levels in caregivers of family members with mild cognitive impairment or early-stage dementia: Secondary analysis of a randomized clinical trial. *Int J Geriatr Psychiatry*. 2019 07;34(7):957-65. doi: <https://dx.doi.org/10.1002/gps.5095>. PMID: 30868641.
215. Garand L, Rinaldo DE, Alberth MM, et al. Effects of problem solving therapy on mental health outcomes in family caregivers of persons with a new diagnosis of mild cognitive impairment or early dementia: a randomized controlled trial. *Am J Geriatr Psychiatry*. 2014;22(8):771-81. doi: 10.1016/j.jagp.2013.07.007. PMID: 107797083. Language: English. Entry Date: 20150417. Revision Date: 20161207. Publication Type: journal article.
216. Goldberg SE, Bradshaw LE, Kearney FC, et al. Care in specialist medical and mental health unit compared with standard care for older people with cognitive impairment admitted to general hospital: randomised controlled trial (NIHR TEAM trial). *Bmj*. 2013 Jul 02;347:f4132. doi: <https://dx.doi.org/10.1136/bmj.f4132>. PMID: 23819964.
217. Gossink F, Pijnenburg Y, Scheltens P, et al. An intervention programme for caregivers of dementia patients with frontal behavioural changes: an explorative study with controlled effect on sense of competence. *Psychogeriatrics*. 2018 Nov;18(6):451-9. doi: <https://dx.doi.org/10.1111/psyg.12351>. PMID: 30066463.
218. Gregory MA, Gill DP, Shellington EM, et al. Group-based exercise and cognitive-physical training in older adults with self-reported cognitive complaints: The Multiple-Modality, Mind-Motor (M4) study protocol. *BMC geriatr*. 2016 Jan 16;16:17. doi: <https://dx.doi.org/10.1186/s12877-016-0190-9>. PMID: 26772171.
219. Gustafsson M, Sjolander M, Pfister B, et al. Pharmacist participation in hospital ward teams and hospital readmission rates among people with dementia: a randomized controlled trial. *Eur J Clin Pharmacol*. 2017 Jul;73(7):827-35. doi: <https://dx.doi.org/10.1007/s00228-017-2249-8>. PMID: 28391409.
220. Gustafsson M, Sjolander M, Pfister B, et al. Effects of Pharmacists' Interventions on Inappropriate Drug Use and Drug-Related Readmissions in People with Dementia-A Secondary Analysis of a Randomized Controlled Trial. *Pharmacy (Basel)*. 2018 Jan 16;6(1):16. doi: <https://dx.doi.org/10.3390/pharmacy6010007>. PMID: 29337859.
221. Han JW, Lee H, Hong JW, et al. Multimodal Cognitive Enhancement Therapy for Patients with Mild Cognitive Impairment and Mild Dementia: A Multi-Center, Randomized, Controlled, Double-Blind, Crossover Trial. *J Alzheimers Dis*. 2017;55(2):787-96. PMID: 120361850. Language: English. Entry Date: 20180725. Revision Date: 20180731. Publication Type: journal article.
222. Herrera C, Chambon C, Michel BF, et al. Positive effects of computer-based cognitive training in adults with mild cognitive impairment. *Neuropsychologia*. 2012 Jul;50(8):1871-81. doi: <https://dx.doi.org/10.1016/j.neuropsychologia.2012.04.012>. PMID: 22525705.
223. Hindle J, Watermeyer T, Roberts J, et al. Cognitive rehabilitation for Parkinson's disease dementia: a study protocol for a pilot randomised controlled trial. *In Trials*
224. Holmes D, Teresi JA, Ramirez M, et al. An evaluation of a monitoring system intervention: falls, injuries, and affect in nursing homes. *Clin Nurs Res*. 2007 Nov;16(4):317-35. PMID: 17991911.
225. Ito T, Meguro K, Akanuma K, et al. A randomized controlled trial of the group reminiscence approach in patients with vascular dementia. *Dement Geriatr Cogn Disord*. 2007;24(1):48-54. PMID: 17565213.
226. Jansen AP, van Hout HP, Nijpels G, et al. Effectiveness of case management among older adults with early symptoms of dementia and their primary informal caregivers: a randomized clinical trial. *Int J Nurs Stud*. 2011 Aug;48(8):933-43. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2011.02.004>. PMID: 21356537.
227. Jansen AP, van Hout HP, van Marwijk HW, et al. (Cost)-effectiveness of case-management by district nurses among primary informal caregivers of older adults with dementia symptoms and the older adults who receive informal care: design of a randomized controlled trial [ISCRTN83135728]. *BMC Public Health*. 2005 Dec 12;5:133. PMID: 16343336.
228. Jha A, Jan F, Gale T, et al. Effectiveness of a recovery-orientated psychiatric intervention package on the wellbeing of people with early dementia: a preliminary randomised controlled trial. *Int J Geriatr Psychiatry*. 2013 Jun;28(6):589-96. doi: <https://dx.doi.org/10.1002/gps.3863>. PMID: 22847712.
229. Jin Z. Efficacy of acupuncture combined with Nimodipine in the treatment of vascular dementia. In *China and foreign medical treatment [zhong wai yi liao]*
230. Keller HH, Gibbs AJ, Boudreau LD, et al. Prevention of weight loss in dementia with comprehensive nutritional treatment. *J Am Geriatr Soc*. 2003 Jul;51(7):945-52. PMID: 12834514.
231. Kiosses DN, Ravdin LD, Gross JJ, et al. Problem adaptation therapy for older adults with major depression and cognitive impairment: a randomized clinical trial. *JAMA Psychiatry*. 2015;72(1):22-30. doi: 10.1001/jamapsychiatry.2014.1305. PMID: 103874772. Language: English. Entry Date: 20150410. Revision Date: 20161222. Publication Type: journal article.

232. Lam LC, Chau RC, Wong BM, et al. Interim follow-up of a randomized controlled trial comparing Chinese style mind body (Tai Chi) and stretching exercises on cognitive function in subjects at risk of progressive cognitive decline. *Int J Geriatr Psychiatry*. 2011 Jul;26(7):733-40. doi: <https://dx.doi.org/10.1002/gps.2602>. PMID: 21495078.
233. Larouche E, Hudon C, Goulet S. Mindfulness mechanisms and psychological effects for aMCI patients: A comparison with psychoeducation. *Complementary therapies in clinical practice*. 2019 01 Feb;34:93-104. doi: <http://dx.doi.org/10.1016/j.ctcp.2018.11.008>. PMID: 626243559.
234. Lautenschlager NT, Cox KL, Flicker L, et al. Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. [Erratum appears in *JAMA*. 2009 Jan 21;301(3):276]. *Jama*. 2008 Sep 03;300(9):1027-37. doi: <https://dx.doi.org/10.1001/jama.300.9.1027>. PMID: 18768414.
235. Lee GJ, Bang HJ, Lee KM, et al. A comparison of the effects between 2 computerized cognitive training programs, Bettercog and COMCOG, on elderly patients with MCI and mild dementia: A single-blind randomized controlled study. *Medicine (Baltimore)*. 2018 Nov;97(45):e13007. doi: <https://dx.doi.org/10.1097/MD.00000000000013007>. PMID: 30407291.
236. Lin HC, Yang YP, Cheng WY, et al. Distinctive effects between cognitive stimulation and reminiscence therapy on cognitive function and quality of life for different types of behavioural problems in dementia. *Scand J Caring Sci*. 2018 Jun;32(2):594-602. doi: <https://dx.doi.org/10.1111/scs.12484>. PMID: 28881430.
237. Littbrand H, Rosendahl E, Lindelöf N, et al. A high-intensity functional weight-bearing exercise program for older people dependent in activities of daily living and living in residential care facilities: evaluation of the applicability with focus on cognitive function. *Phys Ther*. 2006;86(4):489-98. PMID: 106471666. Language: English. Entry Date: 20060714. Revision Date: 20150820. Publication Type: Journal Article.
238. Liu-Ambrose T, Best JR, Davis JC, et al. Aerobic exercise and vascular cognitive impairment: A randomized controlled trial. *Neurology*. 2016;87(20):2082-90. PMID: 119524521. Language: English. Entry Date: 20180723. Revision Date: 20180723. Publication Type: journal article.
239. Low L-F, Goodenough B, Fletcher J, et al. The Effects of Humor Therapy on Nursing Home Residents Measured Using Observational Methods: The SMILE Cluster Randomized Trial. *J Am Med Dir Assoc*. 2014;15(8):564-9. doi: [10.1016/j.jamda.2014.03.017](https://doi.org/10.1016/j.jamda.2014.03.017). PMID: 103984346. Language: English. Entry Date: 20140805. Revision Date: 20150710. Publication Type: Journal Article.
240. Man DW, Chung JC, Lee GY. Evaluation of a virtual reality-based memory training programme for Hong Kong Chinese older adults with questionable dementia: a pilot study. *Int J Geriatr Psychiatry*. 2012 May;27(5):513-20. doi: <https://dx.doi.org/10.1002/gps.2746>. PMID: 21681818.
241. Meinzer M, Lindenberg R, Phan MT, et al. Transcranial direct current stimulation in mild cognitive impairment: Behavioral effects and neural mechanisms. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association*. 2015 Sep;11(9):1032-40. doi: <http://dx.doi.org/10.1016/j.jalz.2014.07.159>. PMID: 2014-52911-001.
242. Middleton LE, Ventura MI, Santos-Modesitt W, et al. The Mental Activity and eXercise (MAX) trial: Effects on physical function and quality of life among older adults with cognitive complaints. *Contemp Clin Trials*. 2018 01;64:161-6. doi: <https://dx.doi.org/10.1016/j.cct.2017.10.009>. PMID: 29066293.
243. Muniz R, Serra CM, Reisberg B, et al. Cognitive-motor intervention in Alzheimer's disease: long-term results from the Maria Wolff trial. *J Alzheimers Dis*. 2015;45(1):295-304. doi: <https://dx.doi.org/10.3233/JAD-142364>. PMID: 25547632.
244. Nakatsuka M, Nakamura K, Hamanoso R, et al. A Cluster Randomized Controlled Trial of Nonpharmacological Interventions for Old-Old Subjects with a Clinical Dementia Rating of 0.5: The Kurihara Project. *Dement Geriatr Cogn Dis Extra*. 2015 May-Aug;5(2):221-32. doi: <https://dx.doi.org/10.1159/000380816>. PMID: 26195978.
245. Okamoto T, Hashimoto Y, Kobayashi R. Effects of interval walking training compared to normal walking training on cognitive function and arterial function in older adults: a randomized controlled trial. *Aging Clinical and Experimental Research*. 2018. doi: <http://dx.doi.org/10.1007/s40520-018-1093-8>. PMID: 625589391.
246. Olazarán J, Muniz R, Reisberg B, et al. Benefits of cognitive-motor intervention in MCI and mild to moderate Alzheimer disease. *Neurology*. 2004 Dec 28;63(12):2348-53. PMID: 15623698.
247. Padala PR, Padala KP, Lensing SY, et al. Repetitive transcranial magnetic stimulation for apathy in mild cognitive impairment: A double-blind, randomized, sham-controlled, cross-over pilot study. *Psychiatry Res*. 2018 03;261:312-8. doi: <https://dx.doi.org/10.1016/j.psychres.2017.12.063>. PMID: 29331848.
248. Park JE, Jeon SY, Kim SA, et al. A Multidomain Intervention for Modifying Lifestyle Habits Reduces the Dementia Risk in Community-Dwelling Older Adults: A Single-Blinded Randomized Controlled Pilot Study. *J Alzheimers Dis*. 2019;70(1):51-60. doi: <https://dx.doi.org/10.3233/JAD-190016>. PMID: 31127782.
249. Quayhagen MP, Quayhagen M, Corbeil RR, et al. Coping with dementia: evaluation of four nonpharmacologic interventions. *Int Psychogeriatr*. 2000 Jun;12(2):249-65. PMID: 10937544.
250. Regan B, Wells Y, Farrow M, et al. MAXCOG-maximizing cognition: a randomized controlled trial of the efficacy of goal-oriented cognitive rehabilitation for people with mild cognitive impairment and early Alzheimer disease. In *Am J Geriatr Psychiatry*
251. Resnick B, Gruber-Baldini AL, Zimmerman S, et al. Nursing home resident outcomes from the Res-Care intervention. *J Am Geriatr Soc*. 2009;57(7):1156-65. doi: [10.1111/j.1532-5415.2009.02327.x](https://doi.org/10.1111/j.1532-5415.2009.02327.x). PMID: 105377579.

- Language: English. Entry Date: 20090904. Revision Date: 20150711. Publication Type: Journal Article.
252. Rovio S, Kareholt I, Helkala EL, et al. Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. *Lancet neurol.* 2005 Nov;4(11):705-11. PMID: 16239176.
253. Sakamoto Y, Ebihara S, Ebihara T, et al. Fall Prevention Using Olfactory Stimulation with Lavender Odor in Elderly Nursing Home Residents: A Randomized Controlled Trial...[corrected][published erratum appears in *J AM GERIATR SOC* 2012 Nov; 60(11): 2193]. *J Am Geriatr Soc.* 2012;60(6):1005-11. doi: 10.1111/j.1532-5415.2012.03977.x. PMID: 104462071. Language: English. Entry Date: 20120615. Revision Date: 20150711. Publication Type: Journal Article.
254. Schaap FD, Dijkstra GJ, Stewart RE, et al. Effects of Dementia Care Mapping on well-being and quality of life of older people with intellectual disability: A quasi-experimental study. *J Appl Res Intellect Disabil.* 2019 Jul;32(4):849-60. doi: <https://dx.doi.org/10.1111/jar.12576>. PMID: 30868692.
255. Schaap FD, Finnema EJ, Stewart RE, et al. Effects of Dementia Care Mapping on job satisfaction and caring skills of staff caring for older people with intellectual disabilities: A quasi-experimental study. *J Appl Res Intellect Disabil.* 2019 Sep;32(5):1228-40. doi: <https://dx.doi.org/10.1111/jar.12615>. PMID: 31087472.
256. Schindel-Martin L, Morden P, Cetinski G, et al. Teaching staff to respond effectively to cognitively impaired residents who display self-protective behaviors. *Am J Alzheimers Dis Other Demen.* 2003;18(5):273-81. PMID: 106741786. Language: English. Entry Date: 20040604. Revision Date: 20150711. Publication Type: Journal Article.
257. Seltzer MM, Litchfield LC, Kapust LR, et al. Professional and family collaboration in case management: a hospital-based replication of a community-based study. *Soc Work Health Care.* 1992;17(1):1-22. PMID: 1440105.
258. Shelton JT, Lee JH, Scullin MK, et al. Improving Prospective Memory in Healthy Older Adults and Individuals with Very Mild Alzheimer's Disease. *J Am Geriatr Soc.* 2016 Jun;64(6):1307-12. doi: <https://dx.doi.org/10.1111/jgs.14134>. PMID: 27321610.
259. Sink KM, Espeland MA, Castro CM, et al. Effect of a 24-Month Physical Activity Intervention vs Health Education on Cognitive Outcomes in Sedentary Older Adults: The LIFE Randomized Trial. *Jama.* 2015 Aug 25;314(8):781-90. doi: <https://dx.doi.org/10.1001/jama.2015.9617>. PMID: 26305648.
260. Sink KM, Espeland MA, Rushing J, et al. The LIFE Cognition Study: design and baseline characteristics. *Clin Interv Aging.* 2014;9:1425-36. doi: <https://dx.doi.org/10.2147/CIA.S65381>. PMID: 25210447.
261. Tappen R, Hain D. The effect of in-home cognitive training on functional performance of individuals with mild cognitive impairment and early-stage Alzheimer's disease. In *Res*
262. Thodberg K, Sorensen LU, Christensen JW, et al. Therapeutic effects of dog visits in nursing homes for the elderly. *Psychogeriatrics.* 2016 Sep;16(5):289-97. doi: <https://dx.doi.org/10.1111/psyg.12159>. PMID: 26510632.
263. Ukawa S, Yuasa M, Ikeno T, et al. Randomised controlled pilot study in Japan comparing a home visit program using a Functioning Improvement Tool with a home visit with conversation alone. *Australas J Ageing.* 2012 Sep;31(3):187-9. doi: <https://dx.doi.org/10.1111/j.1741-6612.2012.00589.x>. PMID: 22950591.
264. Unverzagt FW, Smith DM, Rebok GW, et al. The Indiana Alzheimer Disease Center's Symposium on Mild Cognitive Impairment. Cognitive training in older adults: lessons from the ACTIVE Study. *Curr Alzheimer Res.* 2009 Aug;6(4):375-83. PMID: 19689237.
265. Van Der Lee J, Bakker TJEM, Dros RM. Recovery from burden: Informal caregiver profiles that predict treatment success. *Int Psychogeriatr.* 2019;317-29. doi: <http://dx.doi.org/10.1017/S1041610218000613>. PMID: 622667094.
266. Wang S-C, Yu C-L, Chang S-H. Effect of music care on depression and behavioral problems in elderly people with dementia in Taiwan: A quasi-experimental, longitudinal study. *Aging Ment Health.* 2017 Feb;21(2):156-62. doi: <http://dx.doi.org/10.1080/13607863.2015.1093602>. PMID: 2017-03789-005.
267. Wang Y. Study of treatment for post stroke dementia.
268. Wang Y, Xiao LD, Ullah S, et al. Evaluation of a nurse-led dementia education and knowledge translation programme in primary care: A cluster randomized controlled trial. *Nurse Educ Today.* 2017 Feb;49:1-7. doi: <https://dx.doi.org/10.1016/j.nedt.2016.10.016>. PMID: 27855297.
269. Wilcock J, Iliffe S, Griffin M, et al. Tailored educational intervention for primary care to improve the management of dementia: the EVIDEM-ED cluster randomized controlled trial. *Trials.* 2013 Nov 20;14:397. doi: <https://dx.doi.org/10.1186/1745-6215-14-397>. PMID: 24257429.
270. Wilson KC, Scott M, Abou-Saleh M, et al. Long-term effects of cognitive-behavioural therapy and lithium therapy on depression in the elderly. *Br J Psychiatry.* 1995 Nov;167(5):653-8. PMID: 8564323.
271. Yang BF, Zeng XH, Liu Y, et al. Effect of acupuncture treatment on vascular cognitive impairment without dementia: Study protocol for a randomized controlled trial. *Trials.* 2015;13. doi: <http://dx.doi.org/10.1186/1745-6215-15-442>. PMID: 605062793.
272. Yang JW, Shi GX, Zhang S, et al. Effectiveness of acupuncture for vascular cognitive impairment no dementia: a randomized controlled trial. *Clin Rehabil.* 2019 Jan 23;269215518819050. doi: <https://dx.doi.org/10.1177/0269215518819050>. PMID: 30672317.
273. Young DK, Ng PY, Kwok T, et al. The effects of an expanded cognitive stimulation therapy model on the improvement of cognitive ability of elderly with mild stage Dementia living in a community - a randomized waitlist controlled trial. *Aging Ment Health.* 2018 May 21:1-8. doi: <https://dx.doi.org/10.1080/13607863.2018.1471586>. PMID: 29781725.

274. Yu J, Zhang X, Liu C, et al. Effect of acupuncture treatment on vascular dementia. *Neurol Res.* 2006 Jan;28(1):97-103. PMID: 16464371.
275. Zenthofer A, Meyer-Kuhling I, Hufeland A-L, et al. Carers' education improves oral health of older people suffering from dementia - Results of an intervention study. In *Clin Interv Aging*
276. Zhao J, Li H, Lin R, et al. Effects of creative expression therapy for older adults with mild cognitive impairment at risk of Alzheimer's disease: a randomized controlled clinical trial. *Clin Interv Aging.* 2018;13:1313-20. doi: <https://dx.doi.org/10.2147/CIA.S161861>. PMID: 30087557.
277. Zhu Y, Wu H, Qi M, et al. Effects of a specially designed aerobic dance routine on mild cognitive impairment. *Clin Interv Aging.* 2018;13:1691-700. doi: <https://dx.doi.org/10.2147/CIA.S163067>. PMID: 30237705.

Ineligible publication type

278. Abalan F, Manciet G, Dartigues JF, et al. Nutrition and SDAT. *Biol Psychiatry.* 1992 Jan 01;31(1):103-5. PMID: 1543795.
279. Agar M, Chenoweth L, Mitchell G, et al. Implementing facilitated case conferencing for aged care residents with advanced dementia-development of a palliative care planning coordinator role. In *Palliat Med*
280. Agar M, Chenoweth L, Mitchell G, et al. Cluster randomised controlled trial of facilitated case conferencing for aged care residents with advanced dementia. In *Palliat Med*
281. Amieva H, Dartigues JF. ETNA3, a clinical randomized study assessing three cognitive-oriented therapies in dementia: Rationale and general design. *Revue Neurologique.* 2013 October;169(10):752-6. doi: <http://dx.doi.org/10.1016/j.neurol.2013.07.011>. PMID: 52760976.
282. Ballard C. Video decision support affects patients' care choices in dementia care planning. In *Evidence-based mental health*
283. Brennan PF, Moore SM, Smyth KA. ComputerLink: electronic support for the home caregiver. *ANS Adv Nurs Sci.* 1991 Jun;13(4):14-27. PMID: 2059002.
284. Brodaty H, Arasaratnam C. Meta-analysis of nonpharmacological interventions for neuropsychiatric symptoms of dementia.[Erratum appears in *Am J Psychiatry.* 2012 Sep;169(9):946-53. doi: <https://dx.doi.org/10.1176/appi.ajp.2012.11101529>. PMID: 22952073.
285. Brodaty H, Green A, Koschera A. Meta-analysis of psychosocial interventions for caregivers of people with dementia. *J Am Geriatr Soc.* 2003 May;51(5):657-64. PMID: 12752841.
286. Capisizu A, Zamfirescu A, Aurelian SM, et al. Post intervention outcomes of physical exercise training in elderly with mild to moderate dementia. *Eur J Neurol.* 2019;26:113-. doi: 10.1111/ene.14018. PMID: CN-01979194.
287. Casper GR, Calvitti A, Brennan PF, et al. ComputerLink: the impact of a computer network on Alzheimer's caregivers' decision-making confidence and skill. *Medinfo.* 1995;8 Pt 2:1546. PMID: 8591497.
288. Chan WC, Lautenschlager N, Dow B, et al. A home-based exercise intervention for caregivers of persons with dementia: Study protocol for a randomised controlled trial. *Trials.* 2016 21 Sep;17(1)(no pagination)(460). doi: <http://dx.doi.org/10.1186/s13063-016-1582-z>. PMID: 612244535.
289. Clark ME, Lipe AW, Bilbrey M. Use of music to decrease aggressive behaviors in people with dementia. *J Gerontol Nurs.* 1998 Jul;24(7):10-7. PMID: 9801526.
290. Collier. The use of multi-sensory stimulation to improve functional performance in older people with dementia: a randomised single blind trial. In *British and Irish theses* [<http://ezproxyulsoxacuk:2514/idx/scripts/itasp?Xml=f:\index\idx\docs\all\56\it00525859htm&subfolder=/search>]
291. Crawford SE, Leaver VW, Mahoney SD. Using Reiki to decrease memory and behavior problems in mild cognitive impairment and mild Alzheimer's disease. *J Altern Complement Med.* 2006 Nov;12(9):911-3. PMID: 17109583.
292. Curto PD, Sole SC, Castro M, et al. Effects of music therapy in institutionalized elderly with dementia. In *European geriatric medicine*
293. Dam AEH, de Vugt ME, van Boxtel MPJ, et al. Effectiveness of an online social support intervention for caregivers of people with dementia: The study protocol of a randomised controlled trial. *Trials.* 2017 29 Aug;18(1)(no pagination)(395). doi: <http://dx.doi.org/10.1186/s13063-017-2097-y>. PMID: 618042762.
294. DeGregory C. The effects of multiple gratitude interventions among informal caregivers of persons with dementia and Alzheimer's disease.
295. Eggermont LH, Blankevoort CG, Scherder EJ. Walking and night-time restlessness in mild-to-moderate dementia: a randomized controlled trial. *Age Ageing.* 2010 Nov;39(6):746-9. doi: <https://dx.doi.org/10.1093/ageing/afq115>. PMID: 20823127.
296. Finnegan S, Nichols V, Sheehan B, et al. A randomised controlled trial of moderate to high intensity exercise training for people with dementia: dementia And Physical Activity (DAPA)...The Chartered Society of Physiotherapy UK Conference 2018, Birmingham, UK, 19-20 October 2018. *Physiotherapy.* 2019;105:e104-e. PMID: CN-01690752.
297. Fleiner T, Zijlstra W, Dauth H, et al. Evaluation of a hospital-based day-structuring exercise programme on exacerbated behavioural and psychological symptoms in dementia - the exercise carousel: Study protocol for a randomised controlled trial. *Trials.* 2015 May 26;16(1)(no pagination)(228). doi: <http://dx.doi.org/10.1186/s13063-015-0758-2>. PMID: 604800524.
298. Fortinsky RH, Gitlin LN, Pizzi LT, et al. Translation of the Care of Persons with Dementia in their Environments (COPE) intervention in a publicly-funded home care context: Rationale and research design. *Contemp Clin Trials.* 2016 07;49:155-65. doi: <https://dx.doi.org/10.1016/j.cct.2016.07.006>. PMID: 27394383.

299. Fujii M, Hatakeyama R, Fukuoka Y, et al. Lavender aroma therapy for behavioral and psychological symptoms in dementia patients. *Geriatr Gerontol Int*. 2008 Jun;8(2):136-8. doi: <https://dx.doi.org/10.1111/j.1447-0594.2008.00461.x>. PMID: 18713168.
300. Gerlach LB, Kales HC. Learning their language: the importance of detecting and managing pain in dementia.
301. Goeman D, Comans T, Enticott JC, et al. Evaluating the Efficacy of the "Support for Life" Program for People with Dementia and Their Families and Carers' to Enable Them to Live Well: A Protocol for a Cluster Stepped Wedge Randomized Controlled Trial. *Front*. 2016;4:245. PMID: 27843893.
302. Griffiths AW, Surr CA, Creese B, et al. The development and use of the assessment of dementia awareness and person-centred care training tool in long-term care. *Dementia*. 2018 Jan 01:1471301218768165. doi: <https://dx.doi.org/10.1177/1471301218768165>. PMID: 29631493.
303. Hamilton G, Gallagher M, Pierson K, et al. Palliative care for dementia: a randomized controlled study of a replicable and financially viable model for organizations providing caregiver support. In *Journal of pain and symptom management*
304. Hatakeyama R, Fukushima K, Fukuoka Y, et al. Personal home made digital video disk for patients with behavioral psychological symptoms of dementia. *Geriatr Gerontol Int*. 2010 Jul;10(3):272-4. doi: <https://dx.doi.org/10.1111/j.1447-0594.2010.00623.x>. PMID: 20629763.
305. Hayden LJ, Glynn SM, Hahn TJ, et al. The use of Internet technology for psychoeducation and support with dementia caregivers. *Psychol Serv*. 2012 May;9(2):215-8. doi: <https://dx.doi.org/10.1037/a0027056>. PMID: 22662739.
306. Heinrich S, Berwig M, Simon A, et al. German adaptation of the Resources for Enhancing Alzheimer's Caregiver Health II: study protocol of a single-centred, randomised controlled trial. *BMC geriatr*. 2014 Feb 12;14:21. doi: <https://dx.doi.org/10.1186/1471-2318-14-21>. PMID: 24520910.
307. Hirazakura A, Hatakeyama R, Fukuoka Y, et al. Emotional therapy for patients with dementia. *Geriatr Gerontol Int*. 2008 Dec;8(4):303-6. doi: <https://dx.doi.org/10.1111/j.1447-0594.2008.00486.x>. PMID: 19149844.
308. Ho RT, Cheung JK, Chan WC, et al. A 3-arm randomized controlled trial on the effects of dance movement intervention and exercises on elderly with early dementia. *BMC geriatr*. 2015 Oct 19;15:127. doi: <https://dx.doi.org/10.1186/s12877-015-0123-z>. PMID: 26481870.
309. Hokkanen L, Rantala L, Remes AM, et al. Dance and movement therapeutic methods in management of dementia: a randomized, controlled study. *J Am Geriatr Soc*. 2008 Apr;56(4):771-2. doi: <https://dx.doi.org/10.1111/j.1532-5415.2008.01611.x>. PMID: 18380687.
310. Holle R, Grassel E, Ruckdaschel S, et al. Dementia care initiative in primary practice: study protocol of a cluster randomized trial on dementia management in a general practice setting. *BMC Health Serv Res*. 2009 Jun 06;9:91. doi: <https://dx.doi.org/10.1186/1472-6963-9-91>. PMID: 19500383.
311. Hooghiemstra AM, Eggermont LH, Scheltens P, et al. Study protocol: EXERcise and cognition in sedentary adults with early-ONset dementia (EXERCISE-ON). *BMC Neurol*. 2012 Aug 16;12:75. doi: <https://dx.doi.org/10.1186/1471-2377-12-75>. PMID: 22897903.
312. Huis In Het Veld JG, Willemse BM, van Asch IF, et al. Online Self-Management Support for Family Caregivers to Help Them Manage Behavior Changes in Their Relative With Dementia: Study Protocol for a Randomized Controlled Trial and a Process Evaluation. *JMIR Res Protoc*. 2017 Nov 28;6(11):e239. doi: <https://dx.doi.org/10.2196/resprot.8365>. PMID: 29183870.
313. Huttemann D. Alzheimer's treatment: bearer of hope fails in phase III. In *Pharmazeutische zeitung*
314. Jablonski RA, Kolanowski A, Therrien B, et al. Reducing care-resistant behaviors during oral hygiene in persons with dementia. *BMC Oral Health*. 2011 Nov 19;11:30. doi: <https://dx.doi.org/10.1186/1472-6831-11-30>. PMID: 22100010.
315. Jablonski-Jaudon RA, Kolanowski AM, Winstead V, et al. Maturation of the MOUTH Intervention: From Reducing Threat to Relationship-Centered Care. *J Gerontol Nurs*. 2016 Mar;42(3):15-23; quiz 4-5. doi: <https://dx.doi.org/10.3928/00989134-20160212-05>. PMID: 26934969.
316. Jutkowitz E, Pizzi LT, Popp J, et al. PND38 EVALUATION OF DEMENTIA INFORMAL CAREGIVERS WILLINGNESS TO PAY FOR A NONPHARMACOLOGIC INTERVENTION TO REDUCE BEHAVIORAL SYMPTOMS. *Value Health*. 2019;22:S277-. doi: 10.1016/j.jval.2019.04.1311. PMID: CN-01963978.
317. Kikutani T, Yoneyama T, Nishiwaki K, et al. Effect of oral care on cognitive function in patients with dementia. *Geriatr Gerontol Int*. 2010 Oct;10(4):327-8. doi: <https://dx.doi.org/10.1111/j.1447-0594.2010.00637.x>. PMID: 20887627.
318. Kohler L, Meinke-Franze C, Hein J, et al. Does an interdisciplinary network improve dementia care? Results from the IDemUck-study. *Curr Alzheimer Res*. 2014;11(6):538-48. PMID: 24938504.
319. Korn L, Logsdon R, Polissar N, et al. P02.84. A randomized trial of Polarity therapy for stress and pain reduction in American Indian and Alaska Native family dementia caregivers. *BMC Altern Med*. 2012;12(Suppl 1):1-. doi: 10.1186/1472-6882-12-S1-P140. PMID: 104481508. Language: English. Entry Date: 20121129. Revision Date: 20150711. Publication Type: Journal Article.
320. Kudoh H, Hatakeyama A, Yamamoto T, et al. Foot care using green tea paste for behavioral and psychological symptoms in dementia patients. *Geriatr Gerontol Int*. 2009 Dec;9(4):399-401. doi: <https://dx.doi.org/10.1111/j.1447-0594.2009.00545.x>. PMID: 20002761.
321. Kuo LM, Huang HL, Liang J, et al. Trajectories of health-related quality of life among family caregivers of individuals with dementia: A home-based caregiver-training program matters. *Geriatr Nurs*. 2017 Mar - Apr;38(2):124-32. doi:

- <https://dx.doi.org/10.1016/j.gerinurse.2016.08.017>. PMID: 27720499.
322. Kwak J, Valuch K, Brondino M. Randomized crossover study of music & memory intervention with nursing home residents with dementia. In *Gerontologist* 323. Kwok T, Lam L, Chung J. Case management to improve quality of life of older people with early dementia and to reduce caregiver burden. *Hong Kong Med*. 2012 Dec;18 Suppl 6:4-6. PMID: 23249844.
324. Lai CK, Lai DL, Ho JS, et al. Interdisciplinary collaboration in the use of a music-with-movement intervention to promote the wellbeing of people with dementia and their families: Development of an evidence-based intervention protocol. *Nurs Health Sci*. 2016 Mar;18(1):79-84. doi: <https://dx.doi.org/10.1111/nhs.12238>. PMID: 26354593.
325. Lai CKY, Wong LF, Liu K-H, et al. Online and onsite training for family caregivers of people with dementia: results from a pilot study. In *Int J Geriatr Psychiatry* 326. Lauterberg J, Grossfeld-Schmitz M, Ruckdäschel S, et al. IDA project—concept and implementation of a cluster-randomized trial on dementia management in a general practice setting. In *Zeitschrift für ärztliche fortbildung und qualitätssicherung* 327. Li J, Chang Y, Jungquist C, et al. Sleep in long-term care residents with dementia: pilot of a person-centered care intervention. In *Sleep* 328. Liu W, Wang P, Gray H, et al. Client satisfaction with a stress reduction program for Chinese dementia caregivers. *Hallym International Journal of Aging*. 2008;10(2):91-110. doi: 10.2190/HA.10.2.c. PMID: 105203686. Language: English. Entry Date: 20100625. Revision Date: 20150711. Publication Type: Journal Article.
329. Logsdon RG. Dementia: psychosocial interventions for family caregivers. *Lancet*. 2008 Jul 19;372(9634):182-3. doi: [https://dx.doi.org/10.1016/S0140-6736\(08\)61048-X](https://dx.doi.org/10.1016/S0140-6736(08)61048-X). PMID: 18640438.
330. Logsdon RG, McCurry SM, Teri L. A home health care approach to exercise for persons with Alzheimer's disease. *Care Manag J*. 2005;6(2):90-7. PMID: 16544870.
331. Luk KY, Lai KY, Li CC, et al. The effect of horticultural activities on agitation in nursing home residents with dementia. *Int J Geriatr Psychiatry*. 2011 Apr;26(4):435-6. doi: <https://dx.doi.org/10.1002/gps.2493>. PMID: 21412848.
332. Mahoney DF, Tarlow B, Sandaire J. A computer-mediated intervention for Alzheimer's caregivers. *Comput Nurs*. 1998 Jul-Aug;16(4):208-16; quiz 17-8. PMID: 9675988.
333. Mahoney EK, Trudeau SA, Penyack SE, et al. Challenges to intervention implementation: lessons learned in the Bathing Persons with Alzheimer's Disease at Home study. *Nurs Res*. 2006 Mar-Apr;55(2 Suppl):S10-6. PMID: 16601628.
334. Mariko A, Matsuda H, Takahashi M, et al. Touch on the acupoint of shinchuu of Alzheimer's disease patients. In *Geriatr Gerontol Int* 335. Miskelly F. Electronic tracking of patients with dementia and wandering using mobile phone technology. *Age Ageing*. 2005 Sep;34(5):497-9. PMID: 16107453.
336. Moore J. Familiar physical activity to familiar music: the effects on apathy, agitation, eating ability, and dietary intake in institutionalized older adults with dementia.
337. Nguyen MF, Bonnefoy M, Adrait A, et al. Efficacy of Hearing Aids on the Cognitive Status of Patients with Alzheimer's Disease and Hearing Loss: A Multicenter Controlled Randomized Trial. *J Alzheimers Dis*. 2017;58(1):123-37. doi: <https://dx.doi.org/10.3233/JAD-160793>. PMID: 28387664.
338. Ochmann S, Kasper E, Hoffmann W, et al. Feasibility of a cognitive rehabilitation group program for patients with mild dementia in Alzheimer's disease: a randomized, controlled, single blinded pilot study. In *Alzheimer's dement* 339. Panca M. PMH15 COST-EFFECTIVENESS OF THE MANAGING AGITATION AND RAISING QUALITY OF LIFE IN DEMENTIA (MARQUE) INTERVENTION FOR AGITATION SYMPTOMS IN PEOPLE WITH DEMENTIA IN CARE HOMES. *Value Health*. 2019;22:S228-. doi: 10.1016/j.jval.2019.04.1060. PMID: CN-01963980.
340. Parra-Vidales E, Soto-Perez F, Perea-Bartolome MV, et al. Online interventions for caregivers of people with dementia: a systematic review. *Actas Esp Psiquiatr*. 2017 May;45(3):116-26. PMID: 28594057.
341. Poon P, Hui E, Dai D, et al. Cognitive intervention for community-dwelling older persons with memory problems: telemedicine versus face-to-face treatment. *Int J Geriatr Psychiatry*. 2005 Mar;20(3):285-6. PMID: 15717335.
342. Poptsi E, Kounti F, Samakouri M, et al. Longitudinal cognitive training program in people with mild Alzheimer's disease: A Randomized Controlled Trial. *Hellenic Journal of Nuclear Medicine*. 2017 May-August;20(Supplement 1):233-43. PMID: 621578557.
343. Rollin-Sillaire A, Breuilh L, Salleron J, et al. Reasons that prevent the inclusion of Alzheimer's disease patients in clinical trials. *Br J Clin Pharmacol*. 2013 Apr;75(4):1089-97. doi: <https://dx.doi.org/10.1111/j.1365-2125.2012.04423.x>. PMID: 22891847.
344. Sarkamo T, Laitinen S, Numminen A, et al. Pattern of Emotional Benefits Induced by Regular Singing and Music Listening in Dementia. *J Am Geriatr Soc*. 2016 Feb;64(2):439-40. doi: <https://dx.doi.org/10.1111/jgs.13963>. PMID: 26889849.
345. Satoh M, Ogawa J, Tokita T, et al. Physical exercise with music maintains activities of daily living in patients with dementia: mihama-kiho project part 2. In *Journal of alzheimers disease* 346. Schmid AA, Spangler-Morris C, Beauchamp RC, et al. The Home-Based Occupational Therapy Intervention in the Alzheimer's Disease Multiple Intervention Trial (ADMIT). *Occup Ther Ment Health*. 2015 Jan 01;31(1):19-34. PMID: 26997685.
347. Schulz R, Belle SH, Czaja SJ, et al. Introduction to the special section on Resources for Enhancing Alzheimer's Caregiver Health (REACH). *Psychol Aging*. 2003 Sep;18(3):357-60. PMID: 14518799.
348. Sithinamsuwan P, Saengwanitch S, Pinidbunjerdkool A, et al. The effect of Thai traditional music on cognitive function, psychological health and quality of sleep among older Thai individuals with dementia. In *J Am Geriatr Soc*

349. Slomski A. Short therapy reduces depression in family caregivers of patients with dementia. *Jama*. 2015 Jan 27;313(4):347. doi: <https://dx.doi.org/10.1001/jama.2014.17991>. PMID: 25626027.
350. Snowdon J. Tailored treatment for depression with co-morbid dementia. In *Australian and new zealand journal of psychiatry*
351. Soto M, Abushakra S, Cummings J, et al. Progress in Treatment Development for Neuropsychiatric Symptoms in Alzheimer's Disease: Focus on Agitation and Aggression. A Report from the EU/US/CTAD Task Force. *Jpad*. 2015 Sep;2(3):184-8. PMID: 26413494.
352. Spector A, Woods B, Orrell M. Cognitive stimulation for the treatment of Alzheimer's disease. In *Expert rev*
353. Tanner J, Student M, Black B, et al. Effectiveness of a multicomponent care coordination intervention on dementia caregivers in the community-a randomized control trial Jeremy A. Tanner. In *Am J Geriatr Psychiatry*
354. Thompson CA, Spilsbury K, Hall J, et al. Systematic review of information and support interventions for caregivers of people with dementia. *BMC geriatr*. 2007;7 (no pagination)(18). doi: <http://dx.doi.org/10.1186/1471-2318-7-18>. PMID: 47329461.
355. Thornley J, Hirjee H, Vasudev A. Music therapy in patients with dementia and behavioral disturbance on an inpatient psychiatry unit: results from a pilot randomized controlled study. *Int Psychogeriatr*. 2016 May;28(5):869-71. doi: <https://dx.doi.org/10.1017/S1041610215001866>. PMID: 26572722.
356. Tibaldi V, Aimonino N, Ponzetto M, et al. A randomized controlled trial of a home hospital intervention for frail elderly demented patients: behavioral disturbances and caregiver's stress. *Arch Gerontol Geriatr Suppl*. 2004(9):431-6. PMID: 15207444.
357. Vernooij-Dassen M, Huygen F, Felling A, et al. Home care for dementia patients. *J Am Geriatr Soc*. 1995 Apr;43(4):456-7. PMID: 7706643.
358. Viana J, Vickers J, Cook M, et al. Currents of memory: recent progress, translational challenges, and ethical considerations in fornix deep brain stimulation trials for Alzheimer's disease. In *Neurobiology of aging* (no pagination), 2017
359. Vink A, Zuidersma M, Boersma F, et al. Effect of music therapy versus recreational activities on neuropsychiatric symptoms in elderly adults with dementia: an exploratory randomized controlled trial. In *J Am Geriatr Soc*
360. Voigt-Radloff S, Hüll M. Practice example: neurodegenerative disorders: randomised controlled multicentre trial of home-based occupational therapy for patients with dementia. In *Zeitschrift für evidenz, fortbildung und qualität IM gesundheitswesen*
361. Wells D, Dawson P, Sidani S, et al. The benefits of abilities-focused morning care for residents with dementia and their caregivers. *Perspectives*. 2000;24(1):17. PMID: 12026330.
362. Willemse BM, Smit D, de Lange J, et al. Nursing home care for people with dementia and residents' quality of life, quality of care and staff well-being: design of the Living Arrangements for people with Dementia (LAD)-study. *BMC geriatr*. 2011 Mar 17;11:11. doi: <https://dx.doi.org/10.1186/1471-2318-11-11>. PMID: 21414185.
363. Williams V, Bishop-Fitzpatrick L, Lane J, et al. "Video-based coping skills to reduce health risk and improve psychological and physical well-being in Alzheimer's disease family caregivers": erratum. In *Psychosom Med*
364. Woods B. Promoting well-being and independence for people with dementia. *Int J Geriatr Psychiatry*. 1999 Feb;14(2):97-105; discussion -9. PMID: 10885861.
365. Yoo R, Yeom J, Kim GH, et al. Erratum to: A Multicenter, Randomized Clinical Trial to Assess the Efficacy of a Therapeutic Intervention Program for Caregivers of People with Dementia. *J Clin Neurol*. 2019 Jul;15(3):428. doi: <https://dx.doi.org/10.3988/jcn.2019.15.3.428>. PMID: 31286725.

Ineligible study design

366. The tailored activity program (TAP) to address behavioral disturbances in frontotemporal dementia: a feasibility and pilot study. *Disability and rehabilitation*. 2019;41(3):299-310. doi: <https://doi.org/10.1080/09638288.2017.1387614>. PMID: CN-01966803.
367. Ahn N, Kim K. Effects of an elastic band resistance exercise program on lower extremity muscle strength and gait ability in patients with Alzheimer's disease. In *J Phys Ther Sci*
368. Alcala-Lozano R, Morelos-Santana E, Cortes-Sotres JF, et al. Similar clinical improvement and maintenance after rTMS at 5 Hz using a simple vs. complex protocol in Alzheimer's disease. *Brain Stimul*. 2018 May - Jun;11(3):625-7. doi: <https://dx.doi.org/10.1016/j.brs.2017.12.011>. PMID: 29326021.
369. Anderson K, Bird M, Macpherson S, et al. Findings from a pilot investigation of the effectiveness of a snoezelen room in residential care: should we be engaging with our residents more? *Geriatr Nurs*. 2011 May-Jun;32(3):166-77. doi: <https://dx.doi.org/10.1016/j.gerinurse.2010.12.011>. PMID: 21306798.
370. Arcoverde C, Deslandes A, Rangel A, et al. Role of physical activity on the maintenance of cognition and activities of daily living in elderly with Alzheimer's disease: papel da atividade física na manutenção da cognição e atividades de vida diária em idosos com doença de Alzheimer. In *Arq Neuropsiquiatr*
371. Bach D, Bach M, Bohmer F, et al. Reactivating occupational therapy: a method to improve cognitive performance in geriatric patients. *Age Ageing*. 1995 May;24(3):222-6. PMID: 7645443.
372. Ballard C, Brown R, Fossey J, et al. Brief psychosocial therapy for the treatment of agitation in Alzheimer disease (the CALM-AD trial). *Am J Geriatr Psychiatry*. 2009 Sep;17(9):726-33. doi: <https://dx.doi.org/10.1097/JGP.0b013e3181b0f8c0>. PMID: 19700946.
373. Barnes DE, Mehling W, Wu E, et al. Preventing loss of independence through exercise (PLIE): a pilot clinical trial in older adults with dementia. *PLoS ONE*.

- 2015;10(2):e0113367. doi:
<https://dx.doi.org/10.1371/journal.pone.0113367>. PMID:
 25671576.
374. Bass DM, Hornick T, Kunik M, et al. Findings From a Real-World Translation Study of the Evidence-Based "Partners in Dementia Care". *Innov Aging*. 2019 Jul;3(3):igz031. doi:
<https://dx.doi.org/10.1093/geroni/igz031>. PMID:
 31660442.
375. Boulay M, Benveniste S, Boespflug S, et al. A pilot usability study of MINWii, a music therapy game for demented patients. *Technology & Health Care*. 2011;19(1):233-46. PMID: 104577441. Language: English. Entry Date: 20120323. Revision Date: 20150711. Publication Type: Journal Article.
376. Brooker DJ, Snape M, Johnson E, et al. Single case evaluation of the effects of aromatherapy and massage on disturbed behaviour in severe dementia. *Br J Clin Psychol*. 1997 May;36(Pt 2):287-96. PMID: 9167869.
377. Cadore E, Moneo A, Mensat M, et al. Positive effects of resistance training in frail elderly patients with dementia after long-term physical restraint. In *Age (Dordr)*
378. Callahan CM, Boustani MA, Weiner M, et al. Implementing dementia care models in primary care settings: The Aging Brain Care Medical Home. *Aging Ment Health*. 2011 Jan;15(1):5-12. doi:
<https://dx.doi.org/10.1080/13607861003801052>. PMID:
 20945236.
379. Carmody J, Potter J, Lewis K, et al. Development and pilot testing of a decision aid for drivers with dementia. *BMC Med Inf Decis Mak*. 2014 Mar 19;14:19. doi:
<https://dx.doi.org/10.1186/1472-6947-14-19>. PMID:
 24642051.
380. Chao HC, Kaas M, Su YH, et al. Effects of the Advanced Innovative Internet-Based Communication Education Program on Promoting Communication Between Nurses and Patients With Dementia. *J Nurs Res*. 2016 Jun;24(2):163-72. doi:
<https://dx.doi.org/10.1097/jnr.000000000000109>. PMID:
 26376065.
381. Chenoweth L, Jeon Y-H. Determining the efficacy of dementia care mapping as an outcome measure and a process for change: a pilot study. In *Aging Ment Health*
382. Chew J, Chong MS, Fong YL, et al. Outcomes of a multimodal cognitive and physical rehabilitation program for persons with mild dementia and their caregivers: a goal-oriented approach. *Clin Interv Aging*. 2015;10:1687-94. doi: <https://dx.doi.org/10.2147/CIA.S93914>. PMID: 26543358.
383. Chew J, Chong M-S, Fong Y-L, et al. Outcomes of a multimodal cognitive and physical rehabilitation program for persons with mild dementia and their caregivers: a goal-oriented approach. In *Clin Interv Aging*
384. Churchill M, Safaoui J, McCabe BW, et al. Using a therapy dog to alleviate the agitation and desocialization of people with Alzheimer's disease. *J Psychosoc Nurs Ment Health Serv*. 1999 Apr;37(4):16-22. PMID: 10218187.
385. Clarke A, Chan J, Stott J, et al. An adapted mindfulness intervention for people with dementia in care homes: feasibility pilot study. In *Int J Geriatr Psychiatry*
386. Cohen-Mansfield J, Marx MS, Dakheel-Ali M, et al. Can agitated behavior of nursing home residents with dementia be prevented with the use of standardized stimuli? *J Am Geriatr Soc*. 2010 Aug;58(8):1459-64. doi:
<https://dx.doi.org/10.1111/j.1532-5415.2010.02951.x>. PMID: 20579167.
387. Cohen-Mansfield J, Thein K, Marx MS, et al. The relationships of environment and personal characteristics to agitated behaviors in nursing home residents with dementia. *J Clin Psychiatry*. 2012 Mar;73(3):392-9. doi:
<https://dx.doi.org/10.4088/JCP.10m06605>. PMID:
 22152398.
388. Corbeil RR, Quayhagen MP, Quayhagen M. Intervention effects on dementia caregiving interaction: a stress-adaptation modeling approach. *J Aging Health*. 1999 Feb;11(1):79-95. PMID: 10848143.
389. Cotelli M, Calabria M, Manenti R, et al. Improved language performance in Alzheimer disease following brain stimulation. *J Neurol Neurosurg Psychiatry*. 2011 Jul;82(7):794-7. doi:
<https://dx.doi.org/10.1136/jnnp.2009.197848>. PMID:
 20574108.
390. Cott CA, Dawson P, Sidani S, et al. The effects of a walking/talking program on communication, ambulation, and functional status in residents with Alzheimer disease. *Alzheimer Dis Assoc Disord*. 2002 Apr-Jun;16(2):81-7. PMID: 12040303.
391. Custodio N, Herrera-Perez E, Lira D, et al. Evaluation of the INECO Frontal Screening and the Frontal Assessment Battery in Peruvian patients with Alzheimer's disease and behavioral variant Frontotemporal dementia. In *Eneurologicalsci*
392. Czaja SJ, Gitlin LN, Schulz R, et al. Development of the risk appraisal measure: a brief screen to identify risk areas and guide interventions for dementia caregivers. *J Am Geriatr Soc*. 2009 Jun;57(6):1064-72. doi:
<https://dx.doi.org/10.1111/j.1532-5415.2009.02260.x>. PMID: 19453305.
393. Dabelko-Schoeny H, Phillips G, Darrough E, et al. Equine-assisted intervention for people with dementia. In *Anthrozoos*
394. Davison TE, Nayer K, Coxon S, et al. A personalized multimedia device to treat agitated behavior and improve mood in people with dementia: A pilot study. *Geriatr Nurs*. 2016 Jan-Feb;37(1):25-9. doi:
<https://dx.doi.org/10.1016/j.gerinurse.2015.08.013>. PMID:
 26412509.
395. de Oliveira AM, Radanovic M, Homem de Mello PC, et al. An intervention to reduce neuropsychiatric symptoms and caregiver burden in dementia: Preliminary results from a randomized trial of the tailored activity program-outpatient version. *Int J Geriatr Psychiatry*. 2019 Sep;34(9):1301-7. doi: <https://dx.doi.org/10.1002/gps.4958>. PMID: 30035341.
396. de Oliveira Silva F, Ferreira JV, Placido J, et al. Three months of multimodal training contributes to mobility and executive function in elderly individuals with mild cognitive impairment, but not in those with Alzheimer's disease: A randomized controlled trial. *Maturitas*. 2019 Aug;126:28-33. doi:
<https://dx.doi.org/10.1016/j.maturitas.2019.04.217>. PMID:
 31239114.
397. de Souto Barreto P, Cadroy Y, Kelaiditi E, et al. The prognostic value of body-mass index on mortality in older

- adults with dementia living in nursing homes. *Clin Nutr*. 2017 Apr;36(2):423-8. doi: <https://dx.doi.org/10.1016/j.clnu.2015.12.009>. PMID: 26724185.
398. Deal JA, Betz J, Yaffe K, et al. Hearing Impairment and Incident Dementia and Cognitive Decline in Older Adults: The Health ABC Study. *J Gerontol A Biol Sci Med Sci*. 2017 May 01;72(5):703-9. doi: <https://dx.doi.org/10.1093/gerona/glw069>. PMID: 27071780.
399. Deponte A, Missan R. Effectiveness of validation therapy (VT) in group: preliminary results. *Arch Gerontol Geriatr*. 2007 Mar-Apr;44(2):113-7. PMID: 16712981.
400. Der-Fa L, Hart LK, Lutgendorf SK, et al. Slowing progression of early stages of AD with alternative therapies: A feasibility study. *Geriatr Nurs*. 2013;34(6):457-64. doi: 10.1016/j.gerinurse.2013.07.003. PMID: 107924864. Language: English. Entry Date: 20131224. Revision Date: 20150818. Publication Type: Journal Article.
401. Drummond MF, Mohide EA, Tew M, et al. Economic evaluation of a support program for caregivers of demented elderly. *Int J Technol Assess Health Care*. 1991;7(2):209-19. PMID: 1907599.
402. Duggleby W, Ploeg J, McAiney C, et al. A Comparison of Users and Nonusers of a Web-Based Intervention for Carers of Older Persons With Alzheimer Disease and Related Dementias: Mixed Methods Secondary Analysis. *J Med Internet Res*. 2019 Oct 17;21(10):e14254. doi: <https://dx.doi.org/10.2196/14254>. PMID: 31625947.
403. Duggleby W, Ploeg J, McAiney C, et al. Web-Based Intervention for Family Carers of Persons with Dementia and Multiple Chronic Conditions (My Tools 4 Care): Pragmatic Randomized Controlled Trial. *J Med Internet Res*. 2018 Jun 29;20(6):e10484. doi: <https://dx.doi.org/10.2196/10484>. PMID: 29959111.
404. Dunne TE, Nearing SA, Cipolloni PB, et al. Visual contrast enhances food and liquid intake in advanced Alzheimer's disease. *Clin Nutr*. 2004 Aug;23(4):533-8. PMID: 15297089.
405. Edvardsson D, Sandman PO, Borell L. Implementing national guidelines for person-centered care of people with dementia in residential aged care: effects on perceived person-centeredness, staff strain, and stress of conscience. *Int Psychogeriatr*. 2014 Jul;26(7):1171-9. doi: <https://dx.doi.org/10.1017/S1041610214000258>. PMID: 24576607.
406. Edwards N, Gardiner M, Ritchie D, et al. Effect of exercise on negative affect in residents in special care units with moderate to severe dementia. In *Alzheimer disease and associated disorders*
407. Ferrucci R, Mrakic-Spota S, Gardini S, et al. Behavioral and Neurophysiological Effects of Transcranial Direct Current Stimulation (tDCS) in Fronto-Temporal Dementia. *Front Behav Neurosci*. 2018;12:235. doi: <https://dx.doi.org/10.3389/fnbeh.2018.00235>. PMID: 30420799.
408. Forester BP, Mellen E, Mathias L, et al. Electroconvulsive therapy for the treatment of acute agitation and aggression in alzheimer's dementia (ect-ad). *Am J Geriatr Psychiatry*. 2019;27 (3 Supplement):S125-S6. PMID: CN-01911574.
409. Fusar-Poli L, Bieleninik L, Brondino N, et al. The effect of music therapy on cognitive functions in patients with dementia: a systematic review and meta-analysis. *Aging Ment Health*. 2017 Jul 10;1-10. doi: <https://dx.doi.org/10.1080/13607863.2017.1348474>. PMID: 28691506.
410. Galante E, Venturini G, Fiaccadori C. Computer-based cognitive intervention for dementia: preliminary results of a randomized clinical trial. *G Ital Med Lav Ergon*. 2007 Jul-Sep;29(3 Suppl B):B26-32. PMID: 18575355.
411. Garcia-Alberca JM. Cognitive-behavioral treatment for depressed patients with Alzheimer's disease. An open trial. *Arch Gerontol Geriatr*. 2017 Jul;71:1-8. doi: <https://dx.doi.org/10.1016/j.archger.2017.02.008>. PMID: 28237746.
412. George DR, George DR. Intergenerational volunteering and quality of life: mixed methods evaluation of a randomized control trial involving persons with mild to moderate dementia. *Qual Life Res*. 2011;20(7):987-95. doi: 10.1007/s11136-010-9837-8. PMID: 104578344. Language: English. Entry Date: 20120323. Revision Date: 20171109. Publication Type: journal article.
413. George DR, Whitehouse PJ. Intergenerational volunteering and quality of life for persons with mild-to-moderate dementia: results from a 5-month intervention study in the United States. *J Am Geriatr Soc*. 2010;58(4):796-7. doi: 10.1111/j.1532-5415.2010.02790.x. PMID: 105169520. Language: English. Entry Date: 20100521. Revision Date: 20150711. Publication Type: Journal Article.
414. Germain S, Wojtasik V, Lekeu F, et al. Efficacy of Cognitive Rehabilitation in Alzheimer Disease: A 1-Year Follow-Up Study. *J Geriatr Psychiatry Neurol*. 2019 Jan;32(1):16-23. doi: <https://dx.doi.org/10.1177/0891988718813724>. PMID: 30477375.
415. Glueckauf RL, Davis WS, Willis F, et al. Telephone-based, cognitive-behavioral therapy for African American dementia caregivers with depression: initial findings. *Rehabil Psychol*. 2012 May;57(2):124-39. doi: <https://dx.doi.org/10.1037/a0028688>. PMID: 22686551.
416. Goyder J, Orrell M, Wenborn J, et al. Staff training using STAR: a pilot study in UK care homes. *Int Psychogeriatr*. 2012 Jun;24(6):911-20. doi: <https://dx.doi.org/10.1017/S1041610211002559>. PMID: 22217445.
417. Gray SG, Clair AA. Influence of aromatherapy on medication administration to residential-care residents with dementia and behavioral challenges. *Am J Alzheimers Dis Other Demen*. 2002 May-Jun;17(3):169-74. PMID: 12083347.
418. Greer KL, Pustay KA, Zaun TC, et al. A comparison of the effects of toys versus live animals on the communication of patients with dementia of the Alzheimer's type. *Clinical Gerontologist: The Journal of Aging and Mental Health*. 2001;24(3-4):157-82. doi: http://dx.doi.org/10.1300/J018v24n03_13. PMID: 2002-18267-012.
419. Guo Y, Shi X, Uchiyama H, et al. A study on the rehabilitation of cognitive function and short-term memory in patients with Alzheimer's disease using transcutaneous

- electrical nerve stimulation. *Front Med Biol Eng*. 2002;11(4):237-47. PMID: 12735425.
420. Guzman A, Wenborn J, Swinson T, et al. Evaluation of the 'Ladder to the Moon, Culture Change Studio Engagement Programme' staff training: Two quasi-experimental case studies. *Int J Older People Nurs*. 2017 Sep;12(3). doi: <https://dx.doi.org/10.1111/opn.12147>. PMID: 28185441.
421. Haffmans PM, Sival RC, Lucius SA, et al. Bright light therapy and melatonin in motor restless behaviour in dementia: a placebo-controlled study. *Int J Geriatr Psychiatry*. 2001 Jan;16(1):106-10. PMID: 11180494.
422. Harris JB. The impact of physical versus social activity on the physical and cognitive functioning of seniors with dementia.
423. Hassan S, Aguirre E, Betz A, et al. Evaluating the effect of Brainfood groups for people with mild cognitive impairment and mild dementia: Preliminary mixed-methodology study. *BJPsych Open*. 2018 01 Jul;4(4):208-14. doi: <http://dx.doi.org/10.1192/bjo.2018.29>. PMID: 624229678.
424. Haupt M, Karger A, Jänner M. Improvement of agitation and anxiety in demented patients after psychoeducative group intervention with their caregivers. *Int J Geriatr Psychiatry*. 2000;15(12):1125-9. doi: 10.1002/1099-1166(200012)15:12<1125::AID-GPS257>3.0.CO;2-F. PMID: 104723571. Language: English. Entry Date: 20110610. Revision Date: 20150711. Publication Type: Journal Article.
425. Heyn P, Abreu BC, Ottenbacher KJ. The effects of exercise training on elderly persons with cognitive impairment and dementia: a meta-analysis. *Arch Phys Med Rehabil*. 2004 Oct;85(10):1694-704. PMID: 15468033.
426. Hicks-Moore SL, Robinson BA. Favorite music and hand massage: Two interventions to decrease agitation in residents with dementia. *Dementia: The International Journal of Social Research and Practice*. 2008 Feb;7(1):95-108. doi: <http://dx.doi.org/10.1177/1471301207085369>. PMID: 2008-01605-006.
427. Hilgeman MM, Allen RS, Snow AL, et al. Preserving Identity and Planning for Advance Care (PIPAC): preliminary outcomes from a patient-centered intervention for individuals with mild dementia. *Aging Ment Health*. 2014 May;18(4):411-24. doi: <https://dx.doi.org/10.1080/13607863.2013.868403>. PMID: 24359036.
428. Hindle JV, Watermeyer TJ, Roberts J, et al. Goal-orientated cognitive rehabilitation for dementias associated with parkinson's disease-a pilot randomised controlled trial. *Int J Geriatr Psychiatry*. 2018;33(5):718-28. doi: <http://dx.doi.org/10.1002/gps.4845>. PMID: 623403793.
429. Holm A, Michel M, Stern GA, et al. The outcomes of an inpatient treatment program for geriatric patients with dementia and dysfunctional behaviors. *Gerontologist*. 1999;39(6):668-76. PMID: 107097417. Language: English. Entry Date: 20000301. Revision Date: 20150711. Publication Type: Journal Article.
430. Holm AK, Lepp M, Ringsberg KC. Dementia: involving patients in storytelling--a caring intervention. A pilot study. *J Clin Nurs*. 2005 Feb;14(2):256-63. PMID: 15669935.
431. Holmes C, Hopkins V, Hensford C, et al. Lavender oil as a treatment for agitated behaviour in severe dementia: a placebo controlled study. *Int J Geriatr Psychiatry*. 2002 Apr;17(4):305-8. PMID: 11994882.
432. Hooghiemstra AM, Eggermont LH, Scheltens P, et al. Exercise and early-onset Alzheimer's disease: theoretical considerations. *Dement Geriatr Cogn Dis Extra*. 2012 Jan;2:132-45. doi: <https://dx.doi.org/10.1159/000335493>. PMID: 22590474.
433. Hsu MH, Flowerdew R, Parker M, et al. Individual music therapy for managing neuropsychiatric symptoms for people with dementia and their carers: a cluster randomised controlled feasibility study. *BMC geriatr*. 2015 Jul 18;15:84. doi: <https://dx.doi.org/10.1186/s12877-015-0082-4>. PMID: 26183582.
434. Hwang HR, Choi SH, Yoon DH, et al. The effect of cognitive training in patients with mild cognitive impairment and early Alzheimer's disease: a preliminary study. *J Clin Neurol*. 2012 Sep;8(3):190-7. doi: <https://dx.doi.org/10.3988/jcn.2012.8.3.190>. PMID: 23091528.
435. Ibarria M, Alegret M, Valero S, et al. Beneficial Effects of an Integrated Psychostimulation Program in Patients with Alzheimer's Disease. *J Alzheimers Dis*. 2016;50(2):559-66. doi: <https://dx.doi.org/10.3233/JAD-150455>. PMID: 26757182.
436. Iliffe S, Robinson L, Bamford C, et al. Introducing case management for people with dementia in primary care: a mixed-methods study. *In Br J Gen Pract*
437. Iso-Markku P, Waller K, Kujala UM, et al. Physical activity and dementia: long-term follow-up study of adult twins. *Ann Med*. 2015 Mar;47(2):81-7. doi: <https://dx.doi.org/10.3109/07853890.2014.994675>. PMID: 25613168.
438. Jayalath D, Ashaye K, Kvavilashvili L. Carers' diaries in dementia: Is there a role in clinical practice? *Alzheimer's and Dementia: Diagnosis, Assessment and Disease Monitoring*. 2016;4:94-8. doi: <http://dx.doi.org/10.1016/j.dadm.2016.03.003>. PMID: 612393378.
439. Jia J, Zuo X, Jia XF, et al. Diagnosis and treatment of dementia in neurology outpatient departments of general hospitals in China. *Alzheimer's and Dementia*. 2016 01 Apr;12(4):446-53. doi: <http://dx.doi.org/10.1016/j.jalz.2015.06.1892>.
440. Johannessen A, Povlsen L, Bruvik Fy, et al. Implementation of a multicomponent psychosocial programme for persons with dementia and their families in Norwegian municipalities: experiences from the perspective of healthcare professionals who performed the intervention. *In Scand J Caring Sci*
441. Johnson L, Deatrack EJ, Oriol K. The use of music to improve exercise participation in people with dementia: a pilot study. *In Physical & occupational therapy in geriatrics*
442. Jones C, Moyle W, Murfield J, et al. Does Cognitive Impairment and Agitation in Dementia Influence Intervention Effectiveness? Findings From a Cluster-Randomized-Controlled Trial With the Therapeutic Robot, PARO. *J Am Med Dir Assoc*. 2018 Jul;19(7):623-6. doi: <https://dx.doi.org/10.1016/j.jamda.2018.02.014>. PMID: 29656838.

443. Kales HC, Gitlin LN, Stanislawski B, et al. WeCareAdvisor™: The Development of a Caregiver-focused, Web-based Program to Assess and Manage Behavioral and Psychological Symptoms of Dementia. *Alzheimer Dis Assoc Disord*. 2017 Jul-Sep;31(3):263-70. doi: <https://dx.doi.org/10.1097/WAD.0000000000000177>. PMID: 27849639.
444. Kang HS, Ahn IS, Kim JH, et al. Neuropsychiatric symptoms in Korean patients with Alzheimer's disease: exploratory factor analysis and confirmatory factor analysis of the neuropsychiatric inventory. *Dement Geriatr Cogn Disord*. 2010;29(1):82-7. doi: <https://dx.doi.org/10.1159/000264629>. PMID: 20130406.
445. Kerkhof YJ, Graff MJ, Bergsma A, et al. Better self-management and meaningful activities thanks to tablets? Development of a person-centered program to support people with mild dementia and their carers through use of hand-held touch screen devices. *Int Psychogeriatr*. 2016 11;28(11):1917-29. PMID: 27425002.
446. Kessels RPC, Olde Hensken LMG. Effects of errorless skill learning in people with mild-to-moderate or severe dementia: a randomized controlled pilot study. *NeuroRehabilitation*. 2009;25(4):307-12. doi: 10.3233/NRE-2009-0529. PMID: 105277868. Language: English. Entry Date: 20100205. Revision Date: 20150711. Publication Type: Journal Article.
447. Klages K, Zecevic A, Orange JB, et al. Potential of Snoezelen room multisensory stimulation to improve balance in individuals with dementia: a feasibility randomized controlled trial. *Clin Rehabil*. 2011 Jul;25(7):607-16. doi: <https://dx.doi.org/10.1177/0269215510394221>. PMID: 21382864.
448. Kolanowski AM, Litaker M, Buettner L. Efficacy of theory-based activities for behavioral symptoms of dementia. *Nurs Res*. 2005 Jul-Aug;54(4):219-28. PMID: 16027564.
449. Koltai DC, Welsh-Bohmer KA, Schmechel DE. Influence of anosognosia on treatment outcome among dementia patients. *Neuropsychol*. 2001 Jul;11(3-4):455-75. doi: <http://dx.doi.org/10.1080/09602010042000097>. PMID: 2001-07363-012.
450. La Garce M. Daylight interventions and Alzheimer's behaviors: A twelve-month study. *Journal of Architectural and Planning Research*. 2004 Fal;21(3):257-69. PMID: 2004-19048-003.
451. Lamotte G, Shah RC, Lazarov O, et al. Exercise Training for Persons with Alzheimer's Disease and Caregivers: A Review of Dyadic Exercise Interventions. *J Mot Behav*. 2017 Jul-Aug;49(4):365-77. doi: <https://dx.doi.org/10.1080/00222895.2016.1241739>. PMID: 27870597.
452. Lancioni GE, O'Reilly MF, Singh NN, et al. Assessing the impact and social perception of self-regulated music stimulation with patients with Alzheimer's disease. *Research in Developmental Disabilities*. 2013;34(1):139-46. doi: 10.1016/j.ridd.2012.07.026. PMID: 104401179. Language: English. Entry Date: 20130607. Revision Date: 20150711. Publication Type: Journal Article.
453. Lancioni GE, Singh NN, O'Reilly MF, et al. Persons with mild or moderate Alzheimer's disease learn to use urine alarms and prompts to avoid large urinary accidents. *Research in Developmental Disabilities*. 2011;32(5):1998-2004. doi: 10.1016/j.ridd.2011.04.011. PMID: 104645817. Language: English. Entry Date: 20111028. Revision Date: 20150711. Publication Type: Journal Article.
454. Lancioni GE, Singh NN, O'Reilly MF, et al. Technology-aided verbal instructions to help persons with mild or moderate Alzheimer's disease perform daily activities. *Research in Developmental Disabilities*. 2010;31(6):1240-50. doi: 10.1016/j.ridd.2010.07.021. PMID: 104927845. Language: English. Entry Date: 20110204. Revision Date: 20150711. Publication Type: Journal Article.
455. Lanza G, Centonze SS, Destro G, et al. Shiatsu as an adjuvant therapy for depression in patients with Alzheimer's disease: A pilot study. *Complement Ther Med*. 2018 Jun;38:74-8. doi: <https://dx.doi.org/10.1016/j.ctim.2018.04.013>. PMID: 29857884.
456. Laporte UF, Graske J, Grill S, et al. Regional dementia care networks in Germany: changes in caregiver burden at one-year follow-up and associated factors. *In Int Psychogeriatr*
457. Lee GY, Yip CC, Yu EC, et al. Evaluation of a computer-assisted errorless learning-based memory training program for patients with early Alzheimer's disease in Hong Kong: a pilot study. *Clin Interv Aging*. 2013;8:623-33. doi: <https://dx.doi.org/10.2147/CIA.S45726>. PMID: 23766638.
458. Lee J, Choi BH, Oh E, et al. Treatment of Alzheimer's Disease with Repetitive Transcranial Magnetic Stimulation Combined with Cognitive Training: A Prospective, Randomized, Double-Blind, Placebo-Controlled Study. *J Clin Neurol*. 2016 Jan;12(1):57-64. doi: <https://dx.doi.org/10.3988/jcn.2016.12.1.57>. PMID: 26365021.
459. Lee MM, Camp CJ, Malone ML. Effects of intergenerational Montessori-based activities programming on engagement of nursing home residents with dementia. *Clin Interv Aging*. 2007;2(3):477-83. PMID: 18044197.
460. Libin A, Cohen-Mansfield J. Therapeutic robot for nursing home residents with dementia: preliminary inquiry. *Am J Alzheimers Dis Other Demen*. 2004 Mar-Apr;19(2):111-6. PMID: 15106392.
461. Lin S, Yang Y, Qi Q, et al. The Beneficial Effect of Physical Exercise on Cognitive Function in a Non-dementia Aging Chinese Population. *Front Aging Neurosci*. 2019;11:238. doi: <https://dx.doi.org/10.3389/fnagi.2019.00238>. PMID: 31555125.
462. Livingston G, Lewis-Holmes E, Pitfield C, et al. Improving the end-of-life for people with dementia living in a care home: an intervention study. *Int Psychogeriatr*. 2013 Nov;25(11):1849-58. doi: <https://dx.doi.org/10.1017/S1041610213001221>. PMID: 23924580.
463. Low LF, Carroll S, Merom D, et al. We think you can dance! A pilot randomised controlled trial of dance for nursing home residents with moderate to severe dementia. *Complement Ther Med*. 2016 Dec;29:42-4. doi: <https://dx.doi.org/10.1016/j.ctim.2016.09.005>. PMID: 27912955.

464. Lyketsos CG, Lindell Veiel L, Baker A, et al. A randomized, controlled trial of bright light therapy for agitated behaviors in dementia patients residing in long-term care. *Int J Geriatr Psychiatry*. 1999 Jul;14(7):520-5. PMID: 10440971.
465. Maci T, Pira FL, Quattrocchi G, et al. Physical and cognitive stimulation in Alzheimer Disease. the GAIA Project: a pilot study. *Am J Alzheimers Dis Other Demen*. 2012 Mar;27(2):107-13. doi: <https://dx.doi.org/10.1177/1533317512440493>. PMID: 22495338.
466. Manckoundia P, Taroux M, Kubicki A, et al. Impact of ambulatory physiotherapy on motor abilities of elderly subjects with Alzheimer's disease. In *Geriatr Gerontol Int*
467. Maseda A, Sanchez A, Marante MP, et al. Effects of Multisensory Stimulation on a Sample of Institutionalized Elderly People With Dementia Diagnosis: A Controlled Longitudinal Trial. *Am J Alzheimers Dis Other Demen*. 2014 Aug;29(5):463-73. doi: <https://dx.doi.org/10.1177/1533317514522540>. PMID: 24604894.
468. Matthews J, Lingler J, Campbell G, et al. Usability of a Wearable Camera System for Dementia Family Caregivers. In *Journal of healthcare engineering*
469. McCaffrey R, Park J, Newman D, et al. The effect of chair yoga in older adults with moderate and severe Alzheimer's disease. *Res*. 2014 Jul-Aug;7(4):171-7. doi: <https://dx.doi.org/10.3928/19404921-20140218-01>. PMID: 24568209.
470. McCreedy EM, Mor V, Thomas K, et al. Effects of music and memory on agitated and aggressive behaviors in dementia: results from nursing home pilot study. *J Am Geriatr Soc*. 2019;67:S91-. doi: 10.1111/jgs.15898. PMID: CN-01940453.
471. Mihailidis A, Boger J, Craig T, et al. The COACH prompting system to assist older adults with dementia through handwashing: an efficacy study. In *BMC geriatr*
472. Milev RV, Kellar T, McLean M, et al. Multisensory stimulation for elderly with dementia: a 24-week single-blind randomized controlled pilot study. *Am J Alzheimers Dis Other Demen*. 2008 Aug-Sep;23(4):372-6. doi: <https://dx.doi.org/10.1177/1533317508316681>. PMID: 18453647.
473. Mioshi E, McKinnon C, Savage S, et al. Improving burden and coping skills in frontotemporal dementia caregivers: a pilot study. *Alzheimer Dis Assoc Disord*. 2013;27(1):84-6. doi: 10.1097/WAD.0b013e31824a7f5b. PMID: 108031647. Language: English. Entry Date: 20130823. Revision Date: 20150712. Publication Type: Journal Article.
474. Mittelman MS, Bartels SJ. Translating research into practice: case study of a community-based dementia caregiver intervention. *Health Aff (Millwood)*. 2014 Apr;33(4):587-95. doi: <https://dx.doi.org/10.1377/hlthaff.2013.1334>. PMID: 24711319.
475. Moyle W, Cooke M, Beattie E, et al. Exploring the effect of companion robots on emotional expression in older adults with dementia: a pilot randomized controlled trial. *J Gerontol Nurs*. 2013 May;39(5):46-53. doi: <https://dx.doi.org/10.3928/00989134-20130313-03>. PMID: 23506125.
476. Moyle W, Cooke M, O'Dwyer ST, et al. The effect of foot massage on long-term care staff working with older people with dementia: a pilot, parallel group, randomized controlled trial. *BMC Nurs*. 2013 Feb 18;12:5. doi: <https://dx.doi.org/10.1186/1472-6955-12-5>. PMID: 23414448.
477. Nair BK, Heim C, Krishnan C, et al. The effect of Baroque music on behavioural disturbances in patients with dementia. *Australas J Ageing*. 2011 Mar;30(1):11-5. doi: <https://dx.doi.org/10.1111/j.1741-6612.2010.00439.x>. PMID: 21395934.
478. Nakanishi M, Hirooka K, Imai Y, et al. Dementia Care Competence Among Care Professionals and Reduced Challenging Behavior of Home-Dwelling Persons with Dementia: A Pre- and Post-Intervention Data Analysis. *J Alzheimers Dis*. 2018;64(2):515-23. doi: <https://dx.doi.org/10.3233/JAD-171077>. PMID: 29914021.
479. Nakrem S, Stensvik GT, Skjong RJ, et al. Staff experiences with implementing a case conferencing care model in nursing homes: a focus group study. *BMC Health Serv Res*. 2019 Mar 27;19(1):191. doi: <https://dx.doi.org/10.1186/s12913-019-4034-0>. PMID: 30917815.
480. Narme P, Tonini A, Khatir F, et al. Non pharmacological treatment for Alzheimer's disease: comparison between musical and non-musical interventions. In *Geriatric ET psychologie neuropsychiatrie du vieillissement*
481. Okumura Y, Tanimukai S, Asada T. Effects of short-term reminiscence therapy on elderly with dementia: a comparison with everyday conversation approaches. In *Psychogeriatrics*
482. Onor ML, Trevisiol M, Negro C, et al. Impact of a multimodal rehabilitative intervention on demented patients and their caregivers. *Am J Alzheimers Dis Other Demen*. 2007 Aug-Sep;22(4):261-72. PMID: 17712157.
483. Orrell M, Spector A, Thorgrimsen L, et al. A pilot study examining the effectiveness of maintenance Cognitive Stimulation Therapy (MCST) for people with dementia. *Int J Geriatr Psychiatry*. 2005;20(5):446-51. PMID: 106207035. Language: English. Entry Date: 20070105. Revision Date: 20150711. Publication Type: Journal Article.
484. Ott CH, Kelber ST, Blaylock M. "Easing the way" for spouse caregivers of individuals with dementia: a pilot feasibility study of a grief intervention. *Res*. 2010 Apr;3(2):89-99. doi: <https://dx.doi.org/10.3928/19404921-20100302-01>. PMID: 20415358.
485. Ozdemir L, Akdemir N. Effects of multisensory stimulation on cognition, depression and anxiety levels of mildly-affected Alzheimer's patients. *J Neurol Sci*. 2009 Aug 15;283(1-2):211-3. doi: <https://dx.doi.org/10.1016/j.jns.2009.02.367>. PMID: 19289242.
486. Paddick SM, Mkenda S, Mbowe G, et al. Cognitive stimulation therapy as a sustainable intervention for dementia in sub-Saharan Africa: feasibility and clinical efficacy using a stepped-wedge design. *Int Psychogeriatr*. 2017 Jun;29(6):979-89. doi: <https://dx.doi.org/10.1017/S1041610217000163>. PMID: 28222815.

487. Parker LJ, Gaugler JE, Samus Q, et al. Adult Day Service Use Decreases Likelihood of a Missed Physician's Appointment Among Dementia Caregivers. *J Am Geriatr Soc*. 2019 Jul;67(7):1467-71. doi: <https://dx.doi.org/10.1111/jgs.15995>. PMID: 31219175.
488. Paun O, Farran CJ, Perraud S, et al. Successful caregiving of persons with Alzheimer's disease: skill development over time. *Alzheimer's Care Quarterly*. 2004;5(3):241-51. PMID: 106657982. Language: English. Entry Date: 20041105. Revision Date: 20150711. Publication Type: Journal Article.
489. Ploeg J, Northwood M, Dugleby W, et al. Caregivers of older adults with dementia and multiple chronic conditions: Exploring their experiences with significant changes. *Dementia*. 2019 Mar 06;1471301219834423. doi: <https://dx.doi.org/10.1177/1471301219834423>. PMID: 30841745.
490. Pomeroy VM. The effect of physiotherapy input on mobility skills of elderly people with severe dementing illness. *Clin Rehabil*. 1993;7(2):163-70. PMID: 23161041.
491. Pomykala KL, Silverman DH, Geist CL, et al. A pilot study of the effects of meditation on regional brain metabolism in distressed dementia caregivers. *Aging health*. 2012 Oct 01;8(5):509-16. PMID: 23378856.
492. Rapp T, Grand A, Cantet C, et al. Public financial support receipt and non-medical resource utilization in Alzheimer's disease results from the PLASA study. *Soc Sci Med*. 2011 Apr;72(8):1310-6. doi: <https://dx.doi.org/10.1016/j.socscimed.2011.02.039>. PMID: 21463914.
493. Resnick B, Galik E, Kolanowski A, et al. Reliability and Validity of the Care Plan Checklist for Evidence of Person-Centered Approaches for Behavioral and Psychological Symptoms Associated With Dementia. *J Am Med Dir Assoc*. 2017 Nov 28;28:28. doi: <https://dx.doi.org/10.1016/j.jamda.2017.10.021>. PMID: 29191763.
494. Rokstad AM, Engedal K, Kirkevold O, et al. The association between attending specialized day care centers and the quality of life of people with dementia. *Int Psychogeriatr*. 2017 04;29(4):627-36. doi: <https://dx.doi.org/10.1017/S1041610216002015>. PMID: 27852334.
495. Romo ML, Schooling CM. Examining the Causal Role of Leptin in Alzheimer Disease: A Mendelian Randomization Study. *Neuroendocrinology*. 2017 01 Aug;105(2):182-8. doi: <http://dx.doi.org/10.1159/000475713>. PMID: 615575945.
496. Rosenberg PB, Mielke MM, Lyketsos CG. Caregiver assessment of patients' depression in Alzheimer disease: longitudinal analysis in a drug treatment study. *Am J Geriatr Psychiatry*. 2005 Sep;13(9):822-6. PMID: 16166413.
497. Rosewarne R, Bruce A, McKenna M. Dementia programme effectiveness in long-term care. *Int J Geriatr Psychiatry*. 1997 Feb;12(2):173-82. PMID: 9097210.
498. Rostad H, Puts M, Cvancarova SM, et al. Associations between Pain and Quality of Life in Severe Dementia: a Norwegian Cross-Sectional Study. In *Dement Geriatr Cogn Dis Extra*
499. Rouch I, Pongan E, Trombert B, et al. One-Year Evolution of Behavioral and Psychological Symptoms of Dementia in Patients Initially Hospitalized in Cognitive Behavioral Units: the EVITAL Prospective Cohort. In *J Alzheimers Dis*
500. Rouse HJ, Small BJ, Faust ME. Assessment of Cognitive Training & Social Interaction in People with Mild to Moderate Dementia: A Pilot Study. *Clin Gerontol*. 2019 01 Jul;42(4):421-34. doi: <http://dx.doi.org/10.1080/07317115.2019.1590489>. PMID: 626971925.
501. Rutherford G, Lithgow B, Moussavi Z. Short and long-term effects of rTMS treatment on Alzheimer's disease at different stages: a pilot study. In *Journal of experimental neuroscience*
502. Salotti P, Sanctis B, Clementi A, et al. Evaluation of the efficacy of a cognitive rehabilitation treatment on a group of Alzheimer's patients with moderate cognitive impairment: a pilot study. In *Aging clinical and experimental research*
503. Sanchez A, Marante-Moar MP, Sarabia C, et al. Multisensory Stimulation as an Intervention Strategy for Elderly Patients With Severe Dementia: A Pilot Randomized Controlled Trial. *Am J Alzheimers Dis Other Demen*. 2016 Jun;31(4):341-50. doi: <https://dx.doi.org/10.1177/1533317515618801>. PMID: 26631687.
504. Sánchez A, Maseda A, Marante-Moar MP, et al. Comparing the Effects of Multisensory Stimulation and Individualized Music Sessions on Elderly People with Severe Dementia: A Randomized Controlled Trial. *J Alzheimers Dis*. 2016;52(1):303-15. doi: 10.3233/JAD-151150. PMID: 114968532. Language: English. Entry Date: 20170126. Revision Date: 20180615. Publication Type: journal article.
505. Sanchez-Valdeon L, Fernandez-Martinez E, Loma-Ramos S, et al. Canine-Assisted Therapy and Quality of Life in People With Alzheimer-Type Dementia: Pilot Study. *Front Psychol*. 2019;10:1332. doi: <https://dx.doi.org/10.3389/fpsyg.2019.01332>. PMID: 31244731.
506. Santana-Sosa E, Barriopedro MI, López-Mojares LM, et al. Exercise training is beneficial for Alzheimer's patients. *Int J Sports Med*. 2008;29(10):845-50. PMID: 105568866. Language: English. Entry Date: 20090206. Revision Date: 20150711. Publication Type: Journal Article.
507. Scharre DW, Weichart E, Nielson D, et al. Deep Brain Stimulation of Frontal Lobe Networks to Treat Alzheimer's Disease. *J Alzheimers Dis*. 2018;62(2):621-33. doi: <https://dx.doi.org/10.3233/JAD-170082>. PMID: 29400666.
508. Scherder EJ, Bouma A. Effects of transcutaneous electrical nerve stimulation on memory and behavior in Alzheimer's disease may be stage-dependent. *Biol Psychiatry*. 1999 Mar 15;45(6):743-9. PMID: 10188004.
509. Scherder EJ, Van Someren EJ, Swaab DF. Transcutaneous electrical nerve stimulation (TENS) improves the rest-activity rhythm in midstage Alzheimer's disease. *Behav Brain Res*. 1999 May;101(1):105-7. PMID: 10342404.

510. Schiffman SS, Graham BG, Sattely-Miller EA, et al. Taste, smell and neuropsychological performance of individuals at familial risk for Alzheimer's disease. *Neurobiol Aging*. 2002 May-Jun;23(3):397-404. PMID: 11959402.
511. Schinköthe D, Wilz G. The Assessment of Treatment Integrity in a Cognitive Behavioral Telephone Intervention Study With Dementia Caregivers. *Clin Gerontol*. 2014;37(3):211-34. doi: 10.1080/07317115.2014.886653. PMID: 104064713. Language: English. Entry Date: 20140418. Revision Date: 20151023. Publication Type: Journal Article.
512. Schulz R, Mendelsohn AB, Haley WE, et al. End-of-life care and the effects of bereavement on family caregivers of persons with dementia. *N Engl J Med*. 2003 Nov 13;349(20):1936-42. PMID: 14614169.
513. Schulze J, Bussche H, Kaduskiewicz H, et al. Institutionalization in incident dementia cases in comparison to age- and sex- matched controls: a 5-year follow-up from Germany. In *Social psychiatry and psychiatric epidemiology*
514. Sedeno L, Piguët O, Abrevaya S, et al. Tackling variability: a multicenter study to provide a gold-standard network approach for frontotemporal dementia. In *Human brain mapping*
515. Siegel Jr BV, Buchsbaum MS, Starr A, et al. Glucose metabolic rate and progression of illness in Alzheimer's disease. *Int J Geriatr Psychiatry*. 1995;10(8):659-67. doi: <http://dx.doi.org/10.1002/gps.930100806>. PMID: 25262446.
516. Sindi S, Ngandu T, Hovatta I, et al. Baseline Telomere Length and Effects of a Multidomain Lifestyle Intervention on Cognition: The FINGER Randomized Controlled Trial. *J Alzheimers Dis*. 2017;59(4):1459-70. doi: <https://dx.doi.org/10.3233/JAD-170123>. PMID: 28777749.
517. Snow AL, Hovanec L, Brandt J. A controlled trial of aromatherapy for agitation in nursing home patients with dementia. *J Altern Complement Med*. 2004;10(3):431-7. doi: 10.1089/1075553041323696. PMID: 106568581. Language: English. Entry Date: 20050128. Revision Date: 20150820. Publication Type: Journal Article.
518. Sposito G, Barbosa A, Figueiredo D, et al. Effects of multisensory and motor stimulation on the behavior of people with dementia. *Dementia*. 2017 Apr;16(3):344-59. doi: <https://dx.doi.org/10.1177/1471301215592080>. PMID: 26112166.
519. Spring HJ, Rowe MA, Kelly A. Improving caregivers' well-being by using technology to manage nighttime activity in persons with dementia. *Res*. 2009;2(1):39-48. doi: 10.3928/19404921-20090101-10. PMID: 105631700. Language: English. Entry Date: 20090306. Revision Date: 20150711. Publication Type: Journal Article.
520. Stevens AB, Lancer K, Smith ER, et al. Engaging communities in evidence-based interventions for dementia caregivers. *Fam Community Health*. 2009 Jan-Mar;32(1 Suppl):S83-92. doi: <https://dx.doi.org/10.1097/01.FCH.0000342843.28477.72>. PMID: 19065098.
521. Sutcliffe C, Larner S. Counselling carers of the elderly at home: a preliminary study. *Br J Clin Psychol*. 1988 May;27(Pt 2):177-8. PMID: 3293671.
522. Svensson B, Hansson L. Mental health first aid for the elderly: A pilot study of a training program adapted for helping elderly people. *Aging Ment Health*. 2017 Jun;21(6):595-601. doi: <https://dx.doi.org/10.1080/13607863.2015.1135873>. PMID: 26829477.
523. Teri L, Logsdon RG, Weiner MF, et al. Treatment for agitation in dementia patients: A behavior management approach. *Psychotherapy*. 1998 Winter;35(4):436-43. PMID: 29227715.
524. Thyrian JR, Eichler T, Hertel J, et al. Burden of Behavioral and Psychiatric Symptoms in People Screened Positive for Dementia in Primary Care: Results of the DelpHi-Study. *J Alzheimers Dis*. 2015;46(2):451-9. doi: <https://dx.doi.org/10.3233/JAD-143114>. PMID: 25765916.
525. Thyrian JR, Eichler T, Pooch A, et al. Systematic, early identification of dementia and dementia care management are highly appreciated by general physicians in primary care - results within a cluster-randomized-controlled trial (DelpHi). *J Multidiscip Healthc*. 2016;9:183-90. doi: <https://dx.doi.org/10.2147/JMDH.S96055>. PMID: 27143912.
526. Thyrian JR, Eichler T, Reimann M, et al. Depressive symptoms and depression in people screened positive for dementia in primary care - results of the DelpHi-study. *Int Psychogeriatr*. 2016 06;28(6):929-37. doi: <https://dx.doi.org/10.1017/S1041610215002458>. PMID: 26785854.
527. Travers C. Increasing enjoyable activities to treat depression in nursing home residents with dementia: A pilot study. *Dementia*. 2017 Feb;16(2):204-18. doi: <https://dx.doi.org/10.1177/1471301215586069>. PMID: 25972128.
528. van der Steen JT, Albers G, Licht-Strunk E, et al. A validated risk score to estimate mortality risk in patients with dementia and pneumonia: barriers to clinical impact. *Int Psychogeriatr*. 2011 Feb;23(1):31-43. doi: <https://dx.doi.org/10.1017/S1041610210001079>. PMID: 20653988.
529. van Mierlo LD, Meiland FJ, Dröes RM, et al. Dementelcoach: effect of telephone coaching on carers of community-dwelling people with dementia. *Int Psychogeriatr*. 2012;24(2):212-22. doi: 10.1017/S1041610211001827. PMID: 104511872. Language: English. Entry Date: 20120511. Revision Date: 20161117. Publication Type: journal article.
530. van Paasschen J, Clare L, Yuen KS, et al. Cognitive rehabilitation changes memory-related brain activity in people with Alzheimer disease. *Neurorehabil Neural Repair*. 2013 Jun;27(5):448-59. doi: <https://dx.doi.org/10.1177/1545968312471902>. PMID: 23369983.
531. Van Someren EJ, Scherder EJ, Swaab DF. Transcutaneous electrical nerve stimulation (TENS) improves circadian rhythm disturbances in Alzheimer disease. *Alzheimer Dis Assoc Disord*. 1998 Jun;12(2):114-8. PMID: 9651141.
532. Vidoni ED, Watts AS, Burns JM, et al. Feasibility of a Memory Clinic-Based Physical Activity Prescription Program. *J Alzheimers Dis*. 2016;53(1):161-70. doi: 10.3233/JAD-160158. PMID: 116350716. Language:

English. Entry Date: 20180129. Revision Date: 20180525. Publication Type: journal article.

533. Watson K, Hatcher D, Good A. A randomised controlled trial of Lavender (*Lavandula Angustifolia*) and Lemon Balm (*Melissa Officinalis*) essential oils for the treatment of agitated behaviour in older people with and without dementia. *Complement Ther Med*. 2019 Feb;42:366-73. doi: <https://dx.doi.org/10.1016/j.ctim.2018.12.016>. PMID: 30670268.
534. Wells Y, Jorm AF. Evaluation of a special nursing home unit for dementia sufferers: a randomised controlled comparison with community care. *Aust N Z J Psychiatry*. 1987 Dec;21(4):524-31. PMID: 3449047.
535. Werner C, Rosner R, Wiloth S, et al. Time course of changes in motor-cognitive exergame performances during task-specific training in patients with dementia: identification and predictors of early training response. *J Neuroengineering Rehabil*. 2018 Nov 08;15(1):100. doi: <https://dx.doi.org/10.1186/s12984-018-0433-4>. PMID: 30409202.
536. Wilks SE, Boyd PA, Bates SM, et al. Montessori-based activities among persons with late-stage dementia: Evaluation of mental and behavioral health outcomes. *Dementia*. 2017 Jan 01;1471301217703242. doi: <https://dx.doi.org/10.1177/1471301217703242>. PMID: 28449593.
537. Wiloth S, Werner C, Lemke NC, et al. Motor-cognitive effects of a computerized game-based training method in people with dementia: a randomized controlled trial. *Aging Ment Health*. 2018 Sep;22(9):1124-35. doi: <https://dx.doi.org/10.1080/13607863.2017.1348472>. PMID: 28682124.

538. Woods DL, Dimond M. The effect of therapeutic touch on agitated behavior and cortisol in persons with Alzheimer's disease. *Biol Res Nurs*. 2002 Oct;4(2):104-14. PMID: 12408216.
539. Wu E, Barnes DE, Ackerman SL, et al. Preventing Loss of Independence through Exercise (PLIÉ): qualitative analysis of a clinical trial in older adults with dementia. *Aging Ment Health*. 2015;19(4):353-62. doi: [10.1080/13607863.2014.935290](https://doi.org/10.1080/13607863.2014.935290). PMID: 103879583. Language: English. Entry Date: 20150121. Revision Date: 20160922. Publication Type: Journal Article.
540. Yoon J, Lee S, Lim H, et al. The effects of cognitive activity combined with active extremity exercise on balance, walking activity, memory level and quality of life of an older adult sample with dementia. In *J Phys Ther Sci*
541. Yu F, Nelson NW, Savik K, et al. Affecting cognition and quality of life via aerobic exercise in Alzheimer's disease. *West J Nurs Res*. 2013 Jan;35(1):24-38. doi: <https://dx.doi.org/10.1177/0193945911420174>. PMID: 21911546.
542. Yu F, Savik K, Wyman JF, et al. Maintaining physical fitness and function in Alzheimer's disease: a pilot study. *Am J Alzheimers Dis Other Demen*. 2011 Aug;26(5):406-12. doi: <https://dx.doi.org/10.1177/1533317511414861>. PMID: 21750046.
543. Zhou Y, Jia J. Clinical observation on catgut implantation at acupoint for treatment of Alzheimer's disease. In *Zhongguo zhen jiu [Chinese acupuncture & moxibustion]*
544. Zhu C, Cosentino S, Ornstein K, et al. Interactive Effects of Dementia Severity and Comorbidities on Medicare Expenditures. In *J Alzheimers Dis*

Appendix C. Treatment for BPSD

Assisted Therapy

Table C-1. Risk of bias assessment: assisted therapy

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Moyle 2017 ¹ (28780395) Jones 2018 ² (29656838) Mervin 2018 ³ (29325922) Moyle 2018 ⁴ (29563027)	10 weeks	Low 10 weeks: 4.3% 15 weeks: 7.0%	Low	Medium	Medium	Medium	Medium	Funder	Medium
Peterson 2017 ⁵ (27716673)	3 months	Medium NR	Medium	High	Medium	High	X	NR	High
Olsen 2016 ⁶ (27155968)	12 weeks 6 months	Low 12 weeks: 1% 6 months: 1%	Low	High	Medium	Low	X	Foundation	High
Olsen 2016 ⁷ (26807956)	12 weeks 6 months	Low 12 weeks: 12% 6 months: 17%	Low	High	Medium	Low	X	Foundation	High
Bemelmans 2015 ⁸ (26115817)	4 months	High 21.9%	X	X	X	X	X	NR	High
Joranson 2015 ⁹ (26096582) Joranson 2016 ¹⁰ (27434512)	12 weeks 6 months	Medium 12 weeks: 10% 6 months: 16.7%	Medium	High	High	Low	X	Government	High
Travers 2013 ¹¹ (NA)	11 weeks	Medium 17.9%	Medium	Low	High	Low	X	Foundation	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table C-2. Characteristics of included studies: assisted therapy

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Moyle 2017 ¹ (28780395) Australia Medium Explanatory	Individual, non-facilitated sessions with PARO (robotic seal) for PLWD over 10 weeks; three afternoon sessions per week, 15 minutes/session	1. Standard care for PLWD at respective facilities 2. Push toy (PARO with robotic features disabled) for PLWD over 10 weeks; three afternoon sessions per week, 15 minutes/session	Long-term care facilities approved and accredited by the Australian government Cluster RCT Multisite 18 Clusters 415 PLWD	Any dementia type or severity. RUDAS ≤ 22 and documented dementia diagnosis	N=415 76% Female 85 years Race NR Education NR	None	NA	NA	NA	NA	10 weeks 15 weeks	Agitation (observed) CMAI-SF Positive behavioral engagement Using object for social engagement Positive verbal engagement Visual engagement	NA	

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: Char=characteristics; CMAI-SF=Cohen-Mansfield Agitation Inventory; FC=formal caregiver; IC=informal caregiver; PMID=PubMed Identification Number; PLWD=Persons Living with Dementia; Rob=Risk of Bias; RUDAS=Rowland Universal Dementia Assessment Scale; SES=socioeconomic status; RS=Reporting Status

Table C-3. Modified PRECIS-2: assisted therapy

Study (PMID)	Eligibility Criteria	Recruitment Path	Setting	Intervention Organization	Flexibility of Intervention: Delivery	Flexibility of Intervention: Adherence	Followup	Primary Outcome	Analysis	Overall Category
										Applicability and/or Qualifiers
Moyle 2017 ¹ (28780395) Jones 2018 ² (29656838) Mervin 2018 ³ (29325922) Moyle 2018 ⁴ (29563027)	Balanced	Mostly Explanatory	Mostly Explanatory	Explanatory	Explanatory	Mostly Explanatory	Mostly Explanatory	Mostly Explanatory	Balanced	Explanatory

Abbreviations: NA=Not Assessed; PMID=PubMed Identification Number

Table C-4. PLWD outcomes summary low and medium risk of bias studies: assisted therapy

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p Value
Moyle 2017 ¹ (28780395) PARO vs. usual care Medium Explanatory	Adjusted Mean Difference (95% CI) Reduction in agitation (observed via video recording) 10 weeks	3.3 (0.9, 5.8)	NR	NR	p=0.008
Moyle 2017 ¹ (28780395) PARO vs. usual care Medium Explanatory	Adjusted Mean Difference (95% CI) CMAI-SF 10 weeks	-1.9 (-5.8, 2.0)	NR	NR	p=0.34
Moyle 2017 ¹ (28780395) PARO vs. usual care Medium Explanatory	Mortality % (N) 15 weeks	NR	5.1% (7)	3.6% (5)	NR
Moyle 2017 ¹ (28780395) PARO vs. plush toy Medium Explanatory	Adjusted Mean Difference (95% CI) Reduction in agitation (observed via video recording) 10 weeks	1.3 (-1.7, 4.2)	NR	NR	p=0.39
Moyle 2017 ¹ (28780395) PARO vs. plush toy Medium Explanatory	Adjusted Mean Difference (95% CI) CMAI-SF 10 weeks	-0.99 (-5.8, 3.8)	NR	NR	p=0.68
Moyle 2017 ¹ (28780395) PARO vs. plush toy Medium Explanatory	Adjusted Mean Difference (95% CI) Positive behavioral engagement (observed via video recording) 10 weeks	6.3 (-0.8, 13.5)	NR	NR	p=0.08
Moyle 2017 ¹ (28780395) PARO vs. plush toy Medium Explanatory	Adjusted Mean Difference (95% CI) Using object for social engagement (observed via video recording) 10 weeks	1.2 (-0.6, 3.0)	NR	NR	p=0.18
Moyle 2017 ¹ (28780395) PARO vs. plush toy Medium Explanatory	Adjusted Mean Difference (95% CI) Positive verbal engagement (observed via video recording) 10 weeks	3.6 (0.81, 6.40)	NR	NR	p=0.01
Moyle 2017 ¹ (28780395) PARO vs. plush toy Medium Explanatory	Adjusted Mean Difference (95% CI) Visual engagement (observed via video recording) 10 weeks	13.1 (9.06, 17.05)	NR	NR	p<0.0001
Moyle 2017 ¹ (28780395) PARO vs. plush toy Medium Explanatory	Mortality % (N) 15 weeks	NR	5.1% (7)	10% (14)	NR

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: CI=Confidence interval; NR=Not Reported; PMID=PubMed Identification Number; RoB=Risk of Bias

Table C-5. Summary of strength of evidence for PLWD outcomes: robot-assisted therapy

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
PARO vs. usual care Agitation	10 weeks	1 clustered RCT (n=257)	Evidence was mixed for agitation reduction. Observation via video recordings favored PARO; however, CMAI-SF showed no difference between groups.	Moderate	Unknown	Direct	Imprecise	Insufficient
PARO vs. plush toy Agitation	10 weeks	1 clustered RCT (n=278)	No difference between groups in measures of agitation.	Moderate	Unknown	Direct	Imprecise	Insufficient
PARO vs. plush toy Engagement	10 weeks	1 clustered RCT (n=278)	Observation via video recordings showed improvements in PARO for positive verbal engagement and visual engagement compared to plush toy; however, there was no difference for positive behavioral engagement and using the object for social engagement.	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: n=Number, CMAI-SF=Cohen-Mansfield Agitation Inventory-Short Form; RCT=Randomized controlled trial.

Evidence Map: Assisted Therapy

Table C-6. Characteristics of evidence map studies: assisted therapy

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Pu, 2020 ¹³ (32122797) Pu (31738463) ¹⁴ Australia Pilot	PARO robotic pet, 30 minutes 5 times a week for 6 weeks individually in bedrooms	Usual care	Long-term care facilities RCT 43 PLWD	Probably or possible dementia and chronic pain	N=43 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	6 weeks	PAINAD CMAI-SF Medication use	NA
Balzotti 2019 ¹⁵ (30136743) Italy Small sample	Gesture verbal treatment: develop PLWD interactive use of gestures with other communication strategies. 2 1-hour group sessions weekly for 12 weeks. Doll therapy: 1 hour daily for 12 weeks.	Usual rehabilitative care	AD care home Quasi-experimental 30 PLWDs	Dementia diagnosis based on DSM-V criteria;	N=30 Mean age: yes % Female: yes % majority race: no Education: yes	None	NA	None	NA	NA	12 weeks	NPI-Q	NA
Menna, 2019 ¹⁶ (30740833) Italy Pilot	Dog-assisted group therapy; 12 weekly sessions	Unclear	Adult daycare center RCT 22 PLWD	Mild to moderate AD (per MMSE)	N=22 Age: no Sex: no Race: no Education: no	None	NA	NA	NA	NA	3 months	Cortisol level GDS	NA
Moyle, 2019 ¹⁷ (30474401) Australia Pilot	Life-like baby doll-assisted therapy; 3 30 minute sessions per week for 3 weeks	Usual care	Long-term care facilities RCT 35 PLWD	Dementia with recent history of BPSD	N=35 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	3 months	CMAI-SF OERS	NA
Cantarella 2018 ¹⁸ (622559552) Italy Pilot study	Doll therapy, 60 minute sessions 5 times/week for 1 month	Hand warmers, 60-minute sessions 5 times/week for 1 month	Nursing home RCT, single site 32 PLWD	Severe AD or vascular dementia (≥5 Short Portable Mental Status Questionnaire) with BPSD	N=32 Age: yes Sex: yes Race: no Education: yes	None	NA	NA	N=NR Age: no Sex: no % majority race: no Education: no	None	1 month	NPI Eating Behavior Scale	NPI-Distress

Liang 2017 ¹⁹ (28668664) New Zealand Pilot	PARO robotic pet at day care and at home, 30-minute group sessions 2-3 times a week for 6 weeks at day care	Facilities' standard of care	Dementia day care centers and at home RCT; Multisite 30 PLWD 30 informal caregivers	Not specified	N=30 Age: yes Sex: yes Race: no Education: no	None	N=30 Age: yes Sex: yes % majority race: no Education: no	None	NA	NA	12 weeks	CMAI-SF NPI Brief Questionnaire CSDD	NA
Petersen 2017 ⁵ (27716673) US High RoB	PARO robotic pet, 20 minutes 3 times a week for 3 months	Facilities' standard of care	Senior living facilities RCT; Multisite 61 PLWD	Mild to moderate dementia according to DSM or NIA criteria	N=61 Age: yes Sex: yes Race: yes Education: no	None	NA	NA	NA	NA	3 months	RAID CSDD GDS	NA
Olsen 2016 ⁶ (27155968) Norway High RoB	Group animal therapy sessions with a dog, 2 times/week for 12 weeks	Usual care	Adult day care centers Cluster RCT 16 clusters 80 PLWD	Dementia diagnosis or MMSE <25	N=80 Age: yes Sex: yes Race: no Education: yes	Prior disability: yes	NA	NA	NA	NA	12 weeks 6 months	Berg Balance Scale CDR QUALID	NA
Olsen 2016 ⁷ (26807956) Norway High RoB	Group animal therapy sessions with a dog, 2 times/week, 30 minutes/session, for 12 weeks	Usual care	Nursing homes Cluster RCT 10 clusters 58 PLWD	Dementia diagnosis or MMSE <25	N=58 Age: yes Sex: yes Race: no Education: yes	Prior disability: yes	NA	NA	NA	NA	12 weeks 6 months	CSDD QUALID Brief Agitation Rating Scale CDR	NA
Bemelmans 2015 ⁸ (26115817) Netherlands High RoB	PARO robotic pet to provide comfort individually to distressed PLWD; as needed over 4 months	PARO robotic pet incorporated in daily activities for care support; as needed over 4 months	Care institutions for intramural psychogeriatric care Quasi experimental time series; multisite 91 PLWD	All dementia severity	N=91 Age: no Sex: yes Race: no Education: no	None	NA	NA	NA	NA	4 months	Individually Prioritized Problems Assessment	NA
Joranson 2015 ⁹ (26096582) Norway High RoB Joranson 2016 ¹⁰ (27434512)	PARO robotic pet, 30-minute group sessions 2 times a week for 12 weeks	Usual care	Nursing homes Cluster RCT; Multisite 60 PLWD	Dementia diagnosis with cognitive impairment or score lower than 25/30 on Norwegian MMSE	N=69 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	12 weeks 6 months	Norwegian BARS Norwegian CSDD QUALID	NA

Valenti Soler 2015 ²⁰ (26388764) Spain Pilot	PARO robotic pet, 30-40-minute group sessions 2 times a week for 3 months	1. Facility's standard of care 2. Humanoid robot, 30-40-minute group sessions 2 times a week for 3 months 3. Animal therapy with dogs, 30-40-minute group sessions 2 times a week for 3 months	Public nursing home RCT; Single site 117 PLWD	Diagnosis of neurodegenerative dementia, all dementia types	N=117 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	3 months	GDS NPI APADEM-NH Apathy Inventory QUALID	NA
Moyle 2013 ²¹ (23506125) Australia Pilot	PARO robotic pet, 45-minute group sessions 3 times a week for 5 weeks	Reading group, 45-minute group sessions 3 times a week for 5 weeks	Residential care facility Crossover RCT; Single site 18 PLWD	Diagnosis of mid to late stage dementia or DSM-IV criteria for probable dementia	N=18 Age: yes Sex: no Race: no Education: No	None	NA	NA	NA	NA	5 weeks 10 weeks	QoL-AD RAID AES GDS AWS OERS	NA
Travers 2013 ¹¹ (NA) Australia High RoB	Group animal therapy with a dog, 40-50-minute sessions 2-3 times/week for 11 weeks	Group therapy, 40-50-minute sessions 2-3 times/week for 11 weeks	Nursing home RCT, multisite 67 PLWD	Mild to moderate dementia (any type)	N=67 Age: yes Sex: yes Race: no Education: yes	None	NA	NA	NA	NA	11 weeks	QoL-AD SF-36 GDS-SF Multidimensional Observational Scale for Elderly Subjects	NA
Majic 2013 ²² (23831177) Germany Pilot	Animal therapy with a dog, one 40-minute session/week for 10 weeks	Usual Care	Nursing home Case-Control 75 PLWD	Dementia diagnosis (DSM-IV criteria) and MMSE <25	N=75 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	10 weeks	CMAI Dementia Mood Assessment Scale	NA

Abbreviations: AES=Apathy Evaluation Scale; APADEM-NH=Apathy Scale for Institutionalized Patients with Dementia Nursing Home; AWS=Revised Algase Wandering Scale–Nursing Home version; BARS=Brief Agitation Rating Scale; BPSD=Behavioral and Psychological Symptoms of Dementia; CDR=Clinical Dementia Rating; Char=characteristics; CMAI=Cohen Mansfield Agitation Inventory; CMAI-SF=Cohen-Mansfield Agitation Inventory-Short Form; CSDD=Cornell Scale for Depression in Dementia; DSM IV=Diagnostic and Statistical Manual; EM=Evidence Map; FC=formal caregiver; GDS=Geriatric Depression Scale; GDS=Global Deterioration Scale; GDS-SF=Geriatric Depression Scale Short Form; IC=informal caregiver; MMSE=Mini-mental State Exam; N=number; NA=Not Applicable; NIA=National Institute on Aging; NPI=Neuropsychiatric Inventory; OERS=Observed Emotion Rating Scale; PAINAD=Pain Assessment in Advanced Dementia; PMID=PubMed Identification Number PLWD=Persons Living with Dementia; QoL-AD=Quality of Life in Alzheimer’s Disease; QUALID=Quality of Life in Late-stage Dementia; RAID=Rating for Anxiety in Dementia; RCT=Randomized controlled trial; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; SF-36=Medical Outcomes Study 36-Item Short Form Health Survey;

Multi-Sensory Stimulation/Snoezelen

Table C-7. Risk of bias assessment: multisensory stimulation

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Strom 2017 ²³ (28553314)	24 weeks	Medium 12.5%	Medium	Medium	Medium	Medium	Medium	NR	Medium
Baker 2003 ²⁴ (12919265)	0 weeks 4 weeks	Low 6.62%	Low	Medium	Medium	Medium	High	NR	Medium
Van Weert, 2005 ²⁵ (15667372)	72 weeks	Medium 27.61%	Medium	High	High	Low	Low	Netherlands government	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table C-8. Characteristics of included studies: multisensory stimulation

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Strom, 2017 ²³ (28553314) Republic of Ireland Medium Explanatory	Group Sonas sessions led by a licensed practitioner; two 45-minute sessions per week for 24 weeks	Group newspaper reading sessions; two 45-minute sessions per week for 24 weeks Usual nursing facility care	Nursing facilities approved and accredited by the Republic of Ireland RCT Multisite 120 PLWD	Diagnosis of dementia, moderate to severe cognitive impairment (MMSE ≤ 20)	N=120 Age: Mean age 84.8 Sex: 77.5% female Race: NR Education NR	NR	NA	NA	NR	NR	12 weeks post-intervention 24 weeks post-intervention	Function	None
Baker 2003 ²⁴ (12919265) (11317951) ²⁶ United Kingdom, Netherlands, and Switzerland Medium Explanatory	Individual, non-directed sessions in a Snoezelen room, facilitated by a nursing staff, occupational therapist, or psychology assistants; two 30-minute sessions per week for 4 weeks	Individual activity sessions chosen according to participant's interests, e.g. playing cards, quizzes, looking at cards; facilitated by a nursing staff, occupational therapist, or psychology assistants; two sessions per week for 4 weeks, 30 minutes per session	Day hospital, psychogeriatric ward RCT Multisite 136 PLWD randomized, 120 analyzed	Diagnosis of Alzheimer's, vascular, or mixed dementia, moderate to severe cognitive impairment (MMSE ≤ 17)	N=136 Age: 82 years Sex: not reported Education: NR	NR	NA	NA	NR	NR	0 weeks post intervention 4 weeks post intervention	Neuropsychiatric symptoms	None

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: Char=characteristics; FC=formal caregiver; IC=informal caregiver; MMSE=Mini-mental State Exam; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RCT=Randomized controlled trial RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status

Table C-9. PLWD outcomes summary low and medium risk of bias studies: multisensory stimulation

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p Value
Baker 2003 ²⁴ (12919265) Explanatory	Unadjusted Mean Difference Reduction in neuropsychiatric symptoms (REHAB general behavior scale, UK participants only) 4 weeks (immediately post-intervention)	-1.9 points, CI not reported	Snoezelen	Active control	Not reported, but described as no difference
Baker 2003 ²⁴ (12919265) Explanatory	Unadjusted Mean Difference Reduction in neuropsychiatric symptoms (REHAB deviant behavior subscale, UK participants only) 4 weeks (immediately post-intervention)	0.5 points, CI not reported	Snoezelen	Active control	Not reported, but described as no difference
Baker 2003 ²⁴ (12919265) Explanatory	Unadjusted Mean Difference Reduction in neuropsychiatric symptoms (GIP scale, Dutch participants only) 4 weeks (immediately post-intervention)	-2.4 points, CI not reported	Snoezelen	Active control	Not reported, but described as no difference
Strom 2017 ²³ (28553314) Explanatory	Unadjusted mean difference Reduction in communication difficulty (HCS total score)(higher is worse) 24 weeks	3.0 points, CI not reported	Sonas	Active control	P=0.019
Strom 2017 ²³ (28553314) Explanatory	Unadjusted mean difference Reduction in communication difficulty (HCS total score) 24 weeks	4.0 points, CI not reported	Sonas	Inactive control	P=0.001

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; REHAB=Rehabilitation Evaluation Hall and Baker Scale; GIP=Behavior Observation Scale for Intra-Mural Psycho-Geriatrics; HCS=Holden Communication Scale

Table C-10. Summary of strength of evidence for PLWD outcomes: multisensory stimulation

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
MSS vs. Active control Neuropsychiatric symptoms	4 weeks	1 RCT (n=120)	No significant difference in measures of neuropsychiatric symptoms (UK population)	Moderate	Unknown	Indirect (BPSD were assessed using instruments that are not standard in the ADRD field (appear to be designed for psychiatric settings) and that contain other domains (e.g. mood, physical function))	Imprecise	Insufficient

Abbreviations: ADRD=Alzheimer's disease and related disorders; BPSD=behavioral and psychological symptoms of dementia; MSS=multisensory stimulation; n=number; RCT=randomized controlled trials

Evidence Map: Multi-Sensory Stimulation

Table C-11. Characteristics of evidence map studies: multisensory stimulation

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Maseda 2018 ²⁷ (29843244) Spain Small sample	Snoezelen	Individualized music sessions	Nursing home; single site 21 PLWD	N=21 Diagnosis of dementia and severe cognitive impairment (GDS=6 or 7)	N=21 Sex: yes Age: yes Race: no Education: yes	None	NA	NA	NA	NA	12 weeks	No validated outcomes	NR
Collier 2008 ²⁸ (21131677) United Kingdom Small sample	Multi-sensory environment	Gardening	Nursing homes or continuing care wards; multisite 30 PLWD	N=30 Moderate to severe dementia Mean MMSE 10.0	N=30 Sex: yes Age: yes Race: no Education: no	None	NA	NA	NA	NA	4 weeks	AMPS motor and process scales	NA
Staal 2007 ²⁹ (18441625) US Pilot	Snoezelen	Activity sessions	Geriatric psychiatric unit; single site 24 PLWD	N=24 Admitting diagnosis of dementia, criteria NR Mean MMSE 15.5	N=24 Sex: yes Age: yes Race: no Education: no	None	NA	NA	NA	NA	Unclear	PAS MAI, physical health domain SANS-AD Katz Index of ADLs RADL	NA
Van Weert 2005 ²⁵ (15667372) (15050851) ³⁰ Netherlands High RoB	Snoezelen	Usual care	Nursing home psychogeriatric wards; multisite 125 PLWD	N=125 Moderate to severe dementia, DSM-III criteria	N=125 Sex: yes Age: yes Race: no Education: no	None	NA	NA	NA	NA	18 months	BIP, 8 of 14 subscales CMAI CSDD	NA
Baillon 2004 ³¹ (15481068) United Kingdom Small sample	Snoezelen	Reminiscence therapy	Nursing home; single site	N=25 Median MMSE 4.0 among completers	N=25 Sex: yes Age: yes Race: no Education: no	None	NA	NA	NA	NA	1 day (note: comparisons were done before and after each individual session)	ABMI	NA
Van Diepen 2002 ³² (35396447) United Kingdom Pilot	Snoezelen	Reminiscence therapy	Day hospital; single site 15 PLWD	N=15 Mean MMSE 7.7	N=15 Sex: no Age: no Race: no Education: no	None	NA	NA	NA	NA	8 weeks post intervention	CMAI ABMI	NA

Baker 1997 ³³ (12519587) United Kingdom Small sample	Snoezelen	Activity sessions	Day hospital; multisite 31 PLWD	Mean MMSE approximately 6	N=31 Sex: yes Age: no Race: no Education: no	None	NA	NA	NA	NA	8 weeks post intervention	REHAB deviant behavior REHAB speech BRS social disturbance	NA
--	-----------	-------------------	------------------------------------	---------------------------	--	------	----	----	----	----	---------------------------	--	----

Abbreviations: ABMI=Agitation Behavior Mapping Instrument; ADL=Activities of Daily Living ; AMPS=Assessment of Motor and Process Skills; BIP=Behavior Observation Scale for Psychogeriatric In-Patients; BRS=Behavior Rating Scale; Char=characteristics; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; DSM=Diagnostic and Statistical Manual; EM=Evidence Map; FC=formal caregiver; GDS=Global Deterioration Scale; IC=informal caregiver; MAI=Multi-level Assessment Instrument; MMSE=Mini-mental State Exam; N=number; NA=Not Applicable; NR=Not Reported; PAS=Pittsburgh Agitation Scale; PMID=PubMed Identification Number; PLWD=Persons Living with Dementia; RADL=Refined Activities of Daily Living Assessment Scale; REHAB=Rehabilitation Evaluation Hall and Baker; RS=Reporting Status; SANS-AD=Scale for the Assessment of Negative Symptoms in Alzheimer’s Disease; SES=socioeconomic status;

Complementary and Alternative Medicine (CAM) Therapy

Table C-12. Risk of bias assessment: CAM therapy

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Overall Rating
Kwan 2017 ³⁴ (25346026)	8 weeks	Low 8 weeks: 0%	Medium	High	High	Low	Medium	High
Yang 2016 ³⁵ (27319407)	8 weeks	Low 8 weeks: 5%	Low	Medium	High	Low	Medium	Medium
Yang 2015 ³⁶ (25880034)	3 weeks	High 3 weeks: 33%	X	X	X	X	X	High
Moyle 2014 ³⁷ (24216598)	3 weeks	Low 3 weeks: 3.64%	Low	Low	Low	Low	Low	Low
Burns 2011 ³⁸ (21335973)	4 weeks	Medium 4 weeks: 16%	Low	Medium	Low	Low	Low	Medium
Lin 2007 ³⁹ (17342790)	8 weeks	Low 8 weeks: 0%	Low	Medium	Medium	Low	Low	Medium
Ballard 2002 ⁴⁰ (12143909)	4 weeks	Low 4 weeks: 1.4%	Low	Low	Medium	High	Low	Medium

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NA=Not Applicable; PMID=PubMed Identification

Table C-13. Characteristics of included studies: CAM therapy

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Yang 2016 ³⁵ (27319407) Taiwan Medium Explanatory	PLWD; got 30 minutes of aromatherapy massage with lavender oil and orange oil once per week for 8 weeks f	PLWD, Usual Care	Long-term care facilities in Taiwan RCT 59 PLWD	Mild to severe dementia based on (score ≤8) (SPMSQ; or (MMSE);	N=29 61% Female 92 years Race NR High school education or higher: 38% Use of psychotropic meds: 70%	None	NA	NA	NA	NA	5 weeks 9 weeks	CCMAI CSDD-C	NA	
Moyle 2014 ³⁷ (24216598) Australia Low Explanatory	PLWD, Trained massage therapists provided 5 minutes massage on each foot using unscented sorbolene for 3 weeks	PLWD, Trained assistants sat quietly near the participant's feet for 10 minutes without talking or making physical contact or any deliberate touching or conversation with the participant	Long-term care facilities in South Queensland Crossover RCT 55 PLWD	Moderate to late stage dementia, MMSE score of less than 18 with a recent history of agitation (PAS >3);	N=55 66% Female 86.5 years Race NR Education NR	None	NA	NA	NA	NA	3 weeks	CMAI OERS	NA	

Burns 2011 ³⁸ (21335973) The United Kingdom (UK) Medium Explanatory	PLWD; 10% Melissa (lemon-balm) oil mixed with base lotion was gently massaged on hands and upper-arms twice a day	PLWD; Sunflower oil mixed with base lotion was gently massaged on hands and upper arms twice a day	Clinical centers – Manchester, London and Southampton, UK RCT 77 PLWD	PLWD with agitation for a minimum of 4 weeks, a score of 13.9 on the CMAI, and satisfied the NINCDS/ADRDA criteria for probable or possible Alzheimer's disease; or was a resident in a nursing home or NHS continuing care facility, had a clinical dementia rating of 3, age over 60 years and were free of psychotropic medication (antipsychotics and/ or cholinesterase inhibitors) for at least 2 weeks	N=38 43% Female 85 years Race NR Education NR	None	NA	NA	NA	NA	4 weeks 12 weeks	PAS NPI Blau-QoL Barthel index	NA
Lin 2007 ³⁹ (17342790) Hong Kong Medium Explanatory	PLWD, lavender oil inhalation for 3 weeks. with a washout period of 2 weeks; 2 drops of oil placed on cotton using dropper in an aromatic diffuser. Two such diffusers are placed at each side of pillow for 1 h every night	PLWD, Sunflower oil inhalation for 3 weeks. with a washout period of 2 weeks; 2 drops of oil placed on cotton using dropper in an aromatic diffuser. Two such diffusers are placed at each side of pillow for 1 h every night	Care and attention homes in Hong Kong Crossover RCT 70 PLWD	Dementia diagnosis based on DSM-IV, APA 1994 with clinically significant agitation evaluated using CCMAI	N=35 41% Female 82 years Race NR Education NR Use of psychotropic meds: 51%	None	NA	NA	NA	NA	3 weeks 5 weeks 8 weeks	CCMAI CNPI	NA
Ballard 2002 ⁴⁰ (12143909) The United Kingdom Medium Explanatory	PLWD; 10% Melissa (lemon-balm) oil mixed with base lotion was gently applied (0.16-0.17g) onto PLWD face and arms, twice a day for 6 doses per day by care assistants	PLWD; Sunflower oil mixed with base lotion was gently applied (0.16-0.17g) onto PLWD face and arms, twice a day for 6 doses per day by care assistants	National Health Service nursing homes in UK RCT 72 PLWD	Severe dementia with agitation characterized by CMAI and also screened by NPI and CDR	N=36 60% Female 78.5 years Race NR Education NR Use of psychotropic meds: 92%	None	NA	NA	NA	NA	4 weeks	CMAI NPI	NA

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: APA=American Psychiatric Association; Barthel index=Barthel scale of Activities of Daily Living; char=characteristics; FC=formal caregiver; IC=informal caregiver; CCMAI=Cohen-Mansfield Agitation Inventory-Chinese version; CDR=Clinical dementia rating scale; CNPI=Neuropsychiatric Inventory-Chinese version; CSDD-C=Cornell Scale for Depression in Dementia-Chinese Version; DSM-IV=Diagnostic and Statistical Manual of Mental Disorders, 4th edition; MMSE=Mini-Mental State Examination; g=grams; h=Hour; N=Number; NHS=National Health

Service; NINCDS/ADRDA=National Institute of Neurological and Communicative Disorders and Stroke/ Alzheimer's Disease and Related Disorders Association; NA=Not applicable; NR=Not reported; OERS=Observed Emotion Rating Scale; PAS=Pittsburgh Agitation Scale; PMID=PubMed Identification Number; PLWD=Persons Living with Dementia; QoL=Quality of life; RCT=Randomized controlled trial; RoB=Risk of Bias; RS=Reporting Status; SPMSQ=Short Portable Mental Status Questionnaire

Table C-14. PLWD outcomes summary low and medium risk of bias studies: CAM therapy

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Overall p-Value	Intervention	p-Value	Comparator	p-Value
Yang 2016 ³⁵ (27319407) Lavender and orange oil vs. Usual care Medium Explanatory	Mean Difference (95% CI) CCMAI 5 weeks	NR	0.316	3.65 (NR)	NR	7.08 (NR)	NR
Yang 2016 ³⁵ (27319407) Lavender and orange oil vs. Usual care Medium Explanatory	Mean Difference (95% CI) CCMAI 9 weeks	NR	0.316	3.41 (NR)	NR	6.3 (NR)	NR
Yang 2016 ³⁵ (27319407) Lavender and orange oil vs. Usual care Medium Explanatory	Mean Difference (95% CI) CSDD-C 5 weeks	NR	<0.001	0.51 (NR)	NR	3.04 (NR)	NR
Yang 2016 ³⁵ (27319407) Lavender and orange oil vs. Usual care Medium Explanatory	Mean Difference (95% CI) CSDD-C 9 weeks	NR	<0.001	0.62 (NR)	NR	6.45 (NR)	NR
Lin 2007 ³⁹ (17342790) Lavender oil vs. sunflower oil Medium Explanatory	Mean Difference (95% CI) CNPI 3 weeks	NR	NR	6.91 (NR)	<0.001	-0.08 (NR)	p=0.24
Lin 2007 ³⁹ (17342790) Lavender oil vs. sunflower oil Medium Explanatory	Mean Difference (95% CI) CCMAI 3 weeks	NR	NR	4.4 (NR)	<0.001	0.04 (NR)	p=0.52
Burns 2011 ³⁸ (21335973) Melissa oil vs sunflower oil Medium Explanatory	Median Difference (95% CI) PAS 4 weeks	NR	0.94	0.0 (-1.3, 0.3)	NR	-0.3 (-1.7, 0)	NR
Burns 2011 ³⁸ (21335973) Melissa oil vs sunflower oil Medium Explanatory	Median Difference (95% CI) PAS 12 weeks	NR	0.56	-0.7 (-1.7, 0)	NR	-0.7 (-1.7, 0)	NR
Burns 2011 ³⁸ (21335973) Melissa oil vs sunflower oil Medium Explanatory	Mean Difference (95% CI) NPI 4 weeks	NR	0.54	-4.8 (-11.3, 1.6)	NR	-9.8 (-18.8, 1.3)	NR
Burns 2011 ³⁸ (21335973) Melissa oil vs sunflower oil Medium Explanatory	Mean Difference (95% CI) NPI 12 weeks	NR	0.52	-7.2 (-12.6, -1.7)	NR	-10.0 (-17.2, -3)	NR
Burns 2011 ³⁸ (21335973) Melissa oil vs sunflower oil Medium Explanatory	Mean Difference (95% CI) QoL 4 weeks	NR	0.24	14 (-15, 42)	NR	-12.0 (-42, 18)	NR
Burns 2011 ³⁸ (21335973) Melissa oil vs sunflower oil Medium Explanatory	Mean Difference (95% CI) QoL 12 weeks	NR	0.033	17.0 (-13, 47)	NR	-2.0 (-34, 30)	NR

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Overall p-Value	Intervention	p-Value	Comparator	p-Value
Burns 2011 ³⁸ (21335973) Melissa oil vs sunflower oil Medium Explanatory	Mean Difference (95% CI) Barthel index 4 weeks	NR	0.91	0.2 (-0.7, 1)	NR	-0.1 (-0.9, 0.7)	NR
Burns 2011 ³⁸ (21335973) Melissa oil vs sunflower oil Medium Explanatory	Mean Difference (95% CI) Barthel index 12 weeks	NR	0.72	-0.8 (-1.8, 0.1)	NR	-0.3 (-1.2, 0.6)	NR
Ballard 2002 ⁴⁰ (12143909) Melissa oil vs sunflower oil Medium Explanatory	Mean Difference (95% CI) CMAI 4 weeks	NR	NR	23.1 (NR)	<0.0001	7.3 (NR)	p=0.005
Moyle 2014 ³⁷ (24216598) Foot massage vs quiet presence Low Explanatory	Mean Difference (95% CI) OERS 3 weeks	NR	NR	NR	NR	NR	NR
Moyle 2014 ³⁷ (24216598) Foot massage vs quiet presence Low Explanatory	Mean Difference (95% CI) CMAI 3 weeks	NR	0.03	-1.28 (NR)	NR	-7.79 (NR)	NR

Abbreviations: Barthel index=Barthel scale of Activities of Daily living; CMAI=Cohen-Mansfield Agitation Inventory-Chinese version; Living; CI=Confidence interval; CNPI=Neuropsychiatric Inventory-Chinese version; CSDD-C=Cornell Scale for Depression in Dementia-Chinese Version; NR=Not Reported; NPI=Neuropsychiatric Inventory; OERS=Observed Emotion Rating Scale; PAS=Pittsburgh Agitation Scale; PMID=PubMed Identification Number; QoL=Quality of life; RoB=Risk of Bias

Table C-15. Summary of strength of evidence for PLWD outcomes: CAM therapy

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Lavender vs sunflower oil Agitation	3 weeks	1 RCT (n=70)	Intervention favors for agitation reduction.	Moderate	Unknown	Direct	Imprecise	Insufficient
Lavender vs sunflower oil NPI	3 weeks	1 RCT (n=70)	Intervention favors for reduction of behavioral disturbances measured using NPI.	Moderate	Unknown	Direct	Imprecise	Insufficient
Lavender and orange oil vs. Usual care Agitation	5 weeks 9 weeks	1 RCT (n=59)	No difference between intervention and placebo groups in measures of agitation.	Moderate	Unknown	Direct	Imprecise	Insufficient
Lavender and orange oil vs. Usual care Depression	5 weeks 9 weeks	1 RCT (n=59)	Intervention favors for depression reduction	Moderate	Unknown	Direct	Imprecise	Insufficient
Melissa oil vs sunflower oil Agitation	4 weeks 12 weeks	2 RCT (n=149)	Mixed results with 1 RCT favoring intervention (4 weeks) while the other one (12 weeks) shows no difference between intervention and placebo groups	Moderate	Inconsistent	Direct	Imprecise	Insufficient
Melissa oil vs sunflower oil NPI	12 weeks	1 RCT (n=77)	No difference between intervention and placebo groups in reduction of behavioral disturbances measured using NPI	Moderate	Inconsistent	Direct	Imprecise	Insufficient
Melissa oil vs sunflower oil QoL	12 weeks	1 RCT (n=77)	Intervention favors in improvement of QoL	Moderate	Inconsistent	Direct	Imprecise	Insufficient
Melissa oil vs sunflower oil Barthel index	12 weeks	1 RCT (n=77)	No difference between intervention and placebo groups in Barthel index	Moderate	Inconsistent	Direct	Imprecise	Insufficient
Foot massage vs quiet presence Agitation	3 weeks	1 RCT (n=55)	Both Intervention and comparison causes an increase in agitation, but the increase is greater in the comparator group than in the intervention group (p=0.03)	Low	Unknown	Direct	Imprecise	Insufficient

Abbreviations: Barthel index=Barthel scale of Activities of Daily living; n=Number; NPI=Neuro psychiatry inventory; RCT=Randomized controlled trial; QoL=Quality of life

Evidence Map: CAM Therapy

Table C-16. Characteristics of evidence map studies: CAM therapy

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Kouzuki 2019 ⁴¹ (31578055) Japan Small sample	0.1% aroma bath salts (n=15) 0.5% aroma bath salts (n=14) 1% aroma bath salts (n=14) where aroma oil added to the bath salt was a 2:1 blend of true lavender and sweet orange and baths were taken once daily (>10 minutes) for 24 weeks	NA	Outpatient clinic of Shinsei Hospital (Kurayoshi, Japan) 3-arm RCT 43 PLWD	AD diagnosis according DSM-V or MCI that met Petersen's diagnostic criteria	N=43 Mean age: 79 years Female: 51% % majority race: no Education: no	None	NA	None	NA	NA	4 weeks 16 weeks 28 weeks 32 weeks	TDAS OSIT-J PSQI-J	NA
Takahashi 2019 ⁴² (32037737) Japan Small samples	Aromatherapy: Low pressure, 50% ethanol extract of Akita cedar leaves diffused in rattan sticks and as spray for 4 weeks (The room fragrance type is placed in the resident space (living room and bedroom), and the spray type is used to mist the patients' clothing and bedding.)	Control group: Ethanol without cedar leaves extract	PLWDs RCT 36 PLWD	Diagnosis of AD	N=36 Age: yes Sex: yes Race: no Education: yes	None	NA	None	NA	NA	8 weeks	FAST NPI ADAS-cog J-ZBI	NA

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Watson 2019 (30670268) Australia Small sample	1.Aromatherapy with Lavender oil 2.Aromatherapy with Lemon-balm oil 2 drops of oil was applied to a cotton patch and then placed on the participant's collar area. The patch was removed after 2 hours for 2 weeks. Washout period of 2 weeks were maintained	3. Same procedure was followed with sunflower oil	Residents of residential aged care facility RCT 39 PLWD	Diagnosis of dementia and MMSE score of ≤10; agitated behaviors recorded on at least one of ACFI domains;	N=39 Mean age: no % Female: no % majority race: no Education: no	None	NA	None	NA	NA	0 week 2 weeks 4 weeks 6 weeks 8 weeks 10 weeks	CMAI NPI	NA
Zalomonson 2019 ⁴³ (30328781) Israel Pilot	2 drops of lavender oil were applied by the study nurse for no more than 1-2 minutes, 3 times a day for 16 weeks	Same procedure using sunflower oil	Psychogeriatric long-term care departments at the Center for Mental Health of the Israel Ministry of Health, Beer-Sheva Crossover Placebo-Controlled Study 42 PLWD	Diagnosis of dementia of any type	N=42 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	4 weeks 8 weeks 12 weeks 16 weeks	NPI	NA
Fung 2018 ⁴⁴ (28986942) Hong-Kong Small sample	Aroma-massage with acupressure: Trained carers or therapists provided no more than 20 minutes of aromatherapy, combining	Cognitive training: Trained care staff or therapists provided 20 minutes of cognitive training (n=20)	Long-term non-government geriatric care facilities 3 arm RCT 60 PLWD	BPSD PLWD scored below 18 (illiterate), 19 (1-2 years of education) and 20 (> 2years education) in CMMSE	N=20 Age: yes Sex: yes Race: yes Education: yes	None	NA	NA	NA	NA	Unclear 3-months Follow-up	CMMSE CCMAI NPI BI20	NA

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
	therapeutic massage with blended 2% lavender essential oil plus sunflower-oil. Acupressure on the selected acu-points was integrated into the massage treatment in a biweekly session.	Exercise: 20 minutes of stretching exercises, as outlined by a physiotherapist. (n=20)											
Kwan 2017 ³⁴ (25346026) Hong Kong High RoB Kwan 2014 ⁴⁵ (25346026) Hong Kong Study Protocol	9 minutes acupressure sessions	Sham acupressure; Usual care	Residential care homes 3 arm RCT 119 PLWD	PLWD showing dementia in medical records and displayed agitated behavior	N=119 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	2 weeks	CMAI Salivary cortisol	NA
Jia 2017 ⁴⁶ (29284465) China Pilot	Acupuncture was performed using needles of 0.25 mm and a length of 40 mm at various acu-points for 3 times a week for 12 weeks	5 mg/day of donepezil hydrochloride for the first 4 weeks and 10 mg/day thereafter	Community residents (older PLWD with dementia) who participated in survey from Tianjin City, China RCT 87 PLWD	Diagnosis based on DSM-IV and NINCDS-ADRDA with MMSE score between 10 and 23 and HAMD score<7, a HIS score≤4, and have a reliable caregiver to accompany the participant to all study visits	N=43 Age: yes Sex: yes Race: no Education: yes	None	NA	NA	NA	NA	16 weeks 28 weeks	ADAS-cog CIBIC-Plus ADCS-ADL ₂₃ NPI	NA
Moorman Li 2016 ⁴⁷ (28265482) US Pilot	Lavender aromatherapy twice a day for 20 min during a two-month period during active clinic days.	None	Adult Day Care Center quasi-experimental 23 PLWD	Clinical diagnosis of dementia	N=23 Age: yes Sex: yes Race: yes Education: no	None	NA	NA	NA	NA	Daily for 2 months pre-observation phase and 2 months intervention period	Behavior/Intervention Monthly Flow Record	NA
Yang 2015 ⁴⁸ (26556080) Taiwan High RoB	Aroma-acupressure: each acupoint was pressed for	Control: Received daily routine as usual	Retirement homes for veterans and long-term care	Diagnosis according to DSM-IV criteria	N=82 Age: yes Sex: yes	None	NA	NA	NA	NA	weekly	CCMAI HRV index	NA

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
	2 minutes with 2.5% lavender oil and a warm-up exercise was completed for 5 minutes, once per day for 5 days for 4 weeks Aromatherapy: 2.5% lavender oil was applied at five acupoints with the same operation time as the aroma-acupressure group(n=106)	without intervention (n=88)	facilities 3 arm RCT 276 PLWD	CMAI>35 (extreme agitation)	Race: no Education: no								
Chieh-Yu 2013 ⁴⁹ (23837414) Australia Small sample	Aromatherapy and hand massage (n=22): Three sprays of lavender mist (3% lavender mist) applied on chest twice a day, for 6 weeks and 5 minutes of hand massage twice a day for 10 days: each hand was massaged for 2.5 minutes. Aromatherapy only (n=23): Three sprays of lavender mist (3% lavender mist) applied on chest twice a day, for 6 weeks	Placebo (n=22) water spray similar to aromatherapy	Long term care facilities in Brisbane 3 arm RCT 67 PLWD	MMSE score ≤ 24 and AD diagnosis according DSM-IV with a documented history of a minimum of two weeks of agitation or aggression within the past three months, requiring physical and/or chemical restraint	N=67 Mean age: 84 years Female: 59% % majority race: no Education: no	None	NA	None	NA	NA	6 weeks	MMSE CMAI-SF	NA
Fu 2013 ⁵⁰ (23837414)	3 sprays of lavender mist	Water mist sprayed in	Long-term care facilities	DSM-IV criteria (confirmed by a	N=23 Age: yes	None	NA	NA	NA	NA	3 months	Disruptive behavior:	NA

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Australia Small sample	(3%) twice daily 7 days a week for 6 weeks 3 sprays of lavender mist (3%) Plus 5 minutes of hand massage (n=22)	the same way (n=22).	3 arm RCT 67 PLWD	medical practitioner and a university Alzheimer's clinic) Moderate to severe-MMSE	Sex: yes Race: yes Education: no							CMAI-SF Cognition: (MMSE)	
Lu 2013 ⁵¹ (23972540) USA Pilot	Along with usual medical and care regimen, HT and BTC protocols were given for the 6 months treatment period on weekly basis	PLWD got usual medical and care regimens	Recruited from long-term nursing care facilities and from the community RCT 22 PLWD	Early AD (information regarding their doctor's diagnosis was acquired from the subjects, their caregivers, or nursing staff)	N=12 Age: yes Sex: yes Race: yes Education: no	None	NA	None	NA	NA	3 months 6 months	MoCA POMS-BF PHQ-9	NA
Rodriguez-Mansilla 2013 ⁵² (24187866) Spain Pilot Rodriguez-Mansilla 2015 ⁵³ (25322869) Spain Pilot	20 minutes of back and lower limb massage with moisturizing cream; Ear-acupuncture at various points (n=40)	Control (did not receive any experimental treatment n=40)	Residents institutionalized in 'CARE' elderly residential homes in Extremadura 3-arm-RCT 120 PLWD	PLWD were diagnosed using the DSM-VI criteria	N=40 Age: yes Sex: yes Race: no Education yes	None	NA	NA	NA	NA	12 weeks	Behavioral alteration and sleep disturbances: Self-designed structured questionnaire with closed ended questions GDS MMSE DOLOPLUS2 CSDD Campbell scale	NA
Jimbo 2009 ⁵⁴ (20377818) Japan Pilot	Aroma exposure of 0.04 ml lemon and 0.08 ml rosemary oil in the morning for 2 hours. 0.08 ml Lavender and 0.04 ml orange oil in the evening for 1.5	None	Setting: NR Quasi-experimental (crossover) 28 PLWD	Diagnosis based on DSM-IV and NINCDS-ADRDA or DSM-IV and NINCDS-AIREN.	N=28 Age: yes Sex yes Race: no Education: No	None	NA	NA	NA	NA	4 weeks 9 weeks	HDS-R GBSS-J FAST CT Scans TDAS Zarit Blood and biochemical examination	NA

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
	hours for 4 weeks												
Hawranik 2008 ⁵⁵ (18272750) Canada Small sample	TT once per day for 5 days	Simulated TT once per day for 5 days (n=16) Usual care for 5 days (n=18)	Residents from personal care and special needs units of one long-term care facility 3 arm RCT 51 PLWD	Diagnosis of senile dementia of the Alzheimer type, score 23 or less in MMSE; with a history or consistent agitated behavior during last 30 days	N=17 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	2 hours; 24 hours 1 week 2 weeks	CMAI MMSE	NA
Woods 2005 ⁵⁶ (15712768) Canada Small sample	TT was performed with contact on the neck and shoulders through an experienced practitioner for 5-7 minutes twice daily for 3 days	Sham TT (n=19) Routine care (n=19)	Long-term care facilities 3-arm-RCT 57 PLWD	Diagnosis of moderate to severe AD according to DSM-IV criteria; MMSE<20;	N=19 Age yes Sex: yes Race: yes Education: yes	None	NA	NA	NA	NA	3 days	RPMB ABRS	NA
Smallwood 2001 ⁵⁷ (11607948) UK Small sample	Aromatherapy massage Conversation and aromatherapy Massage only for twice weekly	None	PLWD were recruited from district general hospital ward 3-arm RCT 21 PLWD	Clinical diagnosis of dementia	N=21 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	weekly	Qualitative assessment of behavior using video recorded sessions	NA

Abbreviations: ABRS=Agitated Behavior Rating Scale; ACFI=Aged Care Funding Instrument; ADAS-cog=Alzheimer's disease Assessment Scale-Cognitive; AD=Alzheimer's Disease; ADCS-ADL₂₃=23-Item Alzheimer's disease Cooperative Study Activities of Daily Living Scales; BI20=Barthel Index-20; BPSD=Behavioral and Psychological Symptoms of Dementia; BTC=Body Talk Cortices; CMAI=Cohen-Mansfield Agitation Inventory-Chinese version; CIBIC-Plus=Clinician's Interview-Based Impression of Change-Plus; CMAI-SF=Cohen-Mansfield Agitation Inventory-Short Form; CMMSE=Mini-Mental State Examination -Chinese version; CSDD=Cornell Scale for depression in dementia; CT=Computed Tomography; DOLOPLUS2=Behavioral pain assessment scale for the elderly presenting with verbal communication disorders; DSM-IV=Diagnostic and Statistical Manual of Mental Disorders, 4th edition; FAST=Functional Assessment Staging of Alzheimer's disease; GBSS-J=Gottfries, Brane, Steen Scale; GDS=Global Deterioration Scale; HDS-R=Hasegawa's Dementia Scale; HRV-index=Heart Rate Variability Index; HT=Healing Touch; J-ZBI=Japanese version of Zarit Caregiver Burden interview; mg=milligram; ml=milliliter; mm=millimeter; MMSE=Mini-Mental State Examination; MoCA=Montreal Cognitive Assessment test; N=Number; NINCDS/ADRDA=National Institute of Neurological and Communicative Disorders and Stroke/ Alzheimer's Disease and Related Disorders Association; NINCDS/AIREN=National Institute of Neurological Disorders and Stroke/ Association Internationale pour la Recherche et l'Enseignement en Neurosciences criteria; NPI=Neuro psychiatry inventory; NA=Not Applicable; PHQ-9=Patient health questionnaire-9; PMID=PubMed Identification Number; POMS-BF=Profile of Mood States-Brief Form; PLWD=Persons Living with Dementia; RCT=Randomized controlled trial; RoB=Risk of Bias; RPMB=The Revised Memory and Behavior Checklist; T-DAS=Touch Panel-type Dementia Assessment Scale; TT=Therapeutic touch
Bright Light Therapy

Bright Light Therapy

Table C-17. Risk of bias assessment: bright light

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Onega 2016 ⁵⁸	8 weeks	Medium 15%	Low	High	Low	Medium	X	Government	High
Rixt F. Riemersma-van der Lek 2008 (18544724) ⁵⁹	6 weeks	Low 6 weeks:7.4% High 6 months+: >21%	Low	6 weeks: Medium 6+ Months: High	Low	High	Medium	Government	High
Hickman 2007 ⁶⁰ Barrick 2010 (20104513) ⁶¹	3 weeks	Medium NR	Medium	High	Medium	X	X	Government	High
Ancoli-Israel 2003 ⁶²	18 days	High 23.9%	X	X	X	X	X	Government	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Bright Light Therapy

Table C-18. Characteristics of evidence map studies: bright light

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Figueiro, 2019 ⁶³ (318551610) United States Small sample	Tailored lighting to maximally affect circadian system	Low intensity lights	4 Assisted Living Facilities RCT 52 PLWD	Dementia per MMSE; unspecified	N=60 Age: Yes Sex: Yes Race: no Education: no	None	NA	NA	NA	NA		Sleep and circadian rhythms CSDD CMAI	NA
Lisa Onega 2016 ⁵⁸ United States High ROB	Bright light exposure	Low intensity light	Long-term care facility (N=4) RCT (bright light N=30; low intensity light N=30) PLWD	None	N=60 (treatment N=30 vs control N=30) Age: Yes Sex: Yes Race: Yes Education: Yes	SES Health problems Marital status	NA	NA	NA	NA	8-weeks)	Depression (DSAOA; DMAS-17; CSDD) Agitation (CMAI-F; CMAI-D; PAS; BARS)	NA
Burns 2009 ⁶⁴ United Kingdom Small sample	Full spectrum bright light therapy (BLT)	Normal light	Nursing homes RCT (standard light N=26; BLT N=22) PLWD	Dementia and behavioral disturbances	N=48 (22 treatment vs 26 control) Age: Yes Sex: Yes	Cataracts	NA	NA	NA	NA	Baseline week, and in weeks 4 and 8	Agitation (Cohen-Mansfield Agitation Inventory; CMAI)	NA
Riemersma-van der Lek 2008 ⁵⁹ (18544724) Netherlands Medium Explanatory	Bright light therapy, 8 hours daily, exposure of ±1000 lux	Usual care (double placebo)	Group care facility Cluster RCT 12 Long-term care facilities 94 PLWD	Clinical diagnosis for dementia and dementia sub-types	N=94 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	6 weeks	MMSE CSDD PGCMS PGCARS MOSES, Withdrawn Behavior NPI-Q CMAI NIADL Total Sleep Duration	NA
Dowling 2007 ⁶⁵ Dowling 2005 (16050432) ⁶⁶ Dowling 2005 (16035127) ⁶⁷ United States Small sample	Bright light exposure	Usual indoor light	Nursing home RCT (morning light N=29; afternoon light N=24; or control N=17) PLWD	Rest-activity disruption and were diagnosis with Alzheimer's disease	N=71 (54 treatment vs 17 control) Age: Yes Sex: Yes Race: Yes	None	NA	NA	NA	NA	End of the baseline week and after the last week of intervention (study protocol 11 weeks)	Neuropsychiatric behaviors (Neuropsychiatric Inventory nursing Home version; NPI-NH)	NA

Hickman 2007 ⁶⁰ 20104513 Barrick 2010 ⁶¹ United States High ROB	Ambient bright light (high-intensity, low glare) in public areas, applied morning, evening, or all day	Standard lighting	Geriatric unit in 2 state-operated psychiatric hospitals, Cluster-unit crossover intervention trial with 8 study periods in one, 22 in the other ; PLWD	Severe or very severe dementia	N=66 Age: Yes Sex: Yes Race: Yes Education: Yes	None	NA	NA	NA	NA	NA	Last week of each 3-week period (multiple periods)	CSDD	NA
Ancoli-Israel 2003 ⁶² United States High ROB	Bright light exposure	Morning dim red light	Nursing home Randomized to treatment groups (morning bright light N=30; morning dim red light N=31; or evening bright light N=31) PLWD	Diagnosis of possible or probable Alzheimer's disease	N=92 (61 treatment vs 31 control) Age: Yes Sex: Yes Education: No	Current diagnoses (e.g. eye disease; cataracts, glaucoma)	NA	NA	NA	NA	NA	3 days of baseline, 10 days of treatment (treatment days 1-5 and 6-10) and 5 days of posttreatment follow-up	Sleep and circadian activity rhythms	NA
Paola Fontana Gasio 2003 ⁶⁸ Switzerland Pilot	Low intensity dawn-dusk simulation (DDS)	Dim red light	Nursing homes Randomly assigned (DDS N=9; Control N=4) PLWD	Symptoms/diagnosis of dementia and sleep disturbances	N=13 (treatment N=9 vs control N=4) Age: Yes Sex: Yes	None	NA	NA	NA	NA	NA	After 3 weeks of baseline; DDS or dim red light treatment	Circadian rest-activity cycle, nocturnal sleep, and cognitive function	NA
Kazuo Mishima 1998 ⁶⁹ Japan Small sample	Bright light therapy	None	Facility Randomized crossover PLWD	Vascular dementia (VD) and PLWD with dementia of Alzheimer's type (DAT) with sleep and behavioral disturbances	N=22 (VD N=12; DAT N=10) Age: Yes Sex: Yes	None	NA	NA	NA	NA	NA	Pretreatment, week 1, week 2, and posttreatment	Behavioral and rest-activity (R-A) rhythm disorders	NA

Abbreviations: BARS=Brief Agitation Rating Scale; CMAI=Cohen-Mansfield Agitation Inventory; CMAI-D=Cohen-Mansfield agitation Inventory-Disruptiveness; ; CMAI-F=Cohen-Mansfield Agitation Inventory-Frequency; CSDD=the Cornell Scale for Depression in Dementia; DSAOA=Depressive Symptom Assessment in Older Adults; DMAS-17=Dementia Mood Assessment Scale-17; FC=formal caregiver; IC=informal caregiver; N=number; NPI-Q=the Neuropsychiatric Inventory; NI-ADL=the nurse-informant adaptation; MOSES=the Multidimensional Observation Scale for Elderly Subjects; NPI-NH=Neuropsychiatric Inventory nursing Home version; PAS=Pittsburgh Agitation Scale; PGCMS=the Philadelphia Geriatric Centre Morale Scale; PGCARS=The Philadelphia Geriatric Centre Affect Rating Scale; PLWD=Person Living with Dementia; PMID=PubMed Identification number; SES=socioeconomic status

Psychosocial Interventions for BPSD

Table C-19. Risk of bias assessment: psychosocial interventions for BPSD

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Overall Rating
Richards 2005 (16137280) ⁷⁰	3 weeks	Low 5%	Medium	Medium	Medium	Low	High	Medium
Toseland 1997 (27097884) ⁷¹	3 months, 1 year	Medium 25%	Medium	High (30% missing data for bed/rise time)	Medium	Low	Low	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Psychosocial Interventions for BPSD

Table C-20. Characteristics of evidence map studies: psychosocial interventions for BPSD

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Kiossis 2015 ⁷² (26402009) US Small sample	Problem adaptation therapy based on process model of emotion regulation	Supportive therapy	Community setting, in-home RCT PLWD	Mild to moderate dementia and major depression; dementia unspecified	N=39 Age: no Sex: yes Race: yes Education: no	None	NA	NA	NA	NA	12 weeks	CSDD Suicidal ideation	NA
Spector 2015 ⁷³ (25698766) (26207801) ⁷⁴ UK Pilot	Cognitive behavioral training for anxiety; up to 10 1-hour sessions	Usual care	Community setting, in-home RCT PLWD	Mild to moderate dementia; dementia unspecified	N=50 Age: yes Sex: yes Race: yes Education: yes	None	NA	NA	NA	NA	15 weeks	RAID CSDD HADS QOL-AD NPI QCPR CSRI MMSE	NA
Stanley 2013 ⁷⁵ (25698766) US Pilot	Cognitive behavioral training for anxiety; 12 weekly session followed by 8 brief telephone sessions	Usual care	Community-based, in-home RCT PLWD with caregiver	Mild to moderate dementia; dementia unspecified	N=32 Age: yes Sex: yes Race: yes Education: no	Marital status	NR	NR	NA	NA	6 months	NPI RAID QOL-AD PSWQ-A GAI	Caregiver distress PHQ-9
Richards 2005 ⁷⁰ (16137280) US High ROB	Individualized social activity intervention to improve sleep	Usual care	Nursing home RCT PLWD	Moderate to severe dementia; less than 85% sleep efficiency	N=147 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	3 weeks	Sleep measures	NA

Beck 2002 ⁷⁶ (12131234) US Small sample	ADL: 45-60 minutes of ADL assistance provided by PNA which catered to the psychosocial, cognitive and physical abilities of PLWD PSA: 30+ minutes of PSA intervention provided by PNA, which involved 25 standardized modules designed to meet the psychosocial, cognitive and physical abilities of PLWD CB: consisted of both the ADL and PSA interventions and lasted 90 minutes daily	Placebo: one-to-one interaction between the participant and PNA for 30 minutes per day No intervention: routine care with no scheduled contact between participants and the PNA	Nursing home 5 arm RCT PLWD	Dementia diagnosis with a MMSE score \leq 20 and report of DB in previous 2 weeks	N=127 Age: yes Sex: yes Race: n Education: no	None	NA	NA	NA	NA	Videotape analysis 1-month follow-up 2-month follow-up	DBS MMSE ODAS AARS PVAS	NA
Toseland 1997 ⁷¹ (27097884) US High ROB	Validation therapy or social contact; 4 30-minute sessions per week for 52 weeks	Usual care	Nursing home 3 arm RCT PLWD	Moderate to severe dementia with behavioral problems; dementia unspecified	N=88 Age: yes Sex: yes Race: yes Education: no	None	NA	NA	NA	NA	3 months, 1 year	MOSES CMAI GIPB MDS+	NR

Abbreviations: CMAI=Cohen-Mansfield Agitation Inventory; CSDD=the Cornell Scale for Depression in Dementia; FC=formal caregiver; IC=informal caregiver; GAI=Geriatric Anxiety Inventory; GDS=Geriatric Depression Scale; GIPB=Geriatric Indices of Positive Behavior; HADS=Hospital Anxiety and Depression Scale; MMSE=Mini-Mental State Examination; MOSES=Multidimensional Observation Scale for Elderly Subjects; MDS+=Minimum Data Set-Resident Assessment Protocol; N=number; NPI=the Neuropsychiatric Inventory; PHQ-9=Patient Health Questionnaire; PSWQ-A=Penn State Worry Questionnaire-Abbreviated; QCPR=Quality of Caregiver and Patient Relationship; QOL-AD=Quality of Life – Alzheimer’s Disease; PLWD=Person Living with Dementia; PMID=PubMed Identification number; RAID=Rating Anxiety in Dementia scale; SES=socioeconomic status

Multicomponent Interventions for BPSD

Table C-21. Risk of bias assessment: multicomponent interventions for BPSD

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Overall Rating
Lin 2009 ⁷⁷ (19507295)	6 weeks, 14 weeks; 21-22 weeks	High Unclear	Medium	High	High	Low	Medium	High
Finnema 2005 ⁷⁸ (15799079)	3 months 7 months	High 25%	Medium	X	X	X	X	High
Kovach 2004 ⁷⁹ (15611216)	Unclear	High unclear 24%	Low	X	X	X	X	High
Schrijnemaekers 2002 ⁸⁰ (12325052)	3, 6, and 12 months	Medium 3 month: 9%, 6 months: 20.5%, 12 months: 38%	Medium	Medium	Medium	Low	High	High
Lawton 1998 ⁸¹ (9924833)	6 months 12 months	Low 12 months: 5%	Medium	Medium	High	Low	Low	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Multicomponent Interventions for BPSD

Table C-22. Characteristics of evidence map studies: multicomponent interventions for BPSD

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Law 2019 ⁸² (31390090) Hong Kong Pilot	Osteoarthritis-specific knee exercises plus self-management support, 8 weeks	Routine group exercise	Community setting RCT PLWD	Dementia unspecified plus knee osteoarthritis	N=56 Age: yes Sex: yes Race: no Education: yes	Marital status	N=56 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration	NA	NA	8 weeks	NPI-Q	RSCSE
Bailey 2017 ⁸³ (26912731) US Small sample	Two 30 minutes group activity sessions, including, CBT, reminiscence, environmental supports and individualized activity for 6 weeks	Usual care (n=25)	Privately owned, for-profit urban nursing homes Cluster RCT 51 PLWD	Mild to moderate cognitive impairment (MMSE score 10-24) and symptoms of depression (GDS)	N=51 Age: yes Sex: yes Race: no Education: no	None	NA	None	NA	NA	Twice per week for 6 weeks	ABC CSDD GDS QOL-AD Activity enjoyment rating Barthel Scale MMSE	NA
Fischer-Terworth 2011 ⁸⁴ (No PMID) Germany Small sample (<25 in each arm)	TEACCH- and Music Therapy-Based Interventions (TMI): modified cognitive engagement for autism plus group music therapy for 6 months.	Nonspecific occupational therapy: Participated in nonspecific occupational therapy	Special dementia care unit Non-RCT 49 PLWD	Mild or moderate dementia (All types)	N=49 Age: No Sex: Yes Race: No Education: No	None	NA	NA	NA	NA	6 months	NPI GDS ICEA-D	NA

Lin 2009 ⁷⁷ (19507295) Taiwan High RoB	I1: Acupressure: acupressure at each acupoint for 2 minutes; 15-minute daily sessions 6 days per week; 4 weeks I2: Montessori based activity program: five categories associated with activities of daily living: scooping, pouring, squeezing, fine motor skills, environmental care, plus personal care; 6 days per week; 4 weeks	Attention control (Presence): for a 15-minute period each day for 6 days a week	Long-term care facilities 3 arm crossover RCT 133 PLWD	Diagnosis of dementia, CMAI score 35 or above	N=23 Age: yes Sex: yes Race: yes Education: yes	None	NA	NA	NA	NA	8 weeks 16 weeks 22 weeks	CMAI Ease of care AARS Family-visit restraint	NA
Kovach 2004 ⁷⁹ (15611216) US High RoB	BACE intervention: assessment; diagnose and plan a correction of the arousal imbalance; implement new activity schedule	Unclear	Long-term care facilities Pretest posttest RCT 102 PLWD	MMSE score of 15 or below and FAST stage 6 or 7 with some agitation but no chronic psychiatric diagnosis other than dementia	N=78 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	Direct observation of participants	MMSE FAST ASD Arousal Balance and Imbalance ABMI Therapeutic Activity	NA
Finnema 2005 ⁷⁸ (15799079) (24337328) ⁸⁵ The Netherlands High ROB	Emotion-oriented care: combination validation therapy, reminiscence, sensory stimulation	Usual care	Nursing homes RCT with matched 16 psychogeriatric wards PLWD	Moderate to severe probably Alzheimer-type dementia, mixed vascular dementia, dementia or amnesic syndrome	N=146 Age: yes Sex: yes Race: no Education: no	Duration institutionalized	NA	NA	N=99 Age: yes Sex: yes Race: no Education: yes	Position	7 months	CSDD GRGS CMAI PGCMS	GHQ-28 Organization and stress scale Absenteeism

Beck 2002 ⁷⁶ (12131234) US Small sample	ADL: 45-60 minutes of ADL assistance from PNA which catered to the psychosocial, cognitive and physical abilities of PLWD PSA: 30+ minutes of PSA from PNA; 25 standardized modules designed to meet the psychosocial, cognitive and physical abilities of PLWD CB: both the ADL and PSA interventions; 90 minutes daily	Placebo: one-to-one interaction between the participant and PNA for 30 minutes per day No intervention: routine care with no scheduled contact between participants and the PNA	Nursing home 5 arm RCT PLWD	Dementia diagnosis with a MMSE score \leq 20 and report of DB in previous 2 weeks	N=127 Age: yes Sex: yes Race: yes Education: no	None	NA	None	NA	NA	Videotape analysis 1-month follow-up 2-month follow-up	DBS MMSE ODAS AARS PVAS	NR
Schrijnemaekers 2002 ⁸⁰ (12325052) The Netherlands High ROB	Emotion-oriented care: combination validation therapy, reminiscence, sensory stimulation	Usual care	Residential homes with structured day care units Cluster RCT 16 units	Moderate to severe cognitive impairment and behavioral problems; dementia unspecified	N=151 Age: yes Sex: yes Race: no Education: no	Duration institutionalized	NA	NA	NR	NR	3, 6, and 12 months	Dutch Behavior Observation Scale for Psychogeriatric Inpatients ADL CMAI GRGS	NR
Lawton 1998 ⁸¹ (9924833) US High RoB	Stimulation-retreat model: which diagnosed, prescribed, and applied a package of care according to individual needs for additional stimulation or relief from stimulation ("retreat")	Unclear	Nursing home residents RCT 102 PLWD	NR	N=49 Age: no Sex: no Race: no Education: yes	None	NA	NA	NA	NA	6 months 12 months	GDS PSMS BEHAVE-AD CMAI MOSES TRS AARS	NA

Abbreviations: AARS=Apparent affect rating scale; ABC=Activity Behavior Checklist; ABMI=Agitation Behavior Mapping Instrument; ADL=Activities of daily living; ASD=Arousal States in Dementia; BACE=Balancing Arousal Controls Excesses; BEHAVE-AD=Behavioral Pathology in Alzheimer's Disease scale; CB=Combined; CBT=Cognitive-behavioral therapy; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; DB=Disruptive behavior(s); DBS=Disruptive behavior scale; ESS=Epworth Sleepiness Scale; FAST=Functional Assessment Staging Tool; GDS=Geriatric Depression Scale; GHQ-28=General Health Questionnaire; GRGS=Geriatric Resident Rating Scale; ICEA-D=Inventory to Assess Communication, Emotional Expression and Activity in Dementia; MMSE=Mini-Mental Status Exam; MOSES=Multidimensional Observation Scale for Elderly Subjects; N=Number; NA=Not Applicable; NR=Not reported; ODAS=Observable displays of affect scale; PGCMS=Philadelphia Geriatric Center Morale Scale; PMID=PubMed Identification Number; PNA=Project nursing assistant(s); PSA=Psychosocial activity; PSMS=Physical Self-Maintenance Scale; PLWD=Persons Living with Dementia; PVAS=Positive visual analogue scale; QOL-AD=Quality of Life-Alzheimer's Disease; RCT=Randomized controlled trial; RMBPC=Revised Memory and Behavior Problems Checklist; RSCSE=Revised Scale for Caregiving Self-Efficacy; RoB=Risk of Bias; SD=Standard deviation; TRS=Therapeutic Recreation Specialist

References for Appendix C

1. Moyle W, Jones CJ, Murfield JE, et al. Use of a Robotic Seal as a Therapeutic Tool to Improve Dementia Symptoms: A Cluster-Randomized Controlled Trial. *J Am Med Dir Assoc*. 2017 Sep 01;18(9):766-73. doi: <https://dx.doi.org/10.1016/j.jamda.2017.03.018>. PMID: 28780395.
2. Jones C, Moyle W, Murfield J, et al. Does Cognitive Impairment and Agitation in Dementia Influence Intervention Effectiveness? Findings From a Cluster-RCT With the Therapeutic Robot, PARO. *J Am Med Dir Assoc*. 2018 Apr 13;13:13. doi: <https://dx.doi.org/10.1016/j.jamda.2018.02.014>. PMID: 29656838.
3. Mervin MC, Moyle W, Jones C, et al. The Cost-Effectiveness of Using PARO, a Therapeutic Robotic Seal, to Reduce Agitation and Medication Use in Dementia: Findings from a Cluster-Randomized Controlled Trial. *J Am Med Dir Assoc*. 2018 Jan 09;09:09. doi: <https://dx.doi.org/10.1016/j.jamda.2017.10.008>. PMID: 29325922.
4. Moyle W, Jones C, Murfield J, et al. Effect of a robotic seal on the motor activity and sleep patterns of older people with dementia, as measured by wearable technology: A cluster-randomised controlled trial. *Maturitas*. 2018 01 Apr;110:10-7. doi: <http://dx.doi.org/10.1016/j.maturitas.2018.01.007>. PMID: 2821060108.
5. Petersen S, Houston S, Qin H, et al. The Utilization of Robotic Pets in Dementia Care. *J Alzheimers Dis*. 2017;55(2):569-74. PMID: 27716673.
6. Olsen C, Pedersen I, Bergland A, et al. Effect of animal-assisted activity on balance and quality of life in home-dwelling persons with dementia. *Geriatr Nurs*. 2016 Jul-Aug;37(4):284-91. doi: <https://dx.doi.org/10.1016/j.gerinurse.2016.04.002>. PMID: 27155968.
7. Olsen C, Pedersen I, Bergland A, et al. Effect of animal-assisted interventions on depression, agitation and quality of life in nursing home residents suffering from cognitive impairment or dementia: a cluster randomized controlled trial. *Int J Geriatr Psychiatry*. 2016 Dec;31(12):1312-21. doi: <https://dx.doi.org/10.1002/gps.4436>. PMID: 26807956.
8. Bemelmans R, Gelderblom GJ, Jonker P, et al. Effectiveness of Robot Paro in Intramural Psychogeriatric Care: A Multicenter Quasi-Experimental Study. *J Am Med Dir Assoc*. 2015 Nov 01;16(11):946-50. doi: <https://dx.doi.org/10.1016/j.jamda.2015.05.007>. PMID: 26115817.
9. Joranson N, Pedersen I, Rokstad AM, et al. Effects on Symptoms of Agitation and Depression in Persons With Dementia Participating in Robot-Assisted Activity: A Cluster-Randomized Controlled Trial. *J Am Med Dir Assoc*. 2015 Oct 01;16(10):867-73. doi: <https://dx.doi.org/10.1016/j.jamda.2015.05.002>. PMID: 26096582.
10. Joranson N, Pedersen I, Rokstad AM, et al. Change in quality of life in older people with dementia participating in Paro-activity: a cluster-randomized controlled trial.[Erratum appears in *J Adv Nurs*. 2017 Jun;73(6):1531; PMID: 28485533]. *J Adv Nurs*. 2016 Dec;72(12):3020-33. doi: <https://dx.doi.org/10.1111/jan.13076>. PMID: 27434512.
11. Travers C, Perkins J, Rand J, et al. An evaluation of dog-assisted therapy for residents of aged care facilities with dementia. *Anthrozoos*. 2013;26(2):213-25. PMID: CN-00871982.
12. Jones C, Moyle W, Murfield J, et al. Does Cognitive Impairment and Agitation in Dementia Influence Intervention Effectiveness? Findings From a Cluster-Randomized-Controlled Trial With the Therapeutic Robot, PARO. *J Am Med Dir Assoc*. 2018 Jul;19(7):623-6. doi: <https://dx.doi.org/10.1016/j.jamda.2018.02.014>. PMID: 29656838.
13. Pu L, Moyle W, Jones C, et al. The Effect of Using PARO for People Living With Dementia and Chronic Pain: A Pilot Randomized Controlled Trial. *J Am Med Dir Assoc*. 2020 Feb 28;28:28. doi: <https://dx.doi.org/10.1016/j.jamda.2020.01.014>. PMID: 32122797.
14. Pu L, Moyle W, Jones C. How people with dementia perceive a therapeutic robot called PARO in relation to their pain and mood: A qualitative study. *J Clin Nurs*. 2020 Feb;29(3-4):437-46. doi: <https://dx.doi.org/10.1111/jocn.15104>. PMID: 31738463.
15. Balzotti A, Filograsso M, Altamura C, et al. Comparison of the efficacy of gesture-verbal treatment and doll therapy for managing neuropsychiatric symptoms in older patients with dementia. *Int J Geriatr Psychiatry*. 2019 09;34(9):1308-15. doi: <https://dx.doi.org/10.1002/gps.4961>. PMID: 30136743.
16. Menna LF, Santaniello A, Gerardi F, et al. Efficacy of animal-assisted therapy adapted to reality orientation therapy: measurement of salivary cortisol. *Psychogeriatrics*. 2019 Sep;19(5):510-2. doi: <https://dx.doi.org/10.1111/psyg.12418>. PMID: 30740833.
17. Moyle W, Murfield J, Jones C, et al. Can lifelike baby dolls reduce symptoms of anxiety, agitation, or aggression for people with dementia in long-term care? Findings from a pilot randomised controlled trial. *Aging Ment Health*. 2019;23(10):1442-50. doi: 10.1080/13607863.2018.1498447. PMID: 138051539. Language: English. Entry Date: 20190817. Revision Date: 20190823. Publication Type: Article.
18. Cantarella A, Borella E, Faggian S, et al. Using dolls for therapeutic purposes: A study on nursing home residents with severe dementia. *Int J Geriatr Psychiatry*. 2018 July;33(7):915-25. doi: <http://dx.doi.org/10.1002/gps.4872>. PMID: 622559552.
19. Liang A, Piroth I, Robinson H, et al. A Pilot Randomized Trial of a Companion Robot for People With Dementia Living in the Community. *J Am Med Dir Assoc*. 2017 Oct 01;18(10):871-8. doi: <https://dx.doi.org/10.1016/j.jamda.2017.05.019>. PMID: 28668664.
20. Valenti SM, Aguera-Ortiz L, Olazarán RJ, et al. Social robots in advanced dementia. *Front Aging Neurosci*. 2015;7(Jun). doi: 10.3389/fnagi.2015.00133. PMID: CN-01103263.
21. Moyle W, Cooke M, Beattie E, et al. Exploring the effect of companion robots on emotional expression in older adults with dementia: a pilot randomized controlled trial. *J Gerontol Nurs*. 2013 May;39(5):46-53. doi: <https://dx.doi.org/10.3928/00989134-20130313-03>. PMID: 23506125.
22. Majic T, Gutzmann H, Heinz A, et al. Animal-assisted therapy and agitation and depression in nursing home residents with dementia: a matched case-control trial. *Am J Geriatr Psychiatry*. 2013 Nov;21(11):1052-9. doi: <https://dx.doi.org/10.1016/j.jagp.2013.03.004>. PMID: 23831177.
23. Strom BS, Engedal K, Benth JS, et al. Effect of the Sonas Programme on Communication in People with Dementia: A Randomized Controlled Trial. *Dement Geriatr Cogn Dis Extra*. 2017 Jan-Apr;7(1):122-35. doi: <https://dx.doi.org/10.1159/000468147>. PMID: 28553314.
24. Baker R, Holloway J, Holtkamp CCM, et al. Effects of multi-sensory stimulation for people with dementia. *J Adv Nurs*. 2003;43(5):465-77. doi: 10.1046/j.1365-2648.2003.02744.x. PMID: 106688835. Language: English. Entry Date: 20050425. Revision Date: 20150819. Publication Type: Journal Article.
25. van Weert JC, van Dulmen AM, Spreeuwenberg PM, et al. Effects of snoezelen, integrated in 24 h dementia care, on nurse-patient communication during morning care. *Patient Educ Couns*. 2005 Sep;58(3):312-26. PMID: 16054329.
26. Baker R, Bell S, Baker E, et al. A randomized controlled trial of the effects of multi-sensory stimulation (MSS) for people with dementia. *Br J Clin Psychol*. 2001 Mar;40(Pt 1):81-96. PMID: 11317951.
27. Maseda A, Cibeira N, Lorenzo-Lopez L, et al. Multisensory Stimulation and Individualized Music Sessions on Older Adults with Severe Dementia: Effects on Mood, Behavior, and Biomedical Parameters. *J Alzheimers Dis*. 2018;63(4):1415-25. doi: <https://dx.doi.org/10.3233/JAD-180109>. PMID: 29843244.
28. Collier L, McPherson K, Ellis-Hill C, et al. Multisensory stimulation to improve functional performance in moderate to severe dementia—interim results. *Am J Alzheimers Dis Other Demen*. 2010;25(8):698-703. doi: 10.1177/1533317510387582. PMID: 104960420. Language: English. Entry Date: 20101214. Revision Date: 20150820. Publication Type: Journal Article.
29. Staal JA, Sacks A, Matheis R, et al. The effects of Snoezelen (multi-sensory behavior therapy) and psychiatric care on agitation, apathy, and activities of daily living in dementia patients on a short term geriatric psychiatric inpatient unit. *Int J Psychiatry Med*. 2007;37(4):357-70. PMID: 18441625.
30. van Weert JC, Kerkstra A, van Dulmen AM, et al. The implementation of snoezelen in psychogeriatric care: an evaluation through the eyes of caregivers. *Int J Nurs Stud*. 2004 May;41(4):397-409. PMID: 15050851.
31. Baillon S, Van Diepen E, Prettyman R, et al. A comparison of the effects of Snoezelen and reminiscence therapy on the agitated behaviour of patients with dementia. *Int J Geriatr Psychiatry*. 2004 Nov;19(11):1047-52. PMID: 15481068.
32. Van Diepen E, Baillon SF, Redman J, et al. A pilot study of the physiological and behavioural effects of snoezelen in dementia. *British Journal of Occupational Therapy*. 2002 February;65(2):61-6. PMID: 35396447.
33. Baker R, Dowling Z, Wareing LA, et al. Snoezelen: its Long-Term and Short-Term Effects on Older People with Dementia. *British Journal of Occupational Therapy*. 1997;60(5):213-8. PMID: 104158278. Language: English. Entry Date: 20131114. Revision Date: 20150820. Publication Type: Journal Article.
34. Kwan RYC, Leung MCP, Lai CKY. A Randomized Controlled Trial Examining the Effect of Acupressure on Agitation and Salivary Cortisol in Nursing Home Residents with Dementia. *Dement Geriatr Cogn Disord*. 2017;44(1-2):92-104. doi: <https://dx.doi.org/10.1159/000478739>. PMID: 28768251.
35. Yang YP, Wang CJ, Wang JJ. Effect of Aromatherapy Massage on Agitation and Depressive Mood in Individuals With Dementia. *J Gerontol Nurs*. 2016 Sep 01;42(9):38-46. doi: <https://dx.doi.org/10.3928/00989134-20160615-03>. PMID: 27319407.
36. Yang MH, Lin LC, Wu SC, et al. Comparison of the efficacy of aroma-acupressure and aromatherapy for the treatment of dementia-associated agitation. *BMC Altern Med*. 2015 Mar 29;15:93. doi: <https://dx.doi.org/10.1186/s12906-015-0612-9>. PMID: 25880034.
37. Moyle W, Cooke ML, Beattie E, et al. Foot massage versus quiet presence on agitation and mood in people with dementia: a randomised controlled trial. *Int J Nurs Stud*. 2014 Jun;51(6):856-64. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2013.10.019>. PMID: 24216598.
38. Burns A, Perry E, Holmes C, et al. A double-blind placebo-controlled randomized trial of Melissa officinalis oil and donepezil for the treatment of agitation in Alzheimer's disease. *Dement Geriatr Cogn Disord*. 2011;31(2):158-64. doi: <https://dx.doi.org/10.1159/000324438>. PMID: 21335973.
39. Lin PW, Chan WC, Ng BF, et al. Efficacy of aromatherapy (*Lavandula angustifolia*) as an intervention for agitated behaviours in Chinese older persons with dementia: a cross-over randomized trial. *Int J Geriatr Psychiatry*. 2007 May;22(5):405-10. PMID: 17342790.
40. Ballard CG, O'Brien JT, Reichelt K, et al. Aromatherapy as a safe and effective treatment for the management of agitation in severe dementia: the results of a double-blind, placebo-controlled trial with Melissa. *J Clin Psychiatry*. 2002 Jul;63(7):553-8. PMID: 12143909.
41. Kouzuki M, Kitao S, Kaju T, et al. Evaluation of the effect of aroma oil as a bath salt on cognitive function. *Psychogeriatrics: the official journal of the Japanese Psychogeriatric Society*. 2019;02. doi: <http://dx.doi.org/10.1111/psyg.12481>. PMID: 629513763.
42. Takahashi Y, Shindo S, Kanbayashi T, et al. Examination of the influence of cedar fragrance on cognitive function and behavioral and psychological symptoms of dementia in Alzheimer type dementia. *Neuropsychopharmacology reports*. 2020. PMID: CN-02084429.
43. Zalomonson S, Freud T, Punchik B, et al. The results of a cross-over placebo-controlled study of the effect of lavender oil on behavioral and psychological symptoms of dementia. *Rejuvenation Res*. 2018 Oct 17;17:17. doi: <https://dx.doi.org/10.1089/rej.2018.2123>. PMID: 30328781.
44. Fung JKK, Tsang HW. Management of behavioural and psychological symptoms of dementia by an aroma-massage with acupressure treatment protocol: A randomised clinical trial. *J Clin Nurs*. 2018 May;27(9-10):1812-25. doi: <https://dx.doi.org/10.1111/jocn.14101>. PMID: 28986942.
45. Kwan RY, Leung MC, Lai CK. Acupressure for agitation in nursing home residents with dementia: study protocol for a randomized controlled trial. *Trials*. 2014 Oct 27;15:410. doi: <https://dx.doi.org/10.1186/1745-6215-15-410>. PMID: 25346026.
46. Jia Y, Zhang X, Yu J, et al. Acupuncture for patients with mild to moderate Alzheimer's disease: a randomized controlled trial. *BMC Altern Med*. 2017 Dec 29;17(1):556. doi: <https://dx.doi.org/10.1186/s12906-017-2064-x>. PMID: 29284465.
47. Moorman Li R, Gilbert B, Orman A, et al. Evaluating the effects of diffused lavender in an adult day care center for patients with dementia in an effort to decrease behavioral issues: a pilot study. *J Drug Assess*. 2017;6(1):1-5. doi: <https://dx.doi.org/10.1080/21556660.2016.1278545>. PMID: 28265482.
48. Yang SY, Shan CL, Qing H, et al. The Effects of Aerobic Exercise on Cognitive Function of Alzheimer's Disease Patients. *CNS Neurol Disord Drug Targets*. 2015;14(10):1292-7. PMID: 26556080.

49. Chieh-Yu F, Moyle W, Cooke M. A randomised controlled trial of the use of aromatherapy and hand massage to reduce disruptive behaviour in people with dementia. *BMC Altern Med*. 2013;13(1):165-73. doi: 10.1186/1472-6882-13-165. PMID: 104229104. Language: English. Entry Date: 20130205. Revision Date: 20150711. Publication Type: Journal Article.
50. Fu CY, Moyle W, Cooke M. A randomised controlled trial of the use of aromatherapy and hand massage to reduce disruptive behaviour in people with dementia. *BMC Altern Med*. 2013 Jul 10;13:165. doi: <https://dx.doi.org/10.1186/1472-6882-13-165>. PMID: 23837414.
51. Lu DF, Hart LK, Lutgendorf SK, et al. Slowing progression of early stages of AD with alternative therapies: a feasibility study. *Geriatr Nurs*. 2013 Nov-Dec;34(6):457-64. doi: <https://dx.doi.org/10.1016/j.gerinurse.2013.07.003>. PMID: 23972540.
52. Rodriguez-Mansilla J, Gonzalez-Lopez-Arza MV, Varela-Donoso E, et al. Ear therapy and massage therapy in the elderly with dementia: a pilot study. *J Tradit Chin Med*. 2013 Aug;33(4):461-7. PMID: 24187866.
53. Rodriguez-Mansilla J, Gonzalez Lopez-Arza MV, Varela-Donoso E, et al. The effects of ear acupressure, massage therapy and no therapy on symptoms of dementia: a randomized controlled trial. *Clin Rehabil*. 2015 Jul;29(7):683-93. doi: <https://dx.doi.org/10.1177/0269215514554240>. PMID: 25322869.
54. Jimbo D, Kimura Y, Taniguchi M, et al. Effect of aromatherapy on patients with Alzheimer's disease. *Psychogeriatrics*. 2009;9(4):173-9. doi: 10.1111/j.1479-8301.2009.00299.x. PMID: CN-00750884.
55. Hawranik P, Johnston P, Deatrach J. Therapeutic touch and agitation in individuals with Alzheimer's disease. *West J Nurs Res*. 2008 Jun;30(4):417-34. doi: <https://dx.doi.org/10.1177/0193945907305126>. PMID: 18272750.
56. Woods DL, Craven RF, Whitney J. The effect of therapeutic touch on behavioral symptoms of persons with dementia. *Altern Ther Health Med*. 2005 Jan-Feb;11(1):66-74. PMID: 15712768.
57. Smallwood J, Brown R, Coulter F, et al. Aromatherapy and behaviour disturbances in dementia: a randomized controlled trial. *Int J Geriatr Psychiatry*. 2001 Oct;16(10):1010-3. PMID: 11607948.
58. Onega LL, Pierce TW, Epperly L. Effect of Bright Light Exposure on Depression and Agitation in Older Adults with Dementia. *Issues Ment Health Nurs*. 2016 Sep;37(9):660-7. PMID: 27254531.
59. Riemersma-van der Lek RF, Swaab DF, Twisk J, et al. Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities: a randomized controlled trial. *Jama*. 2008 Jun 11;299(22):2642-55. doi: <https://dx.doi.org/10.1001/jama.299.22.2642>. PMID: 18544724.
60. Hickman SE, Barrick AL, Williams CS, et al. The effect of ambient bright light therapy on depressive symptoms in persons with dementia. *J Am Geriatr Soc*. 2007 Nov;55(11):1817-24. PMID: 17944896.
61. Barrick A, Sloane P, Williams C, et al. Impact of ambient bright light on agitation in dementia. In *Int J Geriatr Psychiatry*
62. Ancoli-Israel S, Martin JL, Gehrman P, et al. Effect of light on agitation in institutionalized patients with severe Alzheimer disease. *Am J Geriatr Psychiatry*. 2003 Mar-Apr;11(2):194-203. PMID: 12611749.
63. Figueiro MG, Plitnick B, Roohan C, et al. Effects of a Tailored Lighting Intervention on Sleep Quality, Rest-Activity, Mood, and Behavior in Older Adults With Alzheimer Disease and Related Dementias: A Randomized Clinical Trial. *J Clin Sleep Med*. 2019 Dec 15;15(12):1757-67. doi: <https://dx.doi.org/10.5664/jcsm.8078>. PMID: 31855161.
64. Burns A, Allen H, Tomenson B, et al. Bright light therapy for agitation in dementia: a randomized controlled trial. *Int Psychogeriatr*. 2009 Aug;21(4):711-21. doi: <https://dx.doi.org/10.1017/S1041610209008886>. PMID: 19323872.
65. Dowling GA, Graf CL, Hubbard EM, et al. Light treatment for neuropsychiatric behaviors in Alzheimer's disease. *West J Nurs Res*. 2007 Dec;29(8):961-75. PMID: 17596638.
66. Dowling GA, Hubbard EM, Mastick J, et al. Effect of morning bright light treatment for rest-activity disruption in institutionalized patients with severe Alzheimer's disease. *Int Psychogeriatr*. 2005 Jun;17(2):221-36. PMID: 16050432.
67. Dowling GA, Mastick J, Hubbard EM, et al. Effect of timed bright light treatment for rest-activity disruption in institutionalized patients with Alzheimer's disease. *Int J Geriatr Psychiatry*. 2005 Aug;20(8):738-43. PMID: 16035127.
68. Fontana Gasio P, Krauchi K, Cajochen C, et al. Dawn-dusk simulation light therapy of disturbed circadian rest-activity cycles in demented elderly. *Exp Gerontol*. 2003 Jan-Feb;38(1-2):207-16. PMID: 12543279.
69. Mishima K, Hishikawa Y, Okawa M. Randomized, dim light controlled, crossover test of morning bright light therapy for rest-activity rhythm disorders in patients with vascular dementia and dementia of Alzheimer's type. *Chronobiol Int*. 1998 Nov;15(6):647-54. PMID: 9844752.
70. Richards KC, Beck C, O'Sullivan PS, et al. Effect of individualized social activity on sleep in nursing home residents with dementia. *J Am Geriatr Soc*. 2005 Sep;53(9):1510-7. PMID: 16137280.
71. Toseland RW, Diehl M, Freeman K, et al. The impact of validation group therapy on nursing home residents with dementia. *J Appl Gerontol*. 1997 March;16(1):31-50. doi: <http://dx.doi.org/10.1177/073346489701600102>. PMID: 27097884.
72. Kiosses DN, Rosenberg PB, McGovern A, et al. Depression and Suicidal Ideation During Two Psychosocial Treatments in Older Adults with Major Depression and Dementia. *J Alzheimers Dis*. 2015;48(2):453-62. doi: <https://dx.doi.org/10.3233/JAD-150200>. PMID: 26402009.
73. Spector A, Charlesworth G, King M, et al. Cognitive-behavioural therapy for anxiety in dementia: pilot randomised controlled trial. *Br J Psychiatry*. 2015 Jun;206(6):509-16. doi: <https://dx.doi.org/10.1192/bjp.bp.113.140087>. PMID: 25698766.
74. Spector A, Orrell M, Charlesworth G, et al. Factors influencing the person-carer relationship in people with anxiety and dementia. In *Aging Ment Health*
75. Stanley MA, Calleo J, Bush AL, et al. The peaceful mind program: a pilot test of a cognitive-behavioral therapy-based intervention for anxious patients with dementia. *Am J Geriatr Psychiatry*. 2013 Jul;21(7):696-708. doi: <https://dx.doi.org/10.1016/j.jagp.2013.01.007>. PMID: 23567399.
76. Beck CK, Vogelpohl TS, Rasin JH, et al. Effects of behavioral interventions on disruptive behavior and affect in demented nursing home residents. *Nurs Res*. 2002 Jul-Aug;51(4):219-28. PMID: 12131234.
77. Lin LC, Yang MH, Kao CC, et al. Using acupressure and Montessori-based activities to decrease agitation for residents with dementia: a cross-over trial. *J Am Geriatr Soc*. 2009 Jun;57(6):1022-9. PMID: 19507295.
78. Finnema E, Drees RM, Ettema T, et al. The effect of integrated emotion-oriented care versus usual care on elderly persons with dementia in the nursing home and on nursing assistants: a randomized clinical trial. *Int J Geriatr Psychiatry*. 2005 Apr;20(4):330-43. PMID: 15799079.
79. Kovach CR, Taneli Y, Dohearty P, et al. Effect of the BACE intervention on agitation of people with dementia. *Gerontologist*. 2004 Dec;44(6):797-806. PMID: 15611216.
80. Schrijnemaekers V, van Rossum E, Candel M, et al. Effects of emotion-oriented care on elderly people with cognitive impairment and behavioral problems. *Int J Geriatr Psychiatry*. 2002 Oct;17(10):926-37. PMID: 12325052.
81. Lawton MP, Van Haitsma K, Klapper J, et al. A stimulation-retreat special care unit for elders with dementing illness. *Int Psychogeriatr*. 1998 Dec;10(4):379-95. PMID: 9924833.
82. Law W, Kwok TCY. Impacts of a multicomponent intervention programme on neuropsychiatric symptoms in people with dementia and psychological health of caregivers: A feasibility pilot study. *Int J Geriatr Psychiatry*. 2019 Aug 07;07:07. doi: <https://dx.doi.org/10.1002/gps.5191>. PMID: 31390090.
83. Bailey EM, Stevens AB, LaRocca MA, et al. A Randomized Controlled Trial of a Therapeutic Intervention for Nursing Home Residents With Dementia and Depressive Symptoms. *J Appl Gerontol*. 2017 Jul;36(7):895-908. doi: <https://dx.doi.org/10.1177/0733464815627956>. PMID: 26912731.
84. Fischer-Terworth C, Probst P. Evaluation of a TEACCH-and music therapy-based psychological intervention in mild to moderate dementia. In *Geropsych: the journal of gerontopsychology and geriatric psychiatry*
85. van der Kooij CH, Drees RM, de Lange J, et al. The implementation of integrated emotion-oriented care: did it actually change the attitude, skills and time spent of trained caregivers? *Dementia*. 2013 Sep;12(5):536-50. doi: <https://dx.doi.org/10.1177/1471301211435187>. PMID: 24337328.

Appendix D. Interventions for PLWD Well-Being

Exercise

Table D-1. Risk of bias assessment: exercise

Author, Year PMID	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funding	Overall Rating
Fleiner 2020 ¹ 32039852	2 weeks	Medium 18%	Medium	Low	High	Low	Medium	None	High
Liu 2020 ² 32084347	4 weeks	Medium 12%	Medium	Low	High	Low	Medium	Government	High
Chen 2019 ³ 30301358	6 months	Low 3%	Medium	Medium	Medium	Low	Low	Government	Medium
Huang 2019 ⁴ 31743998	10 months	Low 8%	Low	High	High	Low	Low	Government	Medium
Karssemeijer 2019 ⁵ 31409559	12 weeks	Low 12%	Medium	High	Medium	Low	Medium	Nonprofit	High
Karssemeijer 2019 ⁵ 31409559	24 weeks	Medium 20%	Medium	High	Medium	Low	Medium	Nonprofit	High
Henskens 2018 ⁶ 29750023	6 months	Medium 25%	Low	High	Medium	Low	High	Unclear	High
Ho 2018 ⁷ 30496547	8 weeks	Low (no attrition reported)	Medium	High	High	Low	Low	Government	High
Lamb 2018 ⁸ 29769247	6 months	Low 10%	Low	Low	Medium	Low	Medium	Government	Low
Chen 2017 ⁹ 27879982	15 months	Medium 15%	Medium	Low	High	Low	Low	Government	High
Bossers 2016 ¹⁰ 27321604	9 week	Low 11%	Low	Low	High	Low	Low	NR	Medium
Cancela 2016 ¹¹ 26087884	15 months	Medium 40%	Low	High	Medium	Low	Low	Government	High
Hoffman 2016 ¹² 26682695	16 weeks	Low 5%	Low	Low	Medium	Low	Medium	Government Foundation	Low
Hoffman 2016 ¹² 26682695	12 months	Low 15%	Low	Low	Medium	Low	Medium	Government Foundation	Low
Toots 2016 ¹³ 26782852	4 months	Low 8%	Medium	Low	Low	Low	Low	Government	Medium
Toots 2016 ¹³ 26782852	7 months	Low 15%	Medium	Low	Low	Low	Low	Government	Medium
Telenius 2015 ¹⁴ 26630910	12 week	Low 9%	Medium	Low	Medium	Low	Low	Foundation	Medium
Telenius 2015 ¹⁴ 26630910	6 months	Low 18%	Medium	Low	Medium	Low	Low	Foundation	Medium
Yang 2015 ¹⁵ 26556080	3 months	Low (no attrition reported)	Medium	High	Medium	Low	Medium	Government	High
Pitkala 2013 ¹⁶ 23589097	6 months	Medium 14%	Low	Low	Medium	Low	Low	Government Foundation	Medium
Hauer 2012 ¹⁷ 22211512	3 months	Medium 12%	Low	Low	Low	Low	Low	Foundations	Medium
Hauer 2012 ¹⁷ 22211512	6 months	Medium 19%	Low	Low	Low	Low	Low	Foundations	Medium
Fan 2011 ¹⁸ 21385519	12 weeks	Medium 13%	High	X	X	X	X	X	X
Roach 2011 ¹⁹ 21937893	16 weeks	Medium 22%	Medium	Low	High	Low	Medium	Government	High
Eggermont 2009 ²⁰ 18926856	12 weeks	Low 9%	Medium	Low	Medium	Low	Medium	Foundation	Medium

Rolland 2007 ²¹ 17302650	6 months	Medium 13%	Medium	Low	Medium	Low	High	Government	High
Rolland 2007 ²¹ 17302650	12 months	Medium 18%	Medium	Low	Medium	Low	High	Government	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table D-2. Characteristics of included studies: exercise

Author, Year PMID Country RoB Type	Intervention: Intervention Target Mode Components Frequency Duration Delivery/Supervision	Comparison: Target Mode Components Frequency Duration Delivery	Setting Design Clusters N Randomized	PLWD Dementia Characteristics Type(s) Severity Diagnostic Criteria Age Diagnosed	PLWD N Age (mean) Female % Race (% majority) Detailed Race Education (mean years)	PLWD SES Prior Disability Household Characteristics Health Insurance	Informal Caregiver N Age (mean) Female % Race (% majority) Education (mean years) Relationship	Informal Caregiver (IC) Duration Live With PLWD IC Paid Health Status Dementia Family History Employment IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)tics	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing	PLWD Outcomes Quality of Life Daily Function Mobility BPSD Adverse Events Services Other	Informal Caregiver Outcomes
Chen 2019 ³ 30301358 China Medium Explanatory	PLWD Individual 30 min., 3x/week 6 months Supervision: 1:1 with trained Master's level RN	PLWD Routine care	Nursing Home RCT Clusters: NA N=60	Type: 100% AD Severity: mild- moderate (mean MMSE=18) Diagnosis: ISCDRHP- 10 th edition Age diagnosed: NR	N=60 82 years 72% female Race: NR Education: NR	NR	NR	NR	NR	NR	6 months	QoL: NR Function: EdFED, Time autonomous eating (min.) Mobility: NR BPSD: NR AE: NR Services: NR Other: Eating metrics	NR
Huang 2019 ⁴ 31743998 China Medium Explanatory	PLWD Group Tai Chi 20 min., 3x/week 10 months Supervision: professional therapists in LTC	PLWD Routine care	Nursing Homes (3) RCT Clusters: NA N=80	Type: NR Severity: mild (mean MMSE=21) Diagnosis: DSM-IV Age diagnosed: NR	N=80 82 years 68% female Race: NR Education: 72 months	NR	NR	NR	NR	NR	5 months, 10 months	QoL: NR Function: Barthel Mobility: NR BPSD: NPI, GDS AE: AE, falls Services: NR Other: cognitive	NR
Lamb 2018 ⁸ 29769247 UK Low Explanatory Petrou 2019 ²² Lamb 2018 ⁸	PLWD Group sessions in gym: moderate-high intensity aerobic (cycling) & strength exercises (dumb bells) + UC 1-1.5 hours, 2x/week 4 months; then advised home exercise Supervision: PT and assistant	Mostly PLWD Individual Usual care (activity advice, prescriptions for symptoms; counseling for IC)	Community: Groups in gym; Individual at home. RCT Clusters: NA N=494	Type: NR Severity: mild- moderate (mean MMSE=22) Diagnosis: DSM-IV Age diagnosed: NR	N=494 77 years 39% female 97% white More race: no Education: NR	NR	N=459 69 years 77% female Race: NR Education: NR Relation: 78% spouse	NR	NA	NA	6 months, 12 months	QoL: EQ-5D-3L, QOL-AD Function: BADLS Mobility: NR BPSD: NPI AE: AE, SAE (fall, hospitalization, fracture, death) Services: health service use Other: ADAS-cog	QoL: EQ- 5D-3L Burden: ZBI

Bossers 2016 ¹⁰ 27321604 Medium Netherlands Explanatory Bossers, 2015 ²³ , Bossers 2014 ²⁴	PLWD Group 1. Aerobic (walking) plus strength training 30 min, 4x/week (2 weeks strength, 2 walking) 9 weeks (total 36) 2. Aerobic (walking) 30 min, 4x/week 9 weeks Supervision: Trained RA (HMS student)	PLWD Group Social activity 4x/week 9 weeks	Nursing Homes; Psycho-geriatric RCT Clusters: NA N=118	Type: AD, VD, mixed AD-VD, NR Severity: mild 22%, moderate 53%, severe 25% (of n=105) Diagnosis: Chart diagnosis Age diagnosed: NR	N=118 86 years 26% female Race: NR More race: no Education: mean NR; 60% ≤ high school	NR	NR	NR	NA	NA	9 weeks	QoL: NR Function: Katz ADL, E-ADL, PPT-7 Mobility: NR BPSD: NR AE: NR Services: NR Other: NR	NR
Hoffman 2016 ¹² 26682695 Low Denmark Explanatory Sobol 2016 ²⁵ Hoffman 2013 ²⁶	PLWD Group Supervised exercise [strength 1 st 4 weeks; aerobic x 16 weeks (bike ergometer, cross trainer, treadmill)] 60 min, 3x/week 16 weeks Supervision: PT	PLWD Usual care Individual As needed (access to memory clinic staff)	NR (PLWD all community-dwelling) Cluster: NA N=200	100% probable AD Severity: mild (MMSE > 19) Diagnosis: NINCDS-ADRDA Diagnosis age: NR	N=200 71 years 44% female Race: NR More race: NA Education: NR	NR	N=NR, assume 200 (inclusion criterion) All: NR	Live together: 72% Other: NR	NA	NA	16 weeks	QoL: EQ-5D-5L Function: ADCS-ADL Mobility: NR BPSD: NPI-12, HAM-D AE: AE, SAE Services: NR Other: ADAS-Cog, MMSE, adherence	NR (ICG proxy for NPI-12, ADCS-ADL and 1 EQ-5D)
Toots 2016 ¹³ 26782852 Sweden Medium Explanatory Sondell 2018 ²⁷ Toots 2018 ²⁸ Toots 2017 ²⁹ Bostrom 2016 ³⁰	PLWD Groups at RC facilities Individualized high-intensity functional (weight-bearing) exercise (HIFE): leg strength, balance, mobility 5 x 45 min per 2 weeks 4 months (40 sessions) Supervision: 2 PTs	PLWD Seated group activity per OT Frequency: NR Duration: NR Supervision: OT or OTA	16 residential care facilities (9 NH units, 10 dementia units) Cluster: RCT Clusters: 36 N=186	86% AD, VD, or mixed AD-VD Severity: mild-moderate (mean MMSE=15) Diagnosis: DSM-IV-TR Age diagnosed: NR	N=186 85 years 76% female Race: NR More race: NA Education: NR	NR	NA	NA	NA	NA	4 months, 7 months	QoL: NR Function: Barthel ADL, FIM Mobility: BBS BPSD: NR AE: collected/NR (except 1 death) Services: NR Other: adherence	NR
Telenius 2015 ¹⁴ 26630910 Norway Medium Explanatory	PLWD Group Individualized high-intensity functional (weight-bearing) exercise (HIFE): 50-60 min, 2x/week 12 weeks Supervision: 1 PT per 3 PLWD	PLWD Group Seated activity (game, music, read, stretch) 50-60 min, 2x/week 12 weeks Supervision: OT or nurse	18 NHs RCT Block randomized (6-12 PLWD/site) N=170	NR Severity: Mild-moderate (CDR 1 or 2) Diagnosis: NR Age diagnosed: NR	N=170 87 years 74% female Race: NR More race: NA Education: NR	NR	NA	NA	NA	NA	12 weeks, 6 months	QoL: QUALID Function: Barthel ADL Mobility: BBS BPSD: NPI, CSD AE: AE Services: NR Other: MMSE, attendance, CST, 6-meter walk	NR
Pitkala 2013 ¹⁶ 23589097 Finland Medium Explanatory Roitto 2018 ³¹ Ohman 2016 ³² Ohman 2016 ³³ Pertilla 2016 ³⁴ Pitkala 2011 ³⁵	PLWD 1. Individual tailored home-based exercise (various) 1 hour, 2x/week 12 weeks 2. Pre-determined group exercise (balance strength, endurance, executive function) 1 hour, 2x/week 12 weeks Supervision: PT; ICG participation optional	PLWD Usual care plus nutrition & exercise advice per study nurses	Home (individual), adult day care centers (group) RCT Clusters: NA N=210	100% AD 67% moderate-severe Diagnosis: NINCDS-ADRDA Age diagnosed: NR	N=210 (dyads) 78 years 39% female 100% white More race: NA Education: mean NR; 41% 8-12 years; 38% < 8 years	NR	N=210 76 years 61% female 100% white Education: mean NR; 54% 8-12 years; 26% < 8 years Relation: 100% spouse	Duration: NR Live together: 100% Paid: NR Health: count of medications, Charlson Index Family history: NR Employed: NR Training: NR	NA	NA	6 months (12 months=high RoB)	QoL: NR Function: FIM Mobility: SPPB BPSD: NR AE: falls, fractures, hospitalizations Services: health & social service use, costs Other: Adherence	NR

Hauer 2012 ¹⁷ 22211512 Germany Medium Explanatory Pomiersky 2020 ³⁶	PLWD Group Progressive resistance & lower extremity functional training (walk, stair climb, sit to stand)) 2 hours, 2x/week 3 months Supervision: "a qualified instructor"	PLWD Group Low-intensity activity (stretch, hand weights, seated ball games, etc.) 1 hour, 2x/week 3 months Supervisor: Yes but NR	Outpatient geriatric rehabilitation RCT Clusters: NA N=122	Type: 50% AD, 34% VD, 16% other AD/VD Severity: mean MMSE 22 (mild) Diagnosis: NINCDS- ADRDA or NINDS- AIREN, CERAD, TMT Diagnosis age: NR	N=122 83 years 74% female Race: NR More race: NA Education: 11 years	84% community- dwelling Other: NR	NR	NR	NA	NA	3 months, 6 months	QoL: NR Function: NR Mobility: TUG, POMA, modified PAQE, modified SPPB (SPPB NR) AE: AE (text) Services: NR Other: strength (lab metrics), adherence	NR
Eggermont 2009 ²⁰ 18926856 Netherlands Medium Explanatory	PLWD Group (by living unit): Hand movement program (fingers, pinch ball, handle rubber ring) 30 min, 5x/week 6 weeks Supervision: RT or Master students	PLWD Group Stories read by RT followed by conversation 30 min, 5x/week 6 weeks Supervision: RT or student	NHs (10 psychogeriatric units) Cluster RCT Clusters: 10 N=66	Type: uncertain Severity: mean MMSE=16 Diagnosis: DSM-IV Age diagnosed: NR	N=66 85 years Female%: NR Race: NR Education: NR but "NSD"	NR	NA	NA	NA	NA	6 weeks, 12 weeks	QoL: NR Function: NR Mobility: NR BPSD: GDS, SCL- 90 (anxiety) AE: NR Services: NR Other: cognition, actigraphy	NR

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: AD=Alzheimer's Disease; ADAS-Cog= Alzheimer's Disease Assessment Scale, cognitive subscale; ADCS-ADL= Alzheimer's Disease Cooperative Study-Activities of Daily Living; ADL=Activities of Daily Living; AE=adverse events; BADLS=Bristol Activities of Daily Living Scale; BBS=Berg Balance Scale; BPSD=Behavioral and Psychiatric Symptoms of Dementia; CERAD=Consortium to Establish a Registry for Alzheimer's Disease; char=characteristics; CSD=Cornell Scale for Depression; CST=Chair Stand Test; DSM=Diagnostic and Statistical Manual of Mental Disorders; E-ADL=Erlangen-ADL scale; EdFED=Edinburgh Feeding Evaluation in Dementia; EQ-5D-5L=EuroQol 5 dimension-5 level; EQ-5D-3L=EuroQol 5 dimension-3 level; ET=exercise therapist; FC=formal caregiver; FIM=Functional Independence Measure; FINALEX=Finnish Alzheimer's Disease Exercise Trial; GDS=Geriatric Depression Scale; GHQ=Global Health Questionnaire; Govt.=Government; HAM-D=Hamilton Depression Rating Scale; HIFE=high-intensity functional exercise; HMS=Human Movement Science; IC=informal caregiver; ISCDRHP=International Statistical Classification of Diseases and Related Health Problems; LTC=long term care; MMSE=Mine Mental State Exam; N=number; NA=not applicable; NH=nursing home; NINDS-AIREN= National Institute of Neurological Disorders and Stroke and Association Internationale pour la Recherche et l'Enseignement en Neurosciences; NINCDS-ADRDA=National Institute of Neurological and Communicative Diseases and Stroke/Alzheimer's Disease and Related Disorders Association; NPI=Neuropsychiatric Inventory; NR=Not Reported; OT=occupational therapist; OTA=occupational therapy assistant; PAQE=Physical Activity Questionnaire for the Elderly; PMID=PubMed Identification Number; POMA=Performance Oriented Motor Assessment; PPT-7=7-item Physical Performance Test; PT=physical therapist/physiotherapist; PLWD=Persons with Dementia; QoL=Quality of Life; QOL-AD=Quality of Life in Alzheimer's Disease; QUALID=Quality of Life in Late-stage Dementia Scale; RA=Research Assistant; RCT=Randomized controlled trial; RT=Recreational Therapist; SAE=serious adverse events; SCL-90=Symptom Checklist; SPPB=Short Physical Performance Battery; TMT=Trail Making Test; TUG=Timed Up and Go; VD=vascular dementia; wk=week; yrs=years; ZBI=Zarit Burden Interview

Table D-3. PWD outcomes summary low and medium risk of bias studies: exercise

Study (PMID) Comparison RoB Category	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Chen 2019 ³ 30301358 China Medium Explanatory	EdFED (proxy) Mean (SD) Time of autonomous eating (minutes) Mean (SD) 6 months	Change from baseline Change from baseline	-0.63 (0.8) 2.13 (3.5)	-0.10 (0.5) 0.29 (0.4)	<0.05 <0.05
Huang 2019 ⁴ 31743998 Tai Chi vs. Usual Care Medium Explanatory	Barthel ADL GDS NPI 10 months	Mean (SD) Mean (SD) Mean (SD)	94.1 (11.6) 2.4 (1.0) 6.4 (4.8)	92.6 (13.3) 5.4 (1.9) 10.2 (5.7)	NR <0.05 <0.05
Lamb 2018 ⁸ (29769247) Group Exercise (aerobic + strength training) vs. Usual Care Low Explanatory	QOL-AD (self-report) Mean (SD) EQ-5D-3L (self-report) Mean (SD) BADLS (proxy) Mean (SD) NPI (proxy) Median (IQR) 6 months, 12 months	Adjusted mean difference (CI) 0.7(-0.2, 1.7) Adjusted mean difference (CI) -0.002 (-0.04, 0.04) Adjusted mean difference (CI) 0.3 (-1.7, 1.2) Adjusted median difference (CI) -2.1 (-4.8, 0.7)	12 months 38.4 (5.8) 0.81 (0.22) 17.0 (10.2) 12 (4 to 23)	12 months 39.1 (5.7) 0.82 (0.25) 15.9 (9.7) 9 (3 to 20)	0.13 0.93 0.70 0.14

Bossers 2016 ¹⁰ (27321604) Group (walking + strength training) vs. Walking vs. Social Activity Medium Explanatory	Katz ADL (proxy) Between group mean change from baseline E-ADL (RA) Between group mean change from baseline PPT-7 (RA) Between group mean change from baseline 9 weeks	Combined vs social -2.79 Walking vs social -0.99 Combined vs social -3.83 Walking vs social -2.92 Combined vs social -3.38 Walking vs social -1.66	NR	NR	0.01 0.97 <0.001 0.01 0.003 0.29
Hoffman 2016 ¹² (26682695) Group Exercise (aerobic + initial strength training) vs. Usual Care Low Explanatory	EQ-5D-3L (self-report) Mean (SD) ADCS-ADL (proxy) Mean (SD) NPI-12 (proxy) Mean (SD) HAM-D: Depression Mean (SD) 16 weeks	Mean difference (CI) -0.1 (-0.02, 0.04) Mean difference (CI) -0.1 (-1.8, 1.5) Mean difference (CI) -3.5 (-5.8, -1.3) Mean difference (CI) -0.1 (-0.7, 0.5)	0.92 (0.11) 64.4 (9.4) 8.8 (8.5) 1.7 (2.5)	0.92 (0.09) 62.7 (10.4) 11.4 (11.0) 1.8 (2.3)	0.40 0.88 0.002 0.79
Toots 2016 ¹³ (26782852) HIFE (leg strength, balance, mobility) vs. seated group activities Medium Explanatory	Barthel ADL Mean (SE) FIM Mean (SE) BBS Mean (SE) 4 month, 7 months	Mean difference (CI) 0.57 (-0.30, 1.43) Mean difference (CI) 0.78 (-2.21, 3.77) Mean difference (CI) -0.02 (-2.53, 2.49)	-1.56 (0.32) -6.77 (1.09) -2.08 (0.91)	-2.12 (0.32) -7.55 (1.08) -2.05 (0.90)	0.20 0.61 0.98
Telenius 2015 ¹⁴ (26630910) HIFE (strength + balance) vs. seated group activity Medium Explanatory	QUALID Between group difference Barthel ADL Between group difference Berg Balance Scale Mean change from baseline NPI Mean (CI) CSD Between group difference 3 months, 6 months	-0.9 points 1.0 points NR Between group difference -1.6 Between group difference 0.2 points	NR NR +2.7 points 4.8 (3.8, 5.8) NR	NR NR -1.4 points 6.4 (5.1, 7.7) NR	NR NR 0.031 0.059 NR
Pitkala 2013 ¹⁶ 23589097 Medium (6 months) Explanatory	FIM Within-group mean change from baseline 6 months FIM Between-group change from baseline 6 months Adverse events: 12 months Falls Incidence rate (95% CI) Fractures (any) Incidence rate (95% CI) Hospitalizations Incidence rate (95% CI) Health service costs (24 months): Adjusted mean (95% CI) per dyad per year, US dollars	NR NR NR NR NR NR	Home exercise: -6.5 (-4.4, -8.6) Group exercise: -8.9 (-6.7, -11.2) Home: 1.35 (1.07, 1.67) Group: 1.86 (1.51, 2.26) Home: 0.06 (0.02, 0.17) Group: 0.09 (0.03, 0.21) Home: 0.47 (0.31, 0.68) Group: 0.54 (0.37, 0.77) Home exercise: \$25,112 (\$17,642, \$32,581) Group exercise: \$22,066 (\$15,931, \$28,199)	-11.8 (-9.7, -14.0) UC: 3.07 (2.63, 3.57) UC: 0.07 (0.02, 0.18) UC: 0.65 (0.46, 0.90) UC: \$34,121 (\$24,599, \$43,681)	Mixed-effect model p=0.003. Home exercise vs. UC: p=0.001 Group exercise vs UC: p=0.07 p=0.005 p=0.88 p=0.63 Home vs. UC: p=0.13 Group vs. UC: p=0.03
Hauer 2012 ¹⁷ (22211512) Group resistance & functional training vs low intensity motor activity Medium Explanatory	TUG Mean percent change from baseline (SD) POMA Mean percent change from baseline (SD) Modified PAQE Mean percent change from baseline (SD) 3 months, 6 months	Effect size 0.07 Effect size 0.15 Effect size 0.002	-11.2 (28.1) 22.2 (28.4) 134.5 (274.5)	-1.4 (26.3) 0.8 (23.5) 101.1 (178.6)	0.009 <0.001 0.64

Eggermont 2009 ²⁰ (18926856) Group hand movement program vs. read aloud control Medium Explanatory	GDS	NR	-1.56	-0.17	NR
	Change in group mean from baseline SCL-90	NR	-2.61	-0.69	NR
	Change in group mean from baseline 12 weeks, 6 months				

Abbreviations: ADCS-ADL= Alzheimer's Disease Cooperative Study-Activities of Daily Living; ADL=Activities of Daily Living; AE= adverse events; BADLS= Bristol Activities of Daily Living Scale; BBS=Berg Balance Scale; CSD=Cornell Scale for Depression; E-ADL=Erlangen-ADL (instrumental ADLs); EdFED=Edinburgh Feeding Evaluation in Dementia;EQ-5D-3L=EuroQol 5 dimension-3 level; EQ-5D-5L=EuroQol 5 dimension-5 level; FIM=Functional Independence Measure; GDS= Geriatric Depression Scale; HAM-D=Hamilton Depression Rating Scale; HIFE=high-intensity functional exercise; N=number; NPI = Neuropsychiatric Inventory (10 items); NPI-12= Neuropsychiatric Inventory-12 items; NR=not reported; PAQE=Physical Activity Questionnaire for the Elderly; PPT-7=7-item Physical Performance Test; POMA=Performance Oriented Motor Assessment (gait and balance); PMID=PubMed Identification Number; QOL-AD=Quality of Life in Alzheimer's Disease; QUALID= quality of life in late-stage dementia scale; SAE=serious adverse event; SCL-90=Symptom Checklist-90; SD=standard deviation; SPPB=Short Physical Performance Battery; TUG=Timed Up and Go; UC=usual care; vs=versus

Table D-4. Summary of strength of evidence for PLWD outcomes: exercise

Outcome Comparison	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Quality of Life: Group exercise vs usual care (2) or seated group activity (1)	3-12 months	3 RCTs ^{8, 12, 14} (1 NH) n=864	No significant difference in Quality of Life (QOL-AD, EQ-5D-5L, EQ-5D-3L or QUALID) for group exercise compared with usual care or seated group activity.	Low	Inconsistent	Direct	Imprecise	Insufficient
Daily function: Group exercise vs usual care (3) or home exercise (1)	4-12 months	3 RCTs ^{8, 12, 16} (0 NH) n=904	Inconsistent findings. Two RCTs found no significant difference in daily functioning (BADLS, ADCS-ADL) for PLWD for group exercise compared with usual care. One RCT found benefits of group exercise vs usual care and for home exercise vs usual care on the FIM at 6 months.	Low	Inconsistent	Direct	Imprecise	Insufficient
Daily function: Group exercise vs seated group activity (3)	3-7 months	3 RCTs ^{10, 13, 14} (3 NH) n=474	Inconsistent findings. Two RCTs found no significant difference in daily functioning (Barthel ADL) for group exercise compared with seated group activity. One RCT found benefits of combined group exercise vs social activity, and for walking vs. social activity on daily functioning [Katz ADL, E-ADL, PPT-7 (combined only)] at 9 weeks.	Medium	Inconsistent	Direct	Imprecise	Insufficient
Daily function: Tai Chi vs usual care	10 months	1 RCT ⁴ (NH) n=80	No significant difference in daily functioning (Barthel ADL) for Tai Chi compared with usual care.	Medium	Unknown	Direct	Imprecise	Insufficient
Daily function: Eating assistance needed. Supervised 1:1 hand exercise program vs. usual care	6 months	1 RCT ³ (NH) n=60	Statistically significant reduction (< 1 point) in feeding assistance needed (EdFED) with individual hand exercise program vs. usual care over 6 months.	Medium	Unknown	Direct	Imprecise	Insufficient
Daily function: Time of autonomous eating (min.) Supervised 1:1 hand exercise program vs. usual care	6 months	1 RCT ³ (NH) n=60	Statistically significant improvement in minutes of autonomous eating per meal with individual hand exercise program vs. usual care over 6 months.	Medium	Unknown	Direct	Imprecise	Insufficient
Balance: HIFE (strength + balance) vs. seated group activity	3-7 months	2 RCTs ^{13, 14} (2 NH) n=356	Inconsistent findings. One RCT found no significant difference in balance (BBS) between HIFE compared with seated group activity at 3 months, but beneficial effect at 6 months. One RCT found a beneficial of HIFE vs. seated group activities at 4 months but not 7 months.	Medium	Inconsistent	Direct	Imprecise	Insufficient
Neuropsychiatric symptoms: Tai Chi vs usual care	10 months	1 RCT ⁴ (NH) n=80	Statistically significant improvement in NPI with group Tai Chi vs. usual care over 10 months but not at 5 months.	Medium	Unknown	Direct	Imprecise	Insufficient
Neuropsychiatric symptoms: Group exercise vs usual care (2) or seated group activity (1)	3-12 months	3 RCTs ^{8, 12, 14} (1 NH) n=864	Inconsistent findings. Two RCTs found no significant difference in neuropsychiatric symptoms (NPI) with group exercise vs. usual care (1 NH), and one RCT reported a benefit of exercise vs. usual care on the NPI-12 at 4 months.	Low	Inconsistent	Direct	Imprecise	Insufficient
Depression: Tai Chi vs usual care	10 months	1 RCT ⁴ (NH) n=80	Statistically significant improvement in GDS with group Tai Chi vs. usual care over 10 months but not at 5 months.	Medium	Unknown	Direct	Imprecise	Insufficient
Depression: Group exercise vs usual care (1) or seated group activity (2)	3-6 months	3 RCTs ^{12, 14, 20} (2 NH) n=436	No significant difference in depression (HAM-D, CSD, GDS) for group exercise compared with usual care or seated group activity.	Medium	Inconsistent	Direct	Imprecise	Insufficient
Anxiety: Group hand movement program vs. attention control	3 months	1 RCT ²⁰ (NH) n=66	No significant difference in anxiety (SCL-90) with a group hand movement program vs. attention control over 3 months.	Medium	Unknown	Direct	Imprecise	Insufficient

Abbreviations: ADCS-ADL=Alzheimer’s Disease Cooperative Study-Activities of Daily Living; ADL=Activities of Daily Living; BADLS=Bristol Activities of Daily Living Scale; BBS=Berg Balance Scale; CSD=Cornell Scale for Depression; E-ADL=Erlangen-ADL scale; EdFED=Edinburgh Feeding Evaluation in Dementia; EQ-5D-5L=EuroQol 5 dimension-5 level; EQ-5D-3L=EuroQol 5 dimension-3 level; FIM=Functional Independence Measure; GDS=Geriatric Depression Scale; HAM-D=Hamilton Depression Rating Scale; HIFE=high-intensity functional exercise; N=number; NH=nursing home; NPI=Neuropsychiatric Inventory; PPT-7=7-item Physical Performance Test; PLWD=Persons with Dementia; QoL=Quality of Life; QOL-AD=Quality of Life in Alzheimer’s Disease; QUALID=Quality of Life in Late-stage Dementia Scale; RCT=Randomized controlled trial; SCL-90=Symptom Checklist

Evidence Map: Exercise

Table D-5. Characteristics of evidence map studies: exercise

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Enette 2020 ³⁷ (31921371) France Small sample	11. Continuous aerobic training (cycling) 30 minutes 2x/week 9 weeks 12. Interval aerobic cycling 30 minutes 2x/week 9 weeks	Interactive information sessions: supervised and structured. Multiple-choice questionnaires that PLWD discussed afterward	Memory clinic of a university hospital RCT 54 PLWD from NH or home	Mild to moderate AD diagnosed according to DSM-IV and MMSE ≥ 16.	N=54 (3 groups) Mean age: 78 years Female: 61% % majority race: no Education: 7-10 years	NR	NR	NR	NR	NR	10 weeks	QoL-AD, 6 meter walk, cognitive, blood tests	NR
Fleiner 2020 ¹ 32039852 Germany High RoB	PLWD Small group exercise (strength and endurance) 20 minutes 4x/day 3 days/week 2 weeks	Social stimulation program (table games per OT) 120 min./week 2 weeks	3 acute dementia care wards of a psychiatric hospital RCT 70 PLWD	PLWD with ICD-10 diagnosis of dementia (AD, VD, mixed, LBD, PD). Mean MMSE=18. Able to do TUG.	N=70 Mean age: 80 years Female: 47% % majority race: no Education: 13 years	NR	NA	NR	All RNs Other: NR	NA	2 weeks	NR	NPI
Lee, 2020 ³⁸ (31347294) Korea Small sample	Fumanet exercise program 1 hour/week 8 weeks	Non-equivalent control group	Community-based Quasi-experimental (segregated by day at center) N=45	Persons with mild dementia (CDR 0.5 to 1). Not visually impaired. Able to ambulate with or without assistance. Age 45 or older. Other exclusions	N=45 (22 exercise, 23 control) Age: Y Sex: Y Education: Y Race & majority: N	Dementia diagnosis: Y	NA	NA	NR	Training: Y	8 weeks	Cognitive function, gait, depression	NA
Liu 2020 ² (32084347) Taiwan High RoB	Isotonic strength training (weight machines) 2 sets of 12 5 days/week 4 weeks	Aerobic exercise (stationary cycling) 30 minutes 5 days/week 4 weeks	PLWDs Veterans NH RCT 80 PLWDs	MMSE 15-26 Other: NR	N=69 analyzed Mean age: 86 years Female: 16% % majority race: no Education: no	NR	NR	NR	NR	NR	4 weeks	Barthel Index, GDS, cognitive, blood test	NR
Sanders 2020 ³⁹ (32192537) Netherlands Pilot	Outdoor walking and lower limb strength training: 12 weeks low-intensity, then 12 weeks high-intensity training 3x/week 24 weeks	Flexibility exercises and recreational activities 3x/week 24 weeks	Recruited from daycare or residential care RCT 91 PLWDs	Dementia diagnosis per physician using DSM-IV; MMSE score >10 (mild-to-moderate dementia)	N=69 Mean age: 82 years Female: 62% % majority race: no Education: yes (3 levels)	None	NA	None	NA	NA	12 weeks 24 weeks 6MWS, leg strength and STROOP (6 weeks) 18 weeks 36 weeks)	6MWT (6-minute walk test) SPPB (Short Physical Performance Battery) Cognitive	NA

Brett 2019 ⁴⁰ (30912690) Australia Small sample	I1. 45-minute exercise program 1x per week I2. 15-minute exercise program 3x per week	Usual care activities (routine group activities)	2 nursing homes RCT N=60	PLWD living in nursing home, able to participate in intervention. Stratified by able to walk (71%) vs. participated sitting	Reported N=55 (17 Group 1, 19 Group 2, 19 usual care) Mean age: 85 Female: 66% Race % majority: Y Education: N	N of comorbidities: Y	NA	NA	NR	Length of service: Y	12 weeks	Falls, Timed Up & Go, Sit to Stand, Gait speed, Functional reach	NA
Karssemeijer 2019 ⁵ (31409559) Netherlands High RoB	I1. Exergame training I2. Aerobic training Both: 3x/week 12 weeks	Active control 12 weeks	Community centers RCT N=115 (92 completed)	Diagnosed with VD, AD or mixed dementia, MMSE ≥ 17, age ≥ 60 years. Additional criteria applied	N=115 (38 exergame, 38 aerobic, 39 controls) Mean age: 79 Female: 54% Education: N Race & majority: N	Living situation: Y	NA	NA	NR	NR	12 weeks	Frailty, adherence	NA
Nyman 2019 ⁴¹ (31819385) UK Pilot	Group Tai Chi 45 min plus discussion Weekly 20 weeks	Usual care	Community-dwelling dyads (PLWD and IC). Classes in church halls N=85	Diagnosis of dementia per MD (VD, AD, mixed, other), Mini Addenbrooke's Cognitive Exam ≥ 10, willing to attend Tai Chi class, able to do standing, other exclusions	N=85 (70 complete data) Mean age: 78 years Female: 40% Education: N Race & majority: N Time since diagnosis: Y	NR	N=85 Mean age: 71 years Female: 79% Other: NR	Living with PLWD: 87%	NA	NA	6 months	Berg Balance, Timed Up & Go, falls, cognitive	IC: TUG, other
Todri 2019 ⁴² (None) Spain Pilot	Global Postural Reeducation 30-40 minutes 2x/week 6 months (total 48 sessions)	Conventional exercises (proprioceptive, equilibrium, aerobic/walking) 30-40 minutes 2x/week 6 months	PLWDs were recruited from nursing homes, 2 countries 174 PLWD	Dementia (mild to moderate) per NINCDS-ADRDA criteria	N=135 analyzed Mean age: 81 years Female: 66% % majority race: no Education: "Medium"	NR	NA	NR	NR	NR	2 weeks	QoL-AD GDS Barthel Index, NPI Tinetti Scale, MMSE	NR
Henskens 2018 ⁶ (29750023) Netherlands High RoB	I1. [Exercise (strength + walking) + ADL training] I2. ADL training only I3. Exercise only All: 3x/week 6 months	Social control	11 Nursing homes 4 group RCT Cluster: 11 N=87	Dementia (AD, VD, Mixed, Unknown) per DSM-IV, living in NH, age ≥ 65 yrs., MMSE 1-24, complete 6-minute walk test	N=87 (66 completed) Age: Y Sex: Y Race: N Education: Y	NR	NA	NA	NR	NR	3 months, 6 months	QoL, function (created composite)	NA
Ho 2018 ⁷ (30496547) Hong Kong High RoB	I1: Dance movement therapy I2. Exercise Both: 2x/week 12 weeks	Waitlist	Outpatient RCT PLWD N=204	Dementia per DSM-IV or neurocognitive disorder (DSM V), CDR 0.5 to 1, age ≥ 65 years. Other exclusion criteria	N=204 (166 completed)	NR	NA	NA	NR	NR	3, 6, and 12 months	Depression, mood, daily function, cortisol	NA
Liu, 2018 ⁴³ (29969916) Hong Kong/ China Pilot	Simplified Tai-chi program to improve motor skills 16 weeks	Social activities	Community-dwelling PLWD; four community health centers RCT	PLWD aged 60 or older with Montreal Cognitive Assessment score ≤ 20; capable of exercise; caregiver able to exercise	N=26 (13 intervention/, 2 centers; 13 controls/2 centers) Age: Y Sex: Y %majority race: N Education: Y	Prior disability: Y Household Characteristics: Y	N=11 exercise group 13 control group Age: Y Sex: Y Education: Y Relation to PLWD: Y	Duration: Y	NR	NR	16 weeks	Motor performance	

Chen 2017 ⁹ (27879982) Taiwan High RoB	Resistance-band exercise program 3x/week x 6 mo., then 9 mo. per DVD	Usual daily activities	Nursing homes Cluster RCT (8 NH) N=150 PLWD	Wheelchair-bound PLWD, lived in nursing home 3 months, dementia per diagnosis or MMSE (cutpoint by education level)	N=127 completed (65 intervention, 62 controls) Age: Y Sex: Y % majority race: N Education: Y	100% wheelchair-bound Other: NR	NR	NR	NR	NR	3, 6, 9, 12 and 15 months	Depression, behavioral problems (BRS)	NR
Dawson, 2017 ⁴⁴ (28252473) USA Small sample	Moderate-Intensity Home-Based Exercise Program	Continuation of current levels of activity Randomized control trial	Community-based RCT PLWD	Dementia unspecified	N=13 intervention group 10 control group Age: Y Sex: Y % majority race: N Education: Y	NR	NA	NA	NR	NR	12 weeks	Measurements of walking and balance Assessment of cognition	NA
De Souto, 2017 ⁴⁵ (28542742) France Pilot	Multicomponent training: coordination, balance, muscle-strengthening & aerobic	Structured social activity	Nursing Homes RCT cluster PLWD	Dementia unspecified	N=47 PLWD (4 nursing homes) exercise; 50 PLWD (3 nursing homes) social activity Age: Y Sex: Y % majority race: N Education: Y	NR	NA	NA	NR	NR	24 weeks	MMSE Physical performance evaluation	NA
Hamilton, 2017 ⁴⁶ (27692024) Australia Small sample	Home-based exercise program	Structured, individually tailored exercise program	Home-based RCT	PLWD diagnosed with mild to moderate dementia	N=33 completed the study; 9 did not Age: Y Sex: Y % majority race: N Education: Y	NR	NA	NA	NR	NR	6 months	Psychological assessment Physical assessment fall risk Adherence	NA
Henwood, 2017 ⁴⁷ (28473006) Australia Small sample	Aquatic exercise for 2x/week 12 weeks	Usual care	Non-RCT: Facility PLWD allocated by ability to transport to swimming facility N=56	PLWD living in residential care facility, moderate to severe dementia	N=56 (analyzed 46; 23 each group) Age: Y Sex: Y % majority race: N Education: N	NR	NA	NA	NR	NR	12 weeks	Assessment of physical condition, balance and physical functions	NA
Lam, 2017 ⁴⁸ (28094873) Hong Kong Small sample	Whole-body vibration plus routine day activity program 2x/week 9 weeks	Routine day activity program 2x/week 9 weeks	PLWD with mild or moderate dementia, recruited from 2 day-care centers RCT	PLWD aged 65 or older with MMSE 10-22; ability to stand; living in the community	N=54 (27 intervention, 27 control) Age: Y Sex: Y % majority race: N Education: N	NR	NA	NA	NR	NR	9 weeks	Ability to stand; balance; quality of life; balance confidence	NA
Morris 2017 ⁴⁹ (28187125) USA Pilot	Supervised aerobic exercise program: 150 minutes per week for 26 weeks	Non-aerobic stretching and toning	Community-based RCT PLWD	PLWD with probable AD; age 55 or older; sedentary; community dwelling, supportive caregiver; other exclusion criteria	N=76 Age: yes Sex: yes % majority race: no Education: no	NR	NR	NR	NA	NA	26 weeks	Functional ability and depression	NA
Padala, 2017 ⁵⁰ (28655135) US Pilot	Home-based Wii-Fit interactive video-game-led physical exercise program, 8 weeks	Walking program	Community-based RCT N=30	PLWD 60 years and older diagnosed with AD; MMSE score \geq 18 with fear of falling; exclusion criteria applied	N=15 exercise group 15 control group Age: yes Sex: yes % majority race: yes Education: yes	Comorbidities: yes Medications: yes	NR	NR	NA	NA	8 weeks, 16 weeks	Balance	NA

Cancela 2016 ¹¹ (26087884) Spain High RoB	Exercise (stationary cycling) 15 minutes/day 15 months	Sedentary recreational activities	Institutional residential care RCT N=189	PLWD age ≥ 65 yrs., dementia per DSM-IV, able to stand and walk 30 meters without assistance, resident of facility	N=114 completed Age: Y Sex: Y % majority race: N Education: Y	Comorbidities: Y Other: NR	NR	NR	NR	NR	15 months	Neuropsychiatric symptoms, cognition, Katz ADL, mobility, depression	NA
Burge, 2016 ⁵¹ (27831462) Switzerland & Belgium Pilot	Group physical exercise 20 sessions 4 weeks	Social visits 20 sessions 4 weeks	5 Hospitals, acute psychiatric wards RCT PLWD	Moderate to severe dementia, on acute psychiatric ward	N=270 Age: yes Sex: no % majority race: no Education: no	NR	NA	NA	NA	NA	4 weeks	Barthel ADL, FIM, depression & behavior	NA
Venturelli 2016 ⁵² (27540967) Italy Small sample	I1:Aerobic exercise and cognitive training I2:aerobic exercise only I3: cognitive training only	No training	Nursing home PLWD Control trial	PLWD diagnosed with dementia, MMSE between 10 and 15; PLWD screened for physical limitations	N=80 (20 each group) Age: yes Sex: yes % majority race: no Education: no	Prior disability	NA	NA	NR	NR	3 months	Cortisol level; level of cognitive function; neuropsychological symptoms	NA
Bosser, 2015 ²³ (25648055) Netherlands High RoB	I1: Combined aerobic and strength training: 36 individualized sessions over 9 weeks I2: Aerobic only: participated in four walking sessions per week, 9 weeks	Four social visits each week	Nursing homes RCT PLWD	Mild-severe vascular dementia	N=123 Age: yes Sex: yes % majority race: no Education: yes	NR	NA	NA	NA	NA	18 weeks	MMSE 6-minute walk test, the 30-second sit-to-stand test	NA
Holthoff, 2015 ⁵³ (25884637) Germany Pilot	Home-based exercise; lower body on a movement trainer with computer control 12 weeks	Usual care Monthly clinical visits and a counselling including specific advice how to change inactive habits and increase the exercise	Community-based RCT PLWD	PLWD with mild to moderate AD per NINCDS-ADRA criteria, age 55 or older and their family caregivers; PLWD evaluated for other conditions that would rule out AD	N=30 Age: yes Sex: yes % majority race: Education:	Age at diagnosis	NR	NR	NR	NR	12 weeks	ADL; Neuropsychiatric symptoms; Executive function, language ability; Motor skills; Caregiver burden; MMSE	NA
Yang 2015 ¹⁵ (26556080) China High RoB	Moderate aerobic exercise (cycling) 3x/week 3 months	Health education	Outpatient RCT PLWD	Mild AD (criteria NR) or vascular dementia, age 65-80 years, MMSE 10-24; other exclusion criteria	N=50 Age: Y Sex: Y % majority race: N Education: Y	NR	NR	NR	NR	NR	3 months	QoL, cognition, neuropsychiatric symptoms	NA
Yu, 2015 ⁵⁴ (24652914) US Pilot	Cycling as exercise	Single group	Community-based Single group Pilot study	Older adults, mild to moderate AD; MMSE 12-24; CDR 1-3; able to perform exercises, qualified family caregivers	N=26 Age: Y Sex: Y % majority race: Y Education: Y	NR	N=26 Age: Y Sex: Y % majority race: Y Education: Y Relation to PLWD: N	NR	NR	Training: Y	6 months	Cognitive function; ADL; BPSD; caregiver distress	Caregiver distress decreased from baseline

Bosser 2014 ²⁴ (24844772) Netherlands Pilot	Evaluation of a combined aerobic and strength training program for institutionalized PLWD; studied effects on cognitive and physical function	Non-randomized pilot study; comparison of training program and social group visit program	PLWD from a single nursing home	Diagnosis of dementia; age 70 or older; not wheelchair bound; able to walk	N=18 PLWD exercise group 18 PLWD control group Age: Y Sex: Y Race: N Education: N	NR	NR	NA	NA	NR	6 weeks	Cognitive function tests Physical function tests	NA
Suttanon, 2013 ⁵⁵ (23117349) Australia Pilot	Individual tailored home-based exercise program; balance, strengthening and walking exercise	Home-based education program	Community-based RCT PLWD	PLWD assessed with mild to moderate AD	N=40 Age: yes Sex: yes % majority race: no Education: no	Prior disability: Y	NA	NA	NA	NA	6 months		Exercise arm PLWD dropped out due to inability of home caregivers to maintain the program
Canonici, 2012 ⁵⁶ (22994617) Brazil Small sample	Evaluation of a motor intervention program for PLWD with AD to promote functionality and reduce caregiver burden	Motor intervention compared to control group	NR	PLWD diagnosed with mild to moderate AD; CDR of 1 to 2	N=16 PLWD, 16 FC exercise arm 11 PLWD, 11 FC control arm Age: Y Sex: Y Race: N Education: Y	NR	NA	NA	NR	NR	6 months	Measurements of physical function, cognition and caregiver burden	NA
Nascimento, 2012 ⁵⁷ (22499405) Brazil Small sample	Supervised exercise program, including aerobic activity, strength, motor coordination, balance 6 months	PLWD who did not participate in the exercise program	Community Non-RCT: assigned by willingness to follow the exercise program	Community-based PLWD diagnosed with mild to moderate dementia and using MMSE; exclusion criteria for other health conditions	N=20 (10 exercise; 10 controls) Age: N Sex: N % majority race: N Education: N	NR	NA	NA	NA	NA	6 months	Neuropsychiatric, ADL	NA
Vreugdenhil, 2012 ⁵⁸ (21564154) Australia Small sample	Home-based exercise program	Usual care	Community-based RCT	PLWD diagnosed with AD	N=80 Age: Y Sex: Y % majority race: N Education: N	NR	NR	NR	NA	NA	4 months	Cognitive function; physical function; ADL; depression; global function	Caregiver burden
Fan 2011 ¹⁸ (21385519) Taiwan High RoB	Yoga training program 3 x 1 hour/week 12 weeks	Usual activities	Long term care facilities Quasi-experimental N=68	Age 60 or older, living in LTC, mild dementia per DSM-IV and MMSE 18-23, no recent exercise	N=68 (33 yoga, 35 usual activities); reported 59 Mean age: 75 Female: 59% Education: ≤ 9 yrs Race & majority: N	SES: N Disability: N Household: NA Marital status: Y N comorbidities: Y Length of residence: N Insurance: NA	NA	NA	NR	NR	12 weeks	Mobility (6 m. walk), balance, strength, depression (CSDD), behavior (BRS)	NR
Roach 2011 ¹⁹ (21937893) USA High RoB	I1: activity-specific exercise group I2. Supervised walking group 5x/week 16 weeks	Social conversation group	Long term care sites (7) RCT PLWD N=105	Residence in long term care, AD per NINCDS-ADRDA, able to walk, dependent in ≥ 1 ADL	N=82 completed Age: Y Sex: N % majority race: N Education: N	Length of stay: Y Other: N	NA	NA	NR	NR	16 weeks?	Walking, transferring, bed mobility	NR

Stella, 2011 ⁵⁹ (21915483) Brazil Small sample	Individual exercise (flexibility, strength, and agility)& functional balance exercise) 60 minutes three times per week.	Motor intervention compared to PLWD who did not receive the intervention	Community-based RCT PLWD	mild or moderate state of probable AD	N=16 PLWD exercise group 16 PLWD control group Age: Y Sex: Y % majority race: N Education: Y	NR	Age: Y Sex: Y % majority race: N Education: Y Relation to PLWD: N	Duration: Y	NA	NA	6 months	Neuropsychiatric inventory; Caregiver burden and stress	Reduction of caregiver stress
Venturelli, 2011 ⁶⁰ (21852281) Italy Small sample	Group exercise, 30 minutes of moderate walking exercise 4 times a week	Daily organized activities like bingo, patchwork sewing, and music therapy	Nursing homes RCT PLWD	PLWD diagnosed with AD, MMSE <=15 and absence of mobility limitations	N=12 PLWD, 12 caregivers walking group 12 PLWD, 12 caregivers control group Age: yes Sex: no % majority race: no Education: no	NR	Age: Y Sex: Y Relation to PLWD: Y	NR	NA	NA	6 months	Reduction of Functional, cognitive and physical decline	NR
Steinberg, 2009 ⁶¹ (19089875) US Pilot	Home-based exercise intervention program: Aerobic fitness, strength, balance & flexibility training	Home safety assessment	Community-based RCT PLWD	PLWD diagnosed with probable AD, MMSE score >=10; community-residing; ambulatory, with caregiver	N=14 exercise group 13 control group Age: yes Sex: yes % majority race: yes Education: no	NR	NR	NR	NA	NA	12 weeks	Functional performance, cognitive functioning, neuropsychiatric symptoms, caregiver burden and quality of life:	NA
Kwak, 2008 ⁶² (18050054) Korea Small sample	Individual home-based exercise 30–60 minutes per day, 2–3 times per week for 12 months	Control: NR	Community-based RCT	Women with AD,, age >60; living with relative, MMSE 10-26; able to exercise	N=30 (15 exercise, 15 controls) Age: yes Sex: yes % majority race: yes Education: no	NR	Not clear –it appears the IC were all women, but it also appears the PLWD were all women	NR	NA	NA	12 months	Cognition. cardiovascular health, and physical condition	NR
Williams, 2008 ⁶³ (17959874) US Small sample	I1: Group exercise (walking plus strength training, balance, and flexibility) I2: Supervised walking	Social conversation	Nursing home RCT	PLWD diagnosed with AD and symptoms of depression	N=116 Age: yes Sex: yes (overall) % majority race: yes (NR by group) Education: no	NR	NA	NA	Education: Y	Training: Y	16 weeks	Observed Affect Scale (OAS), Alzheimer Mood Scale, Dementia Mood Assessment	NA
Rolland 2007 ²¹ (17302650) France High RoB	Group exercise (walk, strength, balance, flexibility) 2x/week 12 months	Routine medical care	Nursing homes (5) RCT	Mild to severe AD	N=134 Age: Y Sex: Y % majority race: N Education: N	Disability: Hearing and visual impairment Other: NR	NA	NA	NR	NR	12 months	Function, Katz ADL, depression, Neuropsychiatric Inventory	NA
Stevens, 2006 ⁶⁴ (16594879) Australia Pilot	Individually prescribed exercise program of 20 to 30 minutes 3x per weeks	Two groups: 1. no intervention 2. social visit of equal length to exercise program	Nursing home-based RCT	PLWD assessed with mild to moderate dementia; MMSE score <9 and <23; not all PLWD assessed for MMSE	N=24 exercise, 21 social visit, 30 control Age: Y Sex: Y % majority race: N Education: N	NR	NA	NA	NA	NA	12 weeks	Assessment of physical and psychiatric problems, dependence on nursing care	NA

Van de Winkel 2004 ⁶⁵ (15137556) Belgium Small sample	Music-based exercises	Social conversation	RCT Institutionalized	PLWD with MMSE score lower than 24; able to respond to verbal and visual commands; capable of responding to music	N=15 PLWD exercise group 10 PLWD control group Age: N Sex: N % majority race: N Education: N	NR	NA	NA	NA	NA	3 months	Measures of cognition; need for help; aggressiveness; depression; level of activity	NA
Toulotte, 2003 ⁶⁶ (12540351) France Small sample	Exercise: chair-assisted or standing supervised exercise 1 hour, 2x/week 16 weeks	Usual daily routine	Home and institutionalized PLWD Crossover RCT	Demented elderly people with a history of falling	N=20 Age: yes Sex: yes % majority race: no Education: no	NR	NA	NA	NA	NA	16 weeks	Walking, mobility, flexibility and static balance	NA

Abbreviations: AD=Alzheimer's Disease; ADL=Activities of Daily Living; AE=adverse events; BADL=Bristol Activities of Daily Living Scale; BBS=Berg Balance Scale; CDR=Clinical Dementia Rating; BPSD=Behavioral and Psychological Syndrome of Dementia; CSD=Cornell Scale for Depression; CST=Chair Stand Test; E-ADL=Erlangen-ADL (instrumental ADLs); EQ-5D-3L=Euroqol 5 dimension-3 level; EQ-5D-5L=Euroqol 5 dimension-5 level; FIM=Functional Independence Measure; FINALEX=Finnish Alzheimer's Disease Exercise Trial; GDS=Geriatric Depression Scale; GHQ=Global Health Questionnaire; HIFE=high-intensity functional exercise; IC=informal caregiver; MMSE=Mini Mental State Exam; N=number; NINCDS-ADRDA=National Institute of Neurological and Communicative Disorders and Stroke/Alzheimer's Disease and Related Disorders Association; NPI=Neuropsychiatric Inventory (10 items); NPI-12=Neuropsychiatric Inventory-12 items (with sleep and appetite); NR=Not Reported; PAQE=Physical Activity Questionnaire for the Elderly; PPT-7=7-item Physical Performance Test; POMA=Performance Oriented Motor Assessment (gait and balance); PMID=PubMed Identification Number; PLWD=Persons with Dementia; QUALID=quality of life in late-stage dementia scale; RCT=Randomized controlled trial; ROB=risk of bias; SAE=serious adverse event; SCL-90=Symptom Checklist-90; SD=standard deviation
SES=socioeconomic status; SPPB=Short Physical Performance Battery; TMT=Trail Making Test; TUG=Timed Up and Go; UC=usual care; vs=versus

Music Intervention

Table D-6. Risk of bias assessment: music intervention

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Ugur, 2019 ⁶⁷ (30497077)	3 months	Medium 14%	Low	Medium	High	Low	High	University	High
Cheung, 2018 ⁶⁸ (30235949)	6 weeks	Low 6 weeks: 19.39% 12 weeks: 24.84%	Low	Low	Moderate	Low	High	No funding	Medium
Ho, 2018 ⁶⁹ (29468887)	8 weeks	Low 0%	Medium	High	High	Low	Low	The Tung Wah Groups of Hospitals	High
Kwak 2018 ⁷⁰ (29871544)	14 weeks	Low 0%	Low	Medium	Medium	Low	Moderate	Nonprofit, University, Government	Medium
Pongan, 2017 ⁷¹ ((28922159)	12 weeks & 20 weeks	Low 20%	Medium	Low	Medium	Low	High	Nonprofit, Government	High
Wang, 2017 ⁷² (26443002)	24 weeks	Medium 13.37%	High	Medium	High	Low	High	Unclear	High
Sarkamo, 2014 ⁷³ (24009169)	3 months	Low 3 months: 5.61% 9 months (6months follow up post intervention): 16.85%	Low	Low	High	Low	High	Nonprofit, Government	Medium
Vink, 2013 ⁷⁴ (23280604)	4 months	Low 18%	Medium	Medium	Medium	Low	High	Nonprofit	High
Sung, 2012 ⁷⁵ (21823174)	6 weeks	Low 8.33%	Low	Medium	High	Low	High	Government	Medium
Lin, 2011 ⁷⁶ (20672256) #2	6 weeks	Low 6 weeks: 3.84% 10 weeks: 3.84%	Low	Medium	High	Low	High	NR	Medium
Chu, 2014 ⁷⁷ (23639952)		Low	Low	Medium	Medium	Low	Low	The Taipei Medical University Hospital	Medium
Raglio, 2008 ⁷⁸ (18525288)	16 weeks 20 weeks post intervention	Low 3.38%	Medium	Low	High	Low	High	NR	High
Ledger, 2007 ⁷⁹ (17558584)	1 Year	High 25%	High	X	X	X	X	No funding	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number; #2=Companion

Table D-7. Characteristics of included studies: music intervention

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Cheung 2018 ⁸⁰ (30235949) Hong Kong Medium Explanatory	PLWD met 30 mins, twice a week for six weeks to receive music intervention that involved listening to their preferred music and moving their limbs and trunk. The first author facilitated the intervention.	(1) Music Listening: PLWD in this group listened to their preferred music (2) Social activity: chatted casually, twice a week for six weeks.	Residential care facilities RCT multisite 165 PLWD	Any type of dementia with moderate severity according to Global Deterioration Scale (stage 5 or 6)	N=165 82.27 years 75.8% Female 54.5% had no formal education	NR	NA	NA	NA	NA	6 weeks 12 weeks	Agitation: CMIA-Nursing home version (Chinese version)	NA
Cheung 2018 ⁶⁸ (27819483) Taiwan Medium Explanatory	PLWD met 30 mins, twice a week for six weeks to receive music intervention that involved listening to their preferred music and moving their limbs and trunk. The first author facilitated the intervention.	(1) Music Listening: PLWD in this group listened to their preferred music (2) Social activity: chatted casually, twice a week for six weeks	Residential care facilities RCT multisite 165 PLWD	Any type of dementia with moderate severity according to Global Deterioration Scale (stage 5 or 6)	N=165 82.27 years 75.8% Female 54.5% had no formal education	NR	NA	NA	NA	NA	6 weeks 12 weeks	Anxiety: The Chinese RAID Depressive symptoms: GDS	NA
Kwak 2018 ⁷⁰ (29871544) USA Low Explanatory	Music & memory: In addition to treatment as usual, personalized music playlists delivered on digital music players over a 6-week period followed by 2-week washout period & 6 weeks of treatment as usual only	Condition 2: 6 weeks of treatment as usual only followed by 2-week washout period and then music and memory for 6 weeks	Nursing homes RCT (Crossover) 59 PLWD	Alzheimer's disease or other dementia	N=59 86.9 years 77.96% Female 93% White Education: No	NR	NA	NA	NA	NA	14 weeks	Agitation: CMAI Dementia-related cognitive-behavioral issues: NPI-NH Medication use: Standardize record form	NA

Sarkamo, 2014 ⁷³ (24009169) Finland Medium Explanatory # 2 Sarkamo, 2016 ⁸¹ (26519435)	Group Music: (1) Singing coaching group: In a group of 10, PLWD participated in primarily singing familiar songs coupled occasionally with vocal exercises and rhythmic movements led by a trained music teacher once a week for 10 weeks (2) Listening coaching group: In a group of 10, led by a therapist, PLWD primarily listened to songs from CD and discussed about the emotions, thoughts, and memories, once a week for 10 weeks	Usual care: PLWD continued with their normal everyday activities and hobbies without & were not given any additional activities throughout the follow-up.	Day activity centers & inpatient centers RCT 89 Dyads (59 family members & 30 nurses)	All types of dementia (diagnosis made by a geriatrician or a primary care Physician) Mild to moderate (CDR)	N=84 Age: 89.6 years 71.42% Female 54.5% had no formal education	NR	59 IC (family members) randomized; No details provided on how many completed or their demographic details	NR	30 nurses randomized No details provided on how many completed and their demographic details	NR	10 weeks 9 months	PLWD Mood & QoL: CBS & QOL-AD Psychological well-being of CGs: GHQ and ZBI	NA
Sung 2012 ⁷⁵ (21823174) Taiwan Medium Explanatory	Group music: PLWD received 30-min music intervention using percussion instruments with familiar music in a group setting, delivered by trained research assistants in midafternoon twice weekly for 6 weeks,	PLWD in the control group received usual care and did not attend the group music interventions.	Residential care facility RCT 60 PLWD	Any type of dementia with mild to severe severity (the Short Portable Mental Status Questionnaire)	N=55 Average age 81.37 years for the control group and 97.5 years for the control group 65.8% Female 76.2% had no formal education	NR	NA	NA	NA	NA	6 weeks	Anxiety: RAID Agitation: CMAI	NA
Lin 2011 ⁷⁶ (20672256) Taiwan Medium Explanatory 2 ⁷⁷ (23639952)	PLWD received a total of twelve 30-min group music intervention sessions, conducted twice a week for six consecutive weeks by the researcher who underwent training	PLWD in control group continued to engage in their normal Daily activities.	Nursing facilities RCT multisite 104	All dementia types (DSM-IV-TR) with mild-severe severity (C-MMSE-<24)	N=100 82 years 53% Female 100% Chinese Education Mean NR	NR	NA	NA	NA	NA	6 weeks 10 weeks	Agitation: Chinese Version of the Cohen-Mansfield Agitation Inventory (C-CMAI). Depression: C-CSDD	NA

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; C-CMAI=The Chinese Version of the Cohen-Mansfield Agitation Inventory; C-CSDD=The Chinese version of Cornell Scale for Depression; GDS=Geriatric Depression Scale; RAID=The Rating Anxiety in Dementia; CBS=the Cornell-Brown Scale for Quality of Life; QOL-AD=the Quality of Life in Alzheimer's Disease; QOL=Quality of Life; GHQ=The General Health Questionnaire; ZBI=Zarit Burden Interview; RoB=Risk of Bias; NPI-NH=the Neuropsychiatric Inventory-Nursing Home Version; CDR=Clinical Dementia Rating; CG=Caregiver; RCT=Randomized controlled trial; MMSE=Mini-Mental State Exam

Table D-8. PLWD outcomes summary low and medium risk of bias studies: music intervention

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Cheung, 2018 ⁶⁸ (27819483) Music with movement vs. music listening and social activity Medium Explanatory	Adjusted Mean Difference (95% CI) Anxiety (RAID- Chinese version) 6 weeks Depressive symptoms (GDS) 6 weeks	No useable outcome reported No useable outcome reported			
Cheung, 2018 ⁸⁰ (30235949) Music with movement vs. music listening and social activity Medium Explanatory	Adjusted Mean Difference (95% CI) Agitation (CMAI-NH) 6 weeks	Group x time interaction do not favor intervention F(df)=1.22 (4, 324)			0.303
Sarkamo, 2014 ⁷³ (24009169) Music vs. usual care Medium Explanatory	Quality of life ((QOL-AD) – administered in an interview format 3 months Adjusted Mean Difference (95% CI) Mood (CBS) – administered in an interview format	Group x time interaction favors intervention F(df)=12.9 (1, 81) Group x time interaction favors intervention F(df)=4.6 (1, 62)			p=.001 p=.036
Sung, 2012 ⁷⁵ (21823174) Group music vs. usual care Medium Explanatory	Adjusted Mean Difference (95% CI) Agitation (CMAI) 6 weeks Adjusted Mean Difference (95% CI) Anxiety (RAID) 6 weeks	Group x time interaction do not favor intervention F(df)=0.33 (1, 51) The outcome favors intervention: Mean difference (95% CI): 3.77 (1.25 to 6.3)			0.95 p=0.004
Lin, 2011 ⁷⁶ (20672256) Group music vs. usual care Medium Explanatory #2 Chu, 2014 ⁷⁷ (23639952)	Adjusted Mean Difference (95% CI) Agitated behavior (C-CMAI). 6 weeks Adjusted Mean Difference (95% CI) Depression (C-CSDD) 6 weeks	The outcome favors music intervention Adjusted Mean Difference (95% CI): -0.47 (-0.74 to -0.19) -----			<0.001 <.001

Abbreviations: CBS=Caregiver Burden Scale; CMAI=Cohen Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; GDS=Geriatric Depression Scale; NR=Not Reported; PMID=PubMed Identification Number; QoL-AD=Quality of life of the person with dementia; RAID=the Rating Anxiety in Dementia Scale

Table D-9. Summary of strength of evidence for PLWD outcomes: music intervention

Outcome Comparison	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Agitation Group music vs. Usual care	6 weeks	2 RCT (n=155)	Inconsistent finding. One study reported significant reduction of agitation (C-CMAI) in the music group compared to usual care group, another reported no significant difference in the reduction of agitation scores (& CMAI) between two groups.	Moderate	Inconsistent	Direct	Imprecise	Insufficient
Agitation Music with movement vs. music listening & social activity	6 weeks	1 RCT (n=165)	No sig difference between groups in terms of reducing agitation measured using CMAI-NH.	Moderate	Unknown	Direct	Imprecise	Insufficient
Depression Group music vs usual care	6 weeks	1 RCT (100) Companion study (Lin,2011)	Significant difference between groups in the change in depression (C-CSDD) level from baseline to posttest	Moderate	Unknown	Direct	Imprecise	Insufficient
Anxiety Music with movement vs. music listening & social activity	6 weeks	1 RCT (n=165)	Results showed that the Music with movement did not differ from ML and SA in changing the symptoms of anxiety measured using RAID.	Moderate	Unknown	Direct	Imprecise	Insufficient
Mood Group music vs. usual care	3 months	1 RCT (n=84)	A significant effect of music on mood (CBS) was observed in the music group. Music was effective in temporarily improving mood.	Moderate	unknown	Direct	Imprecise	Insufficient
Quality of Life Group music vs. usual care	3 months	1 RCT (n=84)	Result was inconsistent. There was a long-term specific effect in the self-report total scores. However, statistical significance could not be established with caregiver-report QOL-AD scores. Music listening than singing, showed long term effect.	Moderate	Inconsistent	Direct	Imprecise	Insufficient

Abbreviations: CBS=Caregiver Burden Scale; CMAI=Cohen Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; GDS=Geriatric Depression Scale; NR=Not Reported; PMID=PubMed Identification Number; QoL-AD=Quality of life of the person with dementia; RAID=the Rating Anxiety in Dementia Scale

Evidence Map: Music Intervention

Table D-10. Characteristics of evidence map studies: music intervention

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Random-ized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Weise 2020 ⁸² Germany Pilot	Individualized music program: Participants listened to personally-relevant music playlists for 30 minutes every other day for four weeks	Wait-list control	NH Residents RCT 20 PLWD	Clinical diagnosis of dementia (mild to severe) and no severe hearing problems	N=20 Mean age: 85 years % Female: 80 % majority race: no Education: no	None	NA	None	NA	NA	4 weeks	CMAI	NA

Ihara 2019 ⁸³ (30460747) US Quasi-experimental	MUSIC & MEMORY program: an individualized music listening system where individuals are given an iPod programmed with their personalized playlist	Participated in daily planned activities, such as exercises and games.	Residents of community-based adult day health centers Quasi-experimental 51 PLWD	Diagnosis of dementia and MMSE score of ≤24	N=51 Mean age: 82 years % Female: 67 % majority race: no Education: no	None	NA	None	NA	NA	During the intervention 6 weeks	CMAI CSDD video recording of Behavioral observations	NA
Park, 2019 ⁸⁴ (31533443) US Pilot	Chair yoga: Practiced Hatha yoga (breathing, physical posture & guided relaxation), in a chair or standing and using a chair as support twice-weekly 45-minute sessions for 12 weeks	Music: Attended music sessions twice-weekly 45-minute sessions for 12 weeks facilitated by music therapy. Chair-based exercise: Attended 45-min gentle exercise session twice weekly for 12 weeks	Community setting RCT (Cluster) 31 PLWD	All types of dementia Any stage of severity	N=31 Age: 84.3 Sex: Yes % majority race: Yes Education: No	PLWD Household Characteristics: Yes PLWD Detailed Race Information: Yes	NA	NA	NA	NA	12 weeks	Psychological symptoms: HADS Behavioral problems: CMAI-SF QoL; sleep problems; Epworth Sleepiness Scale	NA
Ugur, 2019 ⁶⁷ (30497077) Turkey High RoB	Music Therapy: PLWD listened to instrumental classical Turkish music, 3 sessions per week in the mornings in their own rooms for 12 weeks	Control: MT was not provided for the participants in the control	Nursing homes RCT 70 PLWD	Non-specified Mild -severe	N=60 Age: 82.91 years Sex: Yes % majority race: No Education: Yes	NR	N=60 Age: Yes Sex: No % majority race: No Education: Yes	Health status: Yes	NA	NA	12 weeks	CBS & ADLS	Caregiver burden: CBS
Ho, 2018 ⁶⁹ (29468887) Hong Kong High RoB	Music Intervention: 16 half-hour sessions of music intervention with multi-sensory components over eight weeks	Control: Received standard care	Elderly residential home RCT 73	Moderate dementia (non-specified)	N=73 Age: 85.3 years Sex: Yes % majority race: Yes Education: Yes	NR	NA	NA	NA	NA	8 weeks	Behavioral and psychological symptoms: The NPI (Chinese version) Subjective mood: VAMS	NA

Giovagnoli 2017 ⁸⁵ (29550981) Italy Small sample	Active music therapy (AMT) and neuro-education (NE): involving a regulatory action of sound stimulating non-verbal communication. AMT lasted 2 45 minutes group sessions per week for 12 weeks. NE also 12 weeks	cognitive training lasted 2 45 minutes group sessions per week for 12 weeks	Recruited from a center RCT 50 PLWD	Mild to moderate dementia or AD with a MMSE score >15	N=50 Age: 74 years Sex: Yes % majority race: no Education: Yes	NR	NA	NA	NA	NA	12 weeks 24 weeks	Word Fluency on phonemic cue Attentive Matrices Trail Making Test A/B Weigl Sorting Test Short Story test	
Pongan, 2017 ⁷¹ (28922159) France High RoB	Choral singing: participated in 12 weekly two-hour singing sessions	Painting: participated in 12 weekly two-hour painting sessions	Community setting RCT 65 PLWD	Probable AD with mild severity	N=59 Age: Yes Sex: Yes % majority race: No Education: Yes	NR	NA	NA	NA	NA	12 weeks	Chronic pain: NRS, SVS & BPI Anxiety: STAI Depression: GDS Quality of life: EQ-5D	NA
Wang, 2017 ⁷² (26443002) Taiwan High RoB	Music: received Kagayashiki music care (KMC) twice per week for 24 weeks.	Usual care: provided with activities as usual	Long-term care facility Quasi 172 PLWD	Any type of dementia Mild - severe	N=147 Age: Yes Sex: Yes % majority race: No Education: Yes	NR	NA	NA	NA	NA	24 weeks	Behavior problem: CAPE-BRS Depression: CSD	NA
Narme, 2014 ⁸⁶ (23969994) France (<25 in each arm)	Music Intervention: Listened to Classical instrumental; familiar songs from the 1950–80s, 1 hour twice a week for 4 weeks	Cooking: Participants engaged in cooking activities	Nursing home RCT 48 PLWD	Moderate or severe Alzheimer's type or mixed dementia	N=37 Age: Yes Sex: Yes % majority race: No Education: Yes	NR	NA	NA	NA	NA	4 weeks	Emotional state: Interview Mood: STAI-A, CMAI & NPI FC distress: Distress scale of NPI	NA
Ridder, 2013 ⁸⁷ (23621805) Norway Pilot	Individual music therapy: Individual music therapy was given biweekly over a period of six weeks, altogether 12 sessions	Control: Received Standard Care (for some includes group sing-along sessions as usual)	Nursing home RCT(Crossover) 42 PLWD	Moderate to severe dementia	N=42 Age: Yes Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	6 weeks	CMAI-NH; the ADRQL; & psychotropic medication	NA

Sakamoto, 2013 ⁸⁸ (23298693) Japan Small sample	Music therapy - Passive OR interactive: Each intervention was performed for 30 min once a week for 10 weeks (10 sessions in total). Passive: participants passively listened to the selected music via a CD player. Interactive: participants not only listened to the selected music via a CD player but also participated in interactive activities (e.g., clapping, singing, & dancing) guided by a music facilitator	Control- No music: participant spent time with one caregiver in their own room as usual, without any music intervention	Nursing home RCT 39 PLWD	Severe Alzheimer's Type dementia	N=39 Age: Yes Sex: Yes % majority race: Yes Education: No	NR	NA	NA	NA	NA	10 weeks	BPSD changes: BEHAVE-AD	NA
Vink, 2013 ⁷⁴ (23280604) Netherlands High RoB	Music therapy: Each music therapy intervention lasted for 40 min and was provided twice weekly, by a formally trained music therapist with at least 5 years working experience.	General recreational day activities: 40 minutes of general daily recreational activities	Nursing homes RCT 94 PLWD	Any type of dementia with mild to very severe severity	N=77 Age: Yes Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	4 months	CMAI	NA
Ceccato, 2012 ⁸⁹ Italy Small sample	The Sound Training for Attention & Memory in Dementia (STAM-Dem): Music employed as stimulus. Followed the STAM-Dem for 2 weekly sessions of 45 minutes for 12 weeks (in addition to standard care).	Standard care: Continued with the normal "standard care" provided	RCT 51 PLWD	NR	N=50 Age: Yes Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	12 weeks	Cognitive function: MPI, MPD Mood: GDS Aggressive behavior: CMAI Functional status: ADL	NA

McHugh, 2012 ⁹⁰ USA Pilot study	Music therapy - vocal re-creative music therapy (VMT): residents participated in 25 mins of music session just before their mid-day meal, 4 times a week for 3 weeks.	Control wait-list group: Unclear	Care facility RCT 15 PLWD	Moderate to severe dementia (primarily AD)	N=15 Age: Yes Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	3weeks	Nutritional intake: Care-tracker by Ingenix	NA
Chang et. al, 2010 ⁹¹ (20492038) Small sample (<25 in each arm)	Music(background) was administered over the institution's internal broadcast network during the second week of the study over the lunchtime (11:00–12:00).	No music (time series)	Institution housing ITS 47 PLWD	AD, vascular and other dementia Mild	N=41 Age: 81.68 years Sex: Yes % majority race: No Education: Yes	NR	NA	NA	NA	NA	8 weeks	Dementia-related problem behavior: CMAI (Chinese version)	NA
Cooke, 2010 ⁹² (20603300) Australia Small sample #2 Cooke, 2010 ⁹³ (20635236)	Live music: Participated in a 30 mins of musician-led familiar song singing (with guitar accompaniment) and 10 mins of pre-recorded instrumental music for active listening, three times a week for eight weeks,	Reading: Engaged in a range of reading/ social activities including reading local news stories, short stories, telling jokes & undertaking quiz activities.	Aged care facility RCT(crossover) 47 PLWD	Early to mid-stage dementia or probable dementia	N=47 Age: Yes Sex: Yes % majority race: No Education: Yes	NR	NA	NA	NA	NA	8 Weeks	DQOL; GDS; #2 CMAI-SF; RAID;	NA
Han, 2010 ⁹⁴ (21252549) Singapore (<25 in each arm)	Music therapy and activity program: Attended a weekly group MAP conducted by a qualified music therapist and occupational therapist for once a week for 8 weeks.	Waitlist	Outpatient dementia clinic Non-RCT 45 PLWD	Alzheimer's disease or vascular dementia with moderate severity	N=43 Age: Yes Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	8 weeks	Mood & disruptive behavioral: RMPCP & AES	NA

Raglio 2010 ⁷⁸ (20735342) Italy Pilot	Music therapy: received three cycles of 12 sessions each, three times a week. Each session involved a group of three patients and lasted 30 minutes	Standard care (i.e. educational and entertainment activities such as reading a newspaper, performing physical activities, etc.).	Nursing home RCT 60 PLWD	AD diagnosis based on DSM-IV and MMSE between 18 and 30	N=60 Age: Yes Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	4 weeks 8 weeks	MMSE Barthel index NPI	NA
Sung 2010 ⁹⁵ (20492050) Taiwan Quasi-experimental	30-minute music listening intervention based on personal preferences delivered by trained nursing staff, twice a week for six weeks.	Standard care without music	Long term care facility RCT 60 PLWD	Clinical diagnosis of dementia and moderate to severe cognitive decline with GDS score between 4 and 6	N=52 Age: Yes Sex: Yes % majority race: Yes Education: Yes	NR	NA	NA	NA	NA	6 weeks	RAID	NA
Guetin, 2009 ⁹⁶ (19628939) France (<25 in each arm)	Music therapy: Participated in weekly sessions of individual, receptive music therapy	Reading: Participated under the same conditions in reading sessions.	Nursing homes RCT 30 PLWD	AD with no specified severity	N=30 Age: Yes Sex: Yes % majority race: No Education: Yes	Household Characteristics: Yes Age of Diagnosis: Yes Rest: No	NA	NA	NA	NA	16 weeks	Anxiety: the Hamilton Scale Depression: GDS	NA
Choi, 2008 ⁹⁷ South Korea Pilot	Group music: received 50 minutes of music intervention (singing songs, analysis of libretto, making musical instruments, song writing & playing instruments) 3 times per week for 5 consecutive weeks.	Usual care: received usual care, and did not participate in any structured therapeutic programs during the study period.	Special dementia care unit Non-RCT 20 PLWD	AD, vascular and other type	N=20 Age=74.9 years Sex: No % majority race: No Education: No	NR	NA	NA	NA	NA	5 weeks	Depression: GDS Quality of life: GQoL Behavioral disturbances: NPI-Q	NA
Raglio, 2008 ⁷⁸ (18525288) Italy High RoB	Music Therapy: received 3 cycles of 10 MT sessions (30 min/session) over 16 weeks	Control: received educational support or entertainment activities.	Nursing home RCT 59 PLWD	Mild Alzheimer type or vascular or mixed dementia	N=59 Age: Yes Sex: Yes % majority race: No Education: Yes	NR	NA	NA	NA	NA	16 weeks	BPSD	NA
Ledger, 2007 ⁷⁹ (17558584) Australia High RoB	Group music therapy: Received weekly music therapy for	Standard care: Received standard nursing home care	Nursing home Non-RCT 60 PLWD	Mild or moderate or moderately severe Alzheimer's type	N=45 Age: 71-100 years Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	42 weeks	Agitation; CMAI	NA

Holmes, 2006 ⁹⁸ (16805928) UK Pilot	Music Therapy: Participants listened to either live or prerecorded music for 30 mins	Silence: 30 mins of silence	Residential-care or nursing-home facility RCT 32 PLWD	Moderate or severe dementia with apathy	N=32 Age: Yes Sex: No % majority race: No Education=No	NR	NA	NA	NA	NA	Unclear	Engagement: DCM	NA
Sung, 2006 ⁹⁹ (16765849) Taiwan Small sample	Group music with movement intervention: received 30-min group music with movement intervention in the afternoon twice a week over a 4-week period by a nursing researcher and two research assistants trained in music intervention	Control: Received usual care without intervention	Residential care facility RCT 36 PLWD	Moderate to severe dementia	N=36 Age: Yes Sex: Yes % majority race: No Education: Yes	NR	NA	NA	NA	NA	4 weeks	CMAI; & RAID	NA
Svansdottir, 2006 ¹⁰⁰ (16618375) Iceland Small sample	Music therapy: received 18 sessions of music therapy, each lasting 30 minutes, three times a week for 6 weeks	Usual care: had no change of care.	Nursing homes and psychogeriatric wards. RCT 46 PLWD	Moderate or severe AD	N=47 Age: Yes Sex: No % majority race: No Education: No	NR	NA	NA	NA	NA	6 weeks	Behavioral & psychological symptoms: BEHAVE-AD	NA
Gerdner, 2000 ¹⁰¹ (10798453) Australia (<25 in each arm)	G1- Individualized music: Received individualized music for 6 weeks followed by a 2-week="washout" period and 6 weeks of classical"relaxation" music.	G2-Classical "relaxation" music: Received classical relaxation music for 6 weeks followed by a 2-week"washout" period and 6 weeks of individualized music	Long term care facility RCT (crossover) 45 PLWD	Mild to severe dementia	N=39 Age: Yes Sex: Yes % majority race: Yes Education: No	PLWD Detailed Race Information: Yes	NA	NA	NA	NA	6 weeks	Agitation: MCMAI	NA
Groene, 1993 ¹⁰² USA (<25 in each arm)	Mostly Music therapy: Each participant received one session per day for 7 days (five sessions of music followed by two sessions of reading)	Mostly reading session: Each participant received one session per day for 7 days (five sessions of reading followed by two sessions of music)	Health care facility RCT 30 PLWD	Late moderate to severe stages of Alzheimer's disease	N=30 Age: Yes Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	7 days	Wandering Behaviour: researcher and nursing staff recorded the wandering behavior of each participant	NA

Lord, 1993 ¹⁰³ (8483655) USA (<25 in each arm)	(1) Music: given six 30-min. music sessions during which music of the "Big Bands" of the 1920s and 1930s was played each week (2) Puzzle exercise: were given puzzle exercises during their activity sessions	Control: received no special treatment, but instead was involved in the usual recreational activities of drawing, painting, and watching television.	Nursing care facility RCT 60 PLWD	Alzheimer disease	N=60 Age: Yes Sex: Yes % majority race: No Education: No	NR	NA	NA	NA	NA	6 months	Social interaction, mood & Mental state: Researcher generated questionnaire	NA
--	--	--	---	-------------------	--	----	----	----	----	----	----------	---	----

Abbreviations: ADLS=Activities of Daily Living Scale; ADRQL=The Alzheimer's Disease-Related Quality of Life; AES=the Apparent Emotion Scale; BEHAVE-AD=The Behavioral Pathology in Alzheimer's Disease; BPI=The Brief Pain Inventory; BPSD=Behavioral and Psychological Syndrome of Dementia; CAPE-BRS- The Elderly Behavior Rating Scale; CBS=Caregiver Burden Scale; CDR=Clinical Dementia Rating; CMAI=Cohen Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; DCM=Dementia care mapping; DQOL=Dementia Quality of Life; EM=Evidence Map; DSM-IV=Diagnostic and Statistical Manual of Mental Disorders, 4th Edition; EQ-5D=The EuroQol-5 dimensions; GQoL=Geriatric Quality of Life; FC=Formal caregiver; GDS=Geriatric Depression Scale; GDS=Global Deterioration Scale; HADS=the Hospital Anxiety and Depression Scale; IC=Informal caregiver; ICEA-D=the Inventory to Assess Communication, Emotional Expression and Activity in Dementia; MMSE=Mini-mental state examination; N=number; NA=Not Applicable; NIA=National Institute on Aging; NPI=Neuropsychiatric Inventory; NR=Not Reported; PPI & MPD= Immediate and Deferred Prose Memory test; PLWD=Persons with Dementia; RoB=Risk of Bias; RA=Research Assistant; RAID=the Rating Anxiety in Dementia Scale; RCT=Randomized controlled trial; RMBCP=the Revised Memory and Behavioral Problems Checklist; STAI-I=The State-Trait Anxiety Inventory for Adults; VAMS=The Visual Analog Mood Scale; ZBI=Zarit Burden Interview

Reminiscence Therapy

Table D-11. Risk of bias assessment: reminiscence therapy

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Li 2019 ¹⁰⁴ (31645180)	12 weeks 24 weeks	Low 5.6%	Low	Low	Medium	Low	Medium	Government	Medium
Lok 2019 ¹⁰⁵ (30246408)	8 weeks	Medium 15.5%	Medium	High	High	Medium	X	NR	High
Lin 2018 ¹⁰⁶ (28881430)	10 week 22 weeks	Low 10 weeks: 2.6% 22 weeks: 9.3%	High	High	Low	Low	X	Government	High
Amieva 2016 ¹⁰⁷ (26572551)	3 months 24 months	Medium 3 months: 10.8% High 24 months: 30.3%	Low	Low	Low	Medium	Medium	Government	Medium (3 months) High (24 months)
Duru Asiret 2016 ¹⁰⁸ (26251112)	12 weeks	Low 6%	High	X	X	X	X	NR	High
Van Bogaert 2016 ¹⁰⁹ (27511740)	9 weeks	Medium 16.7%	Medium	Low	High	Low	X	Foundation, Government	High
Wu 2016 ¹¹⁰ (25965388)	6 weeks	Low 2.8%	Low	High	Low	Low	X	Government	High
O'Shea 2014 ¹¹¹ (24633858)	2 years	Medium 17%	Low	Low	High	Low	X	Government	High
Seranni Azcurra 2012 ¹¹² (23429813)	3 months 6 months	Low 3.7%	Medium	Low	Low	High	X	NR	High
Woods 2012 ¹¹³ (23211271)	3 months 10 months	Medium 3 months: 19.1% High 10 months: 28.3%	Low	Low	Low	Medium	Medium	Government	Medium (3 months) High (10 months)
Hsieh 2010 ¹¹⁴ (NA)	3 months	High 33.3%	X	X	X	X	X	Government	High
Wang 2007 ¹¹⁵ (17503545)	8 weeks	Medium NR	Medium	Low	Medium	Low	Medium	Government	Medium
Lai 2004 ¹¹⁶ (15190995)	6 weeks 12 weeks	Medium 15%	Medium	Low	Low	High	X	NR	High
Camberg 1999 ¹¹⁷ (10203120)	4 weeks	Low 0%	Medium	High	Low	Low	X	NR	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table D-12. Characteristics of included studies: reminiscence therapy

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Li 2019 ¹⁰⁴ (31645180) China Medium Explanatory	Group reminiscence therapy in an antique setting; 30 to 45-minute sessions, 2 sessions/week for 12 weeks	Usual care	Geriatric hospital RCT, single site 90 PLWD	Diagnosis of AD based on NINCDS-ADRDA criteria; Mild-to-moderate stage of dementia based on CDR score of 1 or 2	N=90 83 years 45% Female Race NR 90.5% Primary education and above	NR	NA	NA	NA	NA	12 weeks 24 weeks	ADAS-Cog NPI CSDD Barthel index	NPI-Caregiver Distress	
Amieva 2016 ¹⁰⁷ (26572551) France Medium Explanatory	Group reminiscence therapy; 90-minute weekly sessions for 3 months and monthly maintenance sessions for 21 months	Usual care	Memory centers or geriatric day care units RCT; multisite 326 PLWD	Mild to moderate AD diagnosis based on NINCDS-ADRDA criteria, MMSE from 16 to 26, and GDS score of 2 to 5	N=326 79 years 60% Female Race NR 49% Secondary education and above	NR	N=326 Age: NR Sex: NR Race: NR Education: NR Relation to PLWD: NR	NR	NA	NA	3 months 24 months	ADAS-Cog NPI DAD Apathy Inventory MADRS QOL-AD RUD Lite	ZBI	
Woods 2012 (23211271) ¹¹³ UK Medium Explanatory Woods 2009 ¹¹⁸ (19642992) Woods 2016 ¹¹⁹ (27093052)	Group reminiscence therapy for PLWD-informal caregiver dyads; 2-hour weekly sessions for 12 weeks, with monthly maintenance for 7 months	Usual care	Community-based setting RCT; multisite 488 PLWD-informal caregiver dyads	Mild to moderate dementia (any type) based on DSM-IV criteria and CDR	N=488 76 years 50% Female 95% White Education: NR	NR	N=488 70 years 67% Female 95% White Education NR 71% Spouse	NR	NA	NA	3 months 10 months	QoL-AD AMIE CSDD RAID Bristol Activities of Daily Living EQ-5-D Resource Utilization	GHQ QCPR HADS RSS EQ5-D	
Wang 2007 ¹¹⁵ (17503545) Medium Taiwan Explanatory Wang 2009 ¹²⁰ (18930560)	Group reminiscence therapy; 60-minute session one time/week for 8 weeks	Usual care	Elderly care facilities RCT; multisite 102 PLWD	Mild to severe dementia, CDR score of 1 to 3	N=102 79 years 51% Female Race NR 5.2 years education	SES: yes	NA	NA	NA	NA	8 weeks	MMSE GDS-SF CSDD	NR	

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; N=number; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; ; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RS=Reporting Status; SES=socioeconomic status; MMSE=Mini-Mental State Exam; ADAS-Cog=The Alzheimer's Disease Assessment Scale-Cognitive Subscale; NPI=Neuropsychiatric Inventory; CSDD=Cornell Scale for Depression in Dementia; DAD=Disability Assessment for Dementia; MADRS=(Montgomery-Asberg Depression Rating Scale; QoL-AD=Quality of Life - Alzheimer's Disease scale; RUD=resource utilization; GHQ-28=General Health Questionnaire-28 item version; QoL-AD=Quality of life of the person with dementia; AMI(E)=of the autobiographical memory interview; QCPR=the quality of carer-patient relationships; HADS=the Hospital Anxiety and Depression Scale; RAID=the Rating Anxiety in Dementia Scale; RSS=Relocation Stress Syndrome; EQ-5D=the European Quality of Life-5 Dimensions; GDS-SF=Geriatric Depression Scale Short Form. ; ZBI=Zarit Burden Interview; NINCDS/ADRDA=National Institute of Neurological and Communicative Disorders and Stroke/ Alzheimer's Disease and Related Disorders Association; RCT=Randomized controlled trial; CDR=Clinical Dementia Rating; CSDD=Cornell Scale for Depression in Dementia; ZBI=Zarit Burden Interview.

Figure D-1. PLWD outcomes summary low and medium risk of bias studies: activities of daily living, BPSD, depression, and quality of life at 3 months




Table D-13. PLWD outcomes summary low and medium risk of bias studies: reminiscence therapy

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Li 2019 ¹⁰⁴ (31645180) Reminiscence Therapy vs. Usual Care Medium Explanatory	Barthel Index Mean (SD) 6 months	No difference between groups.	55.47 (18.96)	57.07 (17.89)	NR
Li 2019 ¹⁰⁴ (31645180) Reminiscence Therapy vs. Usual Care Medium Explanatory	NPI Mean (SD) 6 months	No difference between groups.	22.21 (10.63)	25.57 (12.45)	NR
Li 2019 ¹⁰⁴ (31645180) Reminiscence Therapy vs. Usual Care Medium Explanatory	CSDD Mean (SD) 6 months	Favors intervention.	0.84 (1.41)	2.48 (2.41)	NR
Amieva 2016 (26572551) Reminiscence Therapy vs. Usual Care Medium Explanatory	Apathy Inventory Mean (SD) 3 months	NA	11.8 (13.1)	10.4 (11.8)	0.69
Amieva 2016 ¹⁰⁷ (26572551) Reminiscence Therapy vs. Usual Care Medium Explanatory	AGGIR Mean (SD) 3 months	NA	7.1 (8.7)	6.63 (7.5)	0.49
Amieva 2016 ¹⁰⁷ (26572551) Reminiscence Therapy vs. Usual Care Medium Explanatory	RUD-Lite Mean (SD) 3 months	NA	2230.1 (3301.8)	2259.3 (3078.3)	0.89
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	RAID (Proxy) Mean (SD) 3 months	Pooled Mean Difference (95% CI) 1.2 (-0.0364, 2.807)	8.44 (6.92)	7.87 (6.45)	NR
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	QCPR Mean (SD) 3 months	Pooled Mean Difference (95% CI) 0.99 (-1.460, 2.424)	57.89 (6.52)	57.37 (6.71)	NR

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; QCPR=the quality of carer-patient relationships; RAID=Rating for Anxiety in Dementia; RUD=resource utilization; GAGGIR: Grille d'Autonomie Gérontologique-Groupes Iso-Ressources (standardized dependency scale in France)

Table D-14. Caregiver outcomes summary low and medium risk of bias studies: reminiscence therapy

Study (PMID) Comparison Caregiver Type RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Li 2019 ¹⁰⁴ (31645180) Reminiscence Therapy vs. Usual Care Medium Explanatory	NPI-Caregiver Distress Mean (SD) 3 months, 6 months	No difference between groups.	3 months: 12.19 (5.1) 6 months: 10.02 (4.5)	3 months: 12.36 (4.51) 6 months: 11.43 (4.69)	NR
Amieva 2016 ¹⁰⁷ (26572551) Reminiscence Therapy vs. Usual Care Medium Explanatory	ZBI Mean (SD) 3 months	NA	31.65 (27.5)	30.05 (25.6)	0.70
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	GHQ-28 Mean (SD) 3 months	Pooled Mean Difference (95% CI) 0.021 (-0.080, 0.122)	22.67 (11.8)	22.9 (10.37)	NR
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	EQ 5D-utility Mean (SD) 3 months	Pooled Mean Difference (95% CI) 0.01 (-0.031, 0.051)	0.76 (0.23)	0.75 (0.23)	NR
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	EQ 5D-VAS Mean (SD) 3 months	Pooled Mean Difference (95% CI) -0.96 (-5.06, 3.13)	71.59 (20.17)	70.99 (19.23)	NR
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	RSS Mean (SD) 3 months	Pooled Mean Difference (95% CI) 0.98 (-7.65, 2.716)	22.81 (10.48)	21.14 (10.21)	NR
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	QCPR Mean (SD) 3 months	Pooled Mean Difference (95% CI) 0.82 (-3.114, 0.128)	52.45 (9.01)	53.43 (8.74)	NR
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	HADS-Anxiety Mean (SD) 3 months	Pooled Mean Difference (95% CI) 0.35 (-0.391, 1.083)	6.61 (4.33)	5.91 (4.18)	NR
Woods 2012 ¹¹³ (23211271) Reminiscence Therapy vs. Usual Care Medium Explanatory	HADS-Depression Mean (SD) 3 months	Pooled Mean Difference (95% CI) 0.11 (-0.494, 0.708)	4.4 (3.19)	3.99 (3.09)	NR

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; HADS=Hasegawa dementia scale revised; QCPR=Quality of Caregiver and Patient Relationship; EQ-5D=EuroQoL 5D scale; ZBI=Zarit Burden Interview

Table D-15. Summary of strength of evidence for PLWD outcomes: reminiscence therapy

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Reminiscence Therapy vs. Usual Care Activities of Daily Living	3 months	3 RCTs (n=802)	No difference between groups.	Moderate	Consistent	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care BPSD	3 months	2 RCTs (n=416)	No difference between groups.	Moderate	Consistent	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Depression	2 to 3 months	4 RCTs (n=1,006)	No difference between groups.	Moderate	Consistent	Indirect	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Quality of Life	3 months	2 RCTs (n=814)	No difference between groups.	Moderate	Consistent	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Anxiety	3 months	1 RCT (n=488)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care PLWD-Caregiver Relationship	3 months	1 RCT (n=488)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Apathy	3 months	1 RCT (n=326)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Dependency	3 months	1 RCT (n=326)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient

Abbreviations: RCT=Randomized controlled trial; PLWD=Persons with Dementia

Table D-16. Summary of strength of evidence for caregiver outcomes: reminiscence therapy

Comparison Caregiver Type Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Reminiscence Therapy vs. Usual Care Caregiver Burden	3 months	1 RCT (n=326)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Caregiver Distress	3 months 6 months	1 RCT (n=90)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Mental Health	3 months	1 RCT (n=488)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Quality of Life	3 months	1 RCT (n=488)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Stress	3 months	1 RCT (n=488)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Anxiety	3 months	1 RCT (n=488)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care Depression	3 months	1 RCT (n=488)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Reminiscence Therapy vs. Usual Care PLWD-Caregiver Relationship	3 months	1 RCT (n=488)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient

Abbreviations: n=number RCT=Randomized controlled trial; PLWD=Persons with Dementia

Evidence Map: Reminiscence Therapy

Table D-17. Characteristics of evidence map studies: reminiscence therapy

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)s	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Ching-Ten, 2020 ¹²¹ Taiwan Small sample	Group reminiscence therapy 1 hour per week for 8 weeks	Usual care	Nursing Home Quasi-experimental 24 PLWD	Veterans with mild dementia	N=24 Age: yes Sex: yes % majority race: no Education: yes	NR	NA	NA	NA	NA	3 months	GDS Meaning of life scale	NA
Lok 2019 ¹⁰⁵ (30246408) Turkey High RoB	Group reminiscence therapy; 60-minute sessions once/week for 8 weeks	Usual care	Nursing home RCT; single site 60 PLWD	AD diagnosis based on IWG-2 criteria, MMSE 13 to 24, CSDD > 8	N=60 Age: no Sex: yes % majority race: no Education: yes	NR	NA	NA	NA	NA	8 weeks	MMSE CSDD QOL-AD	NA
Manav 2019 ¹²² (30612511) Turkey Pilot Study	Reminiscence therapy with internet-based videos; 60 minutes a week for 3 months	Unstructured casual conversations; 25-30 minutes once a week for 3 months	Elderly care and rehabilitation centers RCT; multisite	Mild AD (DSM-V); MMSE 18 to 24	N=32 Age: yes Sex: yes % majority race: no Education: yes	SES: yes	NA	NA	NA	NA	3 months	MMSE Apathy Rating Scale	NA
Martinez 2019 ¹²³ (31453826) US Pilot Study	Online cognitive tasks with positive memory recall; 40 tasks taking 5 minutes to complete	Standard reminders without positive reinforcement	Outpatient memory clinic RCT; single site 22 PLWD	Mild AD; MMSE 18 or higher	N=22 Age: yes Sex: yes % majority race: yes Education: no	NR	NA	NA	NA	NA	3 months	Rosenberg Self-Esteem Scale MMSE GDS	NA
Laird 2018 ¹²⁴ (30206053) UK Pilot Study	Home-based personalized reminiscence using iPad app; 19 weeks	NR	Community-based (intervention at home) Quasi experimental 60 PLWD-informal caregiver dyads	Early to moderate dementia	N=60 Age: yes Sex: yes % majority race: no Education: no	NR	N=50 Age: yes Sex: yes % majority race: no Education: no Relation to PLWD: no	Living with PLWD: yes	NA	NA	19 weeks	Mutuality Scale QCPR WHO-5	Mutuality Scale QCPR WHO-5
Lin 2018 ¹⁰⁶ (28881430) Taiwan High RoB	Reminiscence therapy; 50-minute sessions once/week for 10 weeks	Usual care	Long-term care facilities Quasi experimental ; multi-site	Dementia with agitation MMSE<17 for non-high school graduates MMSE<24 for high school graduates	N=75 Age: yes Sex: yes % majority race: no Education: yes	NR	NA	NA	NA	NA	12 weeks 24 weeks	MMSE QoL-AD	NA

Duru Asiret 2016 ¹⁰⁸ (26251112) Turkey High RoB	Group reminiscence therapy; 30 to 25-minute sessions for 12 weeks	Group conversations. 20 to 25 minutes per week for 12 weeks	Quasi experimental ; single site 62 PLWD	Mild to moderate AD; MMSE from 10 to 24	N=62 Age: yes Sex: yes % majority race: no Education: yes	NR	NA	NA	NA	NA	3 months	ADL Observation Form MMSE GDS	NA
Van Bogaert 2016 ¹⁰⁹ (27511740) Belgium High RoB	Individual reminiscence therapy based on SolCos model; 2 times/week for 8 weeks	Usual care	RCT; multisite	Major neurocognitive disorder (DSM-V); MMSE 10 to 24	N=72 Age: yes Sex: yes % majority race: no Education: no	NR	NA	NA	NA	NA	8 weeks	CSDD MMSE NPI	NA
Wu 2016 ¹¹⁰ (25965388) Taiwan High RoB	Spiritual reminiscence therapy group therapy; 6 weeks	Usual care	Medical center RCT; single site 103 PLWD	Mild to moderate AD 21 <MMSE<24 for mild 13<MMSE<20 for moderate	N=103 Age: yes Sex: yes % majority race: no Education: yes	NR	NA	NA	NA	NA	6 weeks	Heath Hope Index Life Satisfaction Scale Spirituality Index of Well-being MMSE	NA
Gonzalez 2015 ¹²⁵ (25765779) Spain Sample Size	Group reminiscence therapy; 10 weekly 60-minute sessions	Wait-list control	Nursing home Quasi experimental 42 PLWD	AD (DSM-IV); MMSE less than 23; GDS 3 to 4	N=42 Age: yes Sex: yes % majority race: no Education: no	NR	NA	NA	NA	NA	10 weeks	MMSE CES-D RSES PWB	NA
Lalanee 2015 ¹²⁶ (25122521) France Sample Size	Autobiographical cognitive training reminiscence; 6 sessions	Cognitive training program	Health institutions RCT; multisite 33 PLWD	Mild to moderate AD (MMSE>18)	N=33 Age: yes Sex: no % majority race: no Education: yes	NR	NA	NA	NA	NA	6 weeks 8 weeks	Semi autographica l Memory Episodic memory GDS	NA
O'Shea 2014 ¹¹¹ (24633858) High RoB	Staff training to incorporate reminiscence into general care	Usual care activities	Nursing home Cluster RCT 18 units	PLWD with AD, and no significant sensory impairment or acute physical illness	N=304 Age: Y Sex: Y Race: Y Education: N	Prior disability	NA	NA	NR	NR	2 years	QOL-AD CMAI-AD CSDD	MZBI nurse MZBI health care assistant
Nakamae 2014 ¹²⁷ (NA) Japan Pilot Study	Six activity sessions (making rice balls) with reminiscence therapy; one session/week	Eating rice balls	Long-term care facilities RCT; multisite 36 PLWD	AD and vascular dementia; MMSE 23 or lower	N=36 Age: yes Sex: no % majority race: no Education: yes	NR	NA	NA	NA	NA	6 weeks	CSDD MOSES MMSE Vitality Index	NA
Subramaniam 2014 ¹²⁸ (24063317) Wales Pilot Study	Individual life review sessions and co-creating a life story book; 12 sessions	Person life story book created by relatives	Care homes RCT; multisite 23 PLWD	Mild to moderate dementia diagnosis (DSM criteria)	N=24 Age: yes Sex: yes % majority race: no Education: no	NR	N=24 Age: yes Sex: yes % majority race: no Education: no Relation to PLWD: yes	NR	N=68 Age: yes Sex: yes % majority race: no Education: yes	Length of services: yes	12 weeks 18 weeks	QOL-AD AMI-E GDS (Residential) QCPR	FC: ADQ
Van Bogaret 2013 ¹²⁹ (23583001) Belgium Pilot Study	Individual thematically-based reminiscence therapy; 4 weeks with two 45 min sessions/week	No intervention	Psychiatric day care, inpatient, and long-term care facilities RCT; 82 PLWD	AD based on NINCDS-ADRDA	N=82 Age: yes Sex: yes % majority race: no Education: no	NR	NA	NA	NA	NA	4 weeks	MMSE FAB NPI GDS-30 CSDD	NA

Seranni Azcurra 2012 ¹¹² (23429813) Argentina High RoB	Life-story reminiscence; biweekly sessions for 1 hour over 12 weeks	Unstructured social contact; biweekly sessions for 1 hour over 12 weeks	Nursing home RCT; multisite	AD (DSM-IV); Folstein Minimental Exam above 10	N=90 Age: yes Sex: yes % majority race: no Education: yes	NR	NA	NA	N=NR Age: no Sex: no % majority race: no Education: no Relation to PLWD: no	NR	12 weeks 6 months	SRQol SES WIB ADL	ZBI
Hsieh 2010 ¹¹⁴ (NA) Taiwan High RoB	Group reminiscence therapy; 12 sessions 40 to 50 minutes per week	No intervention	Nursing homes RCT; multisite 61 PLWD	Mild to moderate dementia (DSM-IV)	N=61 Age: yes Sex: yes % majority race: no Education: yes	NR	NA	NA	NA	NA	12 weeks	CDR GDS Apathy Evaluation Scale NPI	NA
Baillon 2004 ¹³⁰ (15481068) UK Sample Size	Reminiscence therapy; 3 one-on-one sessions, 40 minutes/ session	Snoezelen; 3 one-on-one sessions, 40 minutes/ session	Crossover RCT; 20 PLWD	Dementia diagnosis with BPSD	N=20 Age: yes Sex: yes % majority race: no Education: no	NR	NA	NA	NA	NA	4 weeks	Agitation Behavior Mapping Instrument Interact Scale	NA
Lai 2004 ¹¹⁶ (15190995) China High RoB	Life-story reminiscence; weekly 30-minute session for 6 weeks	Friendly discussions; 30-minute sessions for 6 weeks	Nursing homes RCT; multisite 101 PLWD	Dementia diagnosis (DSM-IV)	N=101 Age: yes Sex: yes % majority race: no Education: no	NR	NA	NA	NA	NA	6 weeks 12 weeks	SES WIB MMSE	NA
Politis 2004 ¹³¹ (15481065) US Pilot Studies	Kit-based activity intervention to reduce apathy; 30-minute sessions for 4 weeks	One-on-one meetings with activity therapist	Long-term care facility for PLWD RCT; single site 37 PLWD	Dementia diagnosis (DSM-IV) with apathy and a GDS score of 3 to 5	N=37 Age: yes Sex: yes % majority race: no Education: yes	NR	NA	NA	NA	NA	4 weeks	NPI-Apathy NPI ADRQRL CRAI	NA
Camberg 1999 ¹¹⁷ (10203120) US High RoB	Stimulated presence continuous play memory audio recordings	Usual care	Nursing homes RCT; multisite 54 PLWD	ARDR with agitation	N=54 Age: yes Sex: yes % majority race: yes Education: no	NR	NA	NA	NA	NA	12 weeks	Observed Agitation Scale Agitation VAS PARS WVAS FACE	NA

Abbreviations: char=characteristics; EM=Evidence Map; FC=formal caregiver; IC=informal caregiver; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RS=Reporting Status; SES=socioeconomic status; MMSE=Mini-Mental State Exam; QoL-AD=Quality of life of the person with dementia; CSDD=Cornell Scale for Depression in Dementia; GDS=Geriatric Depression Scale; QCPR=the quality of carer-patient relationships; WHO-5=The World health organization-Five Well-Being Index; ADL=Activities of Daily Living; NPI=Neuropsychiatric Inventory; DSM-IV=Diagnostic and Statistical Manual for Mental Disorders, Fourth Edition; ; BPSD=Behavioral and psychological symptoms of dementia; QCPR=Quality of Caregiver and Patient Relationship; MOSES=Multidimensional Observation Scale for Elderly Subjects; ZBI=Zarit Burden Interview

Cognitive Rehabilitation

Table D-18. Risk of bias assessment: cognitive rehabilitation

Author, Year PMID	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funding	Overall Rating
Chen 2020 ¹³² (No PMID)	1 month	Low (reported no attrition)	High	X	X	X	X	Not reported	High
Clare 2019 ¹³³ 30724405	9 months	Low 10%	Low	Low	High	Low	Medium	Government	Medium
Voigt-Radloff 2017 ¹³⁴ 28335810	16 week	Low 13%	Low	Medium	Medium	Low	Low	Foundation (Government-funded)	Low
Voigt-Radloff 2017 ¹³⁴ 28335810	26 week	Low 15%	Low	Medium	Medium	Low	Low	Foundation (Government-funded)	Low
Amieva 2016 ¹⁰⁷ 26572551	3 months	Low 10%	Low	Low	Medium	Low	Medium	Government, Foundation	Low
Amieva 2016 ¹⁰⁷ 26572551	24 months	Medium 28%	Low	Low	Medium	Low	Medium	Government, Foundation	Medium
Kumar 2014 ¹³⁵ 24982692	5 weeks	High: attrition NR	Low	High	High	Low	Medium	Government	High
Wu 2014 ¹³⁶ 24444172	6 months	High: 56%	X	X	X	X	X	Government	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table D-19. Characteristics of analytic set studies: cognitive rehabilitation

Author, Year PMID Country RoB* Type Funder	Target Intervention Group vs Not Components Frequency Duration	Target Comparison Group vs Not Components Frequency Duration	Setting Design Clusters N Randomized	PLWD Dementia Characteristics Type(s) Severity Diagnostic Criteria Age Diagnosed	PLWD Age (mean) Female % Race (% majority) Detailed Race Education (mean years)	PLWD SES Prior Disability Household Characteristics Health Insurance	Informal Caregiver N Age (mean) Female % Race (% majority) Education (mean years) Relation to PLWD	Informal Caregiver Duration Live With PLWD IC Paid Health Dementia Family History Employed Training	Outcome Timing	PLWD Outcomes Quality of Life Daily Function BPSD Adverse Events Services Other	Informal Caregiver Outcome
Clare 2019 ¹³³ 30724405 UK Medium Explanatory Government Clare 2019 ¹³⁷	PLWD Individual goal-oriented cognitive rehabilitation 1 hour/week (10 sessions in 3 months) plus 4 maintenance sessions over 6 months. 9 months Directed by trained OT or RN	Usual care	Home-based intervention by therapist RCT N=475	Types: 60% AD, vascular, mixed per ICD-10 code. Early-stage mild- moderate dementia (mean MMSE 24, enrolled MSSE ≥ 18); stable dementia medication acceptable	N=475 Mean age: 79 years 48% female 96% white race Education: 13 years	Married: 70%	N=474 Mean age: 69 yrs Female: 70% 96% white race Education: 14 years Relation: 70% spouse/partner	NR	3 months and 9 months post intervention	which are typically used in cross- sectional studies (which are typically used in cross-sectional studies which are typically used in cross-sectional studies	Stress, Quality of life
Voigt-Radloff 2017 ¹³⁴ 28335810 Germany, Netherlands Low Explanatory Government Foundation Voigt-Radloff 2011 ¹³⁸	PLWD Individual Errorless Learning (feed- forward instructions) at home 9 x 1-hour task-training sessions 8 weeks	PLWD Individual Trial & Error learning at home 9 x 1-hour task- training sessions 8 weeks	Home (community) RCT Cluster: NA N=161	Type: AD or mixed Severity: mild- moderate (MMSE 14-24), with available ICG Diagnosis: NR Diagnosis age: 1.9 years prior	N=161 77 years 57% female Race: NR More race: NA Education: 98% ≥ 9 years	SES: NR Disability: NR Household: NA Insurance: NR (Germany)	N=NR (161) Mean age: 63 Female: 36% Race: NR Education: NR Relation: NR	Duration: 25 months Live together: 67% Paid: NR Health: NR Family history: NR Employed: NR Training: NR	16 weeks, 26 weeks	QoL: QOL-AD Function: IDDD BPSD: NPI-12, AE: NR Services: NR Other: Task performance videos, cognitive, satisfaction	NR
Amieva 2016 ¹⁰⁷ 26572551 France Medium Explanatory Government Foundation ETNA3	PLWD (+ ICG support group): 1. Group cognitive rehabilitation 2. Group reminiscence 3. Individually-tailored cognitive rehabilitation (with psychologist + weekly call to ICG) All: 1.5 hours/week x 3 months, then once every 6 weeks x 21 months (24 months total) Supervision: psychologist or related	PLWD (+ ICG support group): Individual Usual care (+ ICG support group 1x/week x 3 months, then once every 6 weeks x 21 months).	Outpatient (small groups or individual). RCT Clusters: NA N=653	Type: NR Severity: mild-moderate (mean MMSE=22) Diagnosis: NINCDS- ADRDA Age diagnosed: NR 88% anti-dementia drugs:	78 years 60% female Race: NR More race: NR Education: 49% secondary school or higher	SES: NR Disability: NR Household: NR Insurance: NR (France)	N=NR Mean age: NR Female: NR Race: NR Education: NR Relation: NR	Duration: NR Live together: NR Paid: NR Health: NR Family history: NR Employed: NR Training: NR	3, 24 months	QoL: QOL-AD Function: DAD, GAGGIR BPSD: NPI AE: NR (survival rate reported) Services: RUD Lite, rate of institutionalization Other: ADAS-cog, MMSE, % without severe dementia	Apathy, ZBI

Abbreviations: ADL=Activities of Daily Living; AE=adverse events; CDR=Clinical Dementia Rating; COPM=Canadian Occupational Performance Measure; DAD=Disablement Assessment for Dementia; DEMQOL=Dementia-specific health-related quality of life; GAGGIR: Grille d'Autonomie Gérontologique-Groupes Iso-Ressources (standardized dependency scale in France); GSES=Generalized Self-Efficacy Scale; HADS=Hospital Anxiety and Depression Scale; IC=informal caregiver; IDDD=Interview for Deterioration in Daily Living Activities; MMSE=Mini Mental State Exam; N=number; NPI=Neuropsychiatric Inventory; PMID=PubMed Identification Number; OT: occupational therapist; PLWD=Persons with Dementia; RN=Registered Nurse; RoB=Risk of Bias; SES=socioeconomic status; UC=usual care; ZBI=Zarit Burden Interview; QoL-AD: Quality of Life in Alzheimer's Disease scale; E=Apathy Evaluation; NR=NR; ADAS-Cog=Alzheimer's Disease Assessment Scale - Cognition RUD=resource utilization; BPSD=Behavioral and psychological symptoms of dementia

Table D-20. PLWD outcomes summary low and medium risk of bias studies: cognitive rehabilitation

Study (PMID) Comparison RoB Category* Type	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Clare 2019 ¹³³ (30724405) Individual cognitive rehabilitation vs Usual care Medium Explanatory	DEMQOL Mean (SD)	Mean difference (95% CI) 1.08 (-0.6, 2.8)	92.36 (12.0)	92.25 (12.8)	0.22
	HADS depression Mean (SD)	Mean difference (95% CI) 0.12 (-0.4, 0.6)	4.19 (3.2)	3.83 (2.8)	0.61
	HADS anxiety Mean (SD)	Mean difference (95% CI) 0.26 (-0.3, 0.8)	5.63 (3.8) 6.05 (2.2)	4.88 (3.4)	0.33
	Self-rated goal attainment (COPM, 0-10; 2 point change=clinically significant) Mean (SD) 9 months	Mean difference (95% CI) 1.7 (1.4, 2.1)		4.22 (2.0)	<0.001
Amieva 2016 ¹⁰⁷ (26572551) Individual cognitive rehabilitation vs Usual care Medium Explanatory	QOL-AD Mean (SD)	NR	29.05 (9.2)	28.83 (9.5)	0.94
	DAD (ADLs) Mean (SD)	NR	27.04 (11.9)	25.38 (13.4)	0.39
	GAGGIR (dependency) Mean (SD)	NR	12.73 (11.6)	15.21 (11.5)	0.025
	NPI Mean (SD) 24 months	NR	34.44 (32.8)	39.31 (32.3)	0.081
Voigt-Radloff 2017 ¹³⁴ (28335810) Individual: Errorless Learning (feed-forward instructions) vs Trial & Error Learning Low Explanatory	IDDD (ADL performance) Mean (SD)	Difference (CI) 1.4 (-2.1, 4.8)	22.3 (10.2)	23.6 (10.1)	NR
	NPI Mean (SD)	Difference (CI) 0.6 (-1.5, 2.7)	8.0 (5.4)	8.6 (6.5)	NR
	Task performance rating video (A) Mean (SD) (1=worst, 7=best)	Difference (CI) 0.3 (-0.5, 0.8)	3.8 (1.8)	4.0 (1.7)	NR
	Task performance rating video (B) Mean (SD) (1=worst, 7=best) 26 weeks	Difference (CI) -0.2 (-0.9, 0.5)	4.1 (2.0)	3.9 (1.9)	NR

Abbreviations: PMID=PubMed Identification Number; ADL=Activities of Daily Living; CI=Confidence Interval; COPM=Canadian Occupational Performance Measure; DAD=Disablement Assessment for Dementia; DEMQOL=Dementia-specific health-related quality of life; GAGGIR: Grille d'Autonomie Gérontologique-Groupes Iso-Ressources (standardized dependency scale in France); HADS=Hospital Anxiety and Depression Scale; IDDD=Interview for Deterioration in Daily Living Activities; NPI=Neuropsychiatric Inventory; NR=Not Reported; PMID=PubMed Identification Number; QoL-AD: Quality of Life in Alzheimer's Disease scale; RoB=Risk of Bias; SD=Standard Deviation; vs=versus

Table D-21. Summary of strength of evidence for PLWD outcomes: cognitive rehabilitation

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Individual cognitive rehabilitation vs Usual care Quality of Life	9 months	1 RCT ¹³³ (n=475)	No significant difference between groups	Medium	Unknown	Direct	Imprecise	Insufficient
Individual cognitive rehabilitation vs Usual care Quality of Life	24 months	1 RCT ¹⁰⁷ (n=311)	No significant difference between groups	Medium	Unknown	Direct	Imprecise	Insufficient
Individual cognitive rehabilitation vs usual care Daily function	24 months	1 RCT ¹⁰⁷ (n=311)	No significant difference between groups	Medium	Unknown	Direct	Imprecise	Insufficient
Individual cognitive rehabilitation vs usual care Self-rated goal attainment	9 months	1 RCT ¹³³ (n=475)	Statistically but not clinically significant benefit of individual cognitive rehabilitation compared with usual care on function assessed as self-rated goal attainment	Medium	Unknown	Direct	Imprecise	Insufficient
Errorless Learning (feed-forward instructions) vs Trial & Error Learning Daily function	6 months	1 RCT ¹³⁴ (n=161) 6 months	No difference between groups	Low	Unknown	Direct	Imprecise	Insufficient
Individual cognitive rehabilitation vs usual care Dependency	24 months	1 RCT ¹⁰⁷ (n=311)	Less dependency (slower functional decline) with cognitive rehabilitation over 24 months	Medium	Unknown	Direct	Imprecise	Insufficient
Individual cognitive rehabilitation vs usual care Neuropsychiatric symptoms	24 months	1 RCT ¹⁰⁷ (n=311)	No difference between groups	Medium	Unknown	Direct	Imprecise	Insufficient
Errorless Learning (feed-forward instructions) vs Trial & Error Learning Neuropsychiatric symptoms	6 months	1 RCT ¹³⁴ (n=161)	No difference between groups	Low	Unknown	Direct	Imprecise	Insufficient
Individual cognitive rehabilitation vs usual care Depression	9 months	1 RCT ¹³³ (n=475)	No difference between groups	Medium	Unknown	Direct	Imprecise	Insufficient
Individual cognitive rehabilitation vs usual care Anxiety	9 months	1 RCT ¹³³ (n=475)	No difference between groups	Medium	Unknown	Direct	Imprecise	Insufficient

Abbreviations: PLWD=Persons Living With Dementia; RCT=Randomized Controlled Trial; vs=Versus

Evidence Map: Cognitive Rehabilitation

Table D-22. Characteristics of evidence map studies: cognitive rehabilitation

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics N Age (mean) Sex (% female) Race (% majority) Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:

Mbakile-Mahlanza 2020 ¹³⁹ (31762434) Australia Small sample	Montessori-based activities for PLWD implemented by family members 30 minutes 2x/week 2 weeks	Reading a newspaper with family member 30 minutes 2x/week 2 weeks	Nine Nursing Homes. Cluster crossover RCT; 1-week between groups (6-week study). N=20 PLWD and 20 ICG	Chart diagnosis of dementia, ≥ 3 months in NH, family willing to visit at least 2x/week & attend 3-hr. training. Dementia severity: unable to determine	N=20 Age: NR Sex: NR Other: NR	NR	N=20 Mean age: 64 yrs. Female: 85% Race: NR Education: secondary or higher=68% Relation: 50% offspring	Duration: 5 yrs. Not living with PLWD: 100% (70% IC lived with family) Employed: 48% (45% retired)	NA	NA	2 weeks x 2 (after each group)	Affect, engagement	IC: quality of relationship, mutuality, mood, mastery, Carer-QoL
Chen 2020 ¹³² (No PMID) China High RoB	Montessori-method of sensory stimulation and feeding training (re-identify food and utensils, use utensils). Details: NR 1 month	Observation and timely correction by staff, if needed	Hospital. Retrospective (randomized retrospective data); PLWD with complete records, orally fed, informed consent. Not RCT	Clinical diagnosis of AD or VD. Mild to severe (54% moderate, 40% severe)	N=85 Mean age: 71 yrs. Female: 46% Education: 43% junior high or above	NR	NA	NA	NR	NR	1 month	EdFED, EBS, eating time, MMSE	NR
Yuen 2019 ¹⁴⁰ (no PMID) Hong Kong Small sample	DementiAbility: Montessori-based cognitive rehabilitation 6 sessions, 45 minutes each	Structured social activities	Nursing home RCT N=46 PLWD	Dementia diagnosis	N=46 Age: yes Sex: yes Race: no (Hong Kong) Education: no	NR	NR	NR	NA	NA		Agitation; Global deterioration scale	NA
Silva 2017 ¹⁴¹ (28124633) Portugal Small sample Silva 2017 ¹⁴²	Use of a sense-cam as a cognitive intervention	Memo-taking; Diary	RCT Community-based or integrated in day-care center	PLWD with a diagnosis of probable AD; <6 months since diagnosis; MMSE 15-26; other inclusion criteria applied	N=51 (17 sense-cam, 17 memo-writing, 17 diary) Age: N Sex: N Race % majority: N Education: Y	NR	NA	NA	NR	NR	6 weeks	Depression,; functional capacity; quality of life	NA
Tanaka 2017 ¹⁴³ (27612310) Japan Small sample	5-dimension cognitive rehabilitation: 1. Group (1 hour) 2. Individual (20 min.) Both: 2x/week for 12 weeks	Control	One NH RCT N=60	Resident of one geriatric health facility, MMSE 5-23; dementia AD, VD, mixed, other; and completed this rehabilitation program	N=60 (reported 43: 13 group, 20 individual, 20 control) 21 regular care) Age: Y Sex: Y Race: NR Education: N	SES: NR Disability: NR Household: NR Insurance: NR (Japan)	NR	NR	NR	NR	12 weeks	Cognition, mood, QOL (Japan), GDS, communication	NR
Tsuchiya 2016 ¹⁴⁴ (27303064) Japan Small sample	Brain-Activating Rehabilitation 1 hour 3 times weekly plus standard cognitive rehabilitation	Standard cognitive rehabilitation	Hospital daycare RCT N=48	Diagnosed dementia CDR rating 0.5-3	N=201 Age: Y Sex: Y Race: N Education: N	NR	NA	NA	NA	NA	Post intervention (NR)	MOSES	NA
Seyun Kim 2015 ¹⁴⁵ (25729212) Korea Small sample	Cognitive training in the performance of everyday activities	Unstructured conversation and watch health-related videos	RCT PLWD recruited from a day care center	PLWD diagnosed with possible or probable AD; MMSE ≥ 18	N=43 (22 intervention, 21 control) Age: Y Sex: Y % majority race: N Education: Y	NR	NA	NA	NR	NR	8 weeks	Task performance; satisfaction; quality of life; mental status; memory	NA
Kumar 2014 ¹³⁵ (24982692) India High RoB	Novel occupational therapy program + usual care 5 weeks	Usual care	Outpatient Unblinded RCT	Adults with mild (most) to moderate dementia, MMSE 11-23	n=77 (attrition NR) Age: Y Sex: Y Race: N Education: Y	SES: NR Disability: NR Household: NR Insurance: NR	NA	NA	NA	NA	5 weeks	Standard OT assessment, WHO-QOL-BREF	NR

Thivierge 2014 ¹⁴⁶ (23871120) Canada Small sample Brunelle-Hamann 2015 ¹⁴⁷	Cognitive rehabilitation of instrumental ADL activities	Delayed training	Mix of community and nursing home adults. 6 months RCT crossover	PLWD diagnosed with AD; mean MMSE=22; Additional criteria applied	N=20 (reported 17) Age: Y Sex: Y % majority race: N Education: Y	NR	NA	NA	NR	NR	6 months	ADL/IADL, NPI, memory, QOL	caregiver burden
Wu 2014 ¹³⁶ (24444172) Taiwan High RoB Wu 2013 ¹⁴⁸	Individualized Montessori-based eating activities with spaced retrieval 24 sessions over 8 weeks; number of sessions adjusted per PLWD's recall	1. Standardized Montessori-based eating activities 24 sessions over 8 weeks 2. No treatment	Four Veteran's homes 3-group quasi-experimental N=90	Dementia diagnosis	N=205 (reported 90) Age: Y Sex: Y Race: N Education: Y	NR	NA	NA	NA	NA	1, 3, 6 months	Edinburgh Feeding Evaluation in Dementia (Chinese)	NA
Van der Ploeg 2013 ¹⁴⁹ (23237211) Australia Small sample	One-to-one Montessori-based activities 30 min, 2x/week 2 weeks	Non-personalized activities	Nine residential facilities. RCT crossover after 2 weeks	Adults diagnosed with dementia, agitated behavior & resident for > 3 months	N=44 Age: Y Sex: Y % majority race: N Education: N	Lost English fluency: Y	NA	NA	NR	NR	Two periods of 4 weeks	Agitated behavior; engagement; apathy, affect	NA
Kurz 2012 ¹⁵⁰ (21986341) Germany Pilot	CORDIAL 12 individual weekly sessions. Combined 4 established strategies adopted from neurorehabilitation & psychotherapy	Standard care	Home-based RCT N=201 PLWD	Mild AD MMSE >21	N=201 Age: Y Sex: Y Race: N Education: Y	NR	N=201 Age: Y Sex: Y Race: N Education: N Relation to PLWD: Y	NR	NA	NA	3, 9 months	B-ADL AFIB DEMQOL GDS NPI	ZBI BDI WMSRLM TMT ZUF-8
Lin 2011 ¹⁵¹ (20054841) Taiwan Small sample	Use of a Montessori method to increase eating ability ½ hour, 3x/week 8 weeks	Routine activities	RCT crossover with 2-week washout. Randomized by care unit, 2 dementia care facilities	PLWD diagnosed with dementia; EdFED score of >=2 and MMSE 10-23	N=29 Age Y Sex: Y Race majority: N Education: Y (not by group)	Religion: Y (not by group)	NA	NA	NR	NR	8 week intervention with 2 week washout between	Edinburgh Feeding evaluation, Verbal assistance, Physical assistance, nutrition	NA
Clare 2010 ¹⁵² (20808145) UK Small sample	Individualized cognitive rehabilitation	1. Relaxation therapy 2. No treatment	Community-based RCT	PLWD with diagnosis of AD, early stage, with MMSE>=18 Screened for exclusion criteria	N=69 (23 cognitive rehabilitation, 24 relaxation, 22 no treatment) Age: Y Sex: Y Race % majority: N Education: N	Prior disability: Y	NR	Living with PLWD: Y	NR	NR	8 week intervention period; 6 month follow-up	Quality of life, anxiety, depression, memory functioning, verbal fluency	Stress level, anxiety, physical health, social relationships, environment
Lam 2010 ¹⁵³ (19582757) Hong Kong Pilot	Individualized functional skills training for affective disturbances and functional skills	General occupational therapy program	Community and residential center based RCT	PLWD diagnosed with mild or moderate dementia; Screening criteria applied	N=74 (37 skills training, 37 control) Age: Y Sex: Y Race % majority: N Education: Y	NR	NA	NA	NR	NR	8 week, 3 months, 6 months	MMSE Motor skills Depression Apathy	NA
Lin 2010 ¹⁵⁴ (20054841) Taiwan Small sample	1. Spaced retrieval activities 3x/week, 8 weeks 2. Montessori-based activities 3x/week, 8 weeks	Routine activity	Nursing homes (3) RCT Clusters: 3 n=85	Adults with chart diagnosis of dementia, lived in 1 of 3 NH, scored ≥ 2 on EdFED, MMSE 10-23	N=85 (32 spaced retrieval, 29 Montessori, 24 control) Age: Y Sex: Y Race: N Education: N	SES: NR Disability: NR Household: NA Insurance: NR (Taiwan) More race: NR	NA	NA	NR	NR	8 weeks	Eating ability & feeding assistance required (multiple)	NR

Loewenstein 2004 ¹⁵⁵ (15249277) US Small sample	Cognitive rehabilitation program focused on memory and functional performance	Mental stimulation program, using computer games	Setting NR (assume community due to mean MMSE=24) RCT	PLWD diagnosed with probable or possible AD (DSM-IV), all on stable cholinesterase inhibitor	N=44 (25 cognitive rehabilitation, 19 mental stimulation) Age: Y Sex: Y Race: NR Language: Y Education: Y	NR	NR	NR	NA	NA	12 weeks	Performance on trained skills; memory/recall; ability to perform tasks	NA
Beck 2002 ¹⁵⁶ (12131234) US Small sample	ADL and psychosocial activity combinations (3 groups) 1. Combined psychosocial & ADL intervention 2. ADLs 3. psycho-social	2 control groups: 1. attention control 2. usual care	Nursing home (7 sites) RCT (5 groups total in each center) Randomized 179; reported 127	PLWD diagnosed with dementia MMSE <=20 Screened for exclusion criteria	N=179 (reported 129): 28 ADL, 29 psycho-social activity, 22 combined, 29 attention control, 19 no intervention. Age: Y Sex: Y Race % majority: Y Education: N	% widowed: Y	NA	NA	Education: Y	Training: Y	12-week intervention period One- and two-month follow-up post intervention	Disruptive behavior scores	NA
Quayhagen 1995 ¹⁵⁷ 7761291 USA Small sample	Active cognitive stimulation training (dyadic remediation program)	1. Passive activity 2. Waitlist	Community (intervention in home by ICG) n=95 eligible	AD, living at home with caregiver	N=78 completed (25 intervention, 28 passive activity, 25 control) Age: Y Sex: Y Race: Y Education: Y	NR	N=78 Age: Y Sex: Y Race: N Education: Y	NR	NA	NA	9 months	Memory & Behavior Problems Checklist, multiple cognitive	NR
Tappen 1994 ¹⁵⁸ (8184127) USA Small sample	1. Skill training in performance of basic daily activities	1. Traditional situational approach 2. regular care	RCT Nursing home; random selection from entire NH population	PLWD diagnosed with dementia; exclusion criteria applied	N=63 (21 treatment, 21 situational, 21 regular care) Age: Y Sex: Y % majority race: N Education: N	NR	NA	NA	NR	NR	20 weeks	Self-maintenance; skills performance; goal attainment	NA

Abbreviations: AD=Alzheimer's Disease; ADL=Activities of Daily Living; EBS=Eating Behavior Scale; EdFED=Edinburgh Feeding Evaluation in Dementia; FC=formal caregiver; GDS=Geriatric Depression Scale; IC=informal caregiver; MMSE=Mini Mental State Exam; N=number; N=no; NA=Not applicable; NH=nursing home; NR=Not Reported; PMID=PubMed Identification Number; OT=occupational therapist; RoB=risk of bias; UC=usual care; VD=vascular dementia; Y=yes; **PLWD**=Persons with Dementia; RCT=Randomized controlled trial; IGC=Informal Caregiver; IADL=Instrumental Activities of Daily Living; QoL=Quality of Life; RCT=Randomized controlled trial

Cognitive Training

Table D-23. Risk of bias assessment: cognitive training

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Tseng 2019 ¹⁵⁹ (30970666)	6 months	Medium	High	X	X	X	X	NR	High
Kallio 2018 ¹⁶⁰ (29345724) Finland Medium Explanatory	3 months 9 months	Low 3 months: 0% High 9 months: 20.4%	Low	Low	Low	Medium	Medium	Foundation	Medium (3 months) High (9 months)
Amieva 2016 ¹⁰⁷ (26572551)	3 months 24 months	Medium 3 months: 9.9% High 24 months: 28.1%	Low	Low	Low	Medium	Medium	Government	Medium (3 months) High (24 months)
Cavallo 2016 ¹⁶¹ (27600448)	3 months 6 months	Low 5%	Low	Medium	Low	Medium	Medium	NR	Medium
Giuli 2016 ¹⁶² (26952713)	10 weeks	Low 5.9%	Low	High	Medium	Medium	X	Government	High
Hsu 2016 ¹⁶³ (27878873)	6 weeks	Low 2%	Medium	Low	Medium	Low	Medium	NR	Medium
Kao 2016 ¹⁶⁴ (27307717)	6 weeks 10 weeks 18 weeks 30 weeks	Low 6%	Medium	Low	Medium	Medium	Medium	Government	Medium

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported PMID=PubMed Identification Number

Table D-24. Characteristics of included studies: cognitive training

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Kallio 2018 ¹⁶⁰ (29345724) Finland Medium Explanatory Kallio 2017(NA) ¹⁶⁵	Group or individual paper-and-pencil cognitive training focused on executive function; 2 times/week, 45 minutes/session for 12 weeks	Usual care	Adult day care RCT; single site 147 PLWD	AD or other dementia, very mild to moderate; CDR 0.5 to 2	N=147 83 years 72% Female Race NR 46% Education < 8 years	Household Characteristics: yes	NA	NA	NA	NA	3 months	ADAS-Cog 15-D HRQoL	NA	NA

Amieva 2016 ¹⁰⁷ (26572551) France Medium Explanatory	Group cognitive training program focused on ADL and tasks involving memory, attention, language, or executive function; 90-minute weekly sessions for 3 months and monthly maintenance sessions for 21 months	Usual care	Memory centers or geriatric day care units RCT; multisite 324 PLWD	Mild to moderate AD diagnosis based on NINCDS-ADRDA criteria, MMSE from 16 to 26, and GDS score of 2 to 5	N=324 79 years 58% Female Race NR 52% Secondary education and above	NR	N=324 Age: NR Sex: NR Race: NR Education: NR Relation to PLWD: NR	NR	NA	NA	3 months	ADAS-Cog NPI DAD Apathy Inventory MADRS QOL-AD AGGIR RUD-Lite	ZBI
Cavallo 2016 ¹⁶¹ (27600448) Italy Medium Explanatory	Structured rehabilitative cognitive training software aimed at memory, attention, executive function, and language; three 30 minutes sessions /week for 12 weeks	Using computer to read newspaper or play games; three 30 minutes sessions /week for 12 weeks	Assisted health residence RCT; single site 80 PLWD	Early-stage AD; NINCDS-ADRDA criteria	N=80 76 years 64% Female Race NR 8 years	NR	NA	NA	NA	NA	3 months 6 months	HADS-Depression HADS-Anxiety RMBT	NA
Hsu 2016 ¹⁶³ (27878873) Taiwan Medium Explanatory	Individual spatial retrieval training ; 5 times/week, 40 min/session, for 6 weeks	Usual care	Nursing centers and dementia units of Veterans Homes RCT; multisite 97 PLWD	Dementia with hyperphagia behaviors(all severities, vascular and brain injury excluded) with at least 3 points on the DHBS	N=97 82 years 28% Female Race NR 41% Secondary or higher	NR	NA	NA	NA	NA	6 weeks	DHBS Food Intake	NA
Kao 2016 ¹⁶⁴ (27307717) Taiwan Medium Explanatory	Spatial retrieval training with Montessori activities during recall; 5 times/week, 40 min/session, for 6 weeks	Usual care	Dementia special care units RCT; multisite 100 PLWD	Dementia with hyperphagia behaviors(all severities, vascular and brain injury excluded) with at least 3 points on hyperphagia scale	N=100 82 years 28% Female Race NR 44% High school or higher	NR	NA	NA	N=NR Age NR Sex NR Race NR Education NR	NR	6 weeks 10 weeks 18 weeks 30 weeks	Hyperphagic behavior Pica Behavior Change in Eating Habit Short Meal Frequency	Caregiver Distress to Hyperphagic Behavior

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; N=number; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; HADS=Hospital Anxiety and Depression Scale; NR=Not reported; NINCDS/ADRDA=National Institute of Neurological and Communicative Disorders and Stroke/ Alzheimer's Disease and Related Disorders Association; ADAS-cog=Alzheimer's Disease Assessment Scale, cognitive subsection; NPI=Neuropsychiatric Inventory; DAD=Disability Assessment for Dementia; MADRS=Montgomery-Asberg Depression Rating Scale; QoL-AD: Quality of Life in Alzheimer's Disease scale; RUD=resource utilization; ZBI=Zarit Burden Interview;

Table D-25. PLWD outcomes summary low and medium risk of bias studies: cognitive training

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Kallio 2018 ¹⁶⁰ (29345724) Cognitive Training vs. Usual Care Medium Explanatory	15-D HRQoL Mean Change from Baseline (95% CI) 3 months	NR	-0.04 (-0.058, -0.021)	-0.04 (-0.056, -0.018)	p=0.82

Amieva 2016 ¹⁰⁷ (26572551) Cognitive Training vs. Usual Care Medium Explanatory	NPI Mean (SD) 3 months	NR	25.34 (28.8)	23.29 (28.4)	p=0.22
Amieva 2016 ¹⁰⁷ (26572551) Cognitive Training vs. Usual Care Medium Explanatory	DAD Mean (SD) 3 months	NR	27.54 (9.2)	26.94 (9.6)	p=0.62
Amieva 2016 ¹⁰⁷ (26572551) Cognitive Training vs. Usual Care Medium Explanatory	Apathy Inventory Mean (SD) 3 months	NR	10.26 (11.7)	10.4 (11.8)	p=0.97
Amieva 2016 ¹⁰⁷ (26572551) Cognitive Training vs. Usual Care Medium Explanatory	AGGIR Mean (SD) 3 months	NR	7.0 (7.9)	6.63 (7.5)	p=0.67
Amieva 2016 ¹⁰⁷ (26572551) Cognitive Training vs. Usual Care Medium Explanatory	MADRS Mean (SD) 3 months	NR	10.65 (9.9)	8.82 (9.1)	p=0.06
Amieva 2016 ¹⁰⁷ (26572551) Cognitive Training vs. Usual Care Medium Explanatory	QOL-AD Mean (SD) 3 months	NR	31.99 (8.0)	33.28 (7.7)	p=0.22
Amieva 2016 ¹⁰⁷ (26572551) Cognitive Training vs. Usual Care Medium Explanatory	RUD-Lite Mean (SD) 3 months	NR	2559.3 (3078.3)	2199.8(3002.2)	p=0.50
Cavallo 2016 ¹⁶¹ (27600448) Cognitive Training vs. Attention Control Medium Explanatory	HADS-Anxiety Mean (SD) 3 months, 6 months	No difference between groups at either time point	3 months: 7.65 (2.41) 6 months: NR	3 months: 7.57 (1.33) 6 months: NR	NR
Cavallo 2016 ¹⁶¹ (27600448) Cognitive Training vs. Attention Control Medium Explanatory	HADS-Depression Mean (SD) 3 months, 6 months	No difference between groups at either time point	3 months: 6.42 (2.21) 6 months: NR	3 months: 6.35 (2.21) 6 months: NR	NR
Hsu 2016 ¹⁶³ (27878873) Cognitive Training vs. Usual Care Medium Explanatory	DHBS Mean Change from Baseline (SD) 6 weeks	NR	-0.88 (2.48)	1.04 (1.98)	p<0.001
Hsu 2016 ¹⁶³ (27878873) Cognitive Training vs. Usual Care Medium Explanatory	Average Food Intake (g) Mean Change from Baseline (SD) 6 weeks	NR	-38.63 (145.51)	18.28 (127.34)	p=0.046
Kao 2016 ¹⁶⁴ (27307717) Cognitive Training vs. Usual Care Medium Explanatory	Hyperphagic Behavior Group x Time p-value 6 weeks, 10 weeks, 18 weeks, 30 weeks	NR	NR	NR	All time points p<0.05
Kao 2016 ¹⁶⁴ (27307717) Cognitive Training vs. Usual Care Medium Explanatory	Pica Behavior Group X Time p-value 6 weeks, 10 weeks, 18 weeks, 30 weeks	NR	NR	NR	6 weeks, 10 weeks, 18 weeks: p<0.05 30 weeks: p=0.092
Kao 2016 ¹⁶⁴ (27307717) Cognitive Training vs. Usual Care Medium Explanatory	Change in Eating Habit Group X Time p-value 6 weeks, 10 weeks, 18 weeks, 30 weeks	NR	NR	NR	6 weeks, 10 weeks, 18 weeks: p>0.05 30 weeks: p=0.037
Kao 2016 ¹⁶⁴ (27307717) Cognitive Training vs. Usual Care Medium Explanatory	Short Meal Frequency Group X Time p-value 6 weeks, 10 weeks, 18 weeks, 30 weeks	NR	NR	NR	All time points p<0.05

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number

Table D-26. Caregiver outcomes summary low and medium risk of bias studies: cognitive training

Study (PMID) Comparison Caregiver Type RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Amieva 2016 ¹⁰⁷ (26572551) Cognitive Training vs. Usual Care Medium Explanatory	ZBI Mean (SD) 3 months	NR	30.31 (25.9)	30.05 (25.6)	0.88
Kao 2016 ¹⁶⁴ (27307717) Cognitive Training vs. Usual Care Medium Explanatory	Caregiver Distress to Hyperphagic Behavior Group X Time p-value 6 weeks, 10 weeks, 18 weeks, 30 weeks	NR	NR	NR	6 weeks: p=0.004 10 weeks, 18 weeks, 30 weeks: p>0.05

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; ZBI=Zarit Burden Interview

Table D-27. Summary of strength of evidence for PLWD outcomes: cognitive training

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Cognitive Training vs. Usual Care Quality of Life	3 months	2 RCTs (n=471)	No benefit.	Moderate	Consistent	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care or Attention Control Depression	3 to 6 months	2 RCTs (n=404)	No benefit.	Moderate	Consistent	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Hyperphagia Behavior	6 to 30 weeks	2 RCTs (n=197)	Benefit favoring intervention.	Moderate	Consistent	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Food Intake	6 weeks	1 RCT (n=97)	Benefit favoring intervention.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Short meal frequency	6 to 30 weeks	1 RCT (n=100)	Benefit favoring intervention at all time-points.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Pica behavior	6 to 30 weeks	1 RCT (n=100)	Benefit favoring intervention post-intervention, but not sustained 6-months post-intervention.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Change in eating habits	6 to 30 weeks	1 RCT (n=100)	No benefit.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care BPSD	3 months	1 RCT (n=324)	No benefit.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Function/ADLs	3 months	1 RCT (n=324)	No benefit.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Apathy	3 months	1 RCT (n=324)	No benefit.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Dependence	3 months	1 RCT (n=324)	No benefit.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Attention Control Anxiety	3 and 6 months	1 RCT (n=80)	No benefit.	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: RCT=Randomized controlled trial; ADL=Activities of daily living; BPSD=Behavioral and psychological symptoms of dementia

Table D-28. Summary of strength of evidence for caregiver outcomes: cognitive training

Comparison Caregiver Type Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Cognitive Training vs. Usual Care Caregiver Distress Informal caregivers	3 months	1 RCT (n=324)	No benefit.	Moderate	Unknown	Direct	Imprecise	Insufficient
Cognitive Training vs. Usual Care Caregiver Distress, PLWD hyperphagic behavior Formal caregivers	6 to 30 weeks	1 RCT (n=NR)	No benefit.	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: n=number; PLWD=Persons with Dementia; RCT=Randomized controlled trial

Evidence Map: Cognitive Training

Table D-29. Characteristics of evidence map studies: cognitive training

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Tseng 2019 ¹⁵⁹ Taiwan High RoB	Smart Restored Learning exercise	No intervention	Veterans home Quasi-experimental n=68	PLWD diagnosed with dementia, MMSE <25 and living in Veterans Home	N=68 (35 treatment, 33 control) Age: Y Sex: N Race % majority: N Education: Y	NR	NA	NA	NR	NR	6 months	MMSE Neuropsychiatric Inventory Frontal Assessment Battery	NA
Giovagnoli 2017 ¹⁶⁶ Italy Sample Size	Cognitive training	1. Music therapy or 2. neuro-education	One care center RCT single blind	PLWD with mild-moderate dementia or probable AD. Screened for additional criteria	N=50 (17 cognitive training, 17 music, 13 neuro-education) Age: Y Sex: Y Race: N Education: Y	NR	NR	NR	NR	NR	3 months	Word fluency test Neuropsychological and behavioral assessments	NA

Venturelli 2016 ⁵² (27540967) Italy Small sample	Cognitive training based on reality orientation method; 5 days/week for 1 hour before sunset for 3 months	Walking at moderate intensity; 5 days/week for 1 hour before sunset for 3 months	AD care units RCT 40 PLWD	AD with MMSE between 10 and 15 and neurobehavioral symptoms of SDS	N=40 Age: yes Sex: yes Race: no Education: no	NR	NA	NA	NA	NA	3 months	NPI ABS	NA
Chen, 2016 ¹⁶⁷ (27846102) Taiwan Small sample	Individualized learning therapy 30 minutes twice weekly for 3 months	Usual care	Nursing home Quasi-experimental, 2 homes	Dementia diagnosis, MMSE between 10 and 23	N=44 Age: yes Sex: yes Race: no Education: no	Marital status	NA	NA	NA	NA	12 weeks	Chinese NPI MMSE	NA
De Luca 2016 ¹⁶⁸ Italy Small sample	Combined standard cognitive training and web-based rehabilitation 24 sessions x 8 weeks	Standard treatment	Nursing home RCT	Diagnosis of mild to moderate cognitive decline MMSE 25 +/- 3.4); ability to perform training	N=20 (10 experimental 10 control) Age: Y Sex: Y Race: Y Education: Y	NR	NA	NA	NR	NR	8 weeks	MMSE Attention Verbal fluency ADL, ADL Behavioral, Depression	NA
Giuli 2016 ¹⁶² (26952713) Italy High RoB	Comprehensive cognitive training	No intervention	Community based RCT	Three levels of cognitive impairment: healthy elderly, moderate cognitive impairment, mild to moderate AD; Aged 65 or older	N=321 (47 healthy experimental, 53 healthy control, 48 moderate decline experimental, 49 moderate decline control, 48 AD experimental, 47 AD control) Age: Y Sex: Y Race % majority: N Education: Y	NR	NA	NA	NR	NR	10 weeks	MMSE, Orientation, Memory, Attention, Verbal fluency, Depression, ADL, IADL, confidence in own cognition	NA
Kawashima 2015 ¹⁶⁹ US Sample Size	Americanized SAIDO (working memory training program of basic arithmetic & language)	Usual care	Two nursing homes. Prospective non-randomized, by NH n=47 (39 reported)	PLWD diagnosed with dementia (DSM-IV)	N=n=47 (23 intervention, 24 controls) Age: Y Sex: Y Race % majority: N Education: N	NR	NA	NA	NR	NR	6 months	Cognitive scores, mood mental status	NA
Bergamaschi 2013 ¹⁷⁰ (23784727) Italy Sample Size	Repeated cycles of cognitive training x 1 year	Non-specific cognitive exercises	Day center RCT	PLWD diagnosed with mild to moderate AD; MMSE 18-24	N=32 (16 each group) Age: Y Sex: N Race % majority: N Education: Y	NR	NA	NA	NR	NR	1 year	MMSE Memory and recall Verbal fluency Depression ADL	NA
Mapelli 2013 ¹⁷¹ (24052800) Italy Sample Size	Cognitive stimulation focused on memory, language, spatial and temporal orientation, attention and logic	Occupational therapy s placebo or no intervention	Nursing home RCT PLWD with AD, vascular and mixed dementia	PLWD diagnosed with AD, vascular and mixed dementia AD MMSE 19-24; vascular MMSE 14-18	N=30 (10 each group) Age: Y Sex: N Race % majority: N Education: Y	NR	NA	NA	NR	NR	8 weeks	Cognition Behavior	NA

Tarraga 2006 ¹⁷² (16820420) Spain Sample Size	Cognitive stimulation using an interactive multi-media tool plus psycho-stimulation and medication	Integrated psycho-stimulation and medication, or medication only	Community-based RCT PLWD diagnosed as having suspected AD	PLWD diagnosed with suspected AD >65 years old; at least 3 years education. MMMSE 18-24; GDS 3-4.	N=43 (15 experimental, 16 treatment control, 12 medication only control) Age: Y Sex: Y Race % majority: N Education: statistic only	NR	NA	NA	NR	NR	24 weeks	ADAS-Cog Additional tests of cognition, verbal fluency, recall, level of disability	NA
McGilton 2003 ¹⁷³ (12959805) Canada Sample Size	Training in way-finding in a new environment	No intervention	University geriatric care center RCT	PLWD with diagnosis of AD and moderate to severe cognitive decline. Able to ambulate and understand English	N=32 (17 treatment, 15 control) Age: Y Sex: Y Race % information: N Education: N	NR	NA	NA	NR	Training: Y	4 weeks training Re-evaluation at 1 week and 3 months post-test	Find dining room Agitation Spatial orientation	NA
Davis 2001 ¹⁷⁴ (11236819) US Sample Size	Training in face-name association, spaced retrieval and cognitive stimulation with ICGs	Mock placebo intervention	AD research center RCT Crossover study	Diagnosis of probable AD	N=37 (19 intervention, 18 control) Age: Y Sex: Y Race % majority: N Education: Y	NR	NR	NR	NR	NR	10 weeks; Crossover at 5 weeks	MMSE Verbal and visual memory Attention/concentration Word generation Motor speed Depression Quality of life	NR
Zarit 1982 ¹⁷⁵ (13238424) US Sample Size	1. Group didactic training (with imagery) 2. problem solving All groups with ICG	Wait list control	Community-based RCT n=35 PLWD	Community dwelling adults screened for cognitive impairment (Introduction states adults with	N=35 (14 didactic, 11 problem solving, 10 wait list) Age: Y Sex: N Race % majority: N Education: Y	NR	Relation to PLWD: Y	NR	NA	NA	3.5 weeks	Recall trials Recognition trials	Burden Depression Memory problems Behavior problems Distress over problems

Abbreviations: char=characteristics; EM=Evidence Map; FC=formal caregiver; IC=informal caregiver; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RS=Reporting Status; SES=socioeconomic status; MMSE=Mini-mental State Examination; ADAS-cog=Alzheimer's Disease Assessment Scale; AD=Alzheimer's Disease; RCT=Randomized controlled trial; ADL=Activities of Daily Living; DSM-IV=Diagnostic and Statistical Manual for Mental Disorders, Fourth Edition; IADL=Instrumental; Activities of Daily Living; NPI=Neuropsychiatric Inventory

Cognitive Stimulation Therapy

Table D-30. Risk of bias assessment: cognitive stimulation therapy

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Lok, 2020 ¹⁷⁶	7 weeks	High 23.08%	Low	X	X	X	X	X	High
Lin, 2018 ¹⁷⁷ (28881430)	12 weeks 24 weeks	Medium 14.91%	Medium	High	High	Low	Low	Taiwanese government	High
Orgeta, 2015 ¹⁷⁸ (26292178) Orrell, 2017 ¹⁷⁹ (28350796)	13 weeks 26 weeks	Medium 23.31%	Low	Low	Moderate	Low	High	UK government	High
Orrell, 2014 ¹⁸⁰ (24676963)	12 weeks 24 weeks	Low 15.68%	Low	Low	High	Low	Moderate	UK government	Medium
Spector, 2003 ¹⁸¹ (12948999) Knapp, 2006 ¹⁸² (16738349)	8 weeks	Low 5.47%	Low	Low	Moderate	Low	Low	UK government	Low

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table D-31. Characteristics of included studies: cognitive stimulation therapy

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Orrell, 2014 ¹⁸⁰ (24676963) United Kingdom Medium Explanatory	Maintenance of CST after an initial 7-week program of CST. Program defined as group sessions focused on reality orientation and cognitive stimulation, led by trained facilitator; one 45-minute session for 24 weeks	Usual care after an initial 7-week program of CST	Care homes and community centers RCT Multisite 236 PLWD	DSM-IV criteria for dementia, mild to moderate dementia on Clinical Dementia Rating scale	N=236 Age 83 years 63.6% female Race NR Education NR	NR	NR	NR	NR	NR	12 weeks 24 weeks	Quality of life Neuropsychiatric symptoms Function	NR	
Spector, 2003 ¹⁸¹ (12948999) Knapp, 2006 (16738349) United Kingdom Low Explanatory	Group sessions focused on reality orientation and cognitive stimulation, facilitated by researcher; two 45-minute sessions for 7 weeks	Usual care, typically doing nothing, but sometimes including music, games, arts and crafts, and other activities	Residential homes and day care centers RCT Multisite 201 PLWD	DSM-IV criteria for dementia, MMSE between 10 and 24	N=201 Age 85.3 years 78.6% female Race NR Education NR	NR	NR	NR	NR	NR	8 weeks	Quality of life Function Depression Anxiety	NR	

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; N=number; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; CST=Cognitive Stimulation Therapy; NR=Not Reported; DSM-IV=Diagnostic and Statistical Manual for Mental Disorders, Fourth Edition; RCT=Randomized controlled trial

Table D-32. PLWD outcomes summary low and medium risk of bias studies: cognitive stimulation therapy

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Orrell, 2014 ¹⁸⁰ (24676963) Explanatory	Quality of life, self-rated QoL-AD, DEMQOL 12 weeks	0.32 points (95% CI -0.88 to 1.52, QoL-AD) -0.86 points (95% CI -3.45 to 1.73, DEMQOL)	Cognitive stimulation maintenance therapy after initial cognitive stimulation	Usual care after withdrawal of cognitive stimulation	P=0.54 (QoL-AD) P=0.54 (DEMQOL)
Orrell, 2014 ¹⁸⁰ (24676963) Explanatory	Quality of life, caregiver-rated QoL-AD, DEMQOL 12 weeks	1.53 points (95% CI 0.37 to 2.69, QoL-AD) 3.24 points (95% CI 0.29 to 6.19, DEMQOL)	Cognitive stimulation maintenance therapy after initial cognitive stimulation	Usual care after withdrawal of cognitive stimulation	P=0.01 (QoL-AD) P=0.03 (DEMQOL)

Orrell, 2014 ¹⁸⁰ (24676963) Explanatory	Neuropsychiatric symptoms NPI 12 weeks	1.47 points (95% CI -1.59 to 4.53)	Cognitive stimulation maintenance therapy after initial cognitive stimulation	Usual care after withdrawal of cognitive stimulation	P=0.34
Orrell, 2014 ¹⁸⁰ (24676963) Explanatory	Function ADCS-ADL 12 weeks	2.64 points (95% CI 0.08 to 5.20)	Cognitive stimulation maintenance therapy after initial cognitive stimulation	Usual care after withdrawal of cognitive stimulation	P=0.04
Orrell, 2014 ¹⁸⁰ (24676963) Explanatory	Quality of life, self-rated QoL-AD, DEMQOL 24 weeks	1.78 points (95% CI -0.01 to 3.57, QoL-AD) 0.30 points (95% CI -3.45 to 1.73, DEMQOL)	Cognitive stimulation maintenance therapy after initial cognitive stimulation	Usual care after withdrawal of cognitive stimulation	P=0.03 (QoL-AD) P=0.87 (DEMQOL)
Orrell, 2014 ¹⁸⁰ (24676963) Explanatory	Quality of life, caregiver-rated QoL-AD, DEMQOL 24 weeks	0.07 points (95% CI -1.39 to 1.53, QoL-AD) 1.13 points (95% CI -2.24 to 4.51, DEMQOL)	Cognitive stimulation maintenance therapy after initial cognitive stimulation	Usual care after withdrawal of cognitive stimulation	P=0.95 (QoL-AD) P=0.50 (DEMQOL)
Orrell, 2014 ¹⁸⁰ (24676963) Explanatory	Neuropsychiatric symptoms NPI 24 weeks	1.58 points (95% CI -2.67 to 5.84)	Cognitive stimulation maintenance therapy after initial cognitive stimulation	Usual care after withdrawal of cognitive stimulation	P=0.53
Orrell, 2014 ¹⁸⁰ (24676963) Explanatory	Function ADCS-ADL 24 weeks	0.94 points (95% CI -2.04 to 3.92)	Cognitive stimulation maintenance therapy after initial cognitive stimulation	Usual care after withdrawal of cognitive stimulation	P=0.54
Spector, 2003 ¹⁸¹ (12948999) Knapp, 2006 ¹⁸² (16738349) Explanatory	Quality of life QoL-AD 8 weeks	1.64 points (95% CI 0.09 to 3.18)	Cognitive stimulation	Usual care	P=0.028
Spector, 2003 ¹⁸¹ (12948999) Knapp, 2006 ¹⁸² (16738349) Explanatory	Function HCS 8 weeks	2.3 points (95% CI -0.45 to 4.15)	Cognitive stimulation	Usual care	P=0.09
Spector, 2003 ¹⁸¹ (12948999) Knapp, 2006 ¹⁸² (16738349) Explanatory	Neuropsychiatric symptoms CAPE-BRS 8 weeks	0.40 points (95% CI -0.9 to 1.69 points)	Cognitive stimulation	Usual care	P=0.449
Spector, 2003 ¹⁸¹ (12948999) Knapp, 2006 ¹⁸² (16738349) Explanatory	Anxiety RAID 8 weeks	-1.30 points (95% CI -3.48 to 0.87)	Cognitive stimulation	Usual care	P=0.200
Spector, 2003 ¹⁸¹ (12948999) Knapp, 2006 ¹⁸² (16738349) Explanatory	Depression CSDD 8 weeks	0.12 points (95% CI 1.56 to 1.31)	Cognitive stimulation	Usual care	P=0.648

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; QoL-AD: Quality of Life in Alzheimer's Disease scale; HCS: Holden Communication Scale; CAPE-BRS: Clifton Assessment Procedures for the Elderly - Behaviour Rating Scale; RAID: Rating Anxiety in Dementia scale; CSDD: Cornell Scale for Depression in Dementia; DEMQOL: Dementia Quality of Life scale; NPI: Neuropsychiatric Inventory; ADCS-ADL: Alzheimer's Disease Cooperative Study – Activities of Daily Living scale

Table D-33 Summary of strength of evidence for PLWD outcomes: cognitive stimulation therapy

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
CS vs. Active control Quality of life	8 weeks	1 RCT (n=201)	Improvement in quality of life	Low	Unknown	Direct	Imprecise	Insufficient
CS vs. Active control Function	8 weeks	1 RCT (n=201)	No change in function	Low	Unknown	Direct	Imprecise	Insufficient
CS vs. Active control Neuropsychiatric symptoms	8 weeks	1 RCT (n=201)	No change in neuropsychiatric symptoms	Low	Unknown	Direct	Imprecise	Insufficient
CS vs. Active control Anxiety	8 weeks	1 RCT (n=201)	No change in anxiety	Low	Unknown	Direct	Imprecise	Insufficient
CS vs. Active control Depression	8 weeks	1 RCT (n=201)	No change in depression	Low	Unknown	Direct	Imprecise	Insufficient
CS maintenance therapy vs. Usual care after initial CS therapy Quality of life	24 weeks	1 RCT (n=236)	Improvement in quality of life	Moderate	Unknown	Direct	Imprecise	Insufficient
CS maintenance therapy vs. Usual care after initial CS therapy Neuropsychiatric symptoms	24 weeks	1 RCT (n=236)	Improvement in quality of life	Moderate	Unknown	Direct	Imprecise	Insufficient
CS maintenance therapy vs. Usual care after initial CS therapy Function	24 weeks	1 RCT (n=236)	Improvement in quality of life	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: n=number; RCT=Randomized controlled trial; CS=Cognitive Stimulation

Evidence Map: Cognitive Stimulation Therapy

Table D-34. Characteristics of evidence map studies: cognitive stimulation therapy

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Lok, 2020 ¹⁷⁶ Turkey High RoB	Cognitive stimulation therapy	Usual care	Hospital; single site 78 PLWD	MMSE between 13 and 24	N=60 Age NR Sex Race NR Education	Household characteristics	NR	NR	NR	NR	7 weeks	MMSE CAPS QoL-AD	NR
Cheung, 2019 ¹⁸³ Hong Kong Small sample	Cognitive stimulating play intervention	Social activities	Community residents 30 PLWD	Dementia type NR	N=30 Age: yes Sex: yes Race: NR Education: yes	NR	NR	NR	NR	NR	8 weeks	MoCA FOME	

Lin, 2018 ¹⁷⁷ (28881430) Taiwan High RoB	Cognitive stimulation therapy	Reminiscence therapy; usual care	Long-term care institutions; multisite 105 PLWD	MMSE between 17 and 24, with agitation or depressive symptoms present	N=105 Age: no Sex: yes Race: NR Education: NR	Marital status	NR	NR	NR	NR	12 weeks 24 weeks	MMSE QoL-AD	NR
Capotosto, 2017 ¹⁸⁴ (27272538) Italy Small sample	Cognitive stimulation therapy	General activities	Residential homes; multisite 39 PLWD	Mild to moderate dementia	N=39 Mean age Mean education	NR	NR	NR	NR	NR	7 weeks	MMSE ADAS-Cog QoL-AD CSDD DAD NPI	NR
Wong, 2016 ¹⁸⁵ (29717527) Hong Kong Pilot	Cognitive stimulation therapy	No control group	Community and residential care units of major NGOs; multisite 30 PLWD	Clinical diagnosis of dementia, criteria unspecified	N=30 Mean age Race NR Sex Education	Living situation	Sex	NR	NR	NR	7 weeks	QoL-AD ADAS-Cog	NR
Orgeta, 2015 ¹⁷⁸ (26292178) Orrell, 2017 ¹⁷⁹ (28350796) United Kingdom High RoB	Carer-delivered individual cognitive stimulation therapy	Usual care	Unspecified community settings; multisite 356 PLWD	Dementia unspecified	N=356 Age NR Race Sex Education	Marital status Living situation	356 informal caregivers Sex Race	Marital status Living situation	NR	NR	26 weeks	ADAS-Cog QoL-AD DEMQOL NPI BADLs GDS-15 QCPR	QCPR EQ-5D HADS
Cove, 2014 ¹⁸⁶ (25525349) United Kingdom Small sample	Cognitive stimulation therapy plus carer training; cognitive stimulation therapy alone	Waitlist control	Unspecified community settings; multisite 72 PLWD	DSM-IV diagnosis of mild to moderate dementia	N=72 Mean age Race/ethnicity Sex Education NR	Detailed race/ethnicity	NR	NR	NR		15 weeks	MMSE ADAS-Cog QoL-AD QCPR	QCPR
Yamanaka, 2013 ¹⁸⁷ (23550665) Japan Pilot	Cognitive stimulation therapy	Usual care	Japanese residential and nursing homes; multisite 56 PLWD	Mild to moderate AD	N=56 Mean age Sex Education NR Race NR	NR	NR	NR	NR	NR	7 weeks	MMSE COGNISTAT QoL-AD EQ-5D Face scale for mood	NR
Niu, 2010 ¹⁸⁸ (20713437) China Small sample	Cognitive stimulation therapy	Communication exercise	Military sanatorium; single site 32 PLWD	Mild to moderate AD with marked neuropsychiatric symptoms	N=32 Mean age Sex Education	NR	NR	NR	NR	NR	10 weeks	MMSE NPI	NR
Matsuda, 2007 ¹⁸⁹ (17005066) Japan Small sample	Cognitive stimulation therapy plus donepezil	Donepezil alone	Outpatient neuropsychiatric clinic; single site 30 PLWD	DSM-IV diagnosis of dementia	N=30 NR	NR	NR	NR	NR	NR	1 year	MMSE	NR

Abbreviations: char=characteristics; EM=Evidence Map; FC=formal caregiver; IC=informal caregiver; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RS=Reporting Status; SES=socioeconomic status; MMSE=Mini Mental State Examination; CAPS=Coping and Adaptation Processing Scale; MoCA=Montreal Cognitive Assessment; FOME=Full Object Memory Evaluation; NPI=Neuropsychiatric Inventory; COGNISTAT=Neurobehavioral Cognitive Status Examination; QoL-AD: Quality of Life in Alzheimer's Disease scale; QCPR=Quality of Caregiver and Patient Relationship; DEMQOL=Dementia Quality of Life scale; BADLS=Bristol Activities of Daily Living Scale; GDS-15=Geriatric Depression Scale, 15 item; EQ-5D=EuroQoL 5D scale; HADS=Hospital Anxiety and Depression Scale; ADAS-Cog=Alzheimer's Disease Assessment Scale - Cognition; CSDD=Cornell Scale for Depression in Dementia; DAD=Disability Assessment for Dementia

Recreation Therapy

Table D-35. Risk of bias assessment: recreation therapy

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Pedrinolla, 2019 ¹⁹⁰ (31424399)	24 weeks	Low 6.75%	Low	Low	Moderate	Low	High	Foundations	Medium
Tse, 2018 ¹⁹¹ (30255637)	8 weeks 12 weeks	High NR	X	X	X	X	X	Unfunded	High
Lin, 2015 ¹⁹² (607203463)	24 weeks	High NR	X	X	X	X	X	NR	High
Van Haitsma, 2015 ¹⁹³ (24304555)	0 weeks	Low 7.69%	Moderate	Moderate	High	Low	Moderate	US state government, foundation	High
Low, 2013 ¹⁹⁴ (23315520)	13 weeks 26 weeks	Medium 13.82%	Low	Low	Moderate	Low	Low	Australian government	Medium
Kolanowski, 2011 ¹⁹⁵ (21649633)	Up to 3 weeks	Low 4.69%	Low	Medium	Low	Low	Low	National Institutes of Health	Low
Buettner, 2002 ¹⁹⁶ (11954670)	2 weeks 10 weeks	High NR	X	X	X	X	X	US state government	High
Hopman-Rock, 1999 ¹⁹⁷ (29403282)	24 weeks	High 31.3%	X	X	X	X	X	Foundations, health insurer	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table D-36. Characteristics of included studies: recreation therapy

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Pedrinolla, 2019 ¹⁹⁰ (31424399) Italy	Unstructured group interaction with enclosed natural environment including plants and flowers	Usual care in AD unit of a nursing home, including long corridor for walking plus recreation rooms	Nursing home Single site 163 PLWD	N=163 Probable AD diagnosis MMSE ≤ 15	N=163 Age 77 years 74.2% female Race NR Education NR	NR	NR	NR	NR	NR	24 weeks	Neuropsychiatric symptoms Function Antipsychotic dosage	NR	
Low, 2013 ¹⁹⁴ (23315520) Australia Medium Explanatory	Individual, tailored humor therapy sessions for residents focused on humorous interactions; delivered by a nominated staff member who received one day of training; weekly therapy sessions for 9 to 12 weeks	Usual care, not otherwise specified	Nursing homes Cluster RCT Multisite 36 nursing homes 389 PLWD	N=389 Dementia type NR Mean Global Deterioration Scale 5.0	N=389 Age 84.5 years 64.1% female Race NR Education NR	NR	NR	NR	NR	NR	13 weeks 26 weeks	Depression Neuropsychiatric symptoms Social engagement Quality of life	NR	
Kolanowski, 2011 ¹⁹⁵ (21649633) United States Low Explanatory	Individual recreational activities based on Need-driven Dementia-compromised Behavior model, tailored to residents based on either physical/cognitive function, personality as assessed through the Five Factor Model, or both;	Active control: same individual recreational activities targeted opposite to both physical/cognitive function and personality	Nursing homes Multisite 9 nursing homes 128 PLWD	N = 128 Dementia by DSM-IV criteria Mean MMSE 12.68 to 15.78	N = 128 Mean age 85.87 to 87.21 years Percent White 81.25% to 93.55% Mean years of education 11.64 to 13.87	NR	NR	NR	NR	NR	Up to 7 weeks	Agitation Affect Social engagement	NR	

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; N=number; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; NR=NR; RCT=Randomized controlled trial

Table D-37. PLWD outcomes summary low and medium risk of bias studies: recreation therapy

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Pedrinolla, 2019 ¹⁹⁰ (31424399) Explanatory	Neuropsychiatric symptoms NPI 24 weeks	-31.8 points (95% CI -35.1 to -28.5)	Recreational therapy	Usual care	P < 0.001
Pedrinolla, 2019 ¹⁹⁰ (31424399) Explanatory	Antipsychotic use Quetiapine, mg/day 24 weeks	-150 mg/day (95% CI -175 to -120)	Recreational therapy	Usual care	P < 0.001
Pedrinolla, 2019 ¹⁹⁰ (31424399) Explanatory	Function Barthel Index 24 weeks	1.3 points (95% CI -1.9 to 4.4)	Recreational therapy	Usual care	Not significant; exact value NR
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Neuropsychiatric symptoms NPI 13 weeks	0.05 points (95% CI -0.11 to 0.22)	Recreational therapy	Usual care	P=0.52
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Depression CSDD 13 weeks	0.006 points (95% CI -0.19 to 0.20)	Recreational therapy	Usual care	P=0.95
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Agitation CMAI 13 weeks	-0.04 points (95% CI -0.18 to 0.11)	Recreational therapy	Usual care	P=0.61
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Function MOSES, withdrawal subscale 13 weeks	-0.046 points (95% CI -0.21 to 0.12)	Recreational therapy	Usual care	P=0.58
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Quality of life DEMqoL 13 weeks	Self-rated: -0.10 points (95% CI -0.31 to 0.11) Proxy-rated: 0.07 points (95% CI -0.16 to 0.31)	Recreational therapy	Usual care	Self-rated: p=0.34 Proxy-rated: p=0.53
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Neuropsychiatric symptoms NPI 26 weeks	-0.15 points (95% CI -0.34 to 0.04)	Recreational therapy	Usual care	P=0.13
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Depression CSDD 26 weeks	0.046 points (95% CI -0.18 to 0.27)	Recreational therapy	Usual care	P=0.69
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Agitation CMAI 26 weeks	0.17 points (95% CI 0.004 to 0.34)	Recreational therapy	Usual care	P=0.045
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Function MOSES, withdrawal subscale 26 weeks	0.049 points (95% CI -0.13 to 0.22)	Recreational therapy	Usual care	P=0.59
Low, 2013 ¹⁹⁴ (23315520) Explanatory	Quality of life DEMqoL 26 weeks	Self-rated: 0.05 points (95% CI -0.18 to 0.28) Proxy-rated: -0.07 points (95% CI -0.28 to 0.13)	Recreational therapy	Usual care	Self-rated: p=0.67 Proxy-rated: p=0.48
Kolanowski, 2011 ¹⁹⁵ Explanatory	Agitation CMAI Up to 3 weeks	No difference between arms	Recreational therapy	Active control	p=0.607 personality tailoring alone p=0.339 function tailoring alone p=0.923 personality plus function tailoring

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: CMAI=Cohen-Mansfield Agitation Inventory; CSDD: Cornell Scale for Depression in Dementia; DEMqoL=Dementia Quality of Life Scale; MOSES=Multidimensional Observation Scale for Elderly Subjects; NPI= Neuropsychiatric Inventory; ARS = Affect Rating Scale; DMPT = Dementia Mood Picture Test; PDS = Passivity in Dementia Scale; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number

Table D-38. Summary of strength of evidence for PLWD outcomes: recreation therapy

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
RT vs. Usual care Neuropsychological symptoms	24-26 weeks	2 RCTs (n=552)	Reduction in neuropsychological symptoms	Medium	Inconsistent	Direct	Imprecise	Insufficient
RT vs. Usual care Depression	26 weeks	1 RCT (n=389)	Increase in depression symptoms	Medium	Unknown (single study)	Direct	Imprecise	Insufficient
RT vs. Usual care Agitation	Up to 13 weeks	2 RCT (n=517)	No change in agitation symptoms	Medium	Inconsistent	Direct	Imprecise	Insufficient
RT vs. Usual care Function, social	26 weeks	1 RCT (n=389)	No change in social function	Medium	Unknown (single study)	Direct	Imprecise	Insufficient
RT vs. Usual care Function, physical	24 weeks	1 RCT (n=163)	Improvement in physical function	Medium	Unknown (single study)	Direct	Imprecise	Insufficient
RT vs. Usual care Quality of life, self-rated	26 weeks	1 RCT (n=389)	Improvement in self-rated quality of life	Medium	Unknown (single study)	Direct	Imprecise	Insufficient
RT vs. Usual care Quality of life, proxy-rated	26 weeks	1 RCT (n=389)	Decline in proxy-rated quality of life	Medium	Unknown (single study)	Indirect	Imprecise	Insufficient
RT vs. Usual care Antipsychotic use	24 weeks	1 RCT (n=163)	Decrease in antipsychotic dosage	Medium	Unknown (single study)	Direct	Imprecise	Insufficient

Abbreviations: n=number; RT=Recreational Therapy; RCT=Randomized controlled trial

Evidence Map: Recreation Therapy

Table D-39. Characteristics of evidence map studies: recreation therapy

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Laksono, 2019 ¹⁹⁸ Indonesia Small sample	Chess game	NR	Community residents Single site 20 PLWD	NR	NR	NR	NR	NR	NR	NR	2 weeks	MoCA	NR
Lathifah, 2019 ¹⁹⁹ India Small sample	Dhakonan game	NR	Community residents Single site 20 PLWD	NR	NR	NR	NR	NR	NR	NR	0 weeks	GDS	NR

Tse, 2018 ¹⁹¹ (30255637) Hong Kong High RoB	Play activity program	Reading sessions, unstructured group	Nursing homes Multisite 4 clusters 53 PLWD	N=53 MMSE between 10 and 25	N=53 Age Percent female Education Race NR	NR	NR	NR	NR	NR	8 weeks 12 weeks	Visual analog scale for happiness GDS Katz ADL scale Index of Social Engagement MDS Behavior Rating Scale Non-Pharmacological Therapy Experience Scale	NR
Li, 2017 ²⁰⁰ (27345629) China Small sample	Folk recreation intervention involving art, music, and games	Usual care	Long-term care facility Single site 48 PLWD	N=48 Physician diagnosis of dementia, CDR > 0.5	N =48 Age Percent female Education Race NR	NR	NR	NR	NR	NR	16 weeks	Barthel Index NPI	NR1
Pongan, 2017 ⁷¹ (31288544) France Pilot	Directed singing led by a choir conductor	Painting sessions	Memory clinics Multisite 59 PLWD	N=59 Mild AD, DSM-V criteria	N=59 Age Percent female Education Race	NR	NR	NR	NR	NR	16 weeks	STAI GDS EQ-5D RSES	NR
Schall, 2017 ²⁰¹ (28914089) Germany Small sample	Directed visits to art museums plus art-making sessions	Unstructured visits to art museums	Community 44 PLWD	N=44 Diagnosis of mild to moderate dementia, criteria NR	N=44 Age Percent female Education NR Race NR	NR	N=44 NR	NR	NR	NR	12 weeks	GDS QoL-AD NPI FAHW Smiley Scale	NR
Lin, 2015 ¹⁹² (607203463) China High RoB	Strategy board game sessions, 1-2 hours daily	Not stated	Hospital Single site 147 PLWD	N=147 NINCDS-ADRDA diagnosis of dementia	N=147 NR	NR	NR	NR	NR	NR	24 weeks	MADRS HADS RAND-36	NR
Van Haitsma, 2015 ¹⁹³ (24304555) US High RoB	Activity sessions tailored to individual preferences	Standard activity sessions, involving magazine reading and conversation	Nursing home Single site 180 PLWD	N=180	N=180 Age Percent female Education Race NR	NR	NR	NR	NR	NR	3 weeks	Direct observation of resident behavior MOSES MDS ADL scale	NR
Hattori, 2011 ²⁰² (21518170) Japan Small sample	Art therapy in groups	Basic mathematical exercises	Outpatient clinic Single site 39 PLWD	N=39 NINCDS-ADRDA diagnosis of dementia MMSE ≥ 20	N=39 Age Percent female Race NR Education NR	NR	NR	NR	NR	NR	12 weeks	GDS Apathy Scale SF-8 DBD	NR
Buettner, 2002 ¹⁹⁶ (11954670) US High RoB	15-minute wheelchair tandem bicycle ride with a staff member pedaling the bicycle	Not stated	Nursing home and assisted living facility Multisite 70 PLWD	N=70 MMSE ≤ 24	N=70 NR	NR	NR	NR	NR	NR	2 weeks 12 weeks	GDS CMAI	NR

Hopman-Rock, 1999 ¹⁹⁷ Netherlands (29403282) High RoB	Biweekly sporting activity and game sessions, preferably in a separate room, targeting communication, reactivation, resocialization, and function	Usual activities, including light household activities, bible reading, singing	Nursing homes Multisite 12 clusters 134 PLWD	N=134 NR	N=134 Age Percent female Race NR Education NR	NR	NR	NR	NR	NR	24 weeks	BIP SIPO	NR
--	---	--	--	-------------	---	----	----	----	----	----	----------	-------------	----

Abbreviations: char=characteristics; EM=Evidence Map; FC=formal caregiver; IC=informal caregiver; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RS=Reporting Status; SES=socioeconomic status; MoCA=Montreal Cognitive Assessment; CDR=Clinical Dementia Rating; GDS=Geriatric Depression Scale; CMAI=Cohen-Mansfield Agitation Inventory; BIP=Behavioral Observation Scale for Intramural Psychogeriatrics; SIPO=Social Interaction Scale for Psychogeriatric Older People; SF-8=Short-Form 8; DBD=Dementia Behavior Disturbance Scale; MADRS=Montgomery-Asberg Depression Rating Scale; HADS=Hospital Anxiety and Depression Scale; MOSES=Multidimensional Observation Scale for Elderly Subjects; MDS ADL=Minimum Data Set Activities of Daily Living Scale; STAI=State-Trait Anxiety Inventory; EQ-5D=EuroQoL 5 dimension scale; RSES=Rosenberg Self-Esteem Scale; QoL-AD: Quality of Life in Alzheimer's Disease scale; NPI=Neuropsychiatric Inventory; FAHW=Der Fragebogen zum allgemeinen habituellen Wohlbefinden (FAHW) [Questionnaire of General Habitual Well-being] scale; HCSS=Holden Communication Scale; MDS=Minimum Data Set...; ADL=Activities of Daily Living; RoB=Risk of Bias; NPI=Neuropsychiatric Inventory; QoL-AD: Quality of Life in Alzheimer's Disease scale; QCPR=Quality of Caregiver and Patient Relationship; DEMQOL=Dementia Quality of Life scale; CSDD=Cornell Scale for Depression in Dementia

Psychosocial Interventions

Table D-40. Risk of bias assessment: psychosocial interventions

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Onder 2005 ²⁰³ (16260821)	25 weeks	Medium 12%	Low	High	Medium	Low	High	Italian government	High

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Psychosocial Interventions

Table D-41. Characteristics of evidence map studies: psychosocial interventions

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:

Quinn 2016 ²⁰⁴ (26674087) UK Pilot	Self-management group intervention for 8 weeks: Weekly 90-minute sessions covering information sharing on various aspects such as enjoying hobbies, activities staying well, practical ways to manage memory difficulties, maintaining relationships and social networks, planning for the future, coping skills and accessing local resources.	TAU (n=11)	A memory clinic in a semi-rural area of North Wales RCT 24 PLWD	ICD-10 diagnosis of AD, vascular dementia, or mixed AD and vascular dementia with MMSE score \geq 20	N=24 Mean age: 76 years Female: 25% % majority race: no Education: 63% college/university	NR	N=24 Mean age: 67 years Female: 79% % majority race: no Education: :42% college/university Relationship with PLWD: 79% spouse/partner	Living with PLWD: 92%	NA	NA	3 months 6 months	ACE-III GSES HADS CORE-OM EQ-5D-3L ICECAP-O	NR
Quinn 2014 ²⁰⁵ (24606601) UK Pilot													
Young 2014 ²⁰⁶ (25587218) Hong Kong Small sample	10 weekly sessions of support group (structured, time limited, and allowed participants to share on different topics) for 90 minutes	standardized educational written material on dementia that provided basic information on dementia (n=19)	Non-governmental geriatric centers RCT 39 PLWD	Diagnosis of dementia, MMSE score \geq 18	N=39 Mean age: 80 years Female: 44% % majority race: no Education: 33% illiterate	NR	NA	NA	NA	NA	10 weeks	GDS-C RSES-C GSE-C IMMEL	NA
Tappen 2009 ²⁰⁷ (20077983) US Small sample	30 minutes modified counseling sessions, 3 times per week for 16 weeks	Usual care (n=15)	Long-term care facility RCT 36 PLWD	Clinical diagnosis of probable AD using NINCDS-ADRDA; an MMSE score \leq 25	N=30 Mean age=87 years Female: no % majority race: no Education: 39% illiterate	NR	NA	NA	NA	NA	6 months 12 months	DMAS ADRD MADRS	NA
Burns 2005 ²⁰⁸ (16055825) UK Small sample	Six, 50 minutes sessions of psychodynamic interpersonal therapy	TAU (n=20)	Referrals to memory clinic in South Manchester, UK RCT 40 PLWD	Diagnosis of AD according to NINCDS-ADRDA criteria with a clinical dementia rating of 1 and a score of 15 or above on MMSE	N=40 Mean age: 76 years Female: 48% % majority race: no Education: no	NR	NA	NR	NA	NA	6 weeks 3 months	CSDD BADLS RMPBC MMSE CIB-GIC	GHQ BDI WCC

Onder 2005 ²⁰³ (16260821) Italy High RoB	PLWD treated with donepezil got a reality orientation program. CG then trained to offer the group were trained to offer the program at home 3 days a week, 30 minutes/day, for 25 consecutive weeks, and were invited to stimulate and involve PLWD in reality-based communication.	No treatment	Community-dwelling PLWD RCT 156 PLWD	Diagnosis of AD according to NINCDS-ADRA with MMSE between 14 and 27, did not present with major aphasia or blindness and got donepezil treatment for at least 3 months	N=156 Mean age: 75.8 years Female: 72% % majority race: no Education: yes	NR	N=156 Mean age: 57 years Female: 63% % majority race: no Education: no	NA	NA	NA	25 weeks	MMSE ADAS-cog IADL NPI medication use	HRSD SF-36 Burden of care
Cohen 1999 ²⁰⁹ (10495580) US Quasi-experimental	The Buddy Program: Nondemented residents were trained via 8 hours educational sessions and acted as "buddies" to their demented partner and encouraged to spend at least 30 minutes of social interaction (game/music/book/communication)	No training or partners (n=31)	Nursing homes Quasiexperimental 73 PLWD (assigned; not randomized)	MMSE score ≤ 24 were considered as potential dementia participants but did not show memory or verbal impairment	N=73 Mean age: 80 years Female: 70% % majority race: 50% African American Education: yes	NR	NA	NR	NA	NA	3 months 6 months	MMSE GDS CSDD CMAI BEHAVE-AD FAST PGCMS NPD ADL ESS NSS	NA
Wallis 1983 ²¹⁰ (6354248) UK Small Sample	Reality Orientation therapy for 30 minutes, 5 days/week for 3 months	A variety of group and individual activities was offered to them each day (attention control)	High Royds Hospital Quasiexperimental Unclear	No diagnosis criteria mentioned	N=38 Mean age: 70 years Female: 34% % majority race: no Education: no	NR	NA	NR	NA	NA	2 weeks 3 months 4 months	Crichton scale for behavioral functioning Cognitive abilities	NA

Abbreviations: ACE-III=Addenbrooke's Cognitive Examination-III; AD=Alzheimer's disease; ADAS-cog=Alzheimer's Disease Assessment Scale-Cognitive Subscale; ADL=Activities of daily living; ADRD=Alzheimer's Disease and Related Disorders; BADLS=Bristol Activities of Daily Living Scale; BDI=Beck Depression Inventory; BEHAVE-AD=Behavioral Pathology in Alzheimer's Disease scale; char=characteristics; CIB-GIC=Clinician's Interview-Based Global Impression of Change; CG-Caregiver; CMAI=Cohen-Mansfield Agitation Inventory; CORE-OM=Clinical outcomes in routine evaluation-outcome measure; CSDD=Cornell Scale for Depression in Dementia; DMAS=Dementia Mood Assessment Scale; EQ-5D-3=3-level version of EuroQol; ESS=Environmental Satisfaction Scale; FAST=Functional Assessment Staging Tool; GDS=Geriatric Depression Scale; GDS-C=Chinese Geriatric Depression Scale; GHQ=General Health Questionnaire; GSE-C=General Self-Efficacy Chinese; GSES=General self-efficacy scale; HADS=Hospital anxiety and depression scale; HRSA=Hamilton Rating Scale for Anxiety; HRSD=Hamilton Rating Scale for Depression; IADL=Instrumental Activities of Daily Living; ICECAP-O=Icepop Capability measure for Older people; IMMEL=Index for Managing Memory Loss; MADRS=Montgomery-Asberg Depression Rating Scale; MMSE=Mini-Mental Status Exam; N=Number; NA=Not Applicable; NINCDS-ADRDA=National Institute of Neurological and Communicative Disorders and Stroke/Alzheimer's Disease and Related Disorders Association; NPD=Number of Physical Disorders; NPI=Neuropsychiatric Inventory; NR=Not reported; NSS=Nursing Stress Scale; PGCMS=Philadelphia Geriatric Center Morale Scale; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RCT=Randomized controlled trial; RMBPC=Revised Memory and Behavior Problems Checklist; RoB=Risk of Bias; RS=Reporting status; RSES-C=Rosenberg Self-Esteem Scale-Chinese; SF-36=36 Item Short Form Survey; TAU=Treatment as usual; WCC=Ways of Coping Checklist

Creative Expression Therapy

Table D-42. Risk of bias assessment: creative-expression therapy

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Fritsch 2009 ²¹¹ (19363009)		High	X	X	X	X	X	Nonprofit	High

X indicates domain not rated due to high ROB.

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Creative-Expression Therapy

Table D-43. Characteristics of evidence map studies: creative-expression therapy

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Lin 2019 ²¹² (31440055) China Pilot	Creative Expression Therapy (TimeSlip) twice a week for 6 weeks	Standard Cognitive Training	Long-Term Care Facilities, Prospective non-blinded RCT, 100 PLWD	Not specified	N=91 (43 treatment vs 48 control) Age: yes Sex: yes Race: yes Education: yes	NR	NA	NA	NA	NA	6 weeks	Cognitive Function, Quality of Life, Depression, Communication ability, Emotion status	NA
Houser 2014 ²¹³ (23602304) US Small sample	Two 1-hour TimeSlip sessions per week for 6 weeks	Facilities standard of care	Nursing homes Cluster RCT; 1 site 20 PLWD	Not specified	N=20 (10 treatment vs 10 control) Age: yes Sex: yes % majority race: no Education: no	NR	NA	NA	NA	NA	4 months	Mood score Behavior score	NA

Van Dijk 2012 ²¹⁴ (15802911) Netherlands Pilot	Group 1: Veder method living room theater group with trained care professionals Group 2: Veder method living room group with professional actors	Standard Reminiscence group activity	Psychiatric Nursing Home, Quasi- experimental non-equivalent three group, Multisite, 155 PLWD, 20 Trained professional caregivers, 10 professional actors, 11 therapists and nurses	Alzheimer, Vascular dementia, Dementia NOS	N=155 (69 treatment 1, 31 treatment 2 vs 55 control) Age: yes Sex: yes % majority race: no Education: yes	NR	NR	NR	NR	NR	1 week	Behavior Mood Quality of Life Alertness Memory recall Social isolation behaviors	NA
Phillips 2010 ²¹⁵ (21048483) US Pilot	TimeSlip sessions twice a week for 6 weeks	Facilities standard of care	Nursing homes and Assisted Living Facilities, Quasi- experimental repeated measures, Multisite 56 PLWD	Not specified	N=56 (28 treatment vs 28 control) Age Sex Race Education	Household characteristics	NA	NA	NA	NA	6 weeks	Cornell Scale for Depression in Dementia, Neuropsychiatric Inventory- Nursing Home Version, Functional Assessment of Communication Skills, Quality of Life- AD, Observed Emotion Rating Scale	NA
Fritsch 2009 ²¹¹ (19363009) US High RoB	Staff members trained in TimeSlip; TimeSlips group once a week for 1 hour for 10 weeks	Facilities' standard of care	Nursing homes, randomized matched pair, 20 sites, Undefined N PLWD, Undefined N Daytime staff	Not specified	NR	NR	NA	NA	N=not given Age: yes Sex: yes Race: yes Education: no	Health status: no Training: yes Education: no Position: yes Service: yes	Two weeks	Resident Engagement, Resident Affect	FC: Staff- Initiated interactions with Residents, Attitudes towards PLWD, Job Satisfaction, Burnout

Abbreviations: AES=Apathy Evaluation Scale; APADEM-NH=Apathy Scale for Institutionalized Patients with Dementia Nursing Home; AWS=Revised Algate Wandering Scale-Nursing Home version; BARS=Brief Agitation Rating Scale; CMAI-SF=Cohen-Mansfield Agitation Inventory-Short Form; CSDD=Cornell Scale for Depression in Dementia; DSM=Diagnostic and Statistical Manual; EM=Evidence Map; GDS=Global Deterioration Scale; MMSE=Mini-mental State Examination; N=number; NA=Not Applicable; NIA=National Institute on Aging; NPI=Neuropsychiatric Inventory; NR=Not Reported; OERS=Observed Emotion Rating Scale; PMID=PubMed Identification Number; QoL-AD=Quality of Life in Alzheimer's Disease; QUALID=Quality of Life in Late-stage Dementia; RAID=Rating for Anxiety in Dementia; PLWD=Patient with Dementia; NOS=not otherwise specified; FC=Formal Caregiver

Multicomponent for PLWD Well-Being

Table D-44. Risk of bias assessment: multicomponent for PLWD well-being

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Overall Rating
Okamura 2018 ²¹⁶ (30036871)	3 months	Low 3 months: 1%	Medium	High	High	Low	Low	High
Rokstad 2018 ²¹⁷ (30445937)	24 months	NA	High	X	X	X	X	High
Masayuki Satoh 2017 ²¹⁸ (28222531)	6 months	High 6 months:27%	Medium	X	X	X	X	High
Middelstadt 2016 ²¹⁹ (27497474)	14 weeks	Medium 14 weeks:20%	Low	Low	Medium	Low	High	High
Fernandez-Calvo 2015 ²²⁰ (25121567)	16 weeks	Low 16 weeks: 10%	Medium	Low	High	Low	High	High
Luttenberger 2012 ²²¹ (22468985)	6 months	Medium 6 months: 16%	Low	High	Medium	Low	Medium	Medium
Yamagami 2012 ²²² (23300492)	12 weeks	Low 12 weeks: 1%	Medium	High	High	Low	Medium	High
Brooker 2011 ²²³ (21702705)	18 months	High 18 months: 32%	Low	X	X	X	X	High
Ferrero-Arias 2011 ²²⁴ (21346517)	4 weeks 8 weeks	High Unclear	Medium	X	X	X	X	High
McCurry 2011 ²²⁵ (21797835)	2 months	Low 2 months: 13%	Low	Medium	High	Low	Low	Medium
McCurry 2011 ²²⁵ (21797835)	6 months	Low 6 months: 16%	Low	Medium	High	Low	Low	Medium
Femia 2007 ²²⁶ (18192631)	2 months	X	High	X	X	X	X	High
Chapman 2004 ²²⁷ (15603468)	4 months	High 4 months: 24%	Low	X	X	X	X	High
Tadaka 2004 ²²⁸ (No PMID)	3 months	Low 3 months: 8%	Low	Medium	Medium	Low	High	Medium
Tadaka 2004 ²²⁸ (No PMID)	6 months	Medium 6 months: 17%	Low	Medium	Medium	Low	High	Medium
Wimo 1993 ²²⁹ (8356361)	12 months	X	High	X	X	X	X	High

X indicates a domain not assessed due to high ROB.

Abbreviations: PMID=PubMed Identification Number; PLWD=persons with dementia

Table D-45. Characteristics of included studies: multicomponent for PLWD well-being

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Luttenberger 2012 ²²¹ (22468985) Germany Medium Explanatory	MAKS intervention consisting of motor stimulation, ADL activities and cognitive stimulation	Standard care	Nursing homes RCT	PLWD diagnosed with primary degenerative dementia MMSE <24 Exclusion criteria	N=30 Intervention group 22 Control group Age: Y Sex: Y Education: Y Race % majority: N	Marital Status	NA	NA	Education: Y	Training: Y	12 months Additional follow-up after 10 additional months	ADL	NA	
McCurry 2011 ²²⁵ (21797835) US Medium Explanatory	Increased walking and/or bright light exposure to improve sleep	Contact only	Community-based RCT	PLWD diagnosed with probable or possible AD Two or more sleep problems multiple times a week Average wake time of 1 hour per night or greater	N=27 Walking group 29 Light group 33 Insomnia Treatment and Education group Age: Y Sex: Y Education: Y Race % majority: Y	NR	Age: Y Sex: Y Education: Y Race % majority: Y Spouse or other: Y	NR	Education: Y Experience: Y	Training: Y	6 months	Sleep Disorders Inventory Secondary sleep outcomes measurements	Satisfaction survey	
Tadaka 2004 ²²⁸ (PMID NR) Japan Medium Explanatory	Group care program of reminiscence and reality orientation care	Routine care	Community-based RCT	PLWD diagnosed with AD or vascular dementia CDR score of 1 or 2 Exclusion criteria	N=2 Intervention group 26 Control group Age: Y Sex: Y Education: N Race % majority: N	NR	Relation to PLWD: Y	NR	Education: Y Experience: Y	Training: Y	10 weeks with follow-up at 6 months	MMSE MOSES Self-care Disorientation Depression Irritability Withdrawal	NR	

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: AD=Alzheimer's disease; ADL=Activities of Daily Living; -char=characteristics; FC=formal caregiver; IC=informal caregiver; N=number; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; CST=Cognitive Stimulation Therapy; NR=Not Reported; DSM-IV=Diagnostic and Statistical Manual for Mental Disorders, Fourth Edition; PLWD=Person With Dementia; RCT=Randomized controlled trial

Table D-46. PLWD outcomes summary low and medium risk of bias studies: multicomponent for PLWD well-being

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Luttenberger 2012 ²³⁰ (22468985) MAKS therapy vs. Usual care Medium Explanatory	E-ADL test (Range 0 to 30, higher better) Cohen's d 12 months	No difference between groups for ITT analysis (Cohen's d 0.23) Significant difference in PP analysis (d=0.50)	NR	NR	NR
McCurry 2011 ²²⁵ (21797835) NITE-AD (Sleep education) vs. contact control Medium Explanatory	Sleep Disorders Inventory Daily mean (SE) 6 months	No difference between groups	0.8 (0.2)	0.5 (0.1)	0.12

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Tadaka 2004 ²²⁸ (PMID NR) Group care program consisting of reminiscence and reality orientation care methods in addition to a routine day-care service vs. Routine day-care service Medium Explanatory	Self-care category of MOSES LS Mean (SE) adjusted for baseline and clinical dementia rating 6 months	No difference between groups	16.1 (1.0)	18.5 (1.0)	0.05
Tadaka 2004 ²²⁸ (PMID NR) Group care program consisting of reminiscence and reality orientation care methods in addition to a routine day-care service vs. Routine day-care service Medium Explanatory	Disorientation category of MOSES LS Mean (SE) adjusted for baseline and clinical dementia rating 6 months	Favors intervention (lower score better)	16.0 (1.3)	18.3 (1.3)	0.01
Tadaka 2004 ²²⁸ (PMID NR) Group care program consisting of reminiscence and reality orientation care methods in addition to a routine day-care service vs. Routine day-care service Medium Explanatory	Depression category of MOSES LS Mean (SE) adjusted for baseline and clinical dementia rating 6 months	No difference between groups	17.1 (1.2)	15.9 (1.2)	0.97
Tadaka 2004 ²²⁸ (PMID NR) Group care program consisting of reminiscence and reality orientation care methods in addition to a routine day-care service vs. Routine day-care service Medium Explanatory	Irritability category of MOSES LS Mean (SE) adjusted for baseline and clinical dementia rating 6 months	No difference between groups	14.9 (1.0)	15.0 (1.0)	0.34
Tadaka 2004 ²²⁸ (PMID NR) Group care program consisting of reminiscence and reality orientation care methods in addition to a routine day-care service vs. Routine day-care service Medium Explanatory	Withdrawal category of MOSES LS Mean (SE) adjusted for baseline and clinical dementia rating 6 months	Favors intervention (lower score better)	16.6 (1.1)	19.5 (1.1)	0.02

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number

Table D-47. Summary of strength of evidence for PLWD outcomes: multicomponent for PLWD well-being

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Instrumental Activities of Daily Living Multicomponent: motor stimulation, activities of daily living, and cognitive stimulation components contact control	6 months	1 RCT (n=98)	No difference in instrumental activities of daily living	Medium	Unknown	Direct	Imprecise	Insufficient
Sleep Walking, light, combination treatment versus contact control	6 months	1 RCT (n=66)	No difference in sleep disorders inventory	Medium	Unknown	Direct	Imprecise	Insufficient
Cognitive and Social Function Reminiscence and reality orientation care methods, routine day-care service versus usual care	6 months	1 RCT (n=60)	No difference in self-care, depression, or irritability Favors intervention for withdrawal and disorientation	Medium	Unknown	Direct	Imprecise	Insufficient

Evidence Map: Multicomponent for PLWD Well-Being

Table D-48. Characteristics of evidence map studies: multicomponent for PLWD well-being

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Sado 2020 ²³¹ (32116248) Japan Pilot	Combination of cognitive training and stimulation	Unspecified control group volunteers from within nursing home	Nursing homes Controlled Trial (CT) Cluster N=23 N=57 (non-randomized)	Aged 6 years or older Diagnosis of dementia Long-term-care insurance MMSE between 10 and 26 Living in nursing homes; No previous experience of the intervention	N=30 Age: Yes Sex: Yes Race: No Education: No	NR	NA	NA	NR	NR	12 months	ADL	NR

Young 2019 ²³² (29781725) Hong Kong Pilot	Structured cognitive stimulation therapy and tai chi	Treatment as usual; waitlist	Community-based RCT with waitlist for control group 101 PLWD	PLWD diagnosed with mild stage dementia MMSE \geq 18 Aged 60 or older	N=101 Age: Yes Sex: Yes Race % majority: No Education: Yes	Household characteristics: Yes	NR	NR	NA	NA	7 weeks	DSR dementia rating tests MMSE	NA
Chen 2018 ²³³ (29881275) Taiwan Small sample	Music-based dual task training	Non-music-based walking	Community-based RCT 28 PLWD	PLWD diagnosed with mild to moderate dementia CDR score 0.5, 1 or 2; aged 55 or older, ambulatory Exclusion criteria applied	N=28 Age: Yes Sex: Yes % majority race: No Education: No	NR	NR	NR	NR	NR	2 months; up to 3 make-up sessions allowed	Evaluation of walking and stride Ability to stand Fall efficacy measurement Agitation Memory inventory	NA
Okamura 2018 ²¹⁶ (30036871) Japan High RoB	Combined exercise and cognitive training	Pedaling with arm ergometer	Combined day care and residential care RCT 100 PLWD	PLWD with diagnosis of dementia No restriction on type or severity Aged 65 or older and requiring assistance Exclusion criteria applied	N=100 Age: Yes Sex: Yes Race % majority: No Education: No	NR	NA	NA	NR	NR	6 weeks; 1 month 3 months	MMSE Trail making test ADL	NA
Reisberg 2017 ²³⁴ (28122366) US Small sample	Comprehensive, Individualized, Person-Centered Management with following components: CG training, management assessment, therapeutic home visits and CG support groups	Usual community care	Community dwelling PLWD and CG RCT 20 PLWD	Diagnosis based on DSM-IV-TR criteria fulfilled NINCDS-ADRDA criteria for probable AD; moderate-to-severe AD with GDS 5 or 6; FAST \geq 6a; MMSE between 3 and 14	N=20 Age: yes Sex: yes Race % majority: no Education: yes	NR	NR	NA	NA	NA	4 weeks 12 weeks 28 weeks	NYU-CIBIC-Plus ADCS-ADLsev-abv MMSE FAST-DS BEHAVE-AD-FW RMBPC	NA
Rokstad 2018 ²¹⁷ (30445937) Norway High RoB	Attending daycare-based dementia programs at least 2 days per week	Care as usual	Community dwelling PLWD using daycare services and their CG Quasi-experimental 257 PLWD recruited	Dementia diagnosis according to ICD-10; MMSE \geq 15 and consent of CG	N=257 Age: yes Sex: yes Race % majority: no Education: yes	NR	N=257 Age: yes Sex: yes Race % majority: no Education: no Relation to PLWD=yes	NR	NA	NA	24 months	PSMS IADL MADRS NPI-Q REED	RSS MADRS
Kampragkou 2017 ²³⁵ (ID# not available) Greece Small sample	Aerobic exercise, memory games and music therapy	No intervention	Institutional setting RCT 36 PLWD	PLWD diagnosed with moderate AD, ambulatory, aged 65 or older Exclusion criteria applied	N=30 Age: No Sex: Yes Race % majority: No Education: No	NR	NA	NA	NA	NA	12 weeks	MMSE ADAS ADL Ability to stand Balance	NA

Masayuki Satoh 2017 ²³⁶ (28222531) Japan High RoB	Physical exercise with music (muscle training for the upper and lower extremities, hand clapping to music, breath and voice training, and singing) for 40 minutes 1x week for 6 months	Cognitive stimulation using a portable game console (Nintendo DS LL), and drills consisting of easy calculations, mazes, and mistake-searching in pictures for 40 minutes 1x week for 6 months	PLWD using geriatric nursing services RCT 85 PLWD	Neurological specialist diagnosed dementia according to ICD-10; MMSE score between 16 and 26;	N=85 Age: yes Sex: no Race % majority: no Education: yes	NR	NA	NR	NA	NA	6 months	RCPM RBMT WF TMT-A FIM BEHAVE-AD VSRAD	NA
Kim 2016 ²³⁷ (ID# not available) South Korea Small sample	Cognitive stimulation, including art, music, recollection and horticultural therapy	Pharmacological treatment only	Institutional setting RCT 64 PLWD	PLWD diagnosed with probable AD Exclusion criteria applied	N=53 Age: Yes Sex: Yes Race % majority: No Education: Yes	NR	NR	NR	NR	NR	6 months	MMSE Verbal fluency and word recall Depression Judgement and problem solving QoL	QoL
Middelstadt 2016 ²¹⁹ (27497474) Germany High RoB	Cognitive stimulation program twice weekly for eight weeks	Usual care	PLWD recruited from long term care facilities RCT 71 PLWD	Mild to moderate PLWD diagnosed according to ICD-10 and MMSE score 10-25	N=71 Age: Yes Sex: Yes Race % majority: No Education: No	NR	NA	NA	NA	NA	8 weeks 14 weeks	ADAS-cog QoL-AD NPI-NH ADCS-ADL	NA
Fernandez-Calvo 2015 ²²⁰ (25121567) Spain High RoB	Multicomponent intervention including new recreational and cognitive tasks, a module of functional activities based on real-life situations, cognitive training exercises to carry out at home with the caregiver; and psychoeducational activities; 48 sessions (90 minutes duration, and 3 sessions per week) for 16 weeks	Waitlist group	Community dwelling PLWD RCT 61 PLWD	Diagnosis of probable AD using NINCDS/ADRDA; mild dementia; MMSE≥18 and having a CG for at least 6 months	N=61 Age: yes Sex: yes Race % majority: no Education: yes	NR	NA	NR	NA	NA	16 weeks	ADAS-Cog NPI-Q CSDD RDRS-2	NA

Wesson 2013 ²³⁸ (24004682) Australia Pilot	Home hazard reduction program to reduce falls and increase balance and strength	Usual care	Community-based RCT 22 PLWD and CG	PLWD diagnosed with dementia or ACE-R score <=82 Age 65 or older with non-paid caregiver MMSE.82 Exclusion criteria applied	N=22 Age: Yes Sex: Yes Race % majority: No Education: Yes	Household characteristics: Yes Prior disability: Yes	Sex: Yes Relation to PLWD: Yes	Living with PLWD	NA	NA	12 weeks	Fall risk tests Step and balance tests ADL Depression Agitation	Caregiver burden Task strategy management index
Yamagami 2012 ²²² (23300492) Japan High RoB	24 sessions for 1-hour 2x week, for 12-weeks of intervention based on brain-activating rehabilitation combined reality orientation and various activities (e.g. cooking, singing, and sewing) along with reminiscence therapy.	No treatment	Recruited from residential care homes (group living homes) RCT 54 PLWD	Dementia diagnosis without severe visual or auditory impairments	N=54 Age: Yes Sex: Yes Race % majority: No Education: No	NR	NA	NA	NA	NA	12 weeks	CDR MOSES HDS-R TMT-A Staff interview	NA
Brooker 2011 ²²³ (21702705) UK High RoB	Occupational activities to foster well-being and goal achievement	Trained staff employed to try to increase PLWD activities	Assisted living Cluster RCT 293 PLWD	PLWD exhibiting confused behavior, communication, difficulties, challenging behavior or low mood Exclusion criteria applied	N=293 Age: Yes Sex: Yes Race % majority: No Education: No	Household characteristics: Yes	NA	NA	NR	Training: Yes Position: Yes	18 months	QoL Level of social support Quality of relationships Dementia Care Mapping index	NA
Burgener 2011 ²³⁹ (20509596) US Small sample	Taiji exercises, cognitive-behavioral therapies and cognitive training exercises, as well as support group participation	Education program for caregivers	Community-based RCT Self-referred, caregiver referred, or provider referred Unclear	PLWD diagnosed with dementia at early to middle disease stage	N=32 Age: No Sex: No Race% majority: No Education: No	NR	N=32 Age: Yes Sex: Yes Race % majority: No Education: Yes	NR	NA	NA	20 weeks	MMSE ECFR RSS	Stress Quality of family relationship
Ferrero-Arias 2011 ²²⁴ (21346517) Spain High RoB	Occupational therapy sessions (music and art therapy and psychomotor activity)	Control participants remained in the day rooms performing activities of their own choosing	Recruited from centers of the Association of Alzheimer PLWD' Families of Burgos Crossover RCT 146 PLWD	Dementia criteria of CDR 1 or 2, undergoing occupational therapy	N=146 Age: yes Sex: yes Race% majority: No Education: yes	NR	NA	NA	NA	NA	4 weeks 8 weeks	NPI-Q DAIR	NA

Burgener 2008 ²⁴⁰ (18453642) US Small sample	Taiji exercises, cognitive behavioral therapies and support group	Attention control education program	Community-based RCT Self-referred, caregiver referred, or provider referred 43 PLWD	PLWD diagnosed with dementia at early to middle disease stage	N=43 Age: Yes Sex: Yes Race % majority: No Education: Yes	NR	NR	NR	NA	NA	20 weeks; 40 weeks	MMSE Physical functioning Depression Self esteem	NR
Femia 2007 ²²⁶ (18192631) US High RoB	Adult day service 3 days per week for 6 hours for 2 months: 5 to 6 different activities per day in addition to daily routines (i.e., lunch), which included 30 minutes of physical activities on average, 1 to 2 hours of social activities, and about 1 hour of cognitively stimulating activities	Not using adult day services	Community dwelling PLWD and their CG participating in the Dementia Day Care Program of the NJ Statewide Respite Care Program Quasi-experimental 234 PLWD recruited	Dementia diagnosis	N=201 Age: no Sex: yes Race % majority: no Education: yes	Income=yes	N=201 Age: yes Sex: yes Race % majority: yes Education: yes Relation to PLWD:yes	Employment=yes	NA	NA	2 months	BPSD Domains IADL PADL MMSE DRB	NA
Chapman 2004 ²²⁷ (15603468) US High RoB	Cognitive-communication program plus an acetylcholinesterase inhibitor (donepezil): 1.5-hour sessions 1x week for 8 weeks	Donepezil treatment	Community-dwelling participants RCT 54 PLWD	Dementia diagnosis according to NINCDS/ADRDA and MMSE ≥12	N=54 Age: yes Sex: no Race % majority: no Education: yes	NR	NA	NA	NA	NA	4 months 8 months 12 months	ADAS-cog TFLS NPI CIBIC QoL	NA
Tappen 2000 (11186596) US	Skill training program	Traditional stimulation or regular care	Nursing home RCT 3 groups	PLWD diagnosed with dementia Able to stand with assistance Exclusion criteria applied	N=21 skill group 21 stimulation group 21 regular care group Age: NR by group Sex: NR by group Race % majority: N Education: N	NR	NA	NA	Education: Y	NR	20 weeks	Self-maintenance ADL Goal attainment	NA
Wimo 1993 ²²⁹ (8356361) Sweden High RoB	6 hours of daycare activities including everyday routines such as meals, coffee, short walks, and excursions with stimulating elements, e.g. newspaper reading and videos	Not attending daycare	Non-institutionalized PLWD attending daycare in Sundsvall Quasi-experimental 99 PLWD	Dementia diagnosis according to DSM-III-R criteria	N=99 Age: yes Sex: yes Race % majority: no Education: no	Home support=yes	NA	NA	NA	NA	12 months	MMSE MDDAS institutionalization	NA

Abbreviations: AD=Alzheimer's disease; ADAS-cog=Alzheimer's Disease Assessment Scale, cognitive subsection; ADCS-ADL=Alzheimer's Disease Cooperative Study-Activities of Daily Living Inventory; ADCS-ADLsev-abv=Alzheimer's Disease Cooperative Study Activities of Daily Living Inventory modified for more severe dementia, abbreviated first 12 questions version; ADL=Activities of Daily Living; BEHAVE-AD=Behavioral pathology in Alzheimer's disease; BEHAVE-AD=FW=Behavioral Pathology in Alzheimer's Disease Frequency-Weighted Severity Scale; BPSD=Behavioral and psychological symptoms of dementia; CDR=Clinical Dementia Rating; CG=Caregiver; char=characteristics; CIBIC=Clinician Interview-Based Impression of change; CSDD=Cornell Scale for Depression in Dementia; DAIR=Dementia Apathy Interview and Rating; DRB=Daily Record of Behavior; DSM-IV-TR=Diagnostic and Statistical Manual for Mental Disorders, Fourth Edition, Text Revision; DSR=Dementia Rating Scale; ECFR=Elder-Caregiver-Family Relationship scale; EM=Evidence Map; FAST-DS=Functional Assessment Staging-Disability Score; FC=formal caregiver; FIM=Functional independence measure; HADS=Hasegawa dementia scale revised; IADL=Instrumental Activities of Daily Living; ICD-10=International Classification of Diseases, Tenth Revision; IC=informal caregiver; MADRS=Montgomery-Asberg Depression Rating Scale; MDDAS=Multi-Dimensional Dementia Assessment Scale; MMSE=Mini-Mental Status Examination; MOSES=Multidimensional Observation Scale for Elderly Subjects; N=number; NA=Not Applicable; NINCDS-ADRDA=National Institute of Neurological and Communicative Diseases and Stroke/Alzheimer's Disease and Related Disorders Association; NJ=New Jersey; NPI=Neuropsychiatric Inventory; NPI-NH=Neuropsychiatric Inventory Home Version; NPI-Q=Neuropsychiatric Inventory Questionnaire; NR=Not Reported; NYC-CIBIC Plus=New York University Clinician's Interview-Based Impression of Change-Plus Caregiver Input; PADL=Physical Activities of Daily Living Scale; PMID=PubMed Identification Number; PSMS=Physical Self-Maintenance Scale; PLWD=Persons with Dementia; QoL=Quality of Life; QoL-AD: Quality of Life in Alzheimer's Disease scale; RBMT=Rivermead Behavioral Memory Test; RCPM=Raven's Colored Progressive Matrices-Japanese; RDRS-2=Rapid Disability Rating Scale – Version 2; REED=The Anosognosia Rating Scale; RMBPC=Revised Memory and Behavior Problems Checklist; RoB=Risk of Bias; RS=Reporting Status; RSS=Relatives' Stress Scale; SES=socioeconomic status; TFLS=Texas Functional Living Scale; TMT-A=Trail-Making Test A; WF=Word Fluency; VSRAD=Voxel-based specific regional analysis system for Alzheimer's disease

Assistive Technology

Evidence Map: Assistive Technology

Table D-49. Characteristics of evidence map studies: assistive technology

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Lauriks, 2018 ²⁴¹ Amsterdam Pilot	Application of Assistive Home Technology (AHT); life circles, pathway lighting, automated lighting, automated alerts, automated sun blinds	Lived in same type of group home, but w/out AHT	Group homes in residential care facility (N=9) RCT PLWD (N=54) Caregivers (N=25)	Diagnosis of dementia confirmed by a physician	N=54 (30 treatment vs 24 control) Age: yes Sex: yes Race: NR	NR	NA	NA	N=25 (14 treatment vs 11 control) Sex	Training	On average four months after baseline measurement completed and AHT switched on.	Quality of life (Qaulidem instrument)	FC: Job satisfaction and general health
Tchalla, 2013 ²⁴² France Pilot	Home-based technologies coupled with teleassistance service (HBTec-TS); nightlight path and electronic bracelet coupled with a teleassistance service	No HBTec-TS system was implemented	Community Experimental prospective study	Suffering from Alzheimer's disease, living at home and registered on the frail elderly people register Considered high risk of falling and were receiving a county allocation	N=96 (49 treatment vs 47 control) Age: yes Sex: yes Race: NR	Comorbidity	NA	NA	NA	NA	12 months	Incidence of falls	NA
Rowe, 2010 ²⁴³ United States Pilot	Home monitoring system designed to reliably awaken a caregiver when the PLWD left the bed at night	Paid \$15 at each data collection point and provided with some education material unrelated to any study goals	Community Pretest-posttest control group design Caregivers (N=49)	Medical diagnosis of Alzheimer's disease or other dementia	NR	NR	N=49 Age Sex Race Education Relation to PLWD	Living with PLWD Health status Employment status	NA	NA	Post-test months; 2, 3, 4, 5, 6, 8, 10 and 12 (7 consecutive nights at each data collection month)	NA	IC: Subjective sleep, objective sleep and distress
Gaugler 2019 (29982413) ²⁴⁴ United States Pilot	Wearable / on-body monitoring system alerts caregiver to activities, provides reports behavioral patterns associated potential health problems	Usual care	Community RCT PLWD (N=132) Caregivers (N=132)	English speaking Diagnosis of NINCDS/ADRDA Not receiving similar services Age ≥ 55 years	N=132 Age yes Sex yes Race yes Education yes	SES Household Characteristics Health Insurance Race Information	N=132 Age yes Sex yes Race yes Education yes Relation to PLWD: yes	Living with PLWD Employment Status	NA	NA	6 months	NA	Sense of competence (SSCQ), Caregiver Distress (ZBI), Depression (CES-D)

Abbreviations: CES-D=Center for Epidemiologic Studies Depression Scale; NA=Not Applicable; NINCDS/ADRDA= National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's disease and Related Disorders Association; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Patients living with Dementia; SSCQ=Short Sense of Competence Questionnaire; ZBI=Zarit Burden Index

Electrostimulation

Evidence Map: Electrostimulation

Table D-50. Characteristics of evidence map studies: electrostimulation

Study (PMID)	Intervention	Comparison	Setting and Design	PWD Dementia Characteristics	PWD Non-Disease Char	PWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PWD N PWD Age (mean) PWD Sex (% female) PWD Race (% majority) PWD Education (mean years)	PWD SES PWD Prior Disability PWD Household Characteristics PWD Health Insurance PWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PWD (% majority)	IC Duration IC Living With PWD IC Payment Status IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Elder 2019 (30658705) UK Small sample	2 consecutive 20-min sessions of active (0.048mA/cm ²) tDCS separated by a 30-min break, over 5 consecutive days. The anodal electrode was applied to the right parietal cortex (P4) and the cathodal electrode was applied to the occipital cortex.	Sham tDCS	Community dwelling PLWDs RCT 40 PLWD	Clinical diagnosis of Lewy body dementia; experiencing visual hallucinations of a moderate to severe nature and MMSE score ≥ 12	N=40 Mean age: 76 years % Female: no % majority race: no Education: no	None	NA	None	NA	NA	Day 5 1 month 3 months	MMSE CAMCOG GDS-15 TMT A/B ODFAS computerized attentional and visuoperceptual tasks	NPI CAF ODFAS CGI
Khedr 2019 ³⁴⁵ (30940012) Egypt Small sample	tDCS, 20 minutes on each side. 2 weeks, 10 sessions	21 PLWD got tDCS	1 clinical center RCT N=44	Diagnosis of probable AD	N=23 Age: yes Sex: yes Education: yes Race: no	NR	NA	NA	NA	NA	2 weeks	MMSE MoCA Clock drawing test Cornell Depression Scale	NA
Sabbagh 2019 (31879235) US and Israel Safety study	Combines short bursts of rTMS with computerized cognitive training. Daily sessions applied across three targeted regions, with a total of 1300 rTMS pulses at 10 Hz in short bursts of 20 pulses.	Sham rTMS.	Community dwelling PLWDs RCT 129 PLWD	Diagnosis of mild to moderate AD using DSM-IV and MMSE score of between 18 to 26; ADAS-Cog >17;	N=129 Mean age: 77 years % Female: 46 % majority race: no Education: yes	None	NA	None	NA	NA	Week 7 week 12	ADAS-Cog ADCS-CGI-C	NA

Zhang 2019 (xxx) China Small sample	Repetitive transcranial magnetic stimulation combined with cognitive training: rTMS operation lasted approximately 10 minutes at one targeted encephalic region. Every day, the magnetic coil was first placed over the left DLPFC and then over the left temporal lobe. The treatment protocol was applied for 4 weeks (5 times/week) and with no maintenance sessions	Sham rTMS	PLWDs were recruited from Department of Neurology in Tongji Hospital at Huazhong University of Science and Technology (Wuhan, China) RCT 30 PLWD	Diagnosis of AD (mild to moderate) using NINCDS-ADRDA	N=28 Mean age: 69 years % Female: 79 % majority race: no Education: no	None	NA	None	NA	NA	Week 4 week 8	ADAS-Cog MMSE ACE-III ADL NPI	NA
Lozano 2016 ²⁴⁶ (27567810) Canada, US Small sample Leoutsakos 2018 ²⁴⁷ (29914028) Canada, US	12 months continuous Fornix deep brain stimulation post-surgery	21 control PLWD got sham procedure	7 centers in Canada and the US RCT- Phase II N=42	Mild dementia with CDR-SB of 0.5 or 1 or ADAD-Cog 13 of 12 to 24	N=21 Age: yes Sex: yes Race: no Education: no	NR	NA	NA	NA	NA	12 months	Adverse events Cerebral glucose metabolism Effects of age on metabolism	NA
Wu 2015 ²⁴⁸ (26977125) China Pilot	Risperidone 1 mg per day plus rTMS treatments for a total of 20 sessions, 5 days a week for 4 consecutive weeks	Risperidone 1 mg per day plus sham rTMS treatments for a total of 20 sessions, 5 days a week for 4 consecutive weeks	1 clinical center (The Wuxi Mental Health Center) RCT N=54	Diagnosis of probable AD according to NINCDS/ ADRDA criteria with MMSE score less than 24 and BEHAVE-AD score greater than 8	N=27 Age: yes Sex: yes Education: yes Race: no	NR	NA	NA	NA	NA	4 weeks	BEHAVE-AD ADAS-Cog Vital signs TESS Routine blood tests, urine tests, electrocardiogram, blood biochemistry adverse events	NA
Cotelli 2014 ²⁴⁹ (24678298) Italy Small sample	AtDCS (25 minutes) plus IC Memory training (5 days per week) for 2 weeks	12 PLWD got PtDCS (25 minutes) plus IC Memory training (5 days per week) 12 PLWD got AtDCS (25 minutes) plus motor training (5 days per week)	1 clinical center 3 arm RCT N=36	Diagnosis of mild to moderate AD, according to the NINCDS-ADRDA criteria,	N=12 Age: yes Sex: yes Race: no Education: yes	NR	NA	NA	NA	NA	3 months 6 months	MMSE ADL IADL Balance and gait NPI Language Complex figure copy Attention and executive function	NA
Olazaran 2014 ²⁵⁰ (24898637) Spain Pilot study	2 hours of REAC NPO per day for 2 weeks	60 PLWD were treated with sham procedure	Residential care facility Crossover RCT Single site N=60	Diagnosis of AD or probable AD Nursing home PLWD capable of standing alone; day care PLWD GDS level 6 or 7	N=60 Age: yes Sex: yes Race: no Education: no	NR	NA	NA	NA	NA	2 weeks each arm	GDS Study to assess safety of stimulation in AD PLWD	NA
Olazaran 2013 ²⁵¹ (23603397) Spain Pilot,	1 treatment of REAC NPO for 2 hours with follow-up at 2 to 3 weeks	14 control PLWD got sham treatment	Nursing home PLWD RCT N=31	Diagnosis of probable AD, not bedridden, some gait dysfunction	N=17 Age: yes Sex: yes Race: no Education: no N of medications: yes	NR	NA	NA	NA	NA	2 to 3 weeks	Axial movement test RSGE-CD Walking test SMMSE Barthel Index NPI	NA

Suemoto 2013 ²⁵² (24262299) Brazil Small sample	Anodal tDCS for 20 minutes; 2 weeks; 6 sessions	20 PLWD got sham tDCS	1 clinical center RCT-Phase II N=40	Diagnosis of possible or probable AD and score of 14 or more on Starkstein Apathy Scale	N=20 Age: yes Sex: yes Education: yes Race: yes	NR	NA	NA	NA	NA	1 week 2 weeks 3 week	Starkstein Apathy Scale Cornell Depression Scale NPI Caregiver Distress ADAS-Cog Digit Cancellation Task Word list learning task Word recognition task	NA
Ahmed 2012 ²⁵³ (21671144) Egypt Small sample	1 daily treatment of rTMS on DLPFC (20 Hz) with follow-up at 1 month and 3 months	15 PLWD got rTMS on DLPFC (1 Hz) 15 PLWD sham procedure	1 clinical center 3 arm RCT N=45	Diagnosis of probable AD; Dementia assessed using MMSE, IADL, GDS	N=15 Age: yes Sex: yes Education: yes Residence: yes Duration of illness: yes	NR	NA	NA	NA	NA	3 months	MMSE IADL GDS	NA
Scherder 2006 ²⁵⁴ (16788393) The Netherlands Small sample	High frequency CES 30 min per day, 5 days per week for 6 weeks	10 control PLWD treated same way without any current	Institutionalized PLWD; treatment and control groups blindly selected N=21	Diagnosis based on NINCDS-ADRDA for probable AD and stage 5 of GDS	N=11 Age: no Sex: yes Race: no Education: yes	NR	NA	NA	NA	NA	6 weeks	BOP BDI	NA
Van Dijk 2005 ²⁵⁵ (15802911) The Netherlands Pilot	Electrical stimulation was given 30 minutes a day for 7 days a week for 6 weeks	Placebo stimulation	1 clinical center and 1 home care center RCT N=68	Diagnosis of AD according to NINCDS/ ADRDA criteria with MMSE of 26 or lower along with sufficient hearing and vision	N=34 Age: yes Sex: yes Education: yes Race: no	NR	NA	NA	NA	NA	6 weeks of treatment for 30 min per day Additional follow-up for 6 weeks	Cognitive measures Behavioral measures	NA
Hozumi 1996 ²⁵⁶ (9003961) Japan Small sample	Transcranial electro-stimulation was given for 20 minutes for 2 weeks	13 PLWD got placebo transcranial electro-stimulation	1 clinical center RCT N=27	Elderly PLWD with sleep-wake disorder and dementia	N=14 Age: yes Sex: yes Education: no Race: no Severity: yes	NR	NA	NA	NA	NA	2 weeks	Motivation Behavior disorder Sleep disorder Intelligence Emotion Language Neurological signs Subjective complaint Activity level	NA

Abbreviations: AD=Alzheimer's Disease; ADAS-Cog=Alzheimer's Disease Assessment Scale-Cognitive Subscale; ADAS-Cog13=Alzheimer's Disease Assessment Scale-13; ADCS-CGI-C= Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change; ACE-III= Addenbrooke's cognitive examination III; ADL=Activities of daily living; AtDCS=Anodal Transcranial Direct Current Stimulation; BEHAVE-AD=Behavioral Pathology in Alzheimer's Disease Rating Scale; BDI=Beck Depression Inventory; BOP=Beoordelingsschaal voor Oudere Patienten; CAF=Clinical Assessment of Fluctuation; CAMCOG=Cambridge Cognitive Examination; CES=Cranial Electrostimulation; CDR-SB=Clinical Dementia Rating Sun of Boxes; CGI=Clinical Global Impression; DLPFC=Dorsolateral Prefrontal Cortex; DSM-IV=Diagnostic and Statistical Manual of Mental Disorders 4th edn; GDS=Global Deterioration Scale; GDS-15=Geriatric Depression Scale 15; Hz=Hertz; IADL=Instrumental Daily Living Activity Scale; IC=Individualized Computerized; MMSE=Mini-mental State Examination; MoCA=Montreal Cognitive Scale; N=number; NA=Not Applicable; NINCDS/ADRDA=National Institute of Neurological and Communicative Disorders and Stroke/ Alzheimer's Disease and Related Disorders Association; NPI=Neuropsychiatric Inventory; NR=Not Reported; ODFAS=One Day Fluctuation Scale; PMID=PubMed Identification Number; PtDCS=Placebo Transcranial Direct Current Stimulation; REAC-NPO=Radio Electric Asymmetric Conveyer- Neuropostural optimization; RSGE-CD=Rating Scale for Gait Evaluation in Cognitive Deterioration; rTMS=repitive transcranial magnetic stimulation; SMMSE=Severe Mini-Mental State Examination; tDCS=Transcranial Direct Current Stimulation; TESS=Treatment Emergent Symptom Scale; TMT A/B= Trail Making Test A and B

Other Interventions for PLWD Well-Being

Evidence Map: Other Interventions for PLWD Well-Being

Table D-51. Characteristics of evidence map studies: other interventions for PLWD well-being

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Liu 2018 ²⁵⁷ (30464426) China Small sample	Passive finger movement exercise program based on active finger exercise along with routine care	Routine care	Residents from Hangzhou Love Heart Older's home in China RCT 54 PLWD	Diagnosis of dementia by medical doctors with PLWD's fingers free from disease restricting hand movement along with visual and oral communication function and no history of taking anti-dementia drugs	N=18 Mean age=80 years Female: 67% majority race: no Education: 39% illiterate	NR	NA	NA	NA	NA	12 weeks	Grip strength (electrical hand muscle dynamometer) Barthel index	NA

Rostad 2018 ²⁵⁸ (29763832) Norway Pilot	Pain assessment twice a week for 12 weeks using Doloplus-2 pain scale	Usual care	Nursing home Cluster RCT 121 PLWD	Dementia diagnosis in medical record with being non-verbal or inability to self-report pain	N=58 Mean age: 84 years Female: 78% % majority race: no Education: no	NR	NA	NR	NA	NA	Twice a week for 12 weeks	Doloplus-2 pain scale ATC N02A ATC N02B ATC M01A OMEQ	NA
Batchelor-Murphy 2017 ²⁵⁹ (28165618) US Pilot	I1: Direct Hand (DH): CG holds the utensil or cup for the resident without PLWD active involvement I2: Over Hand (OH): CG puts his/her hand over the resident's hand to assist the PLWD. I3: Under Hand (UH): CG holds the utensil or cup and places his/her hand under the resident's hand. This technique theoretically allows the resident to feel as though he/she initiated the movement, and is in control	NR	Nursing Homes 3 arm RCT 53 PLWD	Diagnosis of AD or related dementia, BIMS score between 0-12	N=23 Mean age: 84 years Female: 59% % majority race: yes Education: no	NR	NA	NA	NA	NA	2 years	EdFED	NA
Kim 2017 ²⁶⁰ (27594544) Korea Quasi-experimental	Participated in a suicide prevention program twice a week for 5 weeks with one pretest and 2 posttests.	NR	Recruited from geriatric daycare centers Randomized Pretest Posttest design 66 PLWD	Early-stage and questionable dementia selected based on MMSE—KC (score between 16 to 19); score > 5 on both SIS and GDSSF-K scale	N=32 Mean age: 82 years Female: 77% % majority race: no Education: 55% illiterate	NR	NA	NR	NA	NA	5 weeks 7 weeks	SIS Perceived health status Barthel index SS-A scale GDSSF-K	NA
Hobbelen 2012 ²⁶¹ (22185768) Netherlands Pilot Hobbelen 2007 ²⁶¹ (18093298) Netherlands Study Protocol	PLWD, 20 minutes session of PMT by trained physical therapists, 3 times a week for 4 weeks	No PMT (attention control)	Nursing home residents RCT 102 PLWD	Dementia diagnosis based on DSM-IV-TR and have paratonia with a MAS score of at least 2 in at least 1 limb;	N=35 41% Female 82 years Race NR Education NR Use of psychotropic meds: 51%	NR	NA	NA	NA	NA	2 weeks 4 weeks	MAS CGI PCS PACSLAC-D	NA
Coyne 1997 ²⁶² (9281930 2) US Pilot	Using directed verbal prompts (6 sets) and positive reinforcements to complete eating tasks	Unclear	Dementia unit of SNF 3 arm RCT 24 PLWD	Diagnosis of dementia; eat more than half of the meal without assistance in the communal dining hall of the facility;	N=12 Mean age: 82 years Female: 100% % majority race: no Education: no	NR	NA	NA	NA	NA	Day 6 Day 7 Day 12 Day 13	LEI	NA

Abbreviations: AD=Alzheimer's Disease; ATC=Anatomical Therapeutic Chemical; ATC-M01A=Anti-inflammatory and anti-rheumatic products; ATC-N02A=opioids; ATC-N02B=Other analgesics and antipyretics; BIMS=Brief Interview for Mental Status; Char=Characteristics; CG=Caregiver; CGI=Clinical Global Impression; Doloplus-2=Behavioral pain assessment scale for the elderly presenting with verbal communication disorders; EdFED=Edinburgh Feeding Evaluation in Dementia; GDSSF-K=Korean Geriatric Depression Scale Short Form; LEI=Level of eating independence; MAS=Modified Ashworth scale; MMSE-KC=Korean Mini-Mental Status Exam; N=Number; NA=Not Applicable; OMEQ=Oral morphine equivalents; PACSLAC-D=Pain Assessment Checklist for Elderly with Limited Ability to Communicate, Dutch version; PMID=PubMed Identification Number; PMT=Passive Movement Therapy; PSC=Patient specific complaints-modified; PLWD=Persons with Dementia; RCT=Randomized controlled trial; RoB=Risk of Bias; SIS=Suicidal Ideation Scale; SNF=Skilled Nursing Facility; SS-A=Social support appraisal

References for Appendix D

1. Fleiner T, Dauth H, Zijlstra W, et al. A Structured Physical Exercise Program Reduces Professional Caregiver's Burden Caused by Neuropsychiatric Symptoms in Acute Dementia Care: Randomized Controlled Trial Results. *J Alzheimers Dis.* 2020;74(2):429-33. doi: <https://dx.doi.org/10.3233/JAD-191102>. PMID: 32039852.
2. Liu IT, Lee WJ, Lin SY, et al. Therapeutic Effects of Exercise Training on Elderly Patients With Dementia: A Randomized Controlled Trial. *Arch Phys Med Rehabil.* 2020 Feb 19;19:19. doi: <https://dx.doi.org/10.1016/j.apmr.2020.01.012>. PMID: 32084347.
3. Chen LL, Li H, Chen XH, et al. Effects of Hand Exercise on Eating Action in Patients With Alzheimer's Disease. *Am J Alzheimers Dis Other Demen.* 2019 02;34(1):57-62. doi: <https://dx.doi.org/10.1177/1533317518803722>. PMID: 30301358.
4. Huang N, Li W, Rong X, et al. Effects of a Modified Tai Chi Program on Older People with Mild Dementia: A Randomized Controlled Trial. *J Alzheimers Dis.* 2019;72(3):947-56. doi: <https://dx.doi.org/10.3233/JAD-190487>. PMID: 31743998.
5. Karssemeijer EGA, Bossers WJR, Aaronson JA, et al. Exergaming as a Physical Exercise Strategy Reduces Frailty in People With Dementia: A Randomized Controlled Trial. *J Am Med Dir Assoc.* 2019 Aug 10;10:10. doi: <https://dx.doi.org/10.1016/j.jamda.2019.06.026>. PMID: 31409559.
6. Henskens M, Nauta IM, Drost KT, et al. The effects of movement stimulation on activities of daily living performance and quality of life in nursing home residents with dementia: a randomized controlled trial. *Clin Interv Aging.* 2018;13:805-17. doi: <https://dx.doi.org/10.2147/CIA.S160031>. PMID: 29750023.
7. Ho RTH, Fong TCT, Chan WC, et al. Psychophysiological effects of Dance Movement Therapy and physical exercise on older adults with mild dementia: A randomized controlled trial. *Journals of Gerontology Series B Psychological Sciences & Social Sciences.* 2018 Nov 28;28:28. doi: <https://dx.doi.org/10.1093/geronb/gby145>. PMID: 30496547.
8. Lamb SE, Sheehan B, Atherton N, et al. Dementia And Physical Activity (DAPA) trial of moderate to high intensity exercise training for people with dementia: randomized controlled trial. *Bmj.* 2018 05 16;361:k1675. doi: <https://dx.doi.org/10.1136/bmj.k1675>. PMID: 29769247.
9. Chen KM, Kuo CC, Chang YH, et al. Resistance Band Exercises Reduce Depression and Behavioral Problems of Wheelchair-Bound Older Adults with Dementia: A Cluster-Randomized Controlled Trial. *J Am Geriatr Soc.* 2017 Feb;65(2):356-63. doi: <https://dx.doi.org/10.1111/jgs.14526>. PMID: 27879982.
10. Bossers WJ, van der Woude LH, Boersma F, et al. Comparison of Effect of Two Exercise Programs on Activities of Daily Living in Individuals with Dementia: A 9-Week Randomized, Controlled Trial. *J Am Geriatr Soc.* 2016 Jun;64(6):1258-66. doi: <https://dx.doi.org/10.1111/jgs.14160>. PMID: 27321604.
11. Cancela JM, Ayan C, Varela S, et al. Effects of a long-term aerobic exercise intervention on institutionalized patients with dementia. *J Sci Med Sport.* 2016 Apr;19(4):293-8. doi: <https://dx.doi.org/10.1016/j.jsams.2015.05.007>. PMID: 26087884.
12. Hoffmann K, Sobol NA, Frederiksen KS, et al. Moderate-to-High Intensity Physical Exercise in Patients with Alzheimer's Disease: A Randomized Controlled Trial. *J Alzheimers Dis.* 2016;50(2):443-53. doi: <https://dx.doi.org/10.3233/JAD-150817>. PMID: 112439259. Language: English. Entry Date: 20161122. Revision Date: 20180627. Publication Type: journal article.
13. Toots A, Littbrand H, Lindelof N, et al. Effects of a High-Intensity Functional Exercise Program on Dependence in Activities of Daily Living and Balance in Older Adults with Dementia. *J Am Geriatr Soc.* 2016 Jan;64(1):55-64. doi: <https://dx.doi.org/10.1111/jgs.13880>. PMID: 26782852.
14. Telenius EW, Engedal K, Bergland A. Long-term effects of a 12 weeks high-intensity functional exercise program on physical function and mental health in nursing home residents with dementia: a single blinded randomized controlled trial. *BMC geriatr.* 2015 Dec 03;15:158. doi: <https://dx.doi.org/10.1186/s12877-015-0151-8>. PMID: 26630910.
15. Yang SY, Shan CL, Qing H, et al. The Effects of Aerobic Exercise on Cognitive Function of Alzheimer's Disease Patients. *CNS Neurol Disord Drug Targets.* 2015;14(10):1292-7. PMID: 26556080.
16. Pitkala KH, Poysti MM, Laakkonen ML, et al. Effects of the Finnish Alzheimer disease exercise trial (FINALEX): a randomized controlled trial. *JAMA Intern Med.* 2013 May 27;173(10):894-901. doi: <https://dx.doi.org/10.1001/jamainternmed.2013.359>. PMID: 23589097.
17. Hauer K, Schwenk M, Zieschang T, et al. Physical training improves motor performance in people with dementia: a randomized controlled trial. *J Am Geriatr Soc.* 2012 Jan;60(1):8-15. doi: <https://dx.doi.org/10.1111/j.1532-5415.2011.03778.x>. PMID: 22211512.
18. Fan J, Chen K. Using silver yoga exercises to promote physical and mental health of elders with dementia in long-term care facilities. In *International psychogeriatrics / IPA*
19. Roach KE, Tappen RM, Kirk-Sanchez N, et al. A randomized controlled trial of an activity specific exercise program for individuals with Alzheimer disease in long-term care settings. *J Geriatr Phys Ther.* 2011 Apr-Jun;34(2):50-6. doi: <https://dx.doi.org/10.1519/JPT.0b013e31820a0ab9c>. PMID: 21937893.
20. Eggermont LH, Knol DL, Hol EM, et al. Hand motor activity, cognition, mood, and the rest-activity rhythm in dementia: a clustered RCT. *Behav Brain Res.* 2009 Jan 23;196(2):271-8. doi: <https://dx.doi.org/10.1016/j.bbr.2008.09.012>. PMID: 18926856.
21. Rolland Y, Pillard F, Klapouszczak A, et al. Exercise program for nursing home residents with Alzheimer's disease: a 1-year randomized, controlled trial. *J Am Geriatr Soc.* 2007 Feb;55(2):158-65. PMID: 17302650.
22. Petrou S, Khan K, Mistry D, et al. Does Structured Exercise Improve Cognitive Impairment in People with Mild to Moderate Dementia? A Cost-Effectiveness Analysis from a Confirmatory Randomised Controlled Trial: The Dementia and Physical Activity (DAPA) Trial. *Pharmacoeconomics - Open.* 2019 01 Jun;3(2):215-27. doi: <http://dx.doi.org/10.1007/s41669-018-0097-9>. PMID: 627968385.
23. Bossers WJ, van der Woude LH, Boersma F, et al. A 9-Week Aerobic and Strength Training Program Improves Cognitive and Motor Function in Patients with Dementia: A Randomized, Controlled Trial. *Am J Geriatr Psychiatry.* 2015 Nov;23(11):1106-16. doi: <https://dx.doi.org/10.1016/j.jagp.2014.12.191>. PMID: 25648055.
24. Bossers WJ, Scherder EJ, Boersma F, et al. Feasibility of a combined aerobic and strength training program and its effects on cognitive and physical function in institutionalized dementia patients. A pilot study. *PLoS ONE.* 2014;9(5):e97577. doi: <https://dx.doi.org/10.1371/journal.pone.0097577>. PMID: 24844772.
25. Sobol NA, Hoffmann K, Frederiksen KS, et al. Effect of aerobic exercise on physical performance in patients with Alzheimer's disease. *Alzheimer's Dement.* 2016 12;12(12):1207-15. doi: <https://dx.doi.org/10.1016/j.jalz.2016.05.004>. PMID: 27344641.
26. Hoffmann K, Frederiksen KS, Sobol NA, et al. Preserving cognition, quality of life, physical health and functional ability in Alzheimer's disease: the effect of physical exercise (ADEX trial): rationale and design. *Neuroepidemiology.* 2013;41(3-4):198-207. doi: <https://dx.doi.org/10.1159/000354632>. PMID: 24135720.
27. Sondell A, Rosendahl E, Gustafson Y, et al. The Applicability of a High-Intensity Functional Exercise Program Among Older People With Dementia Living in Nursing Homes. *J Geriatr Phys Ther.* 2018 May 25;25:25. doi: <https://dx.doi.org/10.1519/JPT.000000000000199>. PMID: 29851748.
28. Toots A, Wiklund R, Littbrand H, et al. The Effects of Exercise on Falls in Older People With Dementia Living in Nursing Homes: A Randomized Controlled Trial. *J Am Med Dir Assoc.* 2018 Nov 28;28:28. doi: <https://dx.doi.org/10.1016/j.jamda.2018.10.009>. PMID: 30503589.
29. Toots A, Littbrand H, Bostrom G, et al. Effects of Exercise on Cognitive Function in Older People with Dementia: A Randomized Controlled Trial. *J Alzheimers Dis.* 2017;60(1):323-32. doi: <https://dx.doi.org/10.3233/JAD-170014>. PMID: 28800328.
30. Bostrom G, Conradsson M, Hornsten C, et al. Effects of a high-intensity functional exercise program on depressive symptoms among people with dementia in residential care: a randomized controlled trial. *Int J Geriatr Psychiatry.* 2016 Aug;31(8):868-78. doi: <https://dx.doi.org/10.1002/gps.4401>. PMID: 26644304.
31. Roitto HM, Kautiainen H, Ohman H, et al. Relationship of Neuropsychiatric Symptoms with Falls in Alzheimer's Disease - Does Exercise Modify the Risk? *J Am Geriatr Soc.* 2018 Dec;66(12):2377-81. doi: <https://dx.doi.org/10.1111/jgs.15614>. PMID: 30320427.
32. Ohman H, Savikko N, Strandberg TE, et al. Effects of Exercise on Cognition: The Finnish Alzheimer Disease Exercise Trial: A Randomized, Controlled Trial. *J Am Geriatr Soc.* 2016 Apr;64(4):731-8. doi: <https://dx.doi.org/10.1111/jgs.14059>. PMID: 27037872.
33. Ohman H, Savikko N, Strandberg T, et al. Effects of Exercise on Functional Performance and Fall Rate in Subjects with Mild or Advanced Alzheimer's Disease: Secondary Analyses of a Randomized Controlled Study. *Dement Geriatr Cogn Disord.* 2016;41(3-4):233-41. doi: <https://dx.doi.org/10.1159/000445712>. PMID: 27160164.
34. Perttala N, Ohman H, Strandberg T, et al. Severity of frailty and the outcome of exercise intervention among participants with Alzheimer disease: a sub-group analysis of a randomized controlled trial. In *European geriatric medicine*
35. Pitkala KH, Raivio MM, Laakkonen ML, et al. Exercise rehabilitation on home-dwelling patients with Alzheimer disease: A randomized, controlled trial. Baseline findings and feasibility. *European Geriatric Medicine.* 2011 December;2(6):338-43. doi: <http://dx.doi.org/10.1016/j.eurger.2011.07.012>. PMID: 51607662.
36. Pomiersky R, Abel B, Werner C, et al. Increasing Physical Activity in Persons With Dementia: A Randomized Controlled Trial. *J Aging Phys Activity.* 2020 Jan 01:1-10. doi: <https://dx.doi.org/10.1123/japa.2019-0183>. PMID: 31896078.
37. Enette L, Vogel T, Merle S, et al. Effect of 9 weeks continuous vs. interval aerobic training on plasma BDNF levels, aerobic fitness, cognitive capacity and quality of life among seniors with mild to moderate Alzheimer's disease: a randomized controlled trial. *Eur.* 2020;17:2. doi: <https://dx.doi.org/10.1186/s11556-019-0234-1>. PMID: 31921371.
38. Lee SM, Joung J, Shin SH. Effects of Fumanet exercise on Korean older adults with mild dementia. *Jpn J Nurs Sci.* 2019 Jul 25:e12286. doi: <https://dx.doi.org/10.1111/jjns.12286>. PMID: 31347294.
39. Sanders LMJ, Hortobagyi T, Karssemeijer EGA, et al. Effects of low- and high-intensity physical exercise on physical and cognitive function in older persons with dementia: a randomized controlled trial. *Alzheimers Res Ther.* 2020 Mar 19;12(1):28. doi: <https://dx.doi.org/10.1186/s13195-020-00597-3>. PMID: 32192537.
40. Brett L, Stapley P, Meedya S, et al. Effect of physical exercise on physical performance and fall incidents of individuals living with dementia in nursing homes: a randomized controlled trial. *Physiother.* 2019 Mar 26:1-14. doi: <https://dx.doi.org/10.1080/09593985.2019.1594470>. PMID: 30912690.
41. Nyman SR, Ingram W, Sanders J, et al. Randomised Controlled Trial Of The Effect Of Tai Chi On Postural Balance Of People With Dementia. *Clin Interv Aging.* 2019;14:2017-29. doi: <https://dx.doi.org/10.2147/CIA.S228931>. PMID: 31819385.
42. Todri J, Lena O, Martinez Gil JL. A single blind randomized controlled trial of global postural re-education: Cognitive effects on Alzheimer disease patients. *European Journal of Psychiatry.* 2019 April - June;33(2):83-90.
43. Liu JYW, Kwan RYC, Lai CK, et al. A simplified 10-step Tai-chi programme to enable people with dementia to improve their motor performance: a feasibility study. *Clin Rehabil.* 2018 Dec;32(12):1609-23. doi: <https://dx.doi.org/10.1177/0269215518786530>. PMID: 29969916.
44. Dawson N, Judge KS, Gerhart H. Improved Functional Performance in Individuals With Dementia After a Moderate-Intensity Home-Based Exercise Program: A Randomized Controlled Trial. *J Geriatr Phys Ther.* 2017 Mar 01;01:01. doi: <https://dx.doi.org/10.1519/JPT.000000000000128>. PMID: 28252473.
45. de Souto Barreto P, Cesari M, Denormandie P, et al. Exercise or Social Intervention for Nursing Home Residents with Dementia: A Pilot Randomized, Controlled Trial. *J Am Geriatr Soc.* 2017 Sep;65(9):E123-E9. doi: <https://dx.doi.org/10.1111/jgs.14947>. PMID: 28542742.
46. Hamilton G, Gallagher M, Pierson K, et al. Palliative care for dementia: a randomized controlled study of a replicable and financially viable model for organizations providing caregiver support. In *Journal of pain and symptom management*
47. Henwood T, Neville C, Baguley C, et al. Aquatic exercise for residential aged care adults with dementia: benefits and barriers to participation. *Int Psychogeriatr.* 2017;29(9):1439-49. doi: [10.1017/S104161021700028X](https://dx.doi.org/10.1017/S104161021700028X). PMID: CN-01403352.
48. Lam F, Liao L, Kwok T, et al. Effects of adding whole-body vibration to routine day activity program on physical functioning in elderly with mild or moderate dementia: a randomized controlled trial. In *Int J Geriatr Psychiatry*
49. Morris JK, Vidoni ED, Johnson DK, et al. Aerobic exercise for Alzheimer's disease: A randomized controlled pilot trial. *PLoS ONE.* 2017;12(2):e0170547. doi: <https://dx.doi.org/10.1371/journal.pone.0170547>. PMID: 28187125.
50. Padala KP, Padala PR, Lensing SY, et al. Home-Based Exercise Program Improves Balance and Fear of Falling in Community-Dwelling Older Adults with Mild Alzheimer's Disease: A Pilot Study. *J Alzheimers Dis.* 2017;59(2):565-74. doi: <https://dx.doi.org/10.3233/JAD-170120>. PMID: 28655135.
51. Burge E, Berchtold A, Maupetit C, et al. Does physical exercise improve adl capacities in people over 65 years with moderate or severe dementia hospitalized in an acute psychiatric setting? A multisite randomized clinical trial. *Int Psychogeriatr.* 2016. PMID: CN-01308751.
52. Venturelli M, Sollima A, Ce E, et al. Effectiveness of Exercise- and Cognitive-Based Treatments on Salivary Cortisol Levels and Sundowning Syndrome Symptoms in Patients with Alzheimer's Disease. *J Alzheimers Dis.* 2016;53(4):1631-40. doi: <http://dx.doi.org/10.3233/JAD-160392>. PMID: 611692625.
53. Holthoff VA, Marschner K, Scharf M, et al. Effects of physical activity training in patients with Alzheimer's dementia: results of a pilot RCT study. *PLoS ONE.* 2015;10(4):e0121478. doi: <https://dx.doi.org/10.1371/journal.pone.0121478>. PMID: 25884637.
54. Yu F, Thomas W, Nelson NW, et al. Impact of 6-month aerobic exercise on Alzheimer's symptoms. *J Appl Gerontol.* 2015 Jun;34(4):484-500. doi: <https://dx.doi.org/10.1177/0733464813512895>. PMID: 24652914.
55. Suttanon P, Hill KD, Said CM, et al. Feasibility, safety and preliminary evidence of the effectiveness of a home-based exercise programme for older people with Alzheimer's disease: a pilot randomized controlled trial. *Clin Rehabil.* 2013 May;27(5):427-38. doi: <https://dx.doi.org/10.1177/0269215512460877>. PMID: 23117349.
56. Canonici P, Andrade LP, Gobbi S, et al. Functional dependence and caregiver burden in Alzheimer's disease: a controlled trial on the benefits of motor intervention. *Psychogeriatrics.* 2012;12(3):186-92. doi: [10.1111/j.1479-8301.2012.00407.x](https://dx.doi.org/10.1111/j.1479-8301.2012.00407.x). PMID: CN-00863385.
57. Nascimento C, Teixeira C, Gobbi L, et al. A controlled clinical trial on the effects of exercise on neuropsychiatric disorders and instrumental activities in women with Alzheimer's disease. *Revista brasileira de fisioterapia (sao carlos (sao paulo, brazil)).* 2012;16(3):197-204. PMID: CN-00863837.
58. Vreugdenhil A, Cannell J, Davies A, et al. A community-based exercise programme to improve functional ability in people with Alzheimer's disease: a randomized controlled trial. *Scand J Caring Sci.* 2012 Mar;26(1):12-9. doi: <https://dx.doi.org/10.1111/j.1471-6712.2011.00895.x>. PMID: 21564154.
59. Stella F, Canonici A, Gobbi S, et al. Attenuation of neuropsychiatric symptoms and caregiver burden in Alzheimer's disease by motor intervention: a controlled trial. *Clinics (sao paulo, brazil).* 2011;66(8):1353-60. PMID: CN-00860745.
60. Venturelli M, Scarsini R, Schena F. Six-month walking program changes cognitive and ADL performance in patients with Alzheimer. *Am J Alzheimers Dis Other Demen.* 2011 Aug;26(5):381-8. doi: <https://dx.doi.org/10.1177/1533317511418956>. PMID: 21852281.
61. Steinberg M, Leoutsakos JM, Podewils LJ, et al. Evaluation of a home-based exercise program in the treatment of Alzheimer's disease: the Maximizing Independence in Dementia (MIND) study. *Int J Geriatr Psychiatry.* 2009 Jul;24(7):680-5. doi: <https://dx.doi.org/10.1002/gps.2175>. PMID: 19089875.
62. Kwak YS, Um SY, Son TG, et al. Effect of regular exercise on senile dementia patients. *Int J Sports Med.* 2008 Jun;29(6):471-4. PMID: 18050054.

63. Williams CL, Tappen RM. Exercise training for depressed older adults with Alzheimer's disease. *Aging Ment Health*. 2008 Jan;12(1):72-80. doi: <https://dx.doi.org/10.1080/13607860701529932>. PMID: 18297481.
64. Stevens J, Killeen M. A randomised controlled trial testing the impact of exercise on cognitive symptoms and disability of residents with dementia. *Contemp Nurse*. 2006 Feb-Mar;21(1):32-40. PMID: 16594879.
65. Van de Winckel A, Feys H, De Weerd W, et al. Cognitive and behavioural effects of music-based exercises in patients with dementia. *Clin Rehabil*. 2004 May;18(3):253-60. PMID: 15137556.
66. Toulotte C, Fabre C, Dangremont B, et al. Effects of physical training on the physical capacity of frail, demented patients with a history of falling: a randomised controlled trial. *Age Ageing*. 2003 Jan;32(1):67-73. PMID: 12540351.
67. Ugur HG, Orak OS, Aktas YY, et al. Effects of music therapy on the care burden of in-home caregivers and physiological parameters of their in-home dementia patients: a randomized controlled trial. *Complementary medicine research*. 2019;26(1):22-30. PMID: CN-01911571.
68. Cheung DSK, Lai CKY, Wong FKY, et al. The effects of the music-with-movement intervention on the cognitive functions of people with moderate dementia: a randomized controlled trial. *Aging Ment Health*. 2018 Mar;22(3):306-15. doi: <https://dx.doi.org/10.1080/13607863.2016.1251571>. PMID: 27819483.
69. Ho RT, Fong TC, Sing CY, et al. Managing behavioral and psychological symptoms in Chinese elderly with dementia via group-based music intervention: A cluster randomized controlled trial. *Dementia*. 2018 Jan 01;1471301218760023. doi: <https://dx.doi.org/10.1177/1471301218760023>. PMID: 29468887.
70. Kwak J, Anderson K, O'Connell Valuch K. Findings From a Prospective Randomized Controlled Trial of an Individualized Music Listening Program for Persons With Dementia. *J Appl Gerontol*. 2018 Jun 01;733464818778991. doi: <https://dx.doi.org/10.1177/0733464818778991>. PMID: 29871544.
71. Pongan E, Tillmann B, Leveque Y, et al. Can Musical or Painting Interventions Improve Chronic Pain, Mood, Quality of Life, and Cognition in Patients with Mild Alzheimer's Disease? Evidence from a Randomized Controlled Trial. *J Alzheimers Dis*. 2017;60(2):663-77. doi: <https://dx.doi.org/10.3233/JAD-170410>. PMID: 28922159.
72. Wang SC, Yu CL, Chang SH. Effect of music care on depression and behavioral problems in elderly people with dementia in Taiwan: a quasi-experimental, longitudinal study. *Aging Ment Health*. 2017 02;21(2):156-62. doi: <https://dx.doi.org/10.1080/13607863.2015.1093602>. PMID: 26443002.
73. Sarkamo T, Tervaniemi M, Laitinen S, et al. Cognitive, emotional, and social benefits of regular musical activities in early dementia: randomized controlled study. *Gerontologist*. 2014 Aug;54(4):634-50. doi: <https://dx.doi.org/10.1093/geront/gnt100>. PMID: 24009169.
74. Vink AC, Zuidersma M, Boersma F, et al. The effect of music therapy compared with general recreational activities in reducing agitation in people with dementia: a randomised controlled trial. *Int J Geriatr Psychiatry*. 2013 Oct;28(10):1031-8. doi: <https://dx.doi.org/10.1002/gps.3924>. PMID: 23280604.
75. Sung HC, Lee WL, Li TL, et al. A group music intervention using percussion instruments with familiar music to reduce anxiety and agitation of institutionalized older adults with dementia. *Int J Geriatr Psychiatry*. 2012 Jun;27(6):621-7. doi: <https://dx.doi.org/10.1002/gps.2761>. PMID: 21823174.
76. Lin Y, Chu H, Yang CY, et al. Effectiveness of group music intervention against agitated behavior in elderly persons with dementia. *Int J Geriatr Psychiatry*. 2011 Jul;26(7):670-8. doi: <https://dx.doi.org/10.1002/gps.2580>. PMID: 20672256.
77. Chu H, Yang CY, Lin Y, et al. The impact of group music therapy on depression and cognition in elderly persons with dementia: a randomized controlled study. *Biol Res Nurs*. 2014 Apr;16(2):209-17. doi: <https://dx.doi.org/10.1177/1099800413485410>. PMID: 23639952.
78. Raglio A, Bellelli G, Traficante D, et al. Efficacy of music therapy in the treatment of behavioral and psychiatric symptoms of dementia. *Alzheimer Dis Assoc Disord*. 2008 Apr-Jun;22(2):158-62. doi: <https://dx.doi.org/10.1097/WAD.0b013e3181630b6f>. PMID: 18525288.
79. Ledger A, Baker F. An investigation of long-term effects of group music therapy on agitation levels of people with Alzheimer's disease. *Aging Ment Health*. 2007;11(3):330-8. PMID: CN-00738375.
80. Cheung DSK, Lai CKY, Wong FKY, et al. Is music-with-movement intervention better than music listening and social activities in alleviating agitation of people with moderate dementia? A randomized controlled trial. *Dementia*. 2018 Sep 20;1471301218800195. doi: <https://dx.doi.org/10.1177/1471301218800195>. PMID: 30235949.
81. Sarkamo T, Laitinen S, Numminen A, et al. Clinical and Demographic Factors Associated with the Cognitive and Emotional Efficacy of Regular Musical Activities in Dementia. *J Alzheimers Dis*. 2016;49(3):767-81. doi: <https://dx.doi.org/10.3233/JAD-150453>. PMID: 26519435.
82. Weise L, Töpfer NF, Deux J, et al. Feasibility and effects of individualized recorded music for people with dementia: A pilot RCT study. *Nordic Journal of Music Therapy*. 2020;29(1):39-56. doi: 10.1080/08098131.2019.1661507. PMID: 141134381. Language: English. Entry Date: 20200117. Revision Date: 20200117. Publication Type: Article.
83. Ihara ES, Tompkins CJ, Inoue M, et al. Results from a person-centered music intervention for individuals living with dementia. *Geriatr Gerontol Int*. 2019 Jan;19(1):30-4. doi: <https://dx.doi.org/10.1111/ggi.13563>. PMID: 30460747.
84. Park J, Tolea MI, Sherman D, et al. Feasibility of Conducting Nonpharmacological Interventions to Manage Dementia Symptoms in Community-Dwelling Older Adults: A Cluster Randomized Controlled Trial. *Am J Alzheimers Dis Other Dement*. 2019 Sep 18;1533317519872635. doi: <https://dx.doi.org/10.1177/1533317519872635>. PMID: 31533443.
85. Giovagnoli AR, Manfredi V, Schifano L, et al. Combining drug and music therapy in patients with moderate Alzheimer's disease: a randomized study. *Neurol Sci*. 2018 Mar 17;17:17. doi: <https://dx.doi.org/10.1007/s10072-018-3316-3>. PMID: 29550981.
86. Narme P, Clement S, Ehrle N, et al. Efficacy of musical interventions in dementia: evidence from a randomized controlled trial. *J Alzheimers Dis*. 2014;38(2):359-69. doi: <https://dx.doi.org/10.3233/JAD-130893>. PMID: 23969994.
87. Ridder HM, Stige B, Qvale LG, et al. Individual music therapy for agitation in dementia: an exploratory randomized controlled trial. *Aging Ment Health*. 2013;17(6):667-78. doi: <https://dx.doi.org/10.1080/13607863.2013.790926>. PMID: 23621805.
88. Sakamoto M, Ando H, Tsutou A. Comparing the effects of different individualized music interventions for elderly individuals with severe dementia. *Int Psychogeriatr*. 2013 May;25(5):775-84. doi: <https://dx.doi.org/10.1017/S1041610212002256>. PMID: 23298693.
89. Ceccato E, Vignato G, Bonetto C, et al. STAM protocol in dementia: a multicenter, single-blind, randomized, and controlled trial. *Am J Alzheimers Dis Other Dement*. 2012 Aug;27(5):301-10. doi: <https://dx.doi.org/10.1177/1533317512452038>. PMID: 22815078.
90. McHugh L, Gardstrom S, Hiller J, et al. The effect of pre-meal, vocal re-creative music therapy on nutritional intake of residents with Alzheimer's disease and related dementias: a pilot study. *Music ther perspect*. 2012;30(1):32-42. PMID: CN-01140174.
91. Chang F, Huang H, Lin K, et al. The effect of a music programme during lunchtime on the problem behaviour of the older residents with dementia at an institution in Taiwan. *J Clin Nurs*. 2010;19(7-8):939-48. doi: 10.1111/j.1365-2702.2009.02801.x. PMID: CN-00752730.
92. Cooke M, Moyle W, Shum D, et al. A randomized controlled trial exploring the effect of music on quality of life and depression in older people with dementia. *J Health Psychol*. 2010 Jul;15(5):765-76. doi: <https://dx.doi.org/10.1177/1359105310368188>. PMID: 20603300.
93. Cooke ML, Moyle W, Shum DH, et al. A randomized controlled trial exploring the effect of music on agitated behaviours and anxiety in older people with dementia. *Aging Ment Health*. 2010 Nov;14(8):905-16. doi: <https://dx.doi.org/10.1080/13607861003713190>. PMID: 20635236.
94. Han P, Kwan M, Chen D, et al. A controlled naturalistic study on a weekly music therapy and activity program on disruptive and depressive behaviors in dementia. *Dementia and geriatric cognitive disorders*. 2010;30(6):540-6. PMID: CN-00856782.
95. Sung HC, Chang AM, Lee WL. A preferred music listening intervention to reduce anxiety in older adults with dementia in nursing homes. *J Clin Nurs*. 2010 Apr;19(7-8):1056-64. doi: <https://dx.doi.org/10.1111/j.1365-2702.2009.03016.x>. PMID: 20492050.
96. Guetin S, Portet F, Picot MC, et al. Effect of music therapy on anxiety and depression in patients with Alzheimer's type dementia: randomised, controlled study. *Dement Geriatr Cogn Disord*. 2009;28(1):36-46. doi: <https://dx.doi.org/10.1159/000229024>. PMID: 19628939.
97. Choi A, Lee M, Cheong K, et al. Effects of group music intervention on behavioral and psychological symptoms in patients with dementia: a pilot-controlled trial. *Int J Neurosci*. 2009;119(4):471-81. doi: 10.1080/00207450802328136. PMID: CN-00688678.
98. Holmes C, Knights A, Dean C, et al. Keep music live: music and the alleviation of apathy in dementia subjects. *Int Psychogeriatr*. 2006 Dec;18(4):623-30. PMID: 16805928.
99. Sung HC, Chang SM, Lee WL, et al. The effects of group music with movement intervention on agitated behaviours of institutionalized elders with dementia in Taiwan. *Complement Ther Med*. 2006 Jun;14(2):113-9. PMID: 16765849.
100. Svansdotir HB, Snaedal J. Music therapy in moderate and severe dementia of Alzheimer's type: a case-control study. *Int Psychogeriatr*. 2006 Dec;18(4):613-21. PMID: 16618375.
101. Gerdner LA. Effects of individualized versus classical "relaxation" music on the frequency of agitation in elderly persons with Alzheimer's disease and related disorders. *Int Psychogeriatr*. 2000 Mar;12(1):49-65. PMID: 10798453.
102. Groene RW. Effectiveness of music therapy 1:1 intervention with individuals having Senile Dementia of the Alzheimer's Type. *J Music Ther*. 1993 Fal;30(3):138-57. doi: <http://dx.doi.org/10.1093/jmt/30.3.138>. PMID: 1994-10861-001.
103. Lord TR, Garner JE. Effects of music on Alzheimer patients. *Percept Mot Skills*. 1993 Apr;76(2):451-5. PMID: 8483655.
104. Li M, Lyu JH, Zhang Y, et al. Efficacy of Group Reminiscence Therapy on Cognition, Depression, Neuropsychiatric Symptoms, and Activities of Daily Living for Patients With Alzheimer Disease. *J Geriatr Psychiatry Neurol*. 2019 Oct 23;891988719882099. doi: <https://dx.doi.org/10.1177/0891988719882099>. PMID: 31645180.
105. Lok N, Bademli K, Selcuk-Tosun A. The effect of reminiscence therapy on cognitive functions, depression, and quality of life in Alzheimer patients: Randomized controlled trial. *Int J Geriatr Psychiatry*. 2019 Jan;34(1):47-53. doi: <https://dx.doi.org/10.1002/gps.4980>. PMID: 30246408.
106. Lin H-C, Yang Y-P, Cheng W-Y, et al. Distinctive effects between cognitive stimulation and reminiscence therapy on cognitive function and quality of life for different types of behavioural problems in dementia. In *Scand J Caring Sci*
107. Amieva H, Robert PH, Grandoulier AS, et al. Group and individual cognitive therapies in Alzheimer's disease: the ETNA3 randomized trial. *Int Psychogeriatr*. 2016 May;28(5):707-17. doi: <https://dx.doi.org/10.1017/S1041610215001830>. PMID: 26572551.
108. Duru Asiret G, Kapucu S. The Effect of Reminiscence Therapy on Cognition, Depression, and Activities of Daily Living for Patients With Alzheimer Disease. *J Geriatr Psychiatry Neurol*. 2016 Jan;29(1):31-7. doi: <https://dx.doi.org/10.1177/0891988715598233>. PMID: 26251112.
109. Van Bogaert P, Tolson D, Eerlingen R, et al. SolCos model-based individual reminiscence for older adults with mild to moderate dementia in nursing homes: a randomized controlled intervention study. *J Psychiatr Ment Health Nurs*. 2016 Nov;23(9-10):568-75. doi: <https://dx.doi.org/10.1111/jpm.12336>. PMID: 27511740.
110. Wu LF, Koo M. Randomized controlled trial of a six-week spiritual reminiscence intervention on hope, life satisfaction, and spiritual well-being in elderly with mild and moderate dementia. *Int J Geriatr Psychiatry*. 2016 Feb;31(2):120-7. doi: <https://dx.doi.org/10.1002/gps.4300>. PMID: 25965388.
111. O'Shea E, Devane D, Cooney A, et al. The impact of reminiscence on the quality of life of residents with dementia in long-stay care. *Int J Geriatr Psychiatry*. 2014 Oct;29(10):1062-70. doi: <https://dx.doi.org/10.1002/gps.4099>. PMID: 24633858.
112. Serrani Azcurra DJ. A reminiscence program intervention to improve the quality of life of long-term care residents with Alzheimer's disease: a randomized controlled trial. *Rev Bras Psiquiatr*. 2012 Dec;34(4):422-33. PMID: 23429813.
113. Woods RT, Bruce E, Edwards RT, et al. REMCARE: reminiscence groups for people with dementia and their family caregivers - effectiveness and cost-effectiveness pragmatic multicentre randomised trial. *Health Technol Assess*. 2012;16(48):v-xv, 1-116. doi: <https://dx.doi.org/10.3310/hta16480>. PMID: 23211271.
114. Hsieh C-J, Chang C, Su S-F, et al. Reminiscence group therapy on depression and apathy in nursing home residents with mild-to-moderate dementia. In *Journal of experimental and clinical medicine*
115. Wang JJ. Group reminiscence therapy for cognitive and affective function of demented elderly in Taiwan. *Int J Geriatr Psychiatry*. 2007 Dec;22(12):1235-40. PMID: 17503545.
116. Lai CK, Chi I, Kayser-Jones J. A randomized controlled trial of a specific reminiscence approach to promote the well-being of nursing home residents with dementia. *Int Psychogeriatr*. 2004 Mar;16(1):33-49. PMID: 15190995.
117. Camberg L, Woods P, Ooi WL, et al. Evaluation of Simulated Presence: a personalized approach to enhance well-being in persons with Alzheimer's disease. *J Am Geriatr Soc*. 1999 Apr;47(4):446-52. PMID: 10203120.
118. Woods RT, Bruce E, Edwards RT, et al. Reminiscence groups for people with dementia and their family carers: pragmatic eight-centre randomised trial of joint reminiscence and maintenance versus usual treatment: a protocol. *Trials*. 2009 Jul 30;10:64. doi: <https://dx.doi.org/10.1186/1745-6215-10-64>. PMID: 19642992.
119. Woods RT, Orrell M, Bruce E, et al. REMCARE: Pragmatic Multi-Centre Randomised Trial of Reminiscence Groups for People with Dementia and their Family Carers: Effectiveness and Economic Analysis. *PLoS ONE*. 2016;11(4):e0152843. doi: <https://dx.doi.org/10.1371/journal.pone.0152843>. PMID: 27093052.
120. Wang J, Yen M, OuYang W. Group reminiscence intervention in Taiwanese elders with dementia. In *Archives of gerontology and geriatrics*
121. Ching-Teng Y, Ya-Ping Y, Chia-Ju L, et al. Effect of group reminiscence therapy on depression and perceived meaning of life of veterans diagnosed with dementia at veteran homes. *Soc Work Health Care*. 2020;59(2):75-90. doi: 10.1080/00981389.2019.1710320. PMID: 141877576. Language: English. Entry Date: 20200227. Revision Date: 20200319. Publication Type: Article.
122. Manav AI, Simsek N. The effect of reminiscence therapy with internet-based videos on cognitive status and apathy of older people with mild dementia. *Journal of geriatric psychiatry and neurology*. 2019;32(2):104-13. PMID: CN-01707450.
123. Martinez N, Stutzman SE, Olson DM. Electronic interventions aimed at increasing self-worth in mild dementia may not be feasible. *J Am Assoc Nurse Pract*. 2019 Aug 22;22:22. doi: <https://dx.doi.org/10.1097/JXX.0000000000000280>. PMID: 31453826.
124. Laird EA, Ryan A, McCauley C, et al. Using Mobile Technology to Provide Personalized Reminiscence for People Living With Dementia and Their Carers: Appraisal of Outcomes From a Quasi-Experimental Study. *JMIR Ment Health*. 2018 Sep 11;5(3):e57. doi: <https://dx.doi.org/10.2196/mental.9684>. PMID: 30206053.
125. Gonzalez J, Mayordomo T, Torres M, et al. Reminiscence and dementia: a therapeutic intervention. *Int Psychogeriatr*. 2015 Oct;27(10):1731-7. doi: <https://dx.doi.org/10.1017/S1041610215000344>. PMID: 25765779.
126. Lalanne J, Gallarda T, Piolino P. "The Castle of Remembrance": New insights from a cognitive training programme for autobiographical memory in Alzheimer's disease. *Neuropsychol*. 2015;25(2):254-82. doi: <https://dx.doi.org/10.1080/09602011.2014.949276>. PMID: 25122521.
127. Nakamae T, Yotsumoto K, Tatsumi E, et al. Effects of productive activities with reminiscence in occupational therapy for people with dementia: A pilot randomized controlled study. *Hong Kong Journal of Occupational Therapy*. 2014 June;24(1):13-9. doi: <http://dx.doi.org/10.1016/j.hkjoct.2014.01.003>. PMID: 53104158.
128. Subramaniam P, Woods B, Whitaker C. Life review and life story books for people with mild to moderate dementia: a randomised controlled trial. *Aging Ment Health*. 2014;18(3):363-75. doi: <https://dx.doi.org/10.1080/13607863.2013.837144>. PMID: 24063317.
129. Van Bogaert P, Van Grinsven R, Tolson D, et al. Effects of SolCos model-based individual reminiscence on older adults with mild to moderate dementia due to Alzheimer disease: a pilot study. *J Am Med Dir Assoc*. 2013 Jul;14(7):528.e9-13. doi: <https://dx.doi.org/10.1016/j.jamda.2013.01.020>. PMID: 23583001.
130. Baillon S, Van Diepen E, Prettymann R, et al. A comparison of the effects of Snoezelen and reminiscence therapy on the agitated behaviour of patients with dementia. *Int J Geriatr Psychiatry*. 2004 Nov;19(11):1047-52. PMID: 15481068.
131. Politis AM, Vozzella S, Mayer LS, et al. A randomized, controlled, clinical trial of activity therapy for apathy in patients with dementia residing in long-term care. *Int J Geriatr Psychiatry*. 2004 Nov;19(11):1087-94. PMID: 15481065.
132. Chen Z, Yu H. Effect of intervention guided by montessori method on improving feeding capacity of patients with dementia. *Int J Clin Exp Med*. 2020;13(2):1148-55.
133. Clare L, Kudlicka A, Oyebode JR, et al. Individual goal-oriented cognitive rehabilitation to improve everyday functioning for people with early-stage dementia: A multicentre randomised controlled trial (the GREAT trial). *Int J Geriatr Psychiatry*. 2019 May;34(5):709-21. doi: <http://dx.doi.org/10.1002/gps.5076>. PMID: 626609676.

134. Voigt-Radloff S, de Werd MM, Leonhart R, et al. Structured relearning of activities of daily living in dementia: the randomized controlled REDALI-DEM trial on errorless learning. *Alzheimers Res Ther*. 2017 Mar 23;9(1):22. doi: <https://dx.doi.org/10.1186/s13195-017-0247-9>. PMID: 28335810.
135. Kumar P, Tiwari SC, Goel A, et al. Novel occupational therapy interventions may improve quality of life in older adults with dementia. *Int Arch Med*. 2014;7:26. doi: <https://dx.doi.org/10.1186/1755-7682-7-26>. PMID: 24982692.
136. Wu HS, Lin LC, Wu SC, et al. The effectiveness of spaced retrieval combined with Montessori-based activities in improving the eating ability of residents with dementia. *J Adv Nurs*. 2014 Aug;70(8):1891-901. doi: <https://dx.doi.org/10.1111/jan.12352>. PMID: 24444172.
137. Clare L, Kudlicka A, Oyeboe JR, et al. Goal-oriented cognitive rehabilitation for early-stage Alzheimer's and related dementias: the GREAT RCT. Health technology assessment. 2019;23(10):1-+. PMID: CN-01911576.
138. Voigt-Radloff S, Leonhart R, Rikkert MO, et al. Study protocol of the multi-site randomised controlled REDALI-DEM trial--the effects of structured relearning methods on daily living task performance of persons with dementia. *BMC geriatr*. 2011 Aug 18;11:44. doi: <https://dx.doi.org/10.1186/1471-2318-11-44>. PMID: 21851594.
139. Mbakile-Mahlanza L, van der Ploeg ES, Busija L, et al. A cluster-randomized crossover trial of Montessori activities delivered by family carers to nursing home residents with behavioral and psychological symptoms of dementia. *Int Psychogeriatr*. 2020 Mar;32(3):347-58. doi: <https://dx.doi.org/10.1017/S1041610219001819>. PMID: 31762434.
140. Yuen I, Kwok T. Effectiveness of DementiaAbility Methods: The Montessori Way on agitation in long-term care home residents with dementia in Hong Kong. *Int J Geriatr Psychiatry*. 2019;34(9):1352-8. doi: <http://dx.doi.org/10.1002/gps.5063>. PMID: 626609449.
141. Silva AR, Pinho MS, Macedo L, et al. It is not only memory: effects of sensecam on improving well-being in patients with mild Alzheimer disease. *Int Psychogeriatr*. 2017 May;29(5):741-54. doi: <https://dx.doi.org/10.1017/S104161021600243X>. PMID: 28124633.
142. Silva AR, Pinho MS, Macedo L, et al. The cognitive effects of wearable cameras in mild Alzheimer disease-An experimental study. *Curr Alzheimer Res*. 2017;14(12):1270-82. doi: <http://dx.doi.org/10.2174/1567205014666170531083015>. PMID: 2018-01500-004.
143. Tanaka S, Honda S, Nakano H, et al. Comparison between group and personal rehabilitation for dementia in a geriatric health service facility: single-blinded randomized controlled study. *Psychogeriatrics*. 2017 May;17(3):177-85. doi: <https://dx.doi.org/10.1111/psyg.12212>. PMID: 27612310.
144. Tsuchiya K, Yamaguchi T, Fujita T, et al. A Quasi-Randomized Controlled Trial of Brain-Activating Rehabilitation in an Acute Hospital. *Am J Alzheimers Dis Other Demen*. 2016 Dec;31(8):612-7. PMID: 27303064.
145. Kim S. Cognitive rehabilitation for elderly people with early-stage Alzheimer's disease. *J Phys Ther Sci*. 2015 Feb;27(2):543-6. doi: <https://dx.doi.org/10.1589/jpts.27.543>. PMID: 25729212.
146. Thivierge S, Jean L, Simard M. A randomized cross-over controlled study on cognitive rehabilitation of instrumental activities of daily living in Alzheimer disease. *Am J Geriatr Psychiatry*. 2014 Nov;22(11):1188-99. doi: <https://dx.doi.org/10.1016/j.jagp.2013.03.008>. PMID: 23871120.
147. Brunelle-Hamann L, Thivierge S, Simard M. Impact of a cognitive rehabilitation intervention on neuropsychiatric symptoms in mild to moderate Alzheimer's disease. *Neuropsychol*. 2015;25(5):677-707. doi: <https://dx.doi.org/10.1080/09602011.2014.964731>. PMID: 25312605.
148. Wu HS, Lin LC. The moderating effect of nutritional status on depressive symptoms in veteran elders with dementia: a spaced retrieval combined with Montessori-based activities. *J Adv Nurs*. 2013 Oct;69(10):2229-41. doi: <https://dx.doi.org/10.1111/jan.12097>. PMID: 23444942.
149. van der Ploeg ES, Eppingstall B, Camp CJ, et al. A randomized crossover trial to study the effect of personalized, one-to-one interaction using Montessori-based activities on agitation, affect, and engagement in nursing home residents with Dementia. *Int Psychogeriatr*. 2013 Apr;25(4):565-75. doi: <https://dx.doi.org/10.1017/S1041610212002128>. PMID: 23237211.
150. Kurz A, Thone-Otto A, Cramer B, et al. CORDIAL: cognitive rehabilitation and cognitive-behavioral treatment for early dementia in Alzheimer disease: a multicenter, randomized, controlled trial. *Alzheimer Dis Assoc Disord*. 2012 Jul-Sep;26(3):246-53. PMID: 21986341.
151. Lin LC, Huang YJ, Watson R, et al. Using a Montessori method to increase eating ability for institutionalised residents with dementia: a crossover design. *J Clin Nurs*. 2011 Nov;20(21-22):3092-101. doi: <https://dx.doi.org/10.1111/j.1365-2702.2011.03858.x>. PMID: 21981704.
152. Clare L, Linden DE, Woods RT, et al. Goal-oriented cognitive rehabilitation for people with early-stage Alzheimer disease: a single-blind randomized controlled trial of clinical efficacy. *Am J Geriatr Psychiatry*. 2010 Oct;18(10):928-39. doi: <https://dx.doi.org/10.1097/JGP.0b013e3181d5792a>. PMID: 20808145.
153. Lam LC, Lui VW, Luk DN, et al. Effectiveness of an individualized functional training program on affective disturbances and functional skills in mild and moderate dementia--a randomized control trial. *Int J Geriatr Psychiatry*. 2010 Feb;25(2):133-41. doi: <https://dx.doi.org/10.1002/gps.2309>. PMID: 19582757.
154. Lin LC, Huang YJ, Su SG, et al. Using spaced retrieval and Montessori-based activities in improving eating ability for residents with dementia. *Int J Geriatr Psychiatry*. 2010 Oct;25(10):953-9. doi: <https://dx.doi.org/10.1002/gps.2433>. PMID: 20054841.
155. Loewenstein DA, Acevedo A, Czaja SJ, et al. Cognitive rehabilitation of mildly impaired Alzheimer disease patients on cholinesterase inhibitors. *Am J Geriatr Psychiatry*. 2004 Jul-Aug;12(4):395-402. PMID: 15249277.
156. Beck CK, Vogelpohl TS, Rasin JH, et al. Effects of behavioral interventions on disruptive behavior and affect in demented nursing home residents. *Nurs Res*. 2002 Jul-Aug;51(4):219-28. PMID: 12131234.
157. Quayhagen MP, Quayhagen M, Corbeil RR, et al. A dyadic remediation program for care recipients with dementia. *Nurs Res*. 1995 May-Jun;44(3):153-9. PMID: 7761291.
158. Tappen RM. The effect of skill training on functional abilities of nursing home residents with dementia. *Res Nurs Health*. 1994 Jun;17(3):159-65. PMID: 8184127.
159. Tseng CF, Lee SH, Hsieh TC, et al. Smart Restored by Learning Exercise Alleviates the Deterioration of Cognitive Function in Older Adults with Dementia-A Quasi-Experimental Research. *Int J Environ Res Public Health*. 2019 04 09;16(7):09. doi: <https://dx.doi.org/10.3390/ijerph16071270>. PMID: 30970666.
160. Kallio EL, Ohman H, Hietanen M, et al. Effects of Cognitive Training on Cognition and Quality of Life of Older Persons with Dementia. *J Am Geriatr Soc*. 2018 Apr;66(4):664-70. doi: <https://dx.doi.org/10.1111/jgs.15196>. PMID: 29345724.
161. Cavallo M, Hunter EM, van der Hiele K, et al. Computerized Structured Cognitive Training in Patients Affected by Early-Stage Alzheimer's Disease is Feasible and Effective: A Randomized Controlled Study. *Arch Clin Neuropsychol*. 2016 Sep 06;06:06. PMID: 27600448.
162. Giuli C, Papa R, Lattanzio F, et al. The Effects of Cognitive Training for Elderly: Results from My Mind Project. *Rejuvenation Res*. 2016 Dec;19(6):485-94. PMID: 26952713.
163. Hsu CN, Lin LC, Wu SC. The effects of spaced retrieval training in improving hyperphagia of people living with dementia in residential settings. *J Clin Nurs*. 2017 Oct;26(19-20):3224-31. doi: <https://dx.doi.org/10.1111/jocn.13671>. PMID: 27878873.
164. Kao CC, Lin LC, Wu SC, et al. Effectiveness of different memory training programs on improving hyperphagic behaviors of residents with dementia: a longitudinal single-blind study. *Clin Interv Aging*. 2016;11:707-20. doi: <https://dx.doi.org/10.2147/CIA.S102027>. PMID: 27307717.
165. Kallio EL, Ohman H, Carlson S, et al. Feasibility and baseline findings of a Finnish cognitive training (FINCOG) intervention in a randomised controlled trial among community-dwelling persons with dementia. *European Geriatric Medicine*. 2017 July;8(3):245-9. doi: <http://dx.doi.org/10.1016/j.eurger.2017.04.008>. PMID: 616108233.
166. Giovagnoli AR, Manfredi V, Parente A, et al. Cognitive training in Alzheimer's disease: a controlled randomized study. *Neurol Sci*. 2017 Aug;38(8):1485-93. doi: <https://dx.doi.org/10.1007/s10072-017-3003-9>. PMID: 28577267.
167. Chen HM, Tsai LJ, Chao SY, et al. Study on the Effects of Individualized Learning Therapy on Cognitive Function and Behavioral and Psychological Symptoms of Dementia in the Institutionalized Older Adults. *J Nurs Res*. 2016 Dec;24(4):300-10. PMID: 27846102.
168. De Luca R, Bramanti A, De Cola MC, et al. Cognitive training for patients with dementia living in a sicilian nursing home: a novel web-based approach. *Neurol Sci*. 2016 01 Oct;37(10):1685-91. doi: <http://dx.doi.org/10.1007/s10072-016-2659-x>. PMID: 611200073.
169. Kawashima R, Hiller D, Sereda S, et al. SAIDO Learning as a Cognitive Intervention for Dementia Care: aPreliminary Study. In *J Am Med Dir Assoc*
170. Bergamaschi S, Arcara G, Calza A, et al. One-year repeated cycles of cognitive training (CT) for Alzheimer's disease. *Aging Clin Exp Res*. 2013 Aug;25(4):421-6. doi: <https://dx.doi.org/10.1007/s40520-013-0065-2>. PMID: 23784727.
171. Mapelli D, Di Rosa E, Nocita R, et al. Cognitive stimulation in patients with dementia: randomized controlled trial. *Dement Geriatr Cogn Dis Extra*. 2013;3(1):263-71. doi: <https://dx.doi.org/10.1159/000353457>. PMID: 24052800.
172. Tarraga L, Boada M, Modinos G, et al. A randomised pilot study to assess the efficacy of an interactive, multimedia tool of cognitive stimulation in Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. 2006 Oct;77(10):1116-21. PMID: 16820420.
173. McGilton KS, Rivera TM, Dawson P. Can we help persons with dementia find their way in a new environment?.[Erratum appears in *Aging Ment Health*. 2003 Nov;7(6):following author index]. *Aging Ment Health*. 2003 Sep;7(5):363-71. PMID: 12959805.
174. Davis RN, Massman PJ, Doody RS. Cognitive intervention in Alzheimer disease: a randomized placebo-controlled study. *Alzheimer Dis Assoc Disord*. 2001 Jan-Mar;15(1):1-9. PMID: 11236819.
175. Zarit SH, Zarit JM, Reeve KE. Memory training for severe memory loss: Effects on senile dementia patients and their families. *Gerontologist*. 1982;22(4):373-7. PMID: 13238424.
176. Lok N, Buldukoglu K, Barcin E. Effects of the cognitive stimulation therapy based on Roy's adaptation model on Alzheimer's patients' cognitive functions, coping-adaptation skills, and quality of life: A randomized controlled trial. *Perspect Psychiatr Care*. 2020 Jan 12;12:12. doi: <https://dx.doi.org/10.1111/ppc.12472>. PMID: 31930518.
177. Lin HC, Yang YP, Cheng WY, et al. Distinctive effects between cognitive stimulation and reminiscence therapy on cognitive function and quality of life for different types of behavioural problems in dementia. *Scand J Caring Sci*. 2018 Jun;32(2):594-602. doi: <https://dx.doi.org/10.1111/scs.12484>. PMID: 28881430.
178. Orgeta V, Leung P, Yates L, et al. Individual cognitive stimulation therapy for dementia: a clinical effectiveness and cost-effectiveness pragmatic, multicentre, randomised controlled trial. *Health Technol Assess*. 2015 Aug;19(64):1-108. doi: <https://dx.doi.org/10.3310/hta19640>. PMID: 26292178.
179. Orrell M, Yates L, Leung P, et al. The impact of individual Cognitive Stimulation Therapy (iCST) on cognition, quality of life, caregiver health, and family relationships in dementia: A randomised controlled trial. *PLoS Med*. 2017 Mar;14(3):e1002269. doi: <https://dx.doi.org/10.1371/journal.pmed.1002269>. PMID: 28350796.
180. Orrell M, Aguirre E, Spector A, et al. Maintenance cognitive stimulation therapy for dementia: single-blind, multicentre, pragmatic randomised controlled trial. *Br J Psychiatry*. 2014 Jun;204(6):454-61. doi: <https://dx.doi.org/10.1192/bjp.bp.113.137414>. PMID: 24676963.
181. Spector A, Thorgrimsen L, Woods B, et al. Efficacy of an evidence-based cognitive stimulation therapy programme for people with dementia: randomised controlled trial. *Br J Psychiatry*. 2003 Sep;183:248-54. PMID: 12948999.
182. Knapp M, Thorgrimsen L, Patel A, et al. Cognitive stimulation therapy for people with dementia: cost-effectiveness analysis. *Br J Psychiatry*. 2006 Jun;188:574-80. PMID: 16738349.
183. Cheung DSK, Li B, Lai DWL, et al. Cognitive Stimulating Play Intervention for Dementia: A Feasibility Randomized Controlled Trial. *Am J Alzheimers Dis Other Demen*. 2019 02;34(1):63-71. doi: <https://dx.doi.org/10.1177/1533317518808036>. PMID: 30370782.
184. Capotosto E, Belacchi C, Gardini S, et al. Cognitive stimulation therapy in the Italian context: Its efficacy in cognitive and non-cognitive measures in older adults with dementia. *Int J Geriatr Psychiatry*. 2017 Mar;32(3):331-40. doi: <http://dx.doi.org/10.1002/gps.4521>. PMID: 2017-05902-002.
185. Wong G, Yek O, Zhang A, et al. Cultural adaptation of cognitive stimulation therapy (CST) for Chinese people with dementia: multicentre pilot study. In *Int J Geriatr Psychiatry*
186. Cove J, Jacobi N, Donovan H, et al. Effectiveness of weekly cognitive stimulation therapy for people with dementia and the additional impact of enhancing cognitive stimulation therapy with a carer training program. *Clin Interv Aging*. 2014;9:2143-50. doi: <https://dx.doi.org/10.2147/CIA.S66232>. PMID: 25525349.
187. Yamanaka K, Kawano Y, Noguchi D, et al. Effects of cognitive stimulation therapy Japanese version (CST-J) for people with dementia: a single-blind, controlled clinical trial.[Erratum appears in *Aging Ment Health*. 2013;17(5):654]. *Aging Ment Health*. 2013;17(5):579-86. doi: <https://dx.doi.org/10.1080/13607863.2013.777395>. PMID: 23550665.
188. Niu YX, Tan JP, Guan JQ, et al. Cognitive stimulation therapy in the treatment of neuropsychiatric symptoms in Alzheimer's disease: a randomized controlled trial. *Clin Rehabil*. 2010 Dec;24(12):1102-11. doi: <https://dx.doi.org/10.1177/0269215510376004>. PMID: 20713437.
189. Matsuda O. Cognitive stimulation therapy for Alzheimer's disease: the effect of cognitive stimulation therapy on the progression of mild Alzheimer's disease in patients treated with donepezil. *Int Psychogeriatr*. 2007;19(2):241-52. doi: 10.1017/S1041610206004194. PMID: CN-00640778.
190. Pedrinolla A, Tamburin S, Brasioli A, et al. An Indoor Therapeutic Garden for Behavioral Symptoms in Alzheimer's Disease: A Randomized Controlled Trial. *J Alzheimers Dis*. 2019;71(3):813-23. doi: <https://dx.doi.org/10.3233/JAD-190394>. PMID: 31424399.
191. Tse MMY, Lau JL, Kwan R, et al. Effects of play activities program for nursing home residents with dementia on pain and psychological well-being: Cluster randomized controlled trial. *Geriatr Gerontol Int*. 2018 Oct;18(10):1485-90. doi: <https://dx.doi.org/10.1111/ggi.13509>. PMID: 30255637.
192. Lin Q, Cao Y, Gao J. The impacts of a gO-game (Chinese chess) intervention on Alzheimer disease in a Northeast Chinese population. *Front Aging Neurosci*. 2015;7 (AUG) (no pagination):163. doi: <http://dx.doi.org/10.3389/fnagi.2015.00163>. PMID: 607203463.
193. Van Hantsma KS, Curyto K, Abbott KM, et al. A randomized controlled trial for an individualized positive psychosocial intervention for the affective and behavioral symptoms of dementia in nursing home residents. *J Gerontol B Psychol Sci Soc Sci*. 2015 Jan;70(1):35-45. doi: <https://dx.doi.org/10.1093/geronb/gbt102>. PMID: 24304555.
194. Low LF, Brodaty H, Goodenough B, et al. The Sydney Multisite Intervention of LaughterBosses and ElderClowns (SMILE) study: cluster randomised trial of humour therapy in nursing homes. *BMJ Open*. 2013 Jan 11;3(1):11. doi: <https://dx.doi.org/10.1136/bmjopen-2012-002072>. PMID: 23315520.
195. Kolanowski A, Litaker M, Buettner L, et al. A randomized clinical trial of theory-based activities for the behavioral symptoms of dementia in nursing home residents. *J Am Geriatr Soc*. 2011 Jun;59(6):1032-41. doi: <https://dx.doi.org/10.1111/j.1532-5415.2011.03449.x>. PMID: 21649633.
196. Buettner LL, Fitzsimmons S. AD-venture program: therapeutic biking for the treatment of depression in long-term care residents with dementia. *Am J Alzheimers Dis Other Demen*. 2002 Mar-Apr;17(2):121-7. PMID: 11954670.
197. Hopman-Rock M, Staats PGM, Tak ECPM, et al. The effects of a psychomotor activation programme for use in groups of cognitively impaired people in homes for the elderly. *Int J Geriatr Psychiatry*. 1999 August;14(8):633-42. doi: <http://dx.doi.org/10.1002/%28SIC%291099-1166%28199908%2914:8%3C633::AID-GPS996%3E3.0.CO;2-W>. PMID: 29403282.
198. Laksono WAW, Haryanto J, Bahiyah K. The impact of chess games towards comfortableness of cognitive mind on elderly. *Indian Journal of Public Health Research and Development*. 2019 September;10(9):873-7.
199. Lathifah MN, Haryanto J, Fauziningtyas R. The influence of dhakonan games to impede dementia in elders. *Indian Journal of Public Health Research and Development*. 2019 September;10(9):1440-4.
200. Li DM, Li XX. The effect of folk recreation program in improving symptoms: a study of Chinese elder dementia patients. *Int J Geriatr Psychiatry*. 2017 08;32(8):901-8. doi: <https://dx.doi.org/10.1002/gps.4543>. PMID: 27345629.
201. Schall A, Tesky VA, Adams AK, et al. Art museum-based intervention to promote emotional well-being and improve quality of life in people with dementia: The ARTEMIS project. *Dementia*. 2017 Jan 01;1471301217730451. doi: <https://dx.doi.org/10.1177/1471301217730451>. PMID: 28914089.
202. Hattori H, Hattori C, Hokao C, et al. Controlled study on the cognitive and psychological effect of coloring and drawing in mild Alzheimer's disease patients. *Geriatr Gerontol Int*. 2011 Oct;11(4):431-7. doi: <https://dx.doi.org/10.1111/j.1447-0594.2011.00698.x>. PMID: 21518170.

203. Onder G, Zanetti O, Giacobini E, et al. Reality orientation therapy combined with cholinesterase inhibitors in Alzheimer's disease: randomised controlled trial. *Br J Psychiatry*. 2005 Nov;187:450-5. PMID: 16260821.
204. Quinn C, Toms G, Jones C, et al. A pilot randomized controlled trial of a self-management group intervention for people with early-stage dementia (The SMART study). *Int Psychogeriatr*. 2016 May;28(5):787-800. doi: <https://dx.doi.org/10.1017/S1041610215002094>. PMID: 26674087.
205. Quinn C, Anderson D, Toms G, et al. Self-management in early-stage dementia: a pilot randomised controlled trial of the efficacy and cost-effectiveness of a self-management group intervention (the SMART study). *Trials*. 2014 Mar 08;15:74. doi: <https://dx.doi.org/10.1186/1745-6215-15-74>. PMID: 24606601.
206. Young DK, Kwok TC, Ng PY. A single blind randomized control trial on support groups for Chinese persons with mild dementia. *Clin Interv Aging*. 2014;9:2105-12. doi: <https://dx.doi.org/10.2147/CIA.S68687>. PMID: 25587218.
207. Tappen RM, Williams CL. Therapeutic conversation to improve mood in nursing home residents with Alzheimer's disease. *Res*. 2009 Oct;2(4):267-75. doi: <https://dx.doi.org/10.3928/19404921-20090428-02>. PMID: 20077983.
208. Burns A, Guthrie E, Marino-Francis F, et al. Brief psychotherapy in Alzheimer's disease: randomised controlled trial. *Br J Psychiatry*. 2005 Aug;187:143-7. PMID: 16055825.
209. Cohen CI, Hyland K, Devlin M. An evaluation of the use of the natural helping network model to enhance the well-being of nursing home residents. *Gerontologist*. 1999;39(4):426-33. PMID: 107227653. Language: English. Entry Date: 19991201. Revision Date: 20150711. Publication Type: Journal Article.
210. Wallis GG, Baldwin M, Higginbotham P. Reality orientation therapy--a controlled trial. *Br J Med Psychol*. 1983 Sep;56(Pt 3):271-7. PMID: 6354248.
211. Fritsch T, Kwak J, Grant S, et al. Impact of TimeSlips, a Creative Expression Intervention Program, on Nursing Home Residents with Dementia and their Caregivers. *Gerontologist*. 2009;49(1):117-27. doi: 10.1093/geront/gnp008. PMID: CN-00755001.
212. Lin LC, Yang MH, Kao CC, et al. Using acupressure and Montessori-based activities to decrease agitation for residents with dementia: a cross-over trial. *J Am Geriatr Soc*. 2009 Jun;57(6):1022-9. PMID: 19507295.
213. Houser WS, George DR, Chinchilli VM. Impact of TimeSlips creative expression program on behavioral symptoms and psychotropic medication use in persons with dementia in long-term care: a cluster-randomized pilot study. *Am J Geriatr Psychiatry*. 2014 Apr;22(4):337-40. doi: <https://dx.doi.org/10.1016/j.jagp.2012.12.005>. PMID: 23602304.
214. van Dijk AM, van Weert JC, Droes RM. Does theatre improve the quality of life of people with dementia? *Int Psychogeriatr*. 2012 Mar;24(3):367-81. doi: <https://dx.doi.org/10.1017/S1041610211001992>. PMID: 22040605.
215. Phillips L, Reid-Arndt S, Pak Y. Effects of a creative expression intervention on emotions, communication, and quality of life in persons with dementia. *Nurs Res*. 2010;59(6):417-25. doi: 10.1097/NNR.0b013e3181faff52. PMID: CN-00770848.
216. Okamura H, Otani M, Shimoyama N, et al. Combined Exercise and Cognitive Training System for Dementia Patients: A Randomized Controlled Trial. *Dement Geriatr Cogn Disord*. 2018;45(5-6):318-25. doi: <https://dx.doi.org/10.1159/000490613>. PMID: 30036871.
217. Rokstad AMM, Engedal K, Kirkevold O, et al. The impact of attending day care designed for home-dwelling people with dementia on nursing home admission: a 24-month controlled study. *BMC Health Serv Res*. 2018 Nov 16;18(1):864. doi: <https://dx.doi.org/10.1186/s12913-018-3686-5>. PMID: 30445937.
218. Satoh M, Ogawa J, Tokita T, et al. Physical exercise with music maintains activities of daily living in patients with dementia: mihama-kiho project part 2. In *Journal of alzheimers disease*
219. Middelstadt J, Folkerts AK, Blawath S, et al. Cognitive Stimulation for People with Dementia in Long-Term Care Facilities: Baseline Cognitive Level Predicts Cognitive Gains, Moderated by Depression. *J Alzheimers Dis*. 2016 08 04;54(1):253-68. doi: <https://dx.doi.org/10.3233/JAD-160181>. PMID: 27497474.
220. Fernandez-Calvo B, Contador I, Ramos F, et al. Effect of unawareness on rehabilitation outcome in a randomised controlled trial of multicomponent intervention for patients with mild Alzheimer's disease. *Neuropsychol*. 2015;25(3):448-77. doi: <https://dx.doi.org/10.1080/09602011.2014.948461>. PMID: 25121567.
221. Luttenberger K, Donath C, Uter W, et al. Effects of multimodal nondrug therapy on dementia symptoms and need for care in nursing home residents with degenerative dementia: a randomized-controlled study with 6-month follow-up. *J Am Geriatr Soc*. 2012 May;60(5):830-40. doi: <https://dx.doi.org/10.1111/j.1532-5415.2012.03938.x>. PMID: 22468985.
222. Yamagami T, Takayama Y, Maki Y, et al. A randomized controlled trial of brain-activating rehabilitation for elderly participants with dementia in residential care homes. *Dement Geriatr Cogn Dis Extra*. 2012 Jan;2(1):372-80. doi: <https://dx.doi.org/10.1159/000342614>. PMID: 23300492.
223. Brooker DJ, Argyle E, Scally AJ, et al. The enriched opportunities programme for people with dementia: a cluster-randomised controlled trial in 10 extra care housing schemes. *Aging Ment Health*. 2011 Nov;15(8):1008-17. doi: <https://dx.doi.org/10.1080/13607863.2011.583628>. PMID: 21702705.
224. Ferrero-Arias J, Goni-Imizcoz M, Gonzalez-Bernal J, et al. The efficacy of nonpharmacological treatment for dementia-related apathy. *Alzheimer Dis Assoc Disord*. 2011 Jul-Sep;25(3):213-9. doi: <https://dx.doi.org/10.1097/WAD.0b013e3182087dbc>. PMID: 21346517.
225. McCurry SM, Pike KC, Vitiello MV, et al. Increasing walking and bright light exposure to improve sleep in community-dwelling persons with Alzheimer's disease: results of a randomized, controlled trial. *J Am Geriatr Soc*. 2011 Aug;59(8):1393-402. doi: <https://dx.doi.org/10.1111/j.1532-5415.2011.03519.x>. PMID: 21797835.
226. Femia EE, Zarit SH, Stephens MA, et al. Impact of adult day services on behavioral and psychological symptoms of dementia. *Gerontologist*. 2007 Dec;47(6):775-88. PMID: 18192631.
227. Chapman SB, Weiner MF, Rackley A, et al. Effects of cognitive-communication stimulation for Alzheimer's disease patients treated with donepezil. *J Speech Lang Hear Res*. 2004 Oct;47(5):1149-63. PMID: 15603468.
228. Tadaka E, Kanagawa K. Randomized controlled trial of a group care program for community-dwelling elderly people with dementia. *Japan Journal of Nursing Science*. 2004 Aug;1(1):19-25. doi: <http://dx.doi.org/10.1111/j.1742-7924.2004.00006.x>. PMID: 2007-06056-002.
229. Wimo A, Mattsson B, Adolfsson R, et al. Dementia day care and its effects on symptoms and institutionalization--a controlled Swedish study. *Scand J Prim Health Care*. 1993 Jun;11(2):117-23. PMID: 8356361.
230. Luttenberger K, Hofner B, Graessel E. Are the effects of a non-drug multimodal activation therapy of dementia sustainable? Follow-up study 10 months after completion of a randomised controlled trial. *BMC Neurol*. 2012 Dec 05;12:151. doi: <https://dx.doi.org/10.1186/1471-2377-12-151>. PMID: 23217188.
231. Sado M, Funaki K, Ninomiya A, et al. Does the Combination of the Cognitive Interventions Improve the Function of Daily Living and Save the Long-Term Care Cost? A Pilot Study of Effectiveness and Cost Saving Analysis of "Learning Therapy" for People with Dementia. *J Alzheimers Dis*. 2020 Feb 27;27:27. doi: <https://dx.doi.org/10.3233/JAD-190886>. PMID: 32116248.
232. Young DK, Ng PY, Kwok T, et al. The effects of an expanded cognitive stimulation therapy model on the improvement of cognitive ability of elderly with mild stage Dementia living in a community - a randomized waitlist controlled trial. *Aging Ment Health*. 2019 Jul;23(7):855-62. doi: <https://dx.doi.org/10.1080/13607863.2018.1471586>. PMID: 29781725.
233. Chen YL, Pei YC. Musical dual-task training in patients with mild-to-moderate dementia: a randomized controlled trial. *Neuropsychiatr*. 2018;14:1381-93. doi: <https://dx.doi.org/10.2147/NDT.S159174>. PMID: 29881275.
234. Reisberg B, Shao Y, Golomb J, et al. Comprehensive, Individualized, Person-Centered Management of Community-Residing Persons with Moderate-to-Severe Alzheimer Disease: A Randomized Controlled Trial. *Dement Geriatr Cogn Disord*. 2017;43(1-2):100-17. doi: <https://dx.doi.org/10.1159/000455397>. PMID: 28122366.
235. Kampragkou C, Iakovidis P, Kampragkou E, et al. Effects of a 12-week aerobic exercise program combined with music therapy and memory exercises on cognitive and functional ability in people with middle type of alzheimer's disease. 2017;4(5):262-8. doi: 10.15621/ijphy/2017/v4i5/159420. PMID: CN-01425536.
236. Masayuki S, Jun-ichi O, Tomoko T, et al. Physical Exercise with Music Maintains Activities of Daily Living in Patients with Dementia: Mihama-Kiho Project Part 21. *J Alzheimers Dis*. 2017;57(1):85-96. doi: 10.3233/JAD-161217. PMID: 121747018. Language: English. Entry Date: 20180302. Revision Date: 20180619. Publication Type: journal article.
237. Kim H-J, Yang Y, Oh J-G, et al. Effectiveness of a community-based multidomain cognitive intervention program in patients with Alzheimer's disease. *Geriatr Gerontol Int*. 2016;16(2):191-9. doi: 10.1111/ggi.12453. PMID: CN-01132849.
238. Wesson J, Clemons L, Brodaty H, et al. A feasibility study and pilot randomised trial of a tailored prevention program to reduce falls in older people with mild dementia. *BMC geriatr*. 2013 Sep 03;13:89. doi: <https://dx.doi.org/10.1186/1471-2318-13-89>. PMID: 24004682.
239. Burgener SC, Marsh-Yant S, Nega KK. A Combined, Multimodal Intervention for Individuals with Dementia. *Res*. 2011;4(1):64-75. doi: 10.3928/19404921-20100504-01. PMID: 104897556. Language: English. Entry Date: 20110622. Revision Date: 20150711. Publication Type: Journal Article.
240. Burgener SC, Yang Y, Gilbert R, et al. The effects of a multimodal intervention on outcomes of persons with early-stage dementia. *Am J Alzheimers Dis Other Dement*. 2008 Aug-Sep;23(4):382-94. doi: <https://dx.doi.org/10.1177/1533317508317527>. PMID: 18453642.
241. Lauriks SM, Meiland FJMP, Oste JPM, et al. Effects of Assistive Home Technology on quality of life and falls of people with dementia and job satisfaction of caregivers; results from a pilot randomized controlled trial. *Assist Technol*. 2018 Dec 28:1-8. doi: <https://dx.doi.org/10.1080/10400435.2018.1531952>. PMID: 30592439.
242. Tchalla AE, Lachal F, Cardinaud N, et al. Preventing and managing indoor falls with home-based technologies in mild and moderate Alzheimer's disease patients: Pilot study in a community dwelling. *Dementia and Geriatric Cognitive Disorders*. 2013 September;36(3-4):251-61. doi: <http://dx.doi.org/10.1159/000351863>. PMID: 369900271.
243. Rowe MA, Kairalla JA, McCrae CS. Sleep in dementia caregivers and the effect of a nighttime monitoring system. *J Nurs Scholarsh*. 2010 Sep 01;42(3):338-47. doi: <https://dx.doi.org/10.1111/j.1547-5069.2010.01337.x>. PMID: 20738745.
244. Gaugler JE, Zmora R, Mitchell LL, et al. Six-Month Effectiveness of Remote Activity Monitoring for Persons Living With Dementia and Their Family Caregivers: An Experimental Mixed Methods Study. *The Gerontologist*. 2019 09 Jan;59(1):78-89. doi: <http://dx.doi.org/10.1093/geront/gny078>. PMID: 625875732.
245. Khedr EM, Salama RH, Abdel Hameed M, et al. Therapeutic Role of Transcranial Direct Current Stimulation in Alzheimer Disease Patients: Double-Blind, Placebo-Controlled Clinical Trial. *Neurorehabil Neural Repair*. 2019 05;33(5):384-94. doi: <https://dx.doi.org/10.1177/1545968319840285>. PMID: 30940012.
246. Lozano AM, Fosdick L, Chakravarty MM, et al. A Phase II Study of Fornix Deep Brain Stimulation in Mild Alzheimer's Disease. *J Alzheimers Dis*. 2016 09 06;54(2):777-87. doi: <https://dx.doi.org/10.3233/JAD-160017>. PMID: 27567810.
247. Leoutsakos JS, Yan H, Anderson WS, et al. Deep Brain Stimulation Targeting the Fornix for Mild Alzheimer Dementia (the ADvance Trial): A Two Year Follow-up Including Results of Delayed Activation. *J Alzheimers Dis*. 2018;64(2):597-606. doi: <https://dx.doi.org/10.3233/JAD-180121>. PMID: 29914028.
248. Wu Y, Xu W, Liu X, et al. Adjunctive treatment with high frequency repetitive transcranial magnetic stimulation for the behavioral and psychological symptoms of patients with Alzheimer's disease: a randomized, double-blind, sham-controlled study. *Shanghai Arch Psychiatry*. 2015 Oct;27(5):280-8. doi: <https://dx.doi.org/10.11919/j.issn.1002-0829.215107>. PMID: 26977125.
249. Cotelli M, Manenti R, Brambilla M, et al. Anodal tDCS during face-name associations memory training in Alzheimer's patients. *Front Aging Neurosci*. 2014;6:38. doi: <https://dx.doi.org/10.3389/fnagi.2014.00038>. PMID: 24678298.
250. Olazaran J, Gonzalez B, Osa-Ruiz E, et al. Motor effects of radio electric asymmetric conveyer in Alzheimer's disease: results from a cross-over trial. *J Alzheimers Dis*. 2014;42(1):325-32. doi: <https://dx.doi.org/10.3233/JAD-140417>. PMID: 24898637.
251. Olazaran J, Gonzalez B, Lopez-Alvarez J, et al. Motor effects of REAC in advanced Alzheimer's disease: results from a pilot trial. *J Alzheimers Dis*. 2013;36(2):297-302. doi: <https://dx.doi.org/10.3233/JAD-130077>. PMID: 23603397.
252. Suemoto CK, Apolinario D, Nakamura-Palacios EM, et al. Effects of a non-focal plasticity protocol on apathy in moderate Alzheimer's disease: a randomized, double-blind, sham-controlled trial. *Brain Stimul*. 2014 Mar-Apr;7(2):308-13. doi: <https://dx.doi.org/10.1016/j.brs.2013.10.003>. PMID: 24262299.
253. Ahmed MA, Darwish ES, Khedr EM, et al. Effects of low versus high frequencies of repetitive transcranial magnetic stimulation on cognitive function and cortical excitability in Alzheimer's dementia. *J Neurol*. 2012 Jan;259(1):83-92. doi: <https://dx.doi.org/10.1007/s00415-011-6128-4>. PMID: 21671144.
254. Scherder EJ, van Tol MJ, Swaab DF. High-frequency cranial electrostimulation (CES) in patients with probable Alzheimer's disease. *Am J Phys Med Rehabil*. 2006 Jul;85(7):614-8. PMID: 16788393.
255. van Dijk KR, Scheltens P, Luijpen MW, et al. Peripheral electrical stimulation in Alzheimer's disease. *Dement Geriatr Cogn Disord*. 2005;19(5-6):361-8. PMID: 15802911.
256. Hozumi S, Hori H, Okawa M, et al. Favorable effect of transcranial electrostimulation on behavior disorders in elderly patients with dementia: a double-blind study. *Int J Neurosci*. 1996 Nov;88(1-2):1-10. PMID: 9003961.
257. Liu B, Chen X, Li Y, et al. Effect of passive finger exercises on grip strength and the ability to perform activities of daily living for older people with dementia: a 12-week randomized controlled trial. *Clin Interv Aging*. 2018;13:2169-77. doi: <https://dx.doi.org/10.2147/CIA.S174756>. PMID: 30464426.
258. Rostad HM, Utne I, Grov EK, et al. The impact of a pain assessment intervention on pain score and analgesic use in older nursing home residents with severe dementia: A cluster randomised controlled trial. *Int J Nurs Stud*. 2018 Aug;84:52-60. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2018.04.017>. PMID: 29763832.
259. Batchelor-Murphy M, McConnell E, Amella E, et al. Experimental Comparison of Efficacy for Three Handfeeding Techniques in Dementia. *J Am Geriatr Soc*. 2017;65(4):E89-e94. doi: 10.1111/jgs.14728. PMID: CN-01415879.
260. Kim J, Yang J. Effectiveness of a community-based program for suicide prevention among elders with early-stage dementia: a controlled observational study. In *Geriatric nursing (new york, NY)*
261. Hobbelen JH, Tan FE, Verhey FR, et al. Passive movement therapy in severe paratonia: a multicenter randomized clinical trial. *Int Psychogeriatr*. 2012 May;24(5):834-44. doi: <https://dx.doi.org/10.1017/S1041610211002468>. PMID: 22185768.
262. Coyne ML, Hoskins L. Improving eating behaviors in dementia using behavioral strategies. *Clin Nurs Res*. 1997 Aug;6(3):275-90. PMID: 9281930.

Appendix E. Psychosocial Interventions for Caregiver Well-Being

Psychosocial Interventions for Caregiver Well-Being

Table E-1. Risk of bias assessment: psychosocial interventions for caregiver well-being

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Au 2020 ¹ (no PMID)	8 weeks	Low 8%	Low	Low	Low	Low	Low	University	Low
Au 2019 ² (31279613)	5 months	Low 15%	Low	Medium	Medium	Low	Low	Government	Low
Kunik 2020 ³ (32115311)	3 months 6 months 12 months	Low 18%	Low	Medium	High	Low	Low	National Institute of Nursing	Low
Bjorge 2019 ⁴ (31651321)	3 months	Low 15%	Low	Medium	Medium	Low	Medium	None	Medium
Ghaffari 2019 ⁵ (no PMID)	8 weeks	Low 7%	Low	Medium	Medium	Medium	Medium	University	Medium
Meng 2019 ⁶ (30884961)	3 months	Low 0%	Medium	Medium	Medium	Low	Medium	Government	Medium
Moskowitz 2019 ⁷ (31045422)	6 weeks	Low 9%	Low	Medium	Medium	Low	Medium	Government	Medium
Nordheim 2019 ⁸ (30775982)	6 months	Medium 19%	Low	Medium	Medium	Low	Medium	Government	Medium
Terracciano ⁹ 2019 (31866419)	6 weeks	High 27%	X	X	X	X	X	Government Nonprofit	High
Uyar 2019 ¹⁰ (no PMID)	16 weeks	Medium 12%	Medium	Medium	Medium	Low	Medium	NR	High Very unclear methods and intervention description
Wawrziczny 2019 ¹¹ (29665714)	10 weeks	X	High	X	X	X	X	Government	High
Boots 2018 ¹² (30006327)	8 weeks	Medium 16%	Low	Low	Medium	Low	Low	Alzheimer Nederland and the Alzheimer Research Fund Limburg.	High
Gitlin 2018 ¹³ (29192967)	4 months	Medium 31%	Low	Low	Medium	Low	Low	Government	Medium
Spalding-Wilson 2018 ¹⁴ (30258974)	1, 3, 6 months	Low 9%	Medium	Medium	High	Low	Low	National Science Foundation, private foundations	High
Wilz 2018 ¹⁵ (29190357)	6 months Follow up at 12 month	Medium 17%	Low	High	Medium	Low	Low	The German Federal Ministry of Health	High
Callahan 2017 ¹⁶ (27893087)	2 years	Medium 35%	Low	Low	High	Low	Medium	NIA	High
Whitlatch 2017 ¹⁷ (29171296)	6 months	Medium 15%	Medium	Medium	High	Low	High	Government	High
Charlesworth 2016 ¹⁸ (27521377)	5 months 1 year	Low 5 months: 7% 12 months: 13%	Low	Low	Medium	Low	Medium	Government	Low
Cheng 2016 ¹⁹ (27401052)	2 months	NR	Medium	Medium	Medium	Low	Medium	Government	Medium
Gonyea 2016 ²⁰ (24855313)	3 months 6 months	Low 15%	Medium	Medium	Medium	Low	Medium	Nonprofit	Medium
Laakkonen 2016 ²¹ (27060101)	3 months 9 months	Low 4%	Low	Medium	Medium	Low	Medium	Government Nonprofit University	Medium
Taati 2016 ²² (no PMID)	8 weeks	High 23%	Low	X	X	X	X	No funding	High

Blom 2015 ²³ (no PMID)	5-6 months	Medium 28%	Low	Medium	Low	Low	High	the Alzheimer's Society in the Netherlands, the health care provider Geriant and the VU University	High
Chiu 2015 ²⁴ (25615434)	4 weeks	Medium 16%	High	High	Low	Low	Low	The UHN AMO Innovation Fund	High
Gallagher-Thompson 2015 ²⁵ (25590939)	10 months	High 25%	Medium	X	X	X	X	The National Office of the Alzheimer's Association & the Alzheimer's Disease Center at University of California, Davis	High
Losada 2015 ²⁶ (26075381)	2 % 6 months	High 30%	X	X	X	X	X	NR	High
Otero 2015 ²⁷ (25331992)	1, 3, 6, 12 months	Low 4.62%	Low	Low	Medium	Low	Low	Spanish government	Low
Tremont 2015 ²⁸ (25074341)	6 months	Medium 15%	Low	Medium	Medium	Low	Medium	Government	Medium
Arango-Lasprilla 2014 ²⁹ (24550547)	3 months	High Unclear	X	X	X	X	X	No funding	High
Livingston 2014 ³⁰ (25300037)	4 months 8 months	Low 4 months: 9% 8 months: 13%	Low	Medium	Medium	Low	Low	Government	Low
Martin-Carrasco 2014 ³¹ (24113563)	4 months	Medium 4 months: 26% 8 months: 39%	Low	Low	Medium	Low	Medium	Government	Medium
Passoni 2014 ³² (24614271)	6 months	X	High	X	X	X	X	Not reported	High
Bruvik 2013 ³³ (24348500)	12 months	Medium 13%	Low	Low	Medium	Low	High	Government, foundations	High
Huang 2013 ³⁴ (23933422)	2 weeks, 3 & 6 months	Medium 10%	Medium	Medium	Low	Low	High	Government	High
Judge 2013 ³⁵ (22899427)	15 weeks	Medium 13%	Medium	Medium	High	Low	Low	NR	High
Kajiyama 2013 ³⁶ (23461355)	3 months	High 31%	Medium	X	X	X	X	The National Institute on Aging, part of the National Institutes of Health.	High
Kuo 2013 ³⁷ (22778053)	2 weeks, 3 & 6 months	Low 16%	Medium	Medium	Low	Low	High	Government	High
Moore 2013 ³⁸ (23916631)	6 weeks	Low 14%	Low	Medium	Medium	Low	Low	National Institute on Aging	Low
Joling 2012 ³⁹ (22303473)	12 months	Low 13%	Low	Medium	Medium	Low	Low	Government	Low
Waldorff 2012 ⁴⁰ (22807076)	1 year	Medium 16%	Low	Low	Medium	Low	Medium	Government	Medium
Chu 2011 ⁴¹ (20847363)	1 month	High 29%	X	X	X	X	X	University	High
de Rotrou 2011 ⁴² (20922772)	3 months 6 months	Medium 31%	Low	Medium	Medium	Low	Medium	Government	Medium
Guerra 2011 ⁴³ (20602013)	6 months	Low 3%	Low	Low	High	Low	High	Nonprofit	Medium
Losada 2011 ⁴⁴ (21061414)	3 months	High 29%	X	X	X	X	X	NR	High
Spijker 2011 ⁴⁵ (21358385)	12 months	Medium 36%	Low	Medium	Low	Low	High	The Dutch Organization of Health Research and Development (ZonMw).	High
Voight-Radloff 2011 ⁴⁶ (22021760)	6 weeks 16 weeks 26 weeks 52 weeks	Low 6-16 weeks Medium 26-52 weeks 6 weeks: 9% 16 weeks: 19% 26 weeks: 25% 52 weeks: 26%	Low	Medium	Medium	Low	Medium	Government	Low 6-16 weeks Medium 26-52 weeks

Wang 2011 ⁴⁷ (21752121)	6 months	Low 2%	Medium	Medium	Medium	Low	Medium	University	Medium
Wilz 2011 ⁴⁸ (no PMID)	3 months	Medium 25%	Medium	Medium	Medium	Low	Low	Government	Medium
Gitlin 2010 ⁴⁹ (20662955)	4 months 9 months	High 12%	Low	Medium	Medium	Low	Low	The National Institute on Aging and National Institute	High
Gitlin 2010 ⁵⁰ (20810376)	4 months 9 months	Medium 12% 27% at 9 month	Low	Medium	High	Low	High	In part by funds from the National Institute on Aging & the National Institute on Nursing Research & the Pennsylvania Department of Health, Tobacco Settlement	High
Kurz 2010 ⁵¹ (19946869)	15 months	High Unclear	X	X	X	X	X	German Federal Ministry of Education and Research	High
Williams 2010 ⁵² (20978227)	3, 6 months	X	High	X	X	X	X	National Institutes on Aging	High
Gavrilova 2009 ⁵³ (18814197)		Medium 12%	Low	Low	Medium	Low	High	The World Health Organization (WHO)	High
Martin-Carrasco 2009 ⁵⁴ (18949763)	4 months 10 months	Medium 10%	Low	Medium	Medium	Low	Medium	Industry	Medium
Andr�n 2008 ⁵⁵ (18269429)	6 & 12 months	X	High	X	X	X	X	Swedish Council for Social Research, foundations, university, and municipality	High
Gallagher- Thompson 2008 ⁵⁶ (25067886)	6 months	Medium 15%	Medium	Medium	Medium	Medium	Medium	Government	Medium
Marquez-Gonzalez 2007 ⁵⁷ (18074249)	2 months	High 47%	X	X	X	X	X	Not reported	High
Ulstein 2007 ⁵⁸ (17986818)	1 year	Medium 14%	Low	Medium	Medium	Low	High	Government, foundation	High
Gonyea 2006 ⁵⁹ (17169938)	6 weeks	Medium 12%	Low	Medium	High	Low	High	Foundations	High
Graff 2006 ⁶⁰ (17114212)	6 weeks 3 months	Medium 6 weeks High 12 weeks 6 weeks: 16% 12 weeks: 23%	Low	Medium	Medium	Low	Medium	Nonprofit University	Medium 6 weeks High 12 weeks
Beauchamp 2005 ⁶¹ (no PMID)	1 month	Low 3%	Medium	High	High	Low	Low	Not reported	High
Farran 2004 ⁶² (41552352)	18 months	Medium 17%	Low	Medium	Medium	Low	Medium	Government	Medium
Burgio 2003 ⁶³ (12937335)	6 months	Medium 16%	Medium	High	Medium	Low	Low	National Institute for Nursing Research	High
Burns 2003 ⁶⁴ (12937333)	2 years	High 55%	X	X	X	X	X	National Institute on Aging, National Institute of Nursing Research, Department of Veterans Affairs	High
Coon 2003 ⁶⁵ (14570964)	4, 7 months	High 23%	X	X	X	X	X	National Institute of Mental Health	High
Gallagher Thompson 2003 ⁶⁶ (12937336)	3 months	Low 0%	Medium	Medium	Medium	Medium	Medium	Government	Medium
Gitlin 2003 ⁶⁷ (12937332)	3 months	Medium 15%	Medium	High	Medium	Low	Low	Government	Medium
Hebert 2003 ⁶⁸ (12496309)	4 months	Medium 18%	Low	Medium	High	Low	Low	The Quebec Health Research Fund, the Alzheimer Society of Canada, the FRSQ Network for Geronto-Geriatric Research, and the Quebec Council for Social Research	High

Fung 2002 ⁶⁹ (12037799)	3 months	Medium 13%	Medium	High	High	Low	Low	Not reported	High
Stolley 2002 ⁷⁰ (11954669)	3, 6 & 12 months	High 27%	Medium	X	X	X	X	NR	High
Wright 2001 ⁷¹ (11885210)	2 & 6 weeks, 3, 6 & 12 months	High Unclear	X	X	X	X	X	Foundation	High
Buckwalter 1999 ⁷² (10222636)	6 & 12 months	High 29%	X	X	X	X	X	National Institutes of Health	High
Chang 1999 ⁷³ (10337848)	4, 8 & 12 weeks	Medium 25%	Medium	Medium	Medium	Low	High	NR	High
Ostwald 1999 ⁷⁴ (10396888)	5 months	Low 19.7%	Low	Medium	High	Low	Low	National Institute of Nursing Research	Medium

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table E-2. Characteristics of included studies: psychosocial interventions for caregiver well-being

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training			IC: FC:
Au 2020 ¹ (no PMID) Hong Kong Low Explanatory	Connecting through caregiving Intergenerational perspective-taking reappraisals Aim to promote balance between self-care and caring of others: (1) enhance self-awareness (2) connect with PLWD through empathetic understanding (3) connect with help Initial home visit, then 7 weekly telephone sessions 8 weeks total	Basic skill building intervention Initial home visit, then 7 weekly telephone sessions 8 weeks total	Home-based RCT N=72 dyads	Alzheimer's disease NPI 20 Physician-diagnosed	N=72 79 years 72% Female Race NR Education NR	NR	N=72 52 years 81% Female Race NR Education 11 Years formal education (mean) 100% Child	3 years caregiving (mean) 57% Living with PLWD 53% Employed	8 weeks	NR	CESD ZBI Satisfaction with Life Survey (all Chinese versions)

Au 2019 ² (31279613) Hong Kong Low Explanatory	Telephone-administered psycho-education with behavioral activation intervention. 8 biweekly sessions.	Telephone based psycho-education with general monitoring. 8 biweekly monitoring sessions.	Home-based RCT N=111	Alzheimer's disease NINCDS-ADRDA	NR	NR	N=111 57 years 81% Female Race NR Education 32% Primary 50% Secondary 18% Tertiary 29% Spouse or partner 65% Child	4+ years caregiving (mean) 11 hours daily caregiving	5 months	NR	CESD ZBI RAS Caregiver: self-efficacy for controlling upsetting thoughts 5 item
Kunik 2020 ³ (32115311) United States Low Pragmatic	Psychosocial approach to understanding behavioral problems / Unmet Needs Model including education, 6 to 8 weekly skills sessions, and telephone wrap-up session for PLWD and caregivers for 3 months	booklet on memory problems, community resources and caregivers, eight brief, weekly calls, and encouraged to bring concerns to primary care physician for 3 months	Community N=228	Documented diagnosed dementia	N=228 77 years 42% Female 67% Non-Hispanic white 69% 14 years	48% \$20k-49,999/yr 21% Black 9% Hispanic 2% Other	N=228 68 years 77% Female 66% Non-Hispanic white 78% 14 years 69% Spouse	88% Living with PLWD	3 months 6 months 12 months	Aggression (CMAI) Pain (PGPIS)	Depression (GDS) Pain (PGPIS) Burden (ZBS)
Bjorge 2019 ⁴ (31651321) Norway Medium Explanatory	Psychosocial intervention (education about dementia, counselling and group sessions) over 12 months	Usual care	Community-based RCT N=208 dyads	Non-specified dementia ICD 10	N=208 79 years 53% Female Race NR 9 Years of education	NR	N=208 64 years 76% Female Race NR Education NR 57% Spouse or partner 44% Child	58% Live with PLWD	3 months	MMSE NPI-Q Lawton & Brody's IADL CSDD PLWD nursing home placement	RSS GDS FEERS
Ghaffari 2019 ⁵ (no PMID) Iran Medium Explanatory	Resilience education 8 weekly 45-minute group sessions Training and education program aimed to promote resilience PowerPoint presentations and educational pamphlets	No treatment Received resilience development education after the study finished	Community-based RCT N=54 caregivers	Alzheimer's disease Physician-diagnosed	NR	NR	N=54 43 years 74% Female Race NR Education 22% Primary education 35% High school diploma 35% University education 77% Spouse or partner 23% Child	Months caregiving 4% <6 months 54% 6-24 months 42% >24 months Employment 18% Retired 50% Working 32% Homemaker	8 weeks	NR	GHQ
Meng 2019 ⁶ (30884961) USA Medium Explanatory	Telephone based cognitive behavioral therapy	In-person cognitive behavioral therapy	Home and Community-based RCT N=109 dyads	Alzheimer's disease or progressive dementia Physician-diagnosed	N=109 81 years 66% Female 100% African-American 12 Years education	NR	N=109 59 years 50% Female 100% African-American 15 Years education 23% Spouse or partner 66% Child	4 years caregiving (mean) 66% Live with PLWD	3 months	NR	Caregiver health (physician office visits, mental health visits, prescription medication, ER visits, hospital days, out of pocket expenditures in dollars, total monthly expenditures in dollars)

<p>Moskowitz 2019⁷ (31045422)</p> <p>USA Medium Explanatory</p>	<p>Life Enhancing Activities for Family Caregivers (LEAF). Taught eight emotion-regulating skills over 6 sessions.</p>	<p>Waitlist control, emotion-reporting</p>	<p>Home-based RCT N=170</p>	<p>Alzheimer's disease Frontotemporal Lewy body Parkinson's dementia Nonspecified dementia</p>	<p>NR</p>	<p>NR</p>	<p>N=170 63 years 84% Female 88% White Education 6% High school education or less 40% College degree or some college 40% Postgraduate education 67% Spouse or partner 27% Child</p>	<p>4+ years caregiving (mean)</p>	<p>6 weeks</p>	<p>Dementia Severity Rating Scale</p>	<p>PROMIS (depressive symptoms, mental health, physical health) NeuroQOL ZBI Caregiver Strain Index (CSI) Differential Emotions Scale (DES) the Positive Aspects of Caregiving Scale The Perceived Stress Scale</p>
<p>Nordheim 2019⁸ (30775982)</p> <p>Germany Medium Explanatory</p>	<p>Couple-based interdisciplinary psychosocial intervention 9 sessions over 10-12 weeks</p>	<p>Usual care</p>	<p>Community-based RCT N=108 dyads</p>	<p>Non-specified dementia NIA criteria</p>	<p>N=108 81 years 39% Female Race NR Education NR</p>	<p>NR</p>	<p>N=108 78 years 61% Female Race NR Education NR</p>	<p>NR</p>	<p>6 months</p>	<p>QOL in Alzheimer's Disease (QOLAD) ADL IADL Older Adults Overprotection Scale (OPSA) Geriatric depression scale (GDS) PLWD social support: (F-SOZU) MMSE</p>	<p>WHOQOL-BREF short form Global QOL subscale Sense of Competence Questionnaire (SCQ) (personal life, performance as a caregiver, satisfaction with PLWD) Caregiver social support: (F-SOZU) Perceived stress GDS</p>
<p>Gitlin 2018¹³ (29192967) Gitlin 2013⁷⁵ (29192967)</p> <p>USA Medium Explanatory</p>	<p>Home-based Tailored Activity Program. 8 sessions with occupational therapists to customize activities to the interests and abilities of the veterans and educate their caregivers about dementia and use of customized activity. Delivery by occupational therapist. 4 months.</p>	<p>Attention control using 8 telephone visits.</p>	<p>Home-based RCT N=160 dyads</p>	<p>Veterans with dementia and their carers. MMSE <23 and documented dementia diagnosis.</p>	<p>N=160 80 years 3% Female 79% White <41% High school diploma or less</p>	<p>NR</p>	<p>N=160 72 years 98% Female 82% White <38% High school diploma or less 87% Spouse or partner</p>	<p>100% Living with PLWD</p>	<p>4 months</p>	<p>Neuropsychiatric Inventory – Clinician (NPI-C) - number of behavioral symptoms Caregiver Assessment of Function and Upset Scale (CAFU) (number of ADL/IADL dependencies, level of dependence) PLWD QOL (caregiver perception of affect)</p>	<p>CES-D ZBI Short Form (12 item) Neuropsychiatric Inventory – Clinician (NPI-C) distress Time spent caregiving (time on ADLs and IADLs, hours on duty, hours on doing things)</p>

Charlesworth 2016 ¹⁸ (27521377) Melunsky 2015 ⁷⁶ (24381218) Charlesworth 2011 ⁷⁷ (21917187) UK Low Explanatory	1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP)) 2) Group reminiscence therapy Remembering Yesterday, Caring Today (RYCT) Once a week for the first 7 weeks followed by monthly meetings for the next 5 months (12 months total)	3) Usual care	Community-based RCT N=291 dyads	Alzheimer's disease Vascular dementia Non-specified dementia DSM-IV	N=291 80 years 53% Female 88% White 75% High school education or less	NR	N=291 67 years 67% Female 89% White 62% High school education or less 64% Spouse or partner	4+ years caregiving (mean)	5 months 1 year	QOL-AD (self and proxy reported) Alzheimer's Disease Cooperative Study-ADL NPI	SF-12 (mental, physical component) Hospital Anxiety and Depression Scale (HADS) (anxiety, Depression) EQ-5D global health visual analog scale Carers of Older People in Europe Index (COPE index) positive aspects Caregiver Quality of Caregiver-PLWD Relationship (QCPR)
Cheng 2016 ¹⁹ (27401052) Cheng 2019a ⁷⁸ (31076215) Cheng 2019b ⁷⁹ (31556447) Cheng 2016 ¹⁹ (28287803) Cheng 2014 ⁸⁰ (24688081) Hong Kong Medium Explanatory	1) Benefit finding intervention (BF) 2) Simplified psychoeducation (SIM-PE)	3. Standard psychoeducation (STD-PE)	Community-based Cluster-randomized RCT N=129 dyads	Alzheimer's disease NINCDS-ADRDA	N=129 NR	NR	N=129 55 years 85% Female Race NR Education 28% Primary or less 57% Secondary 15% Tertiary 27% Spouse or partner 73% Child	2 years caregiving (mean) 13 hours per day 31% Employed	2 months	NR	Hamilton Depression Rating Scale ZBI Caregiver role overload Ryff's Psychological Well-being Scale
Gonyea 2016 ²⁰ (24855313) USA Medium Explanatory	Circulo de Cuidado, a culturally-sensitive, cognitive behavioral (CBT) group intervention. (1) 5 weekly 90-minute group sessions (2) Telephone coaching at week 3, 6, 9 and 12	Psychoeducational (PED) control condition	Community-based RCT N=67 dyads	Possible or probable Alzheimer's Disease Criteria NR	N=291 75 years 63% Female Race NR Education NR	NR	N=291 54 years 51% Female 100% Latino 28% Less than high school education 48% High school certificate or GED 24% Spouse or partner 55% Child	Employment 37% Fulltime 24% Part time 63% Living together 12 hours daily caregiving	3 months 6 months	Neuropsychiatric Inventory-Severity scale (NPI-S) (Spanish)	NPI distress Revised Scale for Caregiving Self-Efficacy (RSCSE) CES-D State Anxiety Inventory-State (STAI-S) (all Spanish)

Laakkonen 2016 ²¹ (27060101) Laakkonen 2013 ⁸¹ (52813196) Laakkonen 2012 ⁸² (22871107) Finland Medium Explanatory	Self-management groups. 4-hour group sessions in a day center once a week for an 8-week period.	Usual care	Clinic-based RCT N=136 dyads	Non-specified dementia	N=136 77 years 27% Female Race NR 45% <8 years education	NR	N=136 74 years 63% Female Race NR 40% <8 years education 100% Spouse or partner	NR	3 months 9 months	PLWD quality of life: 15D Verbal Fluency (VF) Clock Drawing Test (CDT) PLWD Health & social services cost	SF-36 Caregiver quality of life (physical, mental) Spousal Sense of Competence Questionnaire (SCQ) Caregiver mastery Pearlin Mastery Scale (PMS) Caregiver Health & Social services cost
Otero 2015 ⁸³ (25331992) Spain High ROB	Group-based cognitive behavioral intervention for problem solving 5 90-minute weekly sessions	Usual care	Community-based RCT N=173 informal caregivers	Dementia diagnosis (47%)	NR	NR	N=173 54 years Sex NR Race NR 75% primary/higher education 51% Daughter or son	Mean of 10 years caring for PLWD 84% unemployed	1, 3, 6, 12 months	NA	CES-D SCID-CV
Tremont 2015 ²⁸ (25074341) Tremont 2017 ⁸⁴ (28008609) Tremont 2013 ⁸⁵ (23916916) USA Medium Explanatory	Telephone Tracking– Caregiver (FITT-C) received 16 15-60min telephone contacts over 6 months. Provided education, emotional support, directing to resources, encouraging caregivers to attend to their physical, emotional, and social needs, and teaching caregivers strategies to cope with ongoing problems	Telephone Support Control: Received non-directive support through empathic and reflective listening and open-ended questioning without directive strategies, such as education, problem-solving, advice-giving, or task directives.	Academic medical center RCT N=250	NR	NR	NR	N=250 63 years 78% Female 96% White 15. years 51% Spouse or partner 42% Child	80% Living together 3.8 years caregiving	6 months	NR	CESD ZBI RMBPC-RT Reaction Score Family Assessment Device (FAD) Self-Efficacy Questionnaire (symptom management, support services) Positive Aspects of Caregiving (PAC) EuroQoL Visual Analog scale
Martin-Carrasco 2014 ³¹ (24113563) Spain Medium Explanatory	Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care. 7 biweekly 90-120 min sessions over 4 months.	Usual care	Clinic-based RCT N=238 dyads	Alzheimer's disease Vascular dementia Non-specified dementia DSM-IV-TR	N=238 78 years 62% Female Race NR Education 61% Primary school 10% Secondary school 7% College	NR	N=238 62 years 77% Female Race NR Education 48% Primary school 30% Secondary school 13% College 59% Spouse or partner 45% Child	5 years providing care 12 hours per day 25% Working 39% Homemaker 27% Retired	4 months	NR	ZBI General Health Questionnaire 28 item (GHQ-28) SF-12 (all Spanish)

Waldorff 2012 ⁴⁰ (22807076) Phung 2013 (370414872) Denmark Medium Explanatory	DAISY plus support (multifaceted and semi-tailored counselling, education, and support). Up to 7 sessions over 12 months.	Support control group	Home and clinic-based RCT N=330 dyads	Alzheimer's disease Vascular dementia Non-specified dementia Mixed-type DSM-IV NINCDS-ADRDA	N=330 76 years 54% Female Race NR Education 36% NR 26% <3 years 38% ≥3 years	NR	N=330 66 years 67% Female Race NR Education 24% NR 33% <3 years 43% ≥3 years 65% Spouse or partner 27% Child	65% Live with PLWD	1 year	MMSE CSDD EQ-VAS (PLWD and proxy rated) QoL-AD (PLWD and proxy rated) NPI ADSC-ADL	Geriatric depression scale
de Rotrou 2011 ⁴² (20922772) France Medium Explanatory	Psycho Education Program. 12 weekly 2-hour sessions. Content included education, problem-solving techniques and emotion-centered coping strategies, management of PLWD behavior, communication skills, crisis management, resource information and practical advice. Both groups also received antedementia drugs.	Usual care	Memory clinic RCT N=167 dyads	Alzheimer's disease DSM-IV	N=167 79 years 60% Female Race NR 9 Years education	NR	N=167 65 years 68% Female Race NR 11 Years education 57% Spouse or partner 29% Daughter	NR	3 months 6 months	NR	MADRS ZBI Caregiver perception of disease understanding Caregiver coping
Guerra 2011 ⁴³ (20602013) Peru Medium Explanatory	10/66 Caregiver Intervention. Provide education and training on managing problem behaviors. Delivered in 3 modules over 5 30-min weekly sessions.	Waitlist	Home-based RCT N=58	Nonspecified dementia DSM-IV NPI-Q 4.6	N=58 Age 82 72% Female Race NR Education NR	NR	N=58 51 years 88% Female Race NR Education NR 12% Spouse 43% Child	NR	6 months	NPI-Q DEMQOL	ZBI Self-Reporting Questionnaire 20 NPI-Q WHOQOL-BREF

Voight-Radloff 2011 ⁴⁶ (22021760) Voight-Radloff 2011 ⁸⁶ (no PMID) Voight-Radloff 2009 ⁸⁷ (355485226) Germany Low 6-16 weeks Medium 26-52 weeks Explanatory	Community Occupational Therapy in Dementia Program. 10 home visits over 5 weeks by an occupational therapist.	Usual care (one session home consultation)	Home-based RCT N=141 dyads	Alzheimer's disease or mixed-type dementia ICD-10	N=141 78 years 58% Female Race NR 22% High school diploma 77% Less than high school education	NR	N=141 65 years 70% Female Race NR Education NR 57% Spouse or partner 37% Child	NR	6 weeks 16 weeks 26 weeks 52 weeks	Interview for Deterioration in Daily Living Activities in Dementia (IDDD) (performance, initiative) Cornell Scale for Depression in Dementia Perceive, Recall, Plan and Perform System of Task Analysis (PRPP) (independence) Dementia Quality of Life Instrument - overall SF-12 (physical, mental) Number of adverse events Resource Utilization in Dementia (nights in nursing home, night in hospital)	Sense of Competence Questionnaire Center for Epidemiologic Depression Scale Dementia QOL Instrument SF-12 (physical, mental) Resource utilization in Dementia - Basic ADL Care by primary carer (hours per day)
Wang 2011 ⁴⁷ (21752121) USA Medium Explanatory	Family Mutual Support Program in Dementia Care (FMSP-DC). 8 2-hour biweekly sessions over 6 months.	Usual care	Home-based N=80 dyads	Alzheimer's disease DSM-IV	N=80 68 years 81% Female Race NR Education NR	NR	N=80 41 years 65% Female Race NR 73% Secondary education 40% Spouse or partner 38% Child	NR	6 months	MMSE (Chinese) PLWD institutionalizations (number, duration)	Family Caregiving Burden Inventory (FCBI) WHOQOL-BREF Six-item Social Support Questionnaire (SSQ6)
Gitlin 2010 ⁴⁹ (20662955) USA Medium Explanatory	Advancing Caregiving Training (ACT) Health professionals helped target behavioral symptoms of dementia, created plan to help caregivers manage 11 home and phone contacts over 4 months	No treatment or contact	Home-based RCT N=272 dyads	MMSE<24	N=272 82 years 53% Female 70% White 30% Nonwhite Education NR	NR	N=272 66 years 82% Female 70% White 30% Nonwhite Education 7% <High school 26% High school degree 67% Higher education 51% Spouse	4 years caregiving	4, 6 months	Frequency of problem behavior being targeted	ZBI CES-D Perceived Change Index Skill enhancement
Martin-Carrasco 2009 ⁵⁴ (18949763) Spain Medium Explanatory	Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care. 8 sessions over 4 months.	Usual care	Hospital and outpatient clinics N=115 dyads RCT	Alzheimer's disease DSM-IV	N=115 77 years 63% Female Race NR Education NR	NR	N=115 77 years 63% Female Race NR Education NR 55% Spouse or partner 37% Child	12 Hours per day caregiving 3 years caregiving 39% Retired 28% Homemaker 25% Working	4 months 10 months	PLWD Caregiver healthcare and social resources use (number of visits, time spent on medical care)	ZBI Spanish SF-36 GHQ-28 Caregiver healthcare and social resources use (number of visits, time spent on medical care)

Gallagher-Thompson 2008 ⁵⁶ (25067886) USA Medium Explanatory	Coping with Caregiving (CWC). Instruction and practice in small groups to learn specific cognitive and behavioral skills. Weekly 2-hour sessions over 4 months.	Minimal telephone-based control condition (TSC). Empathetic support via 15-20 min telephone calls every 2 weeks.	Home-based RCT N=184	NR	NR	NR	N=184 58 years 100% Female 52% White 48% Latina 13 Years education 33% Spouse or partner	10 Hours per day caregiving 65% Live with PLWD	6 months	NR	CES-D Perceived Stress Scale (PSS-10) RMBPC-CB (both)
Graff 2006 ⁶⁰ (17114212) Graff 2008 ⁸⁸ (18171718) Graff 2007 ⁸⁹ (17895439) Netherlands Medium Explanatory	Community based occupational therapy program. 10 sessions over 5 weeks.	Waitlist for occupational therapy	Home-based RCT N=135 dyads	Mild to moderate dementia Diagnosed with DSM-IV	N=135 78 years 56% Female Race NR Education NR	NR	N=135 63 years 70% Female Race NR Education NR 59% Spouse or partner 32% Child	NR	6 weeks 3 months	Assessment of Motor and Process Skills (AMPS) (process) Interview of Deterioration in Daily Activities in Dementia (IDDD) (performance)	Sense of Competence Questionnaire (SCQ)
Farran 2004 ⁶² (41552352) Farran 2007 ⁹⁰ (175) Farran 2004 ⁹¹ USA Medium Explanatory	Caregiver skill building (CSB) 12 weekly sessions (Five group sessions & 7 individual telephone contacts), 2 group booster sessions & as-needed phone contacts	Information and support group (ISO) Information & support oriented (ISO) group: Routine care-related issues were addressed, but did not include individualized skill enhancement	Home-based RCT N=295	NR	NR	NR	N=295 65 years 83% Female 88% White 62% High school education or less 58% Spouse or partner	NR	18 months	RMPBC Time to institutionalization	CES-D Behavior Management Skill-Revised (BMS-R)
Gallagher-Thompson 2003 ⁶⁶ (12937336) Rabinowitz 2006 ⁹² (16861368) USA Medium Explanatory	Coping With Caregiving psychoeducational program (instruction and practice in small groups to learn specific cognitive and behavioral skills) a) Once a week for the first 10 weeks b) once a month during the "booster phase" for the next 8 months.	Enhanced Support Group condition (guided discussion and empathic listening to develop reciprocal support within the group)	Community-based RCT N=213	NR	NR	NR	N=213 57 years 100% Female 57% White 43% Latino 12 years 38% Spouse or partner 62% Child	5 years (mean)	3 months	NR	CES-D Revised Ways of Coping Checklist (RWCC) (positive, negative) Inventory of Socially Supportive Behaviors (ISSP) (satisfaction, negative interaction) RMBPC
Gitlin 2001 ⁹³ (11220813) USA Medium Explanatory	Home Environmental Intervention Five 90-min home visits by occupational therapists who provided education and physical and social environmental modifications.	Usual care	Home-based RCT N=202 dyads	NR	N=202 78 years 66% Female Race NR Education NR	NR	N=202 61 years 73% Female 74% White 14 years 25% Spouse or partner	100% Living with PLWD	3 months	Functional Independence Measure (modified) (ADL, IADL) MBPC	Caregiver self-efficacy and upset (managing ADL, IADL, behaviors, dependency)

*High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: ADL= Activities of Daily Living; ADSC-ADL= the Alzheimer’s Disease Cooperative Study activities of daily living scale; AMPS= Assessment of Motor and Process Skills; char=characteristics; CAFU= Caregiver Assessment of Function and Upset Scale; CES-D= Center for Epidemiological Studies of Depression scale; CSDD= Cornell Scale for Depression in Dementia; DEMQOL= Dementia Quality of Life measure; DSM-IV= Diagnostic and Statistical Manual of Mental Disorders Fourth edition; EQ-5D= the European Quality of Life-5 Dimensions; EQ-VAS= European quality of life visual analogue scale; FC=formal caregiver; FEERS= Felt Expressed Emotion Rating Scale; GDS== Geriatric Depression Scale; GHQ-28= General Health Questionnaire-28 item version; HADS-T= Hospital Anxiety and Depression Scale Total Score; IADL= Instrumental Activities of Daily Living; IC=informal caregiver; IDDD= Interview of Deterioration in Daily Activities in Dementia; MBPC= Memory and Behavior Problems Checklist; MMSE= Mini-Mental State Examination; N=number; NINCDS-ADRDA= National Institute of Neurological and Communicative Diseases and Stroke/Alzheimer’s Disease and Related Disorders Association; NPI-C= Neuropsychiatric Inventory – Clinician; NPI-Q= The brief form of the Neuropsychiatric Inventory; NPI-S= Neuropsychiatric Inventory-Severity scale (Spanish); PMID=PubMed Identification Number; PLWD=Persons with Dementia; QOL-AD= the Quality of Life in Alzheimer’s Disease; RAS= Relationship Assessment Scale; RMBPC= Revised Memory and Behavior Problems Checklist; RSCSE= Revised Scale for Caregiving Self-Efficacy; RoB=Risk of Bias; RS=Reporting Status; Relatives’ Stress Scale (RSS); SCQ= Sense of Competence Questionnaire; SCID-CV=Structured Clinical Interview for Axis 1 Disorders of the DSM-IV; SES=socioeconomic status; SF-12= Short-Form Health Survey 12; SSQ6= Six-item Social Support Questionnaire; STAI-S= State Anxiety Inventory-State (all Spanish); WHOQOL-BREF= World Health Organization Quality of Life Instruments; ZBI= Zarit Burden Interview; ZBI Spanish SF-36= Spanish validated version of the Zarit Caregiver Burden Interview

Figure E-1. SMD calculated from difference between baseline and FU in intervention minus difference between baseline and FU in comparison group




Table E-3. Intervention components of psychosocial interventions for informal caregivers

Study (PMID) Intervention & Comparison Country	Duration of	Interventionist	Dose/Format	Components
Au 2020 ¹ (no PMID) Intergenerational perspective-taking in-home intervention vs. basic skill building in-home intervention Hong Kong	8 weeks	Therapists	1 40-minute in-home session followed by 7 weekly 35-minute telephone sessions Total hours: 5	Connecting through Caregiving an Intergenerational Perspective-Taking Intervention: discussed the following topics through the lens of intergenerational perspective-taking: self-monitoring mood, scheduling pleasant events, monitoring behavioral problems of PLWD, communicating with PLWD, identifying help, creating implementation plans Basic Skill Building Intervention: discussed the same topics more generally
Au 2019 ² (31279613) Telephone-administered psycho-education with behavioral activation intervention vs. telephone-based psycho-education with general monitoring Hong Kong	20 weeks	Social workers, paraprofessionals	4 weekly individual telephone sessions with social worker followed by 8 biweekly sessions with a paraprofessional, each session lasted approximately 20 minutes Total hours: 4	Behavioral Activation Intervention: Updating recent caregiving situation; Reporting health and needs of CG and CR; Reporting daily routines; Reporting family communication; Activity planning; Review to improve on scheduling; Develop new help-seeking skills; Reviewing to improve communication General Monitoring Intervention: Update caregiving situation: Discuss caregiver's health; Update care giving situation; Discuss care-recipient's needs; Update caregiving situation; Discuss daily/weekly routines; Review support from family/ friends/ agencies;
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Norway	1 year	Nurses, occupational therapists	5 1-hour individual counseling sessions followed by 6 2-hour group meetings Total hours: 17	Psychosocial support intervention: Education about dementia; counselling on problem solving and setting limits; and group sessions Usual care intervention: information on available support
Ghaffari 2019 ⁵ (no PMID) Resilience education group classes vs. usual care Iran	8 weeks	Study researchers	8 weekly 45 minute sessions Total hours: 6	Resilience Education Program: education on Alzheimer's disease, internal supportive factors (self-esteem, optimism, etc), external supportive factors and responsibility, resilience promotion solutions (commitment, control, challenging, coping), problem solving Usual care: received resilience development education after the study finished
Meng 2019 ⁶ (30884961) Telephone based cognitive behavioral therapy vs. in-person cognitive behavioral therapy United States	12 weeks	NR	Telephone based cognitive therapy Dose not reported	Cognitive Behavioral Therapy
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control United States	6 weeks	Facilitator discipline not specified	6 weekly live online sessions using a computer tablet	Life Enhancing Activities for Family Caregivers (LEAF): positive emotion regulation skills: noticing positive events; capitalizing on positive events; gratitude; mindfulness, positive reappraisal; personal strengths; setting attainable goals; acts of kindness Waitlist control: completed daily emotion survey for 6 weeks then crossed over to LEAF program.

Nordheim 2019 ⁸ (30775982) Couple-based Interdisciplinary Psychosocial Intervention vs. 1-2 hours of memory care consultation Germany	10-12 weeks	Psychotherapist, social worker	7 home-based couples' sessions with two additional telephone sessions	Sessions included information about dementia, couple communication training, coping and problem-solving strategies, network and activity analysis, counseling for living space adaptations, and relaxation techniques Manualized program
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits United States Veterans	NR	Occupational therapists	8 in-home sessions	Customize activities to the interests and abilities of the veterans and educate their caregivers about dementia and use of customized activity.
Voigt-Radloff 2017 ⁹⁴ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Germany	5 weeks	Occupational therapists	10 1-hour in-home sessions Total hours: 10	Evidence-based manual Educating PLWD in the performance of simplified daily activities and in the use of aids to compensate for cognitive decline; and educating carers in coping with behavior of the PLWD and in giving supervision to the PLWD. Content included: (1) The PLWD preferences and history of daily activities; (2) their ability to perform activities and to use compensatory strategies within the familiar environment; (3) the possibilities of modifying the PLWD home; (4) the carer's activity preferences, problems in care giving, coping strategies and abilities to supervise; and (5) the interaction between carer and PLWD. Carer received practical and emotional support and was coached in effective supervision, problem-solving and coping strategies by means of cognitive-behavioral interventions.
Charlesworth 2016 ¹⁸ (27521377) Carer Supporter Programme (CSP) program vs. Remembering Yesterday Caring Today (RYCT) program England	8 months 10 months	CSP: Volunteer carer supporters RYCT: Unspecified	CSP: 12 one-to-one peer support 1-hour sessions occurring weekly, followed by twice monthly meetings for the next 5 months - Total hours: 22 RYCT: 12 weekly 2-hour sessions followed by 7 months of monthly sessions – 19 sessions over 10 months Total hours: 38	CSP: One-to-One Peer support. Volunteer carer supporters asked to listen, encourage and give moral support. Though they could also signpost to resources and services, we instructed them not to offer tangible support, respite or direct advice. RYCT: Group Reminiscence
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention vs. simplified lecture only psychoeducation vs. standard psychoeducation Hong Kong	8 weeks	Research Assistants with undergraduate degree in psychology or related field.	8 weekly 2-hour group sessions of 7-11 caregivers Total hours: 16	Benefit finding intervention: Benefit finding, positive reappraisal cognitive techniques; Information on dementia, communication skills, and stress management; causes and coping strategies for BPSD; homebased activities; Skills for helping with ADLs, creating an appropriate home environment for the care recipient, and community resources; goal setting Simplified lecture only psychoeducation Intervention: Lectures and discussions, no practical elements Standard psychoeducation intervention: same benefit finding intervention components excluding dysfunctional thoughts, maintenance cycle, and positive reappraisal
Gonyea 2016 ²⁰ (24855313) Circulo de Cuidado, a culturally-sensitive, cognitive behavioral therapy group intervention vs. psychoeducational control condition United States	17 weeks	MSW Social workers bilingual in Spanish	5 weekly 90-minute small group sessions, followed by 10-15-minute individual telephone coaching sessions at 3, 6, 9- and 12-weeks post-group Total hours: 5.5	Culturally-sensitive Cognitive Behavioral Group Intervention; antecedents-behaviors-consequences (A-B-C) problem solving approach to behavior change; relaxation techniques or exercises; increasing engagement in pleasant activities for caregiver and PLWD; improving communication with PLWD; stress management; relaxation techniques; weekly at-home assignments Psychoeducational group control intervention: dementia education; finding community resources; working with physicians; home safety; communication in context of dementia
Laakkonen 2016 ²¹ (27060101) Self-management group rehabilitation Finland	8 weeks	Nurses, Occupational therapists, Physiotherapists	8 weekly 4-hour group sessions Total hours: 32	Group sessions aim to enhance self-efficacy and problem-solving skills and to provide peer support. Self-management capabilities such as problem-solving skills, self-efficacy, and mastery are built gradually during the intervention Usual preferred topics were dementia, medication, nutrition, active lifestyle, marital relationship, caregiving, available social and health services, and advance care planning.

Otero 2015 ⁸³ (25331992) Cognitive behavioral problem solving vs. usual care Spain	5 weeks	Psychologists	5 1.5-hour group sessions occurring weekly Total hours: 7.5	Cognitive behavioral problem solving group-based program including: problem solving module, goal setting, decision making and planning, and relapse prevention.
Tremont 2015 ²⁸ (25074341) Family Intervention Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) attention control. United States	6 months	Master's degree prepared therapists with training in dementia and caregiving	16 telephone contacts averaging 36 minutes each Total hours: 9.6	Providing dementia education, emotional support, directing caregivers to appropriate resources, encouraging caregivers to attend to their physical, emotional, and social needs, and teaching caregivers' strategies to cope with ongoing problems.
Livingston 2014 ³⁰ (25300037) START (STrategies for RelaTives vs. usual care United Kingdom	unspecified	Supervised psychology graduates	8 home-based individual sessions	REACH Palo Alto Coping with Caregiving program modified with fewer sessions and home-based sessions. Coping with Caregiving Manual Individual sessions Topics: Psychoeducation about dementia, carer stress and understanding behaviors, discussion of behaviors or situations the carer finds difficult, incorporating behavioral management, identifying and changing unhelpful thoughts, assertive communication, increasing communication, acceptance, accessing emotional support and positive reframing. future planning, pleasant activities, maintaining skills over time, & stress reduction techniques
Martin-Carrasco 2014 ³¹ (24113563) Group psychoeducational intervention (PIP) vs. usual care Spain	14 weeks	Therapist (unspecified training)	7 90-120-minute group sessions administered biweekly Total hours: 10.5-14 hours	REACH Palo Alto Coping with Caregiving program modified for Latinos living in Spain and group sessions. Modified Martin-Carrasco 2014 intervention from individual to group sessions. Caregivers received standardized information about the clinical course of dementia and were trained on cognitive and behavioral skills and relaxation techniques to increase their care abilities, communicative skills, and emotional control. 7 modules related with strain and well-being, changing maladaptive behaviors, negative thoughts, ways to communicate, planning the future, and planning enjoyable activities.
Waldorff 2012 ⁴⁰ (22807076) Danish Alzheimer's Study (DAISY) vs. control support Denmark	4 months	Nurse trained in counselling approach	Up to 7 counselling sessions including: 2 sessions with the PLWD and care giver; 2 sessions with the PLWD alone; 2 sessions with the care giver alone; and an optional network session with the PLWD, caregiver, and family network. 5 2-hour courses for caregiver 5 2-hour courses for PLWD 5-8 telephone contacts with study coordinator to discuss counselling session and course information.	The counselling was based on a philosophical approach in which each PLWD or care giver was given the possibility of expressing his or her life story and what is of personal importance and of great value. The counsellor offered the participants guidance with common decision making, advice, and activities that help the participants construct a meaningful life. Caregiver courses: formalized education program on Alzheimer's disease, also supported by printed handouts with information on specific topics. PLWD courses: information on key issues of the disease and its consequences supported by printed handouts with information on specific topics.
de Rotrou 2011 ⁴² (20922772) Psych-Educational Program vs. waitlist control group France	12 weeks	Psychologists were group leaders Disease education provided by health professional (geriatrician, psychiatrist, psychologist, social worker, speech therapist, occupational therapist)	12 weekly 2-hour group sessions of 6-10 caregivers Total hours: 24	Education on dementia, problem-solving techniques and emotion-centered coping strategies, management of PLWD behavior, communication skills, crisis management, resource information and practical advice, ecological stimulation - stimulate their relative in daily activities and social situations Waitlist control: PLWD in both groups received cholinesterase-inhibitor pharmacotherapy
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Peru	5 weeks	Junior Psychologists, social workers	5 weekly 30-minute sessions over 5 weeks Total hours: 2.5	Helping carers to care: dementia education, training on managing problem behaviors
Wang 2011 ⁴⁷ (21752121) Family Mutual Support Programme in Dementia Care (FMSP-DC) vs. usual care China	6 months	Psychiatric nurse	8 biweekly 2-hour group sessions Family led Total hours: 16	Community based program addressing (1) information about client's illness, prognosis and current treatment and care; (2) development of social relationships with close relatives and friends and thus a satisfactory extended social support network; (3) sharing and adaptation of emotional impacts of caregiving; (4) learning about self-care and motivation; (5) improvement of interpersonal relationships with family members and the client; (6) establishing support from the community groups and health-care resources; and (7) improvement of problem solving skills in family care.

Gitlin 2010 ⁴⁹ (20662955) Advancing Caregiving Training (ACT) vs. no treatment United States	16-week active phase followed by 8-week maintenance phase	Occupational therapists Advanced Practice Nurses	During active phase, up to 9 OT sessions & 2 nursing sessions (one home visit, one telephone visit) During maintenance phase: 3 telephone calls from occupational therapist Total hours: 16	Advanced Caregiver Training (ACT): identification of PLWD unmet needs, discomfort, & medical problems, caregiver skill building, problem solving, and communication building, caregiver stress reduction & self-care techniques, environmental modifications & provision of low-cost assistive devices No treatment control group
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program (PIP) vs. usual care Spain	4 months	Clinical psychologist, nurse or social worker	8 individual 90-minute sessions at 1-2-week intervals over 4 months Total hours: 12	REACH Palo Alto Coping with Caregiving program modified for Latinos living in Spain. Structured psycho-educational-type psychosocial intervention program Skills training: (a) to help the caregiver control tension and stress deriving from caregiving; (b) to teach the caregiver different strategies for handling their relative's behavioral problems; and (c) to increase their satisfaction with life.
Gallagher-Thompson 2008 ⁵⁶ (25067886) Coping with Caregiving vs. Minimal telephone-based control condition United States	13-16 weeks	Post-doctoral fellows or advanced graduate students in psychology or related fields who had relevant bilingual/bicultural backgrounds. 2 Interventionists led each session	Weekly 2-hour group sessions of 4-8 female caregivers. Total hours: 26-32	REACH Palo Alto Coping with Caregiving Program updated (Gallagher-Thompson 2003 (12937336)) Coping with Caregiving: cognitive behavioral therapy to develop cognitive and behavioral skills to reduce stress and depression; education about dementia, helpful techniques for managing care recipient's problem behaviors; skills to take better care of the caregiver, including changing unhelpful thoughts, increasing assertive communication, and identifying everyday pleasant activities (both for self and shared with care recipient); planning for the care-recipient's future needs; develop action plan for future Minimal telephone-based control intervention: Individual sessions of empathic support (7 biweekly calls lasting 15-20 minutes)
Graff 2006 ⁶⁰ (17114212) Community based occupational therapy program vs. Waitlist for occupational therapy The Netherlands	5 weeks	Occupational therapists	10 1-hour sessions with caregiver and PLWD Total hours: 10	Cognitive and behavioral interventions, to train PLWD in the use of aids to compensate for cognitive decline and care givers in coping behaviors and supervision Diagnostics and goal defining, PLWD and primary care givers learnt to choose and prioritize meaningful activities they wanted to improve optimize these compensatory and environmental strategies to improve their performance of daily activities
Farran 2004 ⁶² (41552352) Caregiver Skill (CSB) intervention vs. Information and Support Oriented (ISO) Intervention United States	1 year	Nurses, social workers	12 weekly sessions consisting of 5 group sessions and 7 individual telephone contact sessions; Followed with 2 booster sessions at 6 and 12 months and as needed telephone sessions.	CSB: Improve caregiver skill in managing behavioral symptoms. 5 major topics: introduction to CSB conceptual model, potential causes and contributing factors to behavioral symptoms, prevention of behavior during cares, management of restless behaviors, management of hallucinations and delusions. ISO: Standardized implementation of psychoeducational methods, generalized information and support, no individualized behavioral symptom management. 5 topics: understanding dementia, dealing with difficult behaviors, providing personal care, caring for the caregivers, and finding/using help.
Gallagher-Thompson 2003 ⁶⁶ (12937336) Coping with Caregiving vs. Enhanced support group condition United States	120 weeks (10 months)	Psychologists, social workers or other helping professionals, and predoctoral or postdoctoral psychology and social work graduate students	First 10 weeks used 2-hour group sessions of 7-9 female caregivers followed by 8 monthly booster sessions. Total hours: 28	Both programs were tailored to be sensitive to the cultural concerns of Anglo and Latino caregivers, and they were delivered in either English or Spanish by trained interventionists. REACH Palo Alto Coping with Caregiving Group: dementia education, cognitive-behavioral mood management skills Enhanced Support Group: dementia education, guided discussion, empathic listening
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care United States	3 months	Occupational therapists	5 90-minute home visits occurring every other month Total hours: 7.5	Education and physical and social environmental modifications. Dementia education; Education on daily care needs specified by caregiver; role-play, direct observation, and interviewing to problem solve; Problem solving; Environmental simplification and task breakdown strategies

Abbreviations: PLWD=person with dementia

Table E-4. PLWD outcomes summary low and medium risk of bias studies: psychosocial interventions for caregiver well-being

Study (PMID) Comparison RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Medium Explanatory Norway	Between-group differences as estimated regression coefficient (95% CI) PLWD Cognition: MMSE 12 weeks	-0.78 (-1.78 to 0.22)	NR	NR	0.12
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Medium Explanatory Norway	Between-group differences as estimated regression coefficient (95% CI) PLWD neuropsychiatric symptoms: Neuropsychiatric Inventory (NPI-Q) 12 weeks	-0.29 (-0.83 to 0.26)	NR	NR	0.31
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Medium Explanatory Norway	Between-group differences as estimated regression coefficient (95% CI) PLWD function IADL: Lawton & Brody's IADL 12 weeks	0.48 (-0.73 to 1.69)	NR	NR	0.44
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Medium Explanatory Norway	Between-group differences as estimated regression coefficient (95% CI) PLWD depression: Cornell scale for depression in dementia. 12 weeks	0.34 (-0.81 to 1.49)	NR	NR	0.57
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Medium Explanatory Norway	N (percent) PLWD nursing home placement 12 months	NR	24 (22.9%)	23 (22.3%)	NR
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) PLWD cognition: Dementia Severity Rating Scale 6 weeks	-0.03 (-0.33, 0.27)	23.12 (1.07) baseline 23.42 (1.09) 6 weeks	22.49 (1.09) baseline 23.06 (1.10) 6 weeks	Interaction 0.71
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months PLWD quality of life: Quality of Life in Alzheimer's Disease (QOLAD) 6 months	Cohens d = -0.4	36.7 (0.8) baseline 34.7 (0.8) 6 months	37.2 (0.8) baseline 36.4 (0.9) 6 months	Interaction NS
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months PLWD ADL: 6 months	Cohens d = 0.1	90.5 (2.1) baseline 86.3 (2.3) 6 months	87.1 (2.2) baseline 83.7 (2.4) 6 months	Interaction $p > 0.05$
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months PLWD IADL: 6 months	Cohens d = 0.1	5.3 (0.3) baseline 4.2 (0.3) 6 months	4.9 (0.3) baseline 3.9 (0.3) 6 months	Interaction $p > 0.05$
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months PLWD function: Older Adults Overprotection Scale (OPSA) 6 months	Cohens d = 0.1	39.5 (0.8) baseline 39.1 (0.8) 6 months	38.8 (0.8) baseline 38.8 (0.9) 6 months	Interaction NS
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months PLWD depressive symptoms: Geriatric depression scale (GDS) 6 months	Cohens d = 0.0	5.6 (0.3) baseline 5.9 (0.3) 6 months	5.4 (0.3) baseline 5.0 (0.3) 6 months	Interaction NS

Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months PLWD cognition: MMSE 6 months	Cohens d = 0.4	23.0 (0.6) baseline 21.6 (0.7) 6 months	22.7 (0.7) baseline 19.6 (0.8) 6 months	Interaction p < 0.05
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months PLWD social support: (F-SOZU) 6 months	Cohens d = -0.1	54.8 (1.4) baseline 54.0 (1.3) 6 months	54.1 (1.4) baseline 55.1 (1.4) 6 months	Interaction NS
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) PLWD neuropsychiatric symptoms: Neuropsychiatric Inventory – Clinician (NPI-C) -number of behavioral symptoms 4 months	-0.68 (-1.23; -0.13)	NR	NR	0.02
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) PLWD neuropsychiatric symptoms: Neuropsychiatric Inventory – Clinician (NPI-C) 4 months	-24.3 (-45.6; -3.1)	NR	NR	0.02
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) PLWD function ADL/IADL: Caregiver Assessment of Function and Upset Scale (CAFU) - number of ADL/IADL dependencies subscale 4 months	-0.80 (-1.41; -0.20)	NR	NR	0.009
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) PLWD function ADL/IADL: Caregiver Assessment of Function and Upset Scale (CAFU) - level of ADL/IADL dependence (total) 4 months	4.09 (1.06; 7.13)	NR	NR	0.009
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) PLWD QOL: affect (caregiver perception) 4 months	0.47 (-1.37; 0.43)	NR	NR	0.30
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) PLWD function ADL: Caregiver Assessment of Function and Upset Scale (CAFU) - number of ADL dependencies subscale 4 months	-0.61 (-1.08; -0.14)	NR	NR	0.01
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) PLWD function IADL: Caregiver Assessment of Function and Upset Scale (CAFU) - number of IADL dependencies subscale 4 months	-0.25 (-0.54; 0.04)	NR	NR	0.09
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) PLWD function ADL: Caregiver Assessment of Function and Upset Scale (CAFU) - level of ADL dependence subscale 4 months	2.37 (0.32; 4.42)	NR	NR	0.02
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI)PLWD function IADL: Caregiver Assessment of Function and Upset Scale (CAFU) - level of IADL dependence subscale 4 months	1.57 (0.05; 3.08)	NR	NR	0.04
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) PLWD health care use: Number of monthly emergency department visits during intervention 6 months	NR	0.08 (0.16)	0.08 (0.17)	0.83

Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) PLWD health care use: Number of monthly doctor visits during intervention 6 months	NR	1.32 (1.35)	1.30 (1.05)	0.88
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) PLWD community support use: Number of community support services used in final month of intervention 6 months	NR	6.79 (13.72)	5.11 (10.68)	0.34
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Frequency (%) PLWD nursing home admission: number of PLWD admitted to nursing home during the intervention	12 (6%)	NR	NR	0.70 (no group difference)
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups PLWD Health related quality of life: QOL-AD self-reported 12 months	-0.162 (-1.87 to 1.54)	Means reported	Means reported	0.85
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups PLWD Health related quality of life: QOL-AD self-reported 12 months	0.702 (-1.05 to 2.45)			0.43
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups PLWD Health related quality of life: QOL-AD proxy reported 12 months	0.050 (-1.21 to 1.31)			0.94
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups PLWD Health related quality of life: QOL-AD proxy reported 12 months	0.0660 (-1.17 to 1.30)			0.92
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups PLWD function: Alzheimer's Disease Cooperative Study—Activities of Daily Living 12 months	-2.18 (-6.07 to 1.71)			0.27
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups PLWD function: Alzheimer's Disease Cooperative Study—Activities of Daily Living 12 months	-2.45 (-5.95 to 1.06)			0.17

Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups PLWD neuropsychiatric symptoms: neuropsychiatric inventory (NPI) 12 months	-1.20 (-6.64 to 4.23)			0.66
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups PLWD neuropsychiatric symptoms: neuropsychiatric inventory (NPI) 12 months	0.236 (-4.83 to 5.30)			0.93
Gonyea 2016 ²⁰ (24855313) Circulo de Cuidado, a culturally-sensitive, cognitive behavioral (CBT) group intervention vs. psychoeducational (PED) control condition Medium Explanatory USA	Mean (SD) PLWD neuropsychiatric symptoms: Spanish version of the Neuropsychiatric Inventory-Severity scale (NPI-S) 12 week 24 week	NR	21.70 (6.90) baseline 20.67 (6.94) 12 weeks 20.52 (6.98) 24 weeks	21.76 (7.98) baseline 22.03 (9.10) 12 weeks 22.09 (8.46) 24 weeks	Repeated measures ANCOVA <0.001
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) PLWD quality of life: 15D 9 months	d=NR	-0.03 (-0.0 to -0.01)	-0.04 (-0.05 to -0.02)	0.55
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) PLWD cognition: Verbal Fluency (VF) 3 months	d=NR	0.26 (-0.38 to 0.91)	-0.60 (-1.26 to 0.07)	0.07
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) PLWD cognition: Verbal Fluency (VF) 9 months	d=NR	-0.38 (-1.03 to 0.27)	-1.60 (-2.26 to -0.94)	0.01
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) PLWD cognition: Clock Drawing Test (CDT) 3 months	d=NR	0.22 (-0.13 to 0.58)	-0.18 (-0.55 to 0.18)	0.12
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) PLWD cognition: Clock Drawing Test (CDT) 9 months	d = 0.28	-0.11 (-0.46 to 0.25)	-0.65 (-1.02 to -0.30)	0.03
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean cost in € per year (95% CI) PLWD Health & social services cost: 24 months	-436 (-4,986; 4,115)	8,947 (6,455–12,415)	9,383 (6,398–13,374)	.035
Livingston 2014 ³⁰ (25300037) START (STrategies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	Adjusted mean difference (95% CI) PLWD QOL: (QOL-AD) 2 years	0.17 (-1.37 to 1.70)	NR	NR	NR

Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD QOL: European quality of life visual analogue scale (EQ-VAS) score 12 months	NR	-0.31 (-3.41 to 2.79)	0.20 (-2.31 to 2.71)	0.7799
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD cognition: MMSE 12 months	NR	-1.55 (-2.19 to -0.91)	-1.80 (-2.56 to -1.04)	0.6502
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD depression: Cornell depression scale 12 months	NR	-0.20 (-1.04 to 0.64)	1.38 (0.49 to 2.27)	0.0103
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD QOL: proxy rated European quality of life visual analogue scale (EQ-VAS) (proxy rated) score 12 months	NR	-3.88 (-7.38 to -0.37)	-6.83 (-10.10 to -3.56)	0.2308
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD QOL: European quality of life visual analogue scale EQ VAS (PLWD rated) 12 months	NR	0.73 (-2.41 to 3.88)	-4.20 (-8.29 to -0.11)	0.0667
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD WOL: Quality of life scale for Alzheimer's disease QoL-AD (PLWD rated) 12 months	NR	-0.82 (-1.77 to 0.12)	-1.04 (-2.04 to -0.04)	0.7213
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD QOL: quality of life scale for Alzheimer's disease QoL-AD (proxy rated) 12 months	NR	-0.82 (-1.77 to 0.12)	-2.61 (-3.58 to -1.64)	0.0013
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD neuropsychiatric symptoms: neuropsychiatric inventory questionnaire NPI-Q 12 months	NR	0.85 (0.16 to 1.53)	0.43 (-0.26 to 1.13)	0.4045
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) PLWD function: Alzheimer's Disease Cooperative Study activities of daily living scale. ADSC-ADL 12 months	NR	-6.39 (-8.51 to -4.27)	-4.63 (-6.92 to -2.53)	0.2689
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Standardized Mean Difference (95% CI) PLWD neuropsychiatric symptoms: Neuropsychiatric Inventory brief form (NPI-Q) 6 months	-0.05 (-0.59 to .52)	NR	NR	NR

Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Mean change (SD) PLWD neuropsychiatric symptoms: Neuropsychiatric Inventory brief form (NPI-Q) 6 months	NR	-1.7 (3.3)	-1.6 (206)	0.87
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) PLWD function: Interview for Deterioration in Daily Living Activities in Dementia (IDDD) – performance subscale 52 weeks	-2.4 (-7.1; 2.3)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) PLWD function: Interview for Deterioration in Daily Living Activities in Dementia (IDDD) – initiative subscale 52 weeks	-1.0 (-5.0; 3.0)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) PLWD depression: Cornell Scale for Depression in Dementia 52 weeks	-2.0 (-5.1; 1.0)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) PLWD function: Perceive, Recall, Plan and Perform System of Task Analysis (PRPP)– independence subscale 26 weeks	3.3 (-8.3; 14.9)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) PLWD QOL: Dementia Quality of Life Instrument - overall 26 weeks	0.2 (-0.1; 0.5)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) PLWD QOL: Short-Form 12 Health Survey Questionnaire -Physical component 26 weeks	3.0 (-1.6; 7.6)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) PLWD QOL: Short-Form 12 Health Survey Questionnaire – mental component 26 weeks	-0.8 (-4.9; 3.3)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	N PLWD harms: number of adverse events 52 weeks	NR	0	0	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	N PLWD nursing home: Resource Utilisation in Dementia, nights in nursing home (except respite care) 52 weeks	NR	2	1	NS

Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	N PLWD hospitalization: Resource Utilisation in Dementia, nights in hospital 52 weeks	NR	14 PLWD were admitted to hospital for an average of 15 nights	10 PLWD were admitted to hospital for an average of 18 nights.	0.1
Wang 2011 ⁴⁷ (21752121) Family Mutual Support Program in Dementia Care (FMSP-DC) vs. usual care Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months PLWD cognition: MMSE (Chinese) 6 months	NR	17.5 (4.7) baseline 18.1 (4.0) 6 months	17.3 (3.9) baseline 17.2 (4.1) 6 months	NS
Wang 2011 ⁴⁷ (21752121) Family Mutual Support Program in Dementia Care (FMSP-DC) vs. usual care Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months PLWD institutionalization: Number 6 months	NR	5.1 (0.9) baseline 3.8 (1.0) 6 months	5.5 (1.2) baseline 5.2 (1.3) 6 months	<0.01
Wang 2011 ⁴⁷ (21752121) Family Mutual Support Program in Dementia Care (FMSP-DC) vs. usual care Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months PLWD institutionalization: duration 6 months	NR	13.8 (4.0) baseline 10.1 (5.1) 6 months	14.2 (3.8) baseline 17.9 (7.1) 6 months	<0.001
Gitlin 2010 ⁴⁹ (20662955) Advancing Caregiver Training (ACT) vs. no treatment control Medium Explanatory USA	Percentage reporting improvement in problem behavior PLWD neuropsychiatric symptoms: number of problem behaviors 4 months	NR	67.5	45.8	0.002
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Frequency (percent) PLWD Caregiver healthcare and social resources use: number of visits 4 months	NR	0 visits: 29/44 (66%) 1-2 visits: 13/44 (36%)	0 visits: 25/38 (66%) 1-2 visits: 13/38 (36%)	NR (NS)
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) PLWD Caregiver healthcare and social resources use: time spent on medical care 4 months	NR	28.5 (25.5)	21.8 (11.0)	NR
Graff 2006 ⁶⁰ (17114212) Community based occupational therapy program vs. Waitlist for occupational therapy Medium Explanatory	Adjusted Mean difference (95% CI) Effect size Number needed to treat PLWD function: Assessment of Motor and Process Skills (AMPS) – process subscale 6 weeks	1.5 (1.3; 1.7) 2.5 1.3 (1.2 to 1.4) NNT	NR	NR	<0.0001
Graff 2006 ⁶⁰ (17114212) Community based occupational therapy program vs. Waitlist for occupational therapy Medium Explanatory	Adjusted Mean difference (95% CI) PLWD function: Interview of Deterioration in Daily Activities in Dementia (IDDD) – performance subscale 6 weeks	- 11.7 (-13.6; -9.7) 2.3 1.5 (1.4 to 1.6) NNT	NR	NR	<0.0001
Graff 2006 ⁶⁰ (17114212) Graff 2007 ⁸⁹ (17895439) Community based occupational therapy program vs. Waitlist for occupational therapy Medium Explanatory	Adjusted treatment difference (95% CI) Effect size Mean (SD) PLWD quality of life: Dementia Quality of Life Instrument 6 weeks	0.8 (0.6 to 1.1) 1.3 (effect size)	4.0 (0.6)	3.1 (0.8)	< 0.0001
Graff 2006 ⁶⁰ (17114212) Community based occupational therapy program vs. Waitlist for occupational therapy Medium Explanatory	Adjusted treatment difference (95% CI) Effect size Mean (SD) PLWD depression: Cornell scale for depression 6 weeks	-2.8 (-4.3 to -1.3) 0.7 effect size	6.5 (5.3)	9.2 (6.4)	< 0.0001

Farran 2004 ⁶² (41552352) Caregiver skill building (CSB) vs. Information and support group (ISO) Medium Explanatory USA	Regression coefficient (SE) PLWD neuropsychiatric symptoms: Revised Memory and Problem Behavior Checklist (RMPBC) 18 months	-0.046 (0.035)	NR	NR	0.193
Farran 2004 ⁶² (41552352) Caregiver skill building (CSB) vs. Information and support group (ISO) Medium Explanatory USA	Log rank from lifetable PLWD Time to institutionalization 18 months	0.11, df=1	NR	NR	p>0.75
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care Medium Explanatory USA	Adjusted mean difference (95% CI) Mean (SD) baseline Mean (SD) 3 month PLWD function ADL: Functional Independence Measure (modified) ADL subscale 3 months	-0.06 (-0.30 to 0.18)	2.93 (1.49) baseline 3.24 (1.59) 3 months	3.23 (1.36) baseline 3.57 (1.38) 3 months	0.599
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care Medium Explanatory USA	Adjusted mean difference (95% CI) Mean (SD) baseline Mean (SD) 3 month PLWD function IADL: Functional Independence Measure (modified) IADL subscale 3 months	-0.13 (-0.24 to -0.01)	5.43 (.62) baseline 5.54 (.60) 3 months	5.56 (.50) baseline 5.75 (.36) 3 months	0.030
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care Medium Explanatory USA	Adjusted mean difference (95% CI) Mean (SD) baseline Mean (SD) 3 month PLWD neuropsychiatric symptoms: Memory and Behavior Problems Checklist (MBPC) 3 months	1.85 (-0.42 to 4.13)	20.25 (5.39) baseline 17.20 (7.73) 3 months	18.74 (6.31) baseline 14.43 (9.82) 3 months	0.110

Abbreviations: ADL= Activities of Daily Living; CI= Confidence Interval; IADL= Instrumental Activities of Daily Living; MMSE= Mini-Mental State Examination; NR=Not Reported; PMID=PubMed Identification Number; QOL-AD= the Quality of Life in Alzheimer's; SD= Standard Deviation; SE= Standard Error

Table E-5. Caregiver outcomes summary low and medium risk of bias studies: psychosocial interventions for caregiver well-being

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Au 2020 ¹ (no PMID) Intergenerational perspective-taking in-home intervention vs. basic skill building in-home intervention Low Explanatory Hong Kong	Mean (SD) Caregiver life satisfaction: Satisfaction With Life Scale 8 weeks	NR	17.20 (3.79) baseline 23.46 (5.90) 8 weeks	18.09 (4.98) baseline 19.66 (5.62) 8 weeks	p<0.001
Au 2020 ¹ (no PMID) Intergenerational perspective-taking in-home intervention vs. basic skill building in-home intervention Low Explanatory Hong Kong	Mean (SD) Caregiver depression: Center for Epidemiological Studies-Depression (CESD) 8 weeks	NR	15.97 (4.02) baseline 10.76 (4.34) 8 weeks	15.02 (3.53) baseline 13.34 (3.76) 8 weeks	p<0.001
Au 2020 ¹ (no PMID) Intergenerational perspective-taking in-home intervention vs. basic skill building in-home intervention Low Explanatory Hong Kong	Mean (SD) Caregiver burden: Zarit Burden Interview 8 weeks	NR	23.32 (7.49) baseline 16.30 (7.14) 8 weeks	22.83 (6.66) baseline 21.14 (6.14) 8 weeks	p<0.001
Au 2019 ² (31279613) Telephone-administered psycho-education with behavioral activation intervention vs. telephone-based psycho-education with general monitoring	Cohen's d (95% CI) Mean (SD) Caregiver depression: Center for Epidemiological Studies-	-0.66 (-1.04, -0.74)	13.29 (7.63) baseline 8.62 (8.53) 20 months	13.92 (9.05) baseline 15.55 (11.35) 20 months	p<0.001

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Low Explanatory Hong Kong	Depression (CESD) 20 weeks				
Au 2019 ² (31279613) Telephone-administered psycho-education with behavioral activation intervention vs. telephone-based psycho-education with general monitoring Low Explanatory Hong Kong	Cohen's d (95% CI) Mean (SD) Caregiver burden: Zarit Burden Interview 20 weeks	-1.14 (-1.54, -0.75)	32.26 (17.43) baseline 21.82 (14.06) 20 months	29.47 (17.06) baseline 37.47 (17.28) 20 months	p<0.001
Au 2019 ² (31279613) Telephone-administered psycho-education with behavioral activation intervention vs. telephone-based psycho-education with general monitoring Low Explanatory Hong Kong	Cohen's d (95% CI) Mean (SD) Caregiver satisfaction with relationship with care recipient: Relationship Assessment Scale (RAS) 20 weeks	0.44 (0.82, 0.07)	27.78 (5.90) baseline 30.05 (4.79) 20 months	27.41 (5.86) baseline 27.60 (6.08) 20 months	p = 0.01
Au 2019 ² (31279613) Telephone-administered psycho-education with behavioral activation intervention vs. telephone-based psycho-education with general monitoring Low Explanatory Hong Kong	Cohen's d (95% CI) Mean (SD) Caregiver : self-efficacy for controlling upsetting thoughts 5 item 20 weeks	0.86 (1.24, 0.46)	36.08 (8.03) baseline 40.15 (9.02) 20 months	33.58 (10.03) baseline 34.05 (10.07) 20 months	p<0.01
Kunik 2020 ³ (32115311) Unmet Needs Psychosocial vs education and information support Low Explanatory United States	Cohen's d (95% CI) Mean (SD) 3 months	-6.9 (2.32, 0.78)	NR	NR	p=0.27
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Medium Explanatory Norway	Between-group differences as estimated regression coefficient (95% CI) Caregiver stress: Relatives' Stress Scale (RSS) 12 months	1.12 (-1.23 to 3.48)	NR	NR	0.35
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Medium Explanatory Norway	Between-group differences as estimated regression coefficient (95% CI) Caregiver depression: Geriatric Depression Scale (GDS) 12 months	1.70 (0.41 to 3.0)	NR	NR	0.01
Bjorge 2019 ⁴ (31651321) Psychosocial intervention vs. usual care Medium Explanatory Norway	Between-group differences as estimated regression coefficient (95% CI) Caregiver emotions: Felt Expressed Emotion Rating Scale (FEERS) 12 months	0.36 (-0.62 to 1.34)	NR	NR	0.47
Ghaffari 2019 ⁵ (no PMID) Resilience education group classes vs. usual care Medium Explanatory Iran	Mean difference (SD) Quality of life: General Health Questionnaire 28 item (GHQ-28) 8 weeks		39.9 (8.4) baseline 16.08 (3.7) 8 weeks	32.0 (7.0) baseline 32.9 (7.2) 8 weeks	p<0.001
Meng 2019 ⁶ (30884961) Telephone based cognitive behavioral therapy vs. in-person cognitive behavioral therapy Medium Explanatory USA	Mean (SD) Caregiver health service use: physician office visits 3 months	NR	3.6 (2.9) baseline 2.9 (2.6) 3 months	2.4 (3.1) baseline 3.6 (4.3) 3 months	NS

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Meng 2019 ⁶ (30884961) Telephone based cognitive behavioral therapy vs. in-person cognitive behavioral therapy Medium Explanatory USA	Mean (SD) Caregiver health service use: mental health visits 3 months	NR	1.7 (0.6) baseline 2 (0) 3 months	0.1 (0.3) baseline 1.5 (.7) 3 months	NS
Meng 2019 ⁶ (30884961) Telephone based cognitive behavioral therapy vs. in-person cognitive behavioral therapy Medium Explanatory USA	Mean (SD) Caregiver health service use: prescription medication 3 months	NR	3.5 (2.2) baseline 3.5 (2.5) 3 months	3.5 (3.1) baseline 4.1 (2.6) 3 months	NS
Meng 2019 ⁶ (30884961) Telephone based cognitive behavioral therapy vs. in-person cognitive behavioral therapy Medium Explanatory USA	Mean (SD) Caregiver health service use: ER visits 3 months	NR	1.8 (1.2) baseline 1.6 (0.9) 3 months	0.1 (0.5) baseline 1.2 (0.4) 3 months	NS
Meng 2019 ⁶ (30884961) Telephone based cognitive behavioral therapy vs. in-person cognitive behavioral therapy Medium Explanatory USA	Mean (SD) Caregiver health service use: hospital days 3 months	NR	1.4 (0.9) baseline 3 (1.4) 3 months	0.1 (0.8) baseline 3.3 (1.5) 3 months	NS
Meng 2019 ⁶ (30884961) Telephone based cognitive behavioral therapy vs. in-person cognitive behavioral therapy Medium Explanatory USA	Mean (SD) Caregiver health service use: out of pocket expenditures in dollars 3 months	NR	104.9 (94.4) baseline 216.3 (504.3) 3 months	41 (72.7) baseline 191.7 (228.8) 3 months	NS
Meng 2019 ⁶ (30884961) Telephone based cognitive behavioral therapy vs. in-person cognitive behavioral therapy Medium Explanatory USA	Mean (SD) Caregiver health service use: total monthly expenditures in dollars 3 months	NR	1019.9 (1412.1) baseline 883.2 (749.5) 3 months	844 (1851) baseline 768.9 (661.2) 3 months	NS
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver depression: PROMIS depressive symptoms scale 6 weeks	-.25 (-.55, .06)	65.85 (2.02) baseline 55.18 (2.08) 6 weeks	65.62 (2.04) baseline 59.65 (2.15) 6 weeks	Interaction 0.02
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver mental health: PROMIS mental health scale 6 weeks	0.21 [-.09, .52]	37.4 (4.10) baseline 39.8 (4.10) 6 weeks	39.8 (4.10) baseline 39.8 (4.10) 6 weeks	Interaction 0.49
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver physical health: PROMIS physical health scale 6 weeks	0.24 [-.07, .54]	42.3 (4.20) baseline 44.9 (4.30) 6 weeks	44.9 (4.30) baseline 44.9 (4.30) 6 weeks	Interaction 0.03
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver anxiety: Quality of Life in Neurological Disorders (NeuroQOL) anxiety measure 6 weeks	-0.32 [-.63, -.02]	52.15 (1.63) baseline 44.64 (1.68) 6 weeks	51.38 (1.65) baseline 48.91 (1.76) 6 weeks	Interaction .009
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver burden: Zarit Burden Interview 6 weeks	-0.16 [-.46, .14]	42.19 (1.58) baseline 38.18 (1.60) 6 weeks	42.18 (1.59) baseline 41.13 (1.65) 6 weeks	Interaction .066

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver strain: Caregiver Strain Index (CSI) 6 weeks	.01 [-.30, .30]	8.40 (.33) baseline 8.46 (.33) 6 weeks	8.061 (.35) baseline 8.55 (.33) 6 weeks	Interaction 0.74
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver positive emotion: Differential Emotions Scale (DES) 6 weeks	0.58 [.27, .88]	4.64 (.16) baseline 4.86 (.16) 6 weeks	4.92 (.17) baseline 4.31 (.17) 6 weeks	Interaction <.001
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver negative emotion: Differential Emotions Scale (DES) 6 weeks	-0.22 [-.52, .08]	3.66 (.13) baseline 2.6 (.13) 6 weeks	3.70 (.13) baseline 2.91 (.14) 6 weeks	Interaction 0.13
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver positive aspects of caregiving: the Positive Aspects of Caregiving Scale 6 weeks	0.35 [.05, .66]	35.92 (1.09) baseline 39.04 (1.12) 6 weeks	26.88 (1.11) baseline 36.41 (1.16) 6 weeks	Interaction 0.001
Moskowitz 2019 ⁷ (31045422) Life Enhancing Activities for Family Caregivers (LEAF) vs. waitlist control Medium Explanatory USA	Cohen's d (95% CI) Mean (SE) Caregiver perceived stress: The Perceived Stress Scale 6 weeks	-0.20 [-.50, .11]	30.64 (.71) baseline 27.48 (.72) 6 weeks	31.08 (.73) baseline 29.24 (.76) 6 weeks	Interaction 0.10
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months Caregiver quality of life: World Health Organization Quality of Life questionnaire (WHOQOL-BREF) short form Global QOL subscale 6 month	Cohens d = -0.2	60.9 (2.4) baseline 53.2 (2.4) 6 months	58.5 (2.4) baseline 54.5 (2.6) 6 months	NS (p-value NR)
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care. Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months Caregiver confidence: Sense of Competence Questionnaire (SCQ) personal life subscale 6 months	Cohens d = -0.5	16.9 (1.1) baseline 19.9 (1.0) 6 months	17.4 (1.1) baseline 16.5 (1.1) 6 months	NS (p-value NR)
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months Caregiver confidence: Sense of Competence Questionnaire (SCQ) performance as a caregiver subscale 6 months	Cohens d = -0.4	17.2 (1.0) baseline 20.9 (0.9) 6 months	18.0 (1.0) baseline 20.0 (1.0) 6 months	NS (p-value NR)
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months Caregiver confidence: Sense of Competence Questionnaire (SCQ) satisfaction with PLWD subscale 6 months	Cohens d = -0.5	8.9 (0.7) baseline 10.4 (0.7) 6 months	10.1 (0.7) baseline 9.7 (0.7) 6 months	NS (p-value NR)
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months Caregiver social support: (F-SOZU) 6 months	Cohens d =-0.1	53.3 (1.1) baseline 50.5 (1.3) 6 months	53.2 (1.4) baseline 53.5 (1.4) 6 months	NS (p-value NR)
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months Caregiver perceived stress 6 months	Cohens d=0.1	25.5 (1.0) baseline 27.3 (1.2) 6 months	27.2 (1.) baseline 26.9 (1.2) 6 months	NS (p-value NR)

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Nordheim 2019 ⁸ (30775982) couple-based interdisciplinary psychosocial intervention vs. usual care Medium Explanatory Germany	Estimated marginal mean (SE) baseline Estimated marginal mean (SE) 6 months Caregiver depressive symptoms: Geriatric depression scale (GDS) 6 months	Cohens d = 0.0	4.9 (0.3) baseline 4.8 (0.3) 6 months	4.5 (0.3) baseline 5.0 (0.3) 6 months	NS (p-value NR)
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) Caregiver depressive symptoms: Centers for Epidemiologic Study Depression Scale (CES-D) 4 months	-0.59 (-1.74 to 0.55)	NR	NR	0.31
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) Caregiver burden: Zarit Burden Short Form (12 item) 4 months	-0.39 (-2.06 to 1.29) .65	NR	NR	0.65
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) Caregiver distress: Neuropsychiatric Inventory – Clinician (NPI-C) distress subscale 4 months	-0.07 (-0.14 to -0.01)	NR	NR	0.03
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) Caregiver time spent caregiving: hours on ADLS 4 months	-0.53 (-1.20 to 0.14)	NR	NR	0.12
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) Caregiver time spent caregiving: hours on IADLs 4 months	-1.09 (-2.34 to 0.15)	NR	NR	0.08
Gitlin 2018 ¹³ (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) Caregiver time spent caregiving: hours on duty 4 months	-0.28 (-2.25 to 1.68)	NR	NR	0.78
Gitlin 2018 ^{13, 20} (29192967) Home-based Tailored Activity Program vs. attention control using telephone visits Medium Explanatory USA	Mean difference (95% CI) Caregiver time spent caregiving: hours on doing things 4 months	-0.99 (-2.28 to 0.30)	NR	NR	0.13
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) Caregiver health service use: Number of monthly emergency department visits during intervention 6 months	NR	0.02 (0.06)	0.05 (0.13)	0.02
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) Caregiver health service use: Number of monthly hospital visits during intervention 6 months	NR	0.01 (0.04)	0.04 (0.10)	0.006
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) Caregiver health service use: number of monthly doctor visits during intervention 6 months	NR	0.85 (0.89)	0.71 (0.61)	0.88

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Absolute risk reduction (95% CI) Caregiver hospital admissions 6 months	11.4% (0.023 to 0.25)	NR	NR	0.01
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Absolute risk reduction (95% CI) Caregiver emergency department visits 6 months	9.5% (0.01 to 0.19)	NR	NR	0.048
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) Caregiver community support use: Number of community support services used in final month of intervention 6 months	NR	8.47 (5.14)	5.14 (7.05)	0.02
Tremont 2017 ⁸⁴ (28008609) companion to Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) Caregiver out of pocket cost on formal care for PLWD for final month of intervention in dollars 6 months	\$274.43 (901.68)	NR	NR	0.73 (no difference between groups)
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups Caregiver Health related Quality of life: SF-12 mental component 12 months	0.52 (-1.28 to 2.32)	NR	NR	0.57
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups Caregiver Health related Quality of life: SF-12 mental component 12 months	0.10 (-1.72 to 1.93)	NR	NR	0.91
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups Caregiver Health related Quality of life: SF-12 physical component 12 months	0.613 (-1.23 to 2.46)	NR	NR	0.52
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups Caregiver Health related Quality of life: SF-12 physical component 12 months	-0.428 (-2.29 to 1.43)	NR	NR	0.65
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups Caregiver anxiety: Hospital Anxiety and Depression Scale (HADS) anxiety subscale 12 months	-0.33 (-1.31 to 0.60)	NR	NR	0.47

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups Caregiver depression: Hospital Anxiety and Depression Scale (HADS) anxiety subscale 12 months	0.298 (-0.65 to 1.25)	NR	NR	0.54
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups Caregiver depression: Hospital Anxiety and Depression Scale (HADS) depression subscale 12 months	0.0144 (-0.85 to 0.88)	NR	NR	0.97
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups Caregiver depression: Hospital Anxiety and Depression Scale (HADS) depression subscale 12 months	-0.0700 (-0.95 to 0.81)	NR	NR	0.88
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups Caregiver Health-related quality of life: EQ-5D global health visual analog scale 12 months	0.384 (-4.20 to 4.96)	NR	NR	0.87
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups Caregiver Health-related quality of life: EQ-5D global health visual analog scale 12 months	0.915 (-3.65 to 5.48)	NR	NR	0.69
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP vs other 2 groups Caregiver Positive Aspects of Caring: Carers of Older People in Europe Index (COPE index) positive aspects subscale 12 months	0.0125 (-0.54 to 0.56)	NR	NR	0.96
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT vs other 2 groups Caregiver Positive Aspects of Caring: Carers of Older People in Europe Index (COPE index) positive aspects subscale 12 months	0.0784 (-0.45 to 0.61)	NR	NR	0.77
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for CSP over other 2 groups Caregiver Quality of Caregiver-PLWD Relationship (QCPR) 12 months	1.416 (-0.47 to 3.30)	NR	NR	0.14

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Charlesworth 2016 ¹⁸ (27521377) 3 groups: 1) One-to-one peer support to family carers from experienced carers (Carer Supporter Programme (CSP), 2) group reminiscence therapy Remembering Yesterday, Caring Today (RYCT), & 3) usual care Low Explanatory UK	Adjusted mean difference (95% CI) for RYCT over other 2 groups Caregiver Quality of Caregiver-PLWD Relationship (QCPR) 12 months	1.869 (-0.02 to 3.75)	NR	NR	0.05
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention (BF) vs. simplified psychoeducation (SIM-PE) vs. standard psychoeducation (STD-PE) Medium Explanatory Hong Kong	d effect size Adjusted mean (SD) Caregiver depressive symptoms: Hamilton Depression Rating Scale 8 weeks	d = -0.77	2.441 (3.499) BF	5.137 (3.476) SIM-PE	Regression p<0.001
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention (BF) vs. simplified psychoeducation (SIM-PE) vs. standard psychoeducation (STD-PE) Medium Explanatory Hong Kong	d effect size Adjusted mean (SD) Caregiver burden: Zarit Caregiver Burden Interview (ZBI) 8 weeks	d = -0.47	27.133 (11.704) BF	32.544 (11.431) SIM-PE	Regression p=0.028
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention (BF) vs. simplified psychoeducation (SIM-PE) vs. standard psychoeducation (STD-PE) Medium Explanatory Hong Kong	d effect size Adjusted mean (SD) Caregiver role overload: 8 weeks	d = -0.43	9.650 (2.424) BF	10.705 (2.478) SIM-PE	Regression p=0.046
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention (BF) vs. simplified psychoeducation (SIM-PE) vs. standard psychoeducation (STD-PE) Medium Explanatory Hong Kong	d effect size Adjusted mean (SD) Caregiver psychological wellbeing: Ryff's Psychological Well-being Scale 8 weeks	d = 0.42	66.652 (6.753) BF	63.867 (6.559) SIM-PE	Regression p=0.048
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention (BF) vs. simplified psychoeducation (SIM-PE) vs. standard psychoeducation (STD-PE) Medium Explanatory Hong Kong	d effect size Adjusted mean (SD) Caregiver depressive symptoms: Hamilton Depression Rating Scale 8 weeks	d = -0.96	2.441 (3.499) BF	5.833 (3.432) STD-PE	Regression p<0.001
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention (BF) vs. simplified psychoeducation (SIM-PE) vs. standard psychoeducation (STD-PE) Medium Explanatory Hong Kong	d effect size Adjusted mean (SD) Caregiver burden: Zarit Caregiver Burden Interview (ZBI) 8 weeks	d = -0.65	27.133 (11.704) BF	34.749 (11.886) STD-PE	Regression p=0.003
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention (BF) vs. simplified psychoeducation (SIM-PE) vs. standard psychoeducation (STD-PE) Medium Explanatory Hong Kong	d effect size Adjusted mean (SD) Caregiver role overload: 8 weeks	d = NR	9.650 (2.424) BF	10.243 (2.387) STD-PE	Regression p=0.234
Cheng 2016 ¹⁹ (27401052) Benefit finding intervention (BF) vs. simplified psychoeducation (SIM-PE) vs. standard psychoeducation (STD-PE) Medium Explanatory Hong Kong	d effect size Adjusted mean (SD) Caregiver psychological wellbeing: Ryff's Psychological Well-being Scale 8 weeks	d = NR	66.652 (6.753) BF	65.027 (6.894) STD-PE	Regression p=0.264

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Gonyea 2016 ²⁰ (24855313) Circulo de Cuidado, a culturally-sensitive, cognitive behavioral (CBT) group intervention vs. psychoeducational (PED) control condition Medium Explanatory USA	Mean (SD) Caregiver distress of PLWD neuropsychiatric symptoms: Spanish version of the Neuropsychiatric Inventory-Distress (NPI-D) 12 week 24 week	NR	17.94 (6.84) baseline 16.64 (7.02) 12 weeks 16.39 (7.02) 24 weeks	17.91 (7.30) baseline 17.85 (7.26) 12 weeks 18.26 (7.37) 24 weeks	Repeated measures ANCOVA < 0.001
Gonyea 2016 ²⁰ (24855313) Circulo de Cuidado, a culturally-sensitive, cognitive behavioral (CBT) group intervention vs. psychoeducational (PED) control condition Medium Explanatory USA	Mean (SD) Caregiver self-efficacy: Spanish version of the Revised Scale for Caregiving Self-Efficacy (RSCSE) 12 week 24 week	NR	69.11 (12.75) baseline 70.36 (12.44) 12 weeks 70.27 (12.43) 24 weeks	69.19 (9.01) baseline 69.29 (8.80) 12 weeks 69.15 (8.88) 24 weeks	Repeated measures ANCOVA < 0.001
Gonyea 2016 ²⁰ (24855313) Circulo de Cuidado, a culturally-sensitive, cognitive behavioral (CBT) group intervention vs. psychoeducational (PED) control condition Medium Explanatory USA	Mean (SD) Caregiver depressive symptoms: Spanish version of the Center for Epidemiological Studies-Depression Scale (CES-D) 12 week 24 week	NR	14.39 (6.14) baseline 13.79 (5.85) 12 weeks 13.82 (5.88) 24 weeks	14.41 (6.31) baseline 14.38 (6.10) 12 weeks 14.44 (6.07) 24 weeks	Repeated measures ANCOVA <0.01
Gonyea 2016 ²⁰ (24855313) Circulo de Cuidado, a culturally-sensitive, cognitive behavioral (CBT) group intervention vs. psychoeducational (PED) control condition Medium Explanatory USA	Mean (SD) Caregiver anxiety: Spanish version of the State Anxiety Inventory-State (STAI-S) 12 week 24 week	NR	37.09 (10.25) baseline 36.85 (9.88) 12 weeks 36.96 (10.00) 24 weeks	37.12 (9.69) baseline 37.09 (9.61) 12 weeks 37.12 (9.68) 24 weeks	Repeated measures ANCOVA p>0.05 NS
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) Caregiver quality of life: physical component SF-36 3 months	d = 0.38	1.0 (-0.5 to 2.5)	-2.0 (-3.5 to -0.5)	.006
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) Caregiver quality of life: physical component SF-36 9 months	d=NR	-0.0 (-1.5 to 1.4)	-1.7 (-3.2 to -0.2)	0.13
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) Caregiver quality of life: mental component SF-36 3 months	d=NR	NR	NR	0.99
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) Caregiver quality of life: mental component SF-36 9 months	d=NR	NR	NR	0.58
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) Caregiver competence: spousal Sense of Competence Questionnaire (SCQ) 3 months	d=NR	NR	NR	0.76
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) Caregiver competence: spousal Sense of Competence Questionnaire (SCQ) 9 months	d=NR	NR	NR	0.38

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) Caregiver mastery Pearlin Mastery Scale (PMS) 3 months	d=NR	NR	NR	0.06
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean change (95% CI) Caregiver mastery Pearlin Mastery Scale (PMS) 9 months	d=NR	NR	NR	0.11
Laakkonen 2016 ²¹ (27060101) Self-management groups vs. usual care Medium Explanatory Finland	Adjusted Mean cost in € per year (95% CI) Caregiver Health & Social services cost: 24 months	-896 (-3,657; 1,864)	2,658 (1,473–4,176)	3,555 (1,674–6,718)	0.51
Otero 2015 ⁸³ (25331992) Cognitive behavioral problem solving vs. usual care Low Explanatory Spain	Relative risk (RR) Number needed to treat (NNT) Caregiver depression: Structured Clinical Interview for Axis 1 Disorders of the DSM-IV (SCID-CV) 12 months	0.40 RR 6.7 NNT			0.010
Otero 2015 ⁸³ (25331992) Cognitive behavioral problem solving vs. usual care Low Explanatory Spain	Standardized mean difference Center for Epidemiologic Depression Scale (CESD) Spanish version 12 months	1.14			NR
Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver depressive symptoms: Center for Epidemiology Studies Depression Scale (CESD) 6 months	NR	17.04 (10.22) baseline 14.15 (10.00) 6 months	15.19 (9.98) baseline 15.62 (10.18) 6 months	0.003
Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver burden: Zarit Burden Interview (ZBI) 6 months	NR	38.61 (13.98) baseline 35.95 (14.34) 6 months	38.82 (14.63) baseline 37.17 (13.93) 6 months	0.485
Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver reactions to care: Revised Memory and Behavior Problem Checklist (RMBPC-RT) Reaction Score 6 months	NR	22.98 (12.89) baseline 20.04 (13.68) 6 months	22.59 (13.94) baseline 22.00 (12.82) 6 months	0.160
Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver : Family Assessment Device (FAD) 6 months	NR	2.11 (0.61) baseline 2.09 (0.63) 6 months	2.05 (0.55) baseline 2.00 (0.50) 6 months	0.798
Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver confidence: Self-Efficacy Questionnaire (SEQ-SM) Symptom Management subscale 6 months	NR	29.35 (10.78) baseline 34.17 (9.55) 6 months	29.74 (11.86) baseline 32.27 (10.04)6 months	0.108
Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver confidence: Self-Efficacy Questionnaire (SEQ-SS) Support Services subscale 6 months	NR	23.39 (10.89) baseline 31.06 (7.77) 6 months	24.00 (9.93) baseline 30.64 (7.28) 6 months	0.467

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver : Positive Aspects of Caregiving (PAC) 6 months	NR	31.51 (7.58) baseline 34.13 (8.31) 6 months	30.35 (9.65) baseline 35.24 (7.55) 6 months	0.041
Tremont 2015 ²⁸ (25074341) Telephone Tracking–Caregiver (FITT-C) vs. Telephone Support (TS) Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver health related quality of life; EuroQoL Visual Analog scale 6 months	NR	80.08 (16.07) baseline 79.87 (15.00) 6 months	77.14 (17.61) baseline 77.59 (15.69) 6 months	0.748
Livingston 2014 ³⁰ (25300037) START (STrAtegies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	Adjusted mean difference (95% CI) Caregiver Affective symptoms [Hospital Anxiety and Depression Scale-total (HADS-T)] 2 years	–2.58 (–4.26 to –0.90)	NR	NR	NR
Livingston 2014 ³⁰ (25300037) START (STrAtegies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	Adjusted mean difference (95% CI) Caregiver Anxiety: Hospital Anxiety and Depression Scale-anxiety (HADS-A) 2 years	–1.16 (–2.15 to –0.18)	NR	NR	NR
Livingston 2014 ³⁰ (25300037) START (STrAtegies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	OR (95% CI) Caregiver Anxiety: Hospital Anxiety and Depression Scale-anxiety (HADS-A) (score ≥ 9) 2 years	0.57 (0.26 to 1.24)	57 (43.2) N (%)	32 (50.0) N (%)	NR
Livingston 2014 ³⁰ (25300037) START (STrAtegies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	Adjusted mean difference (95% CI) Caregiver Depression: Hospital Anxiety and Depression Scale-depression (HADS-D) 2 years	–1.45 (–2.32 to –0.57)	NR	NR	NR
Livingston 2014 ³⁰ (25300037) START (STrAtegies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	OR (95% CI) Caregiver Depression: Hospital Anxiety and Depression Scale-depression (HADS-D) (score ≥ 9) 2 years	0.14 (0.04 to 0.53)	30 (22.7) N (%)	19 (29.7) N (%)	NR
Livingston 2014 ³⁰ (25300037) START (STrAtegies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	Adjusted mean difference (95% CI) Caregiver QOL: Health Status Questionnaire (HSQ) – mental health 2 years	7.47 (2.87 to 12.08)	NR	NR	NR
Livingston 2014 ³⁰ (25300037) START (STrAtegies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	OR (95% CI) Caregiver Abusive behaviors: Modified Conflict Tactic Score (MCTS) (at least one item with score of ≥ 2) 2 years	0.83 (0.36 to 1.93)	28 (29.5) N (%)	11 (23.4) N (%)	NR
Livingston 2014 ³⁰ (25300037) START (STrAtegies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care	Mean difference (95% CI) Caregiver health care cost: Quality-Adjusted Life-Years	£336 (95% CI –£223 to £895)	NR	NR	NR

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Low Explanatory UK	(QALYs) from the EQ-5D 2 years				
Livingston 2014 ³⁰ (25300037) START (STrategies for RelaTives) manual-based coping strategy programme in promoting the mental health of carers of people with dementia vs. usual care Low Explanatory UK	Mean difference (95% CI) Caregiver health care cost: Quality-Adjusted Life-Years (QALYs) from the Hospital Anxiety and Depression Scale-total (HADS-T) 2 years	£303 (95% CI -£206 to £812)	NR	NR	NR
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver burden: Spanish Zarit caregiver burden interview 4 months	-0.55 (-3.64; 2.55) – Mean difference -0.04 (-0.30; 0.21) - SDM	-1.17 (12.3)	-0.63 (12.0)	0.73
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver psychological distress: Spanish General Health Questionnaire 28 item (GHQ-28) 4 months	-2.34 (-5.27; 0.59) mean diff -0.20 (-0.46; 0.05) SDM	-4.76 (12.6)	-2.42 (10.3)	>.05
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver quality of life: Spanish SF-12 physical function 4 months	-1.02 (-11.10 to 9.06) -0.03 (-0.30 to 0.25)	-1.02 (30.0)	0.0 (41.3)	>.05
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver quality of life: Spanish SF-12 role physical 4 months	1.80 (-5.63 to 9.22) 0.07 (-0.21 to 0.34)	3.09 (26.9)	1.30 (26.7)	>.05
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver quality of life: Spanish SF-12 bodily pain 4 months	6.85 (-1.58 to 15.28) 0.22 (-0.05 to 0.50)	6.38 (25.7)	-0.47 (34.3)	0.11
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver quality of life: Spanish SF-12 general health 4 months	-5.01 (-10.48 to 0.45) -0.25 (-0.53 to 0.02)	-3.12 (19.3)	1.89 (20.0)	0.07
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver quality of life: Spanish SF-12 vitality 4 months	3.71 (-4.34 to 11.76) 0.13 (-0.15 to 0.40)	2.04 (28.6)	-1.67 (29.5)	>.05
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium	Mean difference (95% CI) SMD (95% CI) Caregiver quality of life: Spanish SF-12 social functioning 4 months	-0.78 (-9.33 to 7.77) -0.02 (-0.30 to 0.25)	-4.08 (31.6)	-3.30 (30.3)	>.05

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Explanatory Spain					
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver quality of life: Spanish SF-12 role emotional 4 months	-4.25 (-11.48 to 2.98) -0.16 (-0.44 to 0.11)	-3.06 (27.7)	1.19 (24.5)	>.05
Martin-Carrasco 2014 ³¹ (24113563) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean difference (95% CI) SMD (95% CI) Caregiver quality of life: Spanish SF-12 mental health 4 months	-1.23 (-7.22 to 4.75) -0.06 (-0.33 to 0.22)	1.53 (23.8)	2.76 (19.2)	>.05
Waldorff 2012 ⁴⁰ (22807076) DAISY plus support (multifaceted and semi-tailored counselling, education, and support) vs. support control group Medium Explanatory Denmark	Mean change score (95% CI) Caregiver depressive symptom: Geriatric depression scale 12 months	NR	0.96 (0.05 to 1.86)	0.26 (-0.51 to 1.03)	0.2500
de Rotrou 2011 ⁴² (20922772) Psycho Education Program vs. usual care Medium Explanatory France	Imputed mean (SD) Caregiver depressive symptoms: Montgomery and Asberg Depression Rating Scale (MADRS) Baseline 3 months	NR	9.03 (0.79) baseline 8.61 (0.94) 3 months	10.16 (1.22) baseline 10.11 (1.18) 3 months	NS Value not reported
de Rotrou 2011 ⁴² (20922772) Psycho Education Program vs. usual care Medium Explanatory France	Imputed mean (SD) Caregiver burden: Zarit Caregiver Burden Interview (ZBI) Baseline 3 months	NR	22.97 (2.79) baseline 22.23 (1.59) 3 months	24.30 (4.15) baseline 23.53 (2.05) 3 months	NS Value not reported
de Rotrou 2011 ⁴² (20922772) Psycho Education Program vs. usual care Medium Explanatory France	Imputed mean (SD) Caregiver perception of disease understanding: Visual Analogue Scales (VAS) Baseline 3 months	NR	12.52 (0.27) baseline 14.44 (0.47) 3 months	12.05 (0.27) baseline 12.54 (0.53) 3 months	NR
de Rotrou 2011 ⁴² (20922772) Psycho Education Program vs. usual care Medium Explanatory France	Imputed mean (SD) Caregiver coping: Visual Analogue Scales (VAS) Baseline 3 months	NR	12.53 (0.24) baseline 13.47 (0.44) 3 months	12.72 (0.29) baseline 12.63 (0.52) 3 months	NR
de Rotrou 2011 ⁴² (20922772) Psycho Education Program vs. usual care Medium Explanatory France	Imputed mean (SD) Caregiver depressive symptoms: Montgomery and Asberg Depression Rating Scale (MADRS) Baseline 6 months	NR	9.03 (0.79) baseline 9.28 (0.99) 3 months	10.16 (1.22) baseline 11.43 (1.24) 3 months	Interaction 0.373
de Rotrou 2011 ⁴² (20922772) Psycho Education Program vs. usual care Medium Explanatory France	Imputed mean (SD) Caregiver burden: Zarit Caregiver Burden Interview (ZBI) Baseline 6 months	NR	22.97 (2.79) baseline 23.90 (1.86) 3 months	24.30 (4.15) baseline 25.95 (2.09) 3 months	Interaction 0.657
de Rotrou 2011 ⁴² (20922772) Psycho Education Program vs. usual care Medium Explanatory France	Imputed mean (SD) Caregiver perception of disease understanding: Visual Analogue Scales (VAS) Baseline 6 months	NR	12.52 (0.27) baseline 15.03 (0.38) 3 months	12.05 (0.27) baseline 12.52 (0.50) 3 months	Interaction 0.003

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
de Rotrou 2011 ⁴² (20922772) Psycho Education Program vs. usual care Medium Explanatory France	Imputed mean (SD) Caregiver coping: Visual Analogue Scales (VAS) Baseline 6 months	NR	12.53 (0.24) baseline 13.63 (0.44) 3 months	12.72 (0.29) baseline 12.24 (0.57) 3 months	Interaction 0.025
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Standardized mean difference (95%CI) summary finding Mean change score (SD) Caregiver burden: Zarit Caregiver Burden Interview (ZBI) 6 months	-1.05 (-1.60 to -0.48)	-3.6 (4.6)	0.3 (2.9)	<0.001
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Standardized mean difference (95%CI) summary finding Mean change score (SD) Caregiver mental health: Self reporting questionnaire 20 (SRQ-20) psychological morbidity 6 months	-0.01 (-0.55 to 0.55)	-3.1 (4.0)	-3.0 (3.1)	0.97
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Standardized mean difference (95%CI) summary finding Mean change score (SD) Caregiver distress: Neuropsychiatric Inventory (NPI-Q) carer distress score 6 months	0.01 (-0.55 to 0.57)	-2.3 (4.7)	-2.4 (4.6)	0.96
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Standardized mean difference (95%CI) summary finding Mean change score (SD) Caregiver QOL: WHOQOL-BREF physical subscale 6 months	0.36 (-0.19 to 0.92)	-9.7 (18.7)	-15.5 (13.9)	0.19
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Standardized mean difference (95%CI) summary finding Mean change score (SD) Caregiver QOL: WHOQOL-BREF psychological subscale 6 months	0.10 (-0.44 to 0.65)	10.0 (11.5)	8.9 (11.1)	0.71
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Standardized mean difference (95%CI) summary finding Mean change score (SD) Caregiver QOL: WHOQOL-BREF social subscale 6 months	0.39 (-0.15 to 0.94)	7.1 (12.6)	1.7 (15.2)	0.16
Guerra 2011 ⁴³ 10/66 Helping Carers to Care vs waitlist control Medium Explanatory Peru	Standardized mean difference (95%CI) summary finding Mean change score (SD) Caregiver QOL: WHOQOL-BREF environmental subscale 6 months	-0.15 (-0.70 to 0.39)	7.6 (11.4)	9.5 (13.0)	0.58
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) Caregiver interaction with PLWD: Sense of Competence Questionnaire 52 weeks	3.8 (-3.5 to 11.2)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) Caregiver depression: Center for Epidemiologic Depression Scale 52 weeks	-1.4 (-5.1 to 2.3)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation)	Mean difference (95% CI) Caregiver QOL: Dementia Quality of Life Instrument overall 52 weeks	0.2 (-0.1 to 0.5)	NR	NR	NS

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Medium 26-52 weeks Explanatory Germany					
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) Caregiver QOL: Short-Form 12 Health Survey Questionnaire – physical component 52 weeks	-1.0 (-6.1 to 4.0)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) Caregiver QOL: Short-Form 12 Health Survey Questionnaire – mental component 52 weeks	-1.7 (-6.9 to 3.4)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) Caregiver work: Resource utilization in Dementia - Basic Activities of Daily Living care by primary carer (hours per day) subscale 52 weeks	0.1 (-0.8 to 1.0)	NR	NR	NS
Voight-Radloff 2011 ⁴⁶ (22021760) Community Occupational Therapy in Dementia Program vs. usual care (one session home consultation) Medium 26-52 weeks Explanatory Germany	Mean difference (95% CI) Caregiver work: Resource utilization in Dementia - Instrumental Activities of Daily Living -care by primary carer (hours per day) subscale 52 weeks	0.5 (-0.6 to 1.6)	NR	NR	NS
Wang 2011 ⁴⁷ (21752121) Family Mutual Support Program in Dementia Care (FMSP-DC) vs. usual care Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver burden: Family Caregiving Burden Inventory (FCBI) 6 months	NR	68.2 (11.9) baseline 55.2 (15.0) 6 months	68.8 (16.7) baseline 65.0 (18.1) 6 months	<0.001
Wang 2011 ⁴⁷ (21752121) Family Mutual Support Program in Dementia Care (FMSP-DC) vs. usual care Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver quality of life: World Health Organization Quality of Life Scale (Brief H.K. version) (WHOQOL-BREF) 6 months	NR	65.9 (13.0) baseline 78.8 (19.0) 6 months	67.0 (13.5) baseline 68.9 (15.7) 6 months	<0.001
Wang 2011 ⁴⁷ (21752121) Family Mutual Support Program in Dementia Care (FMSP-DC) vs. usual care Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver social support: Six-item Social Support Questionnaire (SSQ6) – number of support persons subscale 6 months	NR	3.1 (1.0) baseline 4.4 (2.0) 6 months	2.9 (1.1) baseline 2.9 (1.3) 6 months	NS
Wang 2011 ⁴⁷ (21752121) Family Mutual Support Program in Dementia Care (FMSP-DC) vs. usual care Medium Explanatory USA	Mean (SD) baseline Mean (SD) 6 months Caregiver social support: Six-item Social Support Questionnaire (SSQ6) – satisfaction subscale 6 months	NR	5.3 (1.8) baseline 6.2 (3.1) 6 months	5.5 (1.1) baseline 5.6 (2.1) 6 months	NS
Gitlin 2010 ⁴⁹ (20662955) Advancing Caregiver Training (ACT) vs. no treatment control Medium Explanatory USA	Adjusted mean difference (95% CI) Cohen's D Caregiver upset with PLWD problem behaviors 4 months	-0.93 (-1.76 to -0.10) 0.30	NR	NR	0.03
Gitlin 2010 ⁴⁹ (20662955) Advancing Caregiver Training (ACT) vs. no treatment control Medium	Adjusted mean difference (95% CI) Cohen's D	0.33 (0.08 to 0.58) 0.30	NR	NR	0.01

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Explanatory USA	Caregiver confidence managing PLWD problem behaviors 4 months				
Gitlin 2010 ⁴⁹ (20662955) Advancing Caregiver Training (ACT) vs. no treatment control Medium Explanatory USA	Adjusted mean difference (95% CI) Caregiver skills: task simplification strategies 4 months	0.06 (-0.03 to 0.15)	NR	NR	0.21
Gitlin 2010 ⁴⁹ (20662955) Advancing Caregiver Training (ACT) vs. no treatment control Medium Explanatory USA	Adjusted mean difference (95% CI) Caregiver burden: Zarit Caregiver Burden Interview (ZBI) 4 months	-1.37 (-2.75 to 0.01)	NR	NR	0.05
Gitlin 2010 ⁴⁹ (20662955) Advancing Caregiver Training (ACT) vs. no treatment control Medium Explanatory USA	Percentage with CES-D score > 8 Caregiver depressive symptoms: Centers for Epidemiologic Study Depression Scale (CES-D) 4 months	NR	53.0	67.8	0.02 for chi-square test
Gitlin 2010 ⁴⁹ (20662955) Advancing Caregiver Training (ACT) vs. no treatment control Medium Explanatory USA	Adjusted mean difference (95% CI) Cohen's D Caregiver affect: 4 months	0.46 (0.27-0.65) 0.58	NR	NR	0.004
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean change score (SD) Caregiver burden: Zarit caregiver burden interview 10 months	NR	-8.1 (17.3)	2.1 (16.5)	.0083
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver quality of life: Spanish SF-36 physical functioning subscale 10 months	NR	80.1 (20.7)	68.8 (25.6)	0.0310
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver quality of life: Spanish SF-36 physical role subscale 10 months	NR	84.7 (34.6)	56.8 (47.0)	0.0074
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver quality of life: Spanish SF-36 bodily pain subscale 10 months	NR	74.0 (18.7)	61.7 (26.9)	0.0157
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver quality of life: Spanish SF-36 General Health subscale 10 months	NR	53.4 (18.0)	40.1 (15.7)	0.0011
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care	Mean (SD) Caregiver quality of life: Spanish SF-36 Vitality 10 months	NR	53.8 (15.9)	38.9 (17.9)	0.0002

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Medium Explanatory Spain					
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver quality of life: Spanish SF-36 Social Functioning 10 months	NR	71.0 (23.4)	58.9 (27.7)	0.0488
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver quality of life: Spanish SF-36 Emotional Role 10 months	NR	73.5 (41.0)	47.4 (48.2)	0.0160
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver quality of life: Spanish SF-36 Mental Health 10 months	NR	63.0 (9.2)	60.9 (8.3)	0.3197
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver mental health status: General Health Questionnaire (GHQ-28) 4 months	NR	4.7 (7.2)	6.3 (6.6)	0.0340
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver mental health status: General Health Questionnaire (GHQ-28) 10 months	NR	2.2 (4.0)	7.8 (7.6)	0.0004
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Frequency (percent) Caregiver healthcare and social resources use: number of visits 4 months	NR	0 visits: 34/44 (77%) 1-2 visits: 10/44 (23%)	0 visits: 29/38 (77%) 1-2 visits: 9/38 (23%)	NR (NS)
Martin-Carrasco 2009 ⁵⁴ (18949763) Psychoeducational Intervention Program to teach strategies for confronting problems of PLWD care vs. usual care Medium Explanatory Spain	Mean (SD) Caregiver healthcare and social resources use: time spent on medical care 4 months	NR	24.6 (19.6)	14.5 (10.1)	NR
Gallagher-Thompson 2008 ⁵⁶ (25067886) Coping with Caregiving (CWC) vs. minimal telephone-based control condition (TSC) Medium Explanatory	Beta (SE) for linear regression model Caregiver depressive symptoms: CESD Center for Epidemiological Studies Depression scale (CESD) 6 months	-2.135 (1.073)	NR	NR	ANOVA p=.048
Gallagher-Thompson 2008 ⁵⁶ (25067886) Coping with Caregiving (CWC) vs. minimal telephone-based control condition (TSC) Medium Explanatory	Beta (SE) for linear regression model Caregiver perceived psychological stress: Perceived Stress Scale (PSS-10) 6 months	-1.530 (0.760)	NR	NR	ANOVA p=.046
Gallagher-Thompson 2008 ⁵⁶ (25067886) Coping with Caregiving (CWC) vs. minimal telephone-based control condition (TSC)	Beta (SE) for linear regression model Caregiver bother of PLWD neuropsychiatric symptoms:	-0.250 (0.092)	NR	NR	ANOVA p=.007

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Medium Explanatory	Subscale of Revised Memory and Behavior Problem Checklist (RMBPC-CB) 6 months				
Graff 2006 ⁶⁰ (17114212) Community based occupational therapy program vs. Waitlist for occupational therapy Medium Explanatory	Adjusted Mean difference (95% CI) Caregiver competence: Sense of Competence Questionnaire (SCQ) 6 weeks	11.0 (9.2; 12.8) 1.2 2.5 (2.3 to 2.7) NNT	NR	NR	<0.0001
Graff 2006 ⁶⁰ (17114212) Graff 2007 ⁸⁹ (17895439) Community based occupational therapy program vs. Waitlist for occupational therapy Medium Explanatory	Adjusted treatment difference (95% CI) Effect size Mean (SD) Caregiver quality of life: Dementia Quality of Life Instrument 6 weeks	0.7 (0.5 to 0.9) 1.2 (effect size)	4.0 (0.6)	3.4 (0.7)	< 0.0001
Graff 2006 ⁶⁰ (17114212) Graff 2007 ⁸⁹ (17895439) Community based occupational therapy program vs. Waitlist for occupational therapy Medium Explanatory	Adjusted treatment difference (95% CI) Effect size Mean (SD) Caregiver depression: Center for Epidemiologic Studies Depression Scale CES-D 6 weeks	7.6 (5.3 to 9.7) 1.3 effect size	5.8 (4.8)	12.6 (8.5)	< 0.0001
Farran 2004 ⁶² (41552352) Caregiver skill building (CSB) vs. Information and support group (ISO) Medium Explanatory USA	Regression coefficient (SE) Caregiver depression: Center for Epidemiological Studies of Depression scale (CES-D) 18 months	0.034 (0.032)	NR	NR	0.707
Farran 2004 ⁶² (41552352) Caregiver skill building (CSB) vs. Information and support group (ISO) Medium Explanatory USA	Regression coefficient (SE) Caregiver skills: Behavior Management Skill-Revised (BMS-R) 18 months	-0.006 (0.010)	NR	NR	0.538
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care Medium Explanatory USA	Adjusted mean difference (95% CI) Mean (SD) baseline Mean (SD) 3 month Caregiver self-efficacy: managing ADL dependency 3 months	.03 (-.03; .08)	.81 (.33) baseline .93 (.18) 3 months	.80 (.34) baseline .90 (.21) 3 months	0.375
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care Medium Explanatory USA	Adjusted mean difference (95% CI) Mean (SD) baseline Mean (SD) 3 month Caregiver self-efficacy: managing IADL dependency 3 months	.01 (-.03; .05)	.87 (.30) baseline .96 (.15) 3 months	.87 (.26) baseline .95 (.14) 3 months	0.704
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care Medium Explanatory USA	Adjusted mean difference (95% CI) Mean (SD) baseline Mean (SD) 3 month Caregiver self-efficacy: managing behaviors 3 months	.03 (-.03; .10)	.77 (.27) baseline .84 (.24) 3 months	.74 (.32) baseline .80 (.27) 3 months	0.314
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care Medium Explanatory USA	Adjusted mean difference (95% CI) Mean (SD) baseline Mean (SD) 3 month Caregiver upset: managing ADL dependency 3 months	-.06 (-.16; .03)	.26 (.35) baseline .25 (.34) 3 months	.29 (.36) baseline .34 (.37) 3 months	0.156
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care Medium Explanatory USA	Adjusted mean difference (95% CI) Mean (SD) baseline Mean (SD) 3 month Caregiver upset: managing IADL dependency 3 months	-.02 (-.10; .07)	.17 (.30) baseline .17 (.29) 3 months	.22 (.33) baseline .22 (.32) 3 months	0.663
Gitlin 2001 ⁹³ (11220813) Home Environmental Intervention vs. usual care	Adjusted mean difference (95% CI) Mean (SD) baseline	-0.02 (-0.09; .05)	.48 (.27) .43 (.31)	.47 (.30) .45 (.29)	0.501

Study (PMID) Comparison Caregiver Type RoB Category* Country	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Medium Explanatory USA	Mean (SD) 3 month Caregiver upset: managing behaviors 3 months				

Abbreviations: ADL= Activities of Daily Living; CI= Confidence Interval; IADL= Instrumental Activities of Daily Living; MMSE= Mini-Mental State Examination; NR=Not Reported; PMID=PubMed Identification Number; QOL-AD= the Quality of Life in Alzheimer's; SD= Standard Deviation; SE= Standard Error

Table E-6. Summary of strength of evidence for PLWD outcomes: psychosocial interventions for caregiver well-being

Outcome Comparison	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Depression Usual care	6-52 weeks	4 RCTs (n=592)	1 found benefit 3 found no difference	Medium	Inconsistent	Indirect	Imprecise	Insufficient
Depression Attention control	52 weeks	1 RCT n=330	1 found benefit 0 found no difference	Medium	Unknown	Indirect	Imprecise	Insufficient
Function Usual care	6-12 weeks	3 RCTs (n=545)	2 found benefit 1 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Function Usual care	6-12 months	3 RCTs (n=540)	0 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Function Attention control	4-12 months	2 RCTs (n=490)	1 found benefit 1 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Neuropsychiatric symptoms Usual care	3-6 months	4 RCTs (n=740)	1 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Neuropsychiatric symptoms Attention control	4-6 months	2 RCTs (n=227)	2 found benefit 0 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Neuropsychiatric symptoms Attention control	12-18 months	3 RCTs (n=916)	0 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Quality of life Usual care	6 weeks	1 RCT (n=135)	1 found benefit 0 found no difference	Medium	Unknown	Indirect	Imprecise	Insufficient
Quality of life Usual care	6-24 months	5 RCTs (n=936)	0 found benefit 5 found no difference	Medium	Inconsistent	Indirect	Imprecise	Insufficient
Quality of life Attention control	4-12 months	2 RCTs (n=490)	1 found benefit 1 found no difference	Medium	Consistent	Indirect	Imprecise	Insufficient
Social support Usual care	6 months	1 RCT (n=108)	0 found benefit 1 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Social support Attention control	6 months	1 RCT (n=250)	0 found benefit 1 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Health care usage Usual care	4-24 months	4 RCTs (n=652)	1 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Health care usage Attention control	6 months	1 RCT (n=250)	0 found benefit 1 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Harms – adverse events Usual care	52 weeks	1 RCT (n=141)	0 found benefit 1 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Institutionalization Usual care	6-12 months	2 RCTs (n=288)	1 found benefit 1 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Institutionalization Attention control	6-18 months	2 RCT (n=545)	0 found benefit 2 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient

Abbreviations: n=Number; RCT= Randomized Controlled Trial

Table E-7. Summary of strength of evidence for caregiver outcomes: psychosocial interventions

Outcome Comparison	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Anxiety Attention control	24-52 weeks	2 RCT N=358	0 found benefit 2 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Anxiety Usual care	6 weeks	1 RCT n=170	1 found benefit 0 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Anxiety Usual care	2 years	1 RCT n=260	0 found benefit 1 found no difference	Low	Unknown	Direct	Imprecise	Insufficient
Burden of care Attention control	8-24 weeks	5 RCT N= 725	3 found benefit 2 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Burden of care Usual care	6-16 weeks	3 RCT N=680	0 found benefit 3 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Burden of care Usual care	24-52 weeks	5 RCT N=561	2 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Depressive symptoms Attention control	8-24 weeks	7 RCT n= 976	6 found benefit 1 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Depressive symptoms Attention control	52-72 weeks	3 RCT n=916	0 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Depressive symptoms Usual care	6-24 weeks	5 RCT N=852	2 found benefit 3 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Depressive symptoms Usual care	52-104 weeks	5 RCT N=1073	2 found benefit 3 found no difference	Medium	Inconsistent	Indirect	Imprecise	Insufficient
Caregiving bother/distress/affect Attention control	4-12 months	5 RCT N=952	3 found benefit 2 found no difference Have means SD	Medium	Inconsistent	Direct	Imprecise	Insufficient
Caregiving bother/distress/affect Usual care	6-24 weeks	4 RCT N=702	1 found benefit 3 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Health care usage Attention control	3-6 months	2 RCT n=359	1 found benefit 1 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Health care usage Usual care	4-24 months	2 RCT n=251	0 found benefit 2 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Caregiving knowledge Usual care	24 weeks	1 RCT n=167	1 found benefit 0 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Quality of life Attention control	8 weeks	1 RCT n=132	1 found benefit 0 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Quality of life Attention control	24-52 weeks	2 RCT n=541	0 found benefit 2 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Quality of life Usual care	6-24 weeks	7 RCT n=843	4 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Quality of life Usual care	9 months to 2 years	4 RCT N=652	1 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Relationship with PLWD Attention control	20-52 weeks	3 RCT n=652	1 found benefit 2 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Caregiving confidence Attention control	20-24 weeks	3 RCT N=428	2 found benefit 1 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Caregiving confidence Usual care	12-24 weeks	3 RCT N=582	1 found benefit 2 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Caregiving skill Attention control	6-18 months	2 RCT N=545	0 found benefit 2 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient

Caregiving skill Usual care	6-16 weeks	2 RCT N=407	1 found benefit 1 found no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Caregiving skill Usual care	24-52 weeks	3 RCT N=444	0 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Stress Attention	6 months	1 RCT N=184	1 found benefit 0 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Stress Usual care	6-52 weeks	3 RCT n=486	0 found benefit 3 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Social support Attention control	6 months	1 RCT N=250	1 found benefit 0 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Social support Usual care	12-24 weeks	2 RCT n=188	0 found benefit 2 found no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient
Caregiving coping Usual care	24 weeks	1 RCT n=167	1 found benefit 0 found no difference	Medium	Unknown	Direct	Imprecise	Insufficient

Abbreviations: n=Number; RCT= Randomized Controlled Trial

Evidence Map: Psychosocial Interventions for Caregiver Well-Being

Table E-8. Characteristics of evidence map studies: psychosocial interventions for caregiver well-being

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Meichsner, 2019 ⁹⁵ (31384663) Germany Small Sample	CBT intervention delivered via internet, weekly message exchanges with therapist, 8 weeks	No treatment	Community 2 x 3 RCT Caregivers	Dementia, unspecified	N=37 Age: yes Sex: yes Race: no Education: no	NR	N=37 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes	NA	NA	Post intervention, 3 months post intervention	NA	CES-D CGS Burden of care Emotional wellbeing
Orgeta, 2019 ⁹⁶ (31609689) England Pilot	Weekly individual behavioral activation intervention delivered at home, 8 weeks, 1-hour sessions	Treatment as usual	Community RCT Caregiver-PLWD Dyads	Diagnosis of dementia of any type, defined by an MMSE score of ≥ 18	N=63 Age: yes Sex: yes Race: yes Education: yes	Household status	N=63 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Health Status: yes	NA	NA	3 months after start of treatment, 6 months after start of treatment	CSDD BADLS DEMQOL (self and proxy) EQ-5D (self and proxy) NPI	SF-12 HADS EQ-5D

Ta Park, 2019 ⁹⁷ (31609689) (Embase 629157769) ⁹⁸ United States Pilot	Weekly group sessions of Vietnamese-language cognitive-behavioral skills evidence-based program, 4 weeks, 120-minute sessions	Received educational materials on dementia	Community RCT Caregivers	Dementia or neurocognitive disorder, memory loss confirmed by Montreal Cognitive Assessment if in doubt	NR	NR	N=60 Sex: yes Race: yes Education: yes Relation to PLWD: Yes	Living with PLWD: yes Employment Status: yes	NA	NA	3 months post enrollment in study	NR	CES-D RMBPC (Conditional Bother Subscale)
Terracciano 2019 ⁹ (31866419) United States High ROB	Powerful Tools for Caregivers educational program Weekly 2-hours classes for 6 weeks	Waitlist control	Community-based RCT N=60 informal caregivers	Dementia diagnosis BPSD symptoms (assessed with RMBPC)	N=60 Age: yes Sex: yes Race: no Education: no	NR	N=60 Age: yes Sex: yes Race: yes Education: no Relation to PLWD: yes	Duration: yes Living with PLWD: no Employment Status: yes	NA	NA	6 weeks	RMBPC CMAI	ZBI CES-D NPI Caregiver satisfaction
Uyar 2019 ¹⁰ (no PMID) Turkey High ROB	Dementia Care and Support Program Multicomponent intervention 8 sessions over 16 weeks	Waitlist control	Community-based RCT N=61 informal caregivers	Dementia diagnosis Physician-diagnosed	NR	NR	N=61 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: no Living with PLWD: no Employment Status: yes	NA	NA	16 weeks	QOL-AD NPI-S	ZBI SF-36 BDI Beck Anxiety Inventory
Wawrziczny 2019 ¹¹ (29665714) France High ROB	Individualized psychoeducation for 7 weeks, 90 min sessions	Usual support	Community Quasi-experimental Caregivers	Physician diagnosed Alzheimer's, mixed, Lewy body, or frontotemporal dementias	N=102 Age: yes Sex: no Race: no Education: no	Household characteristics SES	N=102 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment: yes	NA	NA	7 weeks	NR	Caregiving self-esteem, distress Self-efficacy Depression Impact on finances Self-rated health
Whitlatch 2019 ⁹⁹ (29171296) US High ROB	Psychoeducation Support, Health, Activities, Resources, and Education (SHARE) Program Dyadic counseling for early stage dementia 6 sessions	Education One 90-minute session	Home-based setting RCT N=128 dyads	Dementia diagnosis	N=128 Age: yes Sex: yes Race: no Education: yes	NR	N=128 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: yes Training: no	NA	NA	2 months	NR	Preferences for Care Tasks Scale Service Availability Measure Dyadic Relationship Scale Care-related Agreement Scale Emotional-Intimacy Disruptive Behavior Scale Dementia Quality of Life

Yoo, 2019 ¹⁰⁰ (30938110) Korea Small Sample	Psycho-educational intervention, 4 in-person sessions delivered over 8-10 weeks, 60 minutes sessions, 1 st session delivered to group, subsequent sessions individual	No treatment	Community Multicenter RCT Caregivers	Possible and probable Alzheimer's disease per NINCDS-ADRDA and DSM-V criteria, a CDR score of 0.5-2.0 and an MMSE score of 10-26, with BPSD within last 6 months to one year	NR	NR	N=76 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	NR	NA	NA	2 weeks post intervention	NA	ZBI (Korean), GDS Philadelphia Geriatric Center Morale Scale (PGCMS) CGA-NPI (burden) Positive Affect and Negative Affect Schedule (PANAS), Self-Compassion Scale (SCS)
Boots 2018 ¹² (30006327) (27142676) ¹⁰¹ (29258980) ¹⁰² Netherlands High ROB	8-week, blended care self-management Partner in Balance program Group modified mindfulness-based cognitive therapy, 7 2-hour sessions over 10 weeks	Waitlist control	Community-based RCT N=81 informal caregivers	Dementia diagnosis	NR	NR	N=81 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Living with PLWD: yes	NA	NA	8 weeks	NR	Caregiver Self-Efficacy Scale CESD Pearlin Mastery Scale Investigation Choice Experiments for the Preferences of Older People HADS
De Oliveira, 2018 ¹⁰³ (30035341) Brazil Small Sample	Activity-based intervention (TAP) delivered face-to-face to dementia PLWD and their caregivers, 8 one-hour sessions over a 3-month period	Psychoeducation intervention delivered face-to-face to dementia PLWD and their caregivers, 8 sessions over a 3-month period	Community RCT Caregiver-PLWD dyads	Diagnosis of dementia by physician, MMSE score <24, and presence of at least three types of neuropsychiatric symptoms	N=21 Age: yes Sex: yes Race: no Education: yes	NR	N=21 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	NR	NA	NA	Post intervention	NPI-C	ZBI
Kales, 2018 ¹⁰⁴ (29747583) United States Pilot	Use of WeCareAdvisor tool (for assessment, management, and tracking of BPSD) for one month, with weekly check-in calls of 20 minutes max	No Treatment	Community RCT Caregivers	Clinical diagnosis of dementia or MMSE<24	N=57 Age: yes Sex: yes Race: yes Education: no	NR	N=57 Age: yes Sex: yes Race: yes Education: yes Relationship: yes	NR	NA	NA	Post intervention	Functional Dependence	Confidence in Using Activities, Caregiver Distress (NPI-Q) Caregiver Stress CES-D ZBI Negative Communication Scale

Meichsner, 2018 ¹⁰⁵ (30597537) Germany Pilot	Telephone-based CBT intervention including grief module delivered via 12 50-minute individual therapy sessions over six months	Written information on dementia and caregiving	Community RCT Caregivers	Diagnosis of dementia	N=273 Age: yes Sex: yes Race: no Education: no	NR	N=273 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: yes Living with PLWD: yes	NA	NA	Post intervention, six months post intervention	NR	Caregiver Grief Scale (coping)
Novelli, 2018 ¹⁰⁶ Brazil Pilot	Activity-based intervention (TAP), 8 sessions in the home over 4-month period	Usual care	Community RCT Caregiver-PLWD dyads	Previous diagnosis of dementia per NIAAA criteria and able to perform at least 2 basic ADLs, with presence of ≥ 2 BPSD in last 30 days	N=30 Age: yes Sex: yes Race: no Education: yes	NR	N=30 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	NR	NA	NA	Post Intervention	NPI (frequency, severity, and carer distress related to behaviors), Quality of Life Scale	NPI Distress ZBI Quality of Life Scale
Spalding-Wilson 2018 ¹⁴ (30258974) US High ROB	Two day-intervention using techniques aimed at fostering self-care for caregivers and improving communication between caregivers	Waitlist	Community-based RCT N=104 informal caregivers	Dementia diagnosis	N=104 Age: no Sex: no Race: no Education: no	NR	N=104 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: no	Duration: yes Living with PLWD: yes	NA	NA	1, 3, 6 months	NR	PSS CBI BDI-II Beck Anxiety Inventory
Van Knippenberg, 2018 ¹⁰⁷ (30126766) Netherlands Pilot	Experience sampling method (ESM) self-monitoring 3 days/week for 6 weeks, with personalized face-to-face feedback received every 2 weeks OR ESM self-monitoring 3 days/week for 6 weeks with no ESM feedback (semi-structured well-being interview instead)	Usual Care	Community RCT Caregivers	Dementia of any subtype or stage	NR	NR	N=76 Age: yes Sex: yes Race: no Education: yes Relation to PLWD	Living with PLWD: yes	NA	NA	Post intervention, 2 months post intervention	NEO Five Factor Inventory (neuroticism) Utrecht Coping List NPI	Short Sense of Competence Questionnaire Pearlin Mastery Scale CES-D PSS HADS Affect (positive, negative)
Wilz 2018 ¹⁵ (29190357) (Embase 626130265) ¹⁰⁸ (Embase 626123635) ¹⁰⁹ Germany High ROB	Tele.TAnDem Intervention Telephone-based CBT for family caregivers 12 50-minute sessions	Usual care	Community-based RCT N=273 informal caregivers	Dementia diagnosis	N=273 Age: yes Sex: yes Race: no Education: no	NR	N=273 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: no Training: no	NA	NA	6 months	NR	CESD BDI Emotional wellbeing Gießen Body Complaints List BEHAVE-AD coping item

Callahan 2017 ¹⁶ (27893087) US High ROB	Collaborative care and occupational therapy 24 sessions over 2 years	Usual care	Home-based setting RCT N=180 dyads	Dementia diagnosis	N=180 Age: yes Sex: yes Race: yes Education: yes	NR	N=180 Age: yes Sex: no Race: no Education: no Relation to PLWD: yes	NR	NA	NA	24 months	NR	ADCS ADL SPPB
Lok, 2017 ¹⁰ Turkey Pilot	"First You Should Get Stronger" program delivered to individuals weekly, 7 weeks, 45-minute sessions	No Treatment	Community RCT Caregivers	Diagnosis of dementia per DSM V criteria	N=40 Age: yes Gender: yes Race: no Education: No	NR	N=40 Age: yes Sex: yes Race: no Education: yes Relationship: no	Duration	NA	NA	Post intervention	NR	IC: Zarit Burden Interview, Health Life Style Behavior Scale II
O'Connor, 2017 ¹¹ Australia Pilot	Activity-based intervention (TAP), 8 home visits over 4-month period	Three phone calls to caregivers over 4-month period on dementia education	Community RCT Caregiver-PLWD dyads	Diagnosis of frontotemporal dementia according to current diagnostic criteria, presence of behavioral disturbances over the past month as rated by the carer, score of >3.31 on Informant Questionnaire on Cognitive Decline for the Elderly, and able to perform at least 2 basic ADLs	N=20 Age: yes Sex: yes Race: no Education: yes	no	N=20 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Living with PLWD: yes	NA	NA	Post Intervention	NPI-C (frequency, severity, and carer distress related to behaviors), DAD (ADLs and IADLs), EruoQoI5-D	Vigilance Items (carer time feeling "on duty" and "doing things" for PLWD)
Supiano, 2017 ¹² United States Small Sample	Weekly Complicated Grief Group Therapy (CGGT), 120-minute sessions, 16 weeks, with weekly homework on grief status and emotions	Weekly phone calls, 30 minutes duration, discussing caregiver's grief status and emotions	Community Randomized attention control Caregivers	Dementia, unspecified	NR	NR	N=35 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	NR	NA	NA	Weekly throughout trial, post intervention	NR	Brief Grief Questionnaire (BGQ) Inventory of Complicated Grief-revised (ICG-r) Clinical Global Impressions – Severity Scale (CGI-S) Clinical Global Impressions-Improvement Scale (CGI-I)

Whitlatch 2017 ¹⁷ (29171296) US High ROB	Psychoeducation Support, Health, Activities, Resources, and Education (SHARE) Program Dyadic counseling for early stage dementia 6 sessions	Education One 90-minute session	Home-based setting RCT N=128 dyads	Dementia diagnosis	N=128 Age: yes Sex: yes Race: yes Education: yes	NR	N=128 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: yes Training: no	NA	NA	2 months	NR	Preferences for Care Tasks Scale Service Availability Measure Dyadic Relationship Scale Care-related Agreement Scale Emotional-Intimacy Disruptive Behavior Scale Dementia Quality of Life
Nunez-Naveira, 2016 ¹³ Denmark, Poland, Spain Pilot	UnderstAID application--with information, skills training, social network forum, and daily reminders--made available via mobile device and/or PC for three months with periodic check-ins by research staff	No Treatment	Community, RCT Caregivers	Diagnosis of dementia by a specialist or neurologist per criteria of CMBD-10, DSM-IV, NINDSADRDA	NR	NR	N=77 Age: yes Sex: yes Race: no Education: no Relationship: yes	Duration: yes Payment: yes Health Status: yes Employment Status: yes	NA	NA	Post intervention	NR	CES-D Revised Caregiving Satisfaction Scale
Sepe-Monti, 2016 ¹⁴ Italy Pilot	Six weekly group sessions of psycho-education program delivered by a psychologist, 2-hour sessions	Two group sessions providing medical information and discussing management of care recipient behaviors, 2-hour sessions, one session in week two of trial, one session in week four; informational leaflet provided	Community Multicenter RCT Caregivers	Probable or possible AD according to NINCDS-ADRDA criteria	N=381 Age: yes Sex: yes Race: no Education: no	NR	N=164 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Living with PLWD: yes	NA	NA	Two weeks post intervention, six months post intervention	NR	Coping Orientations to Problems Experienced- New Italian Version (COPE-NIV) Caregiver Burden Inventory (CBI) NPI NPI-Distress Scale SF-12 (physical and mental health) CES-D STAI-Y
Taati 2016 ²² (no PMID) Iran High ROB	Support group for family caregivers 8 ~2-hour sessions	Nonspecified control	Community-based setting RCT N=80 informal caregivers	Dementia diagnosis	NR	NR	N=80 Age: no Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: no Employment Status: yes Training: no	NA	NA	8 weeks	NR	DASS-21 (depression, anxiety, stress level)

Zauszniewski, 2016 ¹¹⁵ (27500286) (26735507) ¹¹⁶ United States Pilot	40-minute individual resourcefulness training with daily journaling or recording as practice/reinforcement methods for four weeks, with random assignment to choice of method vs. assignment of method	Journaling or recording without learning training skills	Community Modified partially randomized preference trial Caregivers	Dementia, any form	N=138 Age: yes Sex: yes Race: no Education: no	NR	N=138 Age: yes Sex: yes Race: yes Education: yes Relationship: no	Duration: yes Living with PLWD: yes Health Status: yes	NA	NA	Post intervention, 6 weeks post intervention	NR	PSS DCS ESC
Au, 2015 ¹¹⁷ (25848237) China Small Sample	Individual psychoeducation by telephone for 4 weeks, 30-minute sessions, followed by 8 biweekly telephone sessions, 15-20 minutes each, of behavioral activation	Individual psychoeducation by telephone for 4 weeks, 30-minute sessions, followed by 8 biweekly telephone sessions, 15-20 minutes each, of general discussion of psychoeducation and related information	Community RCT Caregivers	Alzheimer's diagnosis for at least 3 months	N=59 Age: yes Sex: yes Race: no Education: no	NR	N=62 Age: yes Sex: yes Race: no Education: no Relationship: yes	Duration: yes Living with PLWD: yes	NA	NA	4 weeks (i.e., after first arm), 6 months (i.e., after second arm)	NR	CES-D RAS
Au 2015 ¹¹⁸ (no PMID) Hong Kong High ROB	Psychoeducation with telephone-assisted behavioral activation intervention 8 biweekly sessions	Psychoeducation only	Community-based setting RCT N=96 informal caregivers	Dementia diagnosis	NR	NR	N=96 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	4 months	NR	CES-D
Blom 2015 ²³ (no PMID) (23305463) ¹¹⁹ (25648589) ¹²⁰ Netherlands High ROB	Mastery of Dementia Intervention Problem solving, relaxation, cognitive restructuring, communication 8 lessons + booster session	Education via e-bulletins	Home-based setting RCT N=251 dyads	Dementia diagnosis	NR	NR	N=251 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: no Living with PLWD: yes Employment Status: no Training: no	NA	NA	3, 6 months	NR	CES-D HADS-A

Chen, 2015 ¹²¹ (25515800) Taiwan Small Sample	Biweekly in-person education sessions about problem-solving skills, dementia, social resources, and emotional support, six sessions over 3 months (session length not specified)	Biweekly phone calls for usual clinical management (phone call length not specified)	Community or Clinic RCT Caregivers	Dementia, unspecified	N=unclear Age: yes Sex: yes Race: no Education: no	NR	N=46 Age: yes Sex: yes Race: no Education: yes Relationship: yes	Living with PLWD: yes	NA	NA	2 weeks post intervention	NR	WCCL-R Caregiver Burden Inventory, RMBPC
Chiu 2015 ²⁴ (25615434) Canada High ROB	Problem-solving technique-based intervention 3 1-hour sessions over 3-4 weeks	Nonspecified control	Home-based setting RCT N=56 informal caregivers	Dementia diagnosis	NR	NR	N=56 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: no Employment Status: yes Training: no	NA	NA	4 weeks	NR	ZBI Pearlin Self-Mastery and Competency Scales PSS
Cristancho-Lacroix, 2015 ¹²² (25967983) (25263541) ¹²³ France Small Sample	Web-based psycho-educational program delivered in 15-30-minute weekly sessions for 12 weeks	Usual care only	Day care center geriatric unit RCT Caregivers	Alzheimer's diagnosis per DSM-IV criteria	N=49 Age: no Sex: no Race: no Education: no	NR	N=49 Age: yes Sex: yes Race: no Education: yes Relationship: yes	Living with PLWD: yes Health status: yes	NA	NA	Post intervention, 3 months post intervention	NR	RSCS RMBPC ZBI NHP
Gallagher-Thompson 2015 ²⁵ (25590939) US High ROB	Psychoeducation Fotonovela for reducing stress and depression in Latino caregivers	Usual care with basic information	Home-based setting RCT N=147 dyads	Dementia diagnosis	N=147 Age: yes Sex: no Race: yes Education: yes	NR	N=147 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: yes Training: no	NA	NA	4, 6 months	NR	CES-D RMBPC Reaction
Gaugler, 2015 ¹²⁴ (25751083) United States Pilot	Individual psychosocial intervention (other family members or care decision makers present per participant discretion), six sessions, weekly for three weeks, then monthly for 3 months, 60-120-minute sessions	No Treatment	Community RCT Caregivers	Dementia	N=36 Age: yes Sex: yes Race: yes Education: yes	NR	N=36 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Employment Status: yes	NA	NA	Post intervention, 4 months post intervention	NR	ZBI CES-D GDS NPI-Q Role overload PSS
Kamkhagi, 2015 ¹²⁵ (no PMID) Brazil Small Sample	Psychological intervention delivered via weekly group sessions, 90-minute sessions, 14 weeks	Non-psychologically-oriented intervention delivered via weekly group sessions, 90-minute sessions, 14 weeks	Community RCT Caregivers	Mild or moderate Alzheimer's disease	NR	NR	N=37 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Living with PLWD: yes	NA	NA	Post intervention	NR	ZBI BDI WHO-QoL Scale

Losada 2015 ²⁶ (21061414) Spain High ROB	1. CBT 2. ACT (Acceptance and Commitment Therapy) to reduce depression in caregivers 8 weekly 90-minute sessions	3. Minimal support control	Community-based setting RCT N=135 informal caregivers	Dementia diagnosis	NR	NR	N=135 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment Status: yes Training: yes	NA	NA	2 months	Barthel Index RMBPC	CES-D Profile of Mood States
Paun, 2015 ¹²⁶ (24510968) United States Pilot	12 weekly group sessions of chronic grief management intervention, 60-90-minute sessions	No treatment (Two check-in calls at 1.5 months and 4.5 months)	Long-term care facilities Pre-test/multiple post-test quasi-experimental design Caregiver	Diagnosis of AD or a related dementia	NR	NR	N=93 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	NR	NA	NA	Post intervention, 3 months post intervention	NR	FPCR (Family Perception of Caregiver Role – role stress) MMCGI (grief in caregivers of persons with AD) CES-D
Abouafia-Brakha, 2014 ¹²⁷ (no PMID) Brazil Small Sample	Weekly 90-minute group sessions of psychoeducation and cognitive behavioral therapy, or weekly 60-minute group sessions of psychoeducation alone, 8 weeks	No Control Group	Community Semi-Random Comparative Trial Caregivers	Clinical diagnosis of possible or probable Alzheimer's disease in the moderate or severe stage, with diagnosis and staging based on the DSM-IV, the FAST, and the MMSE (Portuguese version)	NR	NR	N=35 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: yes	NA	NA	Post intervention	NR	PSS ZBI BDI STAI-S STAI-T Portuguese version
Arango-Lasprilla 2014 ²⁹ (24550547) Spain High ROB	Group-based cognitive behavioral intervention for caregiver mental health	Educational control	Community-based setting RCT N=69 informal caregivers	Dementia diagnosis	NR	NR	N=69 Age: yes Sex: yes Race: yes Education: no Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	8 weeks 3 months	NA	PHQ-9 ZBI Satisfaction with Life Scale PSS
Dowling, 2014 ¹²⁸ (24113564) United States Small Sample	Weekly, one-on-one, positive affect skill-building intervention sessions delivered in person or via video-conference, 5 weeks, 1-hour sessions	Weekly, one-on-one, sessions delivered in person or via video-conference consisting of a thematic interview, 5 weeks, 1-hour sessions	Community Randomized attention control Caregivers	Frontotemporal dementia (FTD)	NR	NR	N=26 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes	NA	NA	One-week post intervention, One-month post intervention	NR	DES CES-D PSS NPI ZBI
Gonzalez, 2014 ¹²⁹ (25328754) United States Pilot	Weekly group resourcefulness trainings delivered for 6 weeks, 2 hours per training	Usual Care	Community RCT Caregivers	Probably diagnosis of AC using the NINCDS/ADRD criteria for probable AC	N=102 Age: yes Sex: yes Race: yes Education: no	NR	N=102 Age: yes Sex: yes Race: yes Education: yes Relationship: yes	Health status: yes	NA	NA	Within one-week post intervention, 12 weeks post intervention	RMBPC	STAI CESD CRS

Passoni 2014 ³² (24614271) Italy High ROB	1. Group-based cognitive behavioral intervention for caregiver mental health 2. Manual only	Control	Community-based setting RCT N=100 informal caregivers	Dementia diagnosis	N=100 Age: yes Sex: yes Race: no Education: no	NR	N=100 Age: yes Sex: yes Race: no Education: no Relation to PLWD: no	Duration: no Employment Status: no Training: no	NA	NA	6 months	NA	Anxiety and Depression Scale-Reduced Form Caregiver Need Assessment Family Strain Questionnaire
Bruvik 2013 ³³ (24348500) Norway High ROB	Tailored psychosocial support intervention for depression	Usual care	Home-based setting RCT N=230 dyads	Dementia diagnosis	N=230 Age: yes Sex: yes Race: no Education: yes	NR	N=230 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: yes Training: no	NA	NA	12 months	CSDD	GDS
Huang 2013 ³⁴ (23933422) Taiwan High ROB	Training program for caregivers for skills managing behavioral problems	Written instructions and telephone followup	Home-based setting RCT N=129 dyads	Dementia diagnosis	N=129 Age: yes Sex: yes Race: no Education: no	NR	N=129 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	3, 6 months	CMAI	Agitation Management Self-efficacy Scale Preparedness and Competence Scales
Judge 2013 ³⁵ (22899427) Judge 2011 (19808841) ¹³⁰ Judge 2011 (no PMID) ¹³¹ US High ROB	ANSWERS Intervention (Acquiring New Skills While Enhancing Remaining Strengths) Education and cognitive rehabilitation skills	Educational pamphlets	Home-based setting RCT N=128 dyads	Dementia diagnosis	N=128 Age: yes Sex: yes Race: yes Education: yes	NR	N=128 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: no Employment Status: yes Training: no	NA	NA	11, 19 weeks	NA	Pearlin Caregiver Mastery scales Depression Anxiety QOL Self-Esteem
Kajiyama 2013 ³⁶ (23461355) US High ROB	iCare Stress Management e-Training Program	Information only	Home-based setting RCT N=150 dyads	Dementia diagnosis	N=150 Age: yes Sex: no Race: no Education: no	NR	N=150 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	3 months	NA	CES-D PSS RMBPC Bother PQOL
Kuo 2013 ³⁷ (22778053) (27653753) ¹³² Kuo 2014 (no PMID) ¹³³ Taiwan High ROB	Training program to improve QOL and decrease depression 2 2-3-hour sessions 1 week apart	Attention control	Home-based setting RCT N=129 informal caregivers	Dementia diagnosis	NR	NR	N=129 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	2 weeks 3, 6 months	NA	HRQOL CES-D

Kwok, 2013 ¹³⁴ (24072965) China Small Sample	Psychoeducation for caregivers by telephone, twelve 30-minute weekly sessions, plus DVD containing educational information about dementia caregiving	DVD containing educational information about dementia caregiving	Community RCT Caregivers	Clinical diagnosis of dementia of any stage	N: no Age: no Sex: yes Race: no Education: no	NR	N=42 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	NR	NA	NA	Post intervention	NR	ZBI Revised Scale for Caregiving Self-efficacy
Moore 2013 ³⁸ (23916631) US High ROB	Pleasant Events Program Behavioral Activation therapy for improving physical and mental health in caregivers 6 weeks	Educational manual	Community-based setting RCT N=100 dyads	Dementia diagnosis	N=100 Age: no Sex: no Race: no Education: no	NR	N=100 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: no	Duration: yes Employment Status: no Training: no	NA	NA	6 weeks 12 months	NA	CES-D Health status Social Support Scale PANAS
Joling 2012 ³⁹ (22303473) (22876304) ¹³⁵ (24053631) ¹³⁶ (18208607) ¹³⁷ Netherlands High ROB	Family Meeting Intervention for depression and anxiety in caregivers Four sessions every 2-3 months	Usual care	Home-based setting RCT N=192 caregivers	Dementia diagnosis	N=192 Age: yes Sex: yes Race: no Education: no	NR	N=192 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: no Living with PLWD: yes Employment Status: no Training: no	NA	NA	3, 6, 9, 12 months	NA	CES-D HADS-A Caregiver Reaction Assessment SF-12
Chu 2011 ⁴¹ (20847363) Taiwan High ROB	Support group for caregiver burden and depression 12 weeks	Usual care	Home-based setting RCT N=85 dyads	Dementia diagnosis	NR	NR	N=85 Age: no Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: no Living with PLWD: yes Employment Status: no Training: no	NA	NA	3, 4 months	NA	BDI-II CBI
Davis, 2011 ¹³⁸ (20842759) United States Small Sample	Psychosocial intervention delivered by telephone, 10 contacts over 3 months: initial call and 7 weekly follow-up calls (60 minutes each), and 2 bimonthly termination calls (35-45 minutes each)	No Treatment	Community RCT Caregivers	Diagnosis of dementia	N=53 Age: yes Sex: no Race: no Education: no	NR	N=53 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	NR	NA	NA	Post intervention	NR	CES-D ZBI SF-36
Losada 2011 ⁴⁴ (21061414) Spain High ROB	CBT for caregiver behavioral activation and modifying dysfunctional thoughts 12 sessions	Usual care	Community-based setting RCT N=170 dyads	Dementia diagnosis	N=170 Age: yes Sex: no Race: no Education: no	NR	N=170 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	3 months	RMBPC	CES-D Dysfunctional Thoughts about Caregiving Questionnaire Leisure Time Satisfaction

Spijker 2011 ⁴⁵ (21358385) (19500421) ¹³⁹ (23290200) ¹⁴⁰ Netherlands High ROB	Systematic Care Program for Dementia	Usual care	Community health centers Cluster RCT	PLWD with suspected dementia and their caregivers Exclusion criteria	N=155 Intervention group 140 Control group Age: yes Sex: yes Education: yes Race % majority: yes	Household characteristics: yes	Age: yes Sex: yes Race % majority: yes Education: yes Relation to PLWD: yes	Living with PLWD: yes	Education: yes	NR	12 months	PLWD institutionalization NPI	Sense of competence Depression QoL Distress
Wilz 2011 ⁴⁸ (no PMID) (26311735) ¹⁴¹ (27792398) ¹⁴² (28428730) Germany High ROB	CBT for goal-attainment and treatment compliance 7 60-minute sessions over 3 months	1. Attention control 2. Usual care	Community-based setting RCT N=229 dyads	Dementia diagnosis	N=229 Age: yes Sex: no Race: no Education: no	NR	N=229 Age: yes Sex: no Race: no Education: no Relation to PLWD: no	Duration: yes Employment Status: no Training: no	NA	NA	3, 6 months	NA	Goal Attainment Scaling
Au, 2010 ¹⁴³ (no PMID) China Small Sample	Weekly CBT group intervention, 13 weeks, 2-hour sessions	No Treatment	Community RCT Caregivers	Presence of Alzheimer's Disease	NR	NR	N=37 Age: yes Sex: no Race: no Education: yes Relation to PLWD: yes	Living with PLWD, Employment: yes	NA	NA	Post intervention	NR	CES-D Revised Scale for Caregiving Self-Efficacy Chinese Way of Coping Questionnaire
Gallagher-Thompson, 2010 ¹⁴⁴ (359062574) United States Pilot	CBT skill training program delivered via DVD, 2.5 hours in length	General educational DVD program on dementia, 2.0 hours in length	Community RCT Caregivers	Significant memory lost or deterioration in cognitive abilities	N=68 Age: yes Sex: no Race: no Education: yes	NR	N=76 Age: yes Sex: yes Race: no Education: yes Relationship: yes	Health status: yes	NA	NA	Post intervention	NR	CES-D Positive Affect subscale RMBPC
Gitlin 2010 ¹⁴⁵ (20810376) US High ROB	COPE (Care of Persons with Dementia in their Environments) biobehavioral intervention 12 home or phone contacts over 4 months	Attention control	Home-based setting RCT N=237 dyads	Dementia diagnosis	N=237 Age: yes Sex: yes Race: yes Education: no	NR	N=237 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	4, 9 months	Functional Independence QOL-AD Agitated Behavior in Dementia	Perceived Change Index Caregiver confidence
Kurz 2010 ⁵¹ (19946869) Germany High ROB	Psychoeducation 7 90-minute biweekly sessions + boosters over 15 months	One session	Home-based setting RCT N=292 dyads	Dementia diagnosis	N=292 Age: yes Sex: yes Race: yes Education: no	NR	N=292 Age: yes Sex: yes Race: yes Education: no Relation to PLWD: yes	NR	NA	NA	5, 15 months	MMSE NPI ADCS-ADL	MADRS SF-12 Time spent caregiving Resource use QOL
Williams 2010 ⁵² (20978227) US High ROB	Psychoeducation video on coping skills for caregivers Telephone coaching 2 10-minute modules per week for 5 weeks	Waitlist	Home-based setting RCT N=116 informal caregivers	Dementia diagnosis	NR	NR	N=116 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: no Training: no	NA	NA	3, 6 months	NA	PSS PSQI STAI STAXI CES-D

Gavrilova 2009 ⁵³ (18814197) Russia High ROB	Psychoeducation 10/66 intervention basic education and training for managing problem behaviors 5 weekly 30- minute sessions	Usual care	Community- based setting RCT N=60 dyads	Dementia diagnosis	N=60 Age: yes Sex: yes Race: no Education: no	NR	N=60 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	6 months	NA	ZBI SRQ 20 WHOQOL- BREF NPI-Q DEMQOL
Dias 2008 ¹⁴⁶ (18523642) India Pilot	Support and guidance from Home Care Advisors team	Caregivers received only education and information on dementia; Intervention delayed for 6 months	Home-based RTC	PLWD diagnosed with mild to moderate dementia and their home caregivers Exclusion criteria	N= 33 intervention group 26 control group Age: yes Sex: yes Education: yes Race % majority: N	Household characteristics: yes Availability of paid help: yes Income: yes	Age: yes Sex: yes Rae % majority: no Education: yes Relation to PLWD: yes	NR	NR	Training: yes	6 months	Activities of daily living NPI	Caregiver mental health Caregiver burden
Gitlin, 2008 ¹⁴⁷ (18310553) (20847903) ¹⁴⁸ (19420314) ¹⁴⁹ United States Pilot	Activity-based intervention (TAP), 6 home visits and two 15-minute phone calls over 4 months	Tailored activity intervention administered upon conclusion of main trial (6 home visits and two 15-minutes phone calls over 4 months)	Community RCT Caregiver- PLWD dyads	Diagnosis of dementia by physician or MMSE score <24	N=60 Age: yes Sex: yes Race: yes Education: yes	NR	N=60 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Health status: yes	NA	NA	Post intervention	CSDD Quality of Life-AD Behavioral occurrences	ZBI Subjective burden Objective burden CES-D
Tremont, 2008 ¹⁵⁰ (20228893) (18838742) ¹⁵¹ Pilot	Psychosocial telephone intervention delivered on set schedule, completing 23 calls over one year, calls ranged from 15- 60 minutes, totaling approximately 12 hours	Standard Care	Community RCT Caregivers	Formal dementia diagnosis per DSM-IV criteria and CDR score of 1 or 2	N=60 Age: yes Sex: no Race: no Education: no	NR	N=60 Age: yes Sex: no Race: no Education: yes Relation to PLWD: no	Duration: yes Living with PLWD: yes Health Status: yes	NA	NA	Post intervention	NR	ZBI RMBPC GDS Self-Efficacy Scale SF-36
Gant, 2007 ¹⁵² (17545139) United States Pilot	Psychosocial intervention delivered via 10-session video series with workbook and 12 weekly telephone coaching sessions (phone call length not specified)	Education intervention delivered via booklet and 7 bi-weekly check-in calls (phone call length not specified)	Community RCT Caregivers	Diagnosis of dementia secondary to: Alzheimer's disease, Parkinson's disease, cerebrovascular disease, Huntington's disease, Korsakoff's disease, multiple sclerosis, or other	N=32 Age: yes Sex: no Race: no Education: no	NR	N=32 Age: yes Sex: yes Race: no Education: yes Relationship: yes	Living with PLWD: yes Employment status: yes	NA	NA	Post intervention	NR	RMBPC RSCSE PNAS Target complaints (upset, sad, frustrated, and irritated)

Glueckauf, 2007 ¹⁵³ (no PMID) United States Small Sample	Weekly cognitive-behavioral intervention delivered via at-home video with telephone follow-up: 7 weeks of 45-minute group phone sessions, followed by 5 weeks of one-hour individual phone sessions, 12 weeks	No treatment	Community RCT Caregivers	Medical diagnosis of Alzheimer's disease or another type of progressive dementia	N=24 Age: yes Sex: yes Race: yes Education: no	NR	N=24 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes	NA	NA	One-week post intervention	NR	Caregiver Appraisal Inventory CAI Caregiver Self-Efficacy CES-D
Marquez-Gonzalez 2007 ⁵⁷ (18074249) (22899425) ¹⁵⁴ Spain High ROB	Cognitive behavioral intervention for caregiver coping 8 weekly 2-hour sessions	Waitlist	Community-based setting RCT N=74 informal caregivers	Dementia diagnosis	NR	NR	N=74 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	8 weeks	MBPC (proxy)	CES-D Dysfunctional Thoughts about Caregiving Questionnaire
Ulstein 2007 ⁵⁸ (17986818) Norway High ROB	Psychosocial intervention for education about dementia and structured problem-solving 3-hour education program + 6 2-hour group meetings over 1 year	Usual care	Community-based setting RCT N=180 dyads	Dementia diagnosis	N=180 Age: no Sex: yes Race: no Education: no	NR	N=180 Age: no Sex: yes Race: no Education: no Relation to PLWD: yes	NR	NA	NA	1 year	NPI	Relative Stress Scale
Gonyea 2006 ⁵⁹ (17169938) US High ROB	Behavioral intervention to reduce caregiver distress, burden, and PLWD symptom severity 5 90-minute weekly sessions, 16-20 hours total training	Education only	Community-based setting RCT N=80 informal caregivers	Dementia diagnosis	NR	NR	N=80 Age: yes Sex: yes Race: yes Education: no Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	6 weeks	NPI	ZBI
Beauchamp 2005 ⁶¹ (no PMID) US High ROB	Multimedia support program delivered via internet	Waitlist	Community-based setting RCT N=299 informal caregivers	Dementia diagnosis	NR	NR	N=299 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	1 month	NA	CES-D STAI CSS Positive Aspects of Caregiving

Akkerman, 2004 ¹⁵⁵ (15106393) United States Small Sample	Weekly cognitive-behavioral group intervention, 9 weeks, 2-hour meetings	No Treatment	Community RCT Caregivers	Diagnosis of Alzheimer's	NR	NR	N=38 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes	NA	NA	Post intervention, 6 weeks post intervention	NR	HAMA BAI
Nobili, 2004 ¹⁵⁶ (15249851) Italy Pilot	60-minute home visit from a psychologist, separate 90-minute home visit from occupational therapist	Standard counseling via help line, information about services and supports	Community RCT Caregiver-PLWD Dyads	Clinical diagnosis of dementia made by a neurologist, geriatrician, or psychiatrist	N=69 Age: yes Sex: yes Race: no Education: yes	NR	N=69 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: yes	NA	NA	6 months post intervention, 12 months post intervention	ADLs SBI-C (problem behaviors)	RSS
Burgio 2003 ⁶³ (12937335) (16625937) ¹⁵⁷ US High ROB	Skills training group workshop + 16 home-based treatment sessions over 12 months	Educational materials and telephone support	Home-based setting RCT N=118 dyads	Dementia diagnosis	N=118 Age: yes Sex: yes Race: yes Education: yes	NR	N=118 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Employment Status: yes Training: no	NA	NA	6 months	MMSE RMBPC (proxy)	Positive Aspects of Caregiving RMBPC both Lubben Social Network Index Leisure Time Satisfaction CES-D Desire to Institutionalize
Burns 2003 ⁶⁴ (12937333) (19290751) ¹⁵⁸ US High ROB	REACH PLWD behavior management and caregiver stress management	Targeted educational materials	Home-based setting RCT N=167 dyads	Dementia diagnosis	N=167 Age: yes Sex: yes Race: yes Education: yes	NR	N=167 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Employment Status: yes Training: no	NA	NA	2 years	RMBPC	General wellbeing scale CES-D
Coon 2003 ⁶⁵ (14570964) US High ROB	Psychoeducation skill training for 1. anger or 2. depression management over 3-4 months	Waitlist	Home-based setting RCT N=169 dyads	Dementia diagnosis	N=169 Age: no Sex: no Race: no Education: no	NR	N=169 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	4, 7 months	NA	STAXI Multiple Affect Adjective Checklist Hostility and Depression subscales Ways of Coping Checklist Positive Coping subscales BDI Caregiving Self Efficacy Scale

Gitlin 2003 ⁶⁷ (12937332) (15860476) ¹⁵⁹ (17595426) ¹⁶⁰ (17563191) ¹⁶¹ US High ROB	REACH Home Environmental Skill-Building Program 5 home-based sessions + 1 telephone session	Usual care	Home-based setting RCT N=255 dyads	Dementia diagnosis	N=255 Age: yes Sex: yes Race: no Education: no	NR	N=255 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Employment Status: no Training: no	NA	NA	6 months	NA	RMBPC Upset Caregiving Mastery Index Task Management Strategy Index
Hebert 2003 ⁶⁸ (12496309) (15841829) ¹⁶² Canada High ROB	Psychoeducation Group Program on stress appraisal and coping 15 2-hour weekly sessions	Usual care	Home-based setting RCT N=158 dyads	Dementia diagnosis	NR	NR	N=158 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: yes Training: no	NA	NA	4 months	RMBPC	RMBPC reaction ZBI STAI Bradburn Revised Affective Scale Inventory of Socially Supportive Behavior Ilfeld Psychiatric Symptoms Index
Huang, 2003 ¹⁶³ Taiwan Pilot	Two-session in-home caregiver training, sessions separated by one week, 2-3-hour sessions; training accompanied by follow-up phone calls once a week for two weeks; accompanied by further weekly follow-up consultation calls beginning three weeks after completion of two-session program	Written educational materials with social telephone phone calls once a week for two weeks; accompanied by weekly follow-up social phone calls beginning three weeks after experimental group completed two-session program	Community RCT Caregivers	Diagnosis of dementia by psychiatrist or neurologist	N=59 Age: yes Sex: yes Race: no Education: yes	NR	N=59 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	NR	NA	NA	Three weeks after experimental group completed two-session training program; again three months after training completion	Cohen-Mansfield Agitation Inventory (CMAI; problematic behaviors)	Agitation Management Self-Efficacy Scale (AMSS)
Martin-Cook, 2003 ¹⁶⁴ (14682086) United States Pilot	Weekly group psychoeducational intervention, 4 weeks, 2-hour sessions	Standard Care	Community RCT Caregivers	Diagnosis of dementing illness with behavioral disturbance	NR	NR	N=37 Age: no Sex: no Race: no Education: no Relationship: no	NR	NA	NA	Two weeks post intervention, 10 weeks post intervention	NR	Caregiver Resentment Scale CES-D NPI
Fung 2002 ⁶⁹ (12037799) Hong Kong High ROB	Support group for family caregivers 12 1-hour sessions	Usual care	Community-based setting RCT N=52 dyads	Dementia diagnosis	NR	NR	N=52 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment Status: yes Training: no	NA	NA	3 months	NA	NPI-D WHOQOL-BREF Mental health service use

Garand, 2002 ¹⁶⁵ (12143075) United States Pilot	Psychoeducational nursing intervention delivered through two in-home visits, 3 hours each, followed by twice weekly phone calls for 6 months (phone call length not specified)	Psychological support, traditional dementia information, and referral to community-based services, delivered through two in-home visits, 3 hours each, followed by twice weekly phone calls for 6 months (phone call length not specified)	Community RCT Caregivers	Dementia, diagnosed or undiagnosed	N=39 Age: yes Sex: yes Race: yes Education: no	NR	N=39 Age: yes Sex: yes Race: yes Education: yes IC Relation to PLWD: yes	Health Status: yes Employment Status, Training: yes	NA	NA	One week after in-home phase, again after 6 months of biweekly phone calls	NR	Total mood disturbance
Stolley, 2002 ⁷⁰ (11954669) US High ROB	Psychoeducational based on progressively lowered stress theory	Basic dementia education	Community RCT Caregivers	Physician-diagnosed dementia; 66% AD	NR	NR	N=241 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes	NA	NA	12 months	NA	ZBI Mastery
Wright 2001 ⁷¹ (11885210) US High ROB	Nurse education and counseling program Home visits and phone calls	Attention control	Community-based setting RCT N=93 informal caregivers	Dementia diagnosis	N=93 Age: yes Sex: yes Race: no Education: no	NR	N=93 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Employment Status: no Training: no	NA	NA	2, 6, 12 weeks 6, 12 months	Blessed Dementia Rating Scale CMAI	Caregiving Hassle Scale CES-D Multilevel Assessment Inventory
Marriott, 2000 ¹⁶⁶ (10974962) United Kingdom Small Sample	Cognitive-behavioral family intervention, individual sessions every two weeks, 14 sessions (length of sessions unspecified)	In-depth interview of caregiver OR no interview	Community RCT Caregivers	Satisfaction of DSM-III-R criteria for primary degenerative dementia of the Alzheimer type	N=42 Age: yes Sex: yes Race: no Education: no	NR	N=42 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Duration: yes Living with PLWD: yes	NA	NA	Post intervention, three months post intervention	ADLs CSDD	GHQ (caseness/psychiatric morbidity) BDI
Steffen, 2000 ¹⁶⁷ (no PMID) United States Pilot	Weekly 30-minute psychoeducational videos--viewed either at home with 20-minute weekly telephone sessions or in a class setting with a trained facilitator--with accompanying workbook, 8 weeks, 90-minute classroom sessions	No Treatment	Community RCT Caregivers	Diagnosis of Alzheimer's disease, multi-infarct dementia or some other dementing illness	NR	NR	N=33 Age: yes Sex: yes Race: yes Education: yes Relationship: yes	NR	NA	NA	Post intervention	NR	CgAI BDI Revised Care-Giving Self-Efficacy Scale

Buckwalter 1999 ⁷² (10222636) (12464756) ¹⁶⁸ US High ROB	Psychoeducation for managing problem behaviors Information, case management referrals, service provision, support groups	NR	Community-based setting RCT N=245 dyads	Dementia diagnosis	N=245 Age: no Sex: yes Race: no Education: no	NR	N=245 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Employment Status: yes Training: yes	NA	NA	6, 12 months	NA	POMS GDRS
Chang 1999 ⁷³ (10337848) (15471059) ¹⁶⁹ USA High ROB	Cognitive-behavioral intervention for PLWD and care-giver	Attention only placebo telephone calls	Homebased RCT	PLWD diagnosed with dementia MMSE<21 Significant eating and dressing problems Home caregivers	N=34 Intervention group 31 Control group Age: yes Sex: no Education: yes Race % majority: yes	NR	Age: yes Sex: yes Race % majority: yes Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Training: yes	NR	NR	12 weeks	MMSE	Depression Burden Satisfaction Anxiety
Ostwald 1999 ⁷⁴ (10396888) US High ROB	Psychoeducation group intervention for caregiver burden and depression 7 weekly 2-hour sessions	Waitlist	Community-based setting RCT N=117 dyads	Dementia diagnosis	N=117 Age: yes Sex: yes Race: no Education: yes	PLWD SES: yes	N=117 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: no	Duration: no Employment Status: no Training: no	NA	NA	5 months	RMBPC	ZBI CES-D
McCurry, 1998 ¹⁷⁰ (9520929) United States Pilot	Weekly small group behavioral treatment sessions, 6 weeks (session length not specified) OR weekly individual behavioral treatment, 4 weeks (session length not specified)	No Treatment	Community RCT Caregivers	Diagnosis of senile dementia	N=36 Age: yes Sex: yes Race: no Education: no	NR	N=36 Age: yes Sex: yes Race: yes Education: yes Relationship: yes	Living with PLWD: yes	NA	NA	Post intervention, 3 months post intervention	NR	CES-D SCB RMBPC
Moniz-Cook, 1998 ¹⁷¹ (no PMID) Pilot	Advice pamphlets plus home-based individualized intervention delivered after dementia diagnosis, six to 12 hours in duration, four to fourteen weeks	Advice pamphlets plus usual care (i.e., referred to local psychogeriatric community team upon diagnosis)	Community RCT Caregiver-PLWD Dyads	Probable diagnosis of Alzheimer's disease, multi-infarct dementia, or frontal lobe dementia per neuropsychological profiles and ICD-10 operational criteria for diagnosing dementia	N=30 Age: yes Sex: yes Race: no Education: no	Household characteristics	N=30 Age: no Sex: no Race: no Education: no Relation to PLWD: no	NR	NA	NA	6 months after diagnosis (caregivers), 18 months after diagnosis (PLWD and caregivers)	Memory impairment RBMT	GHQ BDI HADS

Brodsky, 1997 ¹⁷² (9520929) Australia Pilot	Caregiver training program delivered daily for 10 days in residential setting immediately upon enrollment in study, followed by telephone check-ins at decreasing time intervals over the next 12 months OR same treatment and follow-up delivered 6 months after study enrollment	No Treatment (PLWD memory retraining only, delivered to all PLWD in study)	Community RCT Caregivers	Diagnosis of Dementia	N=96 Age: yes Sex: yes Race: no Education: no	NR	N=96 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Living with PLWD: yes	NA	NA	3 months post-intervention, 6 months post-intervention, 12 months post-intervention, and annually thereafter until conclusion of 8-year study	Nursing home admission, time until death	NR
Teri, 1997 ¹⁷³ (9224439) United States Small Sample	Behavior therapy focused on increasing pleasant events OR behavior therapy focused on problem solving situations of concern, both treatments administered weekly for 9 weeks, 60-minute sessions	Typical therapeutic care weekly for 9 weeks, 60-minute sessions OR no treatment (waitlist)	Community RCT Caregiver-PLWD dyads	Probable Alzheimer's disease per NINCDS-ADRDA criteria	N=88 Age: yes Sex: yes Race: no Education: yes	NR	N=88 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	NR	NA	NA	Post intervention, six months post intervention	HDRS CSDD BDI Record of Independent Living (RIL)	HDRS ZBI
Gendron, 1996 ¹⁷⁴ (26250550) Pilot	Weekly cognitive-behavioral skills training group intervention, 8 weeks, 90-minute sessions	Weekly support group emphasizing information and social exchange 8 weeks, 90-minute sessions	Community RCT Caregivers	Diagnosis of dementia	NR	NR	N=35 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Living with PLWD: yes Health Status: yes	NA	NA	Post intervention, 3-month follow-up, 6 month follow up	NR	Hopkins Symptom Checklist (distress) Automatic Thoughts Questionnaire (depressive thinking) ZBI RMBPC
Hebert, 1994 ¹⁷⁵ (24087060) (25107702) ¹⁷⁶ Canada Small Sample	Weekly structured support group sessions for 8 weeks, 2-hour sessions	Referral to informal monthly meetings of the Alzheimer's Society	Community RCT Caregivers	Dementia confirmed by DSM-III-R criteria	NR	NR	N=45 Age: yes Sex: yes Race: no Education: yes Relationship: yes	Living with PLWD: yes Employment status: yes	NA	NA	Post intervention, 6 months post intervention	SMAF	ZBI BSI RMBPC

Tappen, 1994 ¹⁷⁷ United States Small Sample	Group interventions of (a) skill training or (b) mental stimulation delivered 5 days a week, 20 weeks, 2.5 hours per session	No treatment	Nursing home RCT PLWD	Diagnosis of dementia with six or more errors out of 10 items on the Short Portable Mental Status Questionnaire	N=63 Age: yes Gender: yes Race: no Education: no	NR	NA	NA	NA	NA	Post intervention	Physical Self-Maintenance Scale Performance Test of ADLs	NA
--	--	--------------	---------------------------------	---	--	----	----	----	----	----	-------------------	---	----

Abbreviations: BDI= Beck Depression Inventory; CES-D=C Center for Epidemiologic Studies-Depression Scale; char=characteristics; CMAI= Cohen-Mansfield Agitation Inventory; FC=formal caregiver; GDRS= Geriatric Depression Rating Scale; IC=informal caregiver; MCS-12=Mental Health Composite, 12-item Short-Form Health Survey; MBPC= Memory Behavior Problem Checklist; MMSE= Mini Mental State Examination; N=number; NA=Not Applicable; NPI=Neuropsychiatric Inventory; NPI-D=Neuropsychiatric Inventory-Caregiver Distress; NR=Not Reported; PCS-12=Physical Health Composite, 12-item Short-Form Health Survey; POMS= The Profile of Moods States; PMID=PubMed Identification Number; PSS=Perceived Stress Scale; PLWD=Persons with Dementia; RCT=Risk of Bias; RoB=Risk of Bias; RMBPC=Revised Memory Behavior Problem Checklist; RS=Reporting Status; SES=socioeconomic status; STAI= Spielberger State-Trait Anxiety Inventory; STAXI= Spielberger State-Trait Anger Inventory; WHOQOL-BREF= World Health Organization Quality of Life Instruments; ZBI=Zarit Burden Interview

Social Support

Table E-9. Risk of bias assessment: social support

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Huis 2019 ¹⁷⁸ (32130142)	6 weeks 12 weeks	Medium 14%	Low	Medium	High	Low	Medium	Netherlands Organization for Health Research and Development program Tussen Wet en Doen	High
Van Mierlo 2015 ¹⁷⁹ (25872457)	6 months 12 months	Medium 6 months: 12% High 12 months: 33%	Medium	High	Medium	Medium	X	Government	High
Wang 2012 ¹⁸⁰ (22554214)	28 weeks	Low 7.6%	Medium	Low	Low	Low	Medium	NR	Medium
Logsdon 2010 ¹⁸¹ (20693265)	2 months	Low 4%	Medium	Low	Medium	High	X	Government	High
Charlesworth 2008 ¹⁸² (18284895) Charlesworth 2008 ¹⁸³ (18505757) Wilson 2009 ¹⁸⁴ (19101921)	6 months 15 months 24 months	Medium 6 months: 10% 15 months: 16% High 24 months: 21%	Low	Medium	High	Low	X	Government	High
Winter 2006 ¹⁸⁵ (17267370)	6 months	Low 8.7%	Medium	High	High	Medium	X	Non-profit	High
Mahoney 2003 ¹⁸⁶ (12937334) Mahoney 2001 ¹⁸⁷ (11346473)	6 months 12 months 18 months	Low 6 months: 8% Medium 12 months: 14% 18 months: 18%	Low	Low	Medium (6 months) High (12 and 18 months)	Medium	Medium	Government	Medium (6 months) High (12 and 18 months)
Pillemer 2002 ¹⁸⁸ (NA)	6 months	High 22%	X	X	X	X	X	Government	High
Goodman 1990 ¹⁸⁹ (2354800) Goodman 1990 ¹⁸⁹ (2284602)	NR	High 34%	X	X	X	X	X	NR	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table E-10. Characteristics of included studies: social support

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Wang 2012 ¹⁸⁰ (22554214) China Medium Explanatory	In person, peer-led mutual support group for caregivers; bi-weekly 90-minute sessions for 24 weeks (12 total sessions)	Usual care (conventional family services)	Dementia resource and respite centers RCT Multisite 78 Informal Caregivers	All dementia types and severities (primarily AD and cerebrovascular). Dementia diagnosis based on DSM-IV criteria	N=78 81% Age 61+ 53% Female Race NR Education NR	NR	N=78 65% Age 31-70 62% Female 73% Secondary School or Higher Race NR 37% Spouse	Duration: Yes Employment Status: Yes	NA	NA	28 weeks	NA	NPI-D WHOQOL-BREF(HK) Family Supports Services Index	
Mahoney 2003 ¹⁸⁶ (12937334) US Medium Explanatory Mahoney 2001 ¹⁸⁷ (11346473)	Automated telephone support system for caregivers which included a weekly automated conversation, a personal voice mailbox, telephone support group, and an activity-respite module; available 22 hours a day for 12 months	Usual care	Community-based recruitment from health and social agencies. RCT Single site 100 Informal Caregivers	Diagnosis of probable AD or MMSE ≤ 23, with two IADL impairments and one AD-related disturbing behavior	N=100 77 years 48% Female Race NR 70% High School Diploma or Higher	NR	N=100 63 years 78% Female 79% White 82% High School Diploma or Higher 54% Spouse	Employment Status: Yes	NA	NA	6 months	NA	CES-D STAI	

*High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: AD=Alzheimer's Disease; CES-D= Center for Epidemiologic Studies Depression Scale; char=characteristics; DSM-IV= Diagnostic and Statistical Manual of Mental Disorders 4th edition; FC=formal caregiver; IADL=Instrumental Activities of Daily Living; IC=informal caregiver; MMSE=Mini-mental State Exam; N=number; NA=Not Applicable; NPI-D=Neuropsychiatric Inventory Caregiver Distress; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RCT=Randomized Controlled Trial; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status;; STAI= State-Trait Anxiety Inventory; WHOQOL-BREF(HK)= World Health Organization Quality of Life Instruments (Hong Kong Version)

Table E-11. Caregiver outcomes summary low and medium risk of bias studies: social support

Study (PMID) Comparison Caregiver Type RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Wang 2012 ¹⁸⁰ (22554214) In-person support group vs. Usual care Informal Medium Explanatory	Mean (SD) NPI-D Total Score 28 weeks	NR	37.47 (9.68)	43.88 (13.56)	p=0.005
Wang 2012 ¹⁸⁰ (22554214) In-person support group vs. Usual care Informal Medium Explanatory	Mean (SD) WHOQOL-BREF(HK) Total Score 28 weeks	NR	114.02 (13.98)	88.19 (7.56)	p=0.001
Wang 2012 ¹⁸⁰ (22554214) In-person support group vs. Usual care Informal Medium Explanatory	Mean (SD) Mental Health Services Utilization, Family Supports Services Index 28 weeks	No difference between groups in utilization of mental health services.	NR	NR	NR
Mahoney 2003 ¹⁸⁶ (12937334) Automated support vs. Usual care Informal Medium Explanatory Mahoney 2001 ¹⁸⁷ (11346473)	Mean (SD) CES-D 6 months	No difference between groups.	12.3 (9.1)	14.9 (11.7)	NR
Mahoney 2003 ¹⁸⁶ (12937334) Automated support vs. Usual care Informal Medium Explanatory Mahoney 2001 ¹⁸⁷ (11346473)	Mean (SD) STAI 6 months	No difference between groups.	19.8 (5.7)	20.6 (7.7)	NR

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: CES-D= Center for Epidemiologic Studies Depression Scale; NA=Not Applicable; NPI-D=Neuropsychiatric Inventory Caregiver Distress; NR=Not Reported; PMID=PubMed Identification Number; SD=Standard Deviation; STAI= State-Trait Anxiety Inventory; WHOQOL-BREF(HK)= World Health Organization Quality of Life Instruments (Hong Kong Version)

Table E-12. Summary of strength of evidence for caregiver outcomes: social support

Comparison Caregiver Type Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
In-person support vs. Usual care Informal All Outcomes	28 weeks	1 RCT (n=78)	Insufficient evidence to draw conclusions about all outcomes.	Moderate	Unknown	Direct	Imprecise	Insufficient
Automated support vs. Usual care Informal All Outcomes	6 months	1 RCT (n=100)	Insufficient evidence to draw conclusions about all outcomes.	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: n=number; RCT=Randomized Controlled Trial

Evidence Map: Social Support for Caregivers

Table E-13. Characteristics of evidence map studies: social support for caregivers

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Huis 2020 ¹⁷⁸ (32130142) Netherlands High ROB	Email contacts with a specialist dementia nurse and informational videos and e-bulletins vs. informational videos and e-bulletins vs. e-bulletins		Community 3-arm RCT N=81 informal caregivers	AD Vascular dementia Frontotemporal dementia Dementia with Lewy bodies Mixed dementia Dementia type not known PLWD lives at home and has minimum weekly contact with informal caregiver who is partner or family member	N=81 Age: yes Sex: yes Race: no Education: no	Household Characteristics: yes	N=81 Age: yes Sex: yes Education: yes Relation to PLWD: yes	Living with PLWD: yes	NA	NA	6 weeks, 12 weeks	Disruptive behavior (RMBPC)	Self-efficacy (TRUST) Dyadic relationship quality (DRS)
Gustafson 2019 ¹⁹⁰ (31256126) US Pilot study	Dementia–Comprehensive Health Enhancement Support System (D-CHESS) website access for 6 months	Caregiving book	Community-based (intervention at home) RCT; single site 31 informal caregivers	AD, all severities	N=31 Age: yes Sex: no Race: no Education: no	NR	N=31 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Payment: yes	NA	NA	6 months	NA	Caregiver Burden (composite) Family Conflict (Aneshensel Scale) Satisfaction with Decision Scale MOS Social Support Survey UCLA Loneliness Scale Generalized Anxiety Disorder Scale PHQ Coping Competence (Lawton Scale)

Metcalfe 2019 ¹⁹¹ (31111516) Multinational Pilot study	Online information and support program for caregivers of individuals with young onset dementia; access for 12 weeks	Waitlist control	Community-based memory clinics (intervention at home) RCT; multisite 61 informal caregivers	AD or behavioral variant frontotemporal degeneration onset before the age of 65	N=61 Age: yes Sex: yes Race: no Education: no	NR	N=61 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment Status: yes	NA	NA	12 weeks	NA	RSCE PSS Burden Scale for Family Caregivers EQ-5D-5L
Van Mierlo 2015 ¹⁷⁹ (25872457) Netherlands High ROB	DEMENTIA-specific dynamic interactive social chart (DEM-DISC) providing information on support, coping, dementia, and other topics; accessible to both carers and case managers	No intervention	Community-based setting (intervention at home) Cluster RCT; multisite 27 clusters 73 informal caregivers	Dementia (not specified)	N=73 Age: yes Sex: yes Race: no Education: no	NR	N=73 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Living with PLWD: yes Payment: yes Health Status: yes	NA	NA	6 months 12 months	QOL-AD EQ-5D	Camberwell Assessment of Needs for the Elderly NPI-Distress Short Sense of Competence Questionnaire
Torkamani 2014 ¹⁹² (24643137) UK Pilot Study	Computerized social support platform provided information, social networking, and educational materials	No intervention	Hospital outpatients RCT, multisite 60 PLWD-informal caregiver dyads	Dementia with BI score 35 or greater and MMSE score from 9 to 21	N=60 Age: yes Sex: no Race: no Education: no	NR	N=60 Age: yes Sex: no Race: no Education: no Relation to PLWD: no	NR	NA	NA	3 months 6 months	NPI MMSE MDRS BDRS CDR RMBPC GDS BI Lawton ADL	ZBI BDI Zung Depression Self-rating Scale EQ-5D Quality of Life Scale
Logsdon 2010 ¹⁸¹ (20693265) US High ROB	Early-stage memory loss support group for PLWD-caregiver dyad; 90-minute sessions weekly for 9 weeks	Waitlist control	Community-based setting RCT; single site 142 PLWD-informal caregiver dyads	Dementia diagnosis with MMSE score 18 or higher	N=142 Age: yes Sex: yes Race: yes Education: yes	NR	N=142 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	NR	NA	NA	9 weeks	QOL-AD SF-36 GDS RMBPC	GDS Family Assessment Measure, Communication PSS
van der Roest 2010 ¹⁹³ (20455122) Netherlands Pilot Study	DEMENTIA-specific dynamic interactive social chart (DEM-DISC) providing information on support, coping, dementia, and other topics	No intervention	Community-based setting (intervention at home) CCT; single site 29 informal caregivers	Dementia (not specified)	N=29 Age: yes Sex: yes Race: no Education: no	NR	N=29 Age: yes Sex: no Race: no Education: yes Relation to PLWD: yes	Health status: yes	NA	NA	2 months	Camberwell Assessment of Need for the Elderly QOL-AD	Short Sense of Competence Questionnaire Pearlin Mastery Scale

Charlesworth 2008 ¹⁸² (18284895) High ROB UK Charlesworth 2008 ¹⁸³ (18505757) Wilson 2009 (19101921)	Befriending volunteers to provide emotional support to informal caregivers; weekly contact for 6 months	Usual care	Community-based setting RCT; single site 236 informal caregivers	Primary progressive dementia	N=236 Age: yes Sex: no Race: no Education: no	NR	N=236 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Employment: yes	NA	NA	6 months 15 months 24 months	NA	HADS-Depression EuroQoL Positive and negative affectivity scale HADS-Anxiety MSPSS
Winter 2006 ¹⁸⁵ (17267370) US High ROB	Telephone-based support groups; 1 hour weekly for 6 months	No intervention	Community-based setting (intervention at home) RCT; single site 106 female caregivers	AD and related dementias	N=106 Age: yes Sex: yes Race: no Education: no	NR	N=106 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes	NA	NA	6 months	NA	CES-D ZBI Gain Through Group Involvement Scale
Pillemer 2002 ¹⁸⁸ (NA) US High ROB	One-on-one in person peer support for new informal caregivers; weekly 2-hour sessions for 8 weeks	No intervention	Community-based setting (intervention at home) RCT; single site 147 informal caregivers	AD	N=147 Age: yes Sex: yes Race: no Education: no	NR	N=147 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Living with PLWD: yes	NA	NA	6 months	NA	CES-D Index of Disruptive Behaviors (Caregiving Stress)
Goodman 1990 ¹⁸⁹ (2354800) US High ROB Goodman 1990 ¹⁸⁹ (2284602)	Telephone support network; 2 calls a week for 12 weeks	Telephone-accessed taped lectures on AD and care management accessed anytime over 12 weeks	81 informal caregivers	AD and other dementias	N=81 No Sex: no Race: no Education: no	NR	N=81 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Health Status: yes Employment status: yes	NA	NA	12 weeks	MPBC	ZBI Perceived Social Support and Social Conflict Mental Health Index

Abbreviations: AD=Alzheimer's disease; ADL=Activities of Daily Living; BDI=Beck Depression Inventory; BDRS= Blessed Dementia Rating Scale; BI=Barthel Index; CDR=Clinical Dementia Rating; CES-D= Center for Epidemiologic Studies Depression Scale; char=characteristics; EQ-5D=EuroQoL-5D; EQ-5D-5L=5-level EuroQoL-5D; EM=Evidence Map; FC=formal caregiver; GDS=Geriatric Depression Scale; HADS=Hospital Anxiety and Depression Scale; IC=informal caregiver; MDRS=Mattis Dementia Rating Scale; MOS=Medical Outcomes Survey; MMSE=Mini-mental Scale Exam; MPBC= Memory and Behavior Problem Checklist; MSPSS=Multidimensional Scale of Perceived Social Support; N=number; NA=Not Applicable; NPI=Neuropsychiatric Inventory; NR=Not Reported; PHQ=Patient Health Questionnaire; PMID=PubMed Identification Number; PSS=Perceived Stress Scale; PLWD=Persons with Dementia; QoL-AD= Quality of Life in Alzheimer's Disease; RCT=Randomized Controlled Trial; RS=Reporting Status; RMBPC= Revised Memory and Behavior Problem Checklist; ROB=Risk of Bias; RSCE=Revised Scale for Care-giving Self-Efficacy; SES=socioeconomic status; SF-36; 36-Item Short Form Survey; ZBI=Zarit Burden Interview

Lifestyle Interventions

Table E-14. Risk of bias assessment: lifestyle interventions

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Salamizadeh 2017 ¹⁹⁴ (28670585)	3 weeks	Medium 10%	Medium	High	High	Low	X	NR	High
Farran 2016 ¹⁹⁵ (28752016) Faran 2016 ¹⁹⁶ (29147683) Cothran 2017 ¹⁹⁷ (28956706)	12 months	High 27%	X	X	X	X	X	Government	High
Mahdavi 2016 ¹⁹⁸ (28499566)	5 weeks	Medium 11%	Medium	Medium	High	Medium	X	NR	High
Lowery 2014 ¹⁹⁹ (24338799) D'Amico 2016 ²⁰⁰ (26489776)	12 weeks	Medium 11%	Medium	Low	High	High	X	Government	High
Whitebird 2013 ²⁰¹	2 months 6 months	Low 2 months: 8% Medium 6 months:10%	Medium	Low	Low	Medium	Medium	Government	Medium
Connell 2009 ²⁰² (21709757)	6 months 12 months	Medium 17%	Medium	High	High	Low	X	Government	High
King 2002 (11773209) Castro 2002 ²⁰³ (12021419)	12 months	Medium 15%	Low	Low	High	Low	X	Government	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table E-15. Characteristics of included studies: lifestyle interventions

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes	
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service				IC: FC:
Whitebird 2013 ²⁰¹ (23070934) US Medium Explanatory	Group mindfulness-based stress reduction for informal caregivers; one 2.5-hour session/week for 8 weeks and a 5-hour retreat with CDs and written materials	Group community caregiver education and support for informal caregivers; one 2.5-hour session/week for 8 weeks, a 5-hour retreat, and telephone support calls	Community-based setting RCT, single site 78 family caregivers	Memory loss consistent with dementia	N=78 Age NR Sex NR Race NR Education NR	NR	N=78 57 years 89% Female 99% White 56% College or Graduation Education 74% Adult child	Duration: yes Employment status: yes	NA	NA	2 months 6 months	NA	PSS CES-D STAI Montgomery Borgatta Caregiver Burden Scale Medical Outcomes Study Social Support MCS-12 PCS-12	

*High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: CES-D=Center for Epidemiologic Studies-Depression Scale; char=characteristics; FC=formal caregiver; IC=informal caregiver; MCS-12=Mental Health Composite, 12-item Short-Form Health Survey; N=number; NR=Not Reported; PCS-12=Physical Health Composite, 12-item Short-Form Health Survey; PMID=PubMed Identification Number; PSS=Perceived Stress Scale; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; ; MCS-12=Mental Health Composite, 12-item Short-Form Health Survey; NA=Not Applicable; NR=Not Reported; PCS-12=Physical Health Composite, 12-item Short-Form Health Survey; PMID=PubMed Identification Number; PSS=Perceived Stress Scale; PLWD=Persons with Dementia; RoB=Risk of Bias; RCT=Randomized Controlled Trial; RoB=Risk of Bias; STAI=State-Trait Anxiety Inventory

Table E-16. Caregiver outcomes summary low and medium risk of bias studies: lifestyle interventions

Study (PMID) Comparison Caregiver Type RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	PSS Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: -0.61 6 months: -0.47	2 months: 15.2 (5.8) 6 months: 14.0 (4.5)	2 months: 19.3 (7.6) 6 months: 16.7 (7.2)	2 months: 0.007 6 months: 0.07 Group x Time: 0.01
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	CES-D Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: -0.66 6 months: -0.39	2 months: 10.6 (8.4) 6 months: 10.5 (6.5)	2 months: 17.1 (11.2) 6 months: 13.7 (9.5)	2 months: 0.005 6 months: 0.16 Group x Time: 0.07
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	STAI Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: -0.59 6 months: -0.52	2 months: 34.2 (10.7) 6 months: 34.6 (10.4)	2 months: 41.7 (14.4) 6 months: 41.1 (14.2)	2 months: 0.01 6 months: 0.02 Group x Time: 0.98
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	Montgomery Borgatta Caregiver Burden Scale, Objective Burden Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: 0.11 6 months: -0.11	2 months: 23.8 (3.6) 6 months: 22.1 (5.1)	2 months: 23.3 (5.0) 6 months: 22.6 (5.2)	2 months: 0.58 6 months: 0.67 Group x Time: 0.63
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	Montgomery Borgatta Caregiver Burden Scale, Subjective Demand Burden Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: -0.25 6 months: -0.42	2 months: 12.0 (3.2) 6 months: 11.0 (3.5)	2 months: 12.8 (3.5) 6 months: 12.4 (3.0)	2 months: 0.24 6 months: 0.09 Group x Time: 0.80
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	Montgomery Borgatta Caregiver Burden Scale, Subjective Stress Burden Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: -0.23 6 months: -0.20	2 months: 15.0 (2.3) 6 months: 14.0 (3.5)	2 months: 15.6 (2.9) 6 months: 14.7 (3.6)	2 months: 0.32 6 months: 0.26 Group x Time: 0.68
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	Medical Outcomes Study Social Support Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: 0.11 6 months: 0.06	2 months: 71.1 (21.1) 6 months: 74.2 (21.0)	2 months: 68.8 (21.8) 6 months: 73.0 (21.4)	2 months: 0.66 6 months: 0.84 Group x Time: 0.51
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	MCS-12 Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: 0.66 6 months: 0.54	2 months: 47.4 (9.2) 6 months: 49.7 (7.9)	2 months: 40.8 (10.6) 6 months: 44.6 (10.9)	2 months: 0.007 6 months: 0.04 Group x Time: <0.001
Whitebird 2013 ²⁰¹ (23070934) Mindfulness-based stress reduction vs. Education and support Informal caregivers Medium Explanatory	PCS-12 Mean (SD) 2 months 6 months	Cohen's <i>d</i> 2 months: 0.19 6 months: 0.23	2 months: 49.9 (9.1) 6 months: 51.0 (9.1)	2 months: 48.1 (9.7) 6 months: 48.7 (11.0)	2 months: 0.36 6 months: 0.26 Group x Time: 0.35

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: CES-D=Center for Epidemiologic Studies - Depression Scale; NA=Not Applicable; MCS-12=Mental Health Composite, 12-item Short-Form Health Survey; NR=Not Reported; PCS-12=Physical Health Composite, 12-item Short-Form Health Survey; PMID=PubMed Identification Number; PSS=Perceived Stress Scale; RoB=Risk of Bias; SD=Standard Deviation; STAI=State-Trait Anxiety Inventory

Table E-17. Summary of strength of evidence for caregiver outcomes: lifestyle interventions

Comparison Caregiver Type Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Mindfulness-based stress reduction vs. Education and support Informal caregivers All Outcomes	2 months 6 months	1 RCT (n=78)	Insufficient evidence to draw conclusions about the effect of mindfulness-based stress reduction on informal caregivers.	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: n=number; RCT=randomized controlled trial

Evidence Map: Lifestyle Interventions

Table E-18. Characteristics of evidence map studies: lifestyle interventions

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Kor 2019 ²⁰⁴ (30922609) China Pilot Study	Group modified mindfulness-based cognitive therapy, 7 2-hour sessions over 10 weeks	Usual care with brief education on dementia care (7 group sessions)	Community-based setting RCT; single site 36 informal caregivers	Dementia diagnosis	N=36 Age: no Sex: no Race: no Education: no	NR	N=36 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Employment Status: yes Training: yes	NA	NA	10 weeks 3 months	NA	PSS ZBI CED-S HADS-A Brief Resilience Scale SF-12 Five Facets Mindfulness Questionnaire Short Form
Richards 2018 ²⁰⁵ (29653492) US Sample Size	Visual arts education group (producing different artistic work each week with increasing difficulty) for PLWD-informal caregiver dyads; 1 session/week for 8 weeks	Discussion and slide shows about art and painting for PLWD-informal caregiver dyads; 1 session/week for 8 weeks	Community-based setting RCT; single site 35 PLWD-informal caregiver pairs	Mild to moderate AD and related dementias, MMSE 12 to 26	N=35 Age: yes Sex: yes Race: no Education: no	NR	N=35 Age: yes Sex: yes Race: no Education: no Relation to PLWD: no	NR	NA	NA	8 weeks 6 months	Rosenberg Self-Esteem Scale QOL-AD ADL (FAQ)	ZBI Rosenberg Self-Esteem Scale QOL-AD
Salamizadeh 2017 ¹⁹⁴ (28670585) Iran High ROB	Spiritual care education; 5 educational sessions, 45 to 60 minutes	Usual care	Community-based setting Quasi-experimental 60 informal caregivers	AD diagnosis	N=60 Age: no Sex: no Race: no Education: no	NR	N=60 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes	NA	NA	3 weeks	NA	GSE-10
Waelde 2017 ²⁰⁶ (28263398) US Sample Size	Group stress mindfulness and mantra training for female informal caregivers; 10 sessions over 12 weeks	Telephone psychoeducation and telephone supports, biweekly 10-15-minute calls over 12 weeks	Community-based setting RCT; single site 31 informal caregivers	AD or other dementia	N=31 Age: no Sex: no Race: no Education: no	NR	N=31 Age: yes Sex: yes Race: yes Relation to PLWD: no	Living with PLWD: yes Employment: yes	NA	NA	12 weeks	MMSE	Satisfaction with Life Scale CES-D Self-Efficacy for Controlling Upsetting Thoughts, Revised Scale for Caregiving Self-Efficacy

Brown 2016 ²⁰⁷ (26211415) US Sample Size	Group mindfulness-based stress reduction for informal caregivers; 1.5 to 2 hours/week for 8 weeks	Group social support for informal caregivers; 1.5 to 2 hours/week for 8 weeks	Community-based setting RCT, single site 38 informal caregivers	Early stage AD or other dementia, FAST stage 5 or lower	N=38 Age: no Sex: no Race: no Education: no	NR	N=211 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes	NA	NA	3 months	NA	PSS Acceptance and Action Questionnaire II Profile of Mood States SF-36 ZBI Mortality Scale, Family Care Inventory
Farran 2016 ¹⁹⁵ (28752016) US High ROB Faran 2016 ¹⁹⁶ (29147683) Cothran 2017 ¹⁹⁷ (28956706)	Telephone-based intervention to support and encourage informal caregivers to participate in moderate vigorous physical activity, including goal-setting; regular phone calls over 12 months	Caregiver skill building to provide information, support, and problem-solving to caregivers; regular phone calls over 12 months	Community-based setting with intervention received at home RCT; single site 211 Informal Caregivers	AD or related dementia	N=211 Age: yes Sex: yes Race: no Education: no	NR	N=211 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Living with PLWD: yes Health status: yes Employment Status: yes	NA	NA	12 months	NA	2-minute Step Test 30s Chair Stand Test Caregiver Burden (PADL/IADL) PSS
Hirano 2016 ²⁰⁸ (26289496) Japan Pilot Study	Informal caregiver selected leisure activity, 3 times/week for 30 minutes for 24 weeks	Usual care	Community-based setting RCT; single site 54 informal caregivers	AD, DSM-IV criteria	N=54 Age: yes Sex: yes Race: no Education: no	NR	N=54 Age: yes Sex: yes Race: no Education: no Relation to PLWD: no	Living with PLWD: yes Health status: yes	NA	NA	12 months	MMSE NPI	ZBI
Mahdavi 2016 ¹⁹⁸ (28499566) Iran High ROB	Spiritual group therapy for informal caregivers based on Iranian culture/Islam; weekly 45 to 60-minute sessions for 5 weeks	1. Caregiver discussion group; weekly 45 to 60-minute sessions for 5 weeks 2. No intervention	Community-based setting RCT; single site 100 Informal Caregivers	Diagnosis of AD	N=100 Age: yes Sex: no Race: yes Education: no	NR	N=100 Age: Yes Sex: no Race: yes Education: no Relation to PLWD: yes	NR	NA	NA	5 weeks	NA	Caregiver Strain Index
Danucalov 2015 (26685923) Danucalov 2013 (368865412) Brazil Sample Size	Yoga and compassion meditation; in-person and at home; three 75 min sessions/week for 8 weeks	Waitlist control	Community-based setting RCT; single site 53 informal caregivers	Diagnosis of AD	N=53 Age: no Sex: no Race: yes Education: no	NR	N=53 Sex: yes Age: yes Race: no Education: no Relation to PLWD: no	NR	NA	NA	8 weeks	NA	WHOQOL-Bref Subjective Vitality Scales Mindfulness Attention Awareness Scale Self-Compassion Sale

Leach 2015 ²⁰⁹ (25952550) Australia Pilot Study Leach 2014 ²¹⁰ (24044373) Leach 2016 ²¹¹ (27227995)	Transcendental meditation; 14-hour program over 12 weeks with face to face 12-week follow-up	Waitlist control	Community-based setting RCT; single site 17 informal caregivers	Diagnosis of dementia	N=17 Age: no Sex: no Race: yes Education: no	NR	N=17 Sex: yes Age: yes Race: no Education: no Relation to PLWD: yes	Duration: yes Health Status: yes	NA	NA	12 weeks 24 weeks	NA	AQoL-8D WebNeuro Test Battery
Lowery 2014 ¹⁹⁹ (24338799) UK High ROB D'Amico 2016 ²⁰⁰ (26489776)	Individually tailored, progressively intensive walking regimen for PLWD and informal caregiver dyad, 20–30 min at least 5 times/week for 12 weeks	Usual care	Community-based setting with intervention at home RCT; single site 131 PLWD-informal caregiver dyads	Clinical diagnosis of dementia with ICD-10 Diagnostic Criteria for Research and NPI minimum score of 2 and frequency of 2 (hallucinations and delusions excluded)	N=131 Age: yes Sex: yes Race: yes Education: yes	NR	N=131 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	NR	NA	NA	12 weeks	NPI DemQOL-Proxy Change of Domiciliary Status Mortality	ZBI NPI Caregiver Distress GHQ
Lavretsky 2013 ²¹² (22407663) US Pilot Study	Kirtan Kriya yogic meditation, 12 min/day for 8 weeks	Relation, 12 min/day for 8 weeks	RCT, single site	Dementia based on DSM-IV-R without major depression (screened by HAM-D-24)	N=39 Age: no Sex: no Race: no Education: no	NR	N=39 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: yes Health Status: yes	NA	NA	8 weeks	NA	SF-36 HAM-D MMSE Cumulative Illness Rating Scale CVLT II TMT A TMT B
Hirano 2010 (20850878) Japan Sample Size	Regular exercise for informal caregivers with moderate intensity, 3 times a week for 12 weeks	No exercise	Community-based setting RCT; single site 36 informal caregivers	AD according to DSM-IV criteria	N=36 Age: no Sex: no Race: no Education: no	NR	N=36 Age: yes Sex: yes Race: no Education: no Relation to PLWD: no	Health status: yes Living with PLWD: yes	NA	NA	12 weeks	NPI MMSE	ZBI VAS
Carbpneau 2011 ²¹³ (20598758) Canada Sample Size	Adapted leisure education program for informal caregiver-PLWD dyads; 4 to 6 in person sessions with telephone follow-ups	Usual care	Community and nursing home RCT; single site 49 PLWD-informal caregiver dyads	Not specified	N=49 Age: no Sex: no Race: no Education: no	NR	N=49 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Health status: yes	NA	NA	8 weeks 12 weeks	NA	General Well-being Scale Elder Care Scale Self-Efficacy

Oken 2010 ²¹⁴ (20929380) US Pilot Study	Mindfulness-based cognitive therapy, one 90-minute session in-person/week with at home implementation for 7 weeks	1. Educational class based on Powerful Tools for caregivers with weekly assignments, one 90-minute session/week for 7 weeks 2. Respite only, 3 hours of respite/week for 7 weeks	Community-based RCT; single site 31 informal caregivers	Progressive dementia	N=31 Age: no Sex: no Race: no Education: no	NR	N=31 Age: yes Sex: yes Race: no Education: No Relation to PLWD: yes	NR	NA	NA	8 weeks	NPI	RMBPC Caregiver Appraisal Tool PSS CESD General Perceived Self-Efficacy Scale PSQI Epworth Sleep Questionnaire Global Impression of Change
Connell 2009 ²⁰² (21709757) US High ROB	Telephone-based exercise counseling and goal-setting for female spouse caregivers with exercise workbooks and videos; 14 calls over 6 months	No intervention	Community-based setting with intervention at home RCT; single site 157 female spouse caregivers	No specified	N=157 Age: no Sex: no Race: no Education: no	NR	N=157 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Health status: yes Employment status: yes	NA	NA	6 months 12 months	NA	MOS Short-Form General Health Survey RMBPC (Caregiver Burden) CES-D
Wilz 2008 ²¹⁵ (18381838) Germany Pilot study	Assisted vacations (participating in positive activities, improving utilization of support, psychoeducation) based on the needs of the caregiver and PLWD; 1 week in a hotel or rehabilitation clinic	Waitlist control	Hotel or rehabilitation centers Quasi experimental 29 caregiving wives	Medical diagnosis of dementia	N=29 Age: yes Sex: yes Race: no Education: no	NR	N=29 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	Living with PLWD: yes	NA	NA	3 months	NA	Giessen Subjective Complaints List BDI
King 2002 (11773209) Castro 2002 ²⁰³ (12021419) US High ROB	Home-based, telephone supervised moderate intensity exercise training for female caregivers, 30-40-minute sessions 4 times/week for 12 months, 15 telephone contacts over 12 months	Telephone-based nutrition education, 15 telephone contacts over 12 months	Community-based setting with intervention at home RCT; single site 100 female caregivers	AD or another form of dementia	N=100 Age: yes Sex: no Race: no Education: no	NR	N=100 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: yes Employment Status: yes	NA	NA	12 months	NA	Screen for Caregiver Burden TMAS BDI PSS Interpersonal Social Evaluation List RMBPC (Caregiver Burden) PSQI

Abbreviations: AD=Alzheimer's Disease; AQoL-8D= Validity and Reliability of the Assessment of Quality of Life; BDI=Beck Depression Inventory; CES-D=Center for Epidemiologic Studies Depression Scale; char=characteristics; CVLT II=California Verbal Learning Test, 2nd edition; DemQOL-Proxy=Dementia Quality of Life-Proxy; EM=Evidence Map; FAQ=Functional Activities Questionnaire; FC=formal caregiver; GHQ=General Health Questionnaire; GSE-10= General Self-Efficacy Scale; HAM-D=Hamilton Depression Rating Scale; HADS-Anxiety= Hospital Anxiety and Depression Scale-Anxiety; IADL=Instrumental Activities of Daily Living; IC=informal caregiver; MMSE=Mini-mental State Exam; MOS=Medical Outcomes Survey; N=number; NA=Not Applicable; NPI= Neuropsychiatric Inventory; NR=Not Reported; PADL=Personal Activities of Daily Living; PMID=PubMed Identification Number; PSS=Perceived Stress Scale; PSQI=Pittsburgh Sleep Quality Index; PLWD=Persons with Dementia; QoL-AD= Quality of Life in Alzheimer's Disease; RCT=Randomized Controlled Trial; RMBPC=Revised Memory and Behavior Problems

Checklist; ROB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; SF-12= 12-Item Short Form Health Survey; SF-36= 36-Item Short Form Survey; TMAS=Taylor Manifest Anxiety Scale; TMT=Trail Making Test; VAS=Visual Analogue Scale; WHOQOL-Bref=World Health Organization Quality of Life Instruments; ZBI=Zarit Burden Interview

Respite Care

Table E-19. Risk of bias assessment: respite care vs. usual care

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Vandepitte 2019 ²¹⁶ (31243801) Vandepitte 2016 ²¹⁷ (27912740)	6 months	High 24.7%	X	X	X	X	X	Foundation	High
Zarit 1998 ²¹⁸ (9750575) Kim 2012 ²¹⁹ (21322030)	3 months 12 months	High 3 months: 42.7% 12 months:65.9%	X	X	X	X	X	Government	High
Lawton 1989 ²²⁰ (NA)	12 months	High 46%	X	X	X	X	X	Foundation	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number; NA=Not Applicable

Evidence Map: Respite Care

Table E-20. Characteristics of evidence map studies: respite care vs. usual care

Study (PMID) Country EM Reason	Intervention	Comparison	Setting and Design Setting Design Cluster N Participants Randomized N	PLWD Dementia Characteristics	PLWD Non-Disease Char PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD Non-Disease Char Reporting Status (RS) PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	Informal Caregiver (IC) Characteristics IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	Informal Caregiver (IC) Char. RS IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	Formal Caregiver (FC) Characteristics FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	Formal Caregiver (FC) Char. RS FC Health Status FC Training FC Education FC Position FC Length of Service	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes IC: FC:
Vandepitte 2019 ²¹⁶ (31243801) Belgium High ROB Vandepitte 2016 ²¹⁷ (27912740)	In-home respite care for informal caregivers with support diary for additional caregiver support; 5 to 14 days of 24-hour care for PLWD from trained employee	Usual care	Community-setting with in home care Quasi experimental 198 Informal Caregiver-PLWD Dyads	Not specified	N=198 Age: yes Sex: yes Race: no Education: yes	NR	N=198 Sex: yes Race: no Education: yes Relation to PLWD: yes	Living with PLWD: yes Employment Status: yes	NA	NA	6 months	RMBPC	ZBI EQ-5D-5 L Desire-to-Institutionalize scale
Zarit 1998 ²¹⁸ (9750575) US High ROB Kim 2012 ²¹⁹ (21322030)	Adult day care; at least 2 times/week for 12 months	Usual care (no intervention)	Community-based adult day care Quasi experimental 566 informal caregivers	Not specified	N=566 Age: yes Sex: yes Race: no Education: no	NR	N=566 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	NR	NA	NA	3 months 12 months	PADL IADL	Involvement in Caregiving CES-D Positive and Negative Affect Schedule Role Captivity
Lawton 1989 ²²⁰ (NA) US High ROB	Respite program with options for in-home respite services, adult day care, and institutional respite care (limited to 21 days); any combination of services (caregiver discretion) over 12 months	Usual care with list of local services	In-home, adult care, and/or nursing home RCT; Single site 632 Informal caregivers	Not specified	N=632 Age: yes Sex: yes Race: no Education: yes	Household Characteristics: Yes	N=632 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Employment Status: yes	NA	NA	12 months	NA	Multilevel Assessment Instrument CES-D The Affect Balance Scale

Abbreviations: CES-D=Center for Epidemiologic Studies Depression Scale; char=characteristics; EM=Evidence Map EQ-5D-5L= 5-level EQ-5D version; FC=formal caregiver; IADL=Instrumental Activities of Daily Living; IC=informal caregiver; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PADL=Performance Activities of Daily Living; PLWD=Persons with Dementia; RCT=Randomized Controlled Trial; RMBPC=Revised Memory and Behavior Checklist; ROB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; ZBI=Zarit Burden Interview

Multicomponent Interventions

Table E-21. Risk of bias assessment: multicomponent interventions for informal caregivers

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Overall Rating
Luchsinger 2018 ^{221xxxx} (30084133)	6 months	Low 6 months: 5%	Medium	Medium	Medium	Low	Medium	Medium
Berwig 2017 ²²² (29233097)	6 months	Low 6 months: 12%	Low	Medium	Medium	Low	Low	Medium
Berwig 2017 ²²² (29233097)	9 months	High 9 months: 33%	Low	Medium	Medium	Low	Low	High
Brijoux 2016 ²²³ (27839534)	16 weeks	Medium 16 weeks: 17%	Low	High	Medium	Low	Low	High
Koivisto 2016 ²²⁴ (26177825)	36 months	High 36 months: 45%	Low	NA	NA	NA	NA	High
Baglio 2015 ²²⁵ (24788581)	32 weeks	Medium 32 weeks: 17%	Low	Medium	High	Low	Low	High
Prick 2015 ²²⁶ (26004290)	3 months	Low 3 months: 11%	Low	Low	Medium	Medium	Low	Low
Prick 2015 ²²⁶ (26004290)	6 months	Medium 6 months: 21%	Low	Low	Medium	Medium	Low	Medium
Gaugler 2013 ²²⁷ (23339050)	12 months	High Unclear	Low	NA	NA	NA	NA	High
Martindale-Adams 2013 ²²⁸ (24617278)	12 months	Medium months: 11%	Medium	High	Medium	Low	Low	High
Mittleman 2008 ²²⁹ (18978250)	24 months	Low 24 months: 2%	Low	Low	High	Medium	Medium	Medium
Signe 2008 ⁵⁵ (18269429)	6 months 12 months	NA	High	NA	NA	NA	NA	High
Belle 2006 ²³⁰ (17116917)	6 months	Low 6 months: 9%	Low	Medium	High	Low	Low	Medium
Ducharme 2005 ²³¹ (16024406)	6 months	High 6 months: 34%	Medium	NA	NA	NA	NA	High
Mittleman 2004 ²³² (14722336)	4 months	Low 4 months:9%	Low	Low	High	Low	Medium	Medium
Mittleman 2004 ²³² (14722336)	12 months	High 12 months: 35%	Low	NA	NA	NA	NA	High

Mittleman 2004 ²³² (14722336)	24 months	High 24 months: 53%	Low	NA	NA	NA	NA	High
Mittleman 2004 ²³² (14722336)	36 months	High 36 months: 67%	Low	NA	NA	NA	NA	High
Mittleman 2004 ²³² (14722336)	48 months	High 46 months: 74%	Low	NA	NA	NA	NA	High
Eisdorfer 2003 ²³³ (12937331)	18 months	High 18 months: 32%	Low	NA	NA	NA	NA	High
Teri 2003 ²³⁴ (14559955)	3 months	Low 3 months: 9%	Low	Medium	Medium	Low	Low	Medium
Teri 2003 ²³⁴ (14559955)	6 months	Low 6 months: 18%	Low	Medium	Medium	Low	Low	Medium
Teri 2003 ²³⁴ (14559955)	12 months	Medium 12 months: 25%	Low	Medium	Medium	Low	Low	Medium
Teri 2003 ²³⁴ (14559955)	18 months	High 18 months: 37%	Low	NA	NA	NA	NA	High
Teri 2003 ²³⁴ (14559955)	24 months	High 24 months: 42%	Low	NA	NA	NA	NA	High
Mittelman 1993 ²³⁵ (8314099)	12 months	Low 12 months: 2%	Medium	High	High	Low	Medium	High
Mohide 1990 ²³⁶ (2184186)	6 months	Medium 6 months:30%	Low	High	Medium	Low	Low	High

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; CMAI-SF=Cohen-Mansfield Agitation Inventory-Short-Form; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; RUDAS=Rowland Universal Dementia Assessment Scale

Table E-22. Characteristics of included studies: multicomponent interventions for informal caregivers

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:

Teri 2003 ²³⁴ (14559955) US Low Explanatory	Dyad, trained home health care facilitated 12-hour training sessions with PLWD, PLWD caregiver exercise and behavioral management techniques (RDAD) (2 sessions per week for 3 weeks, 1 session per week for 4 weeks, followed by biweekly sessions for 4 weeks.) 12 weeks	Routine medical care in community.	In-person, in-home visits at homes of community dwelling PLWD and PLWD caregivers. RCT 153 PLWD	National Institute of Neurological and Communicative Diseases and Stroke/Alzheimer Disease and Related Disorders Association criteria for Alzheimer disease. Dementia diagnosis for an average of 4.3 years.	N=153 78 years 41% female 89% white 13 years education	Detailed Race Information: yes	N=153 70 years 70% female 87% white 13 years education 80% spouse	Living Status: yes	NA	NA	12 weeks	Physical role functioning scores (SF-36) SIP Mobility Cornell Depression in Dementia Restricted Activity (number of days reported)	NA
Prick 2015 ²²⁶ (26004290) (28120631) ²³⁷ (27099480) ²³⁸ (25336121) ²³⁹ Netherlands Medium Explanatory	During three months, a personal coach visited PLWD/PLWD caregiver dyads for eight one-hour-long home visits. The first month the dyads were visited weekly, followed by biweekly home visits over the next eight weeks. 12 weeks	Monthly mailed bulletins and phone calls.	Community-dwelling PLWD living with their caregivers. RCT 111 PLWD	Diagnosis of dementia, age 55+ years, living at home with a caregiver, no use of antidepressants, no psychotic symptoms, MMSE score < 14, receiving more than two days respite care in a day care facility.	N=111 77 years 41% Female Race NR Education 4.01 (0-6 scale)	None	N=111 72 years 72% female Education 4.08 (0-6 scale) 90.1% spouse	Living with PLWD: yes	NA	NA	12 weeks	NA	Family caregiver mood Self-Perceived Pressure from Family Care (Dutch version) RMBPC (Dutch version)
Belle 2006 ²³⁰ (17116917) (20122038) ²⁴⁰ (28295134) ²⁴¹ (21357811) ²⁴² , ²⁴³ (24652899) ²⁴⁴ (20056684) ²⁴⁵ (27294873) ²⁴⁶ (23983230) ²⁴⁷ (16920998) ²⁴⁸ (14518801) ²⁴⁹ US Medium Explanatory	12 in-home and telephone Structured sessions involving preliminary talk, information booklet, structured telephone-based support groups, and structured evaluation of each support group session. 6 months	Mailed packet of educational materials and provided 2 brief (<15 minute) telephone "check-in" calls at 3 months and 5 months after randomization 6 months	PLWD caregiver living with or sharing cooking facilities with PLWD relative with diagnosed Alzheimer disease or related disorders providing care least 4 hours per day for at least the past 6 months. RCT 642 PLWD	Alzheimer disease or related disorders. MMSE > 0	N=642 77 years 60% female Education < 12 years	SES: yes Prior disability: yes Household characteristics: yes Detailed race: yes	N=642 60 years 83% female 35% white Education 57% > 12 years 43% spouse	Duration: yes Living with PLWD: yes Employment status: yes	NA	NA	6 months		Prevalence of clinical depression for caregivers (CES-D score ≥15)

Berwig 2017 ²²² (NCT01690117) Germany Medium Explanatory	12 individual two-weekly sessions (9 at home with the informal caregiver and 3 via telephone) and combined five modules. 6 months.	Mailed packet of educational materials and provided 2 brief (<15 minute) telephone "check-in" calls at 3 months and 5 months after randomization 6 months	Community/in-home dwelling PLWD and caregivers. RCT 92 PLWD	AD or related disorders. MMSE > 0	N=92 73 years 72% female Race NR Education NR	NR	Education 13 years 89% spouse	92 living with PLWD	NA	NA	6 months	NA	Benefit caregiver depression (PHQ4) Benefit quality of life (SF-12)
Mittelman 2004 ²³² (14722336) (15121650) ²⁵⁰ (17101889) ²⁵¹ (17804831) ²⁵² (16616406) ²⁵³ (19139247) ²⁵⁴ (16420138) ²⁵⁵ (18179495) ²⁵⁶ (21543959) ²⁵⁷ (15197285) ²⁵⁸ US Medium Explanatory	Counseling sessions (in-person, family, and telephone) for caregivers vs. usual care with home-based phone interviews; Pearlín stress process model	Usual care for 4 months	Spouse caregivers living at home with patient RCT 371 PLWD	AD or related disorders.	N=371 Age: NR 90% White Sex NR Education :NR	Detailed race information: yes	N=371 71 years 60% female 90% white	371 spouse caregivers living at home with patient.	NA	NA	4 months	NA	Benefit caregiver stress perceptions Benefit caregiver depression
Mittelman 2008 ²²⁹ (18978250) (21071943) ²⁵⁹ (19705599) ²⁶⁰ Multinational Medium Explanatory	Counseling sessions (in-person, family, and telephone) for caregivers vs. usual care; five sessions individual and family counselling w/in first 3 months, continuous ad hoc telephone counselling Donepezil for all patients Pearlín stress process model	No control.	Community-based spouse caregivers living at home with patient RCT 158 PLWD	AD or related disorders. mild to moderate dementia at enrollment	N=158 Age: NR Sex NR Race NR Education NR	None	N=158 Age: NR Sex NR Race NR Education NR	None	NA	NA	6 months	NA	Depressive symptoms Beck Depression Inventory (revised).
Luchsinger 2018 ²²¹ (14722336) US Medium Explanatory	Counseling sessions (in-person, family, and telephone) for caregivers vs. usual care 6 planned visits of similar duration in 6 months and ad hoc contacts. Pearlín stress process model	No control.	Community-based spouse caregivers living at home with patient RCT 221 PLWD	AD or related disorders.	N=221 58 years 82% female Race NR Education NR	None	N=221 58 years 32% spouses. Race NR Education 12 years	Employment Status: yes	NA	NA	4 months	NA	Depressive symptoms, GDS Caregiver burden ZBI

* High risk of bias studies included in evidence map. This table includes low and medium risk of bias studies only.

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; CES-D=Center for Epidemiologic Studies Depression Scale; CMAI-SF=Cohen-Mansfield Agitation Inventory-Short-Form; GDS=Geriatric Depression Scale; MMSE=Mini-Mental State Exam; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PHQ-4=Patient Health Questionnaire-4; PLWD=Persons with Dementia; RCT=Randomized Controlled Trial; RoB=Risk of Bias; RS=Reporting Status; RUDAS=Rowland Universal Dementia Assessment Scale; RMBPC= Revised Memory and Behavior Problem Checklist; SF-12= 12-Item Short Form Health Survey; SF-36= 36-Item Short Form Survey; SIP=Sickness Impact Profile; ZBI=Zarit Burden Interview

Table E-23. Caregiver outcomes summary low and medium risk of bias studies: multicomponent interventions for informal caregivers

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Belle 2006 ²³⁰ (17116917) 12 in-home and telephone Structured sessions involving preliminary talk, information booklet, structured telephone-based support groups, and structured evaluation of each support group session vs Mailed packet of educational materials and provided 2 brief (<15 minute) telephone “check-in” calls at 3 months and 5 months after randomization Medium Explanatory	Prevalence of clinical depression for caregivers (CES-D score ≥15) n/N (%) 6 months	Overall prevalence of clinical depression at 6 month follow-up was significantly greater among caregivers in the control group than among those in the intervention group	12.6%	22.7%	0.001
Belle 2006 ²²⁴ (17116917) 12 in-home and telephone Structured sessions involving preliminary talk, information booklet, structured telephone-based support groups, and structured evaluation of each support group session vs Mailed packet of educational materials and provided 2 brief (<15 minute) telephone “check-in” calls at 3 months and 5 months after randomization Medium Explanatory	Burden (modified Zarit Burden) Mean (CI) 6 months	No difference	3 (-15, 21)	4.2 (-16.9, 25.7)	NR
Berwig 2017 ²²² (NCT01690117) 12 individual two-weekly sessions (9 at home with the informal caregiver and 3 via telephone) and combined five modules vs. Mailed packet of educational materials and provided 2 brief (<15 minute) telephone “check-in” calls at 3 months and 5 months after randomization Medium Explanatory	ZBI Mean change from baseline (SD) 9 months	Cohen’s d=0.623. Informal caregivers’ burden increased in both groups, but much more strongly in the control group	2.67 (8.86)	8.10 (8.58)	0.017
Berwig 2017 ²²² (NCT01690117) 12 individual two-weekly sessions (9 at home with the informal caregiver and 3 via telephone) and combined five modules vs. Mailed packet of educational materials and provided 2 brief (<15 minute) telephone “check-in” calls at 3 months and 5 months after randomization Medium Explanatory	PHQ-15 Mean change from baseline (SD) 9 months	Cohen’s d=0.502	-0.601 (3.999)	1.098 (2.773)	0.057

Berwig 2017 ²²² (NCT01690117) 12 individual two-weekly sessions (9 at home with the informal caregiver and 3 via telephone) and combined five modules vs. Mailed packet of educational materials and provided 2 brief (<15 minute) telephone "check-in" calls at 3 months and 5 months after randomization Medium Explanatory	SF-12 Psychological Mean change from baseline (SD) 9 months	Cohen's d=0.902	3.868 (10.662)	-4.618 (8.157)	0.001
Luchsinger 2018 ²²¹ (14722336) NYUCI vs. REACH-OUT Medium Explanatory	GDS Mean (SD) 6 months	There was no difference between groups There was no significant change from baseline in either group	REACH-OUT 9.8 (7.2)	NYUCI 9.6 (7.1)	NR
Luchsinger 2018 ²²¹ (14722336) NYUCI vs. REACH-OUT Medium Explanatory	ZCBS Mean (SD) 6 months	There was no difference between groups There was a decrease in the ZCBS for REACH-OUT (5.2 points (2.2–8.1), p<.001) and the NYUCI (4.6 points, (1.7–7.5, p=.002).	REACH-OUT 35.5 (18.0)	NYUCI 36.2 (15.9)	NR
Teri 2003 ²³⁴ (14559955) Reducing Disability in Alzheimer Disease (RDAD) vs. Routine Medical Care (RMC) Low Pragmatic	SF-36 Mean (SD) 12 weeks	Significant difference between groups	72.1 (33.0)	50.7 (39.1)	<0.001
Teri 2003 ²³⁴ (14559955) Reducing Disability in Alzheimer Disease (RDAD) vs. Routine Medical Care (RMC) Low Explanatory	SIP Mobility Mean (SD) 12 weeks	No significant difference between groups	16.0 (17.1)	15.2 (17.1)	0.17
Teri 2003 ²³⁴ (14559955) Reducing Disability in Alzheimer Disease (RDAD) vs. Routine Medical Care (RMC) Low Explanatory	Cornell Depression in Dementia Scale Mean (SD) 12 weeks	Significant difference between groups	5.2 (3.6)	6.2 (3.8)	0.02
Teri 2003 ²³⁴ (14559955) Reducing Disability in Alzheimer Disease (RDAD) vs. Routine Medical Care (RMC) Low Explanatory	Restricted Activity (number of days reported) Mean (SD) 12 weeks	Significant difference between groups	0.1 (0.4)	0.6 (2.5)	<0.001
Prick 2015 ²²⁶ (26004290) During three months, a personal coach visited PLWD/PLWD caregiver dyads for eight one-hour-long home visits. The first month the dyads were visited weekly, followed by biweekly home visits over the next eight weeks vs. Monthly mailed bulletins and phone calls Medium Explanatory	Family caregiver mood (Dutch version CES-D) Mean (SD) 12 weeks	All analyses showed no benefits of the intervention over time on any of the outcomes	13.71 (8.18)	10.94 (8.42)	0.08
Prick 2015 ²²⁶ (26004290) During three months, a personal coach visited PLWD/PLWD caregiver dyads for eight one-hour-long home visits. The first month the dyads were visited weekly, followed by biweekly home visits over the next eight weeks vs. Monthly mailed bulletins and phone calls Medium Explanatory	Dutch Self-Perceived Pressure from Family Care (SPICC) 12 weeks	All analyses showed no benefits of the intervention over time on any of the outcomes	5.67 (2.36)	5.85 (2.13)	0.45

Prick 2015 ²²⁶ (26004290) During three months, a personal coach visited PLWD/PLWD caregiver dyads for eight one-hour-long home visits. The first month the dyads were visited weekly, followed by biweekly home visits over the next eight weeks vs. Monthly mailed bulletins and phone calls Medium Explanatory	Dutch version of Revised Memory and Behavior Problem Checklist (RMBPC) 12 weeks	All analyses showed no benefits of the intervention over time on any of the outcomes	13.06 (10.38)	12.13 (8.55)	0.82
Mittelman 2008 ²²⁹ (18978250) Five sessions of individual and family counselling within 3 months of enrollment and continuous availability of ad hoc telephone counselling and drug treatment vs. drug alone. Medium Explanatory	Beck Depression Inventory (BDI) (revised) Regression coefficients 6 months	The predicted BDI scores decreased for treatment caregivers and increased for control caregivers.	NR	NR	NR
Mittelman 2004 ²³² (14722336) Counseling sessions (in-person, family, and telephone) for caregivers vs. usual care Medium Explanatory	MBPC Regression coefficients 4 months	Intervention did not affect the frequency of patient behavioral problems, it did significantly reduce caregivers' reaction ratings	NR	NR	NR

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: ZBI: Zarit Burden Inventory; GDS: Geriatric Depression Scale; ZCBS: Zarit Caregiver Burden Scale; MBPC: Memory and Behavior Problems Checklist

Table E-24. Summary of strength of evidence for caregiver outcomes: multicomponent interventions for informal caregivers

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Health in PLWD and informal caregiver Combined exercise & support vs. monthly phone calls & mailed bulletins	12 weeks	1 RCT (n=153)	1 showed benefit	Medium	Inconsistent	Direct	Imprecise	Insufficient
Depression Counseling sessions (in-person, family, and phone) for caregivers vs. usual care	4-6 months	1 RCT (n=371)	1 showed benefit	Medium	Unknown	Direct	Imprecise	Insufficient
Caregiver stress perception Counseling sessions (in-person, family, and phone) for caregivers vs. usual care	4-6 months	1 RCT (n=158)	1 showed benefit	Medium	Unknown	Direct	Imprecise	Insufficient
Depression Education, group discussion, in-home and phone support sessions, and caregiver feedback vs usual care**	6 months	3 RCT (n=895)	2 showed benefit 1 showed no difference	Medium	Consistent	Direct	Imprecise	Low REACH II
Caregiver stress, burden Education, group discussion, in-home and phone support sessions, and caregiver feedback vs usual care**	6 months	3 RCT (n=895)	1 showed benefit 2 showed no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Evidence Map: Multicomponent Interventions for Informal Caregivers

Table E-25. Characteristics of evidence map studies: multicomponent interventions for informal caregivers

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Mazurek, 2019 ²⁶¹ (30666097) Poland Small sample	Meeting Centres Support Programme (MCSP) Integrated group support activities (cognitive stimulation, activity groups, music therapy, psychomotor therapy, family support groups, psychoeducation, counseling)	Usual care	Persons with mild-to-moderately severe dementia, living at home, and a caregiver prepared to participate	Mild-to-moderate dementia (Global Deterioration Scale / GDS)	N=42 Age: yes Sex: yes Race: no Education: yes	Household characteristics	N=42 Age: yes Sex: yes Race: yes Education: yes	NR	NA	NA	6 months	NR	caregiver unmet needs
Zhang, 2019 ²⁶² (31466807) China Pilot	6 bi-weekly self-management support sessions; education,	Education material	Community-based Quasi-experimental Caregiver	85% AD, 15% other subtypes	N=41 Age: yes Sex: yes Race: no Education: yes	None	N=41 Age: yes Sex: yes Race: no Education: yes Relationship: yes	Health status Employment	NA	NA	36 weeks	NPI IADL DAD	SF-36 Self-efficacy
Brijoux 2016 ²²³ (27839534) Germany High RoB	Qualified family companions provided support to the caregiver	Caregivers were supported by conventional care companions	Community-dwelling RCT 76 families	Clinical diagnosis of dementia (all severity)	N=73 Age: yes Sex: yes Race: no Education: no	None	N=73 Age: yes Sex: yes Race: no Education: no Relationship: yes	NR	NA	NA	16 weeks	NR	HRQOL Burden reduction Better networking to get support
Koivisto 2016 ²²⁴ (26177825) Finland	Psychosocial interventions, including education, counseling and social support	Annual follow-up	Community-based – 3 clinics RCT	Patients diagnosed with mild AD CDR of 0.5 Or 1.0 Free of interfering comorbidities Family caregiver present	N=54 Intervention group 76 Control group Age: yes Sex: yes Education: yes Race % majority: no	Comorbidities: Y	N=54 Intervention group 76 Control group Age: yes Sex: yes Education: yes Race % majority: no Relation to PLWD: yes	Employment:	NA	NA	2 years 36 months of follow-up	Admission to nursing home CDR-SOB CERAD-NB MMSE NPI BDI QoL	QoL GHQ Sense of coherence BDI

Baglio, 2015 ²²⁵ (24788581) Italy High RoB	Multidimensional Stimulation group Therapy (MST) to improve cognition, behavior, and motor functioning in PWA with informal caregivers.	No comparison	Memory Clinic Outpatients RCT PLWD	Right-handed persons diagnosed with AD with MRI evidence of AD pathophysiological processes and MMSE score 15-24 and Clinical Dementia Rating scale score of 1-2, age 65-85 years, school attendance range 5-17 years	NR	NR	NR	NR	NA	NA	10 weeks (pathophysiological outcomes only)	BPSD, language, physiological brain activation	NR
Marshall, 2015 ²⁶³ (no PMID) UK Pilot	Living Well with Dementia group intervention lasting for 10 weekly sessions delivered by nurses from a memory clinic	Waitlist-control	NHS hospital or community settings or PLWD home RCT	AD, vascular, or Lewy body diagnosed within previous 18 months MMSE 18+	N=58 Age: yes Sex: yes Race: yes Education: yes	Household characteristics	N=58 Age: yes Sex: yes Race: yes Education: yes Relationship: no	NR	NA	NA	10 weeks	QOL-AD CSDD	None
Czaja 2013 ²⁶⁴ (23831174) US Pilot	Technology-based multi-component psychosocial intervention was delivered in-home and via videophone technology over 5 months.	Attention control caregiver group Caregiver received information only	Community-dwelling 3-arm RCT 110 CG	Physician diagnosis of AD or other type of dementia with MMSE score < 24 and at least one limitation in ADL or two in IADL	N=93 Age: yes Sex: no Race: no Education: yes	None	N=93 Age: yes Sex: yes Race: yes Education: no Relationship: yes	Duration: yes	NA	NA	5 months	NR	CESD RMBPC Positive aspects of caregiving. Social Support
Gaugler, 2013 ²²⁷ (23339050) (26238226) ²⁶⁵ (25628299) ²⁶⁶ (29562359) ²⁶⁷ (30009268) ²⁶⁸ US High RoB	Individual and family counseling, support group referral, and ad hoc consultation for adult children of PLWD	Contact-control	Community/independent living to residential setting Adult children of PLWD N=107	NR	NR	NR	N=107 Age: yes Sex: yes Race: yes Education: yes Relationship: no	Employment	NA	NA	2 years	Residential care placement	caregiver subjective health social support
Martindale-Adams 2013 ²²⁸ (24617278) US	Telephone support and periodic group meetings for caregivers	Printed materials for support	Community-based RCT	Veteran patients diagnosed with dementia or dementia patients with veteran caregivers MMSE of 23 or less At least one ADL or 2 IADL limitations Exclusion criteria	N=77 intervention group 77 control group Age: yes Sex: yes Education: yes Race % majority: no	Not reported	Age: yes Sex: yes Education: yes Race % majority: yes Relation to PLWD: yes	Duration Income Employment	NA	NA	1 year	No reported	Bother Burden Depression General well-being
Signe, 2008 ⁵⁵ (18269429) Sweden High ROB	Psychosocial intervention & conversation group, 5 weeks	Control group (comparison NR)	In-home Quasi-experimental Caregiver	Dementia unspecified	N=150 Age: yes Sex: yes Race: no Education: no	Household characteristics	N=155 Age: yes Sex: yes Race: no Education: no Relationship: yes	Employment	NA	NA	6 month and 12 month measures	NR	Caregiver Burden Scale CASI Nottingham Health Profile

Finkel, 2007 ²⁶⁹ (17463195) US Pilot	Technology-based psychoeducational intervention for family caregivers of dementia patients	Information-only	NR RCT Caregivers	Diagnosis AD or other type of dementia and MMSE score 23 (excluding scores of zero).	N=46 Age: yes Sex: yes Race: yes Education: yes	None	N=46 Age: yes Sex: yes Race: yes Education: yes Relationship: yes	None	NA	NA	6 months	NR	CED-S RMBPC
Ducharme 2005 ²³¹ (16024406) Canada High RoB	90 minutes of weekly sessions (10 weeks) for women CG comprising of a psychoeducational program focusing on cognitive appraisal of stress, perception of control over a situation and problem solving, reframing, and stress symptoms management.	Comparison program offered by an Alzheimer Society Control: No program	Long term care and residential centers 3-arm RCT 137 caregivers	Diagnosed with irreversible dementia	N=unclear Age: yes Sex: yes Race: no Education: no	None	N=unclear Age: yes Sex: yes Race: no Education: no Relationship: yes	NR	NA	NA	10 weeks 3 months	NR	PDI Caregiver Overload SAM CAMI Competence in dealing with healthcare

Eisdorfer 2003 ²³³ (12937331) US High RoB	SET: 60-90 minutes structural family therapy sessions for 12 months, with weekly sessions for the first 4 months, biweekly sessions for the subsequent 2 months, and monthly sessions for the final 6 months of treatment. SET+CTIS: CTIS is an information network that utilizes computer–telephone technology facilitating linkages of the caregivers with both their family and with supportive resources outside of the home.	Telephone-based, minimal support condition; biweekly calls for the first 6 months and monthly calls during the next 7–12 months. The duration of the calls ranged from 5 min to 15 min and consisted of active listening and empathic comments when appropriate.	Community-dwelling 3-arm RCT 225 CG	Medical diagnosis of AD or other type of dementia with MMSE score<24 and at least one limitation in ADL or two in IADL	N=148 Age: yes Sex: yes Race: yes Education: no	None	N=148 Age: yes Sex: yes Race: yes Education: yes Relationship: yes	Duration: yes	NA	NA	6 months 18 months	NR	CESD RMBPC Satisfaction with social support MMSE ADL IADL
Mittelman 1993 ²³⁵ (8314099) US	Individual and family counseling, support group and ad hoc consultation	Routine support	Community-based RCT	Patients with a clinical diagnosis of AD Spouse is primary caregiver Additional family support locally Exclusion criteria	N=103 Intervention group 103 control group Age: yes Sex: yes (inferred) Education: no Race % majority: no	Patient income	N=103 Intervention group 103 control group Age: yes Sex: yes Education: yes Race % majority: yes Relation to PLWD: yes	Duration Employment status Religion	NA	NA	12 months	Admission to nursing home Predictors of nursing home placement regression analysis	Not reported
Mohide, 1990 ²³⁶ (2184186) Canada High ROB	Caregiver Support Program (CSP): caregiver-focused health care, education about dementia and caregiving, assistance with problem solving, regularly scheduled in-home respite, self-help family caregiver support group	Conventional community nursing care	In-home RCT Caregiver	Moderate to severe dementia	N=30 Age: yes Sex: yes Race: no Education: no	SES	N=30 Age: yes Sex: yes Race: no Education: no	Duration	NA	NA	6 months	NR	CES-D Self-Anchoring Striving Scale Caregiver Quality of Life Inventory

Abbreviations: AD= Alzheimer’s disease; ADL=Activities of Daily Living; AES=Apathy Evaluation Scale; APADEM-NH=Apathy Scale for Institutionalized Patients with Dementia Nursing Home; AWS== Revised Algase Wandering Scale–Nursing Home version; BARS=Brief Agitation Rating Scale; BPSD=Behavioral and Psychological Symptoms of Dementia; CDR=Clinical Dementia Rating; CES-D= Center for Epidemiological Studies of Depression scale; CMAI-SF=Cohen-Mansfield Agitation Inventory-Short Form; CSDD=Cornell Scale for Depression in Dementia; DSM=Diagnostic and Statistical Manual; EM=Evidence Map; FC=Formal Caregiver; GDS=Global Deterioration Scale; GDS-SF= Geriatric Depression Scale Short Form; IC= Informal Caregiver; MMSE=Mini-mental State Examination; N=number; N= number; NA=Not Applicable; NIA=National Institute on Aging; NPI=Neuropsychiatric Inventory; NR=Not Reported; OERS=Observed Emotion Rating Scale; PMID=PubMed Identification Number; PLWD=People with dementia; QoL-AD=Quality of Life in Alzheimer’s Disease; QUALID=Quality of Life in Late Dementia; RCT= Randomized controlled trial; RMBPC= Revised Memory and Behavior Problems Checklist; ROB= Risk of bias; RS= Reporting status; SES= Socioeconomic status

Other Interventions for Informal Caregiver Well-Being

Table E-26. Risk of bias assessment: other interventions for informal caregiver well-being

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Jutten, 2018 ²⁷⁰ (30631337) Jutten 2017 ²⁷¹ (28827242)	2.5 months 15 months	High 2.5 months: 16% 12 months: 57%	High	X	X	X	X	Government and Nonprofit	High
Stern 2008 ²⁷² (19064472)	8 weeks	Medium NR	Medium	High	Medium	High	X	Government	High
Simpson 2006 ²⁷³ (16734920)	6 months 12 months	High 12 months: 39%	X	X	X	X	X	Government	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Evidence Map: Other Interventions for Informal Caregiver Well-Being

Table E-27. Characteristics of evidence map studies: other interventions for informal caregiver well-being

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Random-ized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Jutten, 2018 ²⁷⁰ (30631337) Netherlands High ROB Jutten 2017 ²⁷¹ (28827242)	Mixed virtual reality simulator (simulate experience of dementia)	No virtual reality	Community-based Quasi-experimental caregivers	Dementia unspecified	N=201 Age: yes Sex: yes Race: no Education: no	Duration	N=201 Age: yes Sex: yes Race: yes Education: no Relation to PLWD: yes	Living with PLWD: yes	NA	NA	2.5 and 15 months	NA	Empathy Hospital Anxiety and Depression Scale Caregiver Reaction Assessment Relationship Quality Index
Lord 2017 ²⁷⁴ (28243460) UK Pilot Study	DECIDE decision aid for informal caregivers about place of care for PLWD; one 45-minute session with a decision coach	Usual care	Community-based memory clinics RCT, single site 41 informal caregivers	Moderate or severe dementia MMSE < 20	N=41 Age: no Sex: no Race: no Education: no	NR	N=41 Age: yes Sex: yes Race: yes Education: no Relation to PLWD: yes	Living with PLWD: yes	NA	NA	10 weeks	NA	Decisional Conflict Scale Hospital Anxiety and Depression Scale
Stirling 2012 ²⁷⁵ (22429384) Australia Pilot Study	Decision aid about community services, including respite care, for informal caregivers; mailed directly to caregivers and completed over a week	Waitlist control	Community-based setting with in-home decision aid RCT; single site 31 informal caregivers	Not specified	N=31 Age: no Sex: no Race: no Education: no	NR	N=31 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration: Yes	NA	NA	2 weeks 3 months	NA	Modified Caregiver Strain Index Decisional Conflict Scale Control Preference Scale

Rose 2009 ²⁷⁶ (19427574) US Pilot Study	At home AlphaStim® cranial electrical stimulation Alzheimer's disease or multi-infarct dementia for spouse caregivers; 60 minutes per day for 4 weeks	Sham	Rural community setting, intervention at home RCT; single site 39 spouse caregivers	AD or multi-infarct dementia	N=39 Age: yes Sex: no Race: no Education: no	NR	N=39 Age: yes Sex: yes Race: no Education: yes Relation to PLWD: yes	Duration: Yes	NA	NA	2 weeks 4 weeks	NA	PSQI General Sleep Disturbance Scale Geriatric Depression Scale Philadelphia Geriatric Center Caregiving Appraisal Scales
Stern 2008 ²⁷² (19064472) US High ROB	Family caregiver psychoeducatio nal support meetings for PLWD driving cessation; 4 weekly 2-hour meetings	1. Written educational information and resource list for PLWD driving cessation. 2. No intervention (waitlist control)	Community- based setting Cluster RCT 14 sites 66 informal caregivers	Caregiver's report of MCI, AD, or other dementia	N=66 Age: no Sex: no Race: no Education: no	NR	N=66 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	NR	NA	NA	8 weeks	NA	Brief COPE Scale Self-Efficacy Questionnaire
Simpson 2006 ²⁷³ (16734920) UK High ROB	Carer held record folder with sections for carer/family, professional feedback, and a carer diary; held at home and maintained for 12 months	Usual Care	Community- based mental health care and at home CCT 84 informal caregiver- PLWD dyads	Not specified	N=84 Age: no Sex: no Race: no Education: no	NR	N=84 Age: no Sex: no Race: no Education: no Relation to PLWD: no	NR	NA	NA	6 months 12 months	Crichton Behavioral Scale Global Deterioration Scale	Dementia Knowledge Questionnaire Caregiver Strain Index GHQ SF-36 Locus of Control Scale

Abbreviations: AD=Alzheimer's disease; CCT=Controlled Clinical Trial; char=characteristics; COPE=Coping Orientation to Problems Experienced; EM=Evidence Map; FC=formal caregiver; GHQ=General Health Questionnaire; IC=informal caregiver; MCI=Mild Cognitive Impairment; MMSE=Mini Mental State Exam; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PSQI= Pittsburgh Sleep Quality Index; PLWD=Persons with Dementia; RCT=Randomized Controlled Trial; ROB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; SF-36=36-item Short Form Survey

Care Interventions for Formal Caregiver Staff Well-Being

Evidence Map: Formal Caregiver Staff

Table E-28. Characteristics of evidence map studies: care interventions for formal caregiver staff well-being

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Barbosa, 2015 ²⁷⁷ (25237132) (263994850) ²⁷⁸ Portugal Pilot	8 weekly sessions of 60 minutes psychoeducation plus 30 minutes relaxation and stress management strategies	Education only	Aged-care residential facilities Cluster RCT 4 facilities	Dementia unspecified (private, non-profit institutions with staff to resident ratio 1:2 or 1:3)	NR	NR	NA	NA	N=58 Age: yes Sex: yes Race: no Education: yes	Length of service	10 weeks	NR	Perceived stress scale MBI Intrinsic satisfaction with job
Visser, 2008 ²⁷⁹ (18297478) (17191270) ²⁸⁰ Australia Pilot	8-week staff education plus peer support group	Waitlist Education only	Aged-care facility Cluster RCT 3 facilities	Residents selected by staff for displaying agitation; dementia unspecified	N=76 Age: yes Sex: yes Race: no Education: no	NR	NA	NA	N=52 Age: yes Sex: yes Race: no Education: no	NR	6 months	ADRQL CMAI	The Staff Attitudes Questionnaire MBI, third edition
Mackenzie, 2003 ²⁸¹ (14569646) Canada Pilot	4 module training and education to decrease formal caregiver stress and burn-out	Usual care activities	Dementia care units Quasi-experimental 2 care units	NR	NR	NR	NA	NA	N=41 Age: yes Sex: yes Race: no Education: yes	Position	3-month post-intervention	NR	Inventory of Geriatric Nursing Self-Efficacy MBI Satisfaction with teamwork

Abbreviations: ADRQL= Alzheimer's Disease Related Quality of Life; char=characteristics; CMAI= Cohen-Mansfield Agitation Inventory; FC=formal caregiver; IC=informal caregiver; MBI= Maslach Burnout Inventory; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RCT=randomized controlled trial; RS=Reporting Status; SES=socioeconomic status

References for Appendix E

1. Au A, Lai DWL, Biggs S, et al. Perspective-taking interventions for intergenerational caregivers of alzheimer's diseases: a randomized controlled trial. *Research on social work practice*. 2020;30(3):306-19. PMID: CN-02084449.
2. Au A, Yip HM, Lai S, et al. Telephone-based behavioral activation intervention for dementia family caregivers: Outcomes and mediation effect of a randomized controlled trial. *Patient Educ Couns*. 2019 Nov;102(11):2049-59. doi: <https://dx.doi.org/10.1016/j.pec.2019.06.009>. PMID: 31279613.
3. Kunik ME, Stanley MA, Shrestha S, et al. Aggression Prevention Training for Individuals With Dementia and Their Caregivers: A Randomized Controlled Trial. *Am J Geriatr Psychiatry*. 2020 Feb 03;03:03. doi: <https://dx.doi.org/10.1016/j.jagp.2020.01.190>. PMID: 32115311.
4. Bjorge H, Kvaal K, Ulstein I. The effect of psychosocial support on caregivers' perceived criticism and emotional over-involvement of persons with dementia: an assessor-blinded randomized controlled trial. *BMC Health Serv Res*. 2019 Oct 24;19(1):744. doi: <https://dx.doi.org/10.1186/s12913-019-4551-x>. PMID: 31651321.
5. Ghaffari F, Rostami M, Fotokian Z, et al. Effectiveness of resilience education in the mental health of family caregivers of elderly patients with Alzheimer's disease. *Iranian Journal of Psychiatry and Behavioral Sciences*. 2019;13 (3) (no pagination)(e69507).
6. Meng H, Marino VR, Conner KO, et al. Effects of in-person and telephone-based cognitive behavioral therapies on health services use and expenditures among African-American dementia caregivers with depressive symptoms. *Ethn Health*. 2019 Mar 19;1-14. doi: <https://dx.doi.org/10.1080/13557858.2019.1590536>. PMID: 30884961.
7. Moskowitz JT, Cheung EO, Snowberg KE, et al. Randomized controlled trial of a facilitated online positive emotion regulation intervention for dementia caregivers. *Health Psychol*. 2019 May;38(5):391-402. doi: <https://dx.doi.org/10.1037/hea0000680>. PMID: 31045422.
8. Nordheim J, Hausler A, Yasar S, et al. Psychosocial Intervention in Couples Coping with Dementia Led by a Psychotherapist and a Social Worker: The DYADEM Trial. *J Alzheimers Dis*. 2019;68(2):745-55. doi: <https://dx.doi.org/10.3233/JAD-180812>. PMID: 30775982.
9. Terracciano A, Artese A, Yeh J, et al. Effectiveness of Powerful Tools for Caregivers on Caregiver Burden and on Care Recipient Behavioral and Psychological Symptoms of Dementia: A Randomized Controlled Trial. *J Am Med Dir Assoc*. 2019 Dec 19;19:19. doi: <https://dx.doi.org/10.1016/j.jamda.2019.11.011>. PMID: 31866419.
10. Uyar F, Ozmen D, Mavioglu H, et al. Assessment of the impact of dementia care and support program in both patient and caregiver outcomes: An intervention study. *Turk Geriatri Dergisi*. 2019;22(4):482-93.
11. Wawrziczny E, Larochette C, Papo D, et al. A Customized Intervention for Dementia Caregivers: A Quasi-Experimental Design. *J Aging Health*. 2019 Aug;31(7):1172-95. doi: <https://dx.doi.org/10.1177/0898264318770056>. PMID: 29665714.
12. Boots LM, de Vugt ME, Kempen GI, et al. Effectiveness of a Blended Care Self-Management Program for Caregivers of People With Early-Stage Dementia (Partner in Balance): Randomized Controlled Trial. *J Med Internet Res*. 2018 Jul 13;20(7):e10017. doi: <https://dx.doi.org/10.2196/10017>. PMID: 30006327.
13. Gitlin LN, Arthur P, Piersol C, et al. Targeting Behavioral Symptoms and Functional Decline in Dementia: A Randomized Clinical Trial. *J Am Geriatr Soc*. 2018 Feb;66(2):339-45. doi: <https://dx.doi.org/10.1111/jgs.15194>. PMID: 29192967.
14. Spalding-Wilson KN, Guzman-Velez E, Angelica J, et al. A novel two-day intervention reduces stress in caregivers of persons with dementia. *Alzheimer's and Dementia: Translational Research and Clinical Interventions*. 2018 01 Jan;4:450-60. doi: <http://dx.doi.org/10.1016/j.trci.2018.08.004>. PMID: 200115918.
15. Wilz G, Reder M, Meichsner F, et al. The Tele.TAnDem Intervention: Telephone-based CBT for Family Caregivers of People With Dementia. *Gerontologist*. 2018 03 19;58(2):e118-e29. doi: <https://dx.doi.org/10.1093/geront/gnx183>. PMID: 29190357.
16. Callahan CM, Boustani MA, Schmid AA, et al. Targeting Functional Decline in Alzheimer Disease: A Randomized Trial. *Ann Intern Med*. 2017 Feb 07;166(3):164-71. doi: <https://dx.doi.org/10.7326/M16-0830>. PMID: 27893087.
17. Whitlatch CJ, Heid AR, Femia EE, et al. The Support, Health, Activities, Resources, and Education program for early stage dementia: Results from a randomized controlled trial. *Dementia*. 2017 Jan 01:1471301217743033. doi: <https://dx.doi.org/10.1177/1471301217743033>. PMID: 29171296.
18. Charlesworth G, Burnell K, Crellin N, et al. Peer support and reminiscence therapy for people with dementia and their family carers: a factorial pragmatic randomised trial. *J Neurol Neurosurg Psychiatry*. 2016 Nov;87(11):1218-28. doi: <https://dx.doi.org/10.1136/jnnp-2016-313736>. PMID: 27521377.
19. Cheng ST, Fung HH, Chan WC, et al. Short-Term Effects of a Gain-Focused Reappraisal Intervention for Dementia Caregivers: A Double-Blind Cluster-Randomized Controlled Trial. *Am J Geriatr Psychiatry*. 2016 Sep;24(9):740-50. doi: <https://dx.doi.org/10.1016/j.jagp.2016.04.012>. PMID: 27401052.
20. Gonyea JG, Lopez LM, Velasquez EH. The Effectiveness of a Culturally Sensitive Cognitive Behavioral Group Intervention for Latino Alzheimer's Caregivers. *Gerontologist*. 2016 Apr;56(2):292-302. doi: <https://dx.doi.org/10.1093/geront/gnu045>. PMID: 24855313.
21. Laakkonen ML, Kautiainen H, Holttä E, et al. Effects of Self-Management Groups for People with Dementia and Their Spouses--Randomized Controlled Trial. *J Am Geriatr Soc*. 2016 Apr;64(4):752-60. doi: <https://dx.doi.org/10.1111/jgs.14055>. PMID: 27060101.
22. Taati F, Bahramnezhad F, Seyedfatemi N, et al. The effect of participation in support groups on depression, anxiety and stress in family caregivers of people with Alzheimer's: randomized clinical trial. *International journal of medical research & health sciences*. 2016;5(7):256-62. PMID: CN-01308282.
23. Blom MM, Zarit SH, Groot Zwaafink RB, et al. Effectiveness of an Internet intervention for family caregivers of people with dementia: results of a randomized controlled trial. *PLoS ONE*. 2015;10(2):e0116622. doi: <https://dx.doi.org/10.1371/journal.pone.0116622>. PMID: 25679228.
24. Chiu M, Pauley T, Wesson V, et al. Evaluation of a problem-solving (PS) techniques-based intervention for informal carers of patients with dementia receiving in-home care. *Int Psychogeriatr*. 2015;27(6):937-48. doi: 10.1017/S1041610214002798. PMID: CN-01102329.
25. Gallagher-Thompson D, Tzuang M, Hinton L, et al. Effectiveness of a fotonovela for reducing depression and stress in Latino dementia family caregivers. *Alzheimer Dis Assoc Disord*. 2015 Apr-Jun;29(2):146-53. doi: <https://dx.doi.org/10.1097/WAD.0000000000000077>. PMID: 25590939.
26. Losada A, Marquez-Gonzalez M, Romero-Moreno R, et al. Cognitive-behavioral therapy (CBT) versus acceptance and commitment therapy (ACT) for dementia family caregivers with significant depressive symptoms: Results of a randomized clinical trial. *J Consult Clin Psychol*. 2015 Aug;83(4):760-72. doi: <https://dx.doi.org/10.1037/ccp0000028>. PMID: 26075381.
27. Otero P, Smit F, Cuijpers P, et al. Long-term efficacy of indicated prevention of depression in non-professional caregivers: randomized controlled trial. *Psychol Med*. 2015;45(7):1401-12. doi: 10.1017/S0033291714002505. PMID: 25331992.
28. Tremont G, Davis JD, Papandonatos GD, et al. Psychosocial telephone intervention for dementia caregivers: A randomized, controlled trial. *Alzheimer's Dement*. 2015 May;11(5):541-8. doi: <https://dx.doi.org/10.1016/j.jalz.2014.05.1752>. PMID: 25074341.
29. Arango-Lasprilla JC, Panyavin I, Merchan EJH, et al. Evaluation of a group cognitive-behavioral dementia caregiver intervention in Latin America. *American Journal of Alzheimer's Disease and Other Dementias*. 2014 Sep;29(6):548-55. doi: <http://dx.doi.org/10.1177/1533317514523668>. PMID: 2014-38564-010.
30. Livingston G, Barber J, Rapaport P, et al. START (STrAtegies for RelaTives) study: a pragmatic randomised controlled trial to determine the clinical effectiveness and cost-effectiveness of a manual-based coping strategy programme in promoting the mental health of carers of people with dementia. *Health Technol Assess*. 2014 Oct;18(61):1-242. doi: <https://dx.doi.org/10.3310/hta18610>. PMID: 25300037.
31. Martin-Carrasco M, Dominguez-Panchon AI, Gonzalez-Fraile E, et al. Effectiveness of a psychoeducational intervention group program in the reduction of the burden experienced by caregivers of patients with dementia: the EDUCA-II randomized trial. *Alzheimer Dis Assoc Disord*. 2014 Jan-Mar;28(1):79-87. doi: <https://dx.doi.org/10.1097/WAD.0000000000000003>. PMID: 24113563.
32. Passoni S, Moroni L, Toraldo A, et al. Cognitive behavioral group intervention for Alzheimer caregivers. *Alzheimer Dis Assoc Disord*. 2014;28(3):275-82. doi: 10.1097/WAD.0000000000000033. PMID: 107799151. Language: English. Entry Date: 20150508. Revision Date: 20150712. Publication Type: Journal Article.
33. Bruvik FK, Allore HG, Ranhoff AH, et al. The effect of psychosocial support intervention on depression in patients with dementia and their family caregivers: an assessor-blinded randomized controlled trial. *Dement Geriatr Cogn Dis Extra*. 2013;3(1):386-97. doi: <https://dx.doi.org/10.1159/000355912>. PMID: 24348500.
34. Huang HL, Kuo LM, Chen YS, et al. A home-based training program improves caregivers' skills and dementia patients' aggressive behaviors: a randomized controlled trial. *Am J Geriatr Psychiatry*. 2013 Nov;21(11):1060-70. doi: <https://dx.doi.org/10.1016/j.jagp.2012.09.009>. PMID: 23933422.
35. Judge KS, Yarry SJ, Looman WJ, et al. Improved Strain and Psychosocial Outcomes for Caregivers of Individuals with Dementia: Findings from Project ANSWERS. *Gerontologist*. 2013 Apr;53(2):280-92. doi: <https://dx.doi.org/10.1093/geront/gns076>. PMID: 22899427.
36. Kajiyama B, Thompson LW, Eto-Iwase T, et al. Exploring the effectiveness of an internet-based program for reducing caregiver distress using the iCare Stress Management e-Training Program.[Erratum appears in *Aging Ment Health*. 2013;17(5):c1]. *Aging Ment Health*. 2013;17(5):544-54. doi: <https://dx.doi.org/10.1080/13607863.2013.775641>. PMID: 23461355.
37. Kuo LM, Huang HL, Huang HL, et al. A home-based training program improves Taiwanese family caregivers' quality of life and decreases their risk for depression: a randomized controlled trial. *Int J Geriatr Psychiatry*. 2013 May;28(5):504-13. doi: <https://dx.doi.org/10.1002/gps.3853>. PMID: 22778053.
38. Moore RC, Chattillion EA, Ceglowski J, et al. A randomized clinical trial of Behavioral Activation (BA) therapy for improving psychological and physical health in dementia caregivers: results of the Pleasant Events Program (PEP). *Behav Res Ther*. 2013 Oct;51(10):623-32. doi: <https://dx.doi.org/10.1016/j.brat.2013.07.005>. PMID: 23916631.
39. Joling KJ, van Marwijk HW, Smit F, et al. Does a family meetings intervention prevent depression and anxiety in family caregivers of dementia patients? A randomized trial. *PLoS ONE*. 2012;7(1):e30936. doi: <https://dx.doi.org/10.1371/journal.pone.0030936>. PMID: 22303473.
40. Waldorff FB, Buss DV, Eckermann A, et al. Efficacy of psychosocial intervention in patients with mild Alzheimer's disease: the multicentre, rater blinded, randomised Danish Alzheimer Intervention Study (DAISY). *Bmj*. 2012 Jul 17;345:e4693. doi: <https://dx.doi.org/10.1136/bmj.e4693>. PMID: 22807076.
41. Chu H, Yang CY, Liao YH, et al. The effects of a support group on dementia caregivers' burden and depression. *Journal of aging and health*. 2011;23(2):228-41. PMID: CN-00858523.
42. de Rotrou J, Cantegreil I, Fauconau V, et al. Do patients diagnosed with Alzheimer's disease benefit from a psycho-educational programme for family caregivers? A randomised controlled study. *Int J Geriatr Psychiatry*. 2011 Aug;26(8):833-42. doi: <https://dx.doi.org/10.1002/gps.2611>. PMID: 20922772.
43. Guerra M, Ferri CP, Fonseca M, et al. Helping carers to care: the 10/66 dementia research group's randomized control trial of a caregiver intervention in Peru. *Rev Bras Psiquiatr*. 2011 Mar;33(1):47-54. PMID: 20602013.
44. Losada A, Marquez-Gonzalez M, Romero-Moreno R. Mechanisms of action of a psychological intervention for dementia caregivers: effects of behavioral activation and modification of dysfunctional thoughts. *Int J Geriatr Psychiatry*. 2011 Nov;26(11):1119-27. doi: <https://dx.doi.org/10.1002/gps.2648>. PMID: 21061414.
45. Spijker A, Wollersheim H, Teerenstra S, et al. Systematic care for caregivers of patients with dementia: a multicenter, cluster-randomized, controlled trial. *Am J Geriatr Psychiatry*. 2011 Jun;19(6):521-31. doi: <https://dx.doi.org/10.1097/JGP.0b013e3182110599>. PMID: 21358385.
46. Voigt-Radloff S, Graff M, Leonhart R, et al. A multicentre RCT on community occupational therapy in Alzheimer's disease: 10 sessions are not better than one consultation. *BMJ Open*. 2011;1. doi: doi:10.1136/bmjopen-2011-000096.

47. Wang LQ, Chien WT. Randomised controlled trial of a family-led mutual support programme for people with dementia. *J Clin Nurs*. 2011 Aug;20(15-16):2362-6. doi: <https://dx.doi.org/10.1111/j.1365-2702.2011.03746.x>. PMID: 21752121.
48. Wilz G, Schinkothe D, Soellner R. Goal attainment and treatment compliance in a cognitive-behavioral telephone intervention for family caregivers of persons with dementia. *GeroPsych: The Journal of Gerontopsychology and Geriatric Psychiatry*. 2011 01 Sep;24(3):115-25. doi: <http://dx.doi.org/10.1024/1662-9647/a000043>. PMID: 362501128.
49. Gitlin LN, Winter L, Dennis MP, et al. Targeting and managing behavioral symptoms in individuals with dementia: a randomized trial of a nonpharmacological intervention. *J Am Geriatr Soc*. 2010 Aug;58(8):1465-74. doi: <https://dx.doi.org/10.1111/j.1532-5415.2010.02971.x>. PMID: 20662955.
50. Gitlin LN, Winter L, Dennis MP, et al. A biobehavioral home-based intervention and the well-being of patients with dementia and their caregivers: the COPE randomized trial. *JAMA: Journal of the American Medical Association*. 2010;304(9):983-91. doi: 10.1001/jama.2010.1253. PMID: 105093294. Language: English. Entry Date: 20100917. Revision Date: 20161116. Publication Type: journal article.
51. Kurz A, Wagenpfeil S, Hallauer J, et al. Evaluation of a brief educational program for dementia carers: the AENEAS study. *Int J Geriatr Psychiatry*. 2010 Aug;25(8):861-9. doi: <https://dx.doi.org/10.1002/gps.2428>. PMID: 19946869.
52. Williams V, Bishop-Fitzpatrick L, Lane J, et al. Video-Based Coping Skills to Reduce Health Risk and Improve Psychological and Physical Well-Being in Alzheimer's Disease Family Caregivers. *Psychosom Med*. 2010;72(9):897-904. doi: 10.1097/PSY.0b013e3181fc2d09. PMID: CN-00892462.
53. Gavrilova SI, Ferri CP, Mikhaylova N, et al. Helping carers to care--the 10/66 dementia research group's randomized control trial of a caregiver intervention in Russia. *Int J Geriatr Psychiatry*. 2009 Apr;24(4):347-54. doi: <https://dx.doi.org/10.1002/gps.2126>. PMID: 18814197.
54. Martin-Carrasco M, Martin MF, Valero CP, et al. Effectiveness of a psychoeducational intervention program in the reduction of caregiver burden in Alzheimer's disease patients' caregivers. *Int J Geriatr Psychiatry*. 2009 May;24(5):489-99. doi: <https://dx.doi.org/10.1002/gps.2142>. PMID: 18949763.
55. Signe A, Elmstahl S. Psychosocial intervention for family caregivers of people with dementia reduces caregiver's burden: development and effect after 6 and 12 months. *Scand J Caring Sci*. 2008;22(1):98-109. doi: 10.1111/j.1471-6712.2007.00498.x. PMID: CN-00630033.
56. Gallagher-Thompson D, Gray HL, Dupart T, et al. Effectiveness of Cognitive/Behavioral Small Group Intervention for Reduction of Depression and Stress in Non-Hispanic White and Hispanic/Latino Women Dementia Family Caregivers: Outcomes and Mediators of Change. *Journal of Rational-Emotive & Cognitive Behavior Therapy*. 2008 Dec 01;26(4):286-303. PMID: 25067886.
57. Marquez-Gonzalez M, Losada A, Izal M, et al. Modification of dysfunctional thoughts about caregiving in dementia family caregivers: description and outcomes of an intervention programme. *Aging Ment Health*. 2007 Nov;11(6):616-25. PMID: 18074249.
58. Ulstein ID, Sandvik L, Wyller TB, et al. A one-year randomized controlled psychosocial intervention study among family carers of dementia patients--effects on patients and carers. *Dement Geriatr Cogn Disord*. 2007;24(6):469-75. PMID: 17986818.
59. Gonyea JG, O'Connor MK, Boyle PA. Project CARE: a randomized controlled trial of a behavioral intervention group for Alzheimer's disease caregivers. *Gerontologist*. 2006 Dec;46(6):827-32. PMID: 17169938.
60. Graff MJ, Vernooij-Dassen MJ, Thijssen M, et al. Community based occupational therapy for patients with dementia and their care givers: randomised controlled trial. *Bmj*. 2006 Dec 09;333(7580):1196. PMID: 17114212.
61. Beauchamp N, Irvine AB, Seeley J, et al. Worksite-based internet multimedia program for family caregivers of persons with dementia. *Gerontologist*. 2005 Dec;45(6):793-801. PMID: 16326661.
62. Farran CJ, Gilley DW, McCann JJ, et al. Psychosocial interventions to reduce depressive symptoms of dementia caregivers: A randomized clinical trial comparing two approaches. *Journal of Mental Health and Aging*. 2004 Winter;10(4):337-50. PMID: 41552352.
63. Burgio L, Stevens A, Guy D, et al. Impact of two psychosocial interventions on white and African American family caregivers of individuals with dementia. *Gerontologist*. 2003 Aug;43(4):568-79. PMID: 12937335.
64. Burns R, Nichols LO, Martindale-Adams J, et al. Primary care interventions for dementia caregivers: 2-year outcomes from the REACH study. *Gerontologist*. 2003 Aug;43(4):547-55. PMID: 12937333.
65. Coon DW, Thompson L, Steffen A, et al. Anger and depression management: psychoeducational skill training interventions for women caregivers of a relative with dementia. *Gerontologist*. 2003 Oct;43(5):678-89. PMID: 14570964.
66. Gallagher-Thompson D, Coon DW, Solano N, et al. Change in indices of distress among Latino and Anglo female caregivers of elderly relatives with dementia: site-specific results from the REACH national collaborative study. *Gerontologist*. 2003 Aug;43(4):580-91. PMID: 12937336.
67. Gitlin LN, Winter L, Corcoran M, et al. Effects of the home environmental skill-building program on the caregiver-care recipient dyad: 6-month outcomes from the Philadelphia REACH Initiative. *Gerontologist*. 2003 Aug;43(4):532-46. PMID: 12937332.
68. Hebert R, Levesque L, Vezina J, et al. Efficacy of a psychoeducative group program for caregivers of demented persons living at home: a randomized controlled trial. *J Gerontol B Psychol Sci Soc Sci*. 2003 Jan;58(1):S58-67. PMID: 12496309.
69. Fung W, Chien W. The effectiveness of a mutual support group for family caregivers of a relative with dementia. *Arch Psychiatr Nurs*. 2002;16(3):134-44. PMID: 106975399. Language: English. Entry Date: 20021108. Revision Date: 20150819. Publication Type: Journal Article.
70. Stolley JM, Reed D, Buckwalter KC. Caregiving appraisal and interventions based on the progressively lowered stress threshold model. *Am J Alzheimers Dis Other Dement*. 2002 Mar-Apr;17(2):110-20. PMID: 11954669.
71. Wright LK, Litaker M, Laraia MT, et al. Continuum of care for Alzheimer's disease: a nurse education and counseling program. *Issues Ment Health Nurs*. 2001 Apr-May;22(3):231-52. PMID: 11885210.
72. Buckwalter KC, Gerdner L, Kohout F, et al. A nursing intervention to decrease depression in family caregivers of persons with dementia. *Arch Psychiatr Nurs*. 1999 Apr;13(2):80-8. PMID: 10222636.
73. Chang BL. Cognitive-behavioral intervention for homebound caregivers of persons with dementia. *Nurs Res*. 1999 May-Jun;48(3):173-82. PMID: 10337848.
74. Ostwald SK, Hepburn KW, Caron W, et al. Reducing caregiver burden: a randomized psychoeducational intervention for caregivers of persons with dementia. *Gerontologist*. 1999 Jun;39(3):299-309. PMID: 10396888.
75. Gitlin LN, Mann WC, Vogel WB, et al. A non-pharmacologic approach to address challenging behaviors of Veterans with dementia: description of the tailored activity program-VA randomized trial. *BMC geriatr*. 2013 Sep 23;13:96. doi: <https://dx.doi.org/10.1186/1471-2318-13-96>. PMID: 24060106.
76. Melunsky N, Crellin N, Dudzinski E, et al. The experience of family carers attending a joint reminiscence group with people with dementia: A thematic analysis. *Dementia*. 2015 Nov;14(6):842-59. doi: <https://dx.doi.org/10.1177/1471301213516332>. PMID: 24381218.
77. Charlesworth G, Burnell K, Beecham J, et al. Peer support for family carers of people with dementia, alone or in combination with group reminiscence in a factorial design: study protocol for a randomised controlled trial. *Trials*. 2011 Sep 15;12:205. doi: <https://dx.doi.org/10.1186/1745-6215-12-205>. PMID: 21917187.
78. Cheng ST, Chan WC, Lam LCW. Long-Term Outcomes of the Benefit-Finding Group Intervention for Alzheimer Family Caregivers: A Cluster-Randomized Double-Blind Controlled Trial. *Am J Geriatr Psychiatry*. 2019 Sep;27(9):984-94. doi: <https://dx.doi.org/10.1016/j.jagp.2019.03.013>. PMID: 31076215.
79. Cheng ST, Mak EPM, Kwok T, et al. Benefit-Finding Intervention Delivered Individually to Alzheimer Family Caregivers: Longer-Term Outcomes of a Randomized Double-Blind Controlled Trial. *Journals of Gerontology Series B Psychological Sciences & Social Sciences*. 2019 Sep 26;26:26. doi: <https://dx.doi.org/10.1093/geronb/gbz118>. PMID: 31556447.
80. Cheng ST, Lau RW, Mak EP, et al. Benefit-finding intervention for Alzheimer caregivers: conceptual framework, implementation issues, and preliminary efficacy. *Gerontologist*. 2014 Dec;54(6):1049-58. doi: <https://dx.doi.org/10.1093/geront/gnu018>. PMID: 24688081.
81. Laakkonen ML, Savikko N, Holtta E, et al. Self-management groups for people with dementia and their spousal caregivers. A randomized, controlled trial. Baseline findings and feasibility. *European Geriatric Medicine*. 2013 December;4(6):389-93. doi: <http://dx.doi.org/10.1016/j.eurger.2013.09.006>. PMID: 52813196.
82. Laakkonen ML, Holtta EH, Savikko N, et al. Psychosocial group intervention to enhance self-management skills of people with dementia and their caregivers: study protocol for a randomized controlled trial. *Trials*. 2012 Aug 07;13:133. doi: <https://dx.doi.org/10.1186/1745-6215-13-133>. PMID: 22871107.
83. Otero P, Smit F, Cuijpers P, et al. Differential response to depression prevention among a sample of informal caregivers: Moderator analysis of longer-term follow-up trial data. *Psychiatry Res*. 2015 Dec 15;230(2):271-8. doi: <https://dx.doi.org/10.1016/j.psychres.2015.09.005>. PMID: 26456895.
84. Tremont G, Davis JD, Ott BR, et al. Randomized Trial of the Family Intervention: Telephone Tracking-Caregiver for Dementia Caregivers: Use of Community and Healthcare Resources. *J Am Geriatr Soc*. 2017 May;65(5):924-30. doi: <https://dx.doi.org/10.1111/jgs.14684>. PMID: 28008609.
85. Tremont G, Davis J, Papandonatos GD, et al. A telephone intervention for dementia caregivers: background, design, and baseline characteristics. *Contemp Clin Trials*. 2013 Nov;36(2):338-47. doi: <https://dx.doi.org/10.1016/j.cct.2013.07.011>. PMID: 23916916.
86. Voigt-Radloff S, Graff M, Leonhart R, et al. Why did an effective Dutch complex psycho-social intervention for people with dementia not work in the German healthcare context? Lessons learnt from a process evaluation alongside a multicentre RCT. *BMJ Open*. 2011;1(1)(no pagination)(e000094). doi: <http://dx.doi.org/10.1136/bmjopen-2011-000094>. PMID: 611053874.
87. Voigt-Radloff S, Graff M, Leonhart R, et al. WHEDA study: Effectiveness of occupational therapy at home for older people with dementia and their caregivers - The design of a pragmatic randomised controlled trial evaluating a Dutch programme in seven German centres. *BMC geriatr*. 2009;9(1)(no pagination)(44). doi: <http://dx.doi.org/10.1186/1471-2318-9-44>. PMID: 355485226.
88. Graff MJ, Adang EM, Vernooij-Dassen MJ, et al. Community occupational therapy for older patients with dementia and their care givers: cost effectiveness study. *Bmj*. 2008 Jan 19;336(7636):134-8. doi: <https://dx.doi.org/10.1136/bmj.39408.481898.BE>. PMID: 18171718.
89. Graff MJ, Vernooij-Dassen MJ, Thijssen M, et al. Effects of community occupational therapy on quality of life, mood, and health status in dementia patients and their caregivers: a randomized controlled trial. *J Gerontol A Biol Sci Med Sci*. 2007 Sep;62(9):1002-9. PMID: 17895439.
90. Farran CJ, Gilley DW, McCann JJ, et al. Efficacy of behavioral interventions for dementia caregivers. *West J Nurs Res*. 2007 Dec;29(8):944-60. PMID: 17596639.
91. Farran CJ, Loukissa D, Perraud S, et al. Alzheimer's disease caregiving information and skills. Part II: family caregiver issues and concerns. *Res Nurs Health*. 2004 Feb;27(1):40-51. PMID: 14745855.
92. Rabinowitz YG, Mausbach BT, Coon DW, et al. The moderating effect of self-efficacy on intervention response in women family caregivers of older adults with dementia. *Am J Geriatr Psychiatry*. 2006 Aug;14(8):642-9. PMID: 16861368.
93. Gitlin LN, Corcoran M, Winter L, et al. A randomized, controlled trial of a home environmental intervention: effect on efficacy and upset in caregivers and on daily function of persons with dementia. *Gerontologist*. 2001 Feb;41(1):4-14. PMID: 11220813.
94. Voigt-Radloff S, de Werd MM, Leonhart R, et al. Structured relearning of activities of daily living in dementia: the randomized controlled REDALI-DEM trial on errorless learning. *Alzheimers Res Ther*. 2017 Mar 23;9(1):22. doi: <https://dx.doi.org/10.1186/s13195-017-0247-9>. PMID: 28335810.
95. Meichsner F, Theurer C, Wilz G. Acceptance and treatment effects of an internet-delivered cognitive-behavioral intervention for family caregivers of people with dementia: A randomized-controlled trial. *J Clin Psychol*. 2019 Apr;75(4):594-613. doi: <https://dx.doi.org/10.1002/jclp.22739>. PMID: 30597537.
96. Orgeta V, Tuijt R, Leung P, et al. Behavioral Activation for Promoting Well-Being in Mild Dementia: Feasibility and Outcomes of a Pilot Randomized Controlled Trial. *J Alzheimers Dis*. 2019 Oct 08;08:08. doi: <https://dx.doi.org/10.3233/JAD-190696>. PMID: 31609689.
97. Ta Park VM, Ton V, Tiet QQ, et al. Promising results from a pilot study to reduce distress in Vietnamese American dementia and memory loss caregivers. *Alzheimers Dement (N Y)*. 2019;5:319-27. doi: <https://dx.doi.org/10.1016/j.trci.2019.05.006>. PMID: 31384663.

98. Ta Park VM, Ton V, Yeo G, et al. Vietnamese American Dementia Caregivers' Perceptions and Experiences of a Culturally Tailored, Evidence-Based Program to Reduce Stress and Depression. *J Gerontol Nurs*. 2019 01 Sep;45(9):39-50. doi: <http://dx.doi.org/10.3928/00989134-20190813-05>. PMID: 629157769.
99. Whitlatch CJ, Heid AR, Femia EE, et al. The Support, Health, Activities, Resources, and Education program for early stage dementia: Results from a randomized controlled trial. *Dementia*. 2019 08;18(6):2122-39. doi: <https://dx.doi.org/10.1177/1471301217743033>. PMID: 29171296.
100. Yoo R, Yeom J, Kim GH, et al. A Multicenter, Randomized Clinical Trial to Assess the Efficacy of a Therapeutic Intervention Program for Caregivers of People with Dementia. *J Clin Neurol*. 2019 Apr;15(2):235-42. doi: <https://dx.doi.org/10.3988/jcn.2019.15.2.235>. PMID: 30938110.
101. Boots LM, de Vugt ME, Kempen GI, et al. Effectiveness of the blended care self-management program "Partner in Balance" for early-stage dementia caregivers: study protocol for a randomized controlled trial. *Trials*. 2016 May 04;17(1):231. doi: <https://dx.doi.org/10.1186/s13063-016-1351-z>. PMID: 27142676.
102. Boots LM, de Vugt ME, Smeets CM, et al. Implementation of the Blended Care Self-Management Program for Caregivers of People With Early-Stage Dementia (Partner in Balance): Process Evaluation of a Randomized Controlled Trial. *J Med Internet Res*. 2017 Dec 19;19(12):e423. doi: <https://dx.doi.org/10.2196/jmir.7666>. PMID: 29258980.
103. de Oliveira AM, Radanovic M, Homem de Mello PC, et al. An intervention to reduce neuropsychiatric symptoms and caregiver burden in dementia: Preliminary results from a randomized trial of the tailored activity program-outpatient version. *Int J Geriatr Psychiatry*. 2018 Jul 23;23:23. doi: <https://dx.doi.org/10.1002/gps.4958>. PMID: 30035341.
104. Kales HC, Gitlin LN, Stanislawski B, et al. Effect of the WeCareAdvisor™ on family caregiver outcomes in dementia: a pilot randomized controlled trial. *BMC geriatr*. 2018 05 10;18(1):113. doi: <https://dx.doi.org/10.1186/s12877-018-0801-8>. PMID: 29747583.
105. Meichsner F, Theurer C, Wilz G. Acceptance and treatment effects of an internet-delivered cognitive-behavioral intervention for family caregivers of people with dementia: A randomized-controlled trial. *J Clin Psychol*. 2018 Dec 31;31:31. doi: <https://dx.doi.org/10.1002/jclp.22739>. PMID: 30597537.
106. Novelli M, Machado SCB, Lima GB, et al. Effects of the Tailored Activity Program in Brazil (TAP-BR) for Persons With Dementia: A Randomized Pilot Trial. *Alzheimer Dis Assoc Disord*. 2018 Apr 25;25:25. doi: <https://dx.doi.org/10.1097/WAD.0000000000000256>. PMID: 29698251.
107. van Knippenberg RJM, de Vugt ME, Ponds RW, et al. An Experience Sampling Method Intervention for Dementia Caregivers: Results of a Randomized Controlled Trial. *Am J Geriatr Psychiatry*. 2018 Dec;26(12):1231-43. doi: <https://dx.doi.org/10.1016/j.jagp.2018.06.004>. PMID: 30126766.
108. Meichsner F, Topfer NF, Reder M, et al. Telephone-Based Cognitive Behavioral Intervention Improves Dementia Caregivers' Quality of Life. *American Journal of Alzheimer's Disease and other Dementias*. 2019. doi: <http://dx.doi.org/10.1177/1533317518822100>. PMID: 626130265.
109. Topfer NF, Wilz G. Tele.Tandem increases the psychosocial resource utilization of dementia caregivers. *GeroPsych: The Journal of Gerontopsychology and Geriatric Psychiatry*. 2018;31(4):173-83. doi: <http://dx.doi.org/10.1024/1662-9647/a000197>. PMID: 626123635.
110. Lok N, Bademli K. Pilot testing of the "First You Should Get Stronger" program among caregivers of older adults with dementia. *Arch Gerontol Geriatr*. 2017 Jan - Feb;68:84-9. doi: <https://dx.doi.org/10.1016/j.archger.2016.09.006>. PMID: 27689315.
111. O'Connor CM, Clemson L, Brodaty H, et al. The tailored activity program (TAP) to address behavioral disturbances in frontotemporal dementia: a feasibility and pilot study. *Disabil Rehabil*. 2017 Oct 15;1-12. doi: <https://dx.doi.org/10.1080/09638288.2017.1387614>. PMID: 29034719.
112. Supiano KP, Haynes LB, Pond V. The Process of Change in Complicated Grief Group Therapy for Bereaved Dementia Caregivers: An Evaluation Using the Meaning of Loss Codebook. *J Gerontol Soc Work*. 2017 Feb-Mar;60(2):155-69. doi: <https://dx.doi.org/10.1080/01634372.2016.1274930>. PMID: 28051926.
113. Nunez-Naveira L, Alonso-Bua B, de Labra C, et al. UnderstAID, an ICT Platform to Help Informal Caregivers of People with Dementia: A Pilot Randomized Controlled Study. *Biomed Res Int*. 2016;2016:5726465. doi: <https://dx.doi.org/10.1155/2016/5726465>. PMID: 28116300.
114. Sepe-Monti M, Vanacore N, Bartorelli L, et al. The Savvy Caregiver Program: A Probe Multicenter Randomized Controlled Pilot Trial in Caregivers of Patients Affected by Alzheimer's Disease. *J Alzheimers Dis*. 2016 10 04;54(3):1235-46. PMID: 27567824.
115. Zauszniewski JA, Lekhak N, Burant CJ, et al. Preliminary Evidence for Effectiveness of Resourcefulness Training in Women Dementia Caregivers. *J Fam Med*. 2016;3(5). PMID: 27500286.
116. Zauszniewski JA, Lekhak N, Yolpant W, et al. Need for Resourcefulness Training for Women Caregivers of Elders with Dementia. *Issues Ment Health Nurs*. 2015;36(12):1007-12. doi: <https://dx.doi.org/10.3109/01612840.2015.1075236>. PMID: 26735507.
117. Au A, Gallagher-Thompson D, Wong MK, et al. Behavioral activation for dementia caregivers: scheduling pleasant events and enhancing communications. *Clin Interv Aging*. 2015;10:611-9. doi: <https://dx.doi.org/10.2147/CIA.S72348>. PMID: 25848237.
118. Au A. Developing Volunteer-Assisted Behavioral Activation Teleprograms to Meet the Needs of Chinese Dementia Caregivers. *Clin Gerontol*. 2015;38(3):190-202. PMID: CN-01071414.
119. Blom MM, Bosmans JE, Cuijpers P, et al. Effectiveness and cost-effectiveness of an internet intervention for family caregivers of people with dementia: design of a randomized controlled trial. *BMC Psychiatry*. 2013 Jan 10;13:17. doi: <https://dx.doi.org/10.1186/1471-244X-13-17>. PMID: 23305463.
120. Pot AM, Blom MM, Willemse BM. Acceptability of a guided self-help Internet intervention for family caregivers: mastery over dementia. *Int Psychogeriatr*. 2015 Aug;27(8):1343-54. doi: <https://dx.doi.org/10.1017/S1041610215000034>. PMID: 25648589.
121. Chen HM, Huang MF, Yeh YC, et al. Effectiveness of coping strategies intervention on caregiver burden among caregivers of elderly patients with dementia. *Psychogeriatrics*. 2015 Mar;15(1):20-5. doi: <https://dx.doi.org/10.1111/psyg.12071>. PMID: 25515800.
122. Cristancho-Lacroix V, Wrobel J, Cantegreil-Kallen I, et al. A web-based psychoeducational program for informal caregivers of patients with Alzheimer's disease: a pilot randomized controlled trial. *J Med Internet Res*. 2015 May 12;17(5):e117. doi: <https://dx.doi.org/10.2196/jmir.3717>. PMID: 25967983.
123. Cristancho-Lacroix V, Moulin F, Wrobel J, et al. A web-based program for informal caregivers of persons with Alzheimer's disease: an iterative user-centered design. *JMIR Res Protoc*. 2014 Sep 15;3(3):e46. doi: <https://dx.doi.org/10.2196/resprot.3607>. PMID: 25263541.
124. Gaugler JE, Reese M, Sauld J. A Pilot Evaluation of Psychosocial Support for Family Caregivers of Relatives with Dementia in Long-Term Care: The Residential Care Transition Module. *Res*. 2015 Jul-Aug;8(4):161-72. doi: <https://dx.doi.org/10.3928/19404921-20150304-01>. PMID: 25751083.
125. Kamkhagi D, Costa A, Kusminsky S, et al. Benefits of psychodynamic group therapy on depression, burden and quality of life of family caregivers to Alzheimer's disease patients. *Revista de psiquiatria clinica*. 2015;42(6):157-60. doi: 10.1590/0101-60830000000067. PMID: CN-01162210.
126. Paun O, Farran CJ, Fogg L, et al. A chronic grief intervention for dementia family caregivers in long-term care. *West J Nurs Res*. 2015 Jan;37(1):6-27. doi: <https://dx.doi.org/10.1177/0193945914521040>. PMID: 24510968.
127. Abouafia-Brakha T, Suchecki D, Gouveia-Paulino F, et al. Cognitive-behavioural group therapy improves a psychophysiological marker of stress in caregivers of patients with Alzheimer's disease. *Aging Ment Health*. 2014;18(6):801-8. doi: <https://dx.doi.org/10.1080/13607863.2014.880406>. PMID: 24499394.
128. Dowling GA, Merrilees J, Mastick J, et al. Life enhancing activities for family caregivers of people with frontotemporal dementia. *Alzheimer Dis Assoc Disord*. 2014 Apr-Jun;28(2):175-81. doi: <https://dx.doi.org/10.1097/WAD.0b013e3182a6b905>. PMID: 24113564.
129. Gonzalez EW, Polansky M, Lippa CF, et al. Enhancing resourcefulness to improve outcomes in family caregivers and persons with Alzheimer's disease: a pilot randomized trial. *Int J Alzheimer Dis*. 2014;2014:323478. doi: <https://dx.doi.org/10.1155/2014/323478>. PMID: 25328754.
130. Judge KS, Yarry SJ, Orsulic-Jeras S. Acceptability and feasibility results of a strength-based skills training program for dementia caregiving dyads. *Gerontologist*. 2010 Jun;50(3):408-17. doi: <https://dx.doi.org/10.1093/geront/gnp138>. PMID: 19808841.
131. Judge K, Bass D, Snow A, et al. Partners in Dementia Care: a care coordination intervention for Individuals with dementia and their family caregivers. *In Gerontologist*
132. Kuo LM, Huang HL, Liang J, et al. A randomized controlled trial of a home-based training programme to decrease depression in family caregivers of persons with dementia. *J Adv Nurs*. 2017 Mar;73(3):585-98. doi: <https://dx.doi.org/10.1111/jan.13157>. PMID: 27653753.
133. Kuo LM, Huang HL, Hsu WC, et al. Health-related quality of life and self-efficacy of managing behavior problems for family caregivers of vascular dementia and Alzheimer's disease patients. *Dementia and Geriatric Cognitive Disorders*. 2014 16 Apr;38(5-6):310-20. doi: <http://dx.doi.org/10.1159/000360414>. PMID: 603852835.
134. Kwok T, Wong B, Ip I, et al. Telephone-delivered psychoeducational intervention for Hong Kong Chinese dementia caregivers: a single-blinded randomized controlled trial. *Clin Interv Aging*. 2013;8:1191-7. doi: <https://dx.doi.org/10.2147/CIA.S48264>. PMID: 24072965.
135. Joling KJ, van Marwijk HW, van der Horst HE, et al. Effectiveness of family meetings for family caregivers on delaying time to nursing home placement of dementia patients: a randomized trial. *PLoS ONE*. 2012;7(8):e42145. doi: <https://dx.doi.org/10.1371/journal.pone.0042145>. PMID: 22876304.
136. Joling KJ, Bosmans JE, van Marwijk HW, et al. The cost-effectiveness of a family meetings intervention to prevent depression and anxiety in family caregivers of patients with dementia: a randomized trial. *Trials*. 2013 Sep 22;14:305. doi: <https://dx.doi.org/10.1186/1745-6215-14-305>. PMID: 24053631.
137. Joling KJ, van Hout HP, Scheltens P, et al. (Cost)-effectiveness of family meetings on indicated prevention of anxiety and depressive symptoms and disorders of primary family caregivers of patients with dementia: design of a randomized controlled trial. *BMC geriatr*. 2008 Jan 21;8:2. doi: <https://dx.doi.org/10.1186/1471-2318-8-2>. PMID: 18208607.
138. Davis JD, Tremont G, Bishop DS, et al. A telephone-delivered psychosocial intervention improves dementia caregiver adjustment following nursing home placement. *Int J Geriatr Psychiatry*. 2011 Apr;26(4):380-7. doi: <https://dx.doi.org/10.1002/gps.2537>. PMID: 20842759.
139. Spijker A, Verhey F, Graff M, et al. Systematic care for caregivers of people with dementia in the ambulatory mental health service: designing a multicentre, cluster, randomized, controlled trial. *BMC geriatr*. 2009 Jun 07;9:21. doi: <https://dx.doi.org/10.1186/1471-2318-9-21>. PMID: 19500421.
140. Spijker A, Teerenstra S, Wollersheim H, et al. Influence of adherence to a systematic care program for caregivers of dementia patients. *Am J Geriatr Psychiatry*. 2013 Jan;21(1):26-36. doi: <https://dx.doi.org/10.1016/j.jagp.2012.10.003>. PMID: 23290200.
141. Meichsner F, Schinkothe D, Wilz G. Managing Loss and Change: Grief Interventions for Dementia Caregivers in a CBT-Based Trial. *Am J Alzheimers Dis Other Demen*. 2016 May;31(3):231-40. doi: <https://dx.doi.org/10.1177/1533317515602085>. PMID: 26311735.
142. Meichsner F, Wilz G. Dementia caregivers' coping with pre-death grief: effects of a CBT-based intervention. *Aging Ment Health*. 2018 Feb;22(2):218-25. doi: <https://dx.doi.org/10.1080/13607863.2016.1247428>. PMID: 27792398.
143. Au A, Li S, Lee K, et al. The Coping with Caregiving Group Program for Chinese caregivers of patients with Alzheimer's disease in Hong Kong. *In Patient education and counseling*
144. Gallagher-Thompson D, Wang PC, Liu W, et al. Effectiveness of a psychoeducational skill training DVD program to reduce stress in Chinese American dementia caregivers: results of a preliminary study. *Aging Ment Health*. 2010 Apr;14(3):263-73. doi: <http://dx.doi.org/10.1080/13607860903420989>. PMID: 359062574.
145. Gitlin LN, Winter L, Dennis MP, et al. A biobehavioral home-based intervention and the well-being of patients with dementia and their caregivers: the COPE randomized trial. *Jama*. 2010 Sep 01;304(9):983-91. doi: <https://dx.doi.org/10.1001/jama.2010.1253>. PMID: 20810376.
146. Dias A, Dewey ME, D'Souza J, et al. The effectiveness of a home care program for supporting caregivers of persons with dementia in developing countries: a randomised controlled trial from Goa, India. *PLoS ONE*. 2008 Jun 04;3(6):e2333. doi: <https://dx.doi.org/10.1371/journal.pone.0002333>. PMID: 18523642.

147. Gitlin LN, Winter L, Burke J, et al. Tailored activities to manage neuropsychiatric behaviors in persons with dementia and reduce caregiver burden: a randomized pilot study. *Am J Geriatr Psychiatry*. 2008 Mar;16(3):229-39. doi: <https://dx.doi.org/10.1097/JGP.0b013e318160da72>. PMID: 18310553.
148. Gitlin LN, Hodgson N, Jutkowitz E, et al. The cost-effectiveness of a nonpharmacologic intervention for individuals with dementia and family caregivers: the tailored activity program. *Am J Geriatr Psychiatry*. 2010 Jun;18(6):510-9. doi: <https://dx.doi.org/10.1097/JGP.0b013e3181c37d13>. PMID: 20847903.
149. Gitlin LN, Winter L, Vause Earland T, et al. The Tailored Activity Program to reduce behavioral symptoms in individuals with dementia: feasibility, acceptability, and replication potential. *Gerontologist*. 2009 Jun;49(3):428-39. doi: <https://dx.doi.org/10.1093/geront/gnp087>. PMID: 19420314.
150. Tremont G, Davis JD, Bishop DS, et al. Telephone-delivered psychosocial intervention reduces burden in dementia caregivers. *Dementia: The International Journal of Social Research and Practice*. 2008 Nov;7(4):503-20. doi: <http://dx.doi.org/10.1177/1471301208096632>. PMID: 2008-16304-006.
151. Epstein-Lubow G, Davis JD, Miller IW, et al. Persisting burden predicts depressive symptoms in dementia caregivers. *J Geriatr Psychiatry Neurol*. 2008 Sep;21(3):198-203. doi: <https://dx.doi.org/10.1177/0891988708320972>. PMID: 18838742.
152. Gant JR, Steffen AM, Lauderdale SA. Comparative outcomes of two distance-based interventions for male caregivers of family members with dementia. *Am J Alzheimers Dis Other Demen*. 2007 Apr-May;22(2):120-8. PMID: 17545139.
153. Glueckauf R, Sharma D, Davis W, et al. Telephone-based cognitive-behavioral intervention for distressed rural dementia caregivers: initial findings. In *Clin Gerontol*
154. Rodriguez-Sanchez E, Patino-Alonso MC, Mora-Simon S, et al. Effects of a psychological intervention in a primary health care center for caregivers of dependent relatives: a randomized trial. *Gerontologist*. 2013 Jun;53(3):397-406. doi: <https://dx.doi.org/10.1093/geront/gns086>. PMID: 22899425.
155. Akkerman RL, Ostwald SK. Reducing anxiety in Alzheimer's disease family caregivers: the effectiveness of a nine-week cognitive-behavioral intervention. *Am J Alzheimers Dis Other Demen*. 2004 Mar-Apr;19(2):117-23. PMID: 15106393.
156. Nobili A, Riva E, Tettamanti M, et al. The effect of a structured intervention on caregivers of patients with dementia and problem behaviors: a randomized controlled pilot study. *Alzheimer Dis Assoc Disord*. 2004 Apr-Jun;18(2):75-82. PMID: 15249851.
157. Burns R, Nichols LO, Graney MJ, et al. Cognitive abilities of Alzheimer's patients: perceptions of Black and White caregivers. *Int J Aging Hum Dev*. 2006;62(3):209-19. PMID: 16625937.
158. Holland JM, Currier JM, Gallagher-Thompson D. Outcomes from the Resources for Enhancing Alzheimer's Caregiver Health (REACH) program for bereaved caregivers. *Psychol Aging*. 2009 Mar;24(1):190-202. doi: <https://dx.doi.org/10.1037/a0014303>. PMID: 19290751.
159. Gitlin LN, Hauck WW, Dennis MP, et al. Maintenance of effects of the home environmental skill-building program for family caregivers and individuals with Alzheimer's disease and related disorders. *J Gerontol A Biol Sci Med Sci*. 2005 Mar;60(3):368-74. PMID: 15860476.
160. Chee YK, Gitlin LN, Dennis MP, et al. Predictors of adherence to a skill-building intervention in dementia caregivers. *J Gerontol A Biol Sci Med Sci*. 2007 Jun;62(6):673-8. PMID: 17595426.
161. Hilgeman MM, Allen RS, Decoster J, et al. Positive aspects of caregiving as a moderator of treatment outcome over 12 months. *Psychol Aging*. 2007;22(2):361-71. PMID: 105991138. Language: English. Entry Date: 20080222. Revision Date: 20161222. Publication Type: journal article.
162. Lavoie JP, Ducharme F, Levesque L, et al. Understanding the outcomes of a psycho-educational group intervention for caregivers of persons with dementia living at home: a process evaluation. *Aging Ment Health*. 2005 Jan;9(1):25-34. PMID: 15841829.
163. Huang HL, Shyu YI, Chen MC, et al. A pilot study on a home-based caregiver training program for improving caregiver self-efficacy and decreasing the behavioral problems of elders with dementia in Taiwan. *Int J Geriatr Psychiatry*. 2003 Apr;18(4):337-45. PMID: 12673611.
164. Martin-Cook K, Remakel-Davis B, Svetlik D, et al. Caregiver attribution and resentment in dementia care. *Am J Alzheimers Dis Other Demen*. 2003 Nov-Dec;18(6):366-74. PMID: 14682086.
165. Garand L, Buckwalter KC, Lubaroff D, et al. A pilot study of immune and mood outcomes of a community-based intervention for dementia caregivers: the PLST intervention. *Arch Psychiatr Nurs*. 2002 Aug;16(4):156-67. PMID: 12143075.
166. Marriott A, Donaldson C, TARRIER N, et al. Effectiveness of cognitive-behavioural family intervention in reducing the burden of care in carers of patients with Alzheimer's disease. *Br J Psychiatry*. 2000 Jun;176:557-62. PMID: 10974962.
167. Steffen AM. Anger management for dementia caregivers: A preliminary study using video and telephone interventions. *Behavior Therapy*. 2000 Spr;31(2):281-99. doi: <http://dx.doi.org/10.1016/S0005-7894%2800%2980016-7>. PMID: 2000-16043-005.
168. Gerdner LA, Buckwalter KC, Reed D. Impact of a psychoeducational intervention on caregiver response to behavioral problems. *Nurs Res*. 2002 Nov-Dec;51(6):363-74. PMID: 12464756.
169. Chang BL, Nitta S, Carter PA, et al. Perceived helpfulness of telephone calls--providing support for caregivers of family members with dementia. *J Gerontol Nurs*. 2004 Sep;30(9):14-21. PMID: 15471059.
170. McCurry SM, Logsdon RG, Vitiello MV, et al. Successful behavioral treatment for reported sleep problems in elderly caregivers of dementia patients: a controlled study. *J Gerontol B Psychol Sci Soc Sci*. 1998 Mar;53(2):P122-9. PMID: 9520929.
171. Moniz-Cook E, Agar S, Gibson G, et al. A preliminary study of the effects of early intervention with people with dementia and their families in a memory clinic. *Aging Ment Health*. 1998;2(3):199-211. PMID: 107163779. Language: English. Entry Date: 20070101. Revision Date: 20150711. Publication Type: Journal Article.
172. Brodaty H, Gresham M, Luscombe G. The Prince Henry Hospital dementia caregivers' training programme. *Int J Geriatr Psychiatry*. 1997 Feb;12(2):183-92. PMID: 9097211.
173. Teri L, Logsdon RG, Uomoto J, et al. Behavioral treatment of depression in dementia patients: a controlled clinical trial. *J Gerontol B Psychol Sci Soc Sci*. 1997 Jul;52(4):P159-66. PMID: 9224439.
174. Gendron C, Poitras L, Dastoor DP, et al. Cognitive-behavioral group intervention for spousal caregivers: Findings and clinical considerations. *Clin Gerontol*. 1996;17(1):3-19. doi: http://dx.doi.org/10.1300/J018v17n01_02. PMID: 26250550.
175. Hebert R, Leclerc G, Bravo G, et al. Efficacy of a support group programme for care-givers of demented patients in the community: A randomized controlled trial. *Archives of Gerontology and Geriatrics*. 1994;18(1):1-14. doi: <http://dx.doi.org/10.1016/0167-4943%2894%2990043-4>. PMID: 24087060.
176. Hebert R, Girouard D, Leclerc G, et al. The impact of a support group programme for care-givers on the institutionalisation of demented patients. *Archives of Gerontology and Geriatrics*. 1995;20(2):129-34. doi: <http://dx.doi.org/10.1016/0167-4943%2894%2900590-4>. PMID: 25107702.
177. Tappen RM. The effect of skill training on functional abilities of nursing home residents with dementia. *Res Nurs Health*. 1994 Jun;17(3):159-65. PMID: 8184127.
178. Huis In Het Veld JG, Willemsse BM, van Asch IF, et al. Online Self-Management Support for Family Caregivers Dealing With Behavior Changes in Relatives With Dementia (Part 2): Randomized Controlled Trial. *J Med Internet Res*. 2020 Feb 25;22(2):e13001. doi: <https://dx.doi.org/10.2196/13001>. PMID: 32130142.
179. Van Mierlo LD, Meiland FJ, Van de Ven PM, et al. Evaluation of DEM-DISC, customized e-advice on health and social support services for informal carers and case managers of people with dementia; a cluster randomized trial. *Int Psychogeriatr*. 2015 Aug;27(8):1365-78. doi: <https://dx.doi.org/10.1017/S1041610215000423>. PMID: 25872457.
180. Wang LQ, Chien WT, Lee IY. An experimental study on the effectiveness of a mutual support group for family caregivers of a relative with dementia in mainland China. *Contemp Nurse*. 2012 Feb;40(2):210-24. doi: <https://dx.doi.org/10.5172/conu.2012.40.2.210>. PMID: 22554214.
181. Logsdon RG, Pike KC, McCurry SM, et al. Early-stage memory loss support groups: outcomes from a randomized controlled clinical trial. *Journals of Gerontology Series B: Psychological Sciences & Social Sciences*. 2010;65B(6):691-7. doi: geronb/gbq054. PMID: 104963362. Language: English. Entry Date: 20110128. Revision Date: 20150820. Publication Type: Journal Article.
182. Charlesworth G, Shepstone L, Wilson E, et al. Does befriending by trained lay workers improve psychological well-being and quality of life for carers of people with dementia, and at what cost? A randomised controlled trial. *Health Technol Assess*. 2008 Mar;12(4):iii, v-ix, 1-78. PMID: 18284895.
183. Charlesworth G, Shepstone L, Wilson E, et al. Befriending carers of people with dementia: randomised controlled trial. *Bmj*. 2008 Jun 07;336(7656):1295-7. doi: <https://dx.doi.org/10.1136/bmj.39549.548831.AE>. PMID: 18505757.
184. Wilson E, Thalanany M, Shepstone L, et al. Befriending carers of people with dementia: a cost utility analysis. *Int J Geriatr Psychiatry*. 2009 Jun;24(6):610-23. doi: <https://dx.doi.org/10.1002/gps.2164>. PMID: 19101921.
185. Winter L, Gitlin LN. Evaluation of a telephone-based support group intervention for female caregivers of community-dwelling individuals with dementia. *Am J Alzheimers Dis Other Demen*. 2006 Dec-2007 Jan;21(6):391-7. PMID: 17267370.
186. Mahoney DF, Tarlow BJ, Jones RN. Effects of an automated telephone support system on caregiver burden and anxiety: findings from the REACH for TLC intervention study. *Gerontologist*. 2003 Aug;43(4):556-67. PMID: 12937334.
187. Mahoney DM, Tarlow B, Jones RN, et al. Factors affecting the use of a telephone-based intervention for caregivers of people with Alzheimer's disease. *J Telemed Telecare*. 2001;7(3):139-48. PMID: 11346473.
188. Pillemer K, Suiitor JJ. Peer support for Alzheimer's caregivers: Is it enough to make a difference? *Research on Aging*. 2002;24(2):171-92. PMID: 34163761.
189. Goodman C. Evaluation of a model self-help telephone program: impact on natural networks. *Soc Work*. 1990 Nov;35(6):556-62. PMID: 2284602.
190. Gustafson DH, Cody OJ, Chih MY, et al. Pilot Test of a Computer-Based System to Help Family Caregivers of Dementia Patients. *J Alzheimers Dis*. 2019;70(2):539-50. doi: <http://dx.doi.org/10.3233/JAD-190052>. PMID: 628746285.
191. Metcalfe A, Jones B, Mayer J, et al. Online information and support for carers of people with young-onset dementia: A multi-site randomised controlled pilot study. *Int J Geriatr Psychiatry*. 2019 01 Oct;34(10):1455-64. doi: <http://dx.doi.org/10.1002/gps.5154>. PMID: 2002845656.
192. Torkamani M, McDonald L, Saez Aguayo I, et al. A randomized controlled pilot study to evaluate a technology platform for the assisted living of people with dementia and their carers. *J Alzheimers Dis*. 2014;41(2):515-23. doi: <https://dx.doi.org/10.3233/JAD-132156>. PMID: 24643137.
193. van der Roest HG, Meiland FJ, Jonker C, et al. User evaluation of the DEMentia-specific Digital Interactive Social Chart (DEM-DISC). A pilot study among informal carers on its impact, user friendliness and usefulness. *Aging Ment Health*. 2010 May;14(4):461-70. doi: <https://dx.doi.org/10.1080/13607860903311741>. PMID: 20455122.
194. Salamizadeh AMS, Mirzaei TP, Ravari AP. The Impact of Spiritual Care Education on the Self-Efficacy of the Family Caregivers of Elderly People with Alzheimer's Disease. *International Journal of Community Based Nursing & Midwifery*. 2017 Jul;5(3):231-8. PMID: 28670585.
195. Farran CJ, Etkin CD, Eisenstein A, et al. Effect of Moderate to Vigorous Physical Activity Intervention on Improving Dementia Family Caregiver Physical Function: A Randomized Controlled Trial. *J Alzheimers Dis Parkinsonism*. 2016 Aug;6(4). doi: <https://dx.doi.org/10.4172/2161-0460.1000253>. PMID: 28752016.
196. Farran CJ, Paun O, Cothran F, et al. Impact of an Individualized Physical Activity Intervention on Improving Mental Health Outcomes in Family Caregivers of Persons with Dementia: A Randomized Controlled Trial. *AIMS med*. 2016;3(1):15-31. doi: <https://dx.doi.org/10.3934/medsci.2016.1.15>. PMID: 29147683.
197. Cothran FA, Paun O, Barnes LL, et al. Comparing the Effect of a Moderate Physical Activity Intervention on the Mental Health Outcomes of African American and Caucasian Dementia Family Caregivers: A Secondary Data Analysis. *Issues Ment Health Nurs*. 2017 Dec;38(12):996-1004. doi: <https://dx.doi.org/10.1080/01612840.2017.1364807>. PMID: 28956706.

198. Mahdavi B, Fallahi-Khoshknab M, Mohammadi F, et al. Effects of spiritual group therapy on caregiver strain in home caregivers of the elderly with Alzheimer's disease. *Arch Psychiatr Nurs*. 2016. PMID: CN-01308784.
199. Lowery D, Cerga-Pashoja A, Iliffe S, et al. The effect of exercise on behavioural and psychological symptoms of dementia: the EVIDEM-E randomised controlled clinical trial. *Int J Geriatr Psychiatry*. 2014 Aug;29(8):819-27. doi: <https://dx.doi.org/10.1002/gps.4062>. PMID: 24338799.
200. D'Amico F, Rehill A, Knapp M, et al. Cost-effectiveness of exercise as a therapy for behavioural and psychological symptoms of dementia within the EVIDEM-E randomised controlled trial. *Int J Geriatr Psychiatry*. 2016 Jun;31(6):656-65. doi: <https://dx.doi.org/10.1002/gps.4376>. PMID: 26489776.
201. Whitebird RR, Kreitzer M, Crain AL, et al. Mindfulness-based stress reduction for family caregivers: a randomized controlled trial. *Gerontologist*. 2013 Aug;53(4):676-86. doi: <https://dx.doi.org/10.1093/geront/gns126>. PMID: 23070934.
202. Connell CM, Janevic MR. Effects of a Telephone-Based Exercise Intervention for Dementia Caregiving Wives: A Randomized Controlled Trial. *J Appl Gerontol*. 2009 Apr;28(2):171-94. PMID: 21709757.
203. Castro CM, Wilcox S, O'Sullivan P, et al. An exercise program for women who are caring for relatives with dementia. *Psychosom Med*. 2002 May-Jun;64(3):458-68. PMID: 12021419.
204. Kor PPK, Liu JYW, Chien WT. Effects of a modified mindfulness-based cognitive therapy for family caregivers of people with dementia: A pilot randomized controlled trial. *Int J Nurs Stud*. 2019 Oct;98:107-17. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2019.02.020>. PMID: 30922609.
205. Richards AG, Tietyen AC, Jicha GA, et al. Visual Arts Education improves self-esteem for persons with dementia and reduces caregiver burden: A randomized controlled trial. *Dementia*. 2018 Jan 01;1471301218769071. doi: <https://dx.doi.org/10.1177/1471301218769071>. PMID: 29653492.
206. Waelde LC, Meyer H, Thompson JM, et al. Randomized Controlled Trial of Inner Resources Meditation for Family Dementia Caregivers. *J Clin Psychol*. 2017 Dec;73(12):1629-41. doi: <https://dx.doi.org/10.1002/jclp.22470>. PMID: 28263398.
207. Brown KW, Coogler CL, Wegelin J. A pilot randomized controlled trial of mindfulness-based stress reduction for caregivers of family members with dementia. *Aging Ment Health*. 2016 Nov;20(11):1157-66. PMID: 26211415.
208. Hirano A, Umegaki H, Suzuki Y, et al. Effects of leisure activities at home on perceived care burden and the endocrine system of caregivers of dementia patients: a randomized controlled study. *Int Psychogeriatr*. 2016 Feb;28(2):261-8. doi: <https://dx.doi.org/10.1017/S1041610215001295>. PMID: 26289496.
209. Leach MJ, Francis A, Ziaian T. Transcendental Meditation for the improvement of health and wellbeing in community-dwelling dementia caregivers [TRANSCENDENT]: a randomised wait-list controlled trial. *BMC Altern Med*. 2015 May 08;15:145. doi: <https://dx.doi.org/10.1186/s12906-015-0666-8>. PMID: 25952550.
210. Leach MJ, Francis A, Ziaian T. Improving the health and well-being of community-dwelling caregivers of dementia sufferers: study protocol of a randomized controlled trial of structured meditation training. *J Altern Complement Med*. 2014 Feb;20(2):136-41. doi: <https://dx.doi.org/10.1089/acm.2013.0170>. PMID: 24044373.
211. Leach MJ, Ziaian T, Francis A, et al. Recruiting Dementia Caregivers Into Clinical Trials: Lessons Learnt From the Australian TRANSCENDENT Trial. *Alzheimer Dis Assoc Disord*. 2016 Oct-Dec;30(4):338-44. PMID: 27227995.
212. Lavretsky H, Epel ES, Siddarth P, et al. A pilot study of yogic meditation for family dementia caregivers with depressive symptoms: effects on mental health, cognition, and telomerase activity. *Int J Geriatr Psychiatry*. 2013 Jan;28(1):57-65. doi: <https://dx.doi.org/10.1002/gps.3790>. PMID: 22407663.
213. Carbonneau H, Caron CD, Desrosiers J. Effects of an adapted leisure education program as a means of support for caregivers of people with dementia. *Arch Gerontol Geriatr*. 2011 Jul-Aug;53(1):31-9. doi: <https://dx.doi.org/10.1016/j.archger.2010.06.009>. PMID: 20598758.
214. Oken BS, Fonareva I, Haas M, et al. Pilot controlled trial of mindfulness meditation and education for dementia caregivers. *J Altern Complement Med*. 2010 Oct;16(10):1031-8. doi: <https://dx.doi.org/10.1089/acm.2009.0733>. PMID: 20929380.
215. Wilz G, Fink-Heitz M. Assisted vacations for men with dementia and their caregiving spouses: evaluation of health-related effects. *Gerontologist*. 2008 Feb;48(1):115-20. PMID: 18381838.
216. Vandepitte S, Putman K, Van Den Noortgate N, et al. Effectiveness of an in-home respite care program to support informal dementia caregivers: A comparative study. *Int J Geriatr Psychiatry*. 2019 Oct;34(10):1534-44. doi: <https://dx.doi.org/10.1002/gps.5164>. PMID: 31243801.
217. Vandepitte S, Van Den Noortgate N, Putman K, et al. Effectiveness and cost-effectiveness of an in-home respite care program in supporting informal caregivers of people with dementia: design of a comparative study. *BMC geriatr*. 2016 12 02;16(1):207. PMID: 27912740.
218. Zarit SH, Stephens MA, Townsend A, et al. Stress reduction for family caregivers: effects of adult day care use. *J Gerontol B Psychol Sci Soc Sci*. 1998 Sep;53(5):S267-77. PMID: 9750575.
219. Kim K, Zarit SH, Femia EE, et al. Kin relationship of caregivers and people with dementia: stress and response to intervention. *Int J Geriatr Psychiatry*. 2012 Jan;27(1):59-66. doi: <https://dx.doi.org/10.1002/gps.2689>. PMID: 21322030.
220. Lawton MP, Brody EM, Saperstein A, et al. Respite services for caregivers: research findings for service planning. *Home Health Care Serv Q*. 1989;10(1-2):5-32. PMID: 10318337.
221. Luchsinger JA, Burgio L, Mittelman M, et al. Comparative Effectiveness of 2 Interventions for Hispanic Caregivers of Persons with Dementia. *J Am Geriatr Soc*. 2018 Sep;66(9):1708-15. doi: <https://dx.doi.org/10.1111/jgs.15450>. PMID: 30084133.
222. Berwig M, Heinrich S, Spahlholz J, et al. Individualized support for informal caregivers of people with dementia - effectiveness of the German adaptation of REACH II. *BMC geriatr*. 2017 Dec 12;17(1):286. doi: <https://dx.doi.org/10.1186/s12877-017-0678-y>. PMID: 29233097.
223. Brijoux T, Kricheldorf C, Hull M, et al. Supporting families living with dementia in rural areas: a randomized controlled trial of quality of life improvement using qualified volunteers. In *Dtsch*
224. Koivisto AM, Hallikainen I, Valimaki T, et al. Early psychosocial intervention does not delay institutionalization in persons with mild Alzheimer disease and has impact on neither disease progression nor caregivers' well-being: ALSOVA 3-year follow-up. *Int J Geriatr Psychiatry*. 2016 Mar;31(3):273-83. doi: <https://dx.doi.org/10.1002/gps.4321>. PMID: 26177825.
225. Baglio F, Griffanti L, Saibene FL, et al. Multistimulation group therapy in Alzheimer's disease promotes changes in brain functioning. *Neurorehabil Neural Repair*. 2015 Jan;29(1):13-24. doi: <https://dx.doi.org/10.1177/1545968314532833>. PMID: 24788581.
226. Prick AE, de Lange J, Twisk J, et al. The effects of a multi-component dyadic intervention on the psychological distress of family caregivers providing care to people with dementia: a randomized controlled trial. *Int Psychogeriatr*. 2015 Dec;27(12):2031-44. doi: <https://dx.doi.org/10.1017/S104161021500071X>. PMID: 26004290.
227. Gaugler JE, Reese M, Mittelman MS. Effects of the NYU caregiver intervention-adult child on residential care placement. *Gerontologist*. 2013 Dec;53(6):985-97. doi: <https://dx.doi.org/10.1093/geront/gns193>. PMID: 23339050.
228. Martindale-Adams J, Nichols LO, Burns R, et al. A trial of dementia caregiver telephone support. *Can J Nurs Res*. 2013 Dec;45(4):30-48. PMID: 24617278.
229. Mittelman MS, Brodaty H, Wallen AS, et al. A three-country randomized controlled trial of a psychosocial intervention for caregivers combined with pharmacological treatment for patients with Alzheimer disease: effects on caregiver depression. *Am J Geriatr Psychiatry*. 2008;16(11):893-904. doi: 10.1097/JGP.0b013e3181898095. PMID: 105627901. Language: English. Entry Date: 20090130. Revision Date: 20161207. Publication Type: journal article.
230. Belle SH, Burgio L, Burns R, et al. Enhancing the quality of life of dementia caregivers from different ethnic or racial groups: a randomized, controlled trial.[Summary for patients in *Ann Intern Med*. 2006 Nov 21;145(10):I39; PMID: 17116914]. *Ann Intern Med*. 2006 Nov 21;145(10):727-38. PMID: 17116917.
231. Ducharme F, Levesque L, Giroux F, et al. Follow-up of an intervention program for caregivers of a relative with dementia living in a long-term care setting: Are there any persistent and delayed effects? *Aging Ment Health*. 2005 Sep;9(5):461-9. PMID: 16024406.
232. Mittelman MS, Roth DL, Haley WE, et al. Effects of a caregiver intervention on negative caregiver appraisals of behavior problems in patients with Alzheimer's disease: results of a randomized trial. *J Gerontol B Psychol Sci Soc Sci*. 2004 Jan;59(1):P27-34. PMID: 14722336.
233. Eisdorfer C, Czaja SJ, Loewenstein DA, et al. The effect of a family therapy and technology-based intervention on caregiver depression. *Gerontologist*. 2003 Aug;43(4):521-31. PMID: 12937331.
234. Teri L, Gibbons LE, McCurry SM, et al. Exercise plus behavioral management in patients with Alzheimer disease: a randomized controlled trial. *Jama*. 2003 Oct 15;290(15):2015-22. PMID: 14559955.
235. Mittelman MS, Ferris SH, Steinberg G, et al. An intervention that delays institutionalization of Alzheimer's disease patients: treatment of spouse-caregivers. *Gerontologist*. 1993 Dec;33(6):730-40. PMID: 8314099.
236. Mohide EA, Pringle DM, Streiner DL, et al. A randomized trial of family caregiver support in the home management of dementia. *J Am Geriatr Soc*. 1990 Apr;38(4):446-54. PMID: 2184186.
237. Prick AE, de Lange J, Scherder E, et al. The Effects of a Multicomponent Dyadic Intervention With Physical Exercise on the Cognitive Functioning of People With Dementia: A Randomized Controlled Trial. *J Aging Phys Activity*. 2017 Oct 01;25(4):539-52. doi: <https://dx.doi.org/10.1123/japa.2016-0038>. PMID: 28120631.
238. Prick AE, de Lange J, Scherder E, et al. The effects of a multicomponent dyadic intervention on the mood, behavior, and physical health of people with dementia: a randomized controlled trial. *Clin Interv Aging*. 2016;11:383-95. doi: <https://dx.doi.org/10.2147/CIA.S95789>. PMID: 27099480.
239. Prick AE, de Lange J, van 't Leven N, et al. Process evaluation of a multicomponent dyadic intervention study with exercise and support for people with dementia and their family caregivers. *Trials*. 2014 Oct 22;15:401. doi: <https://dx.doi.org/10.1186/1745-6215-15-401>. PMID: 25336121.
240. Elliott AF, Burgio LD, Decoster J. Enhancing caregiver health: findings from the resources for enhancing Alzheimer's caregiver health II intervention. *J Am Geriatr Soc*. 2010 Jan;58(1):30-7. doi: <https://dx.doi.org/10.1111/j.1532-5415.2009.02631.x>. PMID: 20122038.
241. Nichols LO, Martindale-Adams J, Zhu CW, et al. Impact of the REACH II and REACH VA Dementia Caregiver Interventions on Healthcare Costs. *J Am Geriatr Soc*. 2017 May;65(5):931-6. doi: <https://dx.doi.org/10.1111/jgs.14716>. PMID: 28295134.
242. Nichols LO, Martindale-Adams J, Burns R, et al. Translation of a dementia caregiver support program in a health care system--REACH VA. *Arch Intern Med*. 2011 Feb 28;171(4):353-9. doi: <https://dx.doi.org/10.1001/archinternmed.2010.548>. PMID: 21357811.
243. Nichols LO, Chang C, Lummus A, et al. The cost-effectiveness of a behavior intervention with caregivers of patients with Alzheimer's disease. *J Am Geriatr Soc*. 2008;56(3):413-20. doi: 10.1111/j.1532-5415.2007.01569.x. PMID: 105900573. Corporate Author: Resources for Enhancing Alzheimer's Caregivers Health II Investigators. Language: English. Entry Date: 20080425. Revision Date: 20150711. Publication Type: Journal Article.
244. Basu R, Hochhalter AK, Stevens AB. The Impact of the REACH II Intervention on Caregivers' Perceived Health. *J Appl Gerontol*. 2015 Aug;34(5):590-608. doi: <https://dx.doi.org/10.1177/0733464813499640>. PMID: 24652899.
245. Lee CC, Czaja SJ, Schulz R. The moderating influence of demographic characteristics, social support, and religious coping on the effectiveness of a multicomponent psychosocial caregiver intervention in three racial ethnic groups. *Journals of Gerontology Series B: Psychological Sciences & Social Sciences*. 2010;65B(2):185-94. doi: geronb/gbp131. PMID: 105127911. Language: English. Entry Date: 20100430. Revision Date: 20150711. Publication Type: Journal Article.
246. Graham-Phillips A, Roth DL, Huang J, et al. Racial and Ethnic Differences in the Delivery of the Resources for Enhancing Alzheimer's Caregiver Health II Intervention. *J Am Geriatr Soc*. 2016 Aug;64(8):1662-7. doi: <https://dx.doi.org/10.1111/jgs.14204>. PMID: 27294873.
247. Hatch DJ, DeHart WB, Norton MC. Subjective stressors moderate effectiveness of a multi-component, multi-site intervention on caregiver depression and burden. *Int J Geriatr Psychiatry*. 2014 Apr;29(4):406-13. doi: <https://dx.doi.org/10.1002/gps.4019>. PMID: 23983230.
248. McGinnis KA, Schulz R, Stone RA, et al. Concordance of race or ethnicity of interventionists and caregivers of dementia patients: relationship to attrition and treatment outcomes in the REACH study. *Gerontologist*. 2006 Aug;46(4):449-55. PMID: 16920998.

249. Wisniewski SR, Belle SH, Coon DW, et al. The Resources for Enhancing Alzheimer's Caregiver Health (REACH): project design and baseline characteristics. *Psychol Aging*. 2003 Sep;18(3):375-84. PMID: 14518801.
250. Mittelman MS, Roth DL, Coon DW, et al. Sustained benefit of supportive intervention for depressive symptoms in caregivers of patients with Alzheimer's disease. *Am J Psychiatry*. 2004 May;161(5):850-6. PMID: 15121650.
251. Mittelman MS, Haley WE, Clay OJ, et al. Improving caregiver well-being delays nursing home placement of patients with Alzheimer disease. *Neurology*. 2006 Nov 14;67(9):1592-9. PMID: 17101889.
252. Mittelman MS, Roth DL, Clay OJ, et al. Preserving health of Alzheimer caregivers: impact of a spouse caregiver intervention. *Am J Geriatr Psychiatry*. 2007 Sep;15(9):780-9. PMID: 17804831.
253. Drentea P, Clay OJ, Roth DL, et al. Predictors of improvement in social support: Five-year effects of a structured intervention for caregivers of spouses with Alzheimer's disease. *Soc Sci Med*. 2006 Aug;63(4):957-67. PMID: 16616406.
254. Haley WE, Bergman EJ, Roth DL, et al. Long-term effects of bereavement and caregiver intervention on dementia caregiver depressive symptoms. *Gerontologist*. 2008 Dec;48(6):732-40. PMID: 19139247.
255. Roth DL, Mittelman MS, Clay OJ, et al. Changes in social support as mediators of the impact of a psychosocial intervention for spouse caregivers of persons with Alzheimer's disease. *Psychol Aging*. 2005 Dec;20(4):634-44. PMID: 16420138.
256. Gaugler JE, Roth DL, Haley WE, et al. Can counseling and support reduce burden and depressive symptoms in caregivers of people with Alzheimer's disease during the transition to institutionalization? Results from the New York University caregiver intervention study. *J Am Geriatr Soc*. 2008 Mar;56(3):421-8. doi: <https://dx.doi.org/10.1111/j.1532-5415.2007.01593.x>. PMID: 18179495.
257. Gaugler JE, Roth DL, Haley WE, et al. Modeling trajectories and transitions: results from the New York University caregiver intervention. *Nurs Res*. 2011 May-Jun;60(3 Suppl):S28-37. doi: <https://dx.doi.org/10.1097/NNR.0b013e318216007d>. PMID: 21543959.
258. Jang Y, Clay OJ, Roth DL, et al. Neuroticism and longitudinal change in caregiver depression: impact of a spouse-caregiver intervention program. *Gerontologist*. 2004 Jun;44(3):311-7. PMID: 15197285.
259. Burns A, Mittelman M, Cole C, et al. Transcultural influences in dementia care: observations from a psychosocial intervention study. *Dement Geriatr Cogn Disord*. 2010;30(5):417-23. doi: <https://dx.doi.org/10.1159/000314860>. PMID: 21071943.
260. Brodaty H, Mittelman M, Gibson L, et al. The effects of counseling spouse caregivers of people with Alzheimer disease taking donepezil and of country of residence on rates of admission to nursing homes and mortality. *Am J Geriatr Psychiatry*. 2009 Sep;17(9):734-43. PMID: 19705519.
261. Mazurek J, Szczesniak D, Lion KM, et al. Does the Meeting Centres Support Programme reduce unmet care needs of community-dwelling older people with dementia? A controlled, 6-month follow-up Polish study. *Clin Interv Aging*. 2019;14:113-22. doi: <https://dx.doi.org/10.2147/CIA.S185683>. PMID: 30666097.
262. Zhang SY, Wu F, Tang DL, et al. Pilot testing the caregiver self-management intervention for caregivers of relatives with dementia. *Geriatr Nurs*. 2019 Aug 26;26:26. doi: <https://dx.doi.org/10.1016/j.gerinurse.2019.08.006>. PMID: 31466807.
263. Marshall A, Spreadbury J, Cheston R, et al. A pilot randomised controlled trial to compare changes in quality of life for participants with early diagnosis dementia who attend a "Living Well with Dementia" group compared to waiting-list control. *Aging Ment Health*. 2015;19(6):526-35. PMID: CN-01071410.
264. Czaja SJ, Loewenstein D, Schulz R, et al. A videophone psychosocial intervention for dementia caregivers. *Am J Geriatr Psychiatry*. 2013 Nov;21(11):1071-81. doi: <https://dx.doi.org/10.1016/j.jagp.2013.02.019>. PMID: 23831174.
265. Gaugler JE, Reese M, Mittelman MS. Effects of the Minnesota Adaptation of the NYU Caregiver Intervention on Depressive Symptoms and Quality of Life for Adult Child Caregivers of Persons with Dementia. *Am J Geriatr Psychiatry*. 2015 Nov;23(11):1179-92. doi: <https://dx.doi.org/10.1016/j.jagp.2015.06.007>. PMID: 26238226.
266. Gaugler JE, Reese M, Mittelman MS. Effects of the Minnesota Adaptation of the NYU Caregiver Intervention on Primary Subjective Stress of Adult Child Caregivers of Persons With Dementia. *Gerontologist*. 2016 Jun;56(3):461-74. doi: <https://dx.doi.org/10.1093/geront/gnu125>. PMID: 25628299.
267. Gaugler JE, Reese M, Mittelman MS. Process Evaluation of the NYU Caregiver Intervention-Adult Child. *Gerontologist*. 2018 03 19;58(2):e107-e117. doi: <https://dx.doi.org/10.1093/geront/gnx048>. PMID: 29562359.
268. Gaugler JE, Reese M, Mittelman MS. The Effects of a Comprehensive Psychosocial Intervention on Secondary Stressors and Social Support for Adult Child Caregivers of Persons With Dementia. *Innov Aging*. 2018 Jun;2(2):igy015. doi: <https://dx.doi.org/10.1093/geroni/igy015>. PMID: 30009268.
269. Finkel S, Czaja SJ, Schulz R, et al. E-care: a telecommunications technology intervention for family caregivers of dementia patients. *Am J Geriatr Psychiatry*. 2007 May;15(5):443-8. PMID: 17463195.
270. Jutten LH, Mark RE, Sitskoorn MM. Can the Mixed Virtual Reality Simulator Into D'mentia Enhance Empathy and Understanding and Decrease Burden in Informal Dementia Caregivers? *Dement Geriatr Cogn Dis Extra*. 2018 Sep-Dec;8(3):453-66. doi: <https://dx.doi.org/10.1159/000494660>. PMID: 30631337.
271. Jutten LH, Mark RE, Maria Janssen BWJ, et al. Testing the effectivity of the mixed virtual reality training Into D'mentia for informal caregivers of people with dementia: protocol for a longitudinal, quasi-experimental study. *BMJ Open*. 2017 Aug 21;7(8):e015702. doi: <https://dx.doi.org/10.1136/bmjopen-2016-015702>. PMID: 28827242.
272. Stern RA, D'Ambrosio LA, Mohyde M, et al. At the crossroads: development and evaluation of a dementia caregiver group intervention to assist in driving cessation. *Gerontol Geriatr Educ*. 2008;29(4):363-82. doi: <https://dx.doi.org/10.1080/02701960802497936>. PMID: 19064472.
273. Simpson R, Wakefield P, Spiers N, et al. Carer-held records for dementia: a controlled trial. *Int Psychogeriatr*. 2006 Jun;18(2):259-68. PMID: 16734920.
274. Lord K, Livingston G, Cooper C. A feasibility randomised controlled trial of the DECIDE intervention: dementia carers making informed decisions. *Bjpsych open*. 2017;3(1):12-4. PMID: CN-01415827.
275. Stirling C, Leggett S, Lloyd B, et al. Decision aids for respite service choices by carers of people with dementia: development and pilot RCT. *BMC Med Inf Decis Mak*. 2012 Mar 19;12:21. doi: <https://dx.doi.org/10.1186/1472-6947-12-21>. PMID: 22429384.
276. Rose KM, Taylor AG, Bourguignon C. Effects of cranial electrical stimulation on sleep disturbances, depressive symptoms, and caregiving appraisal in spousal caregivers of persons with Alzheimer's disease. *Appl Nurs Res*. 2009 May;22(2):119-25. doi: <https://dx.doi.org/10.1016/j.apnr.2007.06.001>. PMID: 19427574.
277. Barbosa A, Nolan M, Sousa L, et al. Supporting direct care workers in dementia care: effects of a psychoeducational intervention. *Am J Alzheimers Dis Other Demen*. 2015 Mar;30(2):130-8. doi: <https://dx.doi.org/10.1177/1533317514550331>. PMID: 25237132.
278. Barbosa A, Nolan M, Sousa L, et al. Effects of a Psychoeducational Intervention for Direct Care Workers Caring for People With Dementia: Results From a 6-Month Follow-Up Study. *Am J Alzheimers Dis Other Demen*. 2016 Mar;31(2):144-55. doi: <https://dx.doi.org/10.1177/1533317515603500>. PMID: 26399485.
279. Visser SM, McCabe MP, Hudgson C, et al. Managing behavioural symptoms of dementia: effectiveness of staff education and peer support. *Aging Ment Health*. 2008 Jan;12(1):47-55. doi: <https://dx.doi.org/10.1080/13607860701366012>. PMID: 18297478.
280. Davison T, McCabe M, Visser S, et al. Controlled trial of dementia training with a peer support group for aged care staff. *Int J Geriatr Psychiatry*. 2007;22(9):868-73. doi: 10.1002/gps.1754. PMID: CN-00627010.
281. Mackenzie CS, Peragine G. Measuring and enhancing self-efficacy among professional caregivers of individuals with dementia. *Am J Alzheimers Dis Other Demen*. 2003 Sep-Oct;18(5):291-9. PMID: 14569646.

Appendix F. Care Delivery Interventions

Consultation Service

Table F-1. Risk of bias assessment: consultation services

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Overall Rating
Fortinsky 2009 ¹ (19347683)	12 months	Medium 12 months: 10.7%	Medium	Medium	Low	Medium	Medium	Medium
Borbasi 2011 ² (21692400)	12 months	X	High	X	X	X	X	High
McSweeney 2012 ³ (22344753)	15 weeks	Medium 11%	Low	Low	High	Low	High	High
Opie 2002 ⁴ (11802224)	4 weeks	Low	Medium 3%	High	High	Low	Medium	High
Orrell 2007 ⁵ (17394129)	20 weeks	Medium 19%	Medium	Low	Medium	Low	High	High

Abbreviations: PMID=PubMed Identification Number

Table F-2. Characteristics of included studies: consultation services

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Fortinsky 2009 ¹ (19347683) US Medium Explanatory	Intervention group family caregivers received the dementia care consultation intervention per protocol as well as educational materials about dementia and community resources over 12 months period	Educational materials	Community-based in-home Cluster RCT 26 practice sites	Diagnosed with AD or other dementia. With following ICD-9 codes: arteriosclerotic dementia (290.40,290.41,290.42,290.43); senile dementia (290.00,290.20,290.21,290.30); pre-senile dementia (290.10,290.11,290.12,290.13); memory loss, mild (310.10); or AD (331.00).	N=84 66% Female 82 years Race NR Education NR	None	NA	N=84 72% Female 61 years Race NR Education NR	NA	NA	12 months	Nursing home admission;	IC: 2 measures of self-efficacy Caregiver Burden CES-D HSC measures of symptom severity CSC PBS

*High risk of bias studies included in evidence map

Abbreviations: AD=Alzheimer's disease; char=characteristics; CES-D=Center for Epidemiological Studies Depression Inventory; FC=formal caregiver; CSC=Cognitive Status Scale; IC=informal caregiver; ICD=International Classification of Diseases; HSC=Hopkins Symptoms Checklist; PBS=Problematic Behavior Scale; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RCT=Randomized Controlled Trial; RoB=Risk of Bias; RS=Reporting Status

Table F-3. PLWD outcomes summary low and medium risk of bias studies: consultation services

Study (PMID) Comparison RoB Category	Outcome Timing					p-Value
		Intervention: Before	Intervention: After 12 months	Comparator: Before	Comparator: After 12 months	
Fortinsky 2009 ¹ (19347683) Consultation vs Control Medium Exploratory	Adjusted Odds Ratio (95% CI) Nursing home admission 12 months	NR	0.4 (0.14, 1.18)	NR	1.0	p=0.10

Abbreviations: CES-D=Center for Epidemiological Studies Depression Inventory; CI=Confidence Interval; NR=Not Reported; PMID=PubMed Identification Number; RoB=Risk of Bias

Table F-4. Informal caregiver outcomes summary low and medium risk of bias studies: consultation services

Study (PMID) Comparison RoB Category	Outcome Timing	Intervention		Comparator		p-Value
		Before	After 12 months	Before	After 12 months	
Fortinsky 2009 ¹ (19347683) Consultation vs Control Medium Exploratory	Adjusted Mean (95% CI) Symptom management self-efficacy score 12 months	33.52 (27.77–39.26)	33.88 (28.04–39.73)	33.51 (27.31–39.71)	34.21 (27.92–40.49)	p=0.89
Fortinsky 2009 ¹ (19347683) Consultation vs Control Medium Exploratory	Adjusted Mean (95% CI) Support service self-efficacy 12 months	27.96 (25.17–30.75)	31.95 (28.90–35.00)	24.63 (20.95–28.31)	27.92 (24.17–31.66)	P=0.80
Fortinsky 2009 ¹ (19347683) Consultation vs Control Medium Exploratory	Adjusted Mean (95% CI) CES-D score 12 months	12.13 (8.90–15.35)	9.80 (6.24–13.36)	15.10 (10.77–19.43)	15.00 (10.51–19.49)	p=0.41
Fortinsky 2009 ¹ (19347683) Consultation vs Control Medium Exploratory	Adjusted Mean (95% CI) Zarit burden score 12 months	30.42 (26.30–34.53)	26.18 (21.81–30.55)	36.02 (30.71–41.33)	30.57 (25.03–36.10)	p=0.73
Fortinsky 2009 ¹ (19347683) Consultation vs Control Medium Exploratory	Adjusted Mean (95% CI) Hopkins symptom checklist score 12 months	2.33 (1.57–3.09)	2.92 (2.10–3.74)	3.23 (2.31–4.17)	3.74 (2.78–4.70)	p=0.87

Abbreviations: CES-D=Center for Epidemiological Studies Depression Inventory; CI=Confidence Interval; NR=Not Reported; PMID=PubMed Identification Number; RoB=Risk of Bias

Table F-5. Summary of strength of evidence for PLWD outcomes: consultation services

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Consultation vs Control Nursing home admission	12 months	1 RCT (n=84)	Nursing home admissions showed no difference between the comparison groups.	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: CES-D=Center for Epidemiological Studies Depression Inventory; n=Number; RCT=Randomized Controlled Trial

Table F-6. Summary of strength of evidence for caregiver outcomes: consultation services

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Consultation vs Control Symptom management self-efficacy score	12 months	1 RCT (n=84)	Symptom management self-efficacy scores were not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control Support service self-efficacy	12 months	1 RCT (n=84)	Support service self-efficacy was not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control CES-D score	12 months	1 RCT (n=84)	CES-D scores were not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control Zarit burden score	12 months	1 RCT (n=84)	Zarit burden scores were not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control Hopkins symptom checklist score	12 months	1 RCT (n=84)	Hopkins symptom checklist scores were not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: CES-D=Center for Epidemiological Studies Depression Inventory; n=Number; RCT=Randomized Controlled Trial

Evidence Map: Consultation Services

Table F-7. Characteristics of evidence map studies: consultation services

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Borbasi 2011 ² (21692400) Australia High RoB	Dementia Outreach Service (DEMOS): comprised a multidisciplinary healthcare team to assess and manage residents experiencing cognitive deterioration as a result of dementia. Conducted for 12 months in 3 stages	NR	Residential aged care facilities Quasi-experimental	Not specified	NR	None	NR	None	N=320 Age: no Sex: yes Race: yes Education: no	NA	12 months	NR	Improved capacity and clinical skills of staff (stress, knowledge and self-confidence) Improvement in timeliness and appropriateness of referrals Satisfaction Barriers to success
McSweeney 2012 ³ (22344753) Australia High RoB	Psychiatrist and psychologist's consultation regarding best-practice management of depression via psychosocial (individually tailored, psychosocial care plan) and medical care plan	Care as usual	Aged care facilities RCT; 44 PLWD	MMSE cut point score of 23 and PAS Informant History≥4 PAS, Cognitive Decline Informant History and CSDD	N=44 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	15 weeks	CSDD RAID MMSE BEHAVE-AD CANE	NA
Opie 2002 ⁴ (11802224) Australia High RoB	Multidisciplinary interventions (early and late groups) encompassing psychosocial strategies, nursing approaches, psychotropic approaches and management of pains	NA	Nursing homes Quasi-experimental	Based on CMAI, BAGS and MMSE scores	N=99 Age: yes Sex: no Race: no Education: no	None	NA	NA	NA	NA	Weekly up to 4 weeks	Restlessness Physical aggression Verbal disruption Inappropriate behavior	NA
Orrell 2007 ⁵ (17394129) UK High RoB	1 hour per week liaison mediated personalized intervention package over 20 weeks to address the unmet needs	Care as usual	residential care homes Cluster RCT; 238 PLWD	Diagnosis based on DSM-IV criteria for dementia	N=238 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	20 weeks	CANE QoL-AD MMSE CDR CAPE-BRS BI CBS CSDD RAID	CANE QoL-AD

Abbreviations: BAGS=Behavior Assessment Graphical System; BEHAVE-AD=Behavioural Pathology in Alzheimer's Disease; BI=Barthel Index; CANE=Camberwell Assessment of Need for the Elderly; CAPE-BRS=Clifton Assessment Procedures for the Elderly-Behavior Rating Scale; CBS=Challenging Behavior Scale; CDR=Clinical Dementia Rating; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; DSM=Diagnostic and Statistical Manual; EM=Evidence Map; MMSE=Mini-mental State Examination; N=number; NA=Not Applicable; NR=Not Reported; PAS=Psychogeriatric Assessment Scale; PMID=PubMed Identification Number; QoL-AD=Quality of Life in Alzheimer's Disease; RAID=Rating for Anxiety in Dementia

Case Management

Table F-8. Risk of bias assessment: case management

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Xiao, 2016 ⁶ (25805891)	12 months	Medium 15%	Low	Medium	High	Low	Low	Government	High
MacNeil Vroomen, 2015 ⁷ (26170035)	24 months	High 40%	Medium	X	X	X	X		High
Chien, 2011 ⁸ (21198803)	12, 18 months	Low 2%	Low	Low	Medium	Low	High	Government University	Medium
Lam, 2010 ⁹ (19606455)	12 months	Low 3%	Low	Medium	Medium	Low	High	Government	Medium
Brodady, 2003 ¹⁰ (12590626)	12 weeks	Medium 16%	Low	Low	High	Low	High		High
Challis, 2002 ¹¹ (11994884)	6 months	Low 0%	High	High	X	X	X		High
Eloniemi-Sulkava, 2001 ¹² (11890485)	2 years	Low 0%	Medium	Low	Medium	Low	Medium	Government	Medium
Weinberger, 1993 ¹³ (8426038)	6 months	Medium 14%	Medium	High	High	Low	High		High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table F-9. Characteristics of included studies: case management

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Chien 2011 ⁸ (21198803) Hong Kong Medium Explanatory	Orientation to dementia care, educational workshop, family role and strength rebuilding, community support resources, program evaluation For 6 months	Routine care Usual services provided by the dementia resources center	Community RCT N=92 dyads	Nonspecified dementia DSM-IV	N=92 Age 68 44% Female Race NR Education NR	Mean monthly household income 14,000 Hong Kong dollars	NR Age 45 66% Female Race NR Education 27% Primary or less 64% Secondary 8% Tertiary	NR	NA	NA	12, 18 months	NPI Institutionalization	Family Caregiving Burden Inventory WHOQOL-BREF Social Support Questionnaire Family Support Services Index
Lam 2010 ⁹ (19606455) Hong Kong Medium Explanatory	Case Management model: Assessment and advice. Home-based advice and training on cognitive stimulation strategies, support sites & telephone calls For 4 months	One home visit for home safety at the beginning of the trial, no case management	Community-based RCT N=102	Nonspecified dementia Chinese Mini-Mental State Examination (CMMSE) 15+ Clinical Dementia Rating Scale 1	N=102 Age 78 58% Female Race NR Education NR	NR	N=102 Age NR 74% Female Race NR Education NR 29% Spouse or partner 51% Child	NR	NA	NA	12 months	CSDD NPI Personal Wellbeing Index-Intellectual Disability	ZBI General Health Questionnaire Personal Well-Being Index Social support use
Eloniemi-Sulkava 2001 ¹⁴ (11890485) Finland Medium Explanatory	Care coordination support program 2 years	Usual care activities	Home-based RCT N=100 dyads	Alzheimer's disease Vascular dementia DSM-III	N=100 Age 79 53% Female Race NR Education NR	NR	N=100 Age 64 69% Female Race NR Education NR 56% Spouse 35% Child	NR	NA	NA	2 years	Rate of institutionalization	General Health Questionnaire

Abbreviations: char=Characteristics; CMMSE=Mini State Mental Examination Chinese version; CSDD=Cornell Scale for Depression in Dementia; DSM-III=Diagnostic and Statistical Manual of Mental Disorders-Third Ed; DSM-IV=Diagnostic and Statistical Manual of Mental Disorders-Fourth Ed; FC=Formal Caregiver; IC=Informal Caregiver; N=Number; NA=Not Applicable; NPI=Neuropsychiatric Inventory; NR=Not Reported; PMID=PubMed Identification Number; RCT=Randomized Controlled Trial; RoB-Risk of Bias; RS=Reporting Status; SES=Socioeconomic Status; WHOQOL-BREF= World Health Organization Quality of Life Measure-Brief Version; ZBI=Zarit Burden Interview

Table F-10. PLWD outcomes summary for low and medium risk of bias studies: case management

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Chien, 2011 ⁸ (21198803) Case management vs usual care Medium Explanatory	NPI 18 months	Favors intervention F(1,90) = 3.70	Baseline 81.8 (SD 9.1) 18 month 76.5 (SD 8.9)	Baseline 80.9 (SD 9.5) 18 month 82.4 (SD 11.0)	<0.01
Chien, 2011 ⁸ (21198803) Case management vs usual care Medium Explanatory	Average number institutionalizations over previous 6 months 18 months	Favors intervention F(1,90) = 4.32	Baseline 5.1 (SD 0.9) 18 month 2.6 (SD 1.9)	Baseline 5.5 (SD 1.2) 18 month 6.0 (SD 2.9)	<0.01
Lam, 2010 ⁹ (19606455) Medium Explanatory	Personal well-being index for intellectually disabled (PWI-ID) difference in difference 12 month	No statistical difference	(reported data must be incorrect for NS finding)	Change from baseline 1.4 (-7.1, 10.7)	NS
Lam, 2010 ⁹ (19606455) Medium Explanatory	Cornell Scale for Depression in Dementia (CSDD) difference in difference 12 month	No statistical difference	Change from baseline -1.0 (-3.5, 1.0)	Change from baseline -1.5 (-4.0, 1.0)	NS
Lam, 2010 ⁹ (19606455) Medium Explanatory	NPI difference in difference 12 month	No statistical difference	Change from baseline -2.0 (-18.5, 6.0)	Change from baseline -7.0 (-19.0, 0.0)	NS
Eloniemi-Sulkava, 2001 ¹² (11890485) Medium Explanatory	Rate of institutionalization 2 years	No statistical difference	32% placement	30% placement	NS

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number

Table F-11. Caregiver outcomes summary low and medium risk of bias studies: case management

Study (PMID) Comparison Caregiver Type RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Chien, 2011 ⁸ (21198803) Case management vs usual care Medium Explanatory	Family caregiving burden inventory 18 months	F(1,90) = 7.09	Baseline 68.0 (SD 14.8) 18 month 45.5 (SD 10.0)	Baseline 66.9.0 (SD 14.8) 18 month 64.1 (SD 11.4)	<0.001
Chien, 2011 ⁸ (21198803) Case management vs usual care Medium Explanatory	WHOQoL 18 months	F(1,90) = 6.81	Baseline 64.8 (SD 13.0) 18 month 82.7 (SD 13.5)	Baseline 64.8 (SD 13.0) 18 month 64.5 (SD 13.1)	<0.001
Lam, 2010 ⁹ (19606455) Medium Explanatory	Personal well-being Index for Adult 12 month	No statistical difference	Change from baseline 2.9 (-15.0, 7.1)	Change from baseline 0.0 (-6.1, 5.0)	NS
Lam, 2010 ⁹ (19606455) Medium Explanatory	ZBI 12 month	No statistical difference	Change from baseline 5.0 (-10.5, 12.0)	Change from baseline 3.5 (-9.3, 12.3)	NS
Lam, 2010 ⁹ (19606455) Medium Explanatory	General Health Questionnaire 12 month	No statistical difference	Change from baseline 1.0 (-2.0, 5.5)	Change from baseline 0.0 (-2.0, 3.0)	NS

Abbreviations: NS=Not Significant; PMID=PubMed Identification Number; RoB-Risk of Bias; SD=Standard deviation; WHOQOL=World Health Organization Quality of Life Measure; ZBI=Zarit Burden Interview

Table F-12. Summary of strength of evidence for PLWD outcomes: consultation services

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Institutionalization Case management vs usual care	18-24 months	2 RCTs (n=192)	Mixed findings for institutionalization	Moderate	Unknown	Direct	Imprecise	Insufficient
NPI Case management vs usual care	12-18 months	2 RCT (n=194)	Mixed findings for NPI	Moderate	Unknown	Direct	Imprecise	Insufficient
Quality of Life Case management vs usual care	12 months	1 RCT (n=102)	No significant difference between groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Depression Case management vs usual care	12 months	1 RCT (n=102)	No significant difference between groups	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: N=Number; NPI=Neuropsychiatric Inventory; RCT=Randomized Controlled Trial

Table F-13. Summary of strength of evidence for caregiver outcomes: consultation services

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Quality of Life Case management vs usual care	12-18 months	2 RCTs (n=194)	Mixed findings for NPI	Moderate	Unknown	Direct	Imprecise	Insufficient
Caregiver Burden Case management vs usual care	12-18 months	2 RCT (n=194)	Mixed findings for NPI	Moderate	Unknown	Direct	Imprecise	Insufficient
Caregiver health status Case management vs usual care	12 months	1 RCT (n=102)	No significant difference between groups	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: N=Number; NPI=Neuropsychiatric Inventory; RCT=Randomized Controlled Trial

Evidence Map: Case Management

Table F-14. Characteristics of evidence map studies: case management

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Xiao, 2016 ⁶ (25805891) Australia High ROB	Case manager, home visit for initial assessment, monthly telephone contact	Usual care activities	Community RCT Caregivers	Dementia unspecified, from minority groups	N=61 Age: Yes Sex: Yes Race: No Education: no	(minority groups not specified)	N=61 Age: Yes Sex: Yes % majority race: No Education: no Relationship: Yes	Duration Living with PLWD Health status	NA	NA	6, 12 months	Blessed dementia score NPI QUORE	SSCQ SF-36v2
MacNeil Vroomen, 2015 ⁷ (26170035) Netherlands High RoB	Case management: (1) provided within one care organization (ICMM), (2) case management where multiple case management organizations are present within one region (LM)	Control: group with no access to case management	Home setting Non-RCT 521 dyads		N=521 Age: Yes Sex: Yes Race: No Education: Yes	Marital status Living condition	N=521 Age: Yes Sex: Yes Race: No Education: Yes Relationship: Yes	Living condition: Yes	NA	NA	2 years	Neuropsychiatric problems: NPI	Psychological health: (GHQ-12)
Brodady, 2003 ¹⁰ (12590626) Australia High RoB	Different models of care: (1) Psychogeriatric case management (2) Psychogeriatric consultation	Control: continued to receive whatever treatment they would have had were the survey not to have taken place.	Nursing home RCT 102 PLWD	Not specified	N=102 Age: Yes Sex: Yes Race: No Education: No	None	NA	NA	NA	NA	12 weeks	Depression: HAM-D; CSD; GDS & EBAS-DEP Psychopathology: (NPI) Behavior pathology: BEHAVE-AD	NA
Challis, 2002 ¹¹ (11994884) UK High ROB	Intensive case management, protected small caseloads per worker	Usual care activities	Community Matched case control 43 matched pairs	Not specified	N=86 Age: Yes Sex: Yes Race: No Education: No	SES	NR	NR	NA	NA	2 years	Unmet needs Service patterns and costs	IC: Caregiver burden
Newcomer, 1999 ¹⁵ (10445896) US Pilot demonstration	Medicare demonstration of case management (high attrition over 3 years)	Higher client to case manager ratio	Community RCT Caregivers	Alzheimer's disease	N=2731 Age: Yes Sex: Yes Race: yes Education: No	None	N=2576 Age: Yes Sex: Yes Race: No Education: no Relationship: Yes	None	NA	NA	3 years (attrition 64%)	NR	Caregiver burden and depression
Weinberger, 1993 ¹³ (8426038) Us High RoB	Social work case management	Usual care activities	Community RCT Caregivers	Progressive dementia unspecified	N=264 Age: Yes Sex: Yes Race: Yes Education: No	SES Household characteristics	N=264 Age: Yes Sex: Yes Race: Yes Education: Yes Relationship: Yes	Duration Health status Employment	NA	NA	6 months	Health utilization	NR

Abbreviations: BEHAVE-AD=Behavioral Pathology in Alzheimer's Disease; char=Characteristics; CSD=Cornell Scale for Depression; EBAS-DEP= Even Briefer Assessment for Depression; EM=Evidence Map; FC=Formal Caregiver; GDS=Geriatric Depression Scale; GHQ-12=General Health Questionnaire-12 Item; HAM-D=Hamilton Rating Scale for Depression; IC=Informal Caregiver; N=Number; NA=Not Applicable; NPI=Neuropsychiatric Inventory; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; QUORE=Quality of Patient's Eyes; RCT=Randomized Controlled Trial; RoB=Risk of Bias; RS=Reporting Status; SES=Socioeconomic Status; SF-36V2=Quality of Life Questionnaire; SSCQ=Sense of Competence Questionnaire

Care Protocols for PLWD

Table F-15. Risk of bias assessment: care protocols for PLWD

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Appelhof, 2019 ¹⁶ (30799167)	18 months stepped wedge	Stepped wedge	X	X	X	X	High		High
Husebo, 2019 ¹⁷ (30630722)	9 months	Medium 27%	Low	X	X	X	High	X	High
Livingston 2019 ¹⁸ (30872010)	8 months	Medium 21%	Medium	Medium	Medium	Low	High	Government	High
Nakanishi 2017 ¹⁹ (28857263)	6 months	Medium 15%	Low	High	Medium	Low	High	Government	High
Pieper 2016 ²⁰ (26804064)		Medium 11%	Low	Low	Low	Low	High	Nonprofit	High
McCabe, 2015 ²¹ (25319535)	3 months	Medium 13%	Low	Medium	Medium	Low	High		High
Zwijssen, 2014 ²² (24878214)	Stepped wedge		X	X	X	X	High		High
Rapp, 2013 ²³ (23827658)	10 months	Medium 15%	Medium	Low	High	Low	High		High
Cohen-Mansfield, 2012 ²⁴ (23059151)	2 weeks	High 46%	X	X	X	X	X		High
Kovach, 2012 ²⁵ (22998656)		Low	High 30%	X	X	X	X		High
Salva, 2011 ²⁶ (22159768)		Low	High 31%	X	X	X	X		High
Kovach, 2006 ²⁷ (16869334)	8 weeks	Medium 11%	Low	Low	High	Low	Low	Government	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Care Protocols for PLWD

Table F-16. Characteristics of evidence map studies: care protocols for PLWD

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Appelhof, 2019 ¹⁶ (30799167) Netherlands High RoB	The care program: Involves five steps: evaluation of psychotropic drug prescription, detection, analysis, treatment, & evaluation of treatment of NPS	Control: consisted of care as usual, without the educational program and use of the care program	Young-onset dementia Special care unit RCT (stepped wedge) 274 PLWD	Mild-severe dementia (All types)	N=274 Age: Yes Sex: yes Race: No Education: No	None	NA	NA	NA	NA	18 months	Agitation & aggression: CMAI (Dutch) NPS: NPI-NH (Dutch)	NA
Husebo, 2019 ¹⁷ (30630722) Norway High ROB	4-month training for multicomponent guideline-based, includes communication, systematic pain management, medication review, and activities	Usual care	Nursing homes Cluster RCT 33 nursing homes, both urban (52%) and non-urban (patients per cluster average 8)	Dementia unspecified	N=545 Age: Yes Sex: yes Race: No Education: No	None	NA	NA	NR	NR	9 months	QUALIDEM QUALID EQ-VAS ADL CGIC total medication	Staff distress

Livingston 2019 ¹⁸ (30872010) UK High RoB	MARQUE:6 sessions of manual-based intervention, followed by an implementation and supervision period (panel)	Treatment as usual	Care homes in UK Cluster RCT 20 Care homes 404 PLWD 492 CG	Clinical diagnosis of dementia	N=404 Age: Yes Sex: yes Race: yes Education: yes	None	NA	NA	N=492 Age: Yes Sex: yes Race: yes Education: yes	Length of service Training	8 months	CMAI NPI DEMQL EQ-5D 5L CDR	MBI SCID STS
Nakanishi 2017 ¹⁹ (28857263) Japan High RoB	BASE: consisted of training course; a web-based tool for ongoing monitoring and assessment for challenging behavior, and a multiagency discussion meeting.	Same as the intervention group, the care professionals' input of measures of their persons with dementia using the web-based tool at base-line. However, the web-based tool for the control group had no assessment of unmet needs nor an action plan.	Home care service providers Cluster RCT 283 PLWD 45 Home care providers 98 case managers and CG	Dementia diagnosis	N=283 Age: Yes Sex: yes Race: No Education: No	None	NA	NA	NR	NR	6 months	NPI-NH Abbey Pain Scale BI SMQ	NR
Chen, 2016 ²⁸ (26584896) Taiwan Training	PRT: Basic pain education + PRT	Basic pain education alone	Dementia special care unit RCT (Cluster) 37 NRs +195 PLWDs	NR	NA	NA	N=195 Age: Yes Sex: Yes Race: No Education: Yes	None	N=37 Age: Yes Sex: No Race: No Education: No	Years of experience: Yes	4 months? (Unclear) 3-month follow up	Pain reduction:	Improving the pain management performance of RNs
Pieper 2016 ²⁹ Netherlands High ROB	STA OP! Stepwise multicomponent intervention for NH healthcare professionals Training on general nursing skills, dementia management and pain	Treatment as usual	Nursing home-based Cluster RCT 21 clusters in 12 nursing homes N=288 PLWD	Nonspecified advanced dementia Reisberg Global Deterioration Scale 5-7	N=288 Age yes Sex: yes Race: no Education no	NR	NA	NA	NR	NR	3, 6 months	CMAI NPI-NH CSDD Minimum Dataset Depression in Dementia Psychotropic medication use	NR
Kovach, 2015 ³⁰ (26250849) USA Feasibility study	T3 Protocol	Usual care	Long term care facility RCT 78 PLWD	NR	N=127 Age: Yes Sex: Yes Race: No Education: No	None	NA	NA	NA	NA	8 weeks	Feasibility, assessments, treatment changes, nurse time, and drug costs	NA
McCabe, 2015 ²¹ (25319535) Australia High RoB	The BPSD protocol training: (1) Training & support condition: Training in the use of a BPSD - structured clinical protocol, plus external clinical support (2)Support condition: a workshop on BPSD, plus external clinical support (3) Training condition: training in the use of the structured clinical protocol alone	Care as usual	Residential care facility RCT 475 (staff=261 + PLWD=214)		N=187 Age: Yes Sex: Yes Race: No Education: No	None	NA	NA	N=204 Age: Yes Sex: No Race: No Education: no	Job position: Yes Service length: Yes Dementia training: Yes	3 months	CMAI	Staff stress: Carer stress scale General strain: SDCS Self-efficacy: Staff attitude toward PLWD: (ADQ)

Zwijzen, 2015 ³¹ (25458804) Netherlands High RoB	A new care programme for challenging behaviour: care staff was educated on how to detect and reflect on signs of challenging behavior, consisting of four steps; detection, analysis, treatment & evaluation	Control (Stepped wedged Cluster)	Dementia special care Unit RCT (Stepped wedge cluster) 380?	NA	NA	NA	NA	NA	N=380 Age: Yes Sex: No Race: No Education: no	Education: Yes Job position: Yes Service length: Yes Dementia training: Yes	4 months	NA	Burnout: MBI Job satisfaction & job demands: Stress: Saliva samples Comorbid burden:
Rapp, 2013 ³³ 23827658) German High ROB (Protocol in German)	Guideline-based intervention includes clinical assessments, nondrug interventions, medication review to reduce agitation	Usual care	Nursing homes Cluster RCT 18 nursing homes PIWD	Dementia not specified	N=304 Age: Yes Sex: Yes Race: No Education: no Marital status: no	None	NA	NA	NR	NR	10 month	CMAI Number of neuroleptics	NR
Cohen-Mansfield, 2012 ³² (23059151) US High ROB	TREA decision tree protocol to assess unmet needs for agitation	Usual care	Nursing homes Cluster RCT 11 nursing homes PLWD	Dementia not specified	N=125 Age: Yes Sex: Yes Race: yes Education: yes Marital status: yes	None	NA	NA	NR	NR	2 weeks	ABMI LMBS	NR
Kovach, 2012 ³³ (22998656) USA High RoB	Serial Trial Intervention (STI): A decision support tool-9-Step STI	5-Step STI	Nursing home Quasi 125 PLWD	Moderate to severe dementia	N=125 Age: Yes Sex: Yes Race: Yes Education: No	None	NA	NA	NA	NA	6 weeks	Discomfort & agitation	NA
Montgomery, 2011 ³⁴ (21840840) US Pilot	TCARE protocol decision algorithm assists care managers create care plans	Usual care activities	Community RCT 266 caregivers	Dementia unspecified; caregivers with high burden such that considering nursing home placement	NR	NR	N=266 Age: Yes Sex: Yes Race: No Education: Yes Relationship: yes	Health status	NA	NA	9 months	NR	Modified Montgomery Borgatta Caregiver Burden Scale; CESD; Intention to place
Salva, 2011 ²⁶ (22159768) Spain High ROB	Health and nutrition promotion, includes nutritional counseling	Usual care activities	Community Cluster RCT 11 Medical centers dyad	Mild to moderate dementia	N=946 Age: Yes Sex: Yes Race: no Education: yes		N=946 Age: no Sex: no Race: No Education: no Relationship: yes	Duration Received payment	NA	NA	1 year	CDR NPI MMSE Eating behavior scale	Zarit burden
Verkaik, 2011 ³⁵ (21495077) Netherlands Training	Introduction of a nursing guideline: Nursing team introduced nursing guideline that entails increasing individualized pleasant activities and decreasing unpleasant events.	Usual care	Nursing home RCT 100 PLWD	moderately severe dementia (All types)	N=195 Age: Yes Sex: Yes Race: No Education: Yes Marital status: Yes	None	NA	NA	NA	NA	11 months	Depression; CSDD (Dutch version Mood	NA
Cohen-Mansfield, 2007 ³⁶ (17702884) US Pilot	TREA decision tree protocol to assess unmet needs for agitation	Usual care	Nursing homes Quasi-experimental PLWD	Dementia not specified	N=167 Age: Yes Sex: Yes Race: yes Education: yes Marital status: yes	None	NA	NA	NR	NR	10 days	ABMI LMBS	NR

Kovach, 2006 ³⁷ (16869334) USA High RoB	STI: an innovative clinical protocol for assessment & management of unmet needs in people with late-stage dementia.	Control: Standard care	Long term care facility RCT 127 PLWD	Late stage dementia	N=114 Age: Yes Sex: Yes Race: No Education: Yes	None	NA	NA	NA	NA	8 weeks	Discomfort-DAT and BEHAVE-AD scales Daily Behavior symptoms, assessment & treatment: Daily logs	NA
---	---	---------------------------	--	---------------------	---	------	----	----	----	----	---------	--	----

Abbreviations: ABMI=Agitation behavior mapping instrument; ADL=Activities of Daily Living; ADQ=Approaches to Dementia Questionnaire; BASE=Behavior Analytics and Support Enhancement; BEHAVE-AD=Behavioral pathology in Alzheimer's disease; BI=Barthel Index; BPSD=Behavioral and psychological symptoms of dementia; CDR=Clinical Dementia Rating; CESD=Center for Epidemiological Studies-Depression scale; Discomfort-DAT= Discomfort-Dementia of the Alzheimer's Type; CG=Caregiver; char=characteristics; CGIC=Clinical Global Impressions of Change; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; Discomfort-DAT= Discomfort-Dementia of the Alzheimer's Type; DEMQOL=Dementia Quality of Life; EM=Evidence Map; EQ-VAS=European Quality of Life visual analog scale; EQ-5D-5L=European quality of life five dimensions questionnaire; FC=formal caregiver; IC=informal caregiver; LMBS=Lawtons modified behavior stream; MBI=Maslach Burnout Inventory; MARQUE=Managing Agitation and Raising Quality of Life; MBI=Maslach Burnout Inventory; MMSE=Mini-Mental Status Examination; N=number; NA=Not Applicable; NPI=Neuropsychiatric Inventory; NPI-NH=Neuropsychiatric Inventory Home Version; NPI-Q=Neuropsychiatric Inventory Questionnaire; NR=Not Reported; NPS=Neuropsychiatric symptoms; PMID=PubMed Identification Number; PRT=Pain Recognition and Treatment; PLWD=Persons Living with Dementia; QUALID=Quality of Life late-stage dementia scale; QUALIDEM=Quality of Life dementia scale; RCT=Randomized Controlled Trial; RN=Registered Nurse; RoB=Risk of Bias; RS=Reporting Status; SCID=sense of Competence in Dementia; SDCS=Strains in Dementia Care Scale; SMQ=Short Memory Questionnaire; STA OP!=Dutch Serial Trial Intervention; STI=Serial Trial Intervention; STS=Staff Tactics Scale; T3=Track and Trigger Treatment; TCARE=Tailored Caregiver Assessment and Referral; TREA=Treatment Routes for Exploring Agitation; YOD=Young-onset Dementia

Advance Care Planning

Table F-17. Risk of bias assessment: advance care planning

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Brazil 2018 ³⁸ (28786323)	6 weeks	Medium 19.3%	Low	Low	High	Low	X	Government, Foundation	High
Mitchell 2018 ³⁹ (29868778) Cohen 2019 ⁴⁰ (30273717)	6 months 12 months	Low 12 months: 2%	Low	Low	Low	Medium	Low	Government	Medium
Hanson 2017 ⁴¹ (27893884) Hanson 2016 ⁴² (27271683)	3 months 6 months 9 months	Low 3 months: 1% 6 months: 8.2% Medium 9 months: 15.6%	Low	Low	3 months, 9 months: Low 6 months: High	High	X	Government	High
Reinhardt 2014 ⁴³ (24835382)	3 months 6 months	Medium 3 months: 12.7% 6 months: 18.1%	Medium	Low	High	Low	X	Foundation	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table F-18. Characteristics of included studies: advance care planning

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Mitchell 2018 ³⁹ (29868778) US Medium Explanatory Cohen 2019 ⁴⁰ (30273717)	Advanced Care Planning 12-minute video for goal-oriented care (advanced care directives, care preferences, etc)	Usual care	Nursing home-based Cluster RCT 64 nursing homes 402 PLWD	Nonspecified advanced dementia Primarily AD Global Deterioration Scale 7	N=402 Age 87 80% Female 87% White Education NR	None	N=402 Age 62 66% Female 87% White 99% High school diploma or more 63% Child 13% Spouse	None	NA	NA	6 months 12 months	Hospitalizations	Documented directives Goal of care discussions Comfort Care Preference Prevalence

*High risk of bias studies included in evidence map

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; N=number; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status

Table F-19. PLWD outcomes summary low and medium risk of bias studies: advance care planning

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Mitchell 2018 ³⁹ (29868778) Decision Support Tool vs. Usual Care Medium Explanatory	Hospitalizations N (%) 12 months	NR	20 (9.5%)	21 (11.1%)	NR
Hanson 2011 ⁴⁴ (22091750) Decision Support Tool vs. Usual Care Medium Explanatory	Assisted Feeding Intervention-Specialized Dysphagia Diet (%) 3 months	NR	89%	76%	0.04

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number

Table F-20. Caregiver outcomes summary low and medium risk of bias studies: advance care planning

Study (PMID) Comparison Caregiver Type RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Mitchell 2018 ³⁹ (29868778) Decision Support Tool vs. Usual Care Medium Explanatory	Documented Do Not Hospitalize Directive N (%) 6 months, 12 months	Adjusted Odds Ratio (95% CI) 6 months: 1.08 (0.69, 1.69) 12 months: 1.07 (0.66, 1.72)	6 months: 133 (63.0%) 12 months: 144 (68.2%)	6 months: 119 (63.0%) 12 months: 126 (66.7%)	NR
Mitchell 2018 ³⁹ (29868778) Decision Support Tool vs. Usual Care Medium Explanatory	Goal of care discussions N (%) 6 months, 12 months	Adjusted Odds Ratio (95% CI) 6 months: 1.70 (0.94, 3.07) 12 months: 1.46 (0.86, 2.70)	6 months: 49 (23.2%) 12 months: 72 (34.1%)	6 months: 29 (15.3%) 12 months: 48 (25.4%)	NR
Mitchell 2018 ³⁹ (29868778) Decision Support Tool vs. Usual Care Medium Explanatory	Proxies Preferring Comfort Care N (%) 6 months, 12 months	Adjusted Odds Ratio (95% CI) 6 months: 1.28 (0.85, 1.94) 12 months: 0.72 (0.38, 1.38)	6 months: 153 (73.2%) 12 months: 159 (76.1%)	6 months: 140 (76.9%) 12 months: 151 (82.1%)	NR

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number

Table F-21. Summary of strength of evidence for PLWD outcomes: advance care planning

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Decision Support Tool vs. Usual Care Hospitalizations	12 months	1 cluster RCT (n=402)	Similar rates of hospitalizations between intervention and control groups.	Moderate	Unclear	Direct	Imprecise	Insufficient

Abbreviations: n=Number; RCT=Randomized Controlled Trial

Table F-22. Summary of strength of evidence for caregiver outcomes: advance care planning

Comparison Caregiver Type Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Decision Support Tool vs. Usual Care Documented Do Not Hospitalize Directive	6 months 12 months	1 cluster RCT (n=402)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Decision Support Tool vs. Usual Care Goal of care discussions	6 months 12 months	1 cluster RCT (n=402)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Decision Support Tool vs. Usual Care Proxies Preferring Comfort Care	6 months 12 months	1 cluster RCT (n=402)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient

Abbreviations: n=Number; RCT=Randomized Controlled Trial

Evidence Map: Advance Care Planning

Table F-23. Characteristics of evidence map studies: advance care planning

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Goossens 2019 ⁴⁵ (31818522) Goossens 2019 ⁴⁶ Belgium High RoB	Knowledge-sharing on shared decision making and internal policies role-play exercises on advance care planning	Waitlist control	Nursing Home Cluster RCT 65 Nursing homes N=311 formal caregivers	None	None	NA	NA	NA	N=311 Age: yes Sex: yes Race: no Education: yes	Training: yes Education: yes Position: yes Length of Service: yes	NR	None	Training
Song, 2019 ⁴⁷ (31373868) US Small sample	Adapt SPIRIT (Sharing Patient's Illness Representation to Increase Trust) for PLWD and surrogates; 60 psycho-education	In-person vs videoconference	Community Mixed-methods Dyad	Early stage dementia	N=23 Age: yes Sex: yes Race: yes Education: yes	None	N=23 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	Duration	NA	NA	3 days	Dyad congruence Patient decisional conflict scale	Surrogate decision-making confidence scale
Brazil 2018 ³⁸ (28786323) Northern Ireland High RoB	Advance care planning intervention (education, meetings documentation); 2 family meetings	Usual care	Nursing homes Cluster RCT 24 nursing homes 197 Informal caregivers	Dementia without capacity to complete ACP	N=197 Age: no Sex: no Race: no Education: no	None	N=197 Age: yes Sex: yes Race: no Education: no Relation to PLWD: yes	None	NA	NA	6 weeks	None	Decisional Conflict Scale FPCS GHQ QOD-LTC
Hanson 2017 ⁴¹ (27893884) US High RoB Hanson 2016 ⁴² (27271683)	Goals of care video decision aid and structured discussion with health care providers	Informational video	Nursing homes Cluster RCT 22 Nursing homes 302 PLWD and informal caregivers	Severe to advanced dementia; GDS 5 to 7	N=302 Age: yes Sex: yes Race: yes Education no	None	N=302 Age: yes Sex: yes Race: yes Education: yes Relation to PLWD: yes	None	NA	NA	3 months 9 months	None	Quality of Communication Concordance of GOC ACP problem score SWC-EOLD SM-EOLD
Bonner, 2014 ⁴⁸ (24381040) US Pilot Study	Advance Care Treatment Plan (ACT-Plan); a group-based education intervention, with AA dementia caregivers.	Attention-control health promotion conditions.	Adult Daycare centers Non-RCT 68 caregivers	NA	NA	NA	N=32 Age: Yes Sex: Yes Race: No Education: No	Income: Yes Living with PLWD: Yes	NA	NA	4 weeks	Feasibility:	Knowledge of dementia, CPR, MV & TF; Knowledge of dementia scale; Self-efficacy: the Confidence in Treatment Decisions Made questionnaire
Einterz, 2014 ⁴⁹ (24508326) US Pilot study	GOC decision aid video viewed by the SDM and structured care plan meeting between the SDM and interdisciplinary NH team.	Usual care	Nursing home RCT 18 Dyads	All types of dementia with moderate to severe (GDS)	N=18 Age: yes Sex: yes Race: yes Education: no	None	N=18 Age: yes Sex: yes Race: yes Education: yes Relationship with PLWD: yes	None	NA	NA	3 months	None	Quality of communication and decision-making

Reinhardt 2014 ⁴³ (24835382) US High RoB	Structured conversations about end of life care with palliative care team	Social contact via telephone	Skilled nursing facility 110 informal caregivers	Advance dementia, CPS score of 4 to 6	N=110 Age: yes Sex: yes Race: yes Education: no	None	N=101 Age: yes Sex: yes Race: yes Education: yes Relationship with PLWD: yes	Employment status: yes	NA	NA	3 months 6 months	None	Satisfaction of care at end-of-life in dementia scale PHQ-9 Satisfaction with Life Scale
Sampson, 2011 ⁵⁰ (21228087) UK Pilot	A palliative care and advance care plan (ACP) intervention	Usual care	Hospital RCT 32 dyads	Severe dementia	N=32 Age: yes Sex: yes Race: yes Education: no	None	N=32 Age: yes Sex: yes Race: yes Education: no Relationship: yes	None	NA	NA	6 weeks 6 months	Number of caregivers making an ACP: Feasibility: Pain: Visual analogue scale distress: KD10 DSI: DCS; STAXI; LSQ	distress: KD10 DSI: DCS; STAXI; LSQ

Abbreviations: char=characteristics; EM=Evidence Map; FC=formal caregiver; IC=informal caregiver; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RS=Reporting Status; SES=socioeconomic status

Palliative Care

Table F-24. Risk of bias assessment: palliative

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Verrault , 2018 ⁵¹ (28731379)	1 year	Medium (response rates for caregivers) 57% control, 73% intervention	Medium	Medium	High	Low	X	Government	High
Boogaard, 2018 ⁵² (29343173)	10 months	Medium 11% nursing homes dropped	Medium	X	X	X	High	Gov't	High
Agar, 2017 ⁵³ (28786995)		Cluster RCT	Medium 64% decline rate	X	X	X	High (could not measure as planned)		High
Ahronheim, 2000 ⁵⁴ (15859668)	Over 3 years	Low 0%	Medium	Low	Medium	Low	High		High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Palliative Care

Table F-25. Characteristics of evidence map studies: palliative care

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia IC Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Hanson, 2019 ⁵⁵ (30342242) USA Pilot	Triggered Palliative Care: Received denetia-specific specialty palliative care consultation plus postacute transitional care.	Control: Received usual care and educational information	Hospital RCT 62 dyads	Late-stage (5-7) dementia	N=62 Age: Yes Sex: Yes Race: Yes Education: No	None	N=62 Age: yes Sex: Yes Race: No Education: no Relation: yes	None	NA	NA	60 days	60-day hospital or emergency department visits Patients comfort, family distress etc.	NR
Boogaard, 2018 ⁵² (29343173) Netherlands High ROB	Generic or personal feedback to staff from PLWD about end of life care	No feedback	Nursing homes Cluster RCT 18 nursing homes	Late stage dementia	N=668 Age: Yes Sex: Yes Race: no Education: No	None (length of stay)	N=193 Age: yes Sex: Yes Race: No Education: yes Relation: yes	None	NA	NA	10 months	NR	End-of Life in Dementia scales
Verrault, 2018 ⁵¹ (28731379) Canada High ROB	End of Life care: training for physicians and staff; pain monitoring; regular mouth care; family communication; nurse facilitator	Usual care activities	Long-term care facilities Quasi-experimental 193 dyads	Late stage dementia	N=193 Age: Yes Sex: Yes Race: no Education: No	None (length of stay) (duration of terminal phase)	N=193 Age: no Sex: Yes Race: No Education: no Relation: yes	None	NA	NA	1 year	NR	Family perception of care scale; Symptom management for EoL in dementia Comfort assessment in dying scale
Agar, 2017 ⁵³ (28786995) Australia High ROB	Facilitated family case conference for end of life care, registered nurse	Usual care activities	Nursing homes Cluster RCT 20 nursing homes	Late stage dementia	N=131 Age: Yes Sex: Yes Race: no Education: No	None (length of stay) (time to death) (visitor frequency)	None	None	NA	NA	4-6 weeks post-death	Nurse-rated QUALID	Family-rated End of Life Care scales
Ahronheim, 2000 ⁵⁴ (15859668) US High ROB	Palliative care team recommendations	Usual care activities	Hospital	Late stage dementia	N=99 Age: Yes Sex: Yes Race: yes Education: no	None (length of stay)	None	None	NA	NA	Time until death	Number of medical interventions, procedures, and readmissions	NR

Abbreviations: char=characteristics; EM=Evidence Map; FC=formal caregiver; IC=informal caregiver; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; QUALID=Quality of Life in Late-stage Dementia; RS=Reporting Status; SES=socioeconomic status

Other Service Provision Interventions

Table F-26. Risk of bias assessment: other service provision interventions

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Horvath 2013 ⁵⁶ (24195007)		Medium 15%	Low	Medium	High	Low	High		High
Hanson 2011 ⁴⁴ (22091750) Hanson 2010 ⁵⁷ (20729251) Snyder 2013 ⁵⁸ (23273855)	3 months 9 months	Low 3 months: 1% High 9 months:	Low	Medium	Low	Low	Medium	Government	Medium

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Table F-27. Characteristics of included studies: other service provision interventions

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Hanson 2011 ⁴⁴ (22091750) US Medium Explanatory Hanson 2010 ⁵⁷ (20729251) Snyder 2013 ⁵⁸ (23273855)	Decision aid about dementia, feeding options (feeding tubes, assisted oral feeding); print format reviewed for ~20 minutes	Usual care	Nursing homes Cluster RCT 24 Nursing homes 256 PLWD-surrogate dyads	Advanced dementia with feeding problems; score of 5 to 6 on Cognitive Performance Scale in the MDS and severity stage 6 to 7 on GDS	N=256 85 years 78% Female 70% White Education NR	Detailed Race Information: yes	N=256 59 years 63% Female 70% White Education NR 45% Daughter	None	NA	NA	3 months	Assisted Feeding Intervention, Specialized dysphagia diet	Decisional Conflict Scale Frequency of Discussion about Feeding Satisfaction with Decision Scale Decisional Regret Index

*High risk of bias studies included in evidence map

Abbreviations: char=characteristics; FC=formal caregiver; IC=informal caregiver; N=number; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status

Table F-28. PLWD outcomes summary low and medium risk of bias studies: other service provision interventions

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Hanson 2011 ⁴⁴ (22091750) Decision Support Tool vs. Usual Care Medium Explanatory	Assisted Feeding Intervention-Specialized Dysphagia Diet (%) 3 months	NR	89%	76%	0.04

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number; RoB=Risk of Bias

Table F-29. Caregiver outcomes summary low and medium risk of bias studies: other service provision interventions

Study (PMID) Comparison Caregiver Type RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Hanson 2011 ⁴⁴ (22091750) Decision Support Tool vs. Usual Care Medium Explanatory	Decisional Conflict Scale, Overall Mean Mean Change from Baseline 3 months	NR	1.65 -0.60	1.97 -0.13	<0.001
Hanson 2011 ⁴⁴ (22091750) Decision Support Tool vs. Usual Care Medium Explanatory	Satisfaction with Decision Scale Mean 3 months	NR	1.61	1.66	0.5
Hanson 2011 ⁴⁴ (22091750) Decision Support Tool vs. Usual Care Medium Explanatory	Decisional Regret Index Mean 3 months	NR	11.9	14.3	0.14
Hanson 2011 ⁴⁴ (22091750) Decision Support Tool vs. Usual Care Medium Explanatory	Feeding Discussions, With Physician, NP, or PA Feeding Discussions, Other Nursing Home Staff % 3 months	NR	45% Other staff: 64%	33% Other staff: 71%	0.04 Other staff: 0.42

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number

Table F-30. Summary of strength of evidence for PLWD outcomes: other service provision interventions

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Decision Support Tool vs. Usual Care Assisted Feeding Intervention-Specialized Dysphagia Diet	3 months	1 cluster RCT (n=256)	At 3 months, more PLWD in intervention group had a specialized dysphagia diet.	Moderate	Unclear	Direct	Imprecise	Insufficient

Abbreviations: N=Number, RCT=Randomized Controlled Trial

Table F-31. Summary of strength of evidence for caregiver outcomes: other service provision interventions

Comparison Caregiver Type Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Decision Support Tool vs. Usual Care Decisional conflict	3 months	1 cluster RCT (n=256)	At 3 months, intervention group had less decisional conflict than control group.	Moderate	Unclear	Direct	Imprecise	Insufficient
Decision Support Tool vs. Usual Care Satisfaction with Decisions	3 months	1 cluster RCT (n=256)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Decision Support Tool vs. Usual Care Decisional Regret	3 months	1 cluster RCT (n=256)	No difference between groups.	Moderate	Unclear	Direct	Imprecise	Insufficient
Decision Support Tool vs. Usual Care Feeding Discussions	3 months	1 cluster RCT (n=256)	At 3 months, more feeding discussions with doctor, PA, or NP in intervention group versus control. No difference between groups in feeding discussions with other nursing home staff.	Moderate	Unclear	Direct	Imprecise	Insufficient

Abbreviations: n=number; RCT=Randomized Controlled Trial

Evidence Map: Other Service Provision Interventions

Table F-32. Characteristics of evidence map studies: other service provision interventions

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Laver 2020 ⁵⁹ 32234275 Australia Pilot	Telehealth	Usual care	N=63	Diagnosed dementia or probable dementia or MMSE < 24/30	N=63 Age: yes Sex: yes Ses: yes	Age: yes Sex: yes Household characteristics: yes	NA	Age: yes Sex: yes Household characteristics: yes	NA	NA	16 weeks		CMI PCS CAPU
Villar 2019 ⁶⁰ (29149789) Spain Pilot	Including PLWD with dementia in care planning meetings	Usual care	Nursing homes Cluster RCT Clusters=4 N=52	PLWD living in a residential facility for at least six months, diagnosed with moderate-to-severe dementia	N=52 Age: yes Sex: yes	NR	NA	NA	NR	NR	20-30 minutes	the GDS and GENCAT	NR
Williams 2019 ⁶¹ (31660443) US Pilot	FamTechCare video recording support	Telephone support/ attention control	Community-based RCT	Patients diagnosed with dementia Living at home Exclusion criteria	N=39 Intervention group with 42 caregivers 32 Control group with 41 caregivers Age: Y Sex: Y Education: Y Race % majority: Y	N caregivers: Y Ethnicity: Y	Age: Y Sex: Y Race % majority: Y Education: Y Relation to PLWD: Y	Ethnicity: Y Marital status: Y Care provided: Y	NR	NR	3 months	NR	Burden Depression Sleep disturbance Competence Desire to institutionalize PLWD Reaction to behavioral symptoms
Horvath 2013 ⁵⁶ (24195007) US High ROB	Self-directed educational program for caregiver competence and patient safety; home-based safety tool kit	Customary care	Community-based Recruited from 2 care centers Block randomization RCT	Patients diagnosed with progressive AD PLWD reads and speaks English MMSE <=24 Primary informal caregiver at least 4 hours per day Exclusion criteria	N=60 Experimental group with 60 caregivers 48 control group with 48 caregivers Age: Y Sex: Y Education: N Race % majority: Y	NR	Age: Y Sex: Y Race % majority Education: N Relation to PLWD: Y	NR	NR	Training: Y	12 weeks	Home safety evaluation Risky behaviors and accidents	Self-efficacy Strain

Abbreviations: AD=Alzheimer's disease; char=characteristics; MMSE=Mini Mental State Examination; N=Number; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Person with dementia; RCT=Randomized Controlled Trials; RoB=Risk of Bias; RS=Reporting Status

Collaborative/Coordinated Care Models

Table F-33. Risk of bias assessment: multidisciplinary integrated team collaborative care

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Halek 2020 ⁶² (32062052)	15 days/unclear	Medium 24%	Low	Low	High	Low	Low	Foundation	High
Possin, 2019 ⁶³ 31566651	12 month	Not high	Not high	X	X	X	Not high	Gov't	Move to PRECIS-2
Laporte Uribe 2017 ⁶⁴ (28249632)	12 months	X	High	X	X	X	X	X	High
Thyrian, 2017 ⁶⁵ 28746708	12 months	Medium 21%	Medium (selection concern at GP level)	X	X	X	Medium	Gov't	Move to PRECIS-2
Chodosh, 2015 ⁶⁶ 25656074	6, 12 month	Medium 36% 6 month	Low	Low	Medium	Low	Low	Gov't	Medium (12 month High)
Leontjevas, 2014 ⁶⁷ 23643110	4 months	Low (stepped wedge)	Low	High	Medium	Low	High (43% adherence to treatment)	Gov't	High (for pragmatic) High (for traditional)
Bass, 2013 ⁶⁸ 23869899	6 months	Medium 18% (12 month almost 40%)	Medium	X	X	X	Low	Gov't	Move to PRECIS-2
Chapman, 2007 ⁶⁹ 8537594	8 weeks	Low 2%	Medium	High	Medium	Low	Medium	Gov't	High
Callahan, 2006 ⁷⁰ 16684985	12 month	Medium 18%	Low	Low	Medium	Low	Medium	Gov't	Medium (18 month High ROB)
Vickrey, 2006 ⁷¹ 17116916	22 months (mean)	Not high	Not high	X	X	X	Not high	Gov't	Move to PRECIS-2
Surr, 2019 ⁷² 31056923	16 months	OK	OK	X	X	X	High (26%)		High ROB for Pragmatic
Ballard, 2018 ⁷³ 29408901	9 months	Medium	Low	Medium	Medium	Low	Medium	Gov't	Medium
Lichtwarck, 2018 ⁷⁴	12 weeks	Medium 12%	Low	Low	Med	Low	Low	Gov't	Medium
Dichter, 2015 ⁷⁵ 26138674		Rolling enrollment	Medium	X	X	X	Medium	Gov't	High
Chenoweth, 2014 ⁷⁶ 24666667	8 months	High 31%	X	X	X	X	X	Gov't	High
van de Ven, 2013 ⁷⁷	8 months	Rolling stable	Low	X	X	X	Medium	Gov't	Move to PRECIS-2
Rokstad, 2013 ⁷⁸ 24022375		Medium 25%	Low	X	X	X	High	Gov't	High
Chenoweth, 2009 ⁷⁹ 19282246	8 months	Low 18%	Low	Low	High	Low	High	Gov't	Medium
Eloniemi-Sulkava, 2009 ⁸⁰ (20121986)	1.6 years	Low 0% (use of census data)	Low	Low	Medium	Low	Medium	Gov't	Low
Fossey, 2006 ⁸¹ 16543297	12 months	Unclear/High	Low	Medium	Medium	Low	High	Gov't	High
Rovner, 1996 ⁸² 8537594	6 months	Low 9%	Low	Medium	High	Low	High	Gov't	Medium

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NR=Not Reported; PMID=PubMed Identification Number

Table F-34. Characteristics of included studies: collaborative/coordinated care models

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:

Possin, 2019 ⁶³ 31566651 [add others] US Medium Pragmatic	Collaborative Care Care team navigator; multidisciplinary team Dyad Telephone/ Internet See Appendix Table F.37	Usual care plus contact information for resources, quarterly newsletters	California, Nebraska, and Iowa participating health systems RCT 780 community-based dyads	Diagnosis by treating physician, over age 45 50% Mild, 28% moderate, 22% advanced	N=780; 512 intervention, 268 control Age: 78 (mean) % Female: 56% Race: 80% white Education: 60% post-secondary	SES Dementia Severity Comorbidity	IC N=780; 512 intervention, 268 control IC Age: 65 (mean) IC: 62% Female IC Race: 80% white IC Education: 60% post-secondary IC Relation to PLWD: 55% Spouse	Language Living with PLWD Health status	NA	NA	6, 12 months	QoL-AD Utilization rates	IC: ZBI PHQ-9 Self-efficacy
Thyrian, 2017 ⁶⁵ 28746708 24152974 ⁸³ 24811145 ⁸⁴ 24225205 ⁸⁵ 27534949 ⁸⁶ 22575023 ⁸⁷ 2012 2012 29156941 ⁸⁸ 31409541 ⁸⁹ Germany Pragmatic	Collaborative care/care coordination model; Nurse care coordinators Dyad In-home See Appendix Table F.37	Usual care activities	Community-based Cluster RCT 136 General practitioners PLWD	Dementia aged 70+; DemTect instrument for dementia screening in Germany	N=516; 348 intervention, 168 control Age: 80 (mean) % Female: 61% Race: NR Education: NR	Household characteristics	N= IC Age: NR IC: NR IC Race: NR IC Education: NR IC Relation to PLWD: 55% Spouse	NR	NA	NA	6 months	QoL-AD NPI ADL Health care resource utilization	Berlin Inventory Caregiver Burden
Chodosh, 2015 ⁶⁶ 25656074 (related to ACCESS ⁷¹) US Medium Balanced	Coordinated care program; guideline-based Social work care managers Dyad In-person plus telephone See Appendix Table F.37	Coordinated care program by telephone contact	Urban underserved community RCT 151 community-based dyads	Dementia Diagnosis by system records, physician, or caregiver-confirmed Dementia severity 11 (0-17 Blessed Roth scale)	N=151; 73 intervention, 71 control Age: 73 (mean) % Female: 63% Race: 74% Hispanic/Latino Education: 69% less than high school	None	IC N=151; 73 intervention, 71 control IC Age: 49 (mean) IC Sex: 65% Female IC Race: 78% Hispanic/Latino IC Education: 36% less than high school IC Relation to PLWD: 54% child	Acculturation scale Living with PLWD	NA	NA	6, 12 months	HUI Utilization rates	IC: ZBI RMBPC
Bass, 2013 ⁶⁸ 23869899 25666216 ⁹⁰ 29854922{ US Medium Pragmatic	Care coordination with Partners in Care coaching model Care coordinators Caregivers Telephone, email, mail See Appendix Table F.37	Usual care activities	Veterans health system Matched system-level pairs randomized	At least one dementia diagnosis in VA medical records receiving primary care, aged 50+	N=508	NA	IC N=486; 299 intervention, 187 control IC Age: 69 (mean) IC Sex: NR IC Race: NR IC Education: NR IC Relation to PLWD: 73% spouse	Duration Employment	NA	NA	6, 12 months	None	IC: Unmet needs Role captivity Physical health strain CESD Caregiver support service use Number informal helpers
Vickrey, 2006 ⁷¹ 17116916 ACCESS 19670955 ⁹¹ 22435836 ⁹² 23320655 ⁹³ US Medium Pragmatic	Coordinated care program; guideline-based Social work care managers Dyad Telephone See Appendix Table F.37	Usual care	Urban California Cluster RCT 18 primary care clinics 408 community-based dyads	Dementia over age 65 receiving Medicare; 76% AD Administrative database codes Dementia severity 5.7 intervention, 6.3 control (0-17 Blessed Roth scale)	N=408; 238 intervention, 170 control Age: 80 (mean) % Female: 55% Race: 86% White Education: 81% at least high school	Health Insurance Comorbidity	IC N=408; 238 intervention, 170 control IC Age: 65 (mean) IC Sex: 69% Female IC Race: 87% White IC Education: 94% at least high school IC Relation to PLWD: 55% spouse	Duration Living with PLWD Comorbidity	NA	NA	22 months (mean follow-up)	Adherence to 23 dementia guidelines recommendations HUI	Caregiver knowledge, mastery, confidence EuroQoL-5D
Callahan, 2006 ⁷⁰ 16684985 US Medium Explanatory	Collaborative care program; integrated in primary care Geriatric nurse practitioner care manager Dyad In-person See Appendix Table F.37	Augmented usual care; physician could pursue any evaluation or treatment deemed appropriate	Urban University-affiliated care systems Cluster RCT 74 primary care physicians 153 community-based dyads	Possible or probably Alzheimer disease; Diagnostic and Statistical Manual of Mental Disorders, 3 rd edition	N=153; 84 intervention, 69 control Age: 77 (mean) % Female: 43% Race: 49% Black Education: 8-9 years (mean)	Medicaid SES Chronic disease score	IC N=153; 84 intervention, 69 control IC Age: 61 (mean) IC Sex: 89% Female IC Race: NR IC Education: NR IC Relation to PLWD: 44% spouse/36% child	Live with patient	NA	NA	6, 12, 18 months	NPI ADL CSDD (proxy) Health care resource use MMSE	IC: PHQ-9
Lichtwarck, 2018 ⁷⁴ 23992241 ⁹⁴ Process eval ⁹⁵ Norway Medium Explanatory	TIME Person centered care and CBT Team Staff training PLWD assessment and tailored plan	Brief education only, usual care activities	Nursing homes Cluster RCT 33 nursing homes PLWD	Probably dementia, 27% moderate, 69% severe CDR 1 or higher; moderate degree of agitation	N=229; 125 intervention, 104 control Age: 83 (mean) % Female: 70% Race: NR Education: NR	None	NA	NA	NA	NA	8, 12 weeks	NPI CMAI CSDD QoL in late stage dementia scale	NR

Ballard, 2018 ⁷³ 29408901 Sustainability ⁹⁶ Cost effectiveness ⁹⁷ UK Medium Balanced	WHELD programme person-centered care and social interaction Team PLWD Training and delivery of WHELD	Usual care activities	Nursing homes Cluster RCT 69 nursing homes PLWD	Probably dementia, 9% moderate, 60% moderately severe, 21% severe CDR 1 or higher; moderate degree of agitation	N=847; 404 intervention, 443 control Age: 88 (mean) % Female: 69% Race: NR Education: NR	None	NA	NA	NR	NR	9 months	DEMQOL (proxy CMAI NPI Serious adverse events Cost differences	NR
van de Ven, 2013 ⁷⁷ 23844003 Netherlands Medium Pragmatic	Dementia care mapping, training staff to be certified dementia-care mappers, care action planning	Usual care activities	Nursing homes Cluster RCT 14 care homes PLWD	Dementia, Diagnostic and Statistical Manual of mental disorders – IV criteria	N=268; 102 dementia care mapping, 166 control Age: 84 (mean) % Female: 74% Race: 97% born in Netherlands Education: NR	None	NA	NA	NA	NA	8 months	CMAI NPI QUALID EuroQoL 5D	GHQ-12
Chenoweth, 2009 ⁷⁹ 19282246 (104505176) ⁹⁸ (22078076) ⁹⁹ Australia Medium Balanced	I1: person-centered care individualized care program; I2: dementia care mapping	Usual care activities	Nursing homes Cluster RCT 15 care sites PLWD	Medical diagnosis of dementia, unspecified	N=289; 109 dementia care mapping, 98 person-centered care, 82 control Age: 84 (mean) % Female: 77% Race: 69% English speakers Education: NR	Resident classification	NA	NA	NA	NA	8 months	CMAI QUALID NPI	NR
Eloniemi-Sulkava, 2009 ⁸⁰ (20121986) Finland Medium Explanatory	Care coordination; coordinator embedded in multidisciplinary team, includes peer support groups, psychoeducation In-person	Usual care	Community-based RCT Dyads	Community-dwelling PLWD; 87% AD, predominately mild to moderate severity	N=125; 63 intervention, 62 control Age: 78 (mean) % Female: 77% Race: NR Education: NR	None	N=125; 63 intervention, 62 control Age: 75 (mean) % Female: 72% Race: NR Education: 28% less than 8 years	None	NA	NA	20 to 24 months (phased recruitment)	Time to institutionalization Use of services Service expense	NA
Rovner, 1996 ⁸² 8537594 US Medium Explanatory	Collaborative dementia care program Unclear delivery person PLWD In-person See Appendix Table F.37	Usual care	Urban Nursing Home RCT 89 PLWD	Degenerative or multi-infarct dementia with behavior disorder Diagnostic and Statistical Manual of Mental Disorders, 3 rd edition	N=81; 42 intervention, 39 control Age: 81 (mean) % Female: 77% Race: 83% White Education: NR	None	NA	NA	NA	NA	6 months	No useable outcomes	NA

*High risk of bias studies included in evidence map

Abbreviations: AD=Alzheimer's disease; ADL=Activities of Daily Living; CBT=Cognitive Behavior Therapy; CDR=Clinical Dementia Rating; CESD=Center for Epidemiologic Studies Depression Scale; char=characteristics; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale of Depression in Dementia; char=Characteristics; DEMQOL=Dementia Quality of Life; CG= Caregiver Quality of Life; char=characteristics; CMAI= Cohen Mansfield Agitation Inventory; EUROQOL-5D=European Quality of Life-5 Dimensions; FC=formal caregiver; HUI=Health utilities index; IC=informal caregiver; N=number; NA=Not Applicable; NPI=Neuropsychiatric Inventory; NR=Not Reported; PHQ-9=Patient health questionnaire-9 items; PMID=PubMed Identification Number; PLWD=Persons with Dementia; QoL=Quality of life; QUALID=Quality of life in late-stage dementia; RCT=Randomized Controlled Trials; RMBPC=Revised memory and behavior problem checklist; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; TIME=Targeted interdisciplinary model for evaluation; WHELD=Well-being and Health for People with Dementia; ZBI=Zarit burden inventory

Table F-35. Intervention details for multidisciplinary integrated team collaborative care

Study (PMID)	Team	Contact Frequency	Delivery	Caregiver Elements	PLWD Elements	Colocation	Other Supports
Possin, 2019 ⁶³ 31566651 HCB Care Ecosystem	Care team navigator (unlicensed; some bilingual Spanish, Cantonese) APN, SW, pharmacist; nurse-supervisor	Monthly by phone; email mail also; CG-initiated for problems	Telephone/ internet, some in-person	Screen for problems, Standardized education; personalized support, manualized; ACP	Medication review, monitor health status	Yes	None
Thyrian, 2017 ⁶⁵ 28746708	Nursing care coordinator embedded in German health systems; general physician and social service professionals; initial weekly meeting with nursing scientist, neurologist/ psychiatrist, psychologist, pharmacist for initial assessment	Monthly 1 hour home visits	In-home	Included in PLWD contacts, if the PLWD had an informal caregiver	Initial assessment, intervention task list	Unclear	Intervention management software system
Chodosh, 2015 ⁶⁶ 25656074 HCB ACCESS	Bilingual social work care managers from health system; in-person included care manager from local Association; In-person included care manager from local Association; Both used local resources from participating organizations	Minimum 7 contacts, average 31 contacts in telephone arm, 22 in in-person arm	In-person vs telephone/ internet	Protocolized problem assessment, self-management counseling, education, referrals and follow-up	Unclear	Unclear. Care manger within County health services with primary care clinics	Help cover cost of phone minutes for informal caregivers; Steering committee for cross-organization collaboration
Bass, 2013 ⁶⁸ 23869899 VA Based on Partners in Dementia Care	Care coordinators, embedded in VA system	Minimum 1 contact per month	Telephone/ internet, mail	Protocolized initial assessment, action plan, ongoing monitoring and reassessment	None noted	Yes	Software system
Vickrey, 2006 ⁷¹ 17116916 HCB ACCESS	Social work care managers from 3 health care organizations, 3 local community agencies could also have care managers; supervisor unclear	Frequency based on as-needed; 6 month reassessments	Telephone	Protocolized problem assessment, self-management counseling, education, referrals and follow-up	Primary care provider training	Unclear	Software systems linked; Steering committee for cross-organization collaboration
Callahan, 2006 ⁷⁰ 16684985 HCB	Geriatric nurse practitioner care manager, Primary care physician; geriatrician, geriatric psychiatrist, psychologist	Bi-weekly initially, then monthly, up to 12 months	In-person	Manualized psychoeducation, coping skills, group sessions, legal/financial advice, exercise guidelines, printed caregiver guide,	Medication, problem assessments, non-drug behavioral interventions; PLWD group exercise sessions	Unclear	Software tracking system
Rovner, 1996 ⁸² 8537594 NH	Unclear; day program created within nursing home; Psychiatrist, Creative arts therapist, nursing aides	Daily	In-person	None	Activity program, medication management, team consulting	Unclear	None

Abbreviations: ACP= advance care practice; APN=Advanced practice nurse; CG= caregiver; HCB= Home and community-based; NH= Nursing Home; PMID=PubMed Identification Number; PLWD= Person With Dementia; SW= Social Worker; VA= Veterans Affairs

Table F-36. Modified PRECIS-2: multidisciplinary integrated team collaborative care

Study (PMID)	Eligibility Criteria	Recruitment Path	Setting	Intervention Organization	Flexibility of Delivery of Intervention	Flexibility of Adherence to Intervention	Followup	Primary Outcome	Analysis	Overall Category
										Applicability and/or Qualifiers
Possin, 2019 ⁶³ 31566651	Mostly Explanatory	Mostly Explanatory	Pragmatic	Pragmatic	Mostly Pragmatic	Mostly Pragmatic	Pragmatic to disease management	Mostly Pragmatic	Mostly Pragmatic	Pragmatic
Thyrian, 2017 ⁶⁵ 28746708	Mostly Pragmatic	Balanced	Pragmatic	Mostly Pragmatic	Mostly Pragmatic	Mostly Pragmatic	Mostly Pragmatic	Mostly Pragmatic	Some reporting issues	Pragmatic
Chodosh, 2015 ⁶⁶ 25656074	Balanced	Mostly Explanatory	Pragmatic	Balanced	Mostly Pragmatic	Mostly Pragmatic	Pragmatic to disease management	Mostly Pragmatic	Balanced	Balanced
Bass, 2013 ⁶⁸ 23869899	Pragmatic	Unclear	Pragmatic	Balanced	Unclear	Unclear	Pragmatic	Mostly Pragmatic	Some reporting issues	Pragmatic
Vickrey, 2006 ⁷¹ 17116916	Pragmatic	Mostly Pragmatic	Pragmatic	Pragmatic	Mostly Pragmatic	Mostly Pragmatic	Pragmatic to disease management	Mostly Pragmatic	Mostly Pragmatic	Pragmatic
New PCC Section van de Ven, 2013 ⁷⁷	Pragmatic	Mostly Pragmatic	Pragmatic	Pragmatic	Mostly Pragmatic	Mostly Pragmatic	Pragmatic	Mostly Pragmatic	Some reporting issues	Pragmatic
Chenoweth, 2009 ⁷⁹ 19282246	X	X	Pragmatic	Mostly Explanatory	X	X	X	Mostly Explanatory	X	Explanatory

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: NA=Not Assessed; PMID=PubMed Identification Number

Table F-37. PLWD outcomes summary low and medium risk of bias studies: multidisciplinary integrated team collaborative care

Study (PMID) Comparison RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Possin, 2019 ⁶³ 31566651 Medium Pragmatic	Unstandardized Beta (CI) QoL-AD (proxy) (13-52; higher is better) 12 months	Favors intervention; 0.53 (0.25 to 1.30)	NA	NA	0.04

Possin, 2019 ⁶³ 31566651 Medium Pragmatic	Unstandardized Beta (CI) ED visits (lower is better) 12 months	Favors intervention; -0.14 (-0.29 to -0.01) NNT for single ED visit = 5 Prevented 120 visits (predicted based on based on usual care data)	NA	NA	0.04
Possin, 2019 ⁶³ 31566651 Medium Pragmatic	Unstandardized Beta (CI) Hospitalization (lower is better) 12 months	No statistical difference; -0.03 (-0.18 to 0.12) Prevented 13 visits (predicted based on based on usual care data)	NA	NA	0.71
Possin, 2019 ⁶³ 31566651 Medium Pragmatic	Unstandardized Beta (CI) Ambulance use (lower is better) 12 months	No statistical difference; -0.10 (-0.23 to 0.03) Prevented 16 uses (predicted based on based on usual care data)	NA	NA	0.12
Possin, 2019 ⁶³ 31566651 Medium Pragmatic	Mean cost estimates Based on national estimates 12 months	\$600 per PLWD across ED, ambulance, and hospitalization, over latter 6 months.	NA	NA	NA
Thyrian, 2017 ⁶⁵ 28746708	Adjusted mean difference (CI) QoL-AD (13-52; higher is better) 12 months	No statistical difference 0.02 (-0.09 to 0.05) Effect size 0.07	NA	NA	0.26
Thyrian, 2017 ⁶⁵ 28746708	Adjusted mean difference (CI) NPI 12 months	High risk of bias for missing data 36% of completers	NA	NA	NA
Thyrian, 2017 ⁶⁵ 28746708	Daily living activities	No statistical difference, no data reported	NA	NA	NA
Thyrian, 2017 ⁶⁵ 28746708	Institutionalization	No statistical difference, no data reported	NA	NA	NA
Chodosh, 2015 ⁶⁶ 25656074 Medium Balanced	Adjusted mean difference (CI) HUI (proxy) (lower is better) 6 month Grouped with QoL	No statistical difference; -0.064 (0.14 to 0.03)	NA	NA	0.19
Chodosh, 2015 ⁶⁶ 25656074 Medium Balanced	19 quality indicators	All 19 indicators improved in both arms; Comparable proportional increases in quality indicators for program, ACCESS (original protocol study) vs this study (Table 6, original publication)	NA	NA	NA
Eloniemi-Sulkava, 2009 ⁹⁰ (20121986) Medium Balanced	Time to nursing home placement	Favors intervention at 1.6 years, but no longer statistically significant at 2 years; Hazard ration 0.53 CI (0.23 to 1.19)	1.6 years 25.8%	1.6 years 11.1%	0.03
Vickrey, 2006 ⁷¹ 17116916 Medium Pragmatic	23 quality indicators	All but 2 showed significant improvement.	NA	NA	≤0.013
Vickrey, 2006 ⁷¹ 17116916 Medium Pragmatic	Adjusted mean difference (CI) HUI 18 month Grouped with QoL	Favors intervention; 0.06 (0.005 to 0.11) (MID 0.03)	NA	NA	0.034
Callahan, 2006 ⁷⁰ 16684985 Medium Explanatory	Adjusted mean difference (CI) NPI (lower is better) 12 months	Favors intervention -5.6 (-9.9 to -1.3)	8.0 (12.0)	16.1 (19.4)	0.01
Callahan, 2006 ⁷⁰ 16684985 Medium Explanatory	Adjusted mean difference (CI) CSDD (lower is better) 12 months	No statistical significance -1.0 (-2.6 to 1.5)	3.5 (3.9)	5.8 (5.9)	0.65
Callahan, 2006 ⁷⁰ 16684985 Medium Explanatory	Adjusted mean difference (CI) ADL (Higher is better) 12 months	No statistical significance 1.4 (-2.3 to 5.1)	48.6 (17.7)	44.6 (17.0)	0.44
Callahan, 2006 ⁷⁰ 16684985 Medium Explanatory	Nursing home placement 12 months	No statistical significance	6%	1.5%	0.22
Lichtwarck, 2018 ⁷⁴	Standardized mean difference NPI-NH-10 12 weeks	Favors intervention 0.25	31.1 (26.7 to 35.6)	41.4 (37.3 to 45.5)	0.053
Lichtwarck, 2018 ⁷⁴	Standardized mean difference CMAI 12 weeks	Favors intervention 0.29	59.4 (55.2 to 63.6)	67.1 (63.3 to 70.9)	0.006
Lichtwarck, 2018 ⁷⁴	Standardized mean difference CSDD 12 weeks	Favors intervention 0.26	10.2 (8.7 to 11.7)	12.4 (10.9 to 13.8)	0.010
Lichtwarck, 2018 ⁷⁴	Standardized mean difference QUALID 12 weeks	Favors intervention 0.17	27.2 (25.3 to 29.1)	29.6 (27.8 to 31.5)	0.044
Ballard, 2018 ⁷³ 29408901	Mean difference (CI) DEMQOL (proxy) 9 months	Favors Intervention, 2.54 (0.81 to 4.28) Cohen's D 0.24; NNT 9			0.004
Ballard, 2018 ⁷³ 29408901	Mean difference (CI) CMAI 9 months	Favors Intervention, 4.27 (-7.39 to -1.15) Cohen's D 0.23; NNT 6			0.008
Ballard, 2018 ⁷³ 29408901	Mean difference (CI) NPI - NH 9 months	Favors Intervention, 4.55 (-7.07 to -2.02) Cohen's D 0.30; NNT 9			<0.001

Ballard, 2018 ⁷³ 29408901	Serious adverse events 9 months	Balanced between groups reported	291	258	
---	------------------------------------	----------------------------------	-----	-----	--

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: ADL=Activities of Daily Living; CI=Confidence Interval; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; DEMQOL=Dementia Quality of Life; ED=Emergency Department; HUI=Health utilities index; N=Number; NPI=Neuropsychiatric Inventory; NPI-NH= Neuropsychiatric Inventory-Nursing Home; QoL-AD=Quality of Life-Alzheimer's disease; QUALID=Quality of Life in Late-stage Dementia; PMID=PubMed Identification Number; ROB=Risk of Bias

Table F-38. Caregiver outcomes summary low and medium risk of bias studies: multidisciplinary integrated team collaborative care

Study (PMID) Comparison Caregiver Type RoB Category*	Outcome Timing	Summary Finding	Intervention	Comparator	p-Value
Possin, 2019 ⁶³ 31566651 Medium Pragmatic	Unstandardized Beta (CI) PHQ-9 (0-27; lower is better) 12 months	Favors intervention; -1.14 (-2.15 to -0.13) NNT=12	NA	NA	0.03
Possin, 2019 ⁶³ 31566651 Medium Pragmatic	Unstandardized Beta (CI) Self-efficacy (higher is better) 12 months	No statistical difference; 0.64 (0.14 to 1.41)	NA	NA	0.11
Possin, 2019 ⁶³ 31566651 Medium Pragmatic	Unstandardized Beta (CI) 12-item ZBI (0-48; lower is better) 12 months	Favors intervention; -1.90 (-3.89 to -0.08)	NA	NA	0.046
Thyrian, 2017 ⁶⁵ 28746708	Adjusted mean difference (CI) Berlin Inventory for Caregiver burden 12 months	Favors intervention; -0.50 (-1.09 to 0.08), one-sided analysis Effect size -0.18	NA	NA	0.045
Bass, 2013 ⁶⁸ 23869899 Medium Pragmatic	Unstandardized B, no CI provided Unmet needs (lower is better, 0-39 items) 6 months	Favors intervention; -2.24	NA	NA	0.01
Bass, 2013 ⁶⁸ 23869899 Medium Pragmatic	Unstandardized B, no CI provided Role captivity (lower is better, 0-12) 6 months	No statistical difference; 0.12	NA	NA	NR
Bass, 2013 ⁶⁸ 23869899 Medium Pragmatic	Unstandardized B, no CI provided Physical health strain (lower is better, 0-12) 6 months	No statistical difference; -0.14	NA	NA	NR
Bass, 2013 ⁶⁸ 23869899 Medium Pragmatic	Unstandardized B, no CI provided Relationship strain (lower is better, 0-24) 6 months	No statistical difference; 0.38	NA	NA	NR
Bass, 2013 ⁶⁸ 23869899 Medium Pragmatic	Unstandardized B, no CI provided 11- item CESD (lower is better, 11-33) 6 months	No statistical difference; -0.69 (above from table, text reported beta -0.08, p=0.047)	NA	NA	NR
Bass, 2013 ⁶⁸ 23869899 Medium Pragmatic	Unstandardized B, no CI provided Number of informal helpers (higher is better) 6 months	No statistical difference; -0.32	NA	NA	NR
Bass, 2013 ⁶⁸ 23869899 Medium Pragmatic	Unstandardized B, no CI provided Caregiver support service use (higher is better) 6 months	Favors intervention; 0.20 (above from table, text reported beta 0.14, p=0.008)	NA	NA	NR
Chodosh, 2015 ⁶⁶ 25656074 Medium Balanced	Adjusted mean difference (CI) ZBI (lower is better) 6 month	No statistical difference; 1.8 (-3.46 to 7.05)	NA	NA	0.5
Chodosh, 2015 ⁶⁶ 25656074 Medium Balanced	Adjusted mean difference (CI) PHQ-9 (lower is better) 6 month	No statistical difference; 0.36 (-2.00 to 2.73)	NA	NA	0.76
Vickrey, 2006 ⁷¹ 17116916 Medium Pragmatic	Adjusted mean difference (CI) EuroQol-5D 18 month	No statistical difference; 0.02 (-0.01 to 0.06)	NA	NA	0.127
Vickrey, 2006 ⁷¹ 17116916 Medium Pragmatic	Adjusted mean difference (CI) Caregiving-attributable health strain 18 month	No statistical difference; 4.3 (-0.3 to 8.8)	NA	NA	0.063
Callahan, 2006 ⁷⁰ 16684985 Medium Explanatory	Adjusted mean difference (CI) PHQ-9 (lower is better) 12 month	No statistical difference; -0.9 (-2.2 to 0.5)	3.1 (3.9)	4.6 (5.6)	0.50
Callahan, 2006 ⁷⁰ 16684985 Medium Explanatory	Adjusted mean difference (CI) Caregiver NPI (lower is better) 12 month	Favors intervention; -2.2 (-4.2 to -0.2)	3.5 (5.8)	7.7 (8.7)	0.03

*Explanatory studies are listed first, followed by pragmatic studies.

Abbreviations: CI=Confidence Interval; CESD=Center for Epidemiologic Studies Depression Scale; EuroQol-5D=European Quality of Life-5 Dimensions; NA=Not Applicable; NNT=Number Needed to Treat; NPI=Neuropsychiatric Inventory; PMID=PubMed Identification Number; RoB=Risk of Bias; ZBI=Zarit Burden Interview

Table F-39. Summary of strength of evidence for PLWD outcomes: multidisciplinary integrated team collaborative care

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Collaborative care vs usual care QoL	6-12 months	4 Pragmatic trials (n=1,746)	QoL-AD benefit, no benefit HUI benefit, no benefit	Medium	Consistent	Direct	Imprecise	Low (Weighted to larger, quality pragmatic trials)
Collaborative care vs usual care NPI	6-12 months	1 Explanatory Cluster Trial (N=152)	One benefit	Medium	Unknown	Direct	Imprecise	Insufficient
Collaborative care vs usual care Utilization	6-12 months	1 Pragmatic Trial (n=780)	ED visits benefit, hospitalization no difference, ambulance use no difference	Medium	Unknown	Direct	precise	Low
Collaborative care vs usual care ADL	6-12 months	1 Pragmatic 1 Explanatory Cluster Trial (n=560)	2 no difference	Medium	Consistent	Direct	Imprecise	Insufficient
Collaborative care vs usual care Quality indicators	6-12 months	2 Pragmatic (n=559)	Quality indicators, 2 benefit	Medium	Consistent	Direct	Imprecise	Low (Weighted to larger, quality pragmatic trials)
Collaborative care vs usual care Depression	6-12 months	1 Explanatory Cluster Trial (N=152)	CSDD, no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Collaborative care vs usual care Nursing home placement	6-12 months	1 Pragmatic 1 Explanatory Cluster Trial (n=560)	2 no difference (not long enough for balance toward mild/moderate)	Medium	Consistent	Direct	Imprecise	Insufficient

Abbreviations: ADL=Activities of Daily Living; CSDD=Cornell Scale for Depression in Dementia; ED=Emergency Department; HUI=Health utilities index; N=Number; NPI=Neuropsychiatric Inventory; QoL=Quality of Life; QoL-AD=Quality of Life-Alzheimer's disease

Table F-40. Summary of strength of evidence for caregiver outcomes: multidisciplinary integrated team collaborative care

Comparison Caregiver Type Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Collaborative care vs usual care Informal QoL	6-12 months	1 Pragmatic Trial (n=408)	EuroQoL-5D no difference	Medium	Unknown	Direct	Imprecise	Insufficient
Collaborative care vs usual care Informal Self-efficacy	6-12 months	1 Pragmatic Trial (n=780)	No benefit	Medium	Unknown	Direct	Imprecise	Insufficient
Collaborative care vs usual care Informal Burden	6-12 months	4 Pragmatic (n=1,719)	ZBI 2 Benefit, 1 Berlin no benefit; Role captivity, 2 physical health strain, relationship strain, all no benefit; Caregiver NPI no difference	Medium	Inconsistent	Direct	Imprecise	Insufficient (Weighted to larger, quality pragmatic trials)
Collaborative care vs usual care Informal Quality	6-12 months	1 Pragmatic Trial (n=486)	Unmet needs, informal helpers, support service use 2 of 3 no difference	Medium	Consistent	Direct	Imprecise	Insufficient (Weighted to larger, quality pragmatic trials)
Collaborative care vs usual care Informal Depression	6-12 months	3 Pragmatic, 1 Explanatory (n=1,570)	1 benefit, 3 No difference	Medium	Consistent	Direct	Imprecise	Insufficient (Weighted to larger, quality pragmatic trials)

Abbreviations: EuroQoL-5D=N=Number; NPI=Neuropsychiatric Inventory; QoL=Quality of Life; ZBI=Zarit Burden Index

Evidence Map: Multidisciplinary Integrated Team Collaborative Care

Table F-41. Characteristics of evidence map studies: multidisciplinary integrated team collaborative care

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Froggat 2020 ¹⁰⁰ UK Pilot	Namaste: structured personalized care	Usual care activities	Nursing home	Advanced dementia	N=32 Age: Yes Sex: Yes Race: no Education: No	None	NA	NA	NA	NA	24 weeks	QUALID Comfort in dying	NA

Halek 2020 ⁶² (32062052) Germany High ROB	Two dementia-specific Case Conference models	Usual care	Nursing homes Cluster not specified N=413	Documented diagnosed dementia from nursing charts, FAST score > 1, living at least 15 days in the unit	N=224 Age: yes Sex: yes	Low	NA	NA	N=189	Position: yes	15 days/unclear		Burnout risk (CBI)
Chen, 2019 ¹⁰¹ China Pilot	Interdisciplinary care team	Usual care activities	Community based, China	Not specified	N=148 Age: Yes Sex: Yes Race: no Education: No	None	None	None	NA	NA	6 month	CDR QoL ADL NPI	ZBI
Saxena, 2018 ¹⁰² 29193721 Singapore Unable to evaluate	Primary care Dementia Clinic (PCDC): Each visit consists of a 15-min consultation with FP & 15 min consultation with a nurse	(1) Standard primary care at other National healthcare Group polyclinic (2) Specialist care at the MC	Clinic Quasi 263 PLWD	Not specified	N=263 Age: Yes Sex: Yes Race: Yes Education: No	Marital status	None	None	NA	NA	12 months		Quality of life: QoL-AD CG satisfaction: Cost effectiveness:
Mavandadi, 2017 ¹⁰³ 28134558 US Pilot	Dementia care management: Telephone-based patient- and caregiver (CG)-centered, collaborative care management program that involves CG education and psychosocial support	Usual care	Community RCT 75 Caregivers	All types dementia and stages of severity	N=75 Age: Yes Sex: no Race: no Education: no	None	N=75 Age: Yes Sex: yes Race: yes Education: no Relationship: yes	Informal caregivers of veterans	NA	NA	6 months	Activities of daily living: ADL; IADL	Bother: RMBPC Distress: NPI-Q, Burden: ZBI Cognitive coping: Perceived mastery over caregiving:
Laporte Uribe 2017 ⁶⁴ (28249632) Germany High RoB	PLWD and informal CG using DCN	None	Community-dwelling Quasi-experimental 560 PLWD and IC	Dementia diagnosis by medical professionals and living with IC using DCN	N=389 Age: Yes Sex: Yes Race: no Education: no Living with CG: Yes	None	N=385 Age: Yes Sex: Yes Race: no Education: no Relationship: Yes				12 months	CMAI IADL EQ-5D L	BIZA-D
Samus, 2014 ¹⁰⁴ US Pilot	Home-based care coordination: to systematically identify & address dementia related care needs- Received the written results of the JHDCNA & 18 months of care coordination	Augmented usual care: Received the written results of the JHDCNA following the BL visit including recommendations for each identified unmet need.	Community RCT 303 PLWD	Type of dementia: not specified Severity: Mild – severe	N=303 Age: Yes Sex: Yes Race: yes Education: Yes Living with CG: Yes	NA	NA	NA	NA	NA	18 months	Time to transfer from home: Study partner report Unmet care needs: JHDC NA	None
Leontjevas, 2013 ¹⁰⁵ 23643110 Netherland High ROB	Act in Case of Depression (AiD) at dementia unit: A multidisciplinary care program that involved 2-step screening & diagnostic procedure; multidiscipline treatment; & monitoring of treatment effects.	AiD at somatic unit:	Nursing home RCT (cluster) 793 PLWD		N=793 Age: Yes Sex: Yes Race: no Education: no	NA	NA	NA	NA	NA	?	Depression: CSDD	None

Chapman, 2007 ⁶⁹ 18232242 US High ROB	Advanced illness care teams (AICTs): holistic approach that focused on four domains: (1) medical, (2) meaningful activities, (3) psychological, and (4) behavioral	Usual care; received all the services typically provided by the facility	Nursing homes RCT 118 NH residents	Advance dementia	N=118 Age: Yes Sex: Yes Race: Yes Education: no Marital status: Yes	SES: Yes	NA	NA	NA	NA	8 weeks	Agitated behavior:CM AI Pain:FLACC Depression: CSDD	None
Bass, 2003 ¹⁰⁶ 12604748 14690867 ¹⁰⁷ US Pilot/ demonstration	Care coordination integrated within health system; average 12 contacts per year	Usual care activities	Community-based RCT 157 Dyads	Dementia diagnosis in health records	N=157 Age: no Sex: no Race: no Education: no Marital status: no	Lives alone	NA	NA	NA	NA	12 month	Utilization	IC: CES-D Caregiver strain
Person-Centered Care	And Dementia Care Mapping												
Surr, 2019 ⁷² 31056923 UK High ROB	Dementia care mapping to reduce agitation in nursing homes	Usual care activities	Dementia care homes Cluster RCT 50 Care homes	Care home residents	N=726 Age: yes Sex: Yes Race: no Education: no Marital status: no	None	NA	NA	Information at care home level	NA	16 months	CMAI Resident behaviors QoL	System-level outcomes: Quality of interactions schedule
Li, 2017 ¹⁰⁸ US Pilot	Person-centered dementia care: included in-class staff training plus supervision and support in practice of individualized care	Control : no intervention	Assisted Living facility RCT 28 PLWDs	All types of dementia except those with Parkinson & in severe stage	N=22 Age: yes Sex: Yes Race: Yes Education: no Marital status: Yes	None	NA	NA	NA	NA	3 months 1 month post intervention follow up	Sleep-wake pattern: Actigraphy social engagement: DCM	None
Ballard, 2016 ¹⁰⁹ 26585409 UK Pilot	Person centered care combined with medication review, social interaction, and exercise		Nursing homes Cluster randomized factorial analysis 16 homes	Dementia residents	N=187 Age: yes Sex: Yes Race: Yes Education: no Marital status: no	None	NA	NA	NA	NA	9 months	DEMQOL-proxy antipsychotic use	None
Moyle, 2016 ¹¹⁰ Australia Pilot	Capabilities model based on a person-centered approach	Usual care activities	Nursing home Quasi-experimental	Confirmed dementia diagnosis	N=48 Age: no Sex: no Race: no Education: no Marital status: no	None	N=75 Age: Yes Sex: yes Race: no Education: yes Relationship: yes	Employment	N=81 Age: Yes Sex: Yes Race: no Education: Yes	Position	6, 12 months	QoL-AD	Staff experience of working with demented residents questionnaire
Dichter, 2015 ⁷⁵ 26138674 Germany High ROB	Dementia Care Mapping: (A) DCM applied since 2009: received two DCM cycles per year until 2009. (B) DCM newly introduced during the intervention (C) received an intervention based on a regular and standardized QoL rating that was integrated into the usual care.		Nursing home Quasi 315 PLWD	Types of dementia: Not specified	N=154 Age: yes Sex: Yes Race: No Education: no	None	NA	NA	NA	NA	18 months	Quality of life: QoL-AD Challenging behavior: NPI-NH Functional ability: PSMS	None

Rokstad, 2013 ⁷⁸ 24022375 Norway High ROB	(1) Dementia Care Mapping (DCM): a process to develop the care staff's skills in delivering PCC to the patients (2) VPM: a weekly consensus meeting in the nursing home ward of 45–60 min using the indicators in the VIPS framework to analyze a challenging patient-nurse interaction.	Control: Education on dementia: received five DVDs with lectures (30 min each) about dementia.	Nursing home RCT 624 PLWD	All types and stages of dementia	N=624 Age: yes Sex: Yes Race: No Education: no	General physical health: yes	NA	NA	NA	NA	10 months	Agitation: BARS NPIQ: Quality of life: QUALID	NA
Fossey, 2006 ⁸¹ 16543297 UK High ROB	A training & support intervention: intervention delivered to nursing home staff over 10 months, focusing on alternatives to drugs for the management of agitated behavior in dementia	Treatment as usual	Nursing homes RCT (cluster) 349 PLWD	Type of dementia unspecified Mild to severe	N=349 Age: yes Sex: Yes Race: No Education: no	None	NA	NA	NA	NA	12 months	Agitated behavior:	NA
Chenoweth, 2014 ⁷⁶ 24666667 Australia High ROB	(1) A person-centered care (PCC) (2) Person centered environment (PCE) (3) PCE +PCC	Usual care + usual environment	Aged care homes RCT (Cluster) 601 PLWD	Only AD & severe/very severe were specified in %	N=601 Age: yes Sex: Yes Race: No Education: no	None	NA	NA	NA	NA	4 months 8 months follow up (4 months post intervention)	Quality of life: DEMQOL Agitation: CMAI Cost of program:	NA

Abbreviations: ADL=Activities of Daily Living; AICT=Advanced illness care teams; AiD=Act in Case of Depression; BARS=Brief Agitation Rating Scale; BIZA-D= Berlin Inventory of Caregivers' Burden with of Dementia Patients; CDR=Clinical Dementia Rating; CESD= Center for Epidemiologic Studies Depression Scale CG=Caregiver; char=Characteristics; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale of Depression in Dementia; DCM=Dementia Care Mapping; DEMQOL=Dementia Quality of Life; DVD=Digital Video Disc; EQ-5D L=Quality of Life Instrument; FLACC=Face Legs Activity Cry Consolability; FP=Family Physicians; IADL=Instrumental Activities of Daily Living; JHDCNA=Johns Hopkins Dementia Care Needs Assessment; NH=Nursing home; NPI=Neuropsychiatric Inventory; NPI-NH= Neuropsychiatric Inventory-Nursing Home; NPI-Q= Neuropsychiatric Inventory Questionnaire; MC=Memory Clinic; min=Minutes; PCDC=Primary care Dementia Clinic; PCC=Person Centered Center; PCE=Person Centered Environment; PSMS= Physical Self-Maintenance Scale; PLWD=Person with Dementia; QoL-AD=Quality of Life-Alzheimer's Disease; QoL=Quality of Life; QUALID=Quality of Life in Late-Stage Dementia; RCT=Randomized Controlled Trial; RMBPC=Revised memory and behavior problem checklist; RoB=Risk of Bias; SES=Socioeconomic Status; VPM=VIPS practice model; ZBI=Zarit Burden Index

Formal Caregiver Training

Table F-42. Risk of bias assessment: informal caregiver training

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Gozalo, 2014 ¹¹¹ (24697702)	2 years	Medium	Low	Medium	High	Low	High	Gov't Developer	High
Schindel Martin, 2016 ¹¹² (27659392)	6 month	X	High	X	X	X	X	Gov't Developer	High
Galik, 2013 ¹¹³ (24092822)	6 months	X	Medium	X	X	X	X	X	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Formal Caregiver Training

Table F-43. Characteristics of included studies: formal caregiver training

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Random-ized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Williams, 2017 ¹¹⁴ (27048705) US Small sample	CHAT: communication training designed to alert nursing staff to elder speak communication and its negative effects	Usual care activities	Nursing Home Cluster RCT 13 Nursing homes	Patients with diagnosis of AD	PLWD N=83 Age: yes Sex: yes Race: yes Education: no	None	NA	NA	N=130 Age: yes Sex: yes Race: yes Education: yes	Training Position	3 months	Resistance to care	FC: Communication
Conway, 2016 ¹¹⁵ (26821868) Australia Small sample	MESSAGE communication intervention to facilitate patient support and staff and patient experience	Usual care activities	Nursing home Cluster RCT 12 Nursing homes	Patients with diagnosis of AD with ability to respond to direct assessment	NR	NR	NA	NA	N=38 Age: yes Sex: yes Race: no Education: yes	Training Position	3 months	Self-efficacy M-NCAS ADQ	FC: Communication
Schindel Martin, 2016 ¹¹² (27659392) Canada High ROB	Standardized dementia education training on gentle persuasion approaches	Usual education supports	Hospitals Quasi-experimental 12 clinical areas at 2 hospitals	AD diagnosis	NR	NR	NA	NA	N= 745 Age: yes Sex: yes Race: no Education: yes	Training Length of service	8 weeks	None	FC: Self-Perceived Behavioral Management Self-Efficacy Profile
Hattink, 2015 ¹¹⁶ (26519106) Netherlands & UK Pilot	Access to a Web-based portal consisting of 8 modules & to online peer and expert communities for support and information exchange	Wait-list	Community setting RCT Caregivers	AD diagnosis	NR	NR	N59 Age: yes Sex: yes Race: no Education: no IC Relation to PLWD: yes	IC Duration IC Employment status	N=24 Age: yes Sex: yes Race: no Education: no	Length of service	4 months	ADKS ADQ IRI QoL Burden: assessed with 1 question SSCQ	Usefulness of training
Gozalo, 2014 ¹¹¹ (24697702) US High ROB	Education program on bathing patients with dementia to improve bating experience of patients	Usual care activities until crossover	Nursing homes Crossover cluster RCT 6 Nursing homes	Dementia unspecified	N=240 Age: yes Sex: yes Race: yes Education: no	Health insurance Race Information	NA	NA	NR	NR	2 years	Bath duration Aggressive activity	Usefulness of BWAB in reducing aggressive patient behavior during bathing
O'Shea 2014 ¹¹⁷ (24633858) Ireland Pilot	Dementia Education Program Incorporating Reminiscence for staff facilitated by experienced nurse educators, delivered over 3 days (2 at the beginning and 1 six weeks later) and augmented by telephone support and one site visit	usual care provided by care staff who did not receive the structured education program	Long term care Cluster RCT 18 Residential units 304 PLWD	Dementia diagnosis	N=304 Age: yes Sex: yes Race: no Education: no	None	NR	NR	NR	NR	18-22 weeks	QoL-AD CSDD	QoL-AD ZBI

Galik, 2013 ¹³ (24092822) US High ROB	Function-focused care training	Usual care activities	Nursing home Cluster RCT 4 Nursing homes	Patients with MMSE <16	N=103 Age: yes Sex: yes Race: yes Education: N	Comorbidity Marriage status	NA	NA	N=77 Age: yes Sex: yes Ethnicity: yes Education: yes	NR	6 months	Barthel ADL CMAI CSDD Apathy Inventory Falls ED transfers Death	Restorative Care Behavior Checklist Knowledge Restorative Care Activities: self-efficacy and outcome expectations
Clare, 2013 ¹⁸ (22840185) UK Pilot	Training in observation of awareness in patients with severe AD	No training	Nursing homes Cluster RCT 8 Nursing homes	Participants with AD moderate to severe	N=32 residents Age: yes Sex: yes Race: yes Education: no		NA	N	N=63 Age: no Sex: yes Ethnicity: yes Qualifications: yes	Training	8 weeks	Measures of resident, family and staff quality of life Resident and staff care measures at baseline and follow-up	Resident quality of life Secondary outcome: resident well-being, behavior and cognition; staff attitudes and well-being
Ghandehari, 2013 ¹⁹ (23457681) Canada Pilot	Education program in pain management	Randomized control trial	N=131 staff Focus group of 28 participants (16 nurses, 12 aides) 29 control subjects (13 nurses, 16 aides) From 2 health care regions	Not reported	Not Reported	Not reported	NA	NA	Not reported	Training: yes	2 weeks	Evaluation of training session and workshop knowledge	Assessment of an expert-based education program on pain management
Wenborn, 2013 ²⁰ (23637069) UK Pilot	Assessment of an occupational therapy program to increase provision of activity in residential care home patients	Randomized control trial with blinded assessment	Intervention group of 8 homes, 104 residents Control group of 8 homes, 106 residents	Patients with AD with DSM_IV score less than 25 Age 60 or older, residing at residential home for at least 2 months with no other serious physical or mental health problems	N=104 intervention group 106 control group Age 84.2 intervention group 84.2 control group Sex 66% female intervention group 75% control group Race 95% white intervention group 88% white control group Education: no	SES: no Prior disability: no Household characteristics: no Health insurance: no Race information: yes	NA	NA	Not reported	Not reported	12 weeks	Measures of resident, family and staff quality of life Resident and staff care measures at baseline and follow-up	Effectiveness of training in promoting activity engagement for patients with AD
Leone, 2012 ²¹ (22700526) France Pilot	Evaluation of staff education in managing apathy in older nursing home patients with AD	Randomization of intervention group and control group by nursing home site; 16 total sites	N=119 patients, 284 caregivers in intervention group 111 patients, 279 caregivers in reference group	Patients with diagnosis of AD, per medical records information, MMSE<24 and presence of apathy per diagnostic criteria	N=119 intervention group 111 control group Age 87.83 intervention group 88.82 reference group Sex: 72.3% intervention group 87.4 reference group Race: no Education: no	Not reported	NA	NA	Not reported	Not reported	17 weeks	Measures of resident emotional blunting, interest, level of initiative	Effectiveness of staff training measures in reducing patient apathy in AD patients
McCurry 2012 (22367233) US Small sample	4 training sessions to develop and implement individualized PLWD behavioral sleep plans	Usual care	Adult family homes residents RCT 47 PLWD	Dementia diagnosis by primary care physicians with mean±SD MMSE score being 8.1±7.6 and has one or more sleep problems on the Sleep Disorders Inventory	N=31 Age yes Sex yes Race: yes Education yes	None	NA	NA	N=37 Mean Age: yes Sex: yes Race: yes Education: yes	None	1 month 6 months	Actigraphy CSDD RMBPC ESS	NA

Zimmerman, 2010 ¹²² (No Id #) US Training pilot	Evaluation of staff education program on dementia care focused on leadership, knowledge about AD and pain reduction	Randomization of care centers receiving training vs control group; 16 residential care and nursing home sites in 4 states	N=213 staff, 78 supervisors in 8 centers for intervention group 278 staff, 93 supervisors for control group	Not reported Intervention center patients with AD=62% Control center patients with AD=48%	Not reported	Not reported	NA	NA	N=490 Age: yes Sex: yes Race: yes Education: yes	Health Status :no Training: yes Education: yes Position: yes Length of Service: yes	3 months	Measures of supervisor leadership and measures of work stress, work satisfaction and knowledge	Benefit of training for staff and supervisory personnel
Deudon,2009 ¹²³ (19370714) France Training pilot	8 week, 12 session staff training on dementia and BPSD; feedback sessions		Nursing homes Cluster RCT 16 sites 306 PLWD		N=306 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	12 week post intervention	NPI CMAI Observation Scale (OS)	NA
Kuske, 2009 ¹²⁴ (19193252) Germany Training pilot	Evaluation of a staff education program on caregiver knowledge and competence; evaluation relaxation training for staff	3 arm cluster randomized control trial clustered in 6 nursing homes	N=68 patients, 89 caregivers in intervention group 68 patients, 90 caregivers in relaxation group 74 patients, 94 caregivers in control group	Patients with AD, per MMST and Barthel Index scores	N=68 intervention group 68 patients relaxation group 74 patients control group Age: yes Sex: yes Race: no Education: no	Not reported	NA	NA	Age: yes Sex: yes Race: no Education: yes	Health Status: yes Training: yes Education: yes Position: yes Length of Service: yes	6 months	Measures of knowledge, competence and emotional/health status	Benefit of training on quality of care for AD patients and in reducing caregivers' burden in caring for patients
Chang, 2005 ¹²⁵ (16238764) Taiwan Pilot	Effects of a training program on feeding skills for nursing assistants (NA) feeding patients with AD Pilot study	2 AD-specialized long term-care treatment centers randomly assigned as training or control group	One patient per NA pre and post training, not matched	Patients diagnosed with AD and evaluated by NA as having eating problems and requiring assistance	N=31 NA in training group 36 NA in control group Patient information not reported; 12 patients matched pre and post training to NA in training group; 8 patients matched in control group	Not reported	NA	NA	Age: yes Sex: yes Race: no Education: no	Health Status: no Training: no Education: no Position: yes Length of Service: yes	One post training assessment	Assessment of NA knowledge, using checklist Food intake Feeding time EdFED score	Caregiver skills and attitude developed through training Effects of caregiver training on patient eating behavior
Sloane, 2004 ¹²⁶ (15507054) US Small sample	Techniques to reduce agitation, aggression and discomfort in AD patients 1) Training for person-centered showering or 2) towel bath	Usual care activities	Nursing homes, Crossover Cluster RCT 15 Cluster N 69 PLWD 37 FC	Patients 65 and older with diagnosis of AD or related dementia who required assistance with bathing	N=69 PLWD, Age: yes Sex: yes Race: yes Education: yes	SES: no Prior disability: no Household characteristics: no Health insurance: no Race information: yes	NA	NA	N: 37 Age: yes Sex: yes Race: no Education: no	Health Status: no Training: no Education: no Position: yes Length of Service: yes	6 weeks	Bathing-associated aggression, agitation, and discomfort	Comparison of bathing behavior difficulty of non-trained staff and person-centered vs towel bath intervention
Magai, 2002 ¹²⁷ (11395344) US Pilot	Training in sensitivity to nonverbal communication, 10 – 1 hr sessions	C1: Waitlist C2: dementia education training	Nursing homes Cluster RCT 3 Cluster N	Dementia unspecified	N=91 Age: yes Sex: yes Race: yes Education: no	None	NA	NA	N=21 Age: yes Sex: yes R: yes Education: no	None	12 weeks	CDS CMAI BEHAVE-AD	BSI
Bourgeois, 2001 ¹²⁸ (No ID) US Pilot	Use of memory aids as augmentation or alternative in communicating with nursing home residents with AD	Sub-set of a larger study of the effectiveness of memory aids for communication with patients with AD Intervention and control groups	7 nursing homes; subset of 66 out of 125 residents	Patients diagnosis of AD, MMSE below 25, remaining in care facility for at least 3months	N=33PLWD, 33FC treatment group 33PLWD, 33FC control group Age: yes Sex: yes Race: yes Education: yes	SES: N Prior disability: no Household characteristics: no Health insurance: no Race information: yes	NA	NA	N: yes Age: yes Sex: yes Race: no Education: no	Health Status: no Training: no Education: no Position: yes Length of Service: no	3 months	Content of patient conversation evaluated for specific content items and qualities	Comparison of conversation between patients and staff; untrained staff vs trained staff using communication aids

Wells, 2000 ¹²⁹ (10798473) Canada Small sample	Evaluation of training program of nursing home patients with AD	Randomized control study with one experimental care unit and 3 control units	Four cognitive support units in a large geriatric care center	Patients with a diagnosis of dementia or AD residing in the memory assistance care units for at least 4 weeks; MMSE <19; Control group selected for age and MMSE equivalency with experimental group	N=20 PLWD, 16 FC experimental group 20PD, 28FC control group Age: yes Sex: yes Race: no Education: no Primary diagnosis: yes	SES: no Prior disability: no Household characteristics: no Health insurance: no Race information: no	NA	NA	N: yes Age: yes Sex: yes Race: no Education: yes	Health Status: no Training: yes Education: yes Position: yes Length of Service: yes	6 months	Measurements of resident interaction behavior, level of agitation, perceived ease of caregiving, level of stress	Level of resident interaction with caregivers with or without training
McCallion, 1999 ¹³⁰ (10568079) US	Evaluation of an education program to improve effectiveness of nursing home assistants with residents with dementia	Randomized partial crossover control study; control group given training after 6 month assessment	Two nursing homes, including 2 care units housing patients with dementia in each	Patients with diagnosis of dementia and MMSE score assessed against educational level; GDS of stage 3 or above	N=49PLWD, 39 FC treatment group 56 PLWD, 49 FC control group Age: yes Sex: yes Race: no Religion: yes Education: yes Primary diagnosis: no	SES: no Prior disability: no Household characteristics: no Health insurance: no Race information: yes	NA	NA	N: yes Age: yes Sex: yes Race: yes Education: yes	Health Status: no Training: yes Education: yes Position: yes Length of Service: yes	9 months	FC interviews, KAT, MHQ; staff turnover rate; Resident assessments of signs of and symptoms of depression and aggressive behaviors	Changes in level of behavioral disturbances in PLWD; Changes in FC knowledge and patient management
Proctor, 1999 ¹³¹ (10406361) UK Pilot	Evaluation of training and education intervention in nursing and residential care facilities	Randomized control trial	Twelve matched care facilities, selected randomly as intervention or control groups; 10 patients with difficult behavioral problems selected by staff at each facility	Patients selected by staff; Centers paired by size and accreditation status and randomly selected to intervention or control arm	N=54 PLWD, 51 PLWD control group Age: yes Sex: yes Race: no Education: no Primary diagnosis: no Counts for intervention vs control group not reported	SES: no Prior disability: no Household characteristics: no Health insurance: no Race information: no	NA	NA	Not reported	Not reported	6 months	Measurements of patient cognitive impairment, depression, behavioral disturbance and functional ability	Changes in patient depression scores, level of cognitive impairment and behavior

Abbreviations: AD=Alzheimer's Disease; ADKS=Alzheimer's Disease Knowledge Scale; ADL=Activities of Daily Living; ADQ=Approaches to Dementia Questionnaire; BEHAVE-AD=behavioral pathology in Alzheimer's disease rating scale; BSI=Brief symptom inventory; BWAB=Bathing without a Battle; CDS=Cornell depression scale; char=characteristics; CMAI=Cohen-Mansfield Agitation Inventory; CPS= Cognitive Performance Scale; CSDD=Cornell Scale for Depression in Dementia DSM-IV=Diagnostic and Statistical Manual of Mental Disorders; EdFED= Watson feeding difficulty scale for AD patients; EM=Evidence Map; ESS=Epworth Sleepiness Scale; FAST=Functional Assessment Staging; FC=formal caregiver; GDS= Global Deterioration Scale; GPA=Gentle Persuasive Approaches; IC=informal caregiver; ICD=International Classification of Diseases; IRI=Individual Reactivity Index; KAT=Knowledge Alzheimer's Test; MHQ= Penn State Mental Health Questionnaire; MMSE= Mini-mental State Examination; MMST=German version of MMSE, Mini-mental State Assessment; M-NCAS=Modified Nursing Care Assessment; MZBI=Modified Zarit Burden Interview; N=number; NA=Not Applicable; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; QOL-AD=Quality of Life - Alzheimer's Disease; RMBPC=Revised Memory and Behavior Problems Checklist; RS=Reporting Status; RTC=Restiveness to Care; SBMSEP=Self-perceived Behavioral Management Self-Efficacy Profile; SES=socioeconomic status; SSCQ=Short Sense of Competence Questionnaire; ZBI=Zarit Burden Interview

Informal Caregiver Training

Table F-44. Risk of bias assessment: informal caregiver training

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Hepburn, 2001 ¹³² (113477900)	5 months	Medium 20%	Low	Medium	High	Low	High	Gov't Developer	High
Hepburn, 2006 ¹³³ (43539927)	6 month	Medium 21%	Low	Medium	High	Low	High	Gov't Developer	High
Hepburn, 2007 ¹³⁴ (17378189)	6 months	High 49%	Medium	X	X	X	X	X	High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Table F-45. Characteristics of evidence map studies: informal caregiver training

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Random-ized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Hattink, 2015 ¹¹⁶ (26519106) Netherlands & UK Pilot	Access to a Web-based portal consisting of 8 modules & to online peer and expert communities for support and information exchange	Wait-list	Community setting RCT Caregivers 142	Dementia unspecified	N= Age: no Sex: no % majority race: no Education: no	None	IC N=59 IC Age: yes IC Sex: yes IC % majority race: no IC Education: no IC Relation to PLWD: yes	IC Duration IC Employment status	FC N=24 FC Age: yes FC Sex: yes FC Race: no FC Education: no	Length of service	4 months	ADKS ADQ IRI QoL SSCQ	FC: Usefulness of training IC: Burden: assessed with 1 question
Liddle, 2012 ¹³⁵ (23092595) Australia Small sample	DVD-based training for memory and communication support skills	No training	Community RCT Caregivers	Dementia (Alzheimers, vascular, frontotemporal, not otherwise specified)	N=29 Age: yes Sex: yes % majority race: no Education: yes	None	N=29 Age: yes Sex: yes % majority race: no Education: yes Relationship: yes	None	NA	NA	3 months	None	ZBI PAC RMBPC CSDD
Klodnicka Kouri, 2011 ¹³⁶ ?? Canada Pilot	Individual communication training, 5 weekly 90-120 minute session	Printed information	Community RCT Caregivers	Mild probable Alzheimer's disease	N= Age: no Sex: no % majority race: no Education: no	None	N=50 Age: no Sex: yes % majority race: yes Education: no Relationship: yes	Duration; Living with PLWD; Employment	NA	NA	6 weeks post intervention	None	IC: Caregiver self-efficacy scale; RMBPC communication knowledge and skills
Neely, 2009 ¹³⁷ (19294562) Sweden Small sample	Trained in and practiced strategies to support everyday mnemonic and occupational performance, home-based, 8 – 1 hour sessions	No training	Community RCT Dyad	Mild to moderate Alzheimer's disease or with vascular dementia	N=30 Age: yes Sex: yes % majority race: no Education: no	None	N=30 Age: yes Sex: yes % majority race: no Education: no Relationship: no	None	NA	NA	8 weeks	None	ZBI Beck depression
Hepburn, 2007 ¹³⁴ (17378189) USA High ROB	Group caregiving role-training and mastery-focused coaching 6 - 2 hour sessions	Wait-list control	Community RCT Caregivers	Dementia unspecified	N= Age: no Sex: no % majority race: no Education: no	None	N=102 Age: yes Sex: yes % majority race: yes Education: yes Relationship: yes	SES Duration	NA	NA	5 months	None	Relational deprivation Role captivity Competence Mastery Loss of self-Distress
Hepburn, 2005 ¹³³ ?? USA High ROB	Group caregiving role-training and mastery-focused coaching 6 weekly 2 hour sessions	Wait-list control	Community RCT NA NA	Dementia unspecified	N=215 Age: NR Sex: NR % majority race: NR Education: NR	NA	IC N=215 IC Age: yes IC Sex: yes IC % majority race: no IC Education: yes IC Relation to PLWD: no	None	NA	NA	6 and 12 months	None	BACS remainder were development of distress measure
Martin-Cook, 2005 Martin-Cook, 2005 #15808} (16136843) US Small sample	Individual communication training through modeling and feedback, 4 sessions	Not specified	Community RCT Caregivers	Mild to moderate dementia, primarily Alzheimer's disease	N= Age: yes Sex: yes % majority race: yes Education: yes	None	N=47 Age: no Sex: yes % majority race: no Education: yes Relationship: yes	Duration	NA	NA	17 weeks	ILS/I-ILS ADCS MMSE; NPI; GDS	FMTCS; GSE
Done, 2001 ¹³⁸ (11536349) UK Small sample	Group communication training 1 hour/week, 2 weeks	Information booklet	Community Cluster RCT Local groups (day center or local home) 45 Caregivers	Dementia unspecified	N= Age: no Sex: no % majority race: no Education: no	None	N=45 Age: no Sex: no % majority race: no Education: no Relationship: no	None	NA	NA	6 weeks	None	IC: The relatives stress scale TACI

Gormley, 2001 ¹³⁹ (11395344) UK High ROB	Brief behavior management training program, 4 home sessions	Attention control: sessions about general care questions	Community RCT Caregivers	Dementia unspecified	N= Age: yes Sex: yes % majority race: no Education: no	None	N=62 Age: yes Sex: yes % majority race: no Education: no Relationship: no	None	NA	NA	8 weeks	BEHAVE-AD; MMSE; Blessed Dementia Rating Scale; Zarit Burden Interview	None
Hepburn, 2001 ¹³² (11347790) USA High ROB	Group caregiving role-training and mastery-focused coaching 7 weekly 2 hour sessions	Wait-list control	Community RCT Caregivers	Dementia unspecified	N=117 Age: NR Sex: NR % majority race: NR Education: NR	None	IC N=117 IC Age: yes IC Sex: yes IC % majority race: yes IC Education: yes IC Relation to PLWD: yes	IC: SES	NA	NA	5 months	None	BACS Revised Zarit burden scale CESD
Burgener, 1998 ¹⁴⁰ (9708136) US Pilot	Individual education I1: dementia education only I2: behavioral education only I3: Both I1 and I2 training	No training	Community RCT Caregivers	Moderate to severe Alzheimer's or multi-infarct dementia	N= Age: yes Sex: yes % majority race: no Education: yes	None	N=54 Age: yes Sex: yes % majority race: no Education: yes	Duration	NA	NA	6 month post-intervention	Dementia Behavior Disturbance; ADL	Alzheimer's Disease Knowledge Test; Relative Stress Scale
Robinson, 1994 ¹⁴¹ (7993133) US Small sample	I1: Behavior management skill training I2: Social skills training to mobilizing a social network. Both 6 90-min sessions over 2 weeks; individual	No training	Community RCT Caregivers	Dementia unspecified	None	None	N=33 Age: yes Sex: no % majority race: no Education: yes Spouse/partner	None	NA	NA	1 month post-intervention		IC: Montgomery objective and subjective burden Attitude toward adult day care, asking for help

Abbreviations: ADCS=Alzheimer's Disease Cooperative Study; ADKS=Alzheimer's Disease Knowledge Scale; ADL=Activities of Daily Living; ADQ=Approaches to Dementia Questionnaire; BEHAVE-AD=behavioral pathology in Alzheimer's disease rating scale; BACS=Beliefs about Caregiving Scale; CESD=Center for Epidemiologic Studies Depression Scale; CSDD=Cornell scale for depression in dementia; char=characteristics; DBD=dementia behavior disturbance scale; EM=Evidence Map; FC=formal caregiver; FMTCS=Finding Meaning through Caregiving Scale; GDS=Geriatric Depression Scale; GSE=General Self-Efficacy; IC=informal caregiver; ILS=Independent Living Scale; ILS/I=Independent Living Scale Informant version; IRI=Individual Reactivity Index; MMSE=Mini Mental State Examination; N=number; NA=Not Applicable; NPI=Neuropsychiatric Inventory; NR=Not Reported; PAC=Positive aspects of caregiving; PMID=PubMed Identification Number; PLWD=Persons with Dementia; QoL=Quality of Life; RCT=randomized controlled trial; RMBPC=Revised memory and behavior problems checklist; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status; SSCQ=Short Sense of Competence Questionnaire; TACI= Thomas Assessment of Communication Inadequacy; ZBI=Zarit caregiver burden interview

Family Involvement

Table F-46. Risk of bias assessment: family involvement in training

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Funder	Overall Rating
Jablonski, 2005 ¹⁴² (16138529)	9 months	High 39%	Medium	X	X	X	X		High
Maas, 2004 ¹⁴³ 15084992	9 month	High 41% by 5 months	High	X (no blinded assessors)	X	X	X		High

X indicates that domain was not assessed due to high risk of bias.

Abbreviations: PMID=PubMed Identification Number

Evidence Map: Family Involvement

Table F-47. Characteristics of evidence map studies: family involvement in training

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Random-ized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Bramble, 2011 ¹⁴⁴ (21702706) Australia Pilot	FIC: An education programme delivered to staff by the research team.	Not reported	Long term care facility Quasi 58 Caregiver dyads	NA	NA	NA	N=57 Age: Yes Sex: Yes Race: No Education: Yes Relationship: Yes	Employment: Yes Marital status: Yes	N=58 Age: Yes Sex: Yes % majority race: No Education: Yes	Occupation: Yes Employment: Yes	9 months	NA	Staff knowledge; Stress; Staff attitude towards family CGs; Caregiving roles;
Robison, 2007 ¹⁴⁵ (17766671) USA Pilot	The partners in caregiving in a special care environment program: Staff and family received training sessions on communication and conflict-resolution techniques	Not reported	Nursing homes Cluster RCT 20 nursing homes	NR	N=388 Age: Yes Sex: Yes Race: Yes Education: Yes	None	NA	NA	N=384 Age: Yes Sex: Yes Race: Yes Education: Yes	None	2 & 6 months follow up	NA	IC: Frequency of conflicts; Staff empathy; Negative staff behavior; Engagement in NH activities; Burden; Depression FC: Frequency of conflicts; Depression; Job burnout; Job satisfaction

Jablonski, 2005 ¹⁴² (16138529) USA High RoB	Family involvement in care protocol to help family negotiate a partnership with formal staff	Not reported	Nursing home RCT 164 PLWD	Not reported	N=164 Age: Yes Sex: Yes Race: Yes Education: Yes	Marital status: Yes Occupation: Yes	NA	NA	NA	NA	2months; 4 months, 6 months	Functional abilities: FAC	NA
Maas, 2004 ¹⁴³ 15084992 US High ROB	Family involvement in care protocol to help family negotiate a partnership with formal staff	Usual Care	Nursing home Quasi-experimental Paired nursing homes assigned	PLWD in special care units for dementia	N=185 Age: no Sex: no Race: no Education: no	Time in NH	N=185 Age: Yes Sex: Yes % majority race: yes Education: Yes Relationship: Yes	None	NA	NA	9 months	None	Family Perceptions of Caregiving Role Family Perceptions of Care Tool Staff Perceptions of Caregiving Role Caregiver Stress Inventory Attitudes Toward Families Checklist
McCallion, 1999 ¹⁴⁶ US Pilot	Family training in nonverbal communication and structuring family visits	Usual care activities	Nursing home RCT 66 dyads	Moderate dementia with behavioral problems	N=66 Age: Yes Sex: Yes Race: Yes Education: no	None	N=66 Age: Yes Sex: Yes Race: yes Education: Yes Relationship: Yes	None	NA	NA	3 and 6 months	MOSES CSDD CMAI GIPB medication use	IC: DMSS CHS-M Visit satisfaction

Abbreviations: char=characteristics; CHS-M= Caregiver Hassel Scale modified; CG=Caregiver; CHS-M=Caregiving hassles scale; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; DMSS= Dementia Management Strategy Scale; EM=Evidence Map; FAC=Functional Abilities Checklist; FC=formal caregiver; FIC=Family involvement in care; GIPB=geriatric indices of positive behavior; IC=informal caregiver; MOSES=Multidimensional Observation Scale for Elderly Subjects; N=number; NA=Not Applicable; NH=Nursing homes; NR=Not Reported; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RCT=Randomized Controlled Trial; RoB=Risk of Bias; RS=Reporting Status; SES=socioeconomic status

Multitier Training Interventions

Table F-48. Risk of bias assessment: multitier training intervention

Study (PMID)	Outcome Timing	Attrition Bias Attrition %	Selection Bias	Detection Bias	Performance Bias	Reporting Bias	Fidelity Bias	Overall Rating
Kunik, 2017 ¹⁴⁷ (27743840)	3, 6, 12 month	Medium	Low	Low	Medium	Low	Low	Medium
Teri 2005 ¹⁴⁸ (16326662)	2 months 6 months	Low 2 months: 12.6%	Medium	Medium	Medium	Low	Low	Medium

Abbreviations: PMID=PubMed Identification Number

Table F-49. Characteristics of included studies: caregiver focused training intervention

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country RoB* Type	Intervention Focus Theoretical Model Delivery Person Intervention Target Recipient Mode Components Frequency Duration	Target Mode Components Frequency Duration	Setting Design Cluster N Participants Randomized N	Dementia Types Dementia Severity Diagnostic Criteria Age of Diagnosis	PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Teri 2005 ¹⁴⁸ (16326662) US Medium Exploratory	STAR-C: 8 weekly sessions between community consultants and caregivers, followed by four monthly phone calls; to train the caregivers to give consultants the freedom to use clinical judgment and modify the order of session topics in response to a caregiver's needs	Routine medical car (RMC)e	resided in a home setting outside of a nursing home or assisted living facility	Primary physician diagnosis of probable or possible AD with mean MMSE score of 14	N=95 66% Female 80 years Race yes Education yes	None	NA	N=95 70% Female 65 years Race yes Education yes	NA	NA	2 months 6 months	NPI RMBPC QoL-AD;	CES-D HDRS Caregiver Sleep Questionnaire PSS SSCQ
			RCT										
Teri 2005 ¹⁴⁹ (16199404)			95 PLWD and caregivers										
Teri 2012 ¹⁵⁰ (22247431)													

*High risk of bias studies included in evidence map

Abbreviations: AD=Alzheimer's disease; char=characteristics; CES-D=Center for Epidemiological Studies Depression Inventory; FC=formal caregiver; CSC=Cognitive Status Scale; IC=informal caregiver; ICD=International Classification of Diseases; HSC=Hopkins Symptoms Checklist; PBS=Problematic Behavior Scale; PMID=PubMed Identification Number; PLWD=Persons with Dementia; RCT=Randomized Controlled Trial; RoB=Risk of Bias; RS=Reporting Status

Table F-50. PLWD outcomes summary low and medium risk of bias studies: caregiver focused training intervention

Study (PMID) Comparison RoB Category	Outcome Timing	Intervention	Comparator	p-Value
Teri 2005 ¹⁴⁸ (16326662) STAR vs RMC Medium Exploratory	Mean difference (SD) Caregiver depression CES-D 6 months	12.5 (7.7)	15.8 (10.5)	0.046
Teri 2005 ¹⁴⁸ (16326662) STAR vs RMC Medium Exploratory	Mean difference (SD) Caregiver depression HDRS 6 months	6.7 (3.9)	8.5 (5.7)	0.284
Teri 2005 ¹⁴⁸ (16326662) STAR vs RMC Medium Exploratory	Mean difference (SD) Caregiver burden 6 months	21.4 (12.5)	25.8 (13.7)	0.011
Teri 2005 ¹⁴⁸ (16326662) STAR vs RMC Medium Exploratory	Mean difference (SD) Caregiver reaction 6 months	21.9 (15.6)	23.4 (14.5)	0.024
Teri 2005 ¹⁴⁸ (16326662) STAR vs RMC Medium Exploratory	Mean difference (SD) Caregiver sleep questionnaire 6 months	9.1 (4.6)	9.1 (5.2)	0.124
Teri 2005 ¹⁴⁸ (16326662) STAR vs RMC Medium Exploratory	Mean difference (SD) PLWD_QoL 6 months	28.4 (5.4)	28.2 (4.6)	0.049
Teri 2005 ¹⁴⁸ (16326662) STAR vs RMC Medium Exploratory	Mean difference (SD) PLWD RMBPC-memory subscale 6 months	2.8 (0.8)	3.1 (1.0)	0.070

Abbreviations: CES-D=Center for Epidemiological Studies Depression Inventory; CI=Confidence Interval; NR=Not Reported; PMID=PubMed Identification Number; RoB=Risk of Bias

Table F-51. Summary of strength of evidence for PLWD outcomes: caregiver focused training intervention

Comparison Outcome	Timing	# Studies/ Design (n analyzed)	Finding or Summary Statistic	Study Limitations	Consistency	Directness	Precision	Overall Grade/ Conclusion
Consultation vs Control Nursing home admission	12 months	1 RCT (n=84)	Nursing home admissions showed no difference between the comparison groups.	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control Symptom management self-efficacy score	12 months	1 RCT (n=84)	Symptom management self-efficacy scores were not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control Support service self- efficacy	12 months	1 RCT (n=84)	Support service self-efficacy was not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control CES-D score	12 months	1 RCT (n=84)	CES-D scores were not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control Zarit burden score	12 months	1 RCT (n=84)	Zarit burden scores were not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient
Consultation vs Control Hopkins symptom checklist score	12 months	1 RCT (n=84)	Hopkins symptom checklist scores were not different between the comparison groups	Moderate	Unknown	Direct	Imprecise	Insufficient

Abbreviations: CES-D=Center for Epidemiological Studies Depression Inventory; n=Number; RCT=Randomized Controlled Trial

Evidence Map: Multitier Training Interventions

Table 52. Characteristics of evidence map studies: multitier training interventions

Study (PMID)	Intervention	Comparison	Setting and Design	PLWD Dementia Characteristics	PLWD Non-Disease Char	PLWD Non-Disease Char Reporting Status (RS)	Informal Caregiver (IC) Characteristics	Informal Caregiver (IC) Char. RS	Formal Caregiver (FC) Characteristics	Formal Caregiver (FC) Char. RS	Outcome Timing(s)	PLWD Outcomes	Caregiver Outcomes
Country EM Reason			Setting Design Cluster N Participants Randomized N		PLWD N PLWD Age (mean) PLWD Sex (% female) PLWD Race (% majority) PLWD Education (mean years)	PLWD SES PLWD Prior Disability PLWD Household Characteristics PLWD Health Insurance PLWD Detailed Race Information	IC N IC Age (mean) IC Sex (% female) IC Race (% majority) IC Education (mean years) IC Relation to PLWD (% majority)	IC Duration IC Living With PLWD IC Payment IC Health Status IC Dementia Family History IC Employment Status IC Training	FC N FC Age (mean) FC Sex (% female) FC Race (% majority) FC Education (mean years)	FC Health Status FC Training FC Education FC Position FC Length of Service			IC: FC:
Livingston 2019 ⁵¹ (30221615) UK Pilot	DREAM-START Intervention: cognitive-behavioral components, including psychoeducation, light therapy, establishing a new sleep-wake schedule (based on actiwatch data), behavioral activation, relaxation, and coping skills for families	Treatment as usual	Community-dwelling RCT 62 PLWD	Clinical diagnosis of dementia and a SDI item score ≥ 4	N=62 Age: yes Sex: yes Race: yes Education: yes	None	N=62 Age: yes Sex: yes Race: yes Education: yes	None	NA	NA	3 months	referral rates; follow-up rates at three months; all psychotropic medication prescription (to define rescue medication's role); reported side effects: co-morbid physical illnesses and patient falls sleep via actigraphy ESS, NPI, DEMQOL	PSQI SCI HADS ZBI HSQ-12
Kinnunen 2018 ⁵² (30538021) UK Pilot Livingston 2018 ⁵³ (30221615)	DREAMS START: 6 sessions of manual-based intervention for caregiver's understanding sleep and dementia, comprising a cognitive-behavioral component and light therapy	Treatment as usual	Memory service Cluster RCT 62 PLWD	Dementia diagnosis (any type and severity) and sleep disturbances (who scored ≥ 4 on at least one question on SDI) Sleep Disorders Inventory)	N=62 Age: yes Sex: yes Race: yes Education: yes	Annual Income=yes	N=62 Age: yes Sex: yes Race: yes Education: yes	None	NA	NA	3 months	Feasibility Acceptability Referral rate Follow-up rate Use of psychotropic medications Adverse effects and comorbid physical illnesses	Caregiver burden depression

Kunik 2017 ¹⁴⁷ (27743840) US High RoB	PAVeD: 6 to 8 weekly sessions of 45-minute home visits. To give instruction to caregivers on recognizing pain, enhancing communication, and making daily activities pleasant and enjoyable, and at least two elective sessions	EU-PC: 8 weekly 15-minute phone calls to query symptom severity, ascertain needs for immediate psychiatric care, and provide minimal support.	Veterans' Health Administration (VHA) outpatient database for persons with a diagnosis of dementia RCT 213 PLWD	Mild-to-moderate dementia (defined using FAST 2-6 Functional Assessment Staging validated clinician-rated measure of functional decline in dementia)	N=203 Age: yes Sex: yes Race: yes Education: yes	None	N=203 Age: yes Sex: yes Race: yes Education: yes	NA	NA of ≥	NA	3 months 6 months 12 months	CMAI PGPIS GDS PES-AD	Caregiver burden Caregiver-patient relationship Caregiver satisfaction and perceptions of usefulness of PAVeD
Suominen 2015 ¹⁵⁴ (26482691) Finland Small sample	Tailored nutritional guidance on the basis of the food diaries, the results of the weight measurement, the home visits and discussions with the PLWD and caregiver	Got a written guide about nutrition in older adults and all community-provided normal care	Community dwelling RCT 99 PLWD	Diagnosis criteria for a probable AD diagnosis based on the NINCDS-ADRDA Alzheimer's criteria	N=99 Age: yes Sex: yes Race: no Education: no	None	NA	NA	NA	NA	6 months 12 months	weight change; changes in protein and other nutrients intake, QoL and rate of falls.	NA
McCurry 2005 ¹⁵⁵ (15877554) US Small Sample	Written materials describing age- and dementia-related changes in sleep and standard principles of good sleep hygiene. CG also received specific recommendations for sleep hygiene program for the dementia patient. PLWDs to walk daily and increase daytime light exposure with the use of a light box	Control PLWD got general dementia education and caregiver support	Community-dwelling RCT 36 PLWD	PLWDs diagnosed for probable or possible AD with mean MMSE score±SD of 11.8±8.4 and at least one sleep problem on the Neuropsychiatric Inventory Nighttime Behavior scale	N=22 Age: yes Sex: yes Race: yes Education: yes	None	N=22 Age: yes Sex: yes Race: yes Education: yes Relationship: yes	NA	NA	NA	2 months 6 months	total night sleep, percentage of time asleep, number of awakenings, and duration of time awake Light exposure outcomes CESD Daily sleep logs RMBPC	sleep-wake activity PSQI ESS SDQ

McCurry 2003 ¹⁵⁶ (14511168) US Pilot	Written materials describing age- and dementia-related changes in sleep and standard principles of good sleep hygiene. CG also received specific recommendations for sleep hygiene program for the dementia patient.	Control PLWD got general dementia education and caregiver support	Community-dwelling RCT 22 PLWD	Dementia diagnosis according to NINCDS-ADRDA criteria for probable or possible AD confirmed in writing by their primary care physicians. With mean MMSE score±SD of 10.7±7.8 and at least one sleep problem on the Neuropsychiatric Inventory Nighttime Behavior scale	N=22 Age: yes Sex: yes Race: no Education: no	None	N=22 Age: yes Sex: yes Race: no Education: no	NA	NA	NA	2 months	sleep-wake activity CESD Daily sleep logs	sleep-wake activity PSQI ESS
---	--	---	-----------------------------------	--	---	------	---	----	----	----	----------	---	------------------------------------

Abbreviations: BAGS=Behavior Assessment Graphical System; BEHAVE-AD=Behavioral Pathology in Alzheimer's Disease; BI=Barthel Index; CANE=Camberwell Assessment of Need for the Elderly; CAPE-BRS=Clifton Assessment Procedures for the Elderly-Behavior Rating Scale; CBS=Challenging Behavior Scale; CDR=Clinical Dementia Rating; CESD=Center for Epidemiologic Studies Depression Scale; CMAI=Cohen-Mansfield Agitation Inventory; CSDD=Cornell Scale for Depression in Dementia; DEMQOL=Dementia Quality of Life; DSM=Diagnostic and Statistical Manual; EM=Evidence Map; ESS=Epworth Sleepiness Scale; HADS=Hospital Anxiety and Depression Scale; HSQ-12=Health Status Questionnaire-12; MMSE=Mini-mental State Examination; N=number; NA=Not Applicable; NPI=neuropsychiatric inventory; NR=Not Reported; PAS=Psychogeriatric Assessment Scale; PMID=PubMed Identification Number; PSQI=Pittsburgh Sleep Quality Index; QoL-AD=Quality of Life in Alzheimer's Disease; RAID=Rating for Anxiety in Dementia; RMBPC=Revised Memory and Behavior Problem Checklist; SCI=Sleep Condition Indicator; SDQ=Sleep Disorders Questionnaire; ZBI=Zarit Burden Interview

References for Appendix F

1. Fortinsky RH, Kuldorff M, Kleppinger A, et al. Dementia care consultation for family caregivers: collaborative model linking an Alzheimer's association chapter with primary care physicians. *Aging Ment Health*. 2009 Mar;13(2):162-70. doi: <https://dx.doi.org/10.1080/13607860902746160>. PMID: 19347683.
2. Borbasi S, Emmanuel E, Farrelly B, et al. Report of an evaluation of a nurse-led dementia outreach service for people with the behavioural and psychological symptoms of dementia living in residential aged care facilities. *Perspectives in public health*. 2011;131(3):124-30. PMID: CN-00850682.
3. McSweeney K, Jeffreys A, Griffith J, et al. Specialist mental health consultation for depression in Australian aged care residents with dementia: a cluster randomized trial. *Int J Geriatr Psychiatry*. 2012 Nov;27(11):1163-71. doi: <https://dx.doi.org/10.1002/gps.3762>. PMID: 22344753.
4. Opie J, Doyle C, O'Connor DW. Challenging behaviours in nursing home residents with dementia: a randomized controlled trial of multidisciplinary interventions. *Int J Geriatr Psychiatry*. 2002 Jan;17(1):6-13. PMID: 11802224.
5. Orrell M, Hancock G, Hoe J, et al. A cluster randomised controlled trial to reduce the unmet needs of people with dementia living in residential care. *Int J Geriatr Psychiatry*. 2007 Nov;22(11):1127-34. PMID: 17394129.
6. Xiao LD, De Bellis A, Kyriazopoulos H, et al. The Effect of a Personalized Dementia Care Intervention for Caregivers From Australian Minority Groups. *Am J Alzheimers Dis Other Demen*. 2016 Feb;31(1):57-67. doi: <https://dx.doi.org/10.1177/1533317515578256>. PMID: 25805891.
7. MacNeil Vroomen J, Bosmans JE, van de Ven PM, et al. Community-dwelling patients with dementia and their informal caregivers with and without case management: 2-year outcomes of a pragmatic trial. *J Am Med Dir Assoc*. 2015 Sep 01;16(9):800.e1-8. doi: <https://dx.doi.org/10.1016/j.jamda.2015.06.011>. PMID: 26170035.
8. Chien WT, Lee IY. Randomized controlled trial of a dementia care programme for families of home-resided older people with dementia. *J Adv Nurs*. 2011 Apr;67(4):774-87. doi: <https://dx.doi.org/10.1111/j.1365-2648.2010.05537.x>. PMID: 21198803.
9. Lam LC, Lee JS, Chung JC, et al. A randomized controlled trial to examine the effectiveness of case management model for community dwelling older persons with mild dementia in Hong Kong. *Int J Geriatr Psychiatry*. 2010 Apr;25(4):395-402. doi: <https://dx.doi.org/10.1002/gps.2352>. PMID: 19606455.
10. Brodaty H, Draper BM, Millar J, et al. Randomized controlled trial of different models of care for nursing home residents with dementia complicated by depression or psychosis. *J Clin Psychiatry*. 2003 Jan;64(1):63-72. PMID: 12590626.
11. Challis D, von Abendorff R, Brown P, et al. Care management, dementia care and specialist mental health services: an evaluation. *Int J Geriatr Psychiatry*. 2002 Apr;17(4):315-25. PMID: 11994884.
12. Eloniemi-Sulkava U, Notkola IL, Hentinen M, et al. Effects of supporting community-living demented patients and their caregivers: a randomized trial. *J Am Geriatr Soc*. 2001 Oct;49(10):1282-7. PMID: 11890485.
13. Weinberger M, Gold DT, Divine GW, et al. Social service interventions for caregivers of patients with dementia: impact on health care utilization and expenditures. *J Am Geriatr Soc*. 1993 Feb;41(2):153-6. PMID: 8426038.
14. Eloniemi-Sulkava U, Notkola I, Hentinen M, et al. Effects of supporting community-living demented patients and their caregivers: a randomized trial. *J Am Geriatr Soc*. 2001;49(10):1282-7. doi: 10.1046/j.1532-5415.2001.49255.x. PMID: 106906247. Language: English. Entry Date: 20020308. Revision Date: 20171010. Publication Type: Journal Article.
15. Newcomer R, Spitalny M, Fox P, et al. Effects of the Medicare Alzheimer's Disease Demonstration on the use of community-based services. *Health Serv Res*. 1999 Aug;34(3):645-67. PMID: 10445896.
16. Appelhof B, Bakker C, de Vugt ME, et al. Effects of a Multidisciplinary Intervention on the Presence of Neuropsychiatric Symptoms and Psychotropic Drug Use in Nursing Home Residents With Young-Onset Dementia: Behavior and Evolution of Young-Onset Dementia Part 2 (BEYOND-II) Study. *Am J Geriatr Psychiatry*. 2019 Jan 04;04:04. doi: <https://dx.doi.org/10.1016/j.jagp.2018.12.032>. PMID: 30799167.
17. Husebo BS, Ballard C, Aarsland D, et al. The Effect of a Multicomponent Intervention on Quality of Life in Residents of Nursing Homes: A Randomized Controlled Trial (COSMOS). *Journal of the American Medical Directors Association*. 2019. doi: <http://dx.doi.org/10.1016/j.jamda.2018.11.006>. PMID: 2001439020.
18. Livingston G, Barber J, Marston L, et al. Clinical and cost-effectiveness of the Managing Agitation and Raising Quality of Life (MARQUE) intervention for agitation in people with dementia in care homes: a single-blind, cluster-randomised controlled trial. *Lancet Psychiatry*. 2019 04;6(4):293-304. doi: [https://dx.doi.org/10.1016/S2215-0366\(19\)30045-8](https://dx.doi.org/10.1016/S2215-0366(19)30045-8). PMID: 30872010.
19. Nakanishi M, Endo K, Hirooka K, et al. Psychosocial behaviour management programme for home-dwelling people with dementia: A cluster-randomized controlled trial. *Int J Geriatr Psychiatry*. 2018 March;33(3):495-503. doi: <http://dx.doi.org/10.1002/gps.4784>. PMID: 620640680 Pubmed 28857263.
20. Pieper MJ, Francke AL, van der Steen JT, et al. Effects of a Stepwise Multidisciplinary Intervention for Challenging Behavior in Advanced Dementia: A Cluster Randomized Controlled Trial.[Erratum appears in *J Am Geriatr Soc*. 2016 Jun;64(6):1383; PMID: 27321637]. *J Am Geriatr Soc*. 2016 Feb;64(2):261-9. doi: <https://dx.doi.org/10.1111/jgs.13868>. PMID: 26804064.
21. McCabe MP, Bird M, Davison TE, et al. An RCT to evaluate the utility of a clinical protocol for staff in the management of behavioral and psychological symptoms of dementia in residential aged-care settings. *Aging Ment Health*. 2015;19(9):799-807. doi: <https://dx.doi.org/10.1080/13607863.2014.967659>. PMID: 25319535.
22. Zwijsen SA, Smalbrugge M, Eefsting JA, et al. Coming to grips with challenging behavior: a cluster randomized controlled trial on the effects of a multidisciplinary care program for challenging behavior in dementia. *J Am Med Dir Assoc*. 2014 Jul;15(7):531.e1-e10. doi: <https://dx.doi.org/10.1016/j.jamda.2014.04.007>. PMID: 24878214.
23. Rapp MA, Mell T, Majic T, et al. Agitation in nursing home residents with dementia (VIDEANT trial): effects of a cluster-randomized, controlled, guideline implementation trial. *J Am Med Dir Assoc*. 2013 Sep;14(9):690-5. doi: <https://dx.doi.org/10.1016/j.jamda.2013.05.017>. PMID: 23827658.
24. Cohen-Mansfield J, Thein K, Marx MS, et al. Efficacy of nonpharmacologic interventions for agitation in advanced dementia: a randomized, placebo-controlled trial. *J Clin Psychiatry*. 2012 Sep;73(9):1255-61. doi: <https://dx.doi.org/10.4088/JCP.12m07918>. PMID: 23059151.
25. Kovach CR, Simpson MR, Joosse L, et al. Comparison of the effectiveness of two protocols for treating nursing home residents with advanced dementia. *Res*. 2012 Oct;5(4):251-63. doi: <https://dx.doi.org/10.3928/19404921-20120906-01>. PMID: 22998656.
26. Salva A, Andrieu S, Fernandez E, et al. Health and nutrition promotion program for patients with dementia (NutriAlz): Cluster randomized trial. *J Nutr Health Aging*. 2011;15(10):822-30. doi: 10.1007/s12603-011-0363-3. PMID: 104625164. Language: English. Entry Date: 20120530. Revision Date: 20150818. Publication Type: Journal Article.
27. Kovach CR, Logan BR, Noonan PE, et al. Effects of the Serial Trial Intervention on discomfort and behavior of nursing home residents with dementia. *Am J Alzheimers Dis Other Demen*. 2006 Jun-Jul;21(3):147-55. PMID: 16869334.
28. Chen YH, Lin LC. Ability of the Pain Recognition and Treatment (PRT) Protocol to Reduce Expressions of Pain among Institutionalized Residents with Dementia: A Cluster Randomized Controlled Trial. *Pain Manag Nurs*. 2016 Feb;17(1):14-24. doi: <https://dx.doi.org/10.1016/j.pmn.2015.08.003>. PMID: 26584896.
29. Pieper M, Francke A, Steen J, et al. Effects of a stepwise approach to behavioural problems in dementia: a cluster randomised controlled trial. In *Nederlands tijdschrift voor geneeskunde*
30. Kovach CR, Hekel B, Rababa M. Feasibility Testing of a Protocol to Stop Ineffective Drug and Nondrug Treatments. *West J Nurs Res*. 2015 Nov;37(11):1404-22. doi: <https://dx.doi.org/10.1177/0193945915599070>. PMID: 26250849.
31. Zwijsen SA, Gerritsen DL, Eefsting JA, et al. Coming to grips with challenging behaviour: a cluster randomised controlled trial on the effects of a new care programme for challenging behaviour on burnout, job satisfaction and job demands of care staff on dementia special care units. *Int J Nurs Stud*. 2015 Jan;52(1):68-74. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2014.10.003>. PMID: 25458804.
32. Cohen-Mansfield J, Thein K, Marx MS, et al. What are the barriers to performing nonpharmacological interventions for behavioral symptoms in the nursing home? *J Am Med Dir Assoc*. 2012 May;13(4):400-5. doi: <https://dx.doi.org/10.1016/j.jamda.2011.07.006>. PMID: 21872537.
33. Kovach CR, Simpson MR, Joosse L, et al. Comparison of the Effectiveness of Two Protocols for Treating Nursing Home Residents with Advanced Dementia. *Res*. 2012;5(4):251-63. doi: 10.3928/19404921-20120906-01. PMID: 104426188. Language: English. Entry Date: 20121025. Revision Date: 20150711. Publication Type: Journal Article.
34. Montgomery RJ, Kwak J, Kosloski K, et al. Effects of the TCARE intervention on caregiver burden and depressive symptoms: preliminary findings from a randomized controlled study. *J Gerontol B Psychol Sci Soc Sci*. 2011 Sep;66(5):640-7. doi: <https://dx.doi.org/10.1093/geronb/gbr088>. PMID: 21840840.
35. Verkaik R, Francke AL, van Meijel B, et al. The effects of a nursing guideline on depression in psychogeriatric nursing home residents with dementia. *Int J Geriatr Psychiatry*. 2011 Jul;26(7):723-32. doi: <https://dx.doi.org/10.1002/gps.2586>. PMID: 21495077.
36. Cohen-Mansfield J, Libin A, Marx MS. Nonpharmacological treatment of agitation: a controlled trial of systematic individualized intervention. *J Gerontol A Biol Sci Med Sci*. 2007 Aug;62(8):908-16. PMID: 17702884.
37. Kovach CR, Cashin JR, Sauer L. Deconstruction of a complex tailored intervention to assess and treat discomfort of people with advanced dementia. *J Adv Nurs*. 2006 Sep;55(6):678-88. PMID: 16925616.
38. Brazil K, Carter G, Cardwell C, et al. Effectiveness of advance care planning with family carers in dementia nursing homes: A paired cluster randomized controlled trial. *Palliat Med*. 2018 Mar;32(3):603-12. doi: <https://dx.doi.org/10.1177/0269216317722413>. PMID: 28786323.
39. Mitchell SL, Shaffer ML, Cohen S, et al. An Advance Care Planning Video Decision Support Tool for Nursing Home Residents With Advanced Dementia: A Cluster Randomized Clinical Trial. *JAMA Intern Med*. 2018 Jul 01;178(7):961-9. doi: <https://dx.doi.org/10.1001/jamainternmed.2018.1506>. PMID: 29868778.
40. Cohen SM, Volandes AE, Shaffer ML, et al. Concordance Between Proxy Level of Care Preference and Advance Directives Among Nursing Home Residents With Advanced Dementia: A Cluster Randomized Clinical Trial. *J Pain Symptom Manage*. 2019 Jan;57(1):37-46.e1. doi: <https://dx.doi.org/10.1016/j.jpainsymman.2018.09.018>. PMID: 30273717.
41. Hanson LC, Zimmerman S, Song MK, et al. Effect of the Goals of Care Intervention for Advanced Dementia: A Randomized Clinical Trial. *JAMA Intern Med*. 2017 Jan 01;177(1):24-31. doi: <https://dx.doi.org/10.1001/jamainternmed.2016.7031>. PMID: 27893884.

42. Hanson LC, Song MK, Zimmerman S, et al. Fidelity to a behavioral intervention to improve goals of care decisions for nursing home residents with advanced dementia. *Clin. 2016 Dec*;13(6):599-604. PMID: 27271683.
43. Reinhardt JP, Chichin E, Posner L, et al. Vital conversations with family in the nursing home: preparation for end-stage dementia care. *J Soc Work End Life Palliat Care. 2014*;10(2):112-26. doi: <https://dx.doi.org/10.1080/15524256.2014.906371>. PMID: 24835382.
44. Hanson LC, Carey TS, Caprio AJ, et al. Improving decision-making for feeding options in advanced dementia: a randomized, controlled trial. *J Am Geriatr Soc. 2011 Nov*;59(11):2009-16. doi: <https://dx.doi.org/10.1111/j.1532-5415.2011.03629.x>. PMID: 22091750.
45. Goossens B, Sevenants A, Declercq A, et al. Improving shared decision-making in advance care planning: Implementation of a cluster randomized staff intervention in dementia care. *Patient Educ Couns. 2019 Nov* 26;26:26. doi: <https://dx.doi.org/10.1016/j.pec.2019.11.024>. PMID: 31818522.
46. Goossens B, Sevenants A, Declercq A, et al. 'We DECide optimized' - training nursing home staff in shared decision-making skills for advance care planning conversations in dementia care: protocol of a pretest-posttest cluster randomized trial. *BMC geriatr. 2019 02* 04;19(1):33. doi: <https://dx.doi.org/10.1186/s12877-019-1044-z>. PMID: 30717700.
47. Song MK, Ward SE, Hepburn K, et al. Can Persons with Dementia Meaningfully Participate in Advance Care Planning Discussions? A Mixed-Methods Study of SPIRIT. *J Palliat Med. 2019 Nov*;22(11):1410-6. doi: <https://dx.doi.org/10.1089/jpm.2019.0088>. PMID: 31373868.
48. Bonner GJ, Wang E, Wilkie DJ, et al. Advance care treatment plan (ACT-Plan) for African American family caregivers: a pilot study. *Dementia. 2014 Jan*;13(1):79-95. doi: <https://dx.doi.org/10.1177/1471301212449408>. PMID: 24381040.
49. Einterz SF, Gilliam R, Lin FC, et al. Development and testing of a decision aid on goals of care for advanced dementia. *J Am Med Dir Assoc. 2014 Apr*;15(4):251-5. doi: <https://dx.doi.org/10.1016/j.jamda.2013.11.020>. PMID: 24508326.
50. Sampson EL, Jones L, Thune-Boyle IC, et al. Palliative assessment and advance care planning in severe dementia: an exploratory randomized controlled trial of a complex intervention. *Palliat Med. 2011 Apr*;25(3):197-209. doi: <https://dx.doi.org/10.1177/0269216310391691>. PMID: 21228087.
51. Verreault R, Arcand M, Misson L, et al. Quasi-experimental evaluation of a multifaceted intervention to improve quality of end-of-life care and quality of dying for patients with advanced dementia in long-term care institutions. *Palliat Med. 2018 03*;32(3):613-21. doi: <https://dx.doi.org/10.1177/0269216317719588>. PMID: 28731379.
52. Boogaard JA, de Vet HC, van Soest-Poortvliet MC, et al. Effects of two feedback interventions on end-of-life outcomes in nursing home residents with dementia: A cluster-randomized controlled three-armed trial. *Palliat Med. 2018 Mar*;32(3):693-702. doi: <https://dx.doi.org/10.1177/0269216317750071>. PMID: 29343173.
53. Agar M, Luckett T, Luscombe G, et al. Effects of facilitated family case conferencing for advanced dementia: A cluster randomised clinical trial. *PLoS ONE. 2017*;12(8):e0181020. doi: <https://dx.doi.org/10.1371/journal.pone.0181020>. PMID: 28786995.
54. Ahronheim JC, Morrison RS, Morris J, et al. Palliative care in advanced dementia: a randomized controlled trial and descriptive analysis. *J Palliat Med. 2000*;3(3):265-73. PMID: 15859668.
55. Hanson LC, Kistler CE, Lavin K, et al. Triggered Palliative Care for Late-Stage Dementia: A Pilot Randomized Trial. *J Pain Symptom Manage. 2019 Jan*;57(1):10-9. doi: <https://dx.doi.org/10.1016/j.jpainsymman.2018.10.494>. PMID: 30342242.
56. Horvath KJ, Trudeau SA, Rudolph JL, et al. Clinical trial of a home safety toolkit for Alzheimer's disease. *Int J Alzheimer Dis. 2013*;2013:913606. doi: <https://dx.doi.org/10.1155/2013/913606>. PMID: 24195007.
57. Hanson LC, Gilliam R, Lee TJ. Successful clinical trial research in nursing homes: the Improving Decision-Making Study. *Clin. 2010 Dec*;7(6):735-43. doi: <https://dx.doi.org/10.1177/1740774510380241>. PMID: 20729251.
58. Snyder EA, Caprio AJ, Wessell K, et al. Impact of a decision aid on surrogate decision-makers' perceptions of feeding options for patients with dementia. *J Am Med Dir Assoc. 2013 Feb*;14(2):114-8. doi: <https://dx.doi.org/10.1016/j.jamda.2012.10.011>. PMID: 23273855.
59. Laver K, Liu E, Clemson L, et al. Does Telehealth Delivery of a Dyadic Dementia Care Program Provide a Noninferior Alternative to Face-To-Face Delivery of the Same Program? A Randomized, Controlled Trial. *Am J Geriatr Psychiatry. 2020 Mar* 02;02:02. doi: <https://dx.doi.org/10.1016/j.jagp.2020.02.009>. PMID: 32234275.
60. Vila-Miravent J, Fernández E, Villar F, et al. Involving institutionalized people with dementia in their care-planning meetings: Impact on their quality of life measured by a proxy method: Innovative Practice. *Dementia (14713012). 2019*;18(5):1936-41. doi: 10.1177/1471301217695381. PMID: 137592911. Language: English. Entry Date: 20190724. Revision Date: 20190724. Publication Type: Article.
61. Williams KN, Perkhounkova Y, Shaw CA, et al. Supporting Family Caregivers With Technology for Dementia Home Care: A Randomized Controlled Trial. *Innov Aging. 2019 Jul*;3(3):igz037. doi: <https://dx.doi.org/10.1093/geroni/igz037>. PMID: 31660443.
62. Halek M, Reuther S, Muller-Widmer R, et al. Dealing with the behaviour of residents with dementia that challenges: A stepped-wedge cluster randomized trial of two types of dementia-specific case conferences in nursing homes (FallDem). *Int J Nurs Stud. 2020 Apr*;104:103435. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2019.103435>. PMID: 32062052.
63. Possin KL, Merrilees JJ, Dulaney S, et al. Effect of Collaborative Dementia Care via Telephone and Internet on Quality of Life, Caregiver Well-being, and Health Care Use: The Care Ecosystem Randomized Clinical Trial. *JAMA Intern Med. 2019 Sep* 30;30:30. doi: <https://dx.doi.org/10.1001/jamainternmed.2019.4101>. PMID: 31566651.
64. Laporte Uribe F, Grasse J, Grill S, et al. Regional dementia care networks in Germany: changes in caregiver burden at one-year follow-up and associated factors. *Int Psychogeriatr. 2017 06*;29(6):991-1004. doi: <https://dx.doi.org/10.1017/S1041610217000126>. PMID: 28249632.
65. Thyrian JR, Hertel J, Wucherer D, et al. Effectiveness and Safety of Dementia Care Management in Primary Care: A Randomized Clinical Trial. *JAMA Psychiatry. 2017 Oct* 01;74(10):996-1004. doi: <https://dx.doi.org/10.1001/jamapsychiatry.2017.2124>. PMID: 28746708.
66. Chodosh J, Colaiaco BA, Connor KI, et al. Dementia Care Management in an Underserved Community: The Comparative Effectiveness of Two Different Approaches. *J Aging Health. 2015 Aug*;27(5):864-93. doi: <https://dx.doi.org/10.1177/0898264315569454>. PMID: 25656074.
67. Leontjevas R, Teerenstra S, Smalbrugge M, et al. More insight into the concept of apathy: a multidisciplinary depression management program has different effects on depressive symptoms and apathy in nursing homes. *Int Psychogeriatr. 2013*;25(12):1941-52. doi: 10.1017/S1041610213001440. PMID: 104115576. Language: English. Entry Date: 20140808. Revision Date: 20180803. Publication Type: journal article.
68. Bass D, Judge K, Snow A, et al. Caregiver outcomes of partners in dementia care: effect of a care coordination program for veterans with dementia and their family members and friends. *J Am Geriatr Soc. 2013*;61(8):1377-86. doi: 10.1111/jgs.12362. PMID: CN-00875084.
69. Chapman DG, Toseland RW. Effectiveness of advanced illness care teams for nursing home residents with dementia. *Soc Work. 2007 Oct*;52(4):321-9. PMID: 18232242.
70. Callahan CM, Boustani MA, Unverzagt FW, et al. Effectiveness of collaborative care for older adults with Alzheimer disease in primary care: a randomized controlled trial. *Jama. 2006 May* 10;295(18):2148-57. PMID: 16684985.
71. Vickrey BG, Mittman BS, Connor KI, et al. The effect of a disease management intervention on quality and outcomes of dementia care: a randomized, controlled trial.[Summary for patients in *Ann Intern Med. 2006 Nov* 21;145(10):I31; PMID: 17116913]. *Ann Intern Med. 2006 Nov* 21;145(10):713-26. PMID: 17116916.
72. Surr CA, Griffiths AW, Kelley R, et al. The Implementation of Dementia Care Mapping in a Randomized Controlled Trial in Long-Term Care: Results of a Process Evaluation. *Am J Alzheimers Dis Other Dement. 2019 Sep*;34(6):390-8. doi: <https://dx.doi.org/10.1177/1533317519845725>. PMID: 31056923.
73. Ballard C, Corbett A, Orrell M, et al. Impact of person-centred care training and person-centred activities on quality of life, agitation, and antipsychotic use in people with dementia living in nursing homes: A cluster-randomised controlled trial. *PLoS Med. 2018 02*;15(2):e1002500. doi: <https://dx.doi.org/10.1371/journal.pmed.1002500>. PMID: 29408901.
74. Lichtwarck B, Selbaek G, Kirkevold O, et al. Targeted Interdisciplinary Model for Evaluation and Treatment of Neuropsychiatric Symptoms: A Cluster Randomized Controlled Trial. *Am J Geriatr Psychiatry. 2018 January*;26(1):25-38. doi: <http://dx.doi.org/10.1016/j.jagp.2017.05.015>. PMID: 617024611.
75. Dichter M, Quasdorf T, Schwab C, et al. Dementia care mapping: effects on residents' quality of life and challenging behavior in German nursing homes. A quasi-experimental trial. *Int Psychogeriatr. 2015*;27(11):1875-92. doi: 10.1017/S1041610215000927. PMID: CN-01215190.
76. Chenoweth L, Forbes I, Fleming R, et al. PerCEN: a cluster randomized controlled trial of person-centered residential care and environment for people with dementia. *Int Psychogeriatr. 2014 Jul*;26(7):1147-60. doi: <https://dx.doi.org/10.1017/S1041610214000398>. PMID: 24666667.
77. van de Ven G, Draskovic I, Adang EM, et al. Effects of dementia-care mapping on residents and staff of care homes: a pragmatic cluster-randomised controlled trial. *PLoS ONE. 2013*;8(7):e67325. doi: <https://dx.doi.org/10.1371/journal.pone.0067325>. PMID: 23844003.
78. Rokstad AM, Rosvik J, Kirkevold O, et al. The effect of person-centred dementia care to prevent agitation and other neuropsychiatric symptoms and enhance quality of life in nursing home patients: a 10-month randomized controlled trial. *Dement Geriatr Cogn Disord. 2013*;36(5-6):340-53. doi: <https://dx.doi.org/10.1159/000354366>. PMID: 24022375.
79. Chenoweth L, King MT, Jeon YH, et al. Caring for Aged Dementia Care Resident Study (CADRES) of person-centred care, dementia-care mapping, and usual care in dementia: a cluster-randomised trial.[Erratum appears in *Lancet Neurol. 2009 May*;8(5):419]. *Lancet neurol. 2009 Apr*;8(4):317-25. doi: [https://dx.doi.org/10.1016/S1474-4422\(09\)70045-6](https://dx.doi.org/10.1016/S1474-4422(09)70045-6). PMID: 19282246.
80. Eloniemi-Sulkava U, Saarenheimo M, Laakkonen ML, et al. Family care as collaboration: effectiveness of a multicomponent support program for elderly couples with dementia. Randomized controlled intervention study. *J Am Geriatr Soc. 2009 Dec*;57(12):2200-8. doi: <https://dx.doi.org/10.1111/j.1532-5415.2009.02564.x>. PMID: 20121986.
81. Fossey J, Ballard C, Juszczak E, et al. Effect of enhanced psychosocial care on antipsychotic use in nursing home residents with severe dementia: cluster randomised trial. *BMJ (Clinical research ed). 2006 1 Apr*;332(7544):756-61. PMID: 43509632.
82. Rovner BW, Steele CD, Shmueli Y, et al. A randomized trial of dementia care in nursing homes. *J Am Geriatr Soc. 1996 Jan*;44(1):7-13. PMID: 8537594.
83. Eichler T, Thyrian JR, Dreier A, et al. Dementia care management: going new ways in ambulant dementia care within a GP-based randomized controlled intervention trial. *Int Psychogeriatr. 2014 Feb*;26(2):247-56. doi: <https://dx.doi.org/10.1017/S1041610213001786>. PMID: 24152974.
84. Eichler T, Thyrian JR, Fredrich D, et al. The benefits of implementing a computerized intervention-management-system (IMS) on delivering integrated dementia care in the primary care setting. *Int Psychogeriatr. 2014 Aug*;26(8):1377-85. doi: <https://dx.doi.org/10.1017/S1041610214000830>. PMID: 24811145.
85. Fis T, Thyrian JR, Wucherer D, et al. Medication management for people with dementia in primary care: description of implementation in the DelpHi study. *BMC geriatr. 2013 Nov* 13;13:121. doi: <https://dx.doi.org/10.1186/1471-2318-13-121>. PMID: 24225205.
86. Thyrian JR, Winter P, Eichler T, et al. Relatives' burden of caring for people screened positive for dementia in primary care : Results of the DelpHi study. *Z Gerontol Geriatr. 2017 Jan*;50(1):4-13. doi: <https://dx.doi.org/10.1007/s00391-016-1119-9>. PMID: 27534949.
87. Thyrian JR, Fis T, Dreier A, et al. Life- and person-centred help in Mecklenburg-Western Pomerania, Germany (DelpHi): study protocol for a randomised controlled trial. *Trials. 2012 May* 10;13:56. doi: <https://dx.doi.org/10.1186/1745-6215-13-56>. PMID: 22575023.

88. Zwingmann I, Hoffmann W, Michalowsky B, et al. Supporting family dementia caregivers: testing the efficacy of dementia care management on multifaceted caregivers' burden. *Aging Ment Health*. 2017 Nov 20;1-8. doi: <https://dx.doi.org/10.1080/13607863.2017.1399341>. PMID: 29156941.
89. Michalowsky B, Xie F, Eichler T, et al. Cost-effectiveness of a collaborative dementia care management-Results of a cluster-randomized controlled trial. *Alzheimer's Dement*. 2019 Oct;15(10):1296-308. doi: <https://dx.doi.org/10.1016/j.jalz.2019.05.008>. PMID: 31409541.
90. Morgan RO, Bass DM, Judge KS, et al. A break-even analysis for dementia care collaboration: Partners in Dementia Care. *JGIM: Journal of General Internal Medicine*. 2015;30(6):804-9. doi: 10.1007/s11606-015-3205-x. PMID: 109738468. Language: English. Entry Date: 20150923. Revision Date: 20160604. Publication Type: journal article.
91. Duru OK, Ettner SL, Vassar SD, et al. Cost evaluation of a coordinated care management intervention for dementia. *Am J Manag Care*. 2009 Aug;15(8):521-8. PMID: 19670955.
92. Chodosh J, Pearson ML, Connor KI, et al. A dementia care management intervention: which components improve quality? *Am J Manag Care*. 2012 Feb;18(2):85-94. PMID: 22435836.
93. Brown AF, Vassar SD, Connor KI, et al. Collaborative care management reduces disparities in dementia care quality for caregivers with less education. *J Am Geriatr Soc*. 2013 Feb;61(2):243-51. doi: <https://dx.doi.org/10.1111/jgs.12079>. PMID: 23320655.
94. Lichtwarck B, Selbaek G, Kirkevold O, et al. TIME - Targeted interdisciplinary model for evaluation and treatment of neuropsychiatric symptoms: protocol for an effectiveness-implementation cluster randomized hybrid trial. *BMC Psychiatry*. 2016 Jul 12;16:233. doi: <https://dx.doi.org/10.1186/s12888-016-0944-0>. PMID: 27406242.
95. Lichtwarck B, Myhre J, Selbaek G, et al. TIME to reduce agitation in persons with dementia in nursing homes. A process evaluation of a complex intervention. *BMC Health Serv Res*. 2019 May 31;19(1):349. doi: <https://dx.doi.org/10.1186/s12913-019-4168-0>. PMID: 31151437.
96. Fossey J, Garrod L, Tolbol Froiland C, et al. What influences the sustainability of an effective psychosocial intervention for people with dementia living in care homes? A 9 to 12-month follow-up of the perceptions of staff in care homes involved in the WHELD randomised controlled trial. *Int J Geriatr Psychiatry*. 2019 May;34(5):674-82. doi: <http://dx.doi.org/10.1002/gps.5066>.
97. Romeo R, Zala D, Knapp M, et al. Improving the quality of life of care home residents with dementia: Cost-effectiveness of an optimized intervention for residents with clinically significant agitation in dementia. *Alzheimer's Dement*. 2019 Feb;15(2):282-91. doi: <https://dx.doi.org/10.1016/j.jalz.2018.08.010>. PMID: 30470592.
98. Stein-Parbury J, Chenoweth L, Jeon YH, et al. Implementing Person-Centered Care in Residential Dementia Care. *Clin Gerontol*. 2012;35(5):404-24. doi: 10.1080/07317115.2012.702654. PMID: 104505176. Language: English. Entry Date: 20120917. Revision Date: 20150820. Publication Type: Journal Article.
99. Jeon YH, Luscombe G, Chenoweth L, et al. Staff outcomes from the caring for aged dementia care resident study (CADRES): a cluster randomised trial. *Int J Nurs Stud*. 2012 May;49(5):508-18. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2011.10.020>. PMID: 22078076.
100. Froggatt K, Best A, Bunn F, et al. A group intervention to improve quality of life for people with advanced dementia living in care homes: The namaste feasibility cluster RCT. *Health Technology Assessment*. 2020 January;24(6):vii-139.
101. Chen C, Huang Y, Liu C, et al. Effects of an Interdisciplinary Care Team on the Management of Alzheimer's Disease in China. *J Gerontol Nurs*. 2019 01 May;45(5):39-45. doi: <http://dx.doi.org/10.3928/00989134-20190318-01>. PMID: 627579646.
102. Saxena N, George PP, Teo KW, et al. Evaluation of an integrated primary care-led dementia shared care program in Singapore: An effectiveness and cost-effectiveness study. *Geriatr Gerontol Int*. 2018 Mar;18(3):479-86. doi: <https://dx.doi.org/10.1111/ggi.13196>. PMID: 29193721.
103. Mavandadi S, Wright EM, Graydon MM, et al. A randomized pilot trial of a telephone-based collaborative care management program for caregivers of individuals with dementia. *Psychol Serv*. 2017 Feb;14(1):102-11. doi: <https://dx.doi.org/10.1037/ser0000118>. PMID: 28134558.
104. Samus QM, Johnston D, Black BS, et al. A multidimensional home-based care coordination intervention for elders with memory disorders: the maximizing independence at home (MIND) pilot randomized trial. *Am J Geriatr Psychiatry*. 2014 Apr;22(4):398-414. doi: <https://dx.doi.org/10.1016/j.jagp.2013.12.175>. PMID: 24502822.
105. Leontjevas R, Gerritsen DL, Smalbrugge M, et al. A structural multidisciplinary approach to depression management in nursing-home residents: a multicentre, stepped-wedge cluster-randomised trial. *Lancet*. 2013 Jun 29;381(9885):2255-64. doi: [https://dx.doi.org/10.1016/S0140-6736\(13\)60590-5](https://dx.doi.org/10.1016/S0140-6736(13)60590-5). PMID: 23643110.
106. Bass DM, Clark PA, Looman WJ, et al. The Cleveland Alzheimer's managed care demonstration: outcomes after 12 months of implementation. *Gerontologist*. 2003 Feb;43(1):73-85. PMID: 12604748.
107. Clark PA, Bass DM, Looman WJ, et al. Outcomes for patients with dementia from the Cleveland Alzheimer's Managed Care Demonstration. *Aging Ment Health*. 2004 Jan;8(1):40-51. PMID: 14690867.
108. Li J, Grandner M, Chang Y, et al. Person-centered dementia care and sleep in assisted living residents with dementia: a pilot study. *Behav Sleep Med*. 2017(2):97-113. doi: 10.1080/15402002.2015.1104686. PMID: CN-01380345.
109. Ballard C, Orrell M, YongZhong S, et al. Impact of Antipsychotic Review and Nonpharmacological Intervention on Antipsychotic Use, Neuropsychiatric Symptoms, and Mortality in People With Dementia Living in Nursing Homes: A Factorial Cluster-Randomized Controlled Trial by the Well-Being and Health for People With Dementia (WHELD) Program. *Am J Psychiatry*. 2016 Mar 01;173(3):252-62. doi: <https://dx.doi.org/10.1176/appi.ajp.2015.15010130>. PMID: 26585409.
110. Moyle W, Venturato L, Cooke M, et al. Evaluating the capabilities model of dementia care: A non-randomized controlled trial exploring resident quality of life and care staff attitudes and experiences. *Int Psychogeriatr*. 2016 01 Jul;28(7):1091-100. doi: <http://dx.doi.org/10.1017/S1041610216000296>. PMID: 608901615.
111. Gozalo P, Prakash S, Qato DM, et al. Effect of the bathing without a battle training intervention on bathing-associated physical and verbal outcomes in nursing home residents with dementia: a randomized crossover diffusion study. *J Am Geriatr Soc*. 2014 May;62(5):797-804. doi: <https://dx.doi.org/10.1111/jgs.12777>. PMID: 24697702.
112. Schindel ML, Gillies L, Coker E, et al. An Education Intervention to Enhance Staff Self-Efficacy to Provide Dementia Care in an Acute Care Hospital in Canada: a Nonrandomized Controlled Study. In *American journal of alzheimer's disease and other dementias*
113. Galik E, Resnick B, Hammersla M, et al. Optimizing function and physical activity among nursing home residents with dementia: testing the impact of function-focused care. *Gerontologist*. 2014 Dec;54(6):930-43. doi: <https://dx.doi.org/10.1093/geront/gnt108>. PMID: 24092822.
114. Williams KN, Perkhounkova Y, Herman R, et al. A Communication Intervention to Reduce Resistiveness in Dementia Care: A Cluster Randomized Controlled Trial. *Gerontologist*. 2017 Aug 01;57(4):707-18. doi: <https://dx.doi.org/10.1093/geront/gnw047>. PMID: 27048705.
115. Conway ER, Chenery HJ. Evaluating the MESSAGE Communication Strategies in Dementia training for use with community-based aged care staff working with people with dementia: a controlled pretest-post-test study. *J Clin Nurs*. 2016 Apr;25(7-8):1145-55. doi: <https://dx.doi.org/10.1111/jocn.13134>. PMID: 26821868.
116. Hattink B, Meiland F, van der Roest H, et al. Web-Based STAR E-Learning Course Increases Empathy and Understanding in Dementia Caregivers: Results from a Randomized Controlled Trial in the Netherlands and the United Kingdom. *J Med Internet Res*. 2015 Oct 30;17(10):e241. doi: <https://dx.doi.org/10.2196/jmir.4025>. PMID: 26519106.
117. O'Shea E, Devane D, Cooney A, et al. The impact of reminiscence on the quality of life of residents with dementia in long-stay care. *Int J Geriatr Psychiatry*. 2014 Oct;29(10):1062-70. doi: <https://dx.doi.org/10.1002/gps.4099>. PMID: 24633858.
118. Clare L, Whitaker R, Woods RT, et al. AwareCare: a pilot randomized controlled trial of an awareness-based staff training intervention to improve quality of life for residents with severe dementia in long-term care settings. *Int Psychogeriatr*. 2013 Jan;25(1):128-39. doi: <https://dx.doi.org/10.1017/S1041610212001226>. PMID: 22840185.
119. Ghandehari OO, Hadjistavropoulos T, Williams J, et al. A controlled investigation of continuing pain education for long-term care staff. *Pain Research & Management*. 2013;18(1):11-8. PMID: 107988179. Language: English. Entry Date: 20130426. Revision Date: 20161117. Publication Type: journal article.
120. Wenborn J, Challis D, Head J, et al. Providing activity for people with dementia in care homes: a cluster randomised controlled trial. *Int J Geriatr Psychiatry*. 2013 Dec;28(12):1296-304. doi: <https://dx.doi.org/10.1002/gps.3960>. PMID: 23637069.
121. Leone E, Deudon A, Piano J, et al. Are Dementia Patient's Engagement Using Tailored Stimuli the Same? The Apathy Dilemma in Nursing Home Residents. *Curr Gerontol Geriatr Res*. 2012;2012:942640. doi: <https://dx.doi.org/10.1155/2012/942640>. PMID: 22956946.
122. Zimmerman S, Mitchell CM, Reed D, et al. Outcomes of a dementia care training program for staff in nursing homes and residential care/assisted living settings. *Alzheimer's Care Today*. 2010;11(2):83-99. PMID: 105194596. Language: English. Entry Date: 20100625. Revision Date: 20150711. Publication Type: Journal Article.
123. Deudon A, Maubourguet N, Gervais X, et al. Non-pharmacological management of behavioural symptoms in nursing homes. *Int J Geriatr Psychiatry*. 2009 Dec;24(12):1386-95. doi: <https://dx.doi.org/10.1002/gps.2275>. PMID: 19370714.
124. Kuske B, Luck T, Hanns S, et al. Training in dementia care: a cluster-randomized controlled trial of a training program for nursing home staff in Germany. *Int Psychogeriatr*. 2009 Apr;21(2):295-308. doi: <https://dx.doi.org/10.1017/S1041610208008387>. PMID: 19193252.
125. Chang CC, Lin LC. Effects of a feeding skills training programme on nursing assistants and dementia patients. *J Clin Nurs*. 2005 Nov;14(10):1185-92. PMID: 16238764.
126. Sloane PD, Hoeffler B, Mitchell CM, et al. Effect of person-centered showering and the towel bath on bathing-associated aggression, agitation, and discomfort in nursing home residents with dementia: a randomized, controlled trial. *J Am Geriatr Soc*. 2004 Nov;52(11):1795-804. PMID: 15507054.
127. Magai C, Cohen CI, Gomberg D. Impact of training dementia caregivers in sensitivity to nonverbal emotion signals. *Int Psychogeriatr*. 2002 Mar;14(1):25-38. PMID: 12094904.
128. Bourgeois MS, Dijkstra K, Burgio L, et al. Memory aids as an augmentative and alternative communication strategy for nursing home residents with dementia. *AAC: Augmentative and Alternative Communication*. 2001 Sep;17(3):196-210. doi: <http://dx.doi.org/10.1080/714043383>. PMID: 2001-11280-006.
129. Wells DL, Dawson P, Sidani S, et al. Effects of an abilities-focused program of morning care on residents who have dementia and on caregivers. *J Am Geriatr Soc*. 2000 Apr;48(4):442-9. PMID: 10798473.
130. McCallion P, Toseland RW, Lacey D, et al. Educating nursing assistants to communicate more effectively with nursing home residents with dementia. *Gerontologist*. 1999 Oct;39(5):546-58. PMID: 10568079.
131. Proctor R, Burns A, Powell HS, et al. Behavioural management in nursing and residential homes: a randomised controlled trial. *Lancet*. 1999 Jul 03;354(9172):26-9. PMID: 10406361.
132. Hepburn KW, Tornatore J, Center B, et al. Dementia family caregiver training: affecting beliefs about caregiving and caregiver outcomes. *J Am Geriatr Soc*. 2001 Apr;49(4):450-7. PMID: 11347790.

133. Hepburn KW, Lewis M, Narayan S, et al. Partners in caregiving: A psychoeducation program affecting dementia family caregivers' distress and caregiving outlook. *Clin Gerontol*. 2005;29(1):53-69. doi: http://dx.doi.org/10.1300/J018v29n01_05. PMID: 43539927.
134. Hepburn K, Lewis M, Tornatore J, et al. The Savvy Caregiver program: the demonstrated effectiveness of a transportable dementia caregiver psychoeducation program. *J Gerontol Nurs*. 2007 Mar;33(3):30-6. PMID: 17378189.
135. Liddle J, Smith-Conway ER, Baker R, et al. Memory and communication support strategies in dementia: effect of a training program for informal caregivers. *Int Psychogeriatr*. 2012;24(9):1927-42. doi: 10.1017/S1041610212001366. PMID: 104376604. Language: English. Entry Date: 20130111. Revision Date: 20161117. Publication Type: journal article.
136. Klodnicka Kouri K, Ducharme FC, Giroux F. A psycho-educational intervention focused on communication for caregivers of a family member in the early stage of Alzheimer's disease: Results of an experimental study. *Dementia: The International Journal of Social Research and Practice*. 2011 Aug;10(3):435-53. doi: <http://dx.doi.org/10.1177/1471301211408124>. PMID: 2011-16142-011.
137. Neely A, Vikstrom S, Josephsson S. Collaborative memory intervention in dementia: caregiver participation matters. In *Neuropsychol*
138. Done DJ, Thomas JA. Training in communication skills for informal carers of people suffering from dementia: a cluster randomized clinical trial comparing a therapist led workshop and a booklet. *Int J Geriatr Psychiatry*. 2001 Aug;16(8):816-21. PMID: 11536349.
139. Gormley N, Lyons D, Howard R. Behavioural management of aggression in dementia: a randomized controlled trial. *Age Ageing*. 2001 Mar;30(2):141-5. PMID: 11395344.
140. Burgener SC, Bakas T, Murray C, et al. Effective caregiving approaches for patients with Alzheimer's disease. *Geriatr Nurs*. 1998 May-Jun;19(3):121-6. PMID: 9708136.
141. Robinson K, Yates K. Effects of two caregiver-training programs on burden and attitude toward help. *Arch Psychiatr Nurs*. 1994 Oct;8(5):312-9. PMID: 7993133.
142. Jablonski RA, Reed D, Maas ML. Care intervention for older adults with Alzheimer's disease and related dementias: effect of family involvement on cognitive and functional outcomes in nursing homes. *J Gerontol Nurs*. 2005 Jun;31(6):38-48. PMID: 16138529.
143. Maas ML, Reed D, Park M, et al. Outcomes of family involvement in care intervention for caregivers of individuals with dementia. *Nurs Res*. 2004 Mar-Apr;53(2):76-86. PMID: 15084992.
144. Bramble M, Moyle W, Shum D. A quasi-experimental design trial exploring the effect of a partnership intervention on family and staff well-being in long-term dementia care. *Aging Ment Health*. 2011 Nov;15(8):995-1007. doi: <https://dx.doi.org/10.1080/13607863.2011.583625>. PMID: 21702706.
145. Robison J, Curry L, Gruman C, et al. Partners in caregiving in a special care environment: cooperative communication between staff and families on dementia units. *Gerontologist*. 2007 Aug;47(4):504-15. PMID: 17766671.
146. McCallion P, Toseland RW, Freeman K. An evaluation of a family visit education program. *J Am Geriatr Soc*. 1999 Feb;47(2):203-14. PMID: 9988292.
147. Kunik ME, Snow AL, Wilson N, et al. Teaching Caregivers of Persons with Dementia to Address Pain. *Am J Geriatr Psychiatry*. 2017 Feb;25(2):144-54. doi: <https://dx.doi.org/10.1016/j.jagp.2016.04.009>. PMID: 27743840.
148. Teri L, McCurry SM, Logsdon R, et al. Training community consultants to help family members improve dementia care: a randomized controlled trial. *Gerontologist*. 2005 Dec;45(6):802-11. PMID: 16326662.
149. Teri L, Huda P, Gibbons L, et al. STAR: a dementia-specific training program for staff in assisted living residences. *Gerontologist*. 2005 Oct;45(5):686-93. PMID: 16199404.
150. Teri L, McKenzie G, Logsdon RG, et al. Translation of two evidence-based programs for training families to improve care of persons with dementia. *Gerontologist*. 2012 Aug;52(4):452-9. doi: <https://dx.doi.org/10.1093/geront/gnr132>. PMID: 22247431.
151. Livingston G, Barber JA, Kinnunen KM, et al. DREAMS-START (Dementia RElAted Manual for Sleep; STrAtegies for RelaTives) for people with dementia and sleep disturbances: a single-blind feasibility and acceptability randomized controlled trial. *Int Psychogeriatr*. 2019 02;31(2):251-65. doi: <https://dx.doi.org/10.1017/S1041610218000753>. PMID: 30221615.
152. Kinnunen KM, Rapaport P, Webster L, et al. A manual-based intervention for carers of people with dementia and sleep disturbances: An acceptability and feasibility RCT. *Health Technology Assessment*. 2018 December;22(71):I-408. doi: <http://dx.doi.org/10.3310/hta22710>. PMID: 626134933.
153. Livingston G, Barber JA, Kinnunen KM, et al. DREAMS-START (Dementia RElAted Manual for Sleep; STrAtegies for RelaTives) for people with dementia and sleep disturbances: a single-blind feasibility and acceptability randomized controlled trial. *Int Psychogeriatr*. 2018 Sep 17:1-15. doi: <https://dx.doi.org/10.1017/S1041610218000753>. PMID: 30221615.
154. Suominen MH, Puranen TM, Jyvakorpi SK, et al. Nutritional Guidance Improves Nutrient Intake and Quality of Life, and May Prevent Falls in Aged Persons with Alzheimer Disease Living with a Spouse (NuAD Trial). *J Nutr Health Aging*. 2015 Nov;19(9):901-7. doi: <https://dx.doi.org/10.1007/s12603-015-0558-0>. PMID: 26482691.
155. McCurry SM, Gibbons LE, Logsdon RG, et al. Nighttime insomnia treatment and education for Alzheimer's disease: a randomized, controlled trial. *J Am Geriatr Soc*. 2005 May;53(5):793-802. PMID: 15877554.
156. McCurry SM, Gibbons LE, Logsdon RG, et al. Training caregivers to change the sleep hygiene practices of patients with dementia: the NITE-AD project. *J Am Geriatr Soc*. 2003 Oct;51(10):1455-60. PMID: 14511168.