

# *Draft Comparative Effectiveness Review*

---

Number xx

## **Interventions Targeting Sensory Challenges in Children with Autism Spectrum Disorder (ASD)—An Update**

**Prepared for:**

Agency for Healthcare Research and Quality  
U.S. Department of Health and Human Services  
5600 Fishers Lane Rockville, MD 20857  
www.ahrq.gov

This information is distributed solely for the purposes of prepublication peer review. It has not been formally disseminated by the Agency for Healthcare Research and Quality. The findings are subject to change based on the literature identified in the interim and peer-review/public comments and should not be referenced as definitive. It does not represent and should not be construed to represent an Agency for Healthcare Research and Quality or Department of Health and Human Services (AHRQ) determination or policy.

**Contract No.** [redacted for peer review]

**Prepared by:**

[redacted for peer review]

**Investigators:**

[redacted for peer review]

**AHRQ Publication No. xx-EHCxxx**  
**<Month Year>**

This report is based on research conducted by an Evidence-based Practice Center (EPC) under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. xxx-xxxx-xxxxx). The findings and conclusions in this document are those of the authors, who are responsible for its contents; the findings and conclusions do not necessarily represent the views of AHRQ. Therefore, no statement in this report should be construed as an official position of AHRQ or of the U.S. Department of Health and Human Services.

**None of the investigators have any affiliations or financial involvement that conflicts with the material presented in this report.**

The information in this report is intended to help health care decisionmakers—patients and clinicians, health system leaders, and policymakers, among others—make well-informed decisions and thereby improve the quality of health care services. This report is not intended to be a substitute for the application of clinical judgment. Anyone who makes decisions concerning the provision of clinical care should consider this report in the same way as any medical reference and in conjunction with all other pertinent information, i.e., in the context of available resources and circumstances presented by individual patients.

This report is made available to the public under the terms of a licensing agreement between the author and the Agency for Healthcare Research and Quality. This report may be used and reprinted without permission except those copyrighted materials that are clearly noted in the report. Further reproduction of those copyrighted materials is prohibited without the express permission of copyright holders.

AHRQ or U.S. Department of Health and Human Services endorsement of any derivative products that may be developed from this report, such as clinical practice guidelines, other quality enhancement tools, or reimbursement or coverage policies may not be stated or implied

This report may periodically be assessed for the currency of conclusions. If an assessment is done, the resulting surveillance report describing the methodology and findings will be found on the Effective Health Care Program Web site at [www.effectivehealthcare.ahrq.gov](http://www.effectivehealthcare.ahrq.gov). Search on the title of the report.

Persons using assistive technology may not be able to fully access information in this report. For assistance contact [EffectiveHealthCare@ahrq.hhs.gov](mailto:EffectiveHealthCare@ahrq.hhs.gov).

**Suggested citation:**

The Agency for Healthcare Research and Quality (AHRQ), through its Evidence-based Practice Centers (EPCs), sponsors the development of systematic reviews to assist public- and private-sector organizations in their efforts to improve the quality of health care in the United States. These reviews provide comprehensive, science-based information on common, costly medical conditions, and new health care technologies and strategies.

Systematic reviews are the building blocks underlying evidence-based practice; they focus attention on the strength and limits of evidence from research studies about the effectiveness and safety of a clinical intervention. In the context of developing recommendations for practice, systematic reviews can help clarify whether assertions about the value of the intervention are based on strong evidence from clinical studies. For more information about AHRQ EPC systematic reviews, see [www.effectivehealthcare.ahrq.gov/reference/purpose.cfm](http://www.effectivehealthcare.ahrq.gov/reference/purpose.cfm)

AHRQ expects that these systematic reviews will be helpful to health plans, providers, purchasers, government programs, and the health care system as a whole. Transparency and stakeholder input are essential to the Effective Health Care Program. Please visit the Web site ([www.effectivehealthcare.ahrq.gov](http://www.effectivehealthcare.ahrq.gov)) to see draft research questions and reports or to join an e-mail list to learn about new program products and opportunities for input.

If you have comments on this systematic review, they may be sent by mail to the Task Order Officer named below at: Agency for Healthcare Research and Quality, 5600 Fishers Lane, Rockville, MD 20857, or by email to [epc@ahrq.hhs.gov](mailto:epc@ahrq.hhs.gov).

Andrew Bindman, M.D.  
Director  
Agency for Healthcare Research and Quality

Arlene Bierman, M.D., M.S.  
Director  
Center for Evidence and Practice Improvement  
Agency for Healthcare Research and Quality

Stephanie Chang, M.D., M.P.H.  
Director, EPC Program  
Center for Evidence and Practice Improvement  
Agency for Healthcare Research and Quality

Laura Pincock, Pharm.D., M.P.H.  
Task Order Officer  
Center for Evidence and Practice Improvement  
Agency for Healthcare Research and Quality

Elisabeth Kato, M.D., M.R.P.  
Task Order Officer  
Center for Evidence and Practice Improvement  
Agency for Healthcare Research and Quality

## **Acknowledgments**

The authors gratefully acknowledge the following individuals for their contributions to this project: [redacted for peer review].

## **Key Informants**

[redacted for peer review]

## **Technical Expert Panel**

[redacted for peer review]

## **Peer Reviewers**

[redacted for peer review]

# Interventions Targeting Sensory Challenges in Children with Autism Spectrum Disorder—An Update

## Structured Abstract

**Objectives.** To evaluate the effectiveness and safety of interventions targeting sensory challenges in children with Autism Spectrum Disorder (ASD).

**Data sources.** We searched MEDLINE, EMBASE, the Cumulative Index of Nursing and Allied Health Literature, and PsycInfo from January 2010–November 2015.

**Review methods.** We included studies comparing interventions incorporating sensory-focused modalities with alternate treatments or no treatment. Studies had to include at least 10 children with ASD between 2 and 12 years old. Two investigators independently screened studies and rated risk of bias. We extracted and summarized data qualitatively given significant heterogeneity. We also assessed strength of the evidence (SOE).

**Results.** We identified 21 unique comparative studies (14 newly published studies and 7 studies addressed in our 2011 review of therapies for children with ASD). Studies included 18 randomized controlled trials (RCTs), one nonrandomized trial, and two retrospective cohort studies (3 low, 11 moderate, and 7 high risk of bias). Populations, intervention approaches, and outcomes assessed varied across studies. Relative to usual care, motor skills, or massage interventions, most studies addressing sensory integration-based approaches reported improvements in sensory and motor skills-related measures in children receiving sensory integration-based treatment in the short-term. Two small RCTs (low and moderate risk of bias) of environmental enrichment reported improvements in cognitive skills in treated children compared with standard care. Three small RCTs (two with moderate and one with high risk of bias) of auditory integration-based approaches reported mixed effects on language outcomes. Two small RCTs and one nonrandomized trial of music therapy reported some significant effects on measures of behavior and joint attention with music-based intervention compared with control interventions in studies using different protocols and addressing different outcomes. Five studies of massage (with likely overlapping populations in four studies) reported improvements in ASD symptom severity and sensory challenges associated with massage versus a waitlist control condition. Additional interventions with sensory-related components (tactile stimulation exercises, weighted blankets) reported few significant differences between treatment groups. Only one study reported harms of interventions and noted a transient rash associated with use of a weighted blanket.

**Conclusions.** Some interventions targeting sensory challenges may lead to modest short-term (< 12 months) improvements primarily in sensory- and ASD symptom severity-related outcomes; however, the evidence base for any category of intervention is small, and durability of effects beyond the immediate intervention period is unclear. Sensory integration-based approaches improved outcomes related to sensory challenges (low strength of evidence) and motor skills (low strength of evidence), and studies of massage reported improvements in sensory responses (low strength of evidence) and ASD symptoms (low strength of evidence). Environmental enrichment was also associated with improvements in cognitive skills in the short-term (low

strength of evidence). Some positive effects were associated with other approaches studied (auditory integration, music therapy, weighted blankets) but findings in these small studies were not consistent (insufficient strength of evidence). Data on harms of interventions and longer-term results are lacking, as are data on characteristics that modify outcomes, effectiveness of interventions across environments or contexts, and components of interventions that may drive effects. In sum, while some therapies hold promise and warrant further study, substantial needs exist for continuing improvements in methodologic rigor in the field.

# Contents

<b>Introduction</b> .....	1
Background.....	1
Treatment of ASD.....	1
Scope and Key Questions .....	2
Scope of Review .....	2
Key Questions (KQ) .....	2
Categorization of Interventions.....	3
Analytic Framework .....	4
Uses of This Evidence Report.....	5
<b>Methods</b> .....	6
Topic Surveillance and Review Protocol.....	6
Literature Search Strategy.....	7
Search Strategy .....	7
Inclusion and Exclusion Criteria.....	7
Study Selection .....	8
Data Extraction .....	8
Data Synthesis.....	8
Risk of Bias Assessment of Individual Studies .....	8
Determining Overall Risk of Bias Ratings .....	9
Strength of the Body of Evidence.....	9
Applicability .....	9
Peer Review and Public Commentary .....	10
<b>Results</b> .....	11
Results of Literature Searches for Key Questions .....	11
Description of Included Studies.....	12
Gray Literature.....	13
Key Question 1. Benefits and Harms of Interventions Targeting Sensory Challenges .....	13
Studies of Sensory Integration-Based Approaches.....	13
Studies of Environmental Enrichment-Based Approaches.....	14
Studies of Auditory Integration-Based Approaches .....	15
Studies of Music Therapy-Base Approaches .....	16
Studies of Touch/Massage .....	17
Additional Studies.....	19
Key Question 2. Modifiers of Treatment Outcomes.....	20
Key Question 3. Time to Effect of Interventions.....	21
Key Question 4. Evidence that Effects Measured at the End of Treatment Predict Long-Term Functional Outcomes .....	21
Key Question 5. Effectiveness Across Environments or Contexts.....	21
Key Question 6. Drivers of Treatment Outcomes .....	21
<b>Discussion</b> .....	22
State of the Literature.....	22
Summary of Key Findings and Strength of the Evidence.....	22
KQ1. Benefits and Harms of Interventions Targeting Sensory Challenges .....	22
Other Key Questions.....	27
Findings in Relation to What is Already Known.....	27

Applicability .....	29
Implications for Clinical and Policy Decisionmaking .....	30
Limitations of the Comparative Effectiveness Review Process .....	30
Limitations of the Evidence Base .....	31
Research Gaps and Areas for Future Research.....	32
Conclusions.....	32
<b>References</b> .....	34
<b>Acronyms and Abbreviations</b> .....	35

**Tables**

Table 1. Inclusion criteria .....	7
Table 2. Overview of studies addressing interventions targeting sensory challenges .....	12
Table 3. Strength of the evidence for sensory integration-based interventions .....	23
Table 4. Strength of the evidence for environmental enrichment interventions .....	24
Table 5. Strength of the evidence for auditory integration-based interventions.....	25
Table 6. Strength of the evidence for massage-based interventions .....	26

**Figures**

Figure 1. Analytic framework for review .....	4
Figure 2. Disposition of studies identified for this review .....	11

**Appendixes**

- Appendix A. Detailed Methods
- Appendix B. Search Strategies
- Appendix C. Screening and Quality Assessment Forms
- Appendix D. Excluded Studies
- Appendix E. Risk of Bias Ratings
- Appendix F. Applicability Tables
- Appendix G. Detailed Tables of Findings

# Introduction

## Background

Autism Spectrum Disorder (ASD) is a neurodevelopmental disorder broadly defined by impaired social communication as well as restricted or repetitive patterns of behavior and interest. As defined by the *Diagnostic and Statistical Manual of Mental Disorders*, Fifth edition (DSM-5), specific features of ASD include deficits in social and emotional reciprocity (e.g., atypical social approaches, conversational impairment, atypical sharing of interests, attention, and affect); deficits in nonverbal communication (e.g., poorly integrated verbal and nonverbal communication, atypical body-language and gesture use, deficits in use and understanding of nonverbal communication), and deficits in maintaining appropriate relationships (e.g., challenges with peer interest, vulnerabilities forming friendships, difficulties adjusting behavior to suit social contexts) as well as restricted and repetitive patterns of behavior such as stereotyped speech, motor movements, or use of objects; excessive adherence to routine or insistence on sameness; intense interest patterns; and atypical sensory interests or responses.<sup>1</sup> Symptoms of the disorder impair and limit everyday functioning and are thought to be evident in early childhood, although they may not be fully evident until later ages. Although not a core symptom, many children with ASD may also have significant cognitive impairment.

Children with ASD may experience impairments in processing sensory stimuli, including intense interests in or aversion to certain types of sensory input; while somewhat challenging to operationalize, quantify, and measure clinically, estimates of impairments related to sensory processing have ranged from 42 percent to 88 percent of people with ASD and include both hyper- and hypo-responsiveness.<sup>2-4</sup> While common and impairing features of ASD for many individuals, the exact nature of sensory integration in the development and lifespan trajectory is less understood with the field historically lacking, accepted frameworks for diagnosing sensory challenges (e.g., not part of DSM diagnostic criteria until DSM-5) and developing responsive interventions.<sup>2, 3, 5, 6</sup>

## Treatment of ASD

The manifestation and severity of symptoms of ASD differ widely, and treatments include a range of behavioral, psychosocial, educational, medical, and complementary approaches<sup>7-10</sup> that vary by a child's age and developmental status. The goals of treatment for ASD typically focus on improving core deficits in communication, social interactions, or restricted behaviors, as changing these fundamental deficits may help children develop greater functional skills and independence.<sup>11</sup> Treatment frequently is complicated by symptoms or comorbidities that may warrant targeted intervention. There is no cure for ASD and no global consensus on which intervention is most effective.<sup>12, 13</sup> Individual goals for treatment vary for different children and may include combinations of behavioral therapies, educational therapies, medical and related therapies, approaches targeting sensory issues, and allied health therapies; parents may also pursue complementary and alternative medicine therapies.

## Interventions Targeting Sensory Challenges

Increasingly, as reflected in their inclusion in the new DSM-5 diagnostic criteria, the sensory challenges associated with ASD have also become a target for specialized assessment and treatment. Sensory symptoms can involve both strong interests as well as strong aversions, with

interventions commonly targeting aversions/challenges, meeting needs for sensory input within adaptive frameworks, or perceived processing deficits with the goal of improving people's abilities to interact with their environments. For example, a child with ASD may have difficulty tolerating bright lights, clothing or food textures, specific noises (such as a baby crying), tasks of daily living (such as brushing hair or teeth), touch, or more idiosyncratic stimuli such as certain colors. These sensitivities can significantly interfere with children's ability to care for themselves, leave the home, participate in school or other interventions, and be involved in social situations. Children may also display a hyperfocus on play or activities that involves a sensory component, sometimes referred to as 'sensory-seeking' or 'stimming' behaviors.

Sensory-focused interventions take a variety of forms and can be implemented by a variety of licensed professionals (such as occupational therapists), teachers, parents, and other providers. Such interventions are not consistently defined but typically involve the incorporation of sensory experiences (e.g., weighted clothing or materials, interventions that provide auditory sensations) to affect a variety of outcomes including adaptive behavior and language.

## **Scope and Key Questions**

### **Scope of Review**

This review updates findings reported in the 2011 Agency for Healthcare Research and Quality (AHRQ) review Therapies for Children with ASD<sup>14</sup> with a focus on studies of interventions targeting sensory challenges. We defined interventions targeting sensory challenges in line with the DSM-5 definition and definitions used in other reviews of sensory-focused interventions.<sup>2,3</sup> DSM-5 classifies sensory challenges as a manifestation of the core symptom of restricted and repetitive patterns of behavior, interests, or activities. The DSM describes sensory challenges as "hyper- or hyporeactivity to sensory input, manifested through extreme responses to specific sounds or textures, excessive smelling or touching of objects, fascination with lights or spinning objects, and sometimes apparent indifference to pain, heat, or cold."<sup>1</sup> Interventions targeting sensory challenges are typically described as designed to provide controlled sensory experiences in order to encourage the modulation and integration of information from the environment, thus promoting adaptive responses to sensory inputs.

Though the field lacks consensus on a definition of sensory-focused approaches, interventions typically use sensory modalities to target behaviors that may be associated with sensory-related impairments.<sup>3,15</sup> We do not include studies of other approaches (e.g., educational interventions) that may address a sensory-related outcome in the current review. A companion review updating findings related to medical interventions is available on the AHRQ Effective Health Care web site.

### **Key Questions (KQ)**

We developed KQs in consultation with Key Informants and the Task Order Officer. KQs were posted for review to the AHRQ Effective Health Care website.

KQs were as follows:

KQ1: Among children ages 2-12 with ASD, what is the comparative effectiveness (benefits and harms) of interventions targeting sensory challenges?

- a) What are the effects on core symptoms (e.g., deficits in social communication and interaction; restricted, repetitive patterns of behavior, interests, or activities including hyper- or hypo- reactivity to sensory input or unusual interest in sensory aspects of the environment) in the short term ( $\leq 6$  months)?
- b) What are the effects on commonly associated symptoms (e.g., motor, medical, mood/anxiety, irritability, and hyperactivity) in the short term ( $\leq 6$  months)?
- c) What are the longer-term effects ( $> 6$  months) on core symptoms (e.g., social deficits, communication deficits, and repetitive behaviors)?
- d) What are the longer-term effects ( $> 6$  months) on commonly associated symptoms (e.g., motor, medical, mood/anxiety, irritability, and hyperactivity)?

KQ2: Among children ages 2-12 with ASD, what are the modifiers of outcome for different interventions targeting sensory challenges?

- a) Is the effectiveness of the therapies reviewed affected by the frequency, duration, intensity, or dose of the intervention?
- b) Is the effectiveness of the therapies reviewed affected by co-interventions or prior treatment or the training and/or experience of the individual providing the therapy?
- c) What characteristics (e.g., age, symptom severity), if any, of the child modify the effectiveness of the therapies reviewed?
- d) What characteristics, if any, of the family modify the effectiveness of the therapies reviewed?

KQ3: What is the time to effect of interventions targeting sensory challenges?

KQ4: What is the evidence that effects measured at the end of the treatment phase predict long-term functional outcomes of interventions targeting sensory challenges?

KQ5: Is the effectiveness of interventions targeting sensory challenges maintained across environments or contexts (e.g., people, places, materials)?

KQ6: What evidence supports specific components of treatment with interventions targeting sensory challenges as driving outcomes, either within a single treatment or across treatments?

## **Categorization of Interventions**

Interventions targeting sensory challenges may be broadly categorized by their core focus (e.g., environmental modification/adaptation, compensatory strategies, sensory processing, auditory integration). While some approaches may combine multiple components, it is important to note that sensory interventions cannot be described using a blanket approach, but rather should be evaluated based on the specific foci of the intervention. As noted, sensory-related approaches may include overlapping targets and intervention constructs, and it is difficult to cleanly identify the category into which an intervention should be placed. Based on the literature meeting criteria for this review, we categorized interventions as:

- Sensory integration-based (interventions using combinations of sensory and kinetic components such as materials with different textures, touch/massage, swinging and

trampoline exercises, and balance and muscle resistance exercises to ameliorate sensory challenges)


- Environmental enrichment-based (interventions incorporating targeted exposure to sensory stimuli to promote tolerance of stimuli in other contexts)
- Auditory integration-based (interventions incorporating auditory components such as filtered sound to ameliorate sensory processing challenges via theorized re-training of aural pathways)
- Music therapy-based (interventions incorporating playing or singing music, or movement to music, to improve challenging behaviors and sensory difficulties)
- Touch/Massage-based (interventions incorporating touch-based approaches by a therapist or caregiver)
- Other (included interventions [tactile-based tasks, weighted blankets] not cleanly fitting into one of the broader categories).

We considered multiple approaches for organizing our results and note that no alternative approaches would have changed our overall findings, either in terms of outcomes or strength of evidence for any category of intervention. We recognize that other approaches to categorizing interventions targeting sensory challenges could be used.

## Analytic Framework

The analytic framework illustrates the population, interventions, outcomes, and adverse effects that guide the literature search and synthesis.

Figure 1. Analytic framework for review



ASD=autism spectrum disorder; KQ=Key Question

## **Uses of This Evidence Report**

We anticipate that the report will be of value to clinicians who treat children with ASD, who can use the report to assess the evidence for different treatment strategies. In addition, this review will be of use to the National Institutes of Health, U.S. Centers for Disease Control and Prevention, Centers for Medicare & Medicaid Services, and the Health Resources and Services Administration—all of which have offices or bureaus devoted to child health issues and who may use the report to compare treatments and determine priorities for funding. This report can bring practitioners up to date about the current state of evidence related to interventions targeting sensory challenges, and it provides an assessment of the risk of bias of studies that aim to determine outcomes of sensory-related options for the management of ASD. It will be of interest to families affected by ASD because of the recurring need for families and their health care providers to make the best possible decisions among numerous options. We also anticipate it will be of use to private sector organizations concerned with ASD; the report can inform such organizations' understanding of the effectiveness of treatments and the amount and quality of evidence available. Researchers can obtain a concise analysis of the current state of knowledge related to interventions targeting sensory challenges in ASD and of areas for future research.

## Methods

In this chapter, we briefly outline the procedures that we used to produce a comparative effectiveness review (CER) update addressing interventions targeting sensory challenges for children with autism spectrum disorder (ASD). Appendix A includes a more detailed discussion of our methods. These procedures follow the methods outlined in the Agency for Healthcare Research and Quality (AHRQ) Effective Health Care Program *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.<sup>16</sup>

### Topic Surveillance and Review Protocol

The topic for the original report<sup>17</sup> was nominated by Autism Speaks in a public process using the Effective Health Care website. AHRQ published an update addressing only behavioral interventions in 2014.<sup>18</sup> We conducted a surveillance process to assess the need to update sections of the original 2011 report by contacting topic experts about the relevance of the Key Questions (KQs) and new evidence that may address them. All members of the research team were required to submit information about potential conflicts of interest before initiation of the work. No members of the review team had any conflicts.

In consultation with clinical experts and stakeholders, and based on our preliminary scan of the literature and surveillance findings, we focused the review update on approaches to address sensory challenges and medical approaches (reported in a separate update). These areas reflect both areas of clinical relevance and sufficient newly published literature for a review update. Given the different major emphases of these interventions (i.e., sensory processing/integration abilities and challenging behaviors) and subsequent differences in study populations, we report findings in two separate reviews.

Based also on the surveillance process and discussions with stakeholders, we revised the KQ addressed in the 2011 report to reflect the focus on medical and sensory approaches specifically. We also eliminated a question on approaches for children at risk for ASD as such children are unlikely to be included in studies in the target areas for this review update.

After review from AHRQ, the questions and framework were posted online for public comment. No changes to the questions or framework were recommended. We identified technical experts on the topic to provide input during the project. The Technical Expert Panel (TEP), representing the fields of pediatrics and developmental pediatrics, psychiatry, family medicine, and occupational therapy and allied health, contributed to AHRQ's broader goals of (1) creating and maintaining science partnerships as well as public-private partnerships and (2) meeting the needs of an array of potential users of its products. Thus, the TEP was both an additional information resource and a sounding board during the project. The TEP included seven members serving as technical or clinical experts. To ensure robust, scientifically relevant work, TEP members participated in conference calls to:

- Help to refine the analytic framework and KQ at the beginning of the project;
- Discuss inclusion/exclusion criteria; and
- Assist with determining key interventions and outcomes of interest.

The final protocol was posted to the AHRQ Effective Health Care web site and registered in the PROSPERO international register of systematic reviews (ID#: CRD42016033941).

# Literature Search Strategy

## Search Strategy

To ensure comprehensive retrieval of relevant studies of therapies for children with ASD, we used four key databases: the MEDLINE<sup>®</sup> medical literature database via the PubMed<sup>®</sup> interface; EMBASE (Excerpta Medica Database), an international biomedical and pharmacological literature database via the Ovid<sup>®</sup> interface; the Cumulative Index of Nursing and Allied Health Literature (CINAHL), and PsycINFO<sup>®</sup>. Search strategies for KQs applied a combination of controlled vocabulary (Medical Subject Headings [MeSH] and Emtree headings) to focus specifically on interventions targeting sensory challenges in children with ASD and harms of interventions (Appendix B). We restricted literature searches for KQs to studies published from 2010 to November 2015 to reflect literature available since the publication of the 2011 review. We will update searches while the report is undergoing peer review.

## Gray Literature

We searched web sites of organizations likely to conduct research, issue guidance, or generate policies for ASD (e.g., Autism Speaks, the American Academy of Child and Adolescent Psychiatry) to inform the review's background and discussion sections. We searched government and regulatory agency web sites for contextual information on benefits and harms of ASD interventions. We searched ClinicalTrials.gov and other trial registries for information about relevant ongoing trials and to confirm that we have obtained available publications of results from completed trials.

## Inclusion and Exclusion Criteria

Table 1 outlines inclusion criteria. We required that eligible RCTs have a total minimum sample size of 10. We required a higher minimum sample size (n=20) for other comparative studies as they typically have fewer controls for bias than RCTs. We recognize that these study design criteria excluded single-subject or single-case experimental designs that have been used to study interventions targeting sensory challenges. These studies are challenging to incorporate in a meaningful way in comparative effectiveness reviews, which attempt to evaluate the effectiveness of interventions at the population level. To mitigate the exclusion of such studies; however, we include summaries of recent reviews that have included such studies and discuss our findings in light of those in other reviews (see Findings in Relation to What is Known). We included studies published in English only. In the opinion of our content experts, much of the relevant literature on ASD is published in English. Eligible studies also reported one or more outcomes of interest and included children at least 2 years of age and up to and including age 12. Studies also included only children with a diagnosis of ASD (or data reported separately for children with ASD).

**Table 1. Inclusion criteria**

Category	Criteria
Study population	Children ages 2-12 with ASD (mean age plus standard deviation is $\leq$ 12 years and 11 months)
Publication languages	English only
Admissible evidence (study design and other)	Admissible designs Randomized controlled trials, prospective and retrospective cohort studies with

Category	Criteria
criteria)	<p>comparison groups, and nonrandomized controlled trials</p> <p>Other criteria Original research studies published from 2010—present and not addressed in prior reviews</p> <p>Studies must have relevant population and ≥20 participants with ASD (non-RCTs) or at least 10 total participants (RCTs)</p> <p>Studies must address one or more of the following for ASD:</p> <ul style="list-style-type: none"> <li>-Outcomes of interest</li> <li>-Treatment modality of interest</li> <li>-Predictors or drivers of treatment outcomes (e.g., biomarkers, clinical changes)</li> <li>-Maintenance of outcomes across environments or contexts</li> <li>-Sufficiently detailed methods and results to enable data extraction</li> <li>-Reporting of outcome data by target population or intervention</li> </ul>

ASD=Autism Spectrum Disorder; RCT=randomized controlled trial

## Study Selection

Two reviewers separately evaluated the abstracts of studies identified in our searches for Key Questions (KQs) for inclusion or exclusion, using an Abstract Review Form (Appendix C). If one reviewer concluded that the article could be eligible for the review based on the abstract, we retained it. Following abstract review, two reviewers independently assessed the full text of each included study using a standardized form (Appendix C) that included questions stemming from our inclusion and exclusion criteria. A senior reviewer resolved disagreements between reviewers. Appendix D includes a list of excluded studies and the reasons for exclusion. Data extracted for each study are available via the Systematic Review Data Repository (<http://srdhr.gov/>).

## Data Extraction

We designed extraction forms to provide sufficient information to enable readers to understand the studies and to determine their quality; we gave particular emphasis to essential information related to the KQs. Team data extractors shared the task of initially entering information into the evidence tables. A second team member also reviewed the articles and edited all initial entries for accuracy, completeness, and consistency. A senior reviewer reconciled disagreements concerning the information reported.

## Data Synthesis

Studies were too heterogeneous to allow for meta-analyses. We summarized data for Key Questions qualitatively using summary tables.

## Risk of Bias Assessment of Individual Studies

We evaluated the overall methodologic risk of bias of individual studies using the ASD-specific assessment approach developed and used in our prior reviews of interventions for ASD and informed by the *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.<sup>16</sup> We developed this tool because standard risk of bias assessment tools (e.g., Cochrane risk of bias assessment) do not fully account for the complexity of interventions and populations represented

in the ASD literature. Specifically, the tool includes questions to address diagnostic approaches and measures of treatment fidelity that may affect outcomes. The tool has not been formally validated.

Two senior investigators assessed each included study independently with disagreements resolved through discussion. Appendix C includes our risk of bias assessment form, and Appendix E includes the risk of bias ratings for each study. Appendix A includes additional details about our risk of bias approach.

## Determining Overall Risk of Bias Ratings

We used the thresholds we established in prior reviews to assess overall high, medium or low risk of bias. We assessed the risk of bias based upon the study-defined primary outcome(s). We assessed each domain described above individually and considered the individual ratings to determine an overall quality assessment of low, moderate, or high risk of bias. Appendix A includes additional details.

## Strength of the Body of Evidence

The assessment of the literature is done by considering both the observed effectiveness of interventions and the confidence that we have in the stability of those effects in the face of future research (see Appendix A for full details). The degree of confidence that the observed effect of an intervention is unlikely to change is presented as strength of evidence. Methods for applying strength of evidence assessments are established in the *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*<sup>16</sup> and are based on consideration study limitations, consistency in direction of the effect, directness in measuring intended outcomes, precision of effect, and reporting bias. Strength of evidence is assessed separately for major intervention-outcome pairs and incorporates data from the entire body of reviewed evidence on behavioral interventions (i.e., comparative studies—both RCTs and prospective and retrospective cohort studies—reported in the 2011 review<sup>17</sup> and studies reported in the current review). The possible strength of evidence grades were:

- High: High confidence that the evidence reflects the true effect. Further research is unlikely to change estimates
- Moderate: Moderate confidence that the evidence reflects the true effect. Further research may change our confidence in the estimate of effect and may change the estimate
- Low: Low confidence that the evidence reflects the true effect. Further research is likely to change confidence in the estimate of effect and is also likely to change the estimate
- Insufficient: Evidence is either unavailable or does not permit a conclusion.

## Applicability

We assessed the applicability of findings reported in the included literature addressing our KQs to the general population of children with ASD by determining the population, intervention, comparator, and setting in each study and developing an overview of these elements for each intervention category. We anticipated that areas in which applicability would be especially important to describe would include ASD severity, comorbidities, age at treatment, and intervention characteristics such provider, dosing/intensity, and setting. Applicability tables for each KQ are in Appendix F.

## **Peer Review and Public Commentary**


Researchers and clinicians with expertise in treating children with ASD and individuals representing stakeholder and user communities will provide external peer review of this report. The draft report will be posted on the AHRQ Web site for 4 weeks to elicit public comment. We will address all reviewer comments, revise the text as appropriate, and document changes and revisions to the report in a disposition of comments report that will be made available 3 months after AHRQ posts the final review on the AHRQ Web site.

# Results

## Results of Literature Searches for Key Questions

We identified 4372 nonduplicative titles or abstracts with potential relevance, with 486 proceeding to full text review (Figure 2). We excluded 373 studies at full text review. We included 14 unique studies (13 publications; one publication reports two separate studies<sup>19</sup>) in the review. These 14 studies included 11 randomized controlled trials (RCTs), one nonrandomized trial, and two retrospective cohort studies. In addition to these 14 studies published since the completion of our original review of therapies for children with Autism Spectrum Disorder (ASD) in 2011,<sup>17</sup> we include seven comparative studies addressed in the 2011 review in order to present a comprehensive assessment of the literature addressing interventions targeting sensory challenges.

**Figure 2. Disposition of studies identified for this review**



Numbers next to each Key Question indicate number of unique studies addressing the question. Studies could address more than one Key Question.

\*Numbers do not tally as studies could be excluded for multiple reasons.

†One paper reports two separate trials. We also include analysis of 7 comparative studies reported in our 2011 review of therapies for children with ASD; thus we describe a total of 21 studies.

Abbreviations: KQ = key question; n = number.

## Description of Included Studies

The 21 studies addressing interventions targeting sensory challenges included 18 RCTs (including one paper that reported two separate studies<sup>19-35</sup> one nonrandomized trial,<sup>36</sup> and two retrospective cohort studies (Table 2).<sup>37, 38</sup> Three studies had low risk of bias,<sup>21, 23, 38</sup> ten had moderate,<sup>20, 22, 24-26, 28, 32-35</sup> and eight had high risk.<sup>19, 25, 29-31, 36, 37</sup> Thirteen studies were conducted in the United States,<sup>19-25, 28, 32, 33, 35, 38</sup> two in the UK,<sup>27, 34</sup> and one each in Japan,<sup>37</sup> Iran,<sup>36</sup> Turkey,<sup>30</sup> Korea,<sup>29</sup> Thailand,<sup>31</sup> and Australia.<sup>26</sup>

In total, studies included approximately 909 children (median 34 total children/study) ranging in age from 2 to 16 years. Overlap of participants in a series of massage-focused studies by one investigative team is unclear. While studies likely included some of the same children, the extent of overlap is not clear.<sup>24, 25, 32, 35, 38</sup>

Severity of ASD and sensory dysfunction varied across studies. Only two studies reported any followup of participants after the completion of treatment,<sup>25, 36</sup> and only one had more than 6 months of treatment.<sup>37</sup> Because few studies addressed sub-questions under Key Questions (KQ) 1 and 2, we present results in the aggregate under each of these KQ. Appendix G includes detailed summary tables summarizing key outcomes.

**Table 2. Overview of studies addressing interventions targeting sensory challenges**

Characteristic	RCTs (n=18)	Nonrandomized trials (n=1)	Retrospective Cohort Studies (n=2)	Total Literature
<b>Intervention category</b>				
Sensory integration-based approaches	3	0	1	4
Environmental enrichment-based approaches	2	0	0	2
Auditory integration-based approaches	4	0	0	4
Music therapy-based approaches	2	1	0	3
Massage-based approaches	5	0	1	6
Additional approaches	2	0	0	2
<b>Treatment duration</b>				
<1-4 weeks	5	1	0	6
5-8 weeks	2	0	0	2
9-12 weeks	3	0	0	3
13-20 weeks	6	0	1	7
≥21 weeks	2	0	1	3
<b>Region of Study Conduct</b>				
Asia	3	1	1	5
Australia	1	0	0	1
Europe	2	0	0	2
North America	12	0	1	13
<b>Risk of Bias</b>				
Low	2	0	0	2
Moderate	10	0	1	11
High	6	1	1	8
<b>Total N participants</b>	<b>733</b>	<b>27</b>	<b>149</b>	<b>909</b>

N = Number; RCT = Randomized Controlled Trial

## Gray Literature

Our searches of ClinicalTrials.gov and other trial registers did not yield additional eligible studies for the review. We used information from organization web sites searched to provide additional context for the discussion section of the report.

## Key Question 1. Benefits and Harms of Interventions Targeting Sensory Challenges

### Studies of Sensory Integration-Based Approaches

#### Key Points

- Four studies addressing sensory integration (SI)-based approaches were small and short-term (< 12 months), with no followup beyond the immediate treatment period. No study reported harms of intervention.
- There is low strength of evidence that sensory-related outcomes improved in children receiving a sensory integration-focused intervention compared with those receiving usual care or other treatment (significant improvements in three of four studies addressing the outcome). There is low strength of evidence that motor skills outcomes were significantly improved in children receiving sensory integration-based treatment compared with those receiving usual care or other treatment (significant improvements in three of three studies addressing the outcome).
- There is low strength of evidence that sensory integration-based treatment did not improve adaptive behavior (no significant effects in two studies addressing the outcome).

#### Overview of the Literature

We identified three RCTs (one low,<sup>21</sup> one moderate,<sup>20</sup> and one high<sup>30</sup> risk of bias) and one retrospective cohort study (high risk of bias<sup>37</sup>) addressing SI-based approaches. These studies included one RCT with high risk of bias addressed in our 2011 review.<sup>30</sup> Studies were conducted in the United States,<sup>20, 21</sup> Japan,<sup>37</sup> and Turkey<sup>30</sup> and included a total of 119 children between the ages of 2 to 12 years. Treatment duration ranged from 6 weeks to 10 months, and no study reported long-term followup after the end of treatment.

#### Detailed Analysis

In these small, short-term studies, outcomes on sensory-related measures and motor skills measures were improved in children receiving a SI-based intervention compared with another intervention in three of four studies, but effects on other outcomes were typically not significantly different between groups (Appendix G). Several outcomes were also parent-reported, and parents were often aware of intervention status.

In one low risk of bias RCT, children with autism or Pervasive Development Disorder-Not Otherwise Specified (PDD-NOS) and a diagnosed sensory processing disorder received treatment focused on SI or treatment focused on building fine motor skills.<sup>21</sup> Intervention for both groups consisted of 18 45-minute sessions over a 6-week span. Both groups improved significantly on blinded parent and teacher ratings of goal attainment related to sensory processing, motor skills, and social functioning, with children receiving SI improving significantly more than those receiving motor skills intervention. Children in the SI group had

significantly fewer parent-rated autistic mannerisms post-treatment than the fine motor group, but other measures of sensory processing, ASD symptoms, or neurological functioning did not differ between groups.

Another RCT with moderate risk of bias compared manualized occupational therapy with sensory integration (OT/SI) to care-as-usual.<sup>20</sup> OT/SI treatment consisted of three weekly sessions over the course of 10 weeks, which were monitored for treatment fidelity. Outcome measurements included parent-generated Goal Attainment Scaling (GAS). After treatment, children receiving OT/SI showed significantly more goals attained and significantly greater improvements in social skills and self-care measures compared with children receiving usual care. Scores on the Vineland Adaptive Behavior Scales (VABS) or other measures related to functional skills or caregiver assistance did not differ between groups.

In a retrospective cohort study (high risk of bias) using previously collected data to compare SI-based therapy in children with high functioning ASD (IQs > 70), both groups received active treatment that included either SI therapy or eclectic group therapy.<sup>37</sup> Treatment lasted for 8 to 10 months. Participants in the SI group improved significantly more than those in the control group in measures of motor abilities, memory and visualization, and combined sensory motor and cognitive skills conducted by an unblinded investigator (all p values < 0.05) but not for measures of spatial positioning and sense of touch or verbal ability. Finally, in a high risk of bias RCT evaluating the effects of an SI protocol on low-functioning children with ASD, children receiving SI intervention had significantly fewer sensory problems at followup than children in the usual care control group using a parent-rated scale.<sup>30</sup>

## **Studies of Environmental Enrichment-Based Approaches**

### **Key Points**

- Two small RCTs addressing environmental enrichment were short term ( $\leq 6$  months). Neither study reported harms of intervention.
- There is low strength of the evidence that environmental enrichment approaches improve cognitive skills (two low and moderate risk of bias RCTs). There is low strength of evidence that these approaches do not affect expressive language.
- Evidence is insufficient to assess effects on atypical sensory responses or receptive language as these outcomes were only addressed in one RCT.

### **Overview of the Literature**

Two RCTs (low and moderate risk of bias), both conducted by the same investigators in the United States, examined environmental enrichment.<sup>22, 23</sup> The 78 children included in studies range in age from 3 to 12 years and received treatment for 6 months, with followup immediately post-treatment in both studies. Children in the studies had specific diagnoses of autism (vs. ASD).

### **Detailed Analysis**

Two small RCTs of environmental enrichment examined the same protocol and reported improvements in ASD symptoms, receptive language, and nonverbal cognitive skills after 6 months of treatment (Appendix G). In one RCT (moderate risk of bias) comparing male children who received standard care plus sensorimotor enrichment to those who received standard care alone, the treatment group received olfactory/tactile stimulation as well as four to seven other

parent-led, sensory-stimulating exercises twice a day over the course of 6 months.<sup>22</sup> Levels of concurrent interventions (e.g., speech, behavioral, physical therapies) were similar across groups and held as stable as possible. Compared with usual care, children receiving environmental enrichment had a more significant decrease in clinician-rated ASD symptoms ( $p=0.03$ ) at the end of treatment, with nearly five times as many participants in the treatment group showing clinically significant drops of five points or more (42% vs. 7%,  $p=0.03$ ). The treatment group also had a 9-point increase in nonverbal cognitive skills as measured by the Leiter-R compared with a decrease of approximately 3 points in the usual care group ( $p=0.008$ ). Both groups improved on expressive language skills, with no significant differences.

A second RCT (low risk of bias) built upon the preliminary work by examining use of the same sensorimotor enrichment regimen over 6 months.<sup>23</sup> Investigators randomized participants to three groups: full treatment, as described in the initial study above; partial treatment, which entailed an abbreviated treatment regimen, and standard care. However, because no differences were found between intervention outcomes across the two treatment groups, the study collapsed findings into a combined treatment group. The treatment groups experienced significant attrition (> 50% across both) that may affect the generalizability of the results. After six months, the treatment group showed more improvement than did the control group in receptive language skills, but both groups improved comparatively for expressive language. The treatment group had significantly more improvement on mean nonverbal IQ scores as well as parent-rated sensory reactivity. Although more children in the treatment group compared with the control group shifted their diagnostic classification on the Autism Diagnostic Observation Schedule-2 (ADOS-2) from “autism” to “autism spectrum,” all children across both groups continued to meet the cut-offs for ASD, making it difficult to interpret the clinical significance of the findings.

## **Studies of Auditory Integration-Based Approaches**

### **Key Points**

- All four RCTs addressing auditory integration-based approaches were small and short term (<6 months). No study reported harms of intervention.
- There is low strength of evidence that auditory integration-based approaches do not improve language outcomes (three RCTs with moderate and high risk of bias).

### **Overview of the Literature**

Four RCTs with moderate<sup>33, 34</sup> and high<sup>19</sup> risk of bias evaluated auditory integration-based approaches. One paper reports two RCTs,<sup>19</sup> and two studies were included in our 2011 review.<sup>33, 34</sup> Studies included a total of 173 children between the ages of 3 and 13 years, and treatment duration ranged from 1 week to 18 weeks, with followup immediately post-treatment.

### **Detailed Analysis**

Two small, short term RCTs of auditory integration-based approaches reported no significant differences between groups in language outcomes assessed on parent, teacher, and clinician observation measures,<sup>33, 34</sup> while two studies reported significant parent-rated improvements in hearing sensitivity and behavior (Appendix G).<sup>19</sup>

One crossover RCT evaluated the effects of Tomatis Sound Therapy on language skills in children with autistic disorder who had not previously had auditory stimulation treatments.<sup>33</sup> In the treatment condition, children listened to music passed through an electronic ear for

attenuation and modulation for two hours per day in accordance with the Tomatis Method protocol. In the placebo condition, children listened to commercially produced music. The study reported no significant group effects. Another RCT of auditory integration therapy including children with significant language delays reported no significant benefits of auditory integration.<sup>34</sup>

Two high risk of bias RCTs (reported in a single publication<sup>19</sup>) examined the use of auditory integration strategies theorized to reduce auditory hypersensitivity by “exercising” the neural regulation abilities of the middle ear muscles. Across both trials, two different sets of participants completed five 45-minute sessions on consecutive days. In Trial 1, participants wore headphones and either listened to filtered music or no sound. In Trial 2, participants either listened to filtered music or unfiltered music. Participants either had at least five words of functional speech or were able to follow one-step instructions. It should be noted that although the ADI-R was used to confirm diagnosis, a subset of participants did not meet full diagnostic cut-offs on this instrument. One week after intervention, parents reported more improvement in the areas of hearing sensitivity, spontaneous speech, listening, and behavioral organization after filtered music compared with children in the control condition (p values <0.01). Children who received filtered music in Trial 2 also had significantly better parent-rated scores on hearing sensitivity and emotional control compared with control children (p values <0.05). Groups in either trial did not differ in the other behavioral domains rated.

## **Studies of Music Therapy-Based Approaches**

### **Key Points**

- All studies addressing music therapy-based approaches were small and short term (< 6 months), and none reported harms of intervention.
- There is insufficient evidence to assess effects of music therapy approaches on any outcome given that two of the three studies addressing these interventions had high risk of bias. Studies also had multiple comparators and addressed different outcomes.

### **Overview of the Literature**

Two RCTs and one nonrandomized trial (one conducted in Australia,<sup>26</sup> one in Iran,<sup>36</sup> and one in Korea<sup>29</sup>) examined music therapy. One nonrandomized trial comparing music therapy and toy play was included in our 2011 review.<sup>29</sup> Studies included a total of 58 children ranging in age from 3 to 12 years, and treatment duration ranged from 45 days to 16 weeks. Followup occurred immediately at the end of treatment in all but one of the studies, which followed up at 2 months post-treatment.<sup>36</sup> One study had moderate risk of bias,<sup>26</sup> and two studies had high risk.<sup>29, 36</sup>

### **Detailed Analysis**

The three small studies addressing music therapy reported some significant effects on measures of behavior (social engagement, behavioral organization), and joint attention with music-based intervention compared with control interventions (Appendix G). Studies used different protocols and addressed different outcomes; thus, drawing conclusions across studies is challenging. In some studies children also received other interventions in addition to music-based approaches.

One RCT (moderate risk of bias) compared family-centered music therapy plus early intervention to early intervention only.<sup>26</sup> Participants had little to no functional verbal

communication and received 2 to 3 hours of community-based intervention per week while participating. Intervention consisted of one hour per week of semi-structured sessions within the home for 16 weeks. Therapists worked with parents and participants to improve selected core social engagement skills through music-based activities. Children who received music therapy had more improvement than controls in parent-rated social engagement ( $p < 0.001$ ) but remained significantly impaired relative to typically developing peers. Groups did not differ on parent-reported autism symptoms, speech and language, or quality of the parent-child relationship.

In another high risk of bias nonrandomized trial comparing 12 1-hour Orff-Schulwerk music therapy sessions over the course of 45 days to no treatment, social skills improved significantly in the treatment group between baseline to post-treatment but not from post-treatment to follow-up.<sup>36</sup> The control group did not improve at any time point, and differences between the treatment and control groups were not significant at the final followup.

In a final crossover RCT with high risk of bias comparing music therapy and toy play, groups did not differ on the Pervasive Development Disorder Behavior Inventory (PDDBI), though both groups improved with time.<sup>29</sup> Results from the Early Social Communication Scales, reflecting growth in joint attention skills, suggested that music therapy was significantly more effective than play sessions. Change scores pre- to post- music therapy were significantly greater than change scores pre- to post- play sessions. In coding for emotional and motivational responsiveness (i.e., joy, emotional synchronicity, initiation of engagement), investigators observed more joy, emotional synchronicity, and initiation of engagement during music therapy than in play sessions. In addition, children had significantly more compliant behavior and significantly fewer episodes of no response behaviors in the music therapy condition.

## Studies of Touch/Massage

### Key Points

- One group of investigators conducted five short-term (< 6 months treatment duration) studies of massage, and participant overlap is unclear. No study reported harms of intervention.
- There is low strength of the evidence that massage improves sensory challenges and ASD symptom severity.
- There is low strength of evidence that massage does not improve maladaptive behavior.
- Evidence is not sufficient to assess effects of massage on language/communication outcomes given inconsistent findings and use of different outcome measures.
- Evidence is insufficient to assess longer-term outcomes (>6 months) as only one study reported longer-term followup. Evidence is insufficient to assess effects on measures of daily living skills, also assessed in only one study. Only one study compared massage plus SI-based treatment to SI-based treatment alone, so evidence is insufficient to assess effects.

### Overview of the Literature

Six studies (5 RCTs and one retrospective cohort study) addressed touch-based therapy. Five studies (four RCTs and one cohort) compared Qigong or traditional massage to no massage treatment.<sup>24, 25, 32, 35, 38</sup> One RCT compared massage plus SI-based treatment with SI-based treatment alone.<sup>31</sup> Studies were conducted in the United States<sup>24, 25, 32, 35, 38</sup> and Thailand<sup>31</sup> and had moderate<sup>24, 32, 35</sup> and high risk of bias.<sup>25, 31, 38</sup> Three RCTs were also reported in our 2011 review.<sup>25, 31, 32</sup>

One team of investigators has published most of the literature in this area, and the extent of overlap among participants in these studies is unclear.<sup>24, 25, 32, 38, 39</sup> Studies included approximately 395 children receiving treatment for 2 to 5 months. One study reported followup of participants 5 months after the end of treatment.<sup>25</sup>

## Detailed Analysis

**Massage vs. No Massage.** Almost all of studies are from one group of investigators, and the participant overlap is unclear (Appendix F). Some used only parent-and teacher-rated outcomes, but others used standardized measures of autism symptoms, language, and adaptive functioning (e.g., Childhood Autism Rating Scale [CARS], Preschool Language Scale [PLS], Vineland Adaptive Behavior Scales [VABS]). Studies generally reported improvements related to sensory processing, autism symptoms, and parent stress in both treatment and control groups over the course of 5 months of either parent or parent + therapist-delivered intervention, with treatment groups improving significantly more than controls. Some studies also examined the moderating influence of baseline variables, such as parent stress or autism severity, which are important confounds to explore. The difficulty differentiating populations in these studies limits the strength of evidence for their findings, although results seem promising regarding a sensory-focused intervention that can be delivered within the home environment with minimal risk of harms.

As noted, four RCTs (three with a moderate risk of bias<sup>24, 35, 38</sup> and one with a high risk of bias<sup>25</sup>) had unclear participant overlap and compared children who received Qigong Sensory Training (QST) to wait-listed controls or usual care. In one study comparing children who received QST to children receiving treatment as usual,<sup>24</sup> parents participated in seven sessions instructing them in the basics of Qigong. Most children concurrently attended early intervention preschools for 5-10 hours a week, and participants included an unknown number of sibling pairs. Children receiving QST significantly improved from baseline on teacher and parent ratings of autism symptoms as well as parent ratings of sensory challenges, self-regulation skills, communication skills, and maladaptive behaviors. When compared with the waitlist control group, improvements were significantly larger for autism symptoms (PDDBI), parent stress, and sensory/self-regulation challenges (Sensory and Self-Regulation Checklist), with medium to large effect sizes. Changes in teacher-rated ASD symptoms (Autism Behavior Checklist [ABC]) were not significantly different between treatment and control groups. Some children (N not clear) received parent + therapist-delivered massage (dual group) while others received only parent-delivered massage. While both groups improved on all measures from baseline, children in the dual group had greater improvements than those in the parent-only group (p values=ns).

Participants in a second RCT included five sibling pairs.<sup>25</sup> Children in the treatment group also had more severe challenging behaviors and sensory impairments at baseline. Children who received QST significantly improved on parent- and teacher-reported measures of autism symptoms, maladaptive behavior, communication skills, and sensory functioning from baseline, whereas children in the control group improved only on teacher measures of maladaptive behaviors. Children in the treatment group improved significantly more on teacher-rated measures of behavior and language (PDDBI, ABC) but not in maladaptive behavior compared with the control group. Children who received QST also improved more on all parent-rated measures of behavior and sensory processing (PDDBI, SSC) than did children in the control

group.<sup>25</sup> Parent-rated data on 19 treatment group participants still available for data collection showed that gains were maintained 5-months after the end of treatment.

A third RCT from the same group compared waitlisted controls to children who received five months of a QST Dual Program (parent + therapist-delivered massage).<sup>35</sup> Baseline variables and attrition rates were similar across groups. Children in both groups showed improvement over time in most domains, although control group improvements were generally of smaller magnitude than the treatment group. Post-hoc analyses revealed specific treatment effects on parent-reported but not clinician-rated measures: autism symptoms, receptive (but not expressive) language, sensory processing, and parent stress improved more in the treatment group compared with control ( $p$  values  $<0.01$ ). Group differences in social and living skills were not significant.

A final study from this group, with high risk of bias and unclear participant overlap with prior studies, retrospectively compared outcomes for children who either received QST for 5-months or were waitlisted controls.<sup>38</sup> Participants in the treatment groups received two varying levels of intervention (Home Program—parent only massage; Dual Program—parent + therapist-delivered massage), and outcome measures were reported by both parents and therapists using different reporting strategies (questionnaire versus observation). Participants in the treatment groups showed significant improvements in tactile defensiveness, self-regulation skills, and parent stress levels compared with controls.

**Massage + SI-based Treatment vs. SI-Based Treatment Alone.** One RCT with high risk of bias (included in the 2011 review) investigated 8 weeks of SI-based therapy compared with SI-based therapy plus traditional Thai massage.<sup>31</sup> Children in the intervention group (but not the control group) had significantly improved parent ratings of anxiety and conduct both relative to baseline and relative to controls ( $p \leq 0.03$ ). Children in both conditions had improved sleep as well as teacher ratings of conduct, attention, and activity level, and these ratings did not significantly differ across treatment groups. Participants in the treatment group had fewer symptoms of hyperactivity and sleep problems at baseline.<sup>31</sup>

## Additional Studies

### Key Points

- One study addressed tactile stimulation exercises and one addressed weighted blankets. These studies provided insufficient data to draw conclusions.

## Overview of the Literature

One RCT of a tactile input task conducted in the United States had moderate risk of bias and included 34 children between 4 and 14 years old.<sup>28</sup> Treatment duration was 24 to 48 hours. Another RCT with low risk of bias conducted in the United Kingdom included 54 children between 5 and 16 years of age.<sup>27</sup> Treatment duration was 2 weeks. Both studies had followup immediately post-treatment.

## Detailed Analysis

Other interventions with sensory-related components reported few significant differences between treatment groups (Appendix G).

**Tactile Input.** One RCT (moderate risk of bias) examined the impact of a tactile-based task on the ability of 34 children with autism to learn a novel task.<sup>28</sup> Participants included children who could not complete standardized cognitive or language evaluations. Children either participated in a tactual-kinesthetic experience (a “hands on” learning activity) or observed someone else performing the activity. Stimuli were presented across two sessions, 24-48 hours apart. Children in the hands-on participation group scored significantly better on ratings of perceived ease of implementing the learning task than children in the control condition ( $p$  values  $\leq 0.05$ ).

**Weighted Blankets.** One crossover RCT with moderate risk of bias examined the impact of a weighted blanket on sleep disturbance in children with severe problems with sleep onset or maintenance.<sup>27</sup> Children used a control or weighted blanket for 12-16 days and then switched. No significant differences emerged on any of the variables of interest related to sleep onset/quality, child behavior, or family functioning. Regardless of baseline factors such as sensory sensitivities, autism severity, and sleep problems, parents were more likely to rate their children as calmer and sleeping better when using the weighted blanket, despite a lack of physiological evidence to support this. Additionally, both children and parents reported preferring the weighted blanket. Investigators reported that one child developed a rash that may have been due to the blanket (resolved in 2 days).

## Key Question 2. Modifiers of Treatment Outcomes

Few studies were likely adequately powered to assess modifiers of effects, and few studies reported potential modifiers. While we sought characteristics of interventions, providers, parents, or children that may modify treatment effects, studies reported only child and family characteristics. We present findings as reported in each study below as potential indicators of characteristics that may affect outcomes.

**Child characteristics.** One study of an environmental enrichment-based approach reported that while more children in the treatment group compared with the control group shifted their diagnostic classification on the ADOS-2 from “autism” to “autism spectrum,” all children across both groups continued to meet the cut-offs for ASD, making it difficult to interpret the clinical significance of the findings. Baseline language and cognitive skills scores contributed to the majority of the variance when predicting which children in the treatment group would shift diagnostic cut-offs on the ADOS.<sup>23</sup> In another RCT comparing filtered and unfiltered music, participants whose parents rated them as having improved hearing sensitivity were more likely to show an increase in their sharing behaviors during a semi-structured play evaluation.<sup>19</sup> An RCT comparing home-based Qigong massage therapy to waitlisted controls found that children with fewer sensory symptoms and fewer difficulties with self-regulation benefited more from the treatment program.<sup>25</sup> In other studies of massage, treatment effects on child behaviors and language were not moderated by baseline ASD severity as measured on the CARS,<sup>35</sup> and interactions between self-regulation skills and tactile sensitivity or tactile sensitivity and parenting stress were not significant.<sup>38</sup> Finally, another study comparing weighted blankets and regular blankets reported that no baseline factors related to autism severity, sleep problems, or sensory sensitivities modified parents’ ratings of children’s sleep quality.<sup>27</sup>

**Family characteristics.** Although many studies collected parent report of child functioning, only one asked parents about their own functioning. One study of massage assessed parental stress levels and found that children in the treatment group had greater improvements in autism symptoms and overall behavior if their parents were less stressed at intake.<sup>24</sup> This would be an important avenue of future research, given that parental and family variables may influence how parents perceive children's functioning and subsequently complete questionnaires. Another study evaluated the quality of parent-child relationships as a potential modifier but did not find any treatment effects.<sup>26</sup>

### **Key Question 3. Time to Effect of Interventions**

No study examined the time to effect for sensory related interventions.

### **Key Question 4. Evidence that Effects Measured at the End of Treatment Predict Long-Term Functional Outcomes**

Only two studies conducted a followup after treatment ended. Followup occurred at two and five months in each study.<sup>25, 36</sup> Additionally, many of the outcome measures were based upon parent reports rather than using standardized interactive assessments. Therefore, little existing evidence at this time contributes to predicting long-term functional outcomes.

### **Key Question 5. Effectiveness Across Environments or Contexts**

We did not identify studies addressing this outcome directly. Many studies collected parent or teacher report of sensory sensitivities as a way of assessing functioning in multiple environments, but these assessments generally did not evaluate functioning within multiple specific contexts.

### **Key Question 6. Drivers of Treatment Outcomes**

We did not identify any studies addressing this KQ.

# Discussion

## State of the Literature

We identified a total of 21 studies (18 randomized controlled trials [RCTs], 1 nonrandomized trial, 2 retrospective cohort studies) addressing interventions targeting sensory challenges in children with Autism Spectrum Disorder (ASD). Three studies had low risk of bias,<sup>21, 23, 38</sup> ten had moderate,<sup>20, 22, 24-26, 28, 32-35</sup> and eight had high risk.<sup>19, 25, 29-31, 36, 37</sup> Most studies were small and used different outcome measures. Treatment length varied from two days to six months, with two studies reporting followup after the immediate intervention period.<sup>25, 36</sup> Protocols involved either parents or therapists engaging in one-on-one or group interactions.

Compared with our previous review, more studies were designed in ways that increased their strengths, including random assignment and tracking of attrition; stratification of assignment or matching based upon key baseline variables; blinded ratings; treatment fidelity protocols; tracking or controlling for concurrent interventions; and using standardized outcome measures or measures that incorporated parent-selected outcomes of importance such as goal attainment scaling. More information is now available and increases the strength of evidence in some cases over the findings of insufficient for all interventions addressed in our 2011 review. The lack of consistency in implementation, combined with generally small sample sizes and limited followup, however, limits our ability to draw strong conclusions regarding treatment efficacy.

Similarly, many studies continued to rely upon parent report of symptoms, although others used unbiased ratings of behavior (such as actigraphy or standardized measures) as well. It will be important for future work to compare sensory-based interventions not only to treatment as usual, but to other interventions that involve engaged and active time with an adult as did some studies in the current review.<sup>21, 23</sup> Additional research is needed that controls for environmental or social factors that could cloud our ability to draw conclusions regarding effects.

## Summary of Key Findings and Strength of the Evidence

### KQ1. Benefits and Harms of Interventions Targeting Sensory Challenges

#### Sensory Integration-Based Approaches

In four small, short-term studies, outcomes on sensory-related measures and motor skills measures were improved in children receiving a sensory-focused intervention compared with a usual care or a motor skills intervention, but effects on other outcomes were typically not significantly different between groups.<sup>20, 21, 30, 37</sup>

We found low strength of evidence for positive effects on measures of sensory challenges. The four studies that looked at this outcome were small and short-term (Table 3). We also found low strength of evidence for positive effects on motor skills, with three small studies showing significant improvements in treatment groups versus controls. We found low strength of evidence for no effect on measures of adaptive functioning. Evidence was insufficient to draw conclusions regarding other outcomes.

**Table 3. Strength of the evidence for sensory integration-based interventions**

Intervention/ Outcome	Study Limitations	Consistency	Directness	Precision	Reporting Bias	Finding  Strength of Evidence Grade
<b>Study Design</b>  <b>Risk of Bias and Number of Studies (N Total)</b>						
<b>Sensory integration-based approaches vs. Control approaches</b>						
Sensory challenges  RCT: 1 low, <sup>21</sup> 1 moderate, <sup>20</sup> 1 high <sup>30</sup> (N=99)  Retrospective cohort: 1 high <sup>37</sup> (N=20)	High	In-consistent	Direct	Imprecise	Undetected	<b>Low SOE for positive effects of sensory integration-based approaches on sensory challenges</b>  Significant improvements in sensory-related behaviors in treatment groups compared with control in 2 RCTs and one cohort study. No group differences in third RCT on parent-reported measure of sensory behaviors but significant improvement in treatment group in sensory-related goals; all studies were small and short-term
Motor skills  RCT: 1 low, <sup>21</sup> 1 moderate <sup>20</sup> (N=69)  Retrospective cohort: 1 high <sup>37</sup> (N=20)	High	Consistent	Direct	Imprecise	Undetected	<b>Low SOE for positive effects of sensory integration on motor skills</b>  Significant improvements in treatment groups vs. control in 3 small studies
Adaptive behavior  RCT: 1 low, <sup>21</sup> 1 moderate <sup>20</sup> (N=69)	High	Consistent	Direct	Imprecise	Undetected	<b>Low SOE for no effect of sensory integration therapies on adaptive behavior</b>  No significant group differences in adaptive behavior in 2 small RCTs with high study limitations. SOE is low given small size and limited followup

N=number; RCT=randomized controlled trial; SOE=strength of the evidence

## Environmental Enrichment-Based Approaches

Two small RCTs of environmental enrichment examined the same protocol involving parent-led sensory stimulation exercises and reported improvements in ASD symptoms, receptive language, and nonverbal cognitive skills after 6 months of treatment.<sup>22, 23</sup>

We considered strength of the evidence to be low for the effects of these approaches on cognitive skills and low for lack of effect on expressive language (Table 4). Strength of evidence was insufficient to assess effects of treatment on atypical sensory responses or receptive language as these outcomes were only addressed in one study.<sup>23</sup>

**Table 4. Strength of the evidence for environmental enrichment interventions**

Intervention/ Outcome	Study Design	Risk of Bias and Number of Studies (N Total)	Study Limitations	Consistency	Directness	Precision	Reporting Bias	Finding	Strength of Evidence Grade
<b>Environmental enrichment vs. Usual care</b>									
Cognitive skills			High	Consistent	Direct	Imprecise	Undetected	<b>Low SOE for positive effects of enrichment on IQ</b>	
RCT: 1 low, <sup>23</sup> 1 moderate <sup>22</sup> (N=78)								Significant improvements in IQ (Leiter) in children receiving enrichment compared with those receiving usual care in 2 small RCTs with short-term followup and high limitations given small sample size	
Expressive language			High	Consistent	Direct	Imprecise	Undetected	<b>Low SOE for lack of effect of enrichment on expressive language</b>	
RCT: 1 low, <sup>23</sup> 1 moderate <sup>22</sup> (N=78)								No group differences in expressive language in 2 small RCTs with short-term followup and high limitations given small sample size	

IQ=intelligence quotient; N=number; RCT=randomized controlled trial; SOE=strength of the evidence

## Auditory Integration-Based Approaches

Two small, short-term RCTs of auditory integration-based approaches assessing language outcomes reported no significant differences between groups in receptive language outcomes;<sup>33, 34</sup> one RCT (in a publication reporting 2 unique studies) reported significant parent-rated improvements in spontaneous speech.<sup>19</sup> Strength of evidence was low for a lack of effect of these approaches on language outcomes (Table 5).

**Table 5. Strength of the evidence for auditory integration-based interventions**

Intervention/ Outcome	Study Limitations	Consistency	Directness	Precision	Reporting Bias	Finding  Strength of Evidence Grade
Study Design  Risk of Bias and Number of Studies (N Total)						
<b>Auditory integration approaches vs. Control</b>						
Language  RCT: 2 moderate, <sup>33, 34</sup> 1 high <sup>19</sup> (N=91)	High	In-consistent	Direct	Imprecise	Undetected	<b>Low SOE for lack of effects of auditory integration approaches on language</b>  No group differences in outcomes in 2 small crossover RCTs with short-term followup; parent-rated improvements in spontaneous speech in treatment group vs. control in a third RCT

N=number; RCT=randomized controlled trial; SOE=strength of the evidence

## Music Therapy-Based Approaches

Three small studies addressing music therapy reported some significant effects on measures of behavior (social engagement, behavioral organization), and joint attention with music-based intervention compared with control interventions; however, studies used different protocols and addressed different outcomes. In some studies children also received interventions in addition to music-based approaches.<sup>26, 29, 36</sup>

We considered the strength of the evidence to be insufficient to assess effects of music therapy approaches on any outcome given that two of the three studies addressing these interventions had high risk of bias. Studies also had multiple comparators and addressed different outcomes.

## Massage-Based Approaches

As noted, five of six massage studies were from one group of investigators, with unclear participant overlap.<sup>24, 25, 31, 32, 35, 38</sup> Studies generally reported improvements related to sensory processing, autism symptoms, and parent stress in both treatment and control groups over the course of 5 months of either parent or parent + therapist-delivered intervention, with treatment groups improving significantly more than controls. The difficulty differentiating between these works limits the strength of evidence for their findings, although results seem promising regarding a sensory-focused intervention that can be delivered within the home environment with minimal risk of harms.

We considered the strength of the evidence to be low for positive effects on sensory challenges and ASD symptom severity with massage compared with control (Table 6). Strength of evidence was low for no effect on maladaptive behavior. The strength of the evidence is low as the extent of overlap among participants in unclear, and study limitations are high (unblinded ratings, diagnostic processes). We considered the strength of the evidence as insufficient to

assess effects of massage on language/communication outcomes given inconsistent findings and use of different outcome measures. Two studies reported greater improvements in the treatment group compared with controls on parent-rated measures of receptive but not expressive language<sup>35</sup> and parent- and clinician-rated composite measures<sup>25</sup> while another reported no significant group differences on a composite measure.<sup>24</sup> We also considered strength of the evidence insufficient to assess longer-term effects (>6 months) as only one study reported longer-term followup<sup>25</sup> and insufficient to assess effects on measures of daily living skills, also assessed in one study.<sup>35</sup> Strength of evidence was insufficient to assess effects of traditional massage plus sensory integration compared with sensory integration alone on any outcomes as only one study addressed this comparison (Table 9).<sup>31</sup>

**Table 6. Strength of the evidence for massage-based interventions**

<b>Intervention/ Outcome</b>	<b>Study Design</b>	<b>Risk of Bias and Number of Studies (N Total)</b>	<b>Study Limitations</b>	<b>Consistency</b>	<b>Directness</b>	<b>Precision</b>	<b>Reporting Bias</b>	<b>Finding</b>	<b>Strength of Evidence Grade</b>
<b>Massage vs. Waitlist Control</b>									
ASD symptom severity	RCT: 2 moderate, <sup>24, 35</sup> 1 high <sup>25</sup> (N=191)	High	Consistent	Direct	Imprecise	Undetected		<b>Low SOE for improvements in ASD symptom severity with massage vs. control in the short-term (&lt;6 months)</b>	Significant group differences in all 3 studies; SOE is low given unclear overlap in participants and high study limitations
Sensory challenges	RCT: 2 moderate, <sup>24, 35</sup> 1 high <sup>25</sup> (N=191) Retrospective cohort: 1 high <sup>38</sup> (N=129)	High	Consistent	Direct	Imprecise	Undetected		<b>Low SOE for positive effects on sensory challenges with massage vs. control in the short-term (&lt;6 months)</b>	Significant group differences in all 4 studies; SOE is low given unclear overlap in participants and high study limitations
Maladaptive behaviors	RCT: 2 moderate, <sup>24</sup> (N=423), 1 high <sup>25</sup> (N=46)	High	Consistent	Direct	Imprecise	Undetected		<b>Low SOE for no effect on maladaptive behaviors with massage vs. control in the short-term (&lt;6 months)</b>	No significant group differences in 2 studies; SOE is low given unclear overlap in participants and high study limitations

N=number; RCT=randomized controlled trial; SOE=strength of the evidence

## **Additional Interventions**

Additional interventions with sensory-related components (tactile stimulation exercises,<sup>28</sup> weighted blankets<sup>27</sup>) reported few significant differences between treatment groups. Strength of evidence was insufficient for all outcomes reported.

## **Other Key Questions**

Few studies were likely adequately powered to assess modifiers of effects, and few studies reported potential modifiers. We did not identify studies addressing the time to effect of interventions; evidence that effects measured at the end of treatment predict long-term functional outcomes; effectiveness of treatments across environments or contexts (e.g., clinic, home, school); and drivers of treatment outcomes.

## **Findings in Relation to What is Already Known**

We identified eleven recent (2010-present) systematic reviews addressing interventions targeting sensory challenges.<sup>6, 10, 40-48</sup> Three reviews were not specific to children with ASD but included studies of individuals with developmental disabilities.<sup>6, 10, 45</sup> As a number of these studies included children with ASD, we retained these reviews.

Reviews addressed comprehensive sensory integration approaches (typically clinic-based interventions using sensory-enhanced modalities to integrate sensory information and potentially ameliorate specific challenges or behaviors); “sensory-based” interventions targeting somatosensory or vestibular symptoms such as therapy balls, massage, and weighted vests; auditory integration; massage; and music therapy. Our findings generally align with these prior review of interventions. Reviews noted low to moderate support for sensory integration-based approaches and limited evidence for other approaches. Reviews consistently noted considerable heterogeneity, limited study quality/high risk of bias, limited followup, and lack of treatment fidelity. Reviews differentiating sensory integration approaches and more general “sensory-based” approaches reported better evidence from those studies that evaluated specific, typically manualized sensory integration modalities compared with sensory-based approaches. One review of auditory integration approaches reported no evidence of effectiveness. One review of music therapy reported promising findings related to improvements in social interaction and communication, and one addressing massage reported that limited evidence precluded conclusions. We provide more detailed summaries of these reviews below.

## **Reviews of Sensory Integration or Sensory-Based Approaches**

One review evaluating interventions for sensory processing disorders in children with ASD included 15 single subject design studies, 2 RCTs, one non-randomized trial, and one case report addressed sensory integration (n=5 studies) or sensory-based interventions (n=14 studies).<sup>3</sup> Six studies had scores of at a least 5 on the 10-point PEDro scale, and four reported higher level evidence on the Center for Evidence-Based Medicine scale. The review reported positive effects on children’s individualized goals (effect sizes ranging from 0.72 to 1.17) associated with sensory integration approaches but noted that durability and generalizability of effects was unclear. Evidence for sensory-based approaches was limited, and the review noted that studies suggest no support for weighted vests and little conclusive evidence for therapy balls or other multisensory inputs.

Another review addressing Ayres sensory integration (4 studies) and more general sensory-based interventions (18 studies) in children and adults with ASD reported similar results.<sup>47</sup> The studies addressing sensory integration included 133 children, and reviewers noted significant improvements in individualized goals, symptom severity, and sleep in studies with low risk of bias. Sensory-based studies (including single subject studies and systematic reviews) reported few positive effects and generally lacked measures of treatment fidelity.

In another review including 25 studies addressing sensory integration therapy for children with ASD, investigators noted positive effects in three studies, mixed findings in eight, and no benefits in 14.<sup>46</sup> The review considered “sensory integration” broadly, and did not differentiate manualized sensory integration from approaches such as weighted vests or brushing. Investigators considered 16 studies to report a “suggestive” (vs. conclusive) level of evidence based on methodologic limitations and concluded that the evidence base does not support use of sensory integration to treat children with ASD.

One review assessing sensory-based treatment in children with disabilities (including 236 of 856 children with ASD) included 15 comparative studies (including 13 RCTs) and 15 single subject design studies.<sup>6</sup> Investigators noted methodologic flaws in randomization (for comparative studies) and blinding and fidelity and reported inconsistent results for sensory integration approaches (no group differences in 3 studies, improvements in sensory behaviors and goal attainment in the sensory treatment arm in 4, and greater improvements with behavioral vs. sensory treatment in 2 other studies). Evidence for positive effects of weighted vests was lacking in six studies and mixed results for therapy balls in two studies. Overall the review concluded that inconclusive evidence supports the efficacy of sensory-based approaches and that such approaches are more likely to be ineffective than effective for the majority of children with developmental disabilities.

Another review of single subject studies of individuals with disabilities assessed “sensory integration” broadly and included 17 studies, six including children with ASD.<sup>45</sup> The investigators considered all studies to have high risk of bias and noted a lack of evidence for the efficacy of interventions: few studies reported positive effects, even those studies with better quality.

A third systematic review also focused on sensory approaches for children with behavioral problems (not necessarily ASD) and included 14 studies (11 included children with ASD, n=185).<sup>10</sup> Seven studies were single-subject design and four were RCTs. Investigators rated all studies as excellent or good quality on the PEDro scale and categorized interventions as tactile-, proprioceptive-, or vestibular-based. The review noted some positive effects associated with tactile approaches, particularly massage, on challenging behaviors including inattention. Proprioceptive studies—all single subject—evaluated weighted vests and reported mixed findings related to on-task, self-stimulatory, and stereotypic behaviors (improved findings in one study, mixed in two, and no effects in another). Vestibular-based approaches included therapy balls/cushions and horseback riding. Investigators noted limited positive effects on engagement in classroom activities associated with therapy balls. The review concluded that evidence for effectiveness of sensory-based interventions remains unclear.

Two other reviews focused broadly on interventions to improve social participation, play, adaptive behavior, education, and repetitive behavior included few studies addressing sensory challenges (n=3) and noted insufficient evidence for effects on social communication<sup>44</sup> and positive effects on self-care in one RCT.<sup>43</sup>

## Other Reviews

**Auditory integration.** One Cochrane review of auditory integration approaches for people with ASD included six RCTs with moderate to high risk of bias.<sup>41</sup> The review reported some improvements in language and challenging behaviors in small studies with disparate outcome measures and limited followup but concluded that interventions overall were ineffective for individuals with ASD.

**Music therapy.** Another Cochrane review of 10 studies (n=165 children with ASD), including theses, single subject studies, and comparative design studies, assessed music therapy and reported significant improvements in social interaction within and outside the therapy context and in communication and social reciprocity in the therapy context.<sup>42</sup> Nonverbal communication in non-therapy contexts was not significantly improved. The review concluded that music therapy may improve social interaction, communication, social reciprocity skills in children with ASD. These results differ from those reported in the current review, likely because we included only comparative studies of music therapy with at least 10 participants.

**Massage.** One review of massage therapy for children with ASD included six comparative studies and reported some positive effects of massage on symptom severity, communication, and sensory outcomes; however, investigators considered all trials to have high risk of bias.<sup>40</sup> The review concluded the evidence for massage as an ASD treatment is limited and methodologic limitations do not allow firm conclusions.

## Applicability

Children in studies meeting our review criteria are similar to the general population of children with ASD and associated sensory challenges in that they represent the heterogeneity of impairments and behavioral challenges associated with ASD. Some studies included children with limited functional speech<sup>30, 33</sup> and/or intellectual disability,<sup>34</sup> others included children with higher cognitive abilities and milder ASD symptom profiles.<sup>21, 37</sup> Differences in severity of expression of ASD or in comorbid conditions may limit applicability of findings to children with levels of symptom expression.

Interventions typically used differing approaches incorporating sensory-focused strategies; thus, findings reported here may not be replicated with interventions using different combinations of strategies or targeting different aspects of sensory functioning. Many of the interventions required expertise in specific sensory-related strategies that likely limit their generalizability to community settings; some lacked manualization, and as a consequence likely have limited replicability/community extension. Others were specifically designed to be conducted by parents, under the supervision of a trainer.

In terms of comparison or control interventions, virtually all “control” participants in the included studies were receiving some level of treatment, aside from participants in one RCT.<sup>36</sup> Some studies attempted to document additional interventions through parent report to demonstrate that baseline rates were comparable across groups, whereas others required that no new interventions be added during the study duration. Outcomes may differ among children receiving different concomitant therapies.

Outcomes also varied across studies, but most studies incorporated commonly used measures of autism symptom severity, behavior, language, and sensory difficulties. Many of these were

based upon parent report, although an increasing number used standardized interactive or observational measurement strategies. Some studies also incorporated outcomes of key importance to parents/caregivers that simultaneously attempted to balance assessing comparability between treatment groups with the heterogeneity inherent in ASD.<sup>20,21</sup> Use of validated, standardized measures improves potential generalizability of findings by helping to establish a shared understanding of progress and to compare outcomes across different intervention modalities and studies. Standardized measures supplemented with more individualized assessment strategies can also help to improve sensitivity by measuring smaller effects and incorporating elements of children's individual family or treatment context.

Treatment duration ranged from a week to 10 months, and most studies reported outcomes immediately following the end of treatment, making durability of effects difficult to assess. Few studies attempted to assess characteristics of the child, family, provider, or intervention approach that may affect outcomes, or whether outcomes extended to other settings and environments. However, the use of control groups, treatment fidelity checks, and replicable and manualized intervention protocols establishes a promising baseline for future investigations.

Given the heterogeneity of these studies, and the heterogeneity of children with ASD, the extent of generalizability to the overall population of children with ASD and sensory challenges is limited and difficult to assess.

## **Implications for Clinical and Policy Decisionmaking**

This review provides limited evidence for decisionmaking about interventions targeting sensory challenges. The small body of comparative literature provides some evidence to support sensory integration-based and touch/massage-based interventions for some children; both interventions positively affected sensory challenges and motor skills. Studies were typically short term and included few children, however, so our confidence in these effects is low.

Decisional dilemmas remain regarding characteristics of the child, family, or intervention that may modify effectiveness or predict which children may be most likely to benefit from a given approach. Similarly, the literature base is currently insufficient to inform our understanding of the time to effect of interventions, longer-term effectiveness of interventions, generalizability of effects outside the treatment context, and components that may drive effectiveness. Though not explicitly assessed in the studies reported here, harms associated with these approaches are likely minimal and caregivers and clinicians must balance the need to ameliorate sensory challenges with the costs of time, effort, and costs and demands of other interventions that a child may receive. As such, caregivers and referring providers should assess the possible benefits of specific sensory-focused intervention modalities based upon the individual needs of the child, broader family goals and capacities, and interventions of more established effectiveness. In this capacity, some practice groups have recommended clear communication regarding the limits of intervention.<sup>49, 50</sup>

## **Limitations of the Comparative Effectiveness Review Process**

We included studies published in English only and did not include unpublished data. We scanned a random sample of 150 non-English abstracts retrieved by our MEDLINE search. Most studies appeared to be case series, narrative reviews, basic science studies, or studies assessing etiology. Only two studies appeared to meet inclusion criteria; thus, given the high percentage of

ineligible items in this scan (99%), we concluded that excluding non-English studies would not introduce significant bias into the review.

We also included only comparative studies of interventions with a sensory-specific focus and including at least 10 children with ASD. This undoubtedly means that most single-subject design studies are not included in this review. Single-subject designs can be helpful in assessing response to treatment in very short timeframes and under very tightly controlled circumstances, but they typically do not provide information on longer-term or functional outcomes. Such studies are useful in demonstrating effects, yielding initial evidence that an intervention merits further study or in identifying whether a particular approach to treatment is likely to be helpful for a specific child. Our goal was to identify and review the best evidence for assessing the effectiveness of interventions targeting sensory challenges in children with ASD, with an eye toward utility in the larger population of children with ASD. By definition, “populations” in single-subject design studies are likely to be idiosyncratic and therefore unlikely to provide information that is generalizable. We also included summaries of other recent reviews to attempt to mitigate any loss of information associated with this criterion.

This review was also focused specifically on children with ASD and specifically on interventions targeting sensory challenges. Sensory approaches may be used with individuals with other impairments, and findings may be generalizable to children with ASD; however, including studies in children with other conditions was beyond the scope of the current review. We also recognize that interventions with a primarily educational or behavioral focus may also address sensory-related outcomes, but inclusion of any intervention approach reporting a sensory-related outcome was outside the scope of the current review. Similarly, we focused on child-related outcomes and did not address measures of parent stress, despite the key importance of the outcome and family context. Finally, we used a non-validated tool to assess risk of bias, though we note that the tool evaluates similar constructs to those assessed in tools such as that used by the Cochrane Collaboration, with the addition of ASD-specific domains.

## **Limitations of the Evidence Base**

As noted, studies in the review had small sample sizes (median 34 total) and typically limited duration of intervention and followup after intervention. Populations across studies were heterogeneous in terms of sensory challenges, ASD severity, age, and intellectual and adaptive functioning. Interventions, even within our broader categories, used differing sensory-specific approaches in differing combinations of components, settings, and duration. Longer-term outcomes are limited as is our ability to determine effects of intervention on the underlying sensory challenges themselves. Potential harms of interventions were addressed in only one study, and few studies assessed potential factors that may modify effectiveness or drive effects of interventions. Studies often used multicomponent strategies, and teasing apart effects of specific components is not currently possible.

Despite these limitations, investigators have made significant improvements in incorporating commonly used measures of symptom severity, behavior, language, and sensory difficulties to facilitate comparisons across studies. Parent-reported outcomes are necessary in this population of children, many of whom may not be able to complete aspects of assessments; however, studies are increasingly incorporating standardized interactive or observational measurement strategies. As noted above, the increasing use of treatment fidelity measures and replicable intervention protocols establishes a promising baseline for future investigations. Investigators in the area are also well-aware of the challenges of conducting research using a disparate and

variously defined set of approaches in a highly heterogeneous population and have made strides in incorporating outcome measures that attempt to balance heterogeneity and comparative effectiveness and measures of intervention fidelity.<sup>15</sup>

## **Research Gaps and Areas for Future Research**

Improving research in this area should also include considerations of power and sample size. Sample size was frequently insufficient to allow firm conclusions. In addition, researchers should continue to provide adequate detail as they describe their interventions to allow for replicable research. Ideally, investigators publish the treatment manuals they develop, which are then referenced in later research; despite gains in this area, many studies made general references to their use of an underlying approach without specifying the ways in which they used or modified the technique. Lack of detail about the intervention makes it difficult to assess the applicability of individual studies, to synthesize groups of studies, or to replicate studies.

Duration of treatment and followup were generally short ( $\leq 10$  months). Few studies provided data on long-term outcomes after cessation of treatment. Future studies should extend the followup period and assess the degree to which outcomes are durable. In addition, few studies adequately accounted for concomitant interventions that might confound observed effectiveness. Accounting for concomitant interventions should be standardized in future research.

As noted, more studies used a common set of outcome measures, but the extent to which these measures assess changes in potential underlying sensory-related impairments is not clear, and understanding whether intervention can alter potential impairments in sensory processing (vs. altering behavioral responses in the short term) is a critical need. Another critical area for further research is identifying which children are likely to benefit from particular interventions. To date, studies have provided limited characterization of the subpopulation of children who experience positive response to intervention and limited characterization of the extent or type of sensory challenges children experience at baseline. Interventions targeting sensory challenges by their nature often employ multiple components, and data on whether specific functional components of the interventions drive effectiveness are currently unavailable. Component analyses in this field would be productive for refining intervention approaches and for assessing applicability and generalizability of the results.

In line with this need, we recommend future consideration of the ways in which the cultural context of the child and family may affect the applicability or effectiveness of specific interventions. The setting of interventions may also influence effects, and understanding the role of context broadly in contributing to effects is an important need. As noted, understanding the extent to which findings from studies of these interventions in children with other conditions are applicable to children with ASD may help to bolster the evidence base.

## **Conclusions**

Some interventions targeting sensory challenges may effect modest improvements primarily in sensory- and ASD symptom severity-related outcomes; however, the evidence base for any category of intervention is small, and durability of effects beyond the immediate intervention period is unclear. Sensory integration-based approaches improved outcomes related to sensory challenges and motor skills, and studies of massage reported improvements in sensory responses and ASD symptoms. Environmental enrichment was also associated with improvements in cognitive skills in the short-term. Some positive effects were associated with other approaches

studied (auditory integration, music therapy, weighted blankets) but findings in these small studies were not consistent. Data on longer-term results are lacking, as are data on characteristics that modify outcomes, generalizability of findings, and components of interventions that may drive effects. In sum, while some therapies hold promise and warrant further study, substantial needs exist for continuing improvements in methodologic rigor in the field.

# References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. V ed. Washington DC: American Psychiatric Association 2013.
2. Baranek GT. Efficacy of sensory and motor interventions for children with autism. *J Autism Dev Disord*. 2002 Oct;32(5):397-422. PMID: 12463517.
3. Case-Smith J, Weaver LL, Fristad MA. A systematic review of sensory processing interventions for children with autism spectrum disorders. *Autism: The International Journal of Research & Practice*. 2015;19(2):133-48 16p. doi: 10.1177/1362361313517762. PMID: 103750467. Language: English. Entry Date: 20150130. Revision Date: 20150710. Publication Type: Journal Article.
4. State of the Science: A Roadmap for Research in Sensory Integration. *American Journal of Occupational Therapy*. 2015;69(6):1-7 p. doi: 10.5014/ajot.2015.019539. PMID: 110629144. Language: English. Entry Date: 20151109. Revision Date: 20151112. Publication Type: Article. Journal Subset: Allied Health.
5. Schaaf RC. Interventions that address sensory dysfunction for individuals with autism spectrum disorders: Preliminary evidence for the superiority of sensory integration compared to other sensory approaches. Evidence-based practices and treatments for children with autism.: Springer Science + Business Media, New York, NY; 2011:245-73.
6. Barton EE, Reichow B, Schnitz A, et al. A systematic review of sensory-based treatments for children with disabilities. *Res Dev Disabil*. 2015 Feb;37:64-80. doi: 10.1016/j.ridd.2014.11.006. PMID: 25460221.
7. Ospina MB, Krebs Seida J, Clark B, et al. Behavioural and developmental interventions for autism spectrum disorder: a clinical systematic review. *PLoS One*. 2008;3(11):e3755. doi: 10.1371/journal.pone.0003755 [doi]. PMID: 19015734.
8. Seida JK, Ospina MB, Karkhaneh M, et al. Systematic reviews of psychosocial interventions for autism: an umbrella review. *Dev Med Child Neurol*. 2009 Feb;51(2):95-104. doi: DMCN3211 [pii]10.1111/j.1469-8749.2008.03211.x [doi]. PMID: 19191842.
9. Parr J. Autism. *BMJ Clin Evid*. 2010;2010 PMID: 21729335.
10. Wan Yunus F, Liu KP, Bissett M, et al. Sensory-Based Intervention for Children with Behavioral Problems: A Systematic Review. *J Autism Dev Disord*. 2015 Nov;45(11):3565-79. doi: 10.1007/s10803-015-2503-9. PMID: 26092640.
11. Myers SM, Johnson CP. Management of children with autism spectrum disorders. *Pediatrics*. 2007 Nov;120(5):1162-82. doi: peds.2007-2362 [pii]10.1542/peds.2007-2362 [doi]. PMID: 17967921.
12. National Autism Center. National Standards Report National Autism Center. Randolph, MA: NAC, 2009. <http://www.nationalautismcenter.org/pdf/NAC%20Standards%20Report.pdf>
13. Committee on Educational Interventions for Children with Autism NRC. Educating Children with Autism Washington DC: National Academies Press; 2001. [http://www.nap.edu/openbook.php?record\\_id=10017&page=R2](http://www.nap.edu/openbook.php?record_id=10017&page=R2)

14. Martin A, Scahill L, Anderson GM, et al. Weight and leptin changes among risperidone-treated youths with autism: 6-month prospective data. *Am J Psychiatry*. 2004 Jun;161(6):1125-7. PMID: 15169706.
15. Schaaf RC, Schoen SA, May-Benson TA, et al. State of the Science: A Roadmap for Research in Sensory Integration. *Am J Occup Ther*. 2015 Nov-Dec;69(6):6906360010p1-7. doi: 10.5014/ajot.2015.019539. PMID: 26565107.
16. Methods Guide for Effectiveness and Comparative Effectiveness Reviews. AHRQ Publication No. 10(14)-EHC063-EF. Rockville, MD: Agency for Healthcare Research and Quality. January 2014. Chapters available at: [www.effectivehealthcare.ahrq.gov](http://www.effectivehealthcare.ahrq.gov)
17. Warren Z, Veenstra-VanderWeele J, Stone W, Bruzek JL, Nahmias AS, Foss-Feig JH, Jerome RN, Krishnaswami S, Sathe NA, Glasser AM, Surawicz T, McPheeters ML. Therapies for Children With Autism Spectrum Disorders. Comparative Effectiveness Review No. 26. (Prepared by the Vanderbilt Evidence-based Practice Center under Contract No. 290-2007-10065-I.) AHRQ Publication No. 11-EHC029-EF. Rockville, MD: Agency for Healthcare Research and Quality. April 2011. Available at: [www.effectivehealthcare.ahrq.gov/reports/final.cfm](http://www.effectivehealthcare.ahrq.gov/reports/final.cfm).
18. Weitlauf AS, McPheeters ML, Peters B, Sathe N, Travis R, Aiello R, Williamson E, Veenstra-VanderWeele J, Krishnaswami S, Jerome R, Warren Z. Therapies for Children With Autism Spectrum Disorder: Behavioral Interventions Update. Comparative Effectiveness Review No. 137. (Prepared by the Vanderbilt Evidence-based Practice Center under Contract No. 290-2012-00009-I.) AHRQ Publication No. 14-EHC036-EF. Rockville, MD: Agency for Healthcare Research and Quality; August 2014. [www.effectivehealthcare.ahrq.gov/reports/final.cfm](http://www.effectivehealthcare.ahrq.gov/reports/final.cfm).
19. Porges SW, Bazhenova OV, Bal E, et al. Reducing auditory hypersensitivities in autistic spectrum disorder: preliminary findings evaluating the listening project protocol. *Front Pediatr*. 2014;2:80. doi: 10.3389/fped.2014.00080. PMID: 25136545.
20. Schaaf RC, Benevides T, Mailloux Z, et al. An intervention for sensory difficulties in children with autism: a randomized trial. *J Autism Dev Disord*. 2014 Jul;44(7):1493-506. doi: 10.1007/s10803-013-1983-8. PMID: 24214165.
21. Pfeiffer BA, Koenig K, Kinnealey M, et al. Effectiveness of sensory integration interventions in children with autism spectrum disorders: a pilot study. *Am J Occup Ther*. 2011 Jan-Feb;65(1):76-85. PMID: 21309374.
22. Woo CC, Leon M. Environmental enrichment as an effective treatment for autism: a randomized controlled trial. *Behav Neurosci*. 2013 Aug;127(4):487-97. doi: 10.1037/a0033010. PMID: 23688137.
23. Woo CC, Donnelly JH, Steinberg-Epstein R, et al. Environmental enrichment as a therapy for autism: A clinical trial replication and extension. *Behav Neurosci*. 2015 Aug;129(4):412-22. doi: 10.1037/bne0000068. PMID: 26052790.
24. Silva LM, Schalock M, Gabrielsen K. Early intervention for autism with a parent-delivered Qigong massage program: a randomized controlled trial. *Am J Occup Ther*. 2011 Sep-Oct;65(5):550-9. PMID: 22026323.
25. Silva LM, Schalock M, Ayres R, et al. Qigong massage treatment for sensory and self-regulation problems in young children with autism: a randomized controlled trial. *Am J Occup Ther*. 2009 Jul-Aug;63(4):423-32. PMID: 19708471.

26. Thompson GA, McFerran KS, Gold C. Family-centred music therapy to promote social engagement in young children with severe autism spectrum disorder: a randomized controlled study. *Child Care Health Dev.* 2014 Nov;40(6):840-52. doi: 10.1111/cch.12121. PMID: 24261547.
27. Gringras P, Green D, Wright B, et al. Weighted blankets and sleep in autistic children--a randomized controlled trial. *Pediatrics.* 2014 Aug;134(2):298-306. doi: 10.1542/peds.2013-4285. PMID: 25022743.
28. Latham SO, Stockman IJ. Effect of augmented sensorimotor input on learning verbal and nonverbal tasks among children with autism spectrum disorders. *J Autism Dev Disord.* 2014 Jun;44(6):1288-302. doi: 10.1007/s10803-013-1990-9. PMID: 24234676.
29. Kim J, Wigram T, Gold C. The effects of improvisational music therapy on joint attention behaviors in autistic children: a randomized controlled study. *J Autism Dev Disord.* 2008 Oct;38(9):1758-66. doi: 10.1007/s10803-008-0566-6 [doi]. PMID: 18592368.
30. Fazlioglu Y, Baran G. A sensory integration therapy program on sensory problems for children with autism. *Percept Mot Skills.* 2008 Apr;106(2):415-22. PMID: 18556898.
31. Piravej K, Tangtrongchitr P, Chandarasiri P, et al. Effects of Thai traditional massage on autistic children's behavior. *J Altern Complement Med.* 2009 Dec;15(12):1355-61. doi: 10.1089/acm.2009.0258 [doi]. PMID: 20001837.
32. Silva LM, Cignolini A, Warren R, et al. Improvement in sensory impairment and social interaction in young children with autism following treatment with an original Qigong massage methodology. *Am J Chin Med.* 2007;35(3):393-406. doi: S0192415X07004916 [pii]. PMID: 17597498.
33. Corbett BA, Shickman K, Ferrer E. Brief report: the effects of Tomatis sound therapy on language in children with autism. *J Autism Dev Disord.* 2008 Mar;38(3):562-6. doi: 10.1007/s10803-007-0413-1 [doi]. PMID: 17610057.
34. Mudford OC, Cross BA, Breen S, et al. Auditory integration training for children with autism: no behavioral benefits detected. *Am J Ment Retard.* 2000 Mar;105(2):118-29. PMID: 10755175.
35. Silva LM, Schalock M, Gabrielsen KR, et al. Early Intervention with a Parent-Delivered Massage Protocol Directed at Tactile Abnormalities Decreases Severity of Autism and Improves Child-to-Parent Interactions: A Replication Study. *Autism Res Treat.* 2015;2015:904585. doi: 10.1155/2015/904585. PMID: 25878901.
36. Ghasemtabar SN, Hosseini M, Fayyaz I, et al. Music therapy: An effective approach in improving social skills of children with autism. *Adv Biomed Res.* 2015;4:157. doi: 10.4103/2277-9175.161584. PMID: 26380242.
37. Iwanaga R, Honda S, Nakane H, et al. Pilot study: efficacy of sensory integration therapy for Japanese children with high-functioning autism spectrum disorder. *Occup Ther Int.* 2014 Mar;21(1):4-11. doi: 10.1002/oti.1357. PMID: 23893373.
38. Silva L, Schalock M. Treatment of tactile impairment in young children with autism: results with qigong massage. *Int J Ther Massage Bodywork.* 2013;6(4):12-20. PMID: 24298297.

39. Patil RR. MMR vaccination and autism: learnings and implications. *Hum Vaccin*. 2011 Feb;7(2):281-2. PMID: 21343697.
40. Lee MS, Kim JI, Ernst E. Massage therapy for children with autism spectrum disorders: a systematic review. *J Clin Psychiatry*. 2011 Mar;72(3):406-11. doi: 10.4088/JCP.09r05848whi. PMID: 21208598.
41. Sinha Y, Silove N, Hayen A, et al. Auditory integration training and other sound therapies for autism spectrum disorders (ASD). *Cochrane Database Syst Rev*. 2011(12):CD003681. doi: 10.1002/14651858.CD003681.pub3. PMID: 22161380.
42. Geretsegger M, Holck U, Gold C. Randomised controlled trial of improvisational music therapy's effectiveness for children with autism spectrum disorders (TIME-A): study protocol. *BMC Pediatr*. 2012;12:2. doi: 10.1186/1471-2431-12-2. PMID: 22221670.
43. Weaver LL. Effectiveness of Work, Activities of Daily Living, Education, and Sleep Interventions for People With Autism Spectrum Disorder: A Systematic Review. *Am J Occup Ther*. 2015 Sep-Oct;69(5):6905180020p1-p11. doi: 10.5014/ajot.2015.017962. PMID: 26356654.
44. Tanner K, Hand BN, O'Toole G, et al. Effectiveness of Interventions to Improve Social Participation, Play, Leisure, and Restricted and Repetitive Behaviors in People With Autism Spectrum Disorder: A Systematic Review. *Am J Occup Ther*. 2015 Sep-Oct;69(5):6905180010p1-p12. doi: 10.5014/ajot.2015.017806. PMID: 26356653.
45. Leong HM, Carter M, Stephenson J. Systematic review of sensory integration therapy for individuals with disabilities: Single case design studies. *Res Dev Disabil*. 2015 Dec;47:334-51. doi: 10.1016/j.ridd.2015.09.022. PMID: 26476485.
46. Lang R, O'Reilly M, Healy O, et al. Sensory integration therapy for autism spectrum disorders: A systematic review. *Research in Autism Spectrum Disorders*. 2012 2015-02-24;6(3):1004-18. doi: <http://dx.doi.org/10.1016/j.rasd.2012.01.006>. PMID: 1022253260; 2012-13564-007.
47. Watling R, Hauer S. Effectiveness of Ayres Sensory Integration((R)) and Sensory-Based Interventions for People With Autism Spectrum Disorder: A Systematic Review. *Am J Occup Ther*. 2015 Sep-Oct;69(5):6905180030p1-p12. doi: 10.5014/ajot.2015.018051. PMID: 26356655.
48. Mistic B, Doesburg SM, Fatima Z, et al. Coordinated Information Generation and Mental Flexibility: Large-Scale Network Disruption in Children with Autism. *Cereb Cortex*. 2015 Sep;25(9):2815-27. doi: 10.1093/cercor/bhu082. PMID: 24770713.
49. Section On C, Integrative M, Council on Children with D, et al. Sensory integration therapies for children with developmental and behavioral disorders. *Pediatrics*. 2012 Jun;129(6):1186-9. doi: 10.1542/peds.2012-0876. PMID: 22641765.
50. Watling RK, KP; Davies PL; Schaaf R. Occupational Therapy Practice Guidelines for Children and Adolescents With Challenges in Sensory Processing and Sensory Integration AOTA. 2011.

## Acronyms and Abbreviations

ABC	Autism Behavior Checklist
ADOS	Autism Diagnostic Observation Schedule
AHRQ	Agency for Healthcare Research And Quality
ASD	Autism Spectrum Disorder
CARS	Childhood Autism Rating Scale
CER	Comparative Effectiveness Review
DSM	<i>Diagnostic and Statistical Manual</i>
G	Group
GAS	Goal Attainment Setting
IQ	Intelligence Quotient
KQ	Key Question
n	Number
NR	Not Reported
NS	Not Significant
OT/SI	Occupational Therapy with Sensory Integration
PDDBI	Pervasive Development Disorder Behavior Inventory
PDD-NOS	Pervasive Developmental Disorder – Not Otherwise Specified
PICOTS	Population, Intervention, Comparator, Outcome, Timing, Setting
PLS	Preschool Language Scale
QST	Qigong Sensory Training
RCT	Randomized Controlled Trial
SI	Sensory Integration
SOE	Strength of Evidence
SSC	Sense And Self-Regulation Checklist
TEP	Technical Expert Panel
VABS	Vineland Adaptive Behavior Scales