



## Effective Health Care Treatments for Hip and Hand Osteoarthritis

### Results of Topic Selection Process & Next Steps

The nominators, the American College of Rheumatology (ACR) and the American College of Physicians (ACP), are interested in an Agency for Healthcare Research and Quality (AHRQ) evidence review to update their 2012 guidelines on pharmacologic and non-pharmacological treatments for hip and hand osteoarthritis. Due to limited program resources, the Effective Health Care (EHC) program will not develop a review at this time. No further activity on this topic will be undertaken by the EHC Program.

### Topic Brief

**Topic Name:** Treatments for Hip and Hand Osteoarthritis

**Topic #:** 0712

**Nomination Date:** October 28, 2016

**Topic Brief Date:** March 2017

#### Authors

Kara Winchell  
Rose Revelo

**Conflict of Interest:** None of the investigators have any affiliations or financial involvement that conflicts with the material presented in this report.

#### Summary of Key Findings

- Appropriateness and importance: The nomination is both appropriate and important.
- Duplication: An AHRQ systematic review on the treatments for hip and hand osteoarthritis (OA) would not be duplicative. We identified 12 complete or in-process reviews/meta-analyses—one review examining physical and occupational therapy for hand OA, and eleven complete or in-process evidence reviews/meta-analyses covering specific interventions for hip OA. The interventions examined in these eleven hip OA evidence reviews are platelet-rich plasma, opioid therapy, exercise and physical therapy, manual therapy, and high- and low-velocity resistance training. Most of the interventions of interest to the nominators are not included in these reviews.
- Impact: The nomination has moderate impact potential. The ACR published guidelines for [nonpharmacological and pharmacologic therapies for hand, hip, and knee osteoarthritis](#) in 2012, using GRADE and a panel consensus. An updated review could inform evidence gaps for recommendations that were not considered strong, particularly related to therapies for hand OA, hip OA. By the time an AHRQ evidence review on this topic will be complete, the ACR guidelines will be more than five years old and in need of updating.

- Feasibility: An AHRQ evidence review is feasible at this time.
  - *Size/scope of review*: We identified 93 studies potentially relevant to the key questions in the nomination.
  - *Clinicaltrials.gov*: We identified 23 ongoing or recently completed trials on ClinicalTrials.gov.
- Value: The nomination has a high value potential, given that the ACR and the ACP will jointly use a new AHRQ systematic review to update their 2012 guidelines.

## Table of Contents

Introduction.....	1
Methods.....	4
Appropriateness and Importance.....	4
Desirability of New Review/Duplication .....	4
Impact of a New Evidence Review .....	4
Feasibility of a New Evidence Review .....	4
Compilation of Findings .....	4
Results .....	5
Appropriateness and Importance.....	5
Desirability of New Review/Duplication .....	5
Impact of a New Evidence Review .....	5
Feasibility of a New Evidence Review .....	5
Value.....	9
Summary of Findings .....	9
References .....	10
Appendices.....	19
Appendix A. Selection Criteria Summary.....	A-1
Appendix B. Search Strategy & Results (Feasibility).....	B-1

## Introduction

Osteoarthritis (OA), the most common form of arthritis, is a degenerative joint disease that affects approximately 30.8 million Americans.<sup>1</sup> Symptoms of OA generally get worse with age and increased “wear and tear.” Treatment for OA varies from exercise/weight loss, to pain medication, to joint replacement surgery.

Topic nomination #0712 was received on October 28, 2016. This nomination was submitted as a joint nomination by the American College of Rheumatology (ACR) and the American College of Physicians (ACP). While the nomination also mentions shoulder OA, we confined the scope to hip and hand OA because these were the focus of the proposed key questions. The key questions for this nomination are:

Key Question 1. What is the clinical effectiveness of therapies in patients with primary or secondary OA of the hip?

- a. Pharmacologic
- b. Non-pharmacologic
- c. By duration and intensity
- d. Subgroups

Key Question 2. What harms are associated with interventions in patients with primary or secondary OA of the hip?

- a. Pharmacologic
- b. Non-pharmacologic
- c. By duration and intensity
- d. Subgroups

Key Question 3. What is the clinical effectiveness of therapies in patients with primary or secondary OA of the hand?

- a. Pharmacologic
- b. Non-pharmacologic
- c. By duration and intensity
- d. Subgroups

Key Question 4. What harms are associated with interventions in patients with primary or secondary OA of the hand?

- a. Pharmacologic
- b. Non-pharmacologic
- c. By duration and intensity
- d. Subgroups

To define the inclusion criteria for the key questions we specify the population, interventions, comparators, and outcomes of interest. See Table 1.

**Table 1. Key Question and PICOs**

<b>Key Question</b>	1. What is the clinical effectiveness of therapies in patients with primary or secondary OA of the hip? a. Pharmacologic b. Non-pharmacologic c. By duration and intensity d. Subgroups	2. What harms are associated with interventions in patients with primary or secondary OA of the hip? a. Pharmacologic b. Non-pharmacologic c. By duration and intensity d. Subgroups	3. What is the clinical effectiveness of therapies in patients with primary or secondary OA of the hand? a. Pharmacologic b. Non-pharmacologic c. By duration and intensity d. Subgroups	4. What harms are associated with interventions in patients with primary or secondary OA of the hand? a. Pharmacologic b. Non-pharmacologic c. By duration and intensity d. Subgroups
<b>Population</b>	a. Adults with OA of the hip b. Adults with OA of the hip c. Adults with OA of the hip d. Adults with OA of the hip, by the following characteristics: sex, disease subtype (lateral, patellofemoral), severity (stage/baseline pain and functional status), weight status (body mass index), baseline fitness (activity level), comorbidities, prior or concurrent treatments (including self-initiated therapies)	a. Adults with OA of the hip b. Adults with OA of the hip c. Adults with OA of the hip d. Adults with OA of the hip, by the following characteristics: sex, disease subtype (lateral, patellofemoral), severity (stage/baseline pain and functional status), weight status (body mass index), baseline fitness (activity level), comorbidities, prior or concurrent treatments (including self-initiated therapies)	a. Adults with OA of the hand b. Adults with OA of the hand c. Adults with OA of the hand d. Adults with OA of the hand, by the following characteristics: sex, disease subtype, severity (stage/baseline pain and functional status), weight status (body mass index), baseline fitness (activity level), comorbidities, prior or concurrent treatments (including self-initiated therapies)	a. Adults with OA of the hand b. Adults with OA of the hand c. Adults with OA of the hand d. Adults with OA of the hand, by the following characteristics: sex, disease subtype, severity (stage/baseline pain and functional status), weight status (body mass index), baseline fitness (activity level), comorbidities, prior or concurrent treatments (including self-initiated therapies)
<b>Interventions</b>	a. Pharmacologic: duloxetine, intraarticular corticosteroids, intraarticular hyaluronic acid, intraarticular platelet-rich plasma/ultrasound-guided intraarticular injections of platelet-rich plasma, intramuscular corticosteroid injection, glucosamine and chondroitin sulfate, NSAIDs, COX-2-selective inhibitors, tramadol, acetaminophen, tapentadol, opioid therapy, tanezumab, alendronate, avocado-soybean unsaponifiable-expanscience (ASU-E), low molecular weight hydrolyzed chicken sternal	a. Pharmacologic: duloxetine, intraarticular corticosteroids, intraarticular hyaluronic acid, intraarticular platelet-rich plasma/ultrasound-guided intraarticular injections of platelet-rich plasma, intramuscular corticosteroid injection, glucosamine and chondroitin sulfate, NSAIDs, COX-2-selective inhibitors, tramadol, acetaminophen, tapentadol, opioid therapy, tanezumab, alendronate, avocado-soybean unsaponifiable-expanscience (ASU-E), low molecular weight hydrolyzed chicken sternal	a. Pharmacologic: duloxetine, topical therapy, DMARDs (Plaquenil, etc.), intraarticular corticosteroids, intraarticular hyaluronic acid, intraarticular platelet-rich plasma/ultrasound-guided intraarticular injections of platelet-rich plasma, doxycycline treatment for erosive osteoarthritis of the hand, glucosamine and chondroitin sulfate, NSAIDs, topical NSAIDs, COX-2-selective inhibitors, diacerein, arnica (topical), hydroxycoumarin content of Sphaeralcea angustifolia, tramadol, acetaminophen,	a. Pharmacologic: duloxetine, topical therapy, DMARDs (Plaquenil, etc.), intraarticular corticosteroids, intraarticular hyaluronic acid, intraarticular platelet-rich plasma/ultrasound-guided intraarticular injections of platelet-rich plasma, doxycycline treatment for erosive osteoarthritis of the hand, glucosamine and chondroitin sulfate, NSAIDs, topical NSAIDs, COX-2-selective inhibitors, diacerein, arnica (topical), hydroxycoumarin content of Sphaeralcea angustifolia, tramadol, acetaminophen,

	<p>cartilage extract, BioCell Collagen, UP446</p> <p>b. Non-pharmacological: thermal baths with a sulfide mineral water, exercise therapy, physical therapy, neuromuscular exercise in patients with severe OA of the hip, gluteus medius muscle activity, high-velocity resistance training/low-velocity resistance training, tailored activity pacing, manual therapy, Basic Body Awareness Therapy (BBAT), cognitive behavioral therapy for chronic pain, internet-based pain coping skills training, self-management program, pulsed radiofrequency, acupuncture, Kneipp hydrotherapy, complex ayurvedic treatment</p> <p>c. Any of the above interventions, stratified by duration and/or intensity</p>	<p>cartilage extract, BioCell Collagen, UP446</p> <p>b. Non-pharmacological: thermal baths with a sulfide mineral water, exercise therapy, physical therapy, neuromuscular exercise in patients with severe OA of the hip, gluteus medius muscle activity, high-velocity resistance training/low-velocity resistance training, tailored activity pacing, manual therapy, Basic Body Awareness Therapy (BBAT), cognitive behavioral therapy for chronic pain, internet-based pain coping skills training, self-management program, pulsed radiofrequency, acupuncture, Kneipp hydrotherapy, complex ayurvedic treatment</p> <p>c. Any of the above interventions, stratified by duration and/or intensity</p>	<p>capsaicin, TNF-inhibitor infliximab, adhesive patches containing Chinese herbal mixtures FNZG and SJG, marhame-aafasel compress (herbal), stinging nettle leaf (topical)</p> <p>b: Non-pharmacological: Occupational therapy (strengthening, splinting, etc), mud-bath treatment, paraffin bath treatment, thermal bath with sulfate-calcium-magnesium-fluoride mineral water, low-level laser therapy (LLLT)</p> <p>c: Any of the above interventions, stratified by duration and/or intensity</p>	<p>capsaicin, TNF-inhibitor infliximab, adhesive patches containing Chinese herbal mixtures FNZG and SJG, marhame-aafasel compress (herbal), stinging nettle leaf (topical)</p> <p>b: Non-pharmacological: Occupational therapy (strengthening, splinting, etc), mud-bath treatment, paraffin bath treatment, thermal bath with sulfate-calcium-magnesium-fluoride mineral water, low-level laser therapy (LLLT)</p> <p>c: Any of the above interventions, stratified by duration and/or intensity</p>
<b>Comparators</b>	Placebo/sham controls, other active interventions	Placebo/sham controls, other active interventions	Placebo/sham controls, other active interventions	Placebo/sham controls, other active interventions
<b>Outcomes</b>	Reduced pain, improved function/mobility, QoL, functional status measures (including Western Ontario and McMaster Universities Arthritis Index [WOMAC] scores), other clinical outcomes	Adverse events	Reduced pain, improved function/mobility, QoL, functional status measures (including Western Ontario and McMaster Universities Arthritis Index [WOMAC] scores), other clinical outcomes	Adverse events

## Methods

To assess topic nomination #0712 *Treatments for Hip and Hand Osteoarthritis* for priority for a systematic review or other AHRQ EHC report, we used a modified process based on established criteria. Our assessment is hierarchical in nature, with the findings of each step in our assessment determining the need for further evaluation of the next step. Details related to our assessment are provided in Appendix A.

1. Determine the *appropriateness* of the nominated topic for inclusion in the EHC program.
2. Establish the overall *importance* of a potential topic as representing a health or healthcare issue in the United States.
3. Determine the *desirability of new evidence review* by examining whether a new systematic review or other AHRQ product would be duplicative.
4. Assess the *potential impact* a new systematic review or other AHRQ product.
5. Assess whether the *current state of the evidence* allows for a systematic review or other AHRQ product (feasibility).
6. Determine the *potential value* of a new systematic review or other AHRQ product.

### Appropriateness and Importance

We assessed the nomination for appropriateness and importance (see Appendix A).

### Desirability of New Review/Duplication

We searched for high-quality, completed or in-process evidence reviews pertaining to the key questions of the nomination. Table 2 includes the citations for the reviews that were determined to address the key questions.

### Impact of a New Evidence Review

The impact of a new evidence review was assessed by analyzing the current standard of care, the existence of potential knowledge gaps, and practice variation. We considered whether a new review could influence the current state of practice through various dissemination pathways (practice recommendation, clinical guidelines, etc.).

### Feasibility of a New Evidence Review

We conducted a literature search in PubMed from November 2011 to March 2017 (Appendix B). From 958 results, studies were separated into therapies for hand OA, therapies for hip OA, and therapies for general OA. All 154 results for hand OA and 37 results for general OA were reviewed. Of the 767 results for hip OA, 430 were removed for examining surgical options. All 337 remaining results for hip OA were reviewed for inclusion. See *Table 2, Feasibility Column, Size/Scope of Review Section* for the citations of included studies.

### Value

We assessed the nomination for value (see Appendix A). We considered whether a partner organization could use the information from the proposed evidence review to facilitate evidence-based change; or the presence of clinical, consumer, or policymaking context that is amenable to evidence-based change.

### Compilation of Findings

We constructed a table outlining the selection criteria as they pertain to this nomination (see Appendix A).

## Results

### Appropriateness and Importance

This is an appropriate and important topic. OA affects approximately 30.8 million Americans, according to the CDC,<sup>1</sup> and the average direct cost of OA per patient per year is \$2,600, with total (direct + indirect) costs estimated around \$5,700 per patient per year.<sup>1</sup>

### Desirability of New Review/Duplication

An AHRQ systematic review on the treatments for hip and hand OA would not be duplicative. We identified 12 complete or in-process reviews/meta-analyses— one review examining physical and occupational therapy for hand OA, and eleven complete or in-process evidence reviews/meta-analyses covering specific interventions for hip OA. The interventions examined in these eleven hip OA evidence reviews are platelet-rich plasma, opioid therapy, exercise and physical therapy, manual therapy, and high- and low-velocity resistance training. Most of the interventions of interest to the nominators are not included in these reviews.

### Impact of a New Evidence Review

The nomination has moderate impact potential. The ACR published guidelines for nonpharmacological and pharmacologic therapies for hand, hip, and knee OA in 2012, using GRADE and a panel consensus. Their recommendations for both pharmacologic and nonpharmacological therapies for hand OA and pharmacologic therapies for hip OA were “not strong” at that time. The ACR guidelines for non-pharmacological therapies for hip were “strong” recommendations. A recommendation of “not strong” may indicate a knowledge gap that a new evidence review may address. By the time an AHRQ evidence review on this topic would be complete, the ACR guidelines will be more than five years old and in need of updating.

### Feasibility of a New Evidence Review

A new evidence review is feasible at this time. Our search of PubMed resulted in 958 unique titles. 430 results were excluded because they examined surgical procedures. Upon title and abstract review of the remaining 528 results, we identified 93 studies relevant to the key questions, 52 of which were RCTs. Among these included studies were several interventions of interest to the nominator that were not found in our search for systematic reviews. These include NSAIDs and COX-2-selective inhibitors, topical therapies, hydrotherapy, splinting, and corticosteroid injections, among others. We found studies examining 18 different pharmacologic treatments for hip OA, 11 non-pharmacological treatments for hip OA, 11 pharmacologic treatments for hand OA, and 10 non-pharmacological treatments for hand OA.

Pharmacologic interventions for hip OA include diclofenac, celecoxib, and injections of platelet-rich plasma and hyaluronic acid. Nonpharmacologic interventions include web-based and in-person physical activity, physical/manual therapies, and acupuncture. Identified studies for pharmacologic interventions for hand OA are like those found for hip—hyaluronic acid injections, diclofenac, and naproxen, to name a few—however, non-pharmacological interventions for hand OA differed vastly from hip. They included splints/orthotics, paraffin and mud baths, immobilization, and hand exercises.

Our search of ClinicalTrials.gov resulted in 23 relevant trials that have been completed in the last two years or are projected to be completed in the next two years. See *Table 2, Feasibility* column for the citations that were determined to address the key questions.

**Table 2.** Key question with the identified corresponding evidence reviews and original research



Key Question	Duplication (Completed or In-Process Evidence Reviews)	Feasibility (Published and Ongoing Research)
1a: Hip OA-- Pharmacologic	Total number of completed or in-process evidence reviews: 4 <ul style="list-style-type: none"> <li>• Cochrane: 1<sup>2</sup></li> <li>• Meta-Analysis of RCTs: 2<sup>3,4</sup></li> <li>• Other Protocol: 1<sup>5</sup></li> </ul>	<u>Size/scope of review</u> Relevant Studies: 24 <ul style="list-style-type: none"> <li>• RCT: 11<sup>6-16</sup></li> <li>• Prospective Cohort: 4<sup>17-20</sup></li> <li>• Prospective Multicenter: 2<sup>21,22</sup></li> <li>• Open-Label: 6<sup>23-28</sup></li> <li>• Survey: 1<sup>29</sup></li> <li>• Retrospective: 1<sup>30</sup></li> </ul> <u>ClinicalTrials.Gov</u> Relevant Trials: 5 <ul style="list-style-type: none"> <li>• Recruiting: 1<sup>31</sup></li> <li>• Active, not recruiting: 1<sup>32</sup></li> <li>• Complete: 3<sup>33-35</sup></li> </ul>
1b: Hip OA— Non-pharmacologic	Total number of completed or in-process evidence reviews: 7 <ul style="list-style-type: none"> <li>• Cochrane: 3<sup>36-38</sup></li> <li>• Other: 1<sup>39</sup></li> <li>• Cochrane Protocol: 1<sup>40</sup></li> <li>• Other Protocol: 2<sup>41,42</sup></li> </ul>	<u>Size/scope of review</u> Relevant Studies: 29 <ul style="list-style-type: none"> <li>• RCT: 18<sup>43-60</sup></li> <li>• Prospective Cohort: 7<sup>61-67</sup></li> <li>• Prognostic Study: 1<sup>68</sup></li> <li>• Controlled Pre-Post: 1<sup>69</sup></li> <li>• Survey: 2<sup>70,71</sup></li> <li>• Post Hoc Analysis: 1<sup>72</sup></li> </ul> <u>ClinicalTrials.Gov</u> Relevant Trials: 11 <ul style="list-style-type: none"> <li>• Recruiting: 6<sup>73-78</sup></li> <li>• Active, not recruiting: 1<sup>79</sup></li> <li>• Complete: 4<sup>80-83</sup></li> </ul>
1c: Hip OA— Duration and Intensity	Total number of completed on in-process evidence reviews: 2 <ul style="list-style-type: none"> <li>• Cochrane: 1<sup>37</sup></li> <li>• Meta-Analysis of RCTs: 1<sup>4</sup></li> </ul>	<u>Size/scope of review</u> Relevant Studies: 17 <ul style="list-style-type: none"> <li>• RCT: 10<sup>7,12-14,45,48,49,51,53,58</sup></li> <li>• Prospective Cohort: 3<sup>18,62,65</sup></li> <li>• Open-Label: 3<sup>25,26,28</sup></li> <li>• Retrospective: 1<sup>30</sup></li> </ul> <u>ClinicalTrials.Gov</u> Relevant Trials: 3 <ul style="list-style-type: none"> <li>• Recruiting: 3<sup>73,76,78</sup></li> <li>• Active, not recruiting: 1<sup>32</sup></li> <li>• Complete: 2<sup>35,80</sup></li> </ul>
1d: Hip OA-- Subgroups	Total number of completed on in-process evidence reviews: 1 <ul style="list-style-type: none"> <li>• Other: 1<sup>39</sup></li> </ul>	<u>Size/scope of review</u> Relevant Studies: 19 <ul style="list-style-type: none"> <li>• RCT: 9<sup>7,9,12,43,46-48,54,58</sup></li> <li>• Prospective Cohort: 6<sup>17,19,61-63,66</sup></li> <li>• Prospective Multicenter: 1<sup>21</sup></li> <li>• Open-Label: 1<sup>24</sup></li> <li>• Prognostic Study: 1<sup>68</sup></li> <li>• Retrospective: 1<sup>30</sup></li> </ul> <u>ClinicalTrials.Gov</u> Relevant Trials: 10 <ul style="list-style-type: none"> <li>• Recruiting: 4<sup>73,75,77,78</sup></li> </ul>

Key Question	Duplication (Completed or In-Process Evidence Reviews)	Feasibility (Published and Ongoing Research)
		<ul style="list-style-type: none"> <li>Active, not recruiting: 2<sup>32,79</sup></li> <li>Complete: 4<sup>33,34,82,83</sup></li> </ul>
2a: Hip OA— Pharmacologic Harms	Total number of completed on in-process evidence reviews: 3 <ul style="list-style-type: none"> <li>Cochrane: 1<sup>2</sup></li> <li>Meta-Analysis of RCTs: 2<sup>3,4</sup></li> </ul>	<u>Size/scope of review</u> Relevant Studies: 20 <ul style="list-style-type: none"> <li>RCT: 7<sup>6,11-16</sup></li> <li>Prospective Cohort: 5<sup>17,18,20,65,84</sup></li> <li>Prospective Multicenter: 2<sup>21,22</sup></li> <li>Open-Label: 5<sup>23,24,26-28</sup></li> <li>Survey: 1<sup>29</sup></li> <li>Retrospective: 1<sup>30</sup></li> </ul> <u>ClinicalTrials.Gov</u> Relevant Trials: 5 <ul style="list-style-type: none"> <li>Recruiting: 1<sup>31</sup></li> <li>Active, not recruiting: 1<sup>32</sup></li> <li>Complete: 3<sup>33-35</sup></li> </ul>
2b: Hip OA— Non-pharmacologic Harms	Total number of completed on in-process evidence reviews: 3 <ul style="list-style-type: none"> <li>Cochrane: 3<sup>36-38</sup></li> </ul>	<u>Size/scope of review</u> Relevant Studies Identified: 4 <ul style="list-style-type: none"> <li>RCT: 3<sup>47,56,85</sup></li> <li>Prospective Cohort: 1<sup>61</sup></li> </ul> <u>ClinicalTrials.Gov</u> Relevant Trials: 3 <ul style="list-style-type: none"> <li>Recruiting: 1<sup>75</sup></li> <li>Active, not recruiting: 1<sup>79</sup></li> <li>Complete: 1<sup>82</sup></li> </ul>
2c: Hip OA— Duration and Intensity Harms	Total number of completed on in-process evidence reviews: 2 <ul style="list-style-type: none"> <li>Cochrane: 1<sup>37</sup></li> <li>Meta-Analysis of RCTs: 1<sup>4</sup></li> </ul>	<u>Size/scope of review</u> Relevant Studies: 8 <ul style="list-style-type: none"> <li>RCT: 3<sup>12-14</sup></li> <li>Prospective Cohort: 2<sup>18,65</sup></li> <li>Open-Label: 2<sup>26,28</sup></li> <li>Retrospective: 1<sup>30</sup></li> </ul> <u>ClinicalTrials.Gov</u> Relevant Trials: 2 <ul style="list-style-type: none"> <li>Active, not recruiting: 1<sup>32</sup></li> <li>Complete: 1<sup>35</sup></li> </ul>
2d: Hip OA— Subgroups Harms	Total number of completed on in-process evidence reviews: 1 <ul style="list-style-type: none"> <li>Other: 1<sup>39</sup></li> </ul>	<u>Size/scope of review</u> Relevant Studies: 8 <ul style="list-style-type: none"> <li>RCT: 2<sup>12,47</sup></li> <li>Prospective Cohort: 3<sup>17,61,62</sup></li> <li>Prospective Multicenter: 1<sup>21</sup></li> <li>Open-Label: 1<sup>24</sup></li> <li>Retrospective: 1<sup>30</sup></li> </ul> <u>ClinicalTrials.Gov</u> Relevant Trials: 6 <ul style="list-style-type: none"> <li>Recruiting: 1<sup>75</sup></li> <li>Active, not recruiting: 2<sup>32,79</sup></li> <li>Complete: 3<sup>33,34,82</sup></li> </ul>
3a: Hand OA-- Pharmacologic	None identified.	<u>Size/scope of review</u> Relevant Studies: 11

Key Question	Duplication (Completed or In-Process Evidence Reviews)	Feasibility (Published and Ongoing Research)
		<ul style="list-style-type: none"> <li>• RCT: 8<sup>86-93</sup></li> <li>• Prospective Cohort: 1<sup>94</sup></li> <li>• Observational Cross-sectional: 1<sup>95</sup></li> <li>• Post Hoc Analysis: 1<sup>96</sup></li> </ul> <p><u>ClinicalTrials.Gov</u> Relevant Trials: 2</p> <ul style="list-style-type: none"> <li>• Recruiting: 1<sup>97</sup></li> <li>• Complete: 1<sup>98</sup></li> </ul>
3b: Hand OA—Non-pharmacologic	Total number of completed or in-process evidence reviews: 1 <ul style="list-style-type: none"> <li>• Other: 1<sup>99</sup></li> </ul>	<p><u>Size/scope of review</u> Relevant Studies: 21</p> <ul style="list-style-type: none"> <li>• RCT: 14<sup>100-113</sup></li> <li>• Prospective Cohort: 4<sup>114-117</sup></li> <li>• Retrospective Cohort Analysis: 1<sup>118</sup></li> <li>• Case Series: 2<sup>119,120</sup></li> <li>• Cross-sectional case series: 1<sup>121</sup></li> <li>• Post Hoc Analysis: 1<sup>122</sup></li> </ul> <p><u>ClinicalTrials.Gov</u> Relevant Trials: 5</p> <ul style="list-style-type: none"> <li>• Recruiting: 2<sup>123,124</sup></li> <li>• Complete: 3<sup>125-127</sup></li> </ul>
3c: Hand OA—Duration and Intensity	Total number of completed or in-process evidence reviews: 1 <ul style="list-style-type: none"> <li>• Other: 1<sup>99</sup></li> </ul>	<p><u>Size/scope of review</u> Relevant Studies: 8</p> <ul style="list-style-type: none"> <li>• RCT: 5<sup>88,100,109,110,112</sup></li> <li>• Prospective Cohort: 2<sup>114,115</sup></li> <li>• Survey: 1<sup>128</sup></li> </ul> <p><u>ClinicalTrials.Gov</u> Relevant Trials: 2</p> <ul style="list-style-type: none"> <li>• Recruiting: 2<sup>123,124</sup></li> </ul>
3d: Hand OA--Subgroups	None identified.	<p><u>Size/scope of review</u> Relevant Studies: 17</p> <ul style="list-style-type: none"> <li>• RCT: 10<sup>86,89,90,92,103,104,106,107,110,111</sup></li> <li>• Prospective Cohort: 2<sup>94,117</sup></li> <li>• Observational Cross-sectional: 1<sup>95</sup></li> <li>• Case Series: 1<sup>119</sup></li> <li>• Post Hoc Analysis: 3<sup>96,122,129</sup></li> </ul> <p><u>ClinicalTrials.Gov</u> Relevant Trials: 2</p> <ul style="list-style-type: none"> <li>• Recruiting: 1<sup>124</sup></li> <li>• Complete: 1<sup>126</sup></li> </ul>
4a: Hand OA—Pharmacologic Harms	None identified.	<p><u>Size/scope of review</u> Relevant Studies: 8</p> <ul style="list-style-type: none"> <li>• RCT: 6<sup>86,88-90,92,93</sup></li> <li>• Observational Cross-sectional: 1<sup>95</sup></li> <li>• Post Hoc Analysis: 1<sup>96</sup></li> </ul> <p><u>ClinicalTrials.Gov</u> None identified.</p>
4b: Hand OA—	None identified.	<u>Size/scope of review</u>

Key Question	Duplication (Completed or In-Process Evidence Reviews)	Feasibility (Published and Ongoing Research)
Non-pharmacologic Harms		Relevant Studies: 8 <ul style="list-style-type: none"> <li>• RCT: 6<sup>105-107,110-112</sup></li> <li>• Case Series: 1<sup>119</sup></li> <li>• Post Hoc analysis: 1<sup>122</sup></li> </ul> <a href="#">ClinicalTrials.Gov</a> None identified.
4c: Hand OA—Duration and Intensity Harms	None identified.	<u>Size/scope of review</u> Relevant Studies: 3 <ul style="list-style-type: none"> <li>• RCT: 3<sup>88,110,112</sup></li> </ul> <a href="#">ClinicalTrials.Gov</a> None identified.
4d: Hand OA—Subgroups Harms	None identified.	<u>Size/scope of review</u> Relevant Studies: 12 <ul style="list-style-type: none"> <li>• RCT: 6<sup>86,89,90,92,106,111</sup></li> <li>• Prospective Cohort: 1<sup>117</sup></li> <li>• Observational Cross-sectional: 1<sup>95</sup></li> <li>• Case Series: 1<sup>119</sup></li> <li>• Post Hoc Analysis: 3<sup>96,122,129</sup></li> </ul> <a href="#">ClinicalTrials.Gov</a> None identified.

*Abbreviations:* OA=Osteoarthritis; RCT=Randomized Controlled Trial

## Value

The nomination has a high value potential, given that the ACR and the ACP will use a new AHRQ systematic review to update their 2012 guidelines.

## Summary of Findings

- Appropriateness and importance: The nomination is both appropriate and important.
- Duplication: An AHRQ systematic review on the treatments for hip and hand osteoarthritis (OA) would not be duplicative. We identified 12 complete or in-process reviews/meta-analyses—one review examining physical and occupational therapy for hand OA, and eleven complete or in-process evidence reviews/meta-analyses covering specific interventions for hip OA. The interventions examined in these eleven hip OA evidence reviews are platelet-rich plasma, opioid therapy, exercise and physical therapy, manual therapy, and high- and low-velocity resistance training. Most of the interventions of interest to the nominators are not included in these reviews.
- Impact: The nomination has moderate impact potential. The ACR published guidelines for [nonpharmacological and pharmacologic therapies for hand, hip, and knee osteoarthritis](#) in 2012, using GRADE and a panel consensus. An updated review could inform evidence gaps for recommendations that were not considered strong, particularly related to therapies for hand OA, hip OA. A recommendation of "not strong" may indicate a knowledge gap that a new evidence review may address. By the time an AHRQ evidence review on this topic will be complete, the ACR guidelines will be more than five years old and in need of updating.
- Feasibility: An AHRQ evidence review is feasible at this time.

- *Size/scope of review:* We identified 93 studies potentially relevant to the key questions in the nomination.
- *Clinicaltrials.gov:* We identified 23 ongoing or recently completed trials on ClinicalTrials.gov.
- *Value:* The nomination has a high value potential, given that the ACR and the ACP will use a new AHRQ systematic review to update their 2012 guidelines.

## References

1. Amano T, Hasegawa Y, Seki T, Takegami Y, Murotani K, Ishiguro N. A pre-operative predictive score for the outcome of eccentric rotational acetabular osteotomy in the treatment of acetabular dysplasia and early osteoarthritis of the hip in adults. *The bone & joint journal*. Oct 2016;98-b(10):1326-1332.
2. da Costa BR, Nuesch E, Kasteler R, et al. Oral or transdermal opioids for osteoarthritis of the knee or hip. *The Cochrane database of systematic reviews*. Sep 17 2014(9):Cd003115.
3. Etropolski M, Kuperwasser B, Flugel M, et al. Safety and tolerability of tapentadol extended release in moderate to severe chronic osteoarthritis or low back pain management: pooled analysis of randomized controlled trials. *Adv Ther*. Jun 2014;31(6):604-620.
4. da Costa BR, Reichenbach S, Keller N, et al. Effectiveness of non-steroidal anti-inflammatory drugs for the treatment of pain in knee and hip osteoarthritis: a network meta-analysis. *Lancet (London, England)*. May 21 2016;387(10033):2093-2105.
5. Ivan Medina PA. The efficacy of platelet-rich plasma injection in the management of hip osteoarthritis: a systematic review. *PROSPERO* 2014;CRD42014010210
6. Balanescu AR, Feist E, Wolfram G, et al. Efficacy and safety of tanezumab added on to diclofenac sustained release in patients with knee or hip osteoarthritis: a double-blind, placebo-controlled, parallel-group, multicentre phase III randomised clinical trial. *Ann Rheum Dis*. Sep 2014;73(9):1665-1672.
7. Battaglia M, Guaraldi F, Vannini F, et al. Efficacy of ultrasound-guided intra-articular injections of platelet-rich plasma versus hyaluronic acid for hip osteoarthritis. *Orthopedics*. Dec 2013;36(12):e1501-1508.
8. Fujii T, Takana K, Orita S, et al. Progressive change in joint degeneration in patients with knee or hip osteoarthritis treated with fentanyl in a randomized trial. *Yonsei Med J*. Sep 2014;55(5):1379-1385.
9. Maheu E, Cadet C, Marty M, et al. Randomised, controlled trial of avocado-soybean unsaponifiable (Piascledine) effect on structure modification in hip osteoarthritis: the ERADIAS study. *Ann Rheum Dis*. Feb 2014;73(2):376-384.
10. Nishii T, Tamura S, Shiomi T, Yoshikawa H, Sugano N. Alendronate treatment for hip osteoarthritis: prospective randomized 2-year trial. *Clinical rheumatology*. Dec 2013;32(12):1759-1766.
11. Prior MJ, Harrison DD, Frustaci ME. A randomized, double-blind, placebo-controlled 12 week trial of acetaminophen extended release for the treatment of signs and symptoms of osteoarthritis. *Curr Med Res Opin*. Nov 2014;30(11):2377-2387.
12. Strand V, Simon LS, Dougados M, et al. Treatment of osteoarthritis with continuous versus intermittent celecoxib. *J Rheumatol*. Dec 2011;38(12):2625-2634.
13. Altman R, Hochberg M, Gibofsky A, Jaros M, Young C. Efficacy and safety of low-dose SoluMatrix meloxicam in the treatment of osteoarthritis pain: a 12-week, phase 3 study. *Curr Med Res Opin*. Dec 2015;31(12):2331-2343.

14. Ekman EF, Gimbel JS, Bello AE, et al. Efficacy and safety of intravenous tanezumab for the symptomatic treatment of osteoarthritis: 2 randomized controlled trials versus naproxen. *J Rheumatol*. Nov 2014;41(11):2249-2259.
15. Gibofsky A, Hochberg MC, Jaros MJ, Young CL. Efficacy and safety of low-dose submicron diclofenac for the treatment of osteoarthritis pain: a 12 week, phase 3 study. *Curr Med Res Opin*. Sep 2014;30(9):1883-1893.
16. Dallari D, Stagni C, Rani N, et al. Ultrasound-Guided Injection of Platelet-Rich Plasma and Hyaluronic Acid, Separately and in Combination, for Hip Osteoarthritis: A Randomized Controlled Study. *The American journal of sports medicine*. Mar 2016;44(3):664-671.
17. Subedi N, Chew NS, Chandramohan M, Scally AJ, Groves C. Effectiveness of fluoroscopy-guided intra-articular steroid injection for hip osteoarthritis. *Clin Radiol*. Nov 2015;70(11):1276-1280.
18. Migliore A, Massafra U, Bizzi E, et al. Intra-articular injection of hyaluronic acid (MW 1,500-2,000 kDa; HyalOne) in symptomatic osteoarthritis of the hip: a prospective cohort study. *Archives of orthopaedic and trauma surgery*. Dec 2011;131(12):1677-1685.
19. Clockaerts S, Van Osch GJ, Bastiaansen-Jenniskens YM, et al. Statin use is associated with reduced incidence and progression of knee osteoarthritis in the Rotterdam study. *Ann Rheum Dis*. May 2012;71(5):642-647.
20. Emadedin M, Ghorbani Liastani M, Fazeli R, et al. Long-Term Follow-up of Intra-articular Injection of Autologous Mesenchymal Stem Cells in Patients with Knee, Ankle, or Hip Osteoarthritis. *Arch Iran Med*. Jun 2015;18(6):336-344.
21. Oteo-Alvaro A, Marin MT, Ruiz-Iban MA, Armada B, Rejas J. Treatment satisfaction after switching to another therapy in Spanish orthopaedic clinic outpatients with knee or hip osteoarthritis previously refractory to paracetamol. *Clin Drug Investig*. Oct 1 2012;32(10):685-695.
22. Zamani O, Bottcher E, Rieger JD, et al. Comparison of safety, efficacy and tolerability of dexibuprofen and ibuprofen in the treatment of osteoarthritis of the hip or knee. *Wien Klin Wochenschr*. Jun 2014;126(11-12):368-375.
23. Migliore A, Tormenta S, Lagana B, et al. Safety of intra-articular hip injection of hyaluronic acid products by ultrasound guidance: an open study from ANTIAGE register. *Eur Rev Med Pharmacol Sci*. Jul 2013;17(13):1752-1759.
24. Mordarski S. Pain management in the elderly: transdermal fentanyl for the treatment of pain caused by osteoarthritis of the knee and hip. *BioMed research international*. 2014;2014:262961.
25. Paoloni M, Di Sante L, Dimaggio M, et al. Kinematic and kinetic modifications in walking pattern of hip osteoarthritis patients induced by intra-articular injections of hyaluronic acid. *Clinical biomechanics (Bristol, Avon)*. Aug 2012;27(7):661-665.
26. Sanchez M, Guadilla J, Fiz N, Andia I. Ultrasound-guided platelet-rich plasma injections for the treatment of osteoarthritis of the hip. *Rheumatology (Oxford)*. Jan 2012;51(1):144-150.
27. Zheng Y, Kostenbader K, Barrett T, et al. Tolerability of Biphasic-Release Hydrocodone Bitartrate/Acetaminophen Tablets (MNK-155): A Phase III, Multicenter, Open-Label Study in Patients With Osteoarthritis or Chronic Low Back Pain. *Clin Ther*. Jun 1 2015;37(6):1235-1247.
28. Altman RD, Strand V, Hochberg MC, et al. Low-dose SoluMatrix diclofenac in the treatment of osteoarthritis: A 1-year, open-label, Phase III safety study. *Postgrad Med*. Jun 2015;127(5):517-528.
29. Migliore A, Massafra U, Bizzi E, Tormenta S, Cassol M, Granata M. Duration of symptom relief after intra-articular injection of hyaluronic acid combined with sorbitol

- (anti-ox-vs) in symptomatic hip osteoarthritis. *Int J Immunopathol Pharmacol*. Apr-Jun 2014;27(2):245-252.
30. Park KD, Kim TK, Bae BW, Ahn J, Lee WY, Park Y. Ultrasound guided intra-articular ketorolac versus corticosteroid injection in osteoarthritis of the hip: a retrospective comparative study. *Skeletal Radiol*. Sep 2015;44(9):1333-1340.
  31. Laval University. Daily Activity and Gait Analysis After Viscosupplement Injection Among Hip Osteoarthritis Patients. *ClinicalTrials.gov*. 2016;NCT02086474.
  32. VA Office of Research and Development. Strategies for Prescribing Analgesics Comparative Effectiveness Trial (SPACE). *ClinicalTrials.gov*. 2016;NCT01583985.
  33. Genzyme. A Study of the Safety and Effectiveness of Synvisc-One® (Hylan G-F 20) in Patients With Primary Osteoarthritis of the Hip. *ClinicalTrials.gov*. 2016;NCT01618708.
  34. Meloxicam. Study of Meloxicam Capsules in Subjects With Osteoarthritis of the Knee or Hip. *ClinicalTrials.gov*. 2015;NCT01801735.
  35. GlaxoSmithKline. A 12-Week Efficacy Study of Paracetamol 1000mg Sustained-release Tablets in Patients With Osteoarthritis. *ClinicalTrials.gov*. 2016;NCT02311881.
  36. Bartels EM, Juhl CB, Christensen R, et al. Aquatic exercise for the treatment of knee and hip osteoarthritis. *The Cochrane database of systematic reviews*. Mar 23 2016;3:CD005523.
  37. Regnaud JP, Lefevre-Colau MM, Trinquart L, et al. High-intensity versus low-intensity physical activity or exercise in people with hip or knee osteoarthritis. *The Cochrane database of systematic reviews*. Oct 29 2015(10):CD010203.
  38. Fransen M, McConnell S, Hernandez-Molina G, Reichenbach S. Exercise for osteoarthritis of the hip. *Cochrane Database of Systematic Reviews*. 2014(4).
  39. Stubbs B, Hurley M, Smith T. What are the factors that influence physical activity participation in adults with knee and hip osteoarthritis? A systematic review of physical activity correlates. *Clin Rehabil*. Jan 2015;29(1):80-94.
  40. Helen French RG, J Haxby Abbott, Marlene Fransen Adjunctive therapies in addition to land-based exercise therapy for osteoarthritis of the hip or knee. *PROSPERO* 2016;CRD42016033609
  41. Qiong Wang TW, Xiaofeng Qi, Qianqian Liang, Xuejun Cui, Min Yao, Yongjun Wang Manual therapy for the relief of chronic pain due to hip osteoarthritis: a systematic review and meta-analysis. *PROSPERO* 2014;CRD42014014851.
  42. Kesava Kovanur Sampath RM, Steve Tumilty Effectiveness of manual therapy (MT) or exercise therapy (ET) or both on pain, physical function and quality of life in patients with hip osteoarthritis (OA): a systematic review and meta-analysis. *PROSPERO* 2013;CRD42013005363
  43. Bossen D, Veenhof C, Van Beek KE, Spreeuwenberg PM, Dekker J, De Bakker DH. Effectiveness of a web-based physical activity intervention in patients with knee and/or hip osteoarthritis: randomized controlled trial. *J Med Internet Res*. Nov 22 2013;15(11):e257.
  44. Brantingham JW, Parkin-Smith G, Cassa TK, et al. Full kinetic chain manual and manipulative therapy plus exercise compared with targeted manual and manipulative therapy plus exercise for symptomatic osteoarthritis of the hip: a randomized controlled trial. *Arch Phys Med Rehabil*. Feb 2012;93(2):259-267.
  45. French HP, Cusack T, Brennan A, et al. Exercise and manual physiotherapy arthritis research trial (EMPART) for osteoarthritis of the hip: a multicenter randomized controlled trial. *Arch Phys Med Rehabil*. Feb 2013;94(2):302-314.

46. Murphy SL, Smith DM, Lyden AK. Type of activity pacing instruction affects physical activity variability in adults with symptomatic knee or hip osteoarthritis. *J Phys Act Health*. Mar 2012;9(3):360-366.
47. Svege I, Nordsletten L, Fernandes L, Risberg MA. Exercise therapy may postpone total hip replacement surgery in patients with hip osteoarthritis: a long-term follow-up of a randomised trial. *Ann Rheum Dis*. Jan 2015;74(1):164-169.
48. Martins F, Kaster T, Schutzler L, Witt CM. Factors influencing further acupuncture usage and a more positive outcome in patients with osteoarthritis of the knee and the hip: a 3-year follow-up of a randomized pragmatic trial. *Clin J Pain*. Nov 2014;30(11):953-959.
49. Poulsen E, Hartvigsen J, Christensen HW, Roos EM, Vach W, Overgaard S. Patient education with or without manual therapy compared to a control group in patients with osteoarthritis of the hip. A proof-of-principle three-arm parallel group randomized clinical trial. *Osteoarthritis Cartilage*. Oct 2013;21(10):1494-1503.
50. Schencking M, Wilm S, Redaelli M. A comparison of Kneipp hydrotherapy with conventional physiotherapy in the treatment of osteoarthritis: a pilot trial. *J Integr Med*. Jan 2013;11(1):17-25.
51. Schepens SL, Braun ME, Murphy SL. Effect of tailored activity pacing on self-perceived joint stiffness in adults with knee or hip osteoarthritis. *Am J Occup Ther*. May-Jun 2012;66(3):363-367.
52. Teirlinck CH, Luijsterburg PA, Dekker J, et al. Effectiveness of exercise therapy added to general practitioner care in patients with hip osteoarthritis: a pragmatic randomized controlled trial. *Osteoarthritis Cartilage*. Jan 2016;24(1):82-90.
53. Tejedor Varillas A, Leon Vazquez F, Lora Pablos D, Perez Martin A, Vargas Negrin F, Gomez de la Camara A. Can an intervention on clinical inertia have an impact on the perception of pain, functionality and quality of life in patients with hip and/or knee osteoarthritis? Results from a cluster randomised trial. *Aten Primaria*. Feb 2012;44(2):65-72.
54. Villadsen A, Overgaard S, Holsgaard-Larsen A, Christensen R, Roos EM. Immediate efficacy of neuromuscular exercise in patients with severe osteoarthritis of the hip or knee: a secondary analysis from a randomized controlled trial. *J Rheumatol*. Jul 2014;41(7):1385-1394.
55. Abbott JH, Robertson MC, Chapple C, et al. Manual therapy, exercise therapy, or both, in addition to usual care, for osteoarthritis of the hip or knee: a randomized controlled trial. 1: clinical effectiveness. *Osteoarthritis Cartilage*. Apr 2013;21(4):525-534.
56. Bennell KL, Egerton T, Martin J, et al. Effect of physical therapy on pain and function in patients with hip osteoarthritis: a randomized clinical trial. *JAMA*. May 21 2014;311(19):1987-1997.
57. Bossen D, Buskermolen M, Veenhof C, de Bakker D, Dekker J. Adherence to a web-based physical activity intervention for patients with knee and/or hip osteoarthritis: a mixed method study. *J Med Internet Res*. Oct 16 2013;15(10):e223.
58. Fukumoto Y, Tateuchi H, Ikezoe T, et al. Effects of high-velocity resistance training on muscle function, muscle properties, and physical performance in individuals with hip osteoarthritis: a randomized controlled trial. *Clin Rehabil*. Jan 2014;28(1):48-58.
59. Krauss I, Steinhilber B, Haupt G, Miller R, Martus P, Janssen P. Exercise therapy in hip osteoarthritis--a randomized controlled trial. *Dtsch Arztebl Int*. Sep 1 2014;111(35-36):592-599.
60. Allen KD, Yancy WS, Jr., Bosworth HB, et al. A Combined Patient and Provider Intervention for Management of Osteoarthritis in Veterans: A Randomized Clinical Trial. *Annals of internal medicine*. Jan 19 2016;164(2):73-83.



61. Angst F, Verra ML, Lehmann S, Benz T, Aeschlimann A. Effects of inpatient rehabilitation in hip and knee osteoarthritis: a naturalistic prospective cohort study with intraindividual control of effects. *Arch Phys Med Rehabil*. Nov 2013;94(11):2139-2145.
62. Smink AJ, Dekker J, Vliet Vlieland TP, et al. Health care use of patients with osteoarthritis of the hip or knee after implementation of a stepped-care strategy: an observational study. *Arthritis Care Res (Hoboken)*. Jun 2014;66(6):817-827.
63. Paans N, van den Akker-Scheek I, Dilling RG, et al. Effect of exercise and weight loss in people who have hip osteoarthritis and are overweight or obese: a prospective cohort study. *Phys Ther*. Feb 2013;93(2):137-146.
64. Skou ST, Odgaard A, Rasmussen JO, Roos EM. Group education and exercise is feasible in knee and hip osteoarthritis. *Danish medical journal*. Dec 2012;59(12):A4554.
65. Anderson ES, Hodell E, Mantuani D, Fahimi J, Pampalone I, Nagdev A. Pilot study of ultrasound-guided corticosteroid hip injections by emergency physicians. *West J Emerg Med*. Nov 2014;15(7):919-924.
66. Holsgaard-Larsen A, Roos EM. Objectively measured physical activity in patients with end stage knee or hip osteoarthritis. *Eur J Phys Rehabil Med*. Dec 2012;48(4):577-585.
67. Jigami H, Sato D, Tsubaki A, et al. Effects of weekly and fortnightly therapeutic exercise on physical function and health-related quality of life in individuals with hip osteoarthritis. *Journal of orthopaedic science : official journal of the Japanese Orthopaedic Association*. Nov 2012;17(6):737-744.
68. French HP, Galvin R, Cusack T, McCarthy GM. Predictors of short-term outcome to exercise and manual therapy for people with hip osteoarthritis. *Phys Ther*. Jan 2014;94(1):31-39.
69. Ageberg E, Nilsson A, Kosek E, Roos EM. Effects of neuromuscular training (NEMEX-TJR) on patient-reported outcomes and physical function in severe primary hip or knee osteoarthritis: a controlled before-and-after study. *BMC musculoskeletal disorders*. Aug 08 2013;14:232.
70. Hinman RS, Nicolson PJ, Dobson FL, Bennell KL. Use of nondrug, nonoperative interventions by community-dwelling people with hip and knee osteoarthritis. *Arthritis Care Res (Hoboken)*. Feb 2015;67(2):305-309.
71. Hofstede SN, Marang-van de Mheen PJ, Vliet Vlieland TP, van den Ende CH, Nelissen RG, van Bodegom-Vos L. Barriers and Facilitators Associated with Non-Surgical Treatment Use for Osteoarthritis Patients in Orthopaedic Practice. *PloS one*. 2016;11(1):e0147406.
72. Eitzen I, Fernandes L, Nordsletten L, Risberg MA. No effects of a 12-week supervised exercise therapy program on gait in patients with mild to moderate osteoarthritis: a secondary analysis of a randomized trial. *J Negat Results Biomed*. Mar 05 2015;14:5.
73. Universidad de Zaragoza. Clinical Effects of Translatory Grades of Movement in Hip Osteoarthritis Patients. *ClinicalTrials.gov*. 2015;NCT02498314.
74. Davis A. STOP OA: A PILOT STUDY (GLA:D Canada) - Education and Targeted, Personalized Exercise for Hip and Knee Osteoarthritis. *ClinicalTrials.gov*. 2016;NCT02693873.
75. University of Southern Denmark. Effect of Exercise as Non-surgical Treatments on Time to Total Hip Replacement Surgery (HipSPORT). *ClinicalTrials.gov*. 2014;NCT01697241.
76. Universidad de Zaragoza. Effects of Grade III Traction-mobilization in End-range Joint Position in Patients With Hip Osteoarthritis. *ClinicalTrials.gov*. 2016;NCT02469025.

77. McMaster University. Resilience for Older Workers With OA Through Exercise. *ClinicalTrials.gov*. 2015;NCT02609672.
78. University of North Carolina, Chapel Hill. Pain Coping Skills Training for African Americans With Osteoarthritis (STAART). *ClinicalTrials.gov*. 2016;NCT02560922.
79. Diakonhjemmet Hospital. Management of Hip and Knee Osteoarthritis in Primary Health Care. *ClinicalTrials.gov*. 2016;NCT02333656.
80. Norwegian University of Science and Technology. Dose-response: Exercise Therapy on Hip Osteoarthritis. *ClinicalTrials.gov*. 2016;NCT01700933.
81. Campelo NMO. Mobilization With Movement in Patients With Osteoarthritis of the Hip (MWM). *ClinicalTrials.gov*. 2015;NCT02390336.
82. University of North Carolina, Chapel Hill. Shoe Lifts for Leg Length Inequality in Adults With Knee or Hip Symptoms. *ClinicalTrials.gov*. 2016;NCT01894100.
83. Duke University. Pilot Testing a Patient and Provider Intervention for Managing Osteoarthritis in Hispanic Adults. (PRIMO-Latino). *ClinicalTrials.gov*. 2015;NCT01782417.
84. Snijders GF, van den Ende CH, van den Bemt BJ, van Riel PL, van den Hoogen FH, den Broeder AA. Treatment outcomes of a Numeric Rating Scale (NRS)-guided pharmacological pain management strategy in symptomatic knee and hip osteoarthritis in daily clinical practice. *Clin Exp Rheumatol*. Mar-Apr 2012;30(2):164-170.
85. Hale LA, Waters D, Herbison P. A randomized controlled trial to investigate the effects of water-based exercise to improve falls risk and physical function in older adults with lower-extremity osteoarthritis. *Arch Phys Med Rehabil*. Jan 2012;93(1):27-34.
86. Chevalier X, Ravaud P, Maheu E, et al. Adalimumab in patients with hand osteoarthritis refractory to analgesics and NSAIDs: a randomised, multicentre, double-blind, placebo-controlled trial. *Ann Rheum Dis*. Sep 2015;74(9):1697-1705.
87. Jahangiri A, Moghaddam FR, Najafi S. Hypertonic dextrose versus corticosteroid local injection for the treatment of osteoarthritis in the first carpometacarpal joint: a double-blind randomized clinical trial. *Journal of orthopaedic science : official journal of the Japanese Orthopaedic Association*. Sep 2014;19(5):737-743.
88. Monfort J, Rotes-Sala D, Segales N, et al. Comparative efficacy of intra-articular hyaluronic acid and corticoid injections in osteoarthritis of the first carpometacarpal joint: results of a 6-month single-masked randomized study. *Joint Bone Spine*. Mar 2015;82(2):116-121.
89. Peniston JH, Gold MS, Wieman MS, Alwine LK. Long-term tolerability of topical diclofenac sodium 1% gel for osteoarthritis in seniors and patients with comorbidities. *Clin Interv Aging*. 2012;7:517-523.
90. Roth SH, Fuller P. Pooled safety analysis of diclofenac sodium topical solution 1.5% (w/w) in the treatment of osteoarthritis in patients aged 75 years or older. *Clin Interv Aging*. 2012;7:127-137.
91. Sanders D, Krause K, O'Muircheartaigh J, et al. Pharmacologic modulation of hand pain in osteoarthritis: a double-blind placebo-controlled functional magnetic resonance imaging study using naproxen. *Arthritis Rheumatol*. Mar 2015;67(3):741-751.
92. polidoro Paschoal Nde O, Natour J, Machado FS, de Oliveira HA, Furtado RN. Effectiveness of Triamcinolone Hexacetonide Intraarticular Injection in Interphalangeal Joints: A 12-week Randomized Controlled Trial in Patients with Hand Osteoarthritis. *J Rheumatol*. Oct 2015;42(10):1869-1877.

93. Wenham CY, Hensor EM, Grainger AJ, et al. A randomized, double-blind, placebo-controlled trial of low-dose oral prednisolone for treating painful hand osteoarthritis. *Rheumatology (Oxford)*. Dec 2012;51(12):2286-2294.
94. Klauser AS, Faschingbauer R, Kupferthaler K, et al. Sonographic criteria for therapy follow-up in the course of ultrasound-guided intra-articular injections of hyaluronic acid in hand osteoarthritis. *Eur J Radiol*. Jul 2012;81(7):1607-1611.
95. Reginato AM, Riera H, Vera M, et al. Osteoarthritis in Latin America: Study of Demographic and Clinical Characteristics in 3040 Patients. *J Clin Rheumatol*. Dec 2015;21(8):391-397.
96. Baraf HS, Gold MS, Petruschke RA, Wieman MS. Tolerability of topical diclofenac sodium 1% gel for osteoarthritis in seniors and patients with comorbidities. *Am J Geriatr Pharmacother*. Feb 2012;10(1):47-60.
97. Bioiberica. Primary Care Study With Chondroitin Sulfate and Glucosamine Hydrochloride in Hand Osteoarthritis (PICASSO). *ClinicalTrials.gov*. 2016;NCT02823548.
98. St George's, University of London. Pain Management in Osteoarthritis Using the Centrally Acting Analgesics Duloxetine and Pregabalin (DUPRO). *ClinicalTrials.gov*. 2016;NCT02612233.
99. Aebischer B, Elsig S, Taeymans J. Effectiveness of physical and occupational therapy on pain, function and quality of life in patients with trapeziometacarpal osteoarthritis - A systematic review and meta-analysis. *Hand therapy*. Mar 2016;21(1):5-15.
100. Villafane JH, Cleland JA, Fernandez-de-Las-Penas C. The effectiveness of a manual therapy and exercise protocol in patients with thumb carpometacarpal osteoarthritis: a randomized controlled trial. *The Journal of orthopaedic and sports physical therapy*. Apr 2013;43(4):204-213.
101. Dilek B, Gozum M, Sahin E, et al. Efficacy of paraffin bath therapy in hand osteoarthritis: a single-blinded randomized controlled trial. *Arch Phys Med Rehabil*. Apr 2013;94(4):642-649.
102. Bani MA, Arazpour M, Kashani RV, Mousavi ME, Hutchins SW. Comparison of custom-made and prefabricated neoprene splinting in patients with the first carpometacarpal joint osteoarthritis. *Disabil Rehabil Assist Technol*. May 2013;8(3):232-237.
103. Becker SJ, Bot AG, Curley SE, Jupiter JB, Ring D. A prospective randomized comparison of neoprene vs thermoplast hand-based thumb spica splinting for trapeziometacarpal arthrosis. *Osteoarthritis Cartilage*. May 2013;21(5):668-675.
104. Dziedzic K, Nicholls E, Hill S, et al. Self-management approaches for osteoarthritis in the hand: a 2x2 factorial randomised trial. *Ann Rheum Dis*. Jan 2015;74(1):108-118.
105. Fioravanti A, Tenti S, Giannitti C, Fortunati NA, Galeazzi M. Short- and long-term effects of mud-bath treatment on hand osteoarthritis: a randomized clinical trial. *Int J Biometeorol*. Jan 2014;58(1):79-86.
106. Hennig T, Haehre L, Hornburg VT, Mowinckel P, Norli ES, Kjekken I. Effect of home-based hand exercises in women with hand osteoarthritis: a randomised controlled trial. *Ann Rheum Dis*. Aug 2015;74(8):1501-1508.
107. Hermann M, Nilsen T, Eriksen CS, Slatkowsky-Christensen B, Haugen IK, Kjekken I. Effects of a soft prefabricated thumb orthosis in carpometacarpal osteoarthritis. *Scand J Occup Ther*. Jan 2014;21(1):31-39.
108. Horvath K, Kulisch A, Nemeth A, Bender T. Evaluation of the effect of balneotherapy in patients with osteoarthritis of the hands: a randomized controlled single-blind follow-up study. *Clin Rehabil*. May 2012;26(5):431-441.

109. Kovacs C, Pecze M, Tihanyi A, Kovacs L, Balogh S, Bender T. The effect of sulphurous water in patients with osteoarthritis of hand. Double-blind, randomized, controlled follow-up study. *Clinical rheumatology*. Oct 2012;31(10):1437-1442.
110. Osteras N, Hagen KB, Grotle M, Sand-Svartrud AL, Mowinckel P, Kjekken I. Limited effects of exercises in people with hand osteoarthritis: results from a randomized controlled trial. *Osteoarthritis Cartilage*. Sep 2014;22(9):1224-1233.
111. Paolillo AR, Paolillo FR, Joao JP, Joao HA, Bagnato VS. Synergic effects of ultrasound and laser on the pain relief in women with hand osteoarthritis. *Lasers Med Sci*. Jan 2015;30(1):279-286.
112. Watt FE, Kennedy DL, Carlisle KE, et al. Night-time immobilization of the distal interphalangeal joint reduces pain and extension deformity in hand osteoarthritis. *Rheumatology (Oxford)*. Jun 2014;53(6):1142-1149.
113. Vegt AE, Grond R, Gruschke JS, et al. The effect of two different orthoses on pain, hand function, patient satisfaction and preference in patients with thumb carpometacarpal osteoarthritis: a multicentre, crossover, randomised controlled trial. *The bone & joint journal*. Feb 2017;99-b(2):237-244.
114. Bani MA, Arazpour M, Hutchins SW, Layeghi F, Bahramizadeh M, Mardani MA. A custom-made neoprene thumb carpometacarpal orthosis with thermoplastic stabilization: an orthosis that promotes function and improvement in patients with the first carpometacarpal joint osteoarthritis. *Prosthet Orthot Int*. Feb 2014;38(1):79-82.
115. Bani MA, Arazpour M, Kashani RV, Mousavi ME, Maleki M, Hutchins SW. The effect of custom-made splints in patients with the first carpometacarpal joint osteoarthritis. *Prosthet Orthot Int*. Apr 2013;37(2):139-144.
116. Hamann N, Heidemann J, Heinrich K, et al. Stabilization effectiveness and functionality of different thumb orthoses in female patients with first carpometacarpal joint osteoarthritis. *Clinical biomechanics (Bristol, Avon)*. Dec 2014;29(10):1170-1176.
117. Umay E, Tezelli MK, Meshur M, Umay S. The effects of balneotherapy on blood pressure and pulse in osteoarthritis patients with hypertension. *Altern Ther Health Med*. Nov-Dec 2013;19(6):16-21.
118. Grenier ML, Mendonca R, Dalley P. The effectiveness of orthoses in the conservative management of thumb CMC joint osteoarthritis: An analysis of functional pinch strength. *Journal of hand therapy : official journal of the American Society of Hand Therapists*. Jul-Sep 2016;29(3):307-313.
119. Villafane JH, Silva GB, Fernandez-Carnero J. Effect of thumb joint mobilization on pressure pain threshold in elderly patients with thumb carpometacarpal osteoarthritis. *J Manipulative Physiol Ther*. Feb 2012;35(2):110-120.
120. Villafane JH, Silva GB, Chiarotto A. Effects of passive upper extremity joint mobilization on pain sensitivity and function in participants with secondary carpometacarpal osteoarthritis: a case series. *J Manipulative Physiol Ther*. Nov-Dec 2012;35(9):735-742.
121. Figueiredo Neto EM, Queluz TT, Freire BF. Physical activity and its association with quality of life in patients with osteoarthritis. *Rev Bras Reumatol*. Dec 2011;51(6):544-549.
122. Villafane JH, Valdes K. Combined thumb abduction and index finger extension strength: a comparison of older adults with and without thumb carpometacarpal osteoarthritis. *J Manipulative Physiol Ther*. May 2013;36(4):238-244.
123. Medical University of Vienna. Short-wave Diathermy in Patients With Osteoarthritis of the Hand. *ClinicalTrials.gov*. 2015;NCT01838954.
124. Federal University of São Paulo. Effectiveness of Trapeziometacarpal Splint. *ClinicalTrials.gov*. 2015;NCT02635932.

125. Amaral DS. Assistive Device for Hand Osteoarthritis. *ClinicalTrials.gov*. 2016;NCT02667145.
126. Federal University of São Paulo. Progressive Strength in Hand Osteoarthritis. *ClinicalTrials.gov*. 2015;NCT02528630.
127. University of Malaga. Effect of Immobilization of the Metacarpophalangeal Joint in Thumb Osteoarthritis. *ClinicalTrials.gov*. 2016;NCT02780999.
128. Kjeker I, Darre S, Slatkowsky-Cristensen B, et al. Self-management strategies to support performance of daily activities in hand osteoarthritis. *Scand J Occup Ther*. Jan 2013;20(1):29-36.
129. Prieto-Alhambra D, Judge A, Javaid MK, Cooper C, Diez-Perez A, Arden NK. Incidence and risk factors for clinically diagnosed knee, hip and hand osteoarthritis: influences of age, gender and osteoarthritis affecting other joints. *Ann Rheum Dis*. Sep 2014;73(9):1659-1664.

## **Appendices**

**Appendix A: Selection Criteria Summary**

**Appendix B: Search Strategy & Results (Feasibility)**

## Appendix A. Selection Criteria Summary

Selection Criteria	Supporting Data
<b>1. Appropriateness</b>	
1a. Does the nomination represent a health care drug, intervention, device, technology, or health care system/setting available (or soon to be available) in the U.S.?	Yes, this topic represents health care drugs and interventions available in the U.S.
1b. Is the nomination a request for a systematic review?	Yes, this topic is a request for an AHRQ systematic review.
1c. Is the focus on effectiveness or comparative effectiveness?	The focus of this review is on both effectiveness and comparative effectiveness.
1d. Is the nomination focus supported by a logic model or biologic plausibility? Is it consistent or coherent with what is known about the topic?	Yes, it is biologically plausible. Yes, it is consistent with what is known about the topic.
<b>2. Importance</b>	
2a. Represents a significant disease burden; large proportion of the population	Yes, this topic represents a significant burden. OA affects approximately 30.8 million Americans, according to the CDC. <sup>1</sup>
2b. Is of high public interest; affects health care decision making, outcomes, or costs for a large proportion of the US population or for a vulnerable population	Yes, this topic affects health care decisions for a large, vulnerable population. An estimated 33.6% of Americans 65 and older suffer from OA. <sup>1</sup>
2c. Represents important uncertainty for decision makers	Yes, this topic represents important uncertainty for decision makers. Guidelines for this rapidly changing field are from 2012, and are in need of a systematic review to update.
2d. Incorporates issues around both clinical benefits and potential clinical harms	Yes, this nomination addresses both benefits and potential harms of treatments for hip and hand OA.
2e. Represents high costs due to common use, high unit costs, or high associated costs to consumers, to patients, to health care systems, or to payers	Yes, according to the CDC, the average direct cost of OA per patient per year is \$2,600, with total (direct + indirect) costs estimated around \$5,700 per patient per year.
<b>3. Desirability of a New Evidence Review/Duplication</b>	
3. Would not be redundant (i.e., the proposed topic is not already covered by available or soon-to-be available high-quality systematic review by AHRQ or others)	An evidence review on the topic would not be redundant. We identified twelve evidence reviews/meta analyses fitting the scope of the nomination. Eleven reviews were about pharmacologic and non-pharmacological interventions for hip OA. The pharmacologic treatments covered include intraarticular platelet-rich plasma injections and opioid therapy. The non-pharmacological options addressed by the identified evidence reviews include exercise therapy, physical therapy, high-velocity resistance training/low-velocity resistance training, and manual therapy. One evidence review examined physical and occupational therapy for hand OA.
<b>4. Impact of a New Evidence Review</b>	

4a. Is the standard of care unclear (guidelines not available or guidelines inconsistent, indicating an information gap that may be addressed by a new evidence review)?	The standard of care is not unclear. The guidelines are four years old, however they are consistent with what is known about treatments for OA. A new evidence review will provide additional evidence to a previously-lacking evidence base.
4b. Is there practice variation (guideline inconsistent with current practice, indicating a potential implementation gap and not best addressed by a new evidence review)?	Yes, there is practice variation due to limited evidence from previous guidelines.
<b>5. Primary Research</b>	
5. Effectively utilizes existing research and knowledge by considering: - Adequacy (type and volume) of research for conducting a systematic review - Newly available evidence (particularly for updates or new technologies)	We identified 93 unique studies relevant across all key questions published in the past 5 years. These studies included a wide variety of interventions. We found studies examining 18 different pharmacologic treatments for hip OA, 11 non-pharmacological treatments for hip OA, 11 pharmacologic treatments for hand OA, and 10 non-pharmacologic treatments for hand OA. All key questions have identified literature.  Additionally, we identified 23 trials on ClinicalTrials.gov—16 related to hip and 7 related to hand.
<b>6. Value</b>	
6a. The proposed topic exists within a clinical, consumer, or policy-making context that is amenable to evidence-based change	The proposed topic exists with a clinical, consumer, and policy-making context that is amendable to evidence-based change. AHRQ has previously published evidence reviews on therapies for OA.
6b. Identified partner who will use the systematic review to influence practice (such as a guideline or recommendation)	Yes, the ACR and ACP will develop evidence-based guidelines based on the results of an AHRQ evidence review.

*Abbreviations:* ACP=American College of Physicians; ACR=American College of Rheumatology; AHRQ=Agency for Healthcare Research and Quality; CDC=Centers for Disease Control and Prevention; OA=Osteoarthritis



## Appendix B. Search Strategy & Results (Feasibility)

Topic: Osteoarthritis of the Hand or Hip Date: November 15, 2016 Database Searched: MEDLINE (PubMed)	
Concept	Search String
Hip Osteoarthritis	((((osteoarthritis[Title]) AND hip[Title])) OR "Osteoarthritis, Hip/therapy"[Majr]
OR	
Hand Osteoarthritis	(((((hand[Title/Abstract] OR finger[Title/Abstract] OR fingers[Title/Abstract] OR thumb[Title/Abstract] OR wrist[Title/Abstract] OR metacarpus[Title/Abstract]))) OR "Hand"[Mesh])) AND ((osteoarthritis[Title]) OR "Osteoarthritis/therapy"[Mesh:NoExp])
AND	
Therapy	((("Therapeutics"[Mesh] OR "therapy" [Subheading])) OR ((therapy[Title/Abstract] OR drug[Title/Abstract] OR pharmacologic[Title/Abstract] OR non-pharmacologic[Title/Abstract] OR treatment[Title/Abstract] OR management[Title/Abstract] OR care[Title/Abstract]))
NOT	
Not Editorials, etc.	(((((("Letter"[Publication Type]) OR "News"[Publication Type]) OR "Patient Education Handout"[Publication Type]) OR "Comment"[Publication Type]) OR "Editorial"[Publication Type])) OR "Newspaper Article"[Publication Type]
Limit to last 5 years ; human ; English ; Adult	Filters activated: published in the last 5 years, Humans, English, Adult: 19+ years
<b>N= 958</b>	
Systematic Review <b>N=36</b>	PubMed subsection: Systematic [sb]
Randomized Controlled Trials <b>N=352</b>	Cochrane Sensitive Search Strategy for RCT's: (((((((groups[tiab]) OR (trial[tiab]) OR (randomly[tiab]) OR (drug therapy[sh]) OR (placebo[tiab]) OR (randomized[tiab]) OR (controlled clinical trial[pt]) OR (randomized controlled trial[pt])
<b>Other N=570</b>	

**138 studies** found for: hand OR hip | **Recruiting** | osteoarthritis | Adult, Senior |

Studies received from 11/15/2011 to 11/15/2016

[https://clinicaltrials.gov/ct2/results?term=hand+OR+hip&recr=Recruiting&type=&rslt=&age\\_v=&age=1&age=2&gndr=&cond=osteoarthritis&intr=&titles=&outc=&spons=&lead=&id=&state1=&cntry1=&state2=&cntry2=&state3=&cntry3=&locn=&rcv\\_s=11%2F15%2F2011&rcv\\_e=11%2F15%2F2016&lup\\_s=&lup\\_e=](https://clinicaltrials.gov/ct2/results?term=hand+OR+hip&recr=Recruiting&type=&rslt=&age_v=&age=1&age=2&gndr=&cond=osteoarthritis&intr=&titles=&outc=&spons=&lead=&id=&state1=&cntry1=&state2=&cntry2=&state3=&cntry3=&locn=&rcv_s=11%2F15%2F2011&rcv_e=11%2F15%2F2016&lup_s=&lup_e=)

**42 studies** found for: hand OR hip | **Active, not recruiting** | osteoarthritis | Adult,

Senior | Studies received from 11/15/2011 to 11/15/2016

[https://clinicaltrials.gov/ct2/results?term=hand+OR+hip&recr=Active%2C+not+recruiting&type=&rslt=&age\\_v=&age=1&age=2&gndr=&cond=osteoarthritis&intr=&titles=&outc=&spons=&lead=&id=&state1=&cntry1=&state2=&cntry2=&state3=&cntry3=&locn=&rcv\\_s=11%2F15%2F2011&rcv\\_e=11%2F15%2F2016&lup\\_s=&lup\\_e=](https://clinicaltrials.gov/ct2/results?term=hand+OR+hip&recr=Active%2C+not+recruiting&type=&rslt=&age_v=&age=1&age=2&gndr=&cond=osteoarthritis&intr=&titles=&outc=&spons=&lead=&id=&state1=&cntry1=&state2=&cntry2=&state3=&cntry3=&locn=&rcv_s=11%2F15%2F2011&rcv_e=11%2F15%2F2016&lup_s=&lup_e=)

**120 studies** found for: hand OR hip | **Completed** | osteoarthritis | Adult, Senior |

Studies received from 11/15/2011 to 11/15/2016

[https://clinicaltrials.gov/ct2/results?term=hand+OR+hip&recr=Completed&type=&rslt=&age\\_v=&age=1&age=2&gndr=&cond=osteoarthritis&intr=&titles=&outc=&spons=&lead=&id=&state1=&cntry1=&state2=&cntry2=&state3=&cntry3=&locn=&rcv\\_s=11%2F15%2F2011&rcv\\_e=11%2F15%2F2016&lup\\_s=&lup\\_e=](https://clinicaltrials.gov/ct2/results?term=hand+OR+hip&recr=Completed&type=&rslt=&age_v=&age=1&age=2&gndr=&cond=osteoarthritis&intr=&titles=&outc=&spons=&lead=&id=&state1=&cntry1=&state2=&cntry2=&state3=&cntry3=&locn=&rcv_s=11%2F15%2F2011&rcv_e=11%2F15%2F2016&lup_s=&lup_e=)