

Living Systematic Review on Cannabis and Other Plant-Based Treatments for Chronic Pain: 2023 Update— Surveillance Report 1

Literature Update Period: Late April 2023 Through Early July 2023

Summary

The findings reported in the systematic review are current. One new observational study meeting the inclusion criteria was identified during the update search of the literature published between late April 2023 and early July 2023.

Overview

This is the first surveillance report since the 2023 annual update of a living systematic review on cannabis and other plant-based treatments for chronic and subacute pain.

The systematic review synthesizes evidence on the benefits and harms of plant-based compounds (PBCs), such as cannabinoids and kratom, used to treat chronic or subacute pain, and addresses concerns about severe adverse effects, abuse, misuse, dependence, and addiction.

The purpose of this surveillance report is to describe new studies identified since the last search (mid-April 2023) and provide a synthesis of the accumulated evidence. Surveillance update reports are planned on a quarterly basis, and the systematic review will be updated annually. The systematic review is available on the Agency for Healthcare Research and Quality (AHRQ) website (<https://effectivehealthcare.ahrq.gov/products/plant-based-chronic-pain-treatment/living-review>). Table 1 provides a summary of the version history.

Table 1. Version history

Search End Date	Report (Publication Date)
July 2021	Systematic Review (Oct. 27, 2021)
August 2021	Surveillance Report 1 (Oct. 27, 2021)
October 2021	Surveillance Report 2 (Jan. 28, 2022)
Mid-January 2022	Surveillance Report 3 (May 2022)
March 2022	Surveillance Report 4 (Aug. 2022)
April 2022	Systematic Review (Aug. 2022)
Early July 2022	Surveillance Report 1 (Sept. 2022)
Mid-October 2022	Surveillance Report 2 (Jan. 2023)
Late January 2023	Surveillance Report 3 (Apr. 2023)
Mid-April 2023	Surveillance Report 4 (July 2023)
Late April 2023	Systematic Review (Aug. 2023)
Early July 2023	Surveillance Report 1 (Sept. 2023)

Main Points

The Key Questions (KQs) for this review focus on the benefits (KQ1) and harms (KQ2) of cannabinoids for treating chronic or subacute pain, as well as the benefits (KQ3) and harms (KQ4) of other PBCs, such as kratom, for treating chronic or subacute pain. Studies of cannabis-related products were grouped based on their tetrahydrocannabinol (THC) to cannabidiol (CBD) ratio using the following categories: high THC to CBD ratio, comparable THC to CBD ratio, and low THC to CBD ratio. One new prospective cohort study comparing various medicinal low THC to CBD ratio products (CBD only or THC to CBD ratio $\leq 1:2$), comparable THC to CBD ratio products, and high THC to CBD (THC only) products in adults with chronic pain was identified for inclusion during this surveillance period.¹ The evidence for low THC to CBD ratio products versus placebo remained insufficient, based on single studies evaluating heterogeneous products. Evidence comparing different cannabis-related products also remained insufficient.

Overall, in patients with chronic (mainly neuropathic) pain with short-term treatment (4 weeks to <6 months):

- Comparable THC to CBD ratio oral spray is probably associated with small improvements in pain severity and overall function versus placebo. There was no increase in risk of serious adverse events or withdrawal due to adverse events. There may be a large increased risk of dizziness and sedation, and a moderate increased risk of nausea.
- Synthetic THC (high THC to CBD ratio) may be associated with small improvement in pain severity but with increased risk of sedation and potential increased risk of nausea versus placebo. Synthetic THC is probably associated with a large increased risk of dizziness.
- Extracted whole-plant high THC to CBD ratio products may be associated with large increases in risk of withdrawal due to adverse events and dizziness versus placebo; outcomes assessing benefit were not reported or insufficient.
- Evidence on whole-plant cannabis (including patient's choice of products), low THC to CBD ratio products (topical, oral, or sublingual CBD), and other cannabinoids (cannabidivarin), and comparisons with other active interventions or different cannabis-related products was insufficient to draw conclusions.
- Other key adverse event outcomes (psychosis, cannabis use disorder, cognitive deficits) and outcomes on the impact on opioid use were not reported or evidence was insufficient to draw conclusions.
- No evidence on other plant-based compounds, such as kratom, met criteria for this review.

Table 2 presents the conclusions from the systematic review, findings from ongoing literature surveillance, and an assessment of the new study on conclusions.

Table 2. Assessment of systematic review conclusions

Key Question^a	Conclusions From Systematic Review (2023)	Findings From Surveillance to Date	Assessment
KQ1 and KQ2. Comparable THC to CBD Ratio Benefits and Harms	Benefits: small improvements in pain severity (7 RCTs) and in function (6 RCTs) (SOE: moderate) Harms: no effect on serious adverse events (SOE: low; 3 RCTs); large increased risk of dizziness and sedation; moderate increased risk of nausea (SOE: low; 6 RCTs)	No new studies	No change in conclusions
KQ1 and KQ2. Synthetic High THC to CBD Ratio Benefits and Harms	Benefits: small improvements in pain severity (SOE: low; 7 RCTs); no effect on overall function/disability (SOE: low; 4 RCTs) Harms: moderate increased risk of sedation (SOE: low; 4 RCTs); large increased risk of nausea (SOE: low; 3 RCTs); and moderate increased risk of dizziness (SOE: moderate; 3 RCTs)	No new studies	No change in conclusions
KQ1 and KQ2. Extracted Whole-Plant High THC to CBD Ratio Benefits and Harms	Benefits: insufficient evidence (2 RCTs) Harms: large increase in risk of dizziness and in study withdrawal due to adverse events (SOE: low; 1 RCT)	No new studies	No change in conclusions
KQ1 and KQ2. Low THC to CBD Ratio Benefits and Harms	Insufficient evidence (3 RCTs ^b)	No new studies	No change in conclusions
KQ1 and KQ2. Whole-Plant Cannabis and Other Cannabinoids Benefits and Harms	Insufficient evidence (2 RCTs)	No new studies	No change in conclusions
KQ1 and KQ2. Comparable THC to CBD Ratio Vs. Synthetic THC Benefits and Harms	Insufficient evidence (1 observational study)	No new studies	No change in conclusions
KQ1 and KQ2. Comparable THC to CBD Ratio Vs. LAOs	Insufficient evidence (1 observational study)	No new studies	Insufficient evidence
KQ1 and KQ2. Whole-Plant High THC to CBD Ratio Flower Vs. Extracted Oils	Insufficient evidence (1 observational study)	No new studies	Insufficient evidence
KQ1 and KQ2. Low Vs. Comparable Vs. High THC to CBD Ratio Products	No studies	1 observational study	Insufficient evidence
KQ3 and KQ4. Kratom or Other Plant-Based Substances Benefits and Harms	Insufficient evidence (0 RCTs)	No new studies	No change in conclusions

Abbreviations: CBD = cannabidiol; KQ = Key Question; LAO = long-acting opioid; RCT = randomized controlled trial; SOE = strength of evidence; THC = tetrahydrocannabinol.

^a For Key Question wording, see the Background section below.

^b Products of varied origin (synthetic or plant derived) and route (oral or topical), resulting in heterogeneity in products and imprecision for specific low THC to CBD ratio product types.

Summary of Findings Tables

The KQs for this review focus on the benefits (KQ1) and harms (KQ2) of cannabinoids for treating chronic pain, as well as the benefits (KQ3) and harms (KQ4) of other PBCs, such as kratom, for treating chronic pain. Tables 3 and 4 summarize benefits and harms of cannabinoids, based on evidence reviewed to date. No evidence was available for other PBCs.

Table 3. Key Question 1: Benefits of cannabinoids for chronic pain compared with placebo in the short term (4 weeks to <6 months)

Product, THC to CBD Ratio Product	Pain Response ^a Effect Size (N Studies) [SOE]	Pain Severity Effect Size (N Studies) [SOE]	Function Effect Size (N Studies) [SOE]
Comparable THC/CBD - Extracted From Whole Plant, Oromucosal Spray	Potential effect (4) ^b [✓]	Small effect (7) [✓✓]	Small effect (6) [✓✓]
High THC – Synthetic, Oral	Insufficient (2)	Small effect (7) [✓]	No effect (4) [✓]
High THC – Extracted From Whole Plant, Oral	No evidence	Insufficient (2)	Insufficient (1)
Low THC – Topical CBD, Extracted From Whole Plant	No evidence	Insufficient (1)	No evidence
Low THC – Oral CBD, Synthetic	No evidence	Insufficient (1)	Insufficient (1)
Low THC – Oral CBD or CBD/THC, Unclear Origin	Insufficient (1)	Insufficient (1)	Insufficient (1)
Low THC – Sublingual CBD/THC, Extracted from Whole Plant	No evidence	Insufficient (1)	No evidence
Other Cannabinoids – CBDV, Oral	Insufficient (1)	Insufficient (1)	No evidence
Whole-Plant Cannabis (12% THC) ^c	No evidence	Insufficient (1)	No evidence

Abbreviations: CBD = cannabidiol; CBDV = cannabidivarin; SOE = strength of evidence; THC = tetrahydrocannabinol.

^a ≥30% improvement from baseline.

^b Potential effect: SOE of low or higher; findings indicate at least a small magnitude of effect but not statistically significant.

^c Comparison was “usual care.”

Effect size: None (i.e., no effect/no statistically significant effect), small, moderate, or large increased benefit; SOE: [✓] = low, [✓✓] = moderate, [✓✓✓] = high.

Table 4. Key Question 2: Harms of cannabinoids for chronic pain compared with placebo in the short term (4 weeks to <6 months)

Product/THC to CBD Ratio	WAE Effect Size (N Studies) [SOE]	SAE Effect Size (N Studies) [SOE]	Dizziness Effect Size (N Studies) [SOE]	Nausea Effect Size (N Studies) [SOE]	Sedation Effect Size (N Studies) [SOE]
Comparable THC/CBD – Extracted From Whole Plant, Oromucosal Spray	No effect (5) [✓]	No effect (3) [✓]	Large effect (6) [✓]	Moderate effect (6) [✓]	Large effect (6) [✓]
High THC – Synthetic, Oral	Potential effect ^a (5) [✓]	Insufficient (1)	Large effect (3) [✓✓]	Potential effect ^a (3) [✓]	Moderate effect (4) [✓]

Product/THC to CBD Ratio	WAE Effect Size (N Studies) [SOE]	SAE Effect Size (N Studies) [SOE]	Dizziness Effect Size (N Studies) [SOE]	Nausea Effect Size (N Studies) [SOE]	Sedation Effect Size (N Studies) [SOE]
High THC – Extracted From Whole Plant, Oral	Large effect (1) [✓]	Insufficient (1)	Large effect (1) [✓]	No evidence	No evidence
Low THC – Topical CBD, Extracted From Whole Plant	No evidence	No evidence	No evidence	No evidence	No evidence
Low THC – Oral CBD, Synthetic	Insufficient (1)	Insufficient (1)	No evidence	No evidence	No evidence
Low THC – Oral CBD or CBD/THC, Unclear Origin	Insufficient (1)	Insufficient (1)	Insufficient (1)	Insufficient (1)	Insufficient (1)
Low THC – Sublingual CBD/THC, Extracted From Whole Plant	Insufficient (1)	Insufficient (1)	No evidence	No evidence	No evidence
Other Cannabinoids – CBDV, Oral	Insufficient (1)	Insufficient (1)	No evidence	No evidence	No evidence
Whole-Plant Cannabis (12% THC) ^b	Insufficient (1)	Insufficient (1)	Insufficient (1)	Insufficient (1)	Insufficient (1)

Abbreviations: CBD = cannabidiol; CBDV = cannabidivarin; SAE = serious adverse event; SOE = strength of evidence; THC = tetrahydrocannabinol; WAE = withdrawal due to adverse event.

^a Potential effect: SOE of low or higher; findings indicate at least a small magnitude of effect but not statistically significant.

^b Comparison was “usual care.”

Effect size: None (i.e., no effect/no statistically significant effect), small, moderate, or large increased risk; SOE: [✓] = low, [✓✓] = moderate, [✓✓✓] = high.

Background

Chronic pain is defined as pain lasting longer than 3 to 6 months or past normal time for tissue healing,^{2,3} and it affects approximately 100 million people in the United States.⁴ Chronic pain adversely affects physical and mental functioning, productivity, and quality of life, and is often refractory to treatment and associated with substantial costs.⁵⁻⁷

While opioids are often prescribed for chronic pain, a recent series of systematic reviews found that opioids,⁸ several nonopioid drugs,⁹ and some nonpharmacologic treatments¹⁰ have small to moderate effects on pain and function, but also frequent adverse effects and some less frequent but serious ones. The 2012 Centers for Disease Control and Prevention *Guideline for Prescribing Opioids for Chronic Pain* recommends that nonopioid therapy is preferred for treatment of chronic pain.¹¹ The limited efficacy of opioids and the ongoing opioid crisis drive a search for alternative pain treatments, including PBCs such as cannabis and related compounds, as some data suggest they may have analgesic properties.¹²

The term *cannabinoid* refers to a group of closely related compounds that are active in cannabis, with the two main cannabinoid compounds being THC and CBD. THC has demonstrated analgesic properties,^{13,14} although its psychoactive effects and abuse potential may limit its suitability as an analgesic. It may also be associated with serious harms, including those related to potential for use disorder or physiological withdrawal. CBD may also have some analgesic or anti-inflammatory properties and is thought to be less intoxicating and not addictive.^{15,16} While not derived from plants, two synthetic cannabinoid products, dronabinol (a synthetic THC) and nabilone (a THC analog), have also been studied for treating chronic pain.

Other PBCs with effects similar to opioids or cannabis, such as kratom, have been considered to treat chronic pain. These may also have serious harms, including dependence, addiction, and physiological withdrawal potential.¹⁷

Although the original review and prior surveillance reports and update focused on chronic pain in adults, subacute pain and adolescents are also relevant. Subacute pain, often defined as pain lasting for 4 to 12 weeks, represents a transitional state between acute (<4 weeks) pain, which often resolves, and chronic pain, which is more likely to persist.¹⁸ Effective treatments for reducing the likelihood of subacute pain becoming chronic are also needed. Adolescents also experience chronic pain and have a high prevalence of cannabis use (recreational or medical^{19,20}).

Four KQs guide the review:

KQ1: In adults or adolescents with chronic or subacute pain, what are the benefits of cannabinoids for treatment of chronic or subacute pain?

KQ2: In adults or adolescents with chronic or subacute pain, what are the harms of cannabinoids for treatment of chronic or subacute pain?

KQ3: In adults or adolescents with chronic or subacute pain, what are the benefits of kratom or other plant-based substances for treatment of chronic or subacute pain?

KQ4: In adults or adolescents with chronic or subacute pain, what are the harms of kratom or other plant-based substances for treatment of chronic or subacute pain?

The protocol for the systematic review can be found on the AHRQ website (<https://effectivehealthcare.ahrq.gov/products/plant-based-chronic-pain-treatment/protocol>) and on the PROSPERO systematic reviews registry (registration number CRD42021229579). The scope of the review was reviewed with a Technical Expert Panel (TEP) following the prior annual update, including considerations for expansion of scope.²¹ With TEP input, the protocol was amended to include adolescents and subacute pain. An updated protocol was submitted to PROSPERO,²² and the title, Key Questions, and inclusion and exclusion criteria were revised to reflect the changes.

Methods

In brief, we searched Ovid[®] MEDLINE[®], PsycINFO[®], Embase[®], the Cochrane Library, and SCOPUS[®] databases monthly through early July 2023 for studies of adults or adolescents with chronic or subacute pain with at least 4 weeks of treatment or followup. Studies of cannabis, kratom, and similar PBCs compared with a placebo, no treatment, each other, or another treatment were eligible. For the period covered by this surveillance report (late April 2023 to early July 2023), one new observational study that compared various low-THC to CBD ratio (CBD only or THC to CBD ratio $\leq 1:2$), comparable THC to CBD ratio, and high-THC to CBD ratio products in patients with chronic pain (various causes) was identified.¹ Pain is the primary outcome for this review; details on the search strategies are in [Appendix A](#). Briefly, we included randomized controlled trials (RCTs) and observational studies with a concurrent control group with a minimum of 4 weeks' followup assessing cannabis and other plant-based interventions in

adults or adolescents with noncancer chronic or subacute pain. The full inclusion and exclusion criteria for all primary and secondary outcomes for this report are in [Appendix B](#).

We followed the methods guidance in the AHRQ Methods Guide,²³ and abstracted key information and conducted risk-of-bias assessments using the Cochrane Back Pain Group’s version of the Cochrane guidance for randomized trials²⁴ and criteria developed by the U.S. Preventive Services Task Force²⁵ for observational studies for each included study. Our methods included categorizing studies based on the duration of followup as short-, intermediate-, and long-term. Studies that assessed the cannabinoids THC and/or CBD were grouped based on their THC to CBD ratios and categorized as high THC to CBD ratio, comparable THC to CBD ratio, and low THC to CBD ratio (Table 5). We also grouped studies by whether the product was a whole-plant product (cannabis), cannabinoids extracted or purified from a whole plant, or synthetic. When studies were similar enough to provide a meaningful combined estimate, we conducted meta-analyses using the profile likelihood random effects model and assessed between-study heterogeneity using Cochran’s Q statistic chi square and the I² test for inconsistency. Magnitude of benefit was categorized into no effect or small, moderate, and large effects. (See [Appendix B](#), Table B-2.)

Table 5. Organizing principle of cannabis-related studies based on ratios of THC to CBD

Intervention Category (Definition)	Source	Possible Derivatives	Example Products	U.S. Availability
High THC (THC to CBD ratio equals $\geq 2:1$ ratio)	Synthetic	Synthetic THC (100% THC or analog)	Dronabinol (Marinol®) or nabilone (Cesamet®)	Available via prescription ^a
	Synthetic	Purified from whole-plant with close to 100% THC	Purified dronabinol (Namisol®) ^{b,c}	Not available in the U.S.
	Plant-based	Commercially marketed product extracted from whole-plant with known high ratio of THC/CBD	THC/CBD extracts with high THC/CBD ratio	Unknown – may be available at dispensaries where allowed
	Plant-based	Whole-plant with known high concentration of THC	Whole-plant cannabis with known high THC concentration	Unknown – may be available at dispensaries where allowed
Comparable THC to CBD (THC to CBD ratio is $< 2:1$ and $> 1:2$)	Plant-based	Extracted from whole-plant with comparable ratio of THC/CBD	Nabiximols (Sativex®) ^d	Not available in the U.S.
	Plant-based	Extracted from whole-plant with comparable ratio of THC/CBD	Oral tinctures with similar ratio of THC/CBD	Unknown – may be available at dispensaries where allowed
	Plant-based	Whole-plant with known comparable ratio of THC/CBD	Whole-plant with known comparable ratio of THC/CBD	Unknown – may be available at dispensaries where allowed
Low THC (THC to CBD ratio equals $\leq 1:2$)	Plant-based	Extracted from whole plant with low ratio of THC/CBD	CBD topical or oral	Unknown – may be available at dispensaries where allowed
Low THC (THC to CBD ratio is $\leq 1:2$)	Synthetic	Synthetic CBD	CBD oral tablets	Unknown

Intervention Category (Definition)	Source	Possible Derivatives	Example Products	U.S. Availability
Whole-Plant Cannabis Products (THC to CBD ratio categorized based on information provided [potentially unknown])	Plant-based	Whole-plant products	Cannabis flowers, resins, buds, leaves, hashish	Unknown – may be available at dispensaries where allowed.
Other Cannabinoids (Cannabinoids other than THC or CBD)	Plant-based	Extracted from whole-plant	Cannabidivarin (CBDV) extracted oil (oral)	Unknown – may be available at dispensaries where allowed

Abbreviations: CBD = cannabidiol; FDA = Food and Drug Administration; THC = tetrahydrocannabinol.

^aThese products are approved by the FDA for non-pain indications (anorexia related to HIV infection, nausea related to chemotherapy).

^bNamisol[®] is chemically identical to dronabinol, and is therefore grouped together with synthetic dronabinol.

^cManufactured in The Netherlands; may be available in some European countries. Not currently FDA-approved.

^dManufactured and available in Canada and some European countries; not FDA-approved.

A more detailed discussion of methods can be found in the [protocol](#) and in [Appendix B](#).

Results to Date

Results Overview

Across all of the monthly literature searches to date, 5,314 citations were screened, from which we included 34 studies.^{1,26-58} For the period covered by this surveillance report, 135 citations were screened

One new observational study (n=296 for primary analysis) that compared various cannabis products met inclusion criteria for this update period. [Appendix C](#) contains a list of included studies, and a literature flow diagram can be found in [Appendix D](#). [Appendix E](#) contains summary tables of individual study data for all included studies and the results of synthesis (i.e., forest plots). [Appendix F](#) contains detailed evidence tables of included studies, and [Appendix G](#) contains risk-of-bias assessments. [Appendix H](#) contains details on strength-of-evidence ratings. A list of studies excluded after reviewing the full manuscripts can be found in [Appendix I](#) along with reasons for their exclusion. [Appendix J](#) provides a funnel plot of high THC ratio studies included in the meta-analysis for pain severity.

Table 6 summarizes the characteristics of included RCTs, and Table 7 summarizes the characteristics of included observational studies.

Table 6. Characteristics of included randomized controlled trials to date

Characteristic	THC/CBD	THC	Synthetic THC	CBD	CBDV
THC to CBD Ratio	Comparable ^a	High	High	Low	NA - other cannabinoids
Source	Plant-extracted	Plant-extracted	Synthetic Nabilone Dronabinol Dronabinol/Namisol ^{®b}	Plant-extracted (2) ^c Synthetic (1) Unclear (1)	Plant-extracted
N Studies	7	2	10	4 ^d (1 topical, 1 sublingual, 2 oral)	1

Characteristic	THC/CBD	THC	Synthetic THC	CBD	CBDV
Comparator (Study Count)	Placebo (7)	Placebo (2)	Placebo (7); Ibuprofen (1); Diphenhydramine (1); Dihydrocodeine (1); Low-THC to CBD ratio (CBD or Dronabinol/CBD ^e) (1)	Placebo (4); Dronabinol (1); Dronabinol/CBD (1); ≥97% purified low-THC to CBD (1:6) sublingual oil (1)	Placebo
Route of Administration, Formulation (Study Count)	Sublingual oromucosal spray, 2.7 mg THC/2.5 mg CBD per 100 mcl	Sublingual oil drops, 24 mg/ml THC/0.51 mg/ml CBD (1) Oral capsule, 2.5 mg THC/0.8 – 1.8 mg CBD extract (1)	Nabilone oral 0.25 mg capsule (1); Nabilone oral 0.5 mg capsule (5); Dronabinol 2.5 mg oral capsule (2); Dronabinol 5 mg oral capsule (1); Namisol ^{®a} 3 mg oral tablet (1)	Topical oil, 83 mg CBD/fluid ounce (1), Oral tablet, 10 mg CBD (1) Oral capsule, 5 mg CBD (1) Oral capsule, 5 mg CBD/2.5 mg dronabinol (1) Sublingual oil, 24.5 mg/mL THC, 147 mg/mL CBD (1)	Oral oil, 50 mg/ml CBDV
Dosing Regimen	Final mean dose 23 mg THC/21 mg CBD daily.	Sublingual drops: 1.2 mg daily, titrated. Final dose 4.4 mg THC daily. Capsule: 2.5 - 12.5 mg THC twice daily, titrated. Final dose NR Oral oil: 1.2 mg daily	Nabilone 0.25 - 2 mg twice daily, titrated. Final mean dose 1.84 Dronabinol capsules: 2.5 -15 mg once or twice daily, titrated. Final dose range 15 - 25 mg/day Namisol ^{®a} tablet: 3 - 8 mg 3 times daily, titrated. Final dose NR.	Topical oil: applied locally 1-4 times/day (volume/dose, final dose NR). Oral tablet: 10 mg daily, titrated (max 3 times daily). Final dose NR. Oral CBD capsule: 5 mg twice daily, titrated. Final median dose 50 mg CBD daily. Oral dronabinol/CBD capsule: 2.5 mg THC/5 mg CBD twice daily, titrated. Final median dose 15 mg THC/30 mg CBD daily. Sublingual oil, titrated to max daily dose of 6 drops 3 times daily (15 mg THC/90 mg CBD)	400 mg CBDV daily. Final dose NR.
Risk of Bias	29% high, 57% moderate, 14% low	50% moderate, 50% low	20% high, 40% moderate, 40% low	50% high, 25% moderate, 25% low	100% moderate

Characteristic	THC/CBD	THC	Synthetic THC	CBD	CBDV
Total Randomized	882	297	592	267	34
Age, Mean Years	53	52	53	65	50
Female, %	66%	89%	61%	40%	3%
Non-White,^e %	1.6% (2)	1% (1)	5.4% (3)	NR	NR
Primary Pain Type (Study Count)	NPP (6); Inflammatory arthritis (1)	NPP (1); Fibromyalgia (1)	NPP (7); Fibromyalgia (1); Headache (1); Visceral pain (1)	NPP (2); OA (1); Unspecified (1)	NPP (1)
Baseline Pain Score, Mean (Range)^f	6.59 (5.3 to 7.3)	8.47 (8.25 to 8.67)	6.48 (4 to 8.1) ^g	5.87 (4.67 to 7.4) ^h	6.28 (6.12 to 6.44)
Study Duration	4 to 15 weeks	8 to 12 weeks	4 to 47 weeks	4 to 16 weeks	4 weeks

Abbreviations: CBD = cannabidiol; CBDV = cannabidivarin; NA = not applicable; NPP = neuropathic pain; NR = not reported; OA = osteoarthritis; RCT = randomized controlled trial; THC = tetrahydrocannabinol.

^a All products were nabixiomols.

^b Namisol[®] is a purified, plant-based product, but grouped with synthetic dronabinol because they are chemically identical.

^c One trial evaluated a plant-extracted low THC to CBD product and a product that underwent further purification.

^d One trial is included in both the synthetic THC and CBD columns, as it compared THC to CBD, CBD/THC, and placebo.

^e (n) = number of studies reporting this characteristic at baseline.

^f Scores were standardized to a 0 to 10 scale.

^g Weighted mean includes median scores for 1 study (6 vs. 6).

^h Weighted mean includes median scores for 1 study (5.2 vs. 6.1).

Table 7. Characteristics of included observational studies to date

Characteristic	THC/CBD ^a	THC	Synthetic THC	THC/CBD Versus Synthetic THC	THC/CBD Versus LAOs
THC to CBD Ratio	Unclear or mixed	High	High	Comparable vs. high	Comparable
Source	Any cannabis product (patient's choice) or mixed cannabinoids	Plant-based	Synthetic (nabilone)	Plant-based vs. synthetic	Plant-based
N Studies	5 (unclear THC to CBD ratio); 2 (mixed THC to CBD ratios) ^b	1	1	1	1
Comparator (Study Count)	No cannabis use (3); usual care (1); no medical cannabis authorization (1); other cannabinoids (2)	Usual care (1)	Gabapentin only; gabapentin + nabilone (1)	Active comparator; oral mucosal spray vs. dronabinol	Long-acting opioids (MME 69.4 [SD 38.9] mg/day)
Route of Administration, Formulation	Unreported (any available allowed, patient's choice) (5); inhaled, sublingual, or oral prescribed cannabis (2)	Whole-plant cannabis, "certified 12.5% THC" (CBD NR) route determined by patient: smoking 27%, oral 8%, vaporization 4%, combination 61%	Nabilone 0.5 mg oral capsule	Nabiximols sublingual oromucosal spray, 2.7 mg THC/2.5 mg CBD per 100 mcl Dronabinol oral capsule (strength NR)	Nabiximols sublingual oromucosal spray, 2.7 mg THC/2.5 mg CBD per 100 mcl Oral long-acting opioids (dose varied)

Characteristic	THC/CBD ^a	THC	Synthetic THC	THC/CBD Versus Synthetic THC	THC/CBD Versus LAOs
Dosing Regimen	None specified. Final dose NR	None specified; titrated to max dose 5 g/day. Final median dose 2.5 g/day	None specified; final mean dose 3 mg/day	None specified; final mean dose 16.6/15.4 mg THC/CBD/day vs. 17.2 mg THC/day	None specified; based on individual patient needs; final mean dose 16.7/15.5 mg THC/CBD/day vs. MME 69.4 mg/day
ROB	57% high, 43% moderate	100% high	100% moderate	100% moderate	100% moderate
N Total	13,565	431	156	674	1,310
Age, Mean Years	53	49	61	46	51
Female, %	56%	57%	59%	57%	57%
% Non-White (Study Count)	54% (1); NR (6)	NR	NR	NR	NR
Primary Pain Type(s)	Mixed musculoskeletal, chronic non-cancer pain	Chronic non-cancer pain	NPP	Peripheral NPP	Peripheral neuropathic back pain
Baseline Pain Score, Mean (Range)^c	5.75 (4.56 to 7.00)	6.35 (6.1 to 6.6)	4.98 (4.58 to 5.31)	4.4 (4.39 to 4.41)	4.32 (4.31 to 4.33)
Study Duration, Weeks (Range)	12 to 208	52	26	24	24

Abbreviations: CBD = cannabidiol; LAO = long-acting opioid; MME = morphine milligram equivalents; NPP = neuropathic pain; NR = not reported; ROB = risk of bias; SD = standard deviation; THC = tetrahydrocannabinol.

^a Patients could choose any medicinal product they preferred in these studies.

^b Includes one new study for this review.

^c Scores were standardized to a 0 to 10 scale.

KQs 1 and 2: Benefits and Harms of Cannabis

One new high risk of bias prospective cohort study (n=296 for primary analysis) by Schubert et al. evaluated patients with chronic pain (mixed conditions).¹ It compared various medicinal low THC to CBD ratio (CBD only or THC to CBD ratio $\leq 1:2$), comparable THC to CBD ratio, and high THC to CBD (THC only) products. Method of administration appeared to be oral; product types (synthetic or plant-based) were not specified, although cannabis flower was described as one type of product. The mean age was 56, and 62 percent were female. Race was not reported. Methodological limitations included unclear patient selection methods and failure to control for confounders.

The study found low, comparable, and high THC to CBD ratio products associated with very similar outcomes on the Patient-Reported Outcomes Measurement Information System-29 (PROMIS-29) pain interference, pain intensity, pain impact, and physical function domains at 13 weeks median follow-up. Outcomes were also similar across cannabis product categories on the PROMIS-29 anxiety and depression domains and for rates of somnolence, anxiety, nausea, vomiting, and dizziness. With the addition of this one new, high risk of bias observational study, the strength of evidence for comparing different cannabis products for pain, function, harms, or other outcomes remained insufficient.

Conclusion

One new prospective cohort study with methodological limitations comparing different cannabis products was identified for this surveillance report. It found different cannabis product categories (low, comparable, or high THC to CBD ratio) associated with similar effects on pain, function, and harms. Due to methodological limitations in the new study, the strength of evidence for comparisons of different cannabis products remained insufficient.

Overall, including previously reviewed evidence, this surveillance report found that evidence on cannabis-related interventions remains restricted to short-term outcomes, primarily in patients with neuropathic pain. Improvement in pain appears small with high and comparable THC to CBD ratio products. Compared with placebo, cannabis-related interventions are associated with greater risk of common adverse events (dizziness, nausea, sedation) and study withdrawal due to adverse events. No studies evaluated adolescents or persons with subacute pain, and evidence for other interventions, including kratom, was insufficient or not found. Additional studies are needed to improve confidence in these findings and to provide evidence on longer term followup, other outcomes, and other interventions, including whole-plant cannabis.

Next Report

The next surveillance report is scheduled for winter 2023.

References

1. Schubert EA, Johnstone MT, Benson MJ, et al. Medicinal cannabis for Australian patients with chronic refractory pain including arthritis. *Br J Pain*. 2023;17(2):206-17. doi:10.1177/20494637221147115. PMID: 37057257.
2. Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain— United States, 2016. *JAMA*. 2016 Apr 19;315(15):1624-45. doi: 10.1001/jama.2016.1464. PMID: 26977696.
3. Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain - United States, 2016. *MMWR Recomm Rep*. 2016 Mar 18;65(1):1-49. doi: 10.15585/mmwr.r6501e1. PMID: 26987082.
4. Dahlhamer J, Lucas J, Zelaya C, et al. Prevalence of chronic pain and high-impact chronic pain among adults — United States, 2016. . *MMWR Morb Mortal Wkly Rep*. 2018;67(36):1001-6. doi: 10.15585/mmwr.mm6736a2. PMID: 30212442.
5. Institute of Medicine (US) Committee on Advancing Pain Research C, and Education. *Relieving pain in America: a blueprint for transforming prevention, care, education, and research*. Washington, DC: National Academies Press; National Academy of Sciences; 2011.
6. Ballantyne JC, Shin NS. Efficacy of opioids for chronic pain: a review of the evidence. *The Clinical Journal of Pain*. 2008 Jul-Aug;24(6):469-78. doi: 10.1097/AJP.0b013e31816b2f26. PMID: 18574357.
7. Eriksen J, Sjogren P, Bruera E, et al. Critical issues on opioids in chronic non-cancer pain: an epidemiological study. *Pain*. 2006 Nov;125(1):172-9. doi: 10.1016/j.pain.2006.06.009. PMID: 16842922.
8. Chou R, Hartung D, Turner J, et al. Opioid treatments for chronic pain. Rockville, MD: Agency for Healthcare Research and Quality (US); 2020 Apr. Report No.: 20-EHC011. PMID: 32338848.
9. McDonagh MS, Selph SS, Buckley DI, et al. *Nonopioid pharmacologic treatments for chronic pain*. Rockville, MD: Agency for Healthcare Research and Quality; 2020. doi: 10.23970/AHRQEPCCER228. PMID: 32338847.
10. Skelly AC, Chou R, Dettori JR, et al. *Noninvasive nonpharmacological treatment for chronic pain: a systematic review update*. Rockville, MD: Agency for Healthcare Research and Quality; 2020. doi: 10.23970/AHRQEPCCER227. PMID: 32338846.
11. Dowell D, Ragan KR, Jones CM, et al. CDC guideline for prescribing opioids for chronic pain— United States, 2022. *MMWR Recomm Rep* 2022. 2022;71((No. RR-3)):1-95. doi: 10.15585/mmwr.rr7103a1.
12. Stockings E, Campbell G, Hall WD, et al. Cannabis and Cannabinoids for the treatment of people with chronic noncancer pain conditions: a systematic review and meta-analysis of controlled and observational studies. *Pain*. 2018 Oct;159(10):1932-54. doi: 10.1097/j.pain.0000000000001293. PMID: 29847469.
13. Elikkottil J, Gupta P, Gupta K. The analgesic potential of cannabinoids. *J Opioid Manag*. 2009 Nov-Dec;5(6):341-57. doi: 10.5055/jom.2009.0034. PMID: 20073408.
14. Whiting PF, Wolff RF, Deshpande S, et al. Cannabinoids for medical use: a systematic review and meta-analysis. *Jama*. 2015 Jun 23-30;313(24):2456-73. doi: 10.1001/jama.2015.6358. PMID: 26103030.
15. Vučković S, Srebro D, Vujović KS, et al. Cannabinoids and pain: new insights from old molecules. *Front Pharmacol*. 2018 Nov 13;9:1259. doi: 10.3389/fphar.2018.01259. PMID: 30542280.
16. Morales P, Hurst DP, Reggio PH. Molecular targets of the phytocannabinoids: a complex picture. *Phytocannabinoids*. 2017:103-31.
17. White CM. Pharmacologic and clinical assessment of Kratom: an update. *Am J Health-Syst Pharm*. 2019 Nov 13;76(23):1915-25. doi: 10.1093/ajhp/zxz221. PMID: 31626272.

18. Marin TJ, Van Eerd D, Irvin E, et al. Multidisciplinary biopsychosocial rehabilitation for subacute low back pain. *Cochrane Database Syst Rev*. 2017 Jun 28;6(6):Cd002193. doi: 10.1002/14651858.CD002193.pub2. PMID: 28656659.
19. Feldman DE, Nahin RL. National Estimates of Chronic Musculoskeletal Pain and Its Treatment in Children, Adolescents, and Young Adults in the United States: Data From the 2007-2015 National Ambulatory Medical Care Survey. *J Pediatr*. 2021 Jun;233:212-9.e1. doi: 10.1016/j.jpeds.2021.01.055. PMID: 33524388.
20. Zuckermann AME, Battista KV, Bélanger RE, et al. Trends in youth cannabis use across cannabis legalization: Data from the COMPASS prospective cohort study. *Prev Med Rep*. 2021 Jun;22:101351. doi: 10.1016/j.pmedr.2021.101351. PMID: 33816088.
21. Chou R, Ahmed AY, Iyer S, et al. Living Systematic Reviews: Practical Considerations for the Agency for Healthcare Research and Quality Evidence-based Practice Center Program from Year 2 of an Ongoing Review [White Paper in Progress]. *Pacific Northwest Evidence-based Practice Center*; 2023.
22. Chou R, Ahmed AY, Morasco BJ, et al. Living Systematic Review on Cannabis and Other Plant-based Treatments for Chronic and Subacute Pain. PROSPERO 2021 CRD42021229579 Available from: . https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021229579.
23. *Methods guide for effectiveness and comparative effectiveness reviews*. Rockville, MD: Agency for Healthcare Research and Quality; 2018. <https://effectivehealthcare.ahrq.gov/products/collections/cer-methods-guide>. Accessed June 1, 2019.
24. Higgins JP, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011 Oct 18;343:d5928. doi: 10.1136/bmj.d5928. PMID: 22008217.
25. Berger ML, Mamdani M, Atkins D, et al. Good research practices for comparative effectiveness research: defining, reporting and interpreting nonrandomized studies of treatment effects using secondary data sources: the ISPOR Good Research Practices for Retrospective Database Analysis Task Force Report--part I. *Value Health*. 2009 Nov-Dec;12(8):1044-52. doi: 10.1111/j.1524-4733.2009.00600.x. PMID: 19793072.
26. Kliuk-Ben Bassat O, Schechter M, Ashtamker N, et al. Medical cannabis for pain management in patients undergoing chronic hemodialysis: randomized, double-blind, cross-over, feasibility study. *Clinical Kidney Journal*. 2023;16(4):701-10. doi: 10.1093/ckj/sfac275. PMID: 37007688.
27. Tait J, Erridge S, Holvey C, et al. Clinical outcome data of chronic pain patients treated with cannabis-based oils and dried flower from the UK Medical Cannabis Registry. *Expert Rev Neurother*. 2023;1-11. doi: 10.1080/14737175.2023.2195551. PMID: 37021592.
28. Zubcevic K, Petersen M, Bach FW, et al. Oral capsules of tetra-hydro-cannabinol (THC), cannabidiol (CBD) and their combination in peripheral neuropathic pain treatment. *Eur J Pain*. 2022doi: 10.1002/ejp.2072. PMID: 36571471.
29. Bestard JA, Toth CC. An open-label comparison of nabilone and gabapentin as adjuvant therapy or monotherapy in the management of neuropathic pain in patients with peripheral neuropathy. *Pain Pract*. 2011 Jul-Aug;11(4):353-68. doi: 10.1111/j.1533-2500.2010.00427.x. PMID: 21087411.
30. Blake DR, Robson P, Ho M, et al. Preliminary assessment of the efficacy, tolerability and safety of a cannabis-based medicine (Sativex) in the treatment of pain caused by rheumatoid arthritis. *Rheumatology (Oxford)*. 2006 Jan;45(1):50-2. PMID: 16282192.
31. Chaves C, Bittencourt PCT, Pelegrini A. Ingestion of a THC-Rich Cannabis Oil in People with Fibromyalgia: A Randomized, Double-Blind, Placebo-Controlled Clinical Trial. *Pain Med*. 2020;21(10):2212-8. doi: 10.1093/pm/pnaa303. PMID: 33118602.

32. de Vries M, van Rijckevorsel DCM, Vissers KCP, et al. Tetrahydrocannabinol Does Not Reduce Pain in Patients With Chronic Abdominal Pain in a Phase 2 Placebo-controlled Study. *Clin Gastroenterol Hepatol*. 2017 Jul;15(7):1079-86.e4. doi: 10.1016/j.cgh.2016.09.147. PMID: 27720917.
33. Eibach L, Scheffel S, Cardebring M, et al. Cannabidivarin for HIV-Associated Neuropathic Pain: A Randomized, Blinded, Controlled Clinical Trial. *Clin Pharmacol Ther*. 2020 Aug 08;109(4):1055-62. doi: 10.1002/cpt.2016. PMID: 32770831.
34. Frank B, Serpell MG, Hughes J, et al. Comparison of analgesic effects and patient tolerability of nabilone and dihydrocodeine for chronic neuropathic pain: randomised, crossover, double blind study. *BMJ*. 2008 Jan 26;336(7637):199-201. doi: 10.1136/bmj.39429.619653.80. PMID: 18182416.
35. Langford RM, Mares J, Novotna A, et al. A double-blind, randomized, placebo-controlled, parallel-group study of THC/CBD oromucosal spray in combination with the existing treatment regimen, in the relief of central neuropathic pain in patients with multiple sclerosis. *J Neurol*. 2013 Apr;260(4):984-97. doi: 10.1007/s00415-012-6739-4. PMID: 23180178.
36. Lynch ME, Cesar-Rittenberg P, Hohmann AG. A double-blind, placebo-controlled, crossover pilot trial with extension using an oral mucosal cannabinoid extract for treatment of chemotherapy-induced neuropathic pain. *J Pain Symptom Manage*. 2014 Jan;47(1):166-73. doi: 10.1016/j.jpainsymman.2013.02.018. PMID: 23742737.
37. Nurmikko TJ, Serpell MG, Hoggart B, et al. Sativex successfully treats neuropathic pain characterised by allodynia: a randomised, double-blind, placebo-controlled clinical trial. *Pain*. 2007 Dec 15;133(1-3):210-20. PMID: 17997224.
38. Pini LA, Guerzoni S, Cainazzo MM, et al. Nabilone for the treatment of medication overuse headache: results of a preliminary double-blind, active-controlled, randomized trial. *J Headache Pain*. 2012 Nov;13(8):677-84. doi: 10.1007/s10194-012-0490-1. PMID: 23070400.
39. Rintala DH, Fiess RN, Tan G, et al. Effect of dronabinol on central neuropathic pain after spinal cord injury: a pilot study. *Am J Phys Med Rehabil*. 2010 Oct;89(10):840-8. doi: 10.1097/PHM.0b013e3181f1c4ec. PMID: 20855984.
40. Rog DJ, Nurmikko TJ, Friede T, et al. Randomized, controlled trial of cannabis-based medicine in central pain in multiple sclerosis. *Neurology*. 2005 Sep 27;65(6):812-9. PMID: 16186518.
41. Schimrigk S, Marziniak M, Neubauer C, et al. Dronabinol Is a Safe Long-Term Treatment Option for Neuropathic Pain Patients. *Eur Neurol*. 2017;78(5-6):320-9. doi: 10.1159/000481089. PMID: 29073592.
42. Selvarajah D, Gandhi R, Emery CJ, et al. Randomized placebo-controlled double-blind clinical trial of cannabis-based medicinal product (Sativex) in painful diabetic neuropathy: depression is a major confounding factor. *Diabetes Care*. 2010 Jan;33(1):128-30. doi: 10.2337/dc09-1029. PMID: 19808912.
43. Serpell M, Ratcliffe S, Hovorka J, et al. A double-blind, randomized, placebo-controlled, parallel group study of THC/CBD spray in peripheral neuropathic pain treatment. *Eur J Pain*. 2014 Aug;18(7):999-1012. doi: 10.1002/j.1532-2149.2013.00445.x. PMID: 24420962.
44. Skrabek RQ, Galimova L, Ethans K, et al. Nabilone for the treatment of pain in fibromyalgia. *J Pain*. 2008 Feb;9(2):164-73. PMID: 17974490.
45. Toth C, Mawani S, Brady S, et al. An enriched-enrolment, randomized withdrawal, flexible-dose, double-blind, placebo-controlled, parallel assignment efficacy study of nabilone as adjuvant in the treatment of diabetic peripheral neuropathic pain. *Pain*. 2012 Oct;153(10):2073-82. doi: 10.1016/j.pain.2012.06.024. PMID: 22921260.
46. Turcotte D, Doupe M, Torabi M, et al. Nabilone as an adjunctive to gabapentin for multiple sclerosis-induced neuropathic pain: a randomized controlled trial. *Pain Med*. 2015 Jan;16(1):149-59. doi: 10.1111/pme.12569. PMID: 25288189.

47. Vigil JM, Stith SS, Adams IM, et al. Associations between medical cannabis and prescription opioid use in chronic pain patients: A preliminary cohort study. *PLoS ONE*. 2017;12(11):e0187795. doi: 10.1371/journal.pone.0187795. PMID: 29145417.
48. Ware MA, Wang T, Shapiro S, et al. Cannabis for the Management of Pain: Assessment of Safety Study (COMPASS). *J Pain*. 2015 Dec;16(12):1233-42. doi: 10.1016/j.jpain.2015.07.014. PMID: 26385201.
49. Wissel J, Haydn T, Muller J, et al. Low dose treatment with the synthetic cannabinoid Nabilone significantly reduces spasticity-related pain : a double-blind placebo-controlled cross-over trial. *J Neurol*. 2006 Oct;253(10):1337-41. PMID: 16988792.
50. Xu DH, Cullen BD, Tang M, et al. The Effectiveness of Topical Cannabidiol Oil in Symptomatic Relief of Peripheral Neuropathy of the Lower Extremities. *Curr Pharm Biotechnol*. 2020;21(5):390-402. doi: 10.2174/1389201020666191202111534. PMID: 31793418.
51. Zajicek JP, Hobart JC, Slade A, et al. Multiple sclerosis and extract of cannabis: results of the MUSEC trial. *J Neurol Neurosurg Psychiatry*. 2012 Nov;83(11):1125-32. doi: 10.1136/jnnp-2012-302468. PMID: 22791906.
52. Gruber SA, Smith RT, Dahlgren MK, et al. No pain, all gain? Interim analyses from a longitudinal, observational study examining the impact of medical cannabis treatment on chronic pain and related symptoms. *Exp Clin Psychopharmacol*. 2021;29(2):147-56. doi: 10.1037/pha0000435. PMID: 33764103.
53. Campbell G, Hall WD, Peacock A, et al. Effect of cannabis use in people with chronic non-cancer pain prescribed opioids: findings from a 4-year prospective cohort study. *Lancet Public Health*. 2018 Jul;3(7):e341-e50. doi: 10.1016/s2468-2667(18)30110-5. PMID: 29976328.
54. Lee C, Lin M, Martins KJB, et al. Opioid use in medical cannabis authorization adult patients from 2013 to 2018: Alberta, Canada. *BMC Public Health*. 2021;21(1):843. doi: 10.1186/s12889-021-10867-w. PMID: 33933061.
55. Merlin JS, Long D, Becker WC, et al. Marijuana Use Is Not Associated With Changes in Opioid Prescriptions or Pain Severity Among People Living With HIV and Chronic Pain. *J Acquir Immune Defic Syndr*. 2019 06 01;81(2):231-7. doi: 10.1097/QAI.0000000000001998. PMID: 30865181.
56. Vela J, Dreyer L, Petersen KK, et al. Cannabidiol treatment in hand osteoarthritis and psoriatic arthritis: a randomized, double-blind placebo-controlled trial. *Pain*. 2021 Jun 1;163(6):1206-14. PMID: 34510141.
57. Ueberall MA, Vila Silvan C, Essner U, et al. Effectiveness, Safety, and Tolerability of Nabiximols Oromucosal Spray vs Typical Oral Long-Acting Opioid Analgesics in Patients with Severe Neuropathic Back Pain: Analysis of 6-Month Real-World Data from the German Pain e-Registry. *Pain Med*. 2022b;23(4):745-60. doi: 10.1093/pm/pnab263. PMID: 34480564.
58. Ueberall MA, Essner U, Silván CV, et al. Comparison of the Effectiveness and Tolerability of Nabiximols (THC:CBD) Oromucosal Spray versus Oral Dronabinol (THC) as Add-on Treatment for Severe Neuropathic Pain in Real-World Clinical Practice: Retrospective Analysis of the German Pain e-Registry. *J Pain Res*. 2022a;15:267-86. doi: 10.2147/JPR.S340968. PMID: 35140513.

Authors

Roger Chou, M.D., FACP
Azrah Y. Ahmed, B.A.
Christina Bougatsos, M.P.H.
Benjamin J. Morasco, Ph.D.
Tracy Dana, M.L.S.
Rongwei Fu, Ph.D.

Acknowledgments

The authors gratefully acknowledge the following individuals for their contributions to this project: Pacific Northwest Evidence-based Practice Center Professor Emerita Marian McDonagh, Pharm.D., informatician Connor Smith, M.S., for work on data visualization, and research associate and librarian Andrew Hamilton, M.L.S., all from Oregon Health & Science University; Jesse Wagner, M.A.; and Task Order Officer Suchitra Iyer, Ph.D., at the Agency for Healthcare Research and Quality.

Disclaimers

This report is based on research conducted by the Pacific Northwest Evidence-based Practice Center under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. 75Q80120D00006). The findings and conclusions in this document are those of the authors, who are responsible for its contents; the findings and conclusions do not necessarily represent the views of AHRQ. Therefore, no statement in this report should be construed as an official position of AHRQ or of the U.S. Department of Health and Human Services.

None of the investigators have any affiliations or financial involvement that conflicts with the material presented in this report.

The information in this report is intended to help healthcare decision makers—patients and clinicians, health system leaders, and policymakers, among others—make well-informed decisions and thereby improve the quality of health care services. This report is not intended to be a substitute for the application of clinical judgment. Anyone who makes decisions concerning the provision of clinical care should consider this report in the same way as any medical reference and in conjunction with all other pertinent information, i.e., in the context of available resources and circumstances presented by individual patients.

This report is made available to the public under the terms of a licensing agreement between the author and the Agency for Healthcare Research and Quality. Most AHRQ documents are publicly available to use for noncommercial purposes (research, clinical or patient education, quality improvement projects) in the United States, and do not need specific permission to be reprinted and used unless they contain material that is copyrighted by others. Specific written permission is needed for commercial use (reprinting for sale, incorporation into software, incorporation into for-profit training courses) or for use outside of the U.S. If organizational policies require permission to adapt or use these materials, AHRQ will provide such permission in writing.

AHRQ or U.S. Department of Health and Human Services endorsement of any derivative products that may be developed from this report, such as clinical practice guidelines, other quality enhancement tools, or reimbursement or coverage policies, may not be stated or implied.

AHRQ appreciates appropriate acknowledgment and citation of its work. Suggested language for acknowledgment: This work is the first surveillance update of a living systematic evidence report, Living Systematic Review on Cannabis and Other Plant-Based Treatments for Chronic Pain: 2023 Update, by the Evidence-based Practice Center Program at the Agency for Healthcare Research and Quality (AHRQ) following the second yearly full systematic review.

Suggested citation: Chou R, Ahmed AY, Bougatsos C, Morasco BJ, Dana T, Fu R. Living Systematic Review on Cannabis and Other Plant-Based Treatments for Chronic Pain: 2023 Update—Surveillance Report 1. (Prepared by

the Pacific Northwest Evidence-based Practice Center under Contract No. 75Q80120D00006.) AHRQ Publication No. 23-EHC032. Rockville, MD: Agency for Healthcare Research and Quality; September 2023. DOI: <https://doi.org/10.23970/AHRQEPCCER250.2023UPDATESR1>. Posted final reports are located on the Effective Health Care Program [search page](#).

Afterword

The Agency for Healthcare Research and Quality (AHRQ), through its Evidence-based Practice Centers (EPCs), sponsors the development of systematic reviews to assist public- and private-sector organizations in their efforts to improve the quality of healthcare in the United States. These reviews provide comprehensive, science-based information on common, costly medical conditions, and new healthcare technologies and strategies.

Systematic reviews are the building blocks underlying evidence-based practice; they focus attention on the strength and limits of evidence from research studies about the effectiveness and safety of a clinical intervention. In the context of developing recommendations for practice, systematic reviews can help clarify whether assertions about the value of the intervention are based on strong evidence from clinical studies. For more information about AHRQ EPC systematic reviews, see <https://effectivehealthcare.ahrq.gov/about/epc/evidence-synthesis>.

This and future quarterly surveillance reports will provide up-to-date information following the last full systematic review about the evidence base to inform health plans, providers, purchasers, government programs, and the healthcare system as a whole on the state of the science. Transparency and stakeholder input are essential to the Effective Health Care Program. Please visit the website (www.effectivehealthcare.ahrq.gov) to see draft research questions and reports or to join an email list to learn about new program products and opportunities for input.

If you have comments on this report, they may be sent by mail to the Task Order Officer named below at: Agency for Healthcare Research and Quality, 5600 Fishers Lane, Rockville, MD 20857, or by email to epc@ahrq.hhs.gov. They will be considered in the next version of the report.

Robert Otto Valdez, Ph.D., M.H.S.A.
Director
Agency for Healthcare Research and Quality

Therese Miller, Dr.P.H.
Director
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Christine Chang M.D., M.P.H.
Acting Director
Evidence-based Practice Center Program
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Suchitra Iyer, Ph.D.
Task Order Officer
Evidence-based Practice Center Program
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Appendix Contents

Appendix A. Literature Search Strategies	A-1
Appendix B. Methods	B-1
Inclusion and Exclusion Criteria.....	B-1
Data Extraction.....	B-3
Risk of Bias Assessment of Individual Studies.....	B-3
Data Synthesis and Analysis	B-3
Grading the Strength of the Body of Evidence	B-5
Peer Review and Public Commentary.....	B-6
Assessing Applicability	B-6
Appendix B References.....	B-1
Appendix C. Included Studies List	C-1
Appendix D. Literature Flow Diagram.....	D-1
Appendix E. Results.....	E-1
Individual Study Summary Tables	E-1
Meta-Analysis Results.....	E-15
Comparable THC to CBD Ratio Studies.....	E-15
High THC to CBD Ratio Studies	E-20
Appendix F. Evidence Tables	F-1
Appendix G. Risk of Bias Assessment	G-1
Appendix H. Details on Strength of Evidence.....	H-1
Appendix H References.....	H-12
Appendix I. Excluded Studies List	I-1
Appendix J. Funnel Plot of High THC Ratio Studies Included in Meta-Analysis for Pain Severity	J-1

Appendix A. Literature Search Strategies

Database: Ovid MEDLINE(R) ALL 1946 to June 29, 2023

- 1 Chronic Pain/
- 2 exp arthralgia/ or exp back pain/ or exp headache/ or exp musculoskeletal pain/ or neck pain/ or exp neuralgia/ or exp nociceptive pain/ or pain, intractable/ or fibromyalgia/ or myalgia/
- 3 Pain/
- 4 chronic or subacute* or sub-acute*).ti,ab,kw.
- 5 3 and 4
- 6 ((chronic or persistent or intractable or refractory or subacute* or sub-acute) adj3 pain).ti,ab,kw.
- 7 (((back or spine or spinal or leg or musculoskeletal or neuropathic or nociceptive or radicular) adj1 pain) or headache or arthritis or fibromyalgia or osteoarthritis).ti,ab,kw.
- 8 1 or 2 or 5 or 6 or 7
- 9 Cannabis/
- 10 exp Cannabinoids/
- 11 Medical Marijuana/
- 12 Mitragyna/
- 13 (cannabis or cannabinoid* or cannabiniol or marijuana or cannabidiol or phytocannabinoid* or tetrahydrocannabinol or dronabinol or nabilone or sativex or "CBD" or "THC" or kratom or khat or qat or psilocybin or hemp or hydroxymitragynine).ti,ab,kf.
- 14 or/9-13
- 15 8 and 14
- 16 limit 15 to english language
- 17 (Animals/ or Models, Animal/ or Disease Models, Animal/) not Humans/
- 18 ((animal or animals or avian or bird or birds or bovine or canine or cow* or dog or dogs or cat or cats or feline or hamster* or horse* or lamb or lamb* or mouse or mice or monkey or monkeys or murine or pig or piglet* or pigs or porcine or primate* or rabbit* or rat or rats or rodent* or songbird* or veterinar*) not (human* or patient*)).ti,kf,jw.
- 19 or/17-18
- 20 16 not 19

Database: EBM Reviews - Cochrane Central Register of Controlled Trials May 2023

- 1 Chronic Pain/
- 2 exp arthralgia/ or exp back pain/ or exp headache/ or exp musculoskeletal pain/ or neck pain/ or exp neuralgia/ or exp nociceptive pain/ or pain, intractable/ or fibromyalgia/ or myalgia/
- 3 Pain/
- 4 (chronic or subacute* or sub-acute*).ti,ab,kw.
- 5 3 and 4
- 6 ((chronic or persistent or intractable or refractory or subacute* or sub-acute*) adj3 pain).ti,ab,hw.
- 7 (((back or spine or spinal or leg or musculoskeletal or neuropathic or nociceptive or radicular) adj1 pain) or headache or arthritis or fibromyalgia or osteoarthritis).ti,ab,hw.
- 8 1 or 2 or 5 or 6 or 7

- 9 (cannabis or cannabinoid* or cannabinal or marijuana or cannabidiol or phytocannabinoid* or tetrahydrocannabinol or dronabinol or nabilone or sativex or "CBD" or "THC" or kratom or khat or qat or psilocybin or hemp or hydroxymitragynine).ti,ab,hw.
- 10 8 and 9
- 11 conference abstract.pt.
- 12 "journal: conference abstract".pt.
- 13 "journal: conference review".pt.
- 14 "http://.www.who.int/trialsearch*".so.
- 15 "https://clinicaltrials.gov*".so.
- 16 11 or 12 or 13 or 14 or 15
- 17 10 not 16

Database: APA PsycInfo 1806 to June Week 3, 2023

- 1 Chronic Pain/
- 2 exp arthralgia/ or exp back pain/ or exp headache/ or exp musculoskeletal pain/ or neck pain/ or exp neuralgia/ or exp nociceptive pain/ or pain, intractable/ or fibromyalgia/ or myalgia/
- 3 Pain/
- 4 (chronic or subacute* or sub-acute*).ti,ab.
- 5 3 and 4
- 6 ((chronic or persistent or intractable or refractory or subacute* or sub-acute*) adj3 pain).ti,ab.
- 7 (((back or spine or spinal or leg or musculoskeletal or neuropathic or nociceptive or radicular) adj1 pain) or headache or arthritis or fibromyalgia or osteoarthritis).ti,ab.
- 8 1 or 2 or 5 or 6 or 7
- 9 Cannabis/
- 10 exp Cannabinoids/
- 11 (cannabis or cannabinoid* or cannabinal or marijuana or cannabidiol or phytocannabinoid* or tetrahydrocannabinol or dronabinol or nabilone or sativex or "CBD" or "THC" or kratom or khat or qat or psilocybin or hemp or hydroxymitragynine).ti,ab.
- 12 or/9-11
- 13 8 and 12
- 14 limit 13 to english language

Database: Elsevier Embase to July 2, 2023

('cannabis'/exp OR cannabis OR cannabinoid* OR 'cannabinal'/exp OR cannabinal OR 'marijuana'/exp OR marijuana OR 'cannabidiol'/exp OR cannabidiol OR phytocannabinoid* OR 'tetrahydrocannabinol'/exp OR tetrahydrocannabinol OR 'dronabinol'/exp OR dronabinol OR 'nabilone'/exp OR nabilone OR 'sativex'/exp OR sativex OR 'cbd' OR 'thc' OR 'kratom'/exp OR kratom OR 'khat'/exp OR khat OR 'qat'/exp OR qat OR 'psilocybin'/exp OR psilocybin OR 'hemp'/exp OR hemp OR hydroxymitragynine) AND ('chronic pain'/exp OR 'subacute pain'/exp OR 'subacute pain' OR arthralgia OR 'back pain' OR headache OR 'musculoskeletal pain' OR 'neck pain' OR neuralgia OR 'nociceptive pain' OR 'intractable pain' OR fibromyalgia OR myalgia OR arthritis OR osteoarthritis) NOT ((animal OR animals OR avian OR bird OR birds OR bovine OR canine OR cow* OR dog OR dogs OR cat OR cats OR feline OR hamster* OR horse* OR lamb OR lamb* OR mouse OR mice OR monkey OR monkeys OR murine OR pig OR piglet* OR pigs OR porcine OR primate* OR rabbit* OR rat OR rats OR rodent* OR songbird* OR veterinar*) NOT (human* OR patient*)) AND ('article'/it OR 'article in press'/it

OR 'conference paper'/it OR 'preprint'/it OR 'review'/it) AND [english]/lim AND [embase]/lim
NOT ([embase]/lim AND [medline]/lim)

Database: Elsevier Scopus June 11, 2023

((TITLE (cannabis OR cannabinoid* OR cannabinoL OR marijuana OR cannabidiol OR phytocannab
inoid* OR tetrahydrocannabinol OR dronabinol OR nabilone OR sativex OR "CBD" OR
"THC" OR kratom OR khat OR qat OR psilocybin OR hemp OR hydroxymitragynine)
) AND (TITLE ("chronic pain" OR "subacute pain" OR arthralgia OR "back
pain" OR headache OR "musculoskeletal pain" OR "neck
pain" OR neuralgia OR "nociceptive pain" OR "intractable
pain" OR fibromyalgia OR myalgia OR arthritis OR osteoarthritis OR "neuropathic pain")
) AND NOT (TITLE-ABS-KEY (animal OR animals OR avian OR bird OR birds OR bovine OR canine OR cow* OR d
og OR dogs OR cat OR cats OR feline OR hamster* OR horse* OR lamb OR lamb*
OR mouse OR mice OR monkey OR monkeys OR murine OR pig OR piglet* OR pigs
OR porcine OR primate* OR rabbit* OR rat OR rats OR rodent* OR songbird* OR vet
erinar*)) AND (LIMIT-TO (LANGUAGE , "English"))

Appendix B. Methods

Inclusion and Exclusion Criteria

Table B-1 outlines the inclusion and exclusion criteria related to populations, interventions, comparators, outcomes, timing, and settings (PICOTS), and study designs of interest for each Key Question (KQ). In the winter of 2022, the protocol was amended to include adolescents and subacute pain.¹ These changes were documented on in a revised protocol submitted to PROSPERO,² the AHRQ Protocol, and the title, key questions, and inclusion and exclusion criteria were edited to reflect said changes. The changes expanded inclusion criteria to include subacute pain and adolescents.

KQ1: In adults or adolescents with chronic or subacute pain, what are the benefits of cannabinoids for treatment of chronic or subacute pain?

KQ2: In adults or adolescents with chronic or subacute pain, what are the harms of cannabinoids for treatment of chronic or subacute pain?

KQ3: In adults or adolescents with chronic or subacute pain, what are the benefits of kratom or other plant-based substances for treatment of chronic or subacute pain?

KQ4: In adults or adolescents with chronic or subacute pain, what are the harms of kratom or other plant-based substances for treatment of chronic or subacute pain?

Table B-1. PICOTS

PICOTS Element	Inclusion Criteria	Exclusion Criteria
Population	All KQs: Adults or adolescents (including pregnant or breastfeeding women) with noncancer chronic pain (>12 weeks or pain persisting past the time for normal tissue healing) or subacute pain (pain lasting 4 weeks to 3 months). See categorization of specifically included pain populations below.	All KQs: Children; adults with acute pain; patients at end of life or in palliative care (e.g., with late stage cancer-related pain)
Interventions	KQs 1 and 2: Cannabinoids (including synthetics) using different delivery mechanisms such as oral, buccal, inhalational, topical, or other administration routes KQs 3 and 4: Kratom or other plant-based substances; co-use of kratom or other plant-based substances and opioids All KQs: Co-use of other drugs for pain	All KQs: Non-plant-based interventions, capsaicin, herbal supplements
Comparators	All KQs: Any comparator or usual care	All KQs: No comparison

PICOTS Element	Inclusion Criteria	Exclusion Criteria
Outcomes	All KQs: Primary efficacy outcomes (i.e., pain, general function [e.g., Short-Form 36 Physical Functioning Scale] or pain-related [e.g., Oswestry Disability Index or Roland-Morris Disability Questionnaire, for low back pain] or disability, including pain interference ^a); harms and adverse effects (e.g., dizziness, nausea, sedation, development of cannabis use disorder, serious adverse events as defined by study); secondary outcomes (i.e., psychological distress including depression and anxiety, quality of life, opioid use, sleep quality, sleep disturbance, healthcare utilization)	All KQs: Other outcomes
Time of followup	All KQs: short term (4 weeks to <6 months), intermediate term (6 to <12 months), long term (≥1 year)	All KQs: Studies with <1-month (4 weeks) of treatment or followup after treatment
Setting	All KQs: Any nonhospital setting or setting of self-directed care	All KQs: Hospital care, hospice care, emergency department care
Study design	All KQs: RCTs; observational studies with a concurrent control group for harms, and to fill gaps in the evidence for benefits	All KQs: Other study designs

Abbreviations: KQ = Key Question; PICOTS = populations, interventions, comparators, outcomes, timing, and settings; RCT = randomized controlled trial.

^aThe degree to which pain directly interferes with patients' ability to participate in their daily activities (challenges in performing daily, social, or work-related tasks due to pain).

Important subgroups to consider in evaluating this evidence are:

- Specific types of pain: neuropathic pain (including nociceptive and centralized; patients with multiple sclerosis and painful skin disorders are included in this category), musculoskeletal pain (including low-back pain), visceral pain, fibromyalgia, inflammatory arthritis, headache disorders, sickle cell disease, and cancer pain (non-end of life)
- Degree of nociplasticity/central sensitization
- Patient demographics (e.g., age, race, ethnicity, sex, socioeconomic status)
- Comorbidities, including past or current substance use disorders, mental health disorders, medical comorbidities, and high risk for opioid use disorder)
- Plant-based compound characteristics: route of administration, frequency of administration, potency of product, dose or estimated dose, specific compounds (e.g. tetrahydrocannabinol, cannabidiol, terpenes, flavonoids), and specific formulations used
- Co-use of other interventions for pain: opioids, nonopioids (e.g., nonsteroidal anti-inflammatory drugs, acetaminophen, gabapentin, pregabalin)

Below are additional details on the scope of this project:

Study Design: For all Key Questions, we included randomized controlled trials (RCTs) of at least 4 weeks duration. Initially, in the base-year of this living systematic review, we included observational studies for both benefits (to address gaps in evidence where RCTs are not available) and harms. Eligible observational studies must have assessed a mean duration of treatment of at least 4 weeks, and have concurrent controls (e.g., cohort and case-control studies). Those controlling for potential confounders were prioritized. As the evidence grows, and more RCTs become available throughout the project, we will reassess the need to include observational studies, specifically to address benefits. A decision to discontinue including them

will be made based on the strength of the RCT evidence. When the RCT evidence on a given Key Question and outcome is insufficient, we will include observational studies that meet inclusion criteria. When the strength of evidence is low, moderate, or high based on RCTs, we will update our protocol to exclude observational studies. We do not anticipate excluding observational studies assessing harms. For all Key Questions, we excluded uncontrolled observational studies, case series, and case reports. Systematic reviews were used to supplement searches and identify primary studies.

Non-English-Language Studies: We restricted to English-language articles, but reviewed English-language abstracts of non-English language articles to identify studies that would otherwise meet inclusion criteria in order to help assess for the likelihood of language bias.

Data Extraction

After studies were selected for inclusion, data were abstracted into categories that included but are not limited to: study design, year, setting, country, sample size, eligibility criteria, population and clinical characteristics, intervention characteristics, and results relevant to each Key Question as outlined in the previous inclusion and exclusion criteria section. Information that was abstracted that was relevant for assessing applicability included the number of patients randomized relative to the number of patients enrolled, use of run-in or wash-out periods, and characteristics of the population, intervention, and care settings. All study data were verified for accuracy and completeness by a second team member. On a quarterly basis, any newly identified studies were abstracted and evidence tables updated. Quarterly reports were published to the Agency for Healthcare Research and Quality (AHRQ) website, and evidence tables are updated in AHRQ's Systematic Review Data Repository Plus (SRDR+).

Risk of Bias Assessment of Individual Studies

Predefined criteria were used to assess the risk of bias of individual controlled trials, systematic reviews, and observational studies. RCTs were evaluated using criteria and methods developed by the Cochrane Back Review Group,³ and cohort and case-control studies were evaluated using criteria developed by the U.S. Preventive Services Task Force.⁴ These criteria and methods were used in accordance with the approach recommended in the chapter, *Assessing the Risk of Bias of Individual Studies When Comparing Medical Interventions in the Methods Guide for Effectiveness and Comparative Effectiveness Reviews* developed by AHRQ.⁵ Studies were given an overall rating of “low,” “medium,” or “high” risk of bias. We used DistillerSR[®] software to conduct these assessments, using dual review by two independent reviewers. Disagreements identified by DistillerSR[®] were resolved through consensus. Assessments and final ratings were converted to evidence tables, and will be uploaded on a quarterly basis to SRDR+.

Data Synthesis and Analysis

We constructed evidence tables showing study characteristics (as discussed above), results, and risk of bias ratings for all included studies, and summary tables to highlight the main findings. Data were qualitatively summarized in tables, using ranges and descriptive analysis and interpretation of the results. Studies identified in prior AHRQ chronic pain reports^{6,7} that meet inclusion criteria are included in this review. We evaluated the persistence of benefits or harms

by evaluating the three periods identified in prior AHRQ pain reports (3 to <6 months, 6 to 12 months, and ≥ 12 months).⁶⁻¹⁰

Meta-analyses were conducted to summarize data and obtain more precise estimates on outcomes for which studies were homogeneous enough to provide a meaningful combined estimate.¹¹ The decision to conduct quantitative synthesis depends on presence of at least two studies, completeness of reported outcomes and a lack of heterogeneity among the reported results. To determine whether meta-analyses were indicated, we considered the risk of bias of the studies and the heterogeneity among studies in design, patient population, interventions, and outcomes. Meta-analyses were conducted using a random effects model based on the profile likelihood method,¹² and statistical heterogeneity was assessed using the I^2 method. Publication bias (small sample size bias) was assessed using funnel plots when there are eight or more studies in meta-analyses. To evaluate subgroup effects, we summarized within-study analyses of subgroup differences and performed study-level analyses on key demographic and clinical factors. Sensitivity analyses were conducted on study risk of bias.

The magnitude of effects for pain and function is classified using the same system used in other recent AHRQ Evidence-based Practice Center (EPC) reviews conducted on chronic pain⁶⁻¹⁰ to provide a consistent benchmark for comparing results of pain interventions across reviews. Table B-2 provides thresholds for determining the magnitude of effect. A small effect is defined for pain as a mean between-group difference following treatment of 5 to 10 points on a 0- to 100-point visual analog scale (VAS), 0.5 to 1.0 points on a 0- to 10-point numeric rating scale, or equivalent; for function as a mean difference of 5 to 10 points on the 0- to 100-point Oswestry Disability Index (ODI) or 1 to 2 points on the 0- to 24-point Roland-Morris Disability Questionnaire (RDQ), or equivalent; and for any outcome as a standardized mean difference (SMD) of 0.2 to 0.5. A moderate effect is defined for pain as a mean difference of 10 to 20 points on a 0- to 100-point VAS, for function as a mean difference of 10 to 20 points on the ODI or 2 to 5 points on the RDQ, and for any outcome as an SMD of 0.5 to 0.8. Large effects are defined as greater than moderate. We apply similar thresholds to other outcomes measures. Small effects using this system may be below published thresholds for clinically meaningful effects; however, there is variability across individual patients regarding what constitutes a clinically meaningful effect, which is influenced by a number of factors such as preferences, duration and type of chronic pain, baseline symptom severity, harms, and costs. For some patients a small improvement in pain or function using a treatment with low cost or no serious harms may be important.

Table B-2. Definitions of effect sizes

Effect Size	Definition
Small effect	<ul style="list-style-type: none"> • MD 0.5 to 1.0 points on a 0 to 10-point scale, 5 to 10 points on a 0 to 100-point scale • SMD 0.2 to 0.5 • RR/OR 1.2 to 1.4
Moderate effect	<ul style="list-style-type: none"> • MD >1 to 2 points on a 0 to 10-point scale, >10 to 20 points on a 0 to 100-point scale • SMD >0.5 to 0.8 • RR/OR 1.5 to 1.9
Large effect	<ul style="list-style-type: none"> • MD >2 points on a 0 to 10-point scale, >20 points on a 0 to 100-point scale • SMD >0.8 • RR/OR ≥ 2.0

Abbreviations: MD = mean difference; OR = odds ratio; RR = risk ratio; SMD = standardized mean difference.

Findings that were not statistically significant were interpreted as follows:

- In determining the strength of evidence (SOE), the precision of evidence was downgraded two levels if inadequate sample size (optimal information size) *and* the 95% confidence interval includes both potentially meaningful benefit and harm (e.g. for a relative effect, the lower bound is ≤ 0.75 *and* the upper bound is ≥ 1.25)¹³
- If the magnitude of effect is below the threshold for a small effect, the finding is considered to have “No effect”⁶
- If the magnitude of effect is small or greater, and SOE is at least Low, the finding is considered to have a “Potential effect, not statistically significant”
- If the magnitude of effect is small or greater, and SOE is insufficient, the finding is considered to have “failed to demonstrate or exclude a beneficial/detrimental effect.”¹⁴

Grading the Strength of the Body of Evidence

We assessed the SOE for all primary comparisons and outcomes listed in Table B-1. Regardless of whether evidence is synthesized quantitatively or qualitatively, the strength of evidence for each Key Question/body of evidence is initially assessed by one researcher for each clinical outcome by using the approach described in the AHRQ Methods Guide.⁵ To ensure consistency and validity of the evaluation, the strength of evidence is reviewed by the entire team of investigators prior to assigning a final grade on the following factors:

- Study limitations (low, medium, or high level of study limitations)
- Consistency (consistent, inconsistent, or unknown/not applicable)
- Directness (direct or indirect)
- Precision (precise or imprecise)
- Reporting/publication bias (suspected or undetected)

The SOE was assigned an overall grade of high, moderate, low, or insufficient according to a four-level scale by evaluating and weighing the combined results of the above domains:

- High—We are very confident that the estimate of effect lies close to the true effect for this outcome. The body of evidence has few or no deficiencies. We believe that the findings are stable, i.e., another study would not change the conclusions.
- Moderate—We are moderately confident that the estimate of effect lies close to the true effect for this outcome. The body of evidence has some deficiencies. We believe that the findings are likely to be stable, but some doubt remains.
- Low—We have limited confidence that the estimate of effect lies close to the true effect for this outcome. The body of evidence has major or numerous deficiencies (or both). We believe that additional evidence is needed before concluding either that the findings are stable or that the estimate of effect is close to the true effect.
- Insufficient—We have no evidence, we are unable to estimate an effect, or we have no confidence in the estimate of effect for this outcome. No evidence is available or the body of evidence has unacceptable deficiencies, precluding reaching a conclusion.

Plain-language statements are used in the Main Points and the Results to Date sections to convey the SOE. High SOE is described as "is associated with" or simply "reduces/increases;" moderate SOE is described as "probably;" and low SOE is described as "may be."¹⁵

Peer Review and Public Commentary

Peer reviewers are invited to provide written comments on the draft report based on their clinical, content, or methodological expertise. The EPC considers all peer review comments on the draft report in preparation of the final report. Peer reviewers do not participate in writing or editing of the final report or other products. The final report does not necessarily represent the views of individual reviewers. The EPC will complete a disposition of all peer review comments. The disposition of comments for systematic reviews and technical briefs will be published 3 months after the publication of the evidence report.

Potential Peer Reviewers must disclose any financial conflicts of interest greater than \$5,000 and any other relevant business or professional conflicts of interest. Invited Peer Reviewers may not have any financial conflict of interest greater than \$5,000. Peer reviewers who disclose potential business or professional conflicts of interest may submit comments on draft reports through the public comment mechanism.

Assessing Applicability

Applicability is assessed in accordance with the AHRQ Methods Guide,¹⁶ which is based on the PICOTS framework. Applicability addresses the extent to which outcomes associated with an intervention are likely to be similar across different patients and settings in clinical practice based on the populations, interventions, comparisons, and outcomes evaluated in the studies. For example, exclusion of chronic pain patients with psychiatric comorbidities reduces applicability to clinical practice since many patients with chronic pain have such comorbidities and may respond more poorly to treatment. Similarly, trials that use active run-in periods evaluate highly selected populations who tolerated and responded well to the study intervention, rather than the general population of chronic pain patients being considered for the intervention. Factors that may affect applicability which we have identified a priori include eligibility criteria and patient factors (e.g., demographic characteristics, duration or severity of pain, underlying pain condition, presence of medical and psychiatric comorbidities, event rates and symptom severity in treatment and control groups), intervention factors (e.g., dose and duration of therapy, intensity and frequency of monitoring, level of adherence, use of co-interventions), comparisons (e.g., type and dosing of comparison), outcomes (e.g., use of unvalidated or nonstandardized outcomes, measurement of short-term or surrogate outcomes), settings (e.g., primary care vs. specialty setting, country), and study design features (e.g., use of run-in periods) relevant to applicability. We use this information to assess the situations in which the evidence is most relevant and to evaluate applicability to real-world clinical practice in typical U.S. settings, summarizing applicability assessments qualitatively.

Appendix B References

1. Chou R, Ahmed AY, Iyer S, et al. Living Systematic Reviews: Practical Considerations for the Agency for Healthcare Research and Quality Evidence-based Practice Center Program from Year 2 of an Ongoing Review [White Paper in Progress]. In: Quality AfHRA, editor: Pacific Northwest Evidence-based Practice Center; 2023.
2. Chou R, Ahmed AY, Morasco BJ, et al. Living Systematic Review on Cannabis and Other Plant-based Treatments for Chronic and Subacute Pain. PROSPERO 2021 CRD42021229579 Available from: . https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021229579.

3. Furlan AD, Pennick V, Bombardier C, et al. 2009 updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine*. 2009 Aug 15;34(18):1929-41. doi: 10.1097/BRS.0b013e3181b1c99f. PMID: 19680101.
4. U.S. Preventive Services Task Force. *Methods and processes*. 2019. <https://www.uspreventiveservicestaskforce.org/uspstf/about-uspstf/methods-and-processes>.
5. *Methods guide for effectiveness and comparative effectiveness reviews*. Rockville, MD: Agency for Healthcare Research and Quality; 2018. <https://effectivehealthcare.ahrq.gov/products/collections/ceer-methods-guide>. Accessed June 1, 2019.
6. Chou R, Hartung D, Turner J, et al. *Opioid treatments for chronic pain*. Rockville, MD; 2020.
7. McDonagh MS, Selph SS, Buckley DI, et al. *Nonopioid pharmacologic treatments for chronic pain*. Rockville, MD: Agency for Healthcare Research and Quality; 2020. doi: 10.23970/AHRQEPCCER228. PMID: 32338847.
8. Skelly AC, Chou R, Dettori JR, et al. *Noninvasive nonpharmacological treatment for chronic pain: a systematic review update*. Rockville, MD: Agency for Healthcare Research and Quality; 2020. doi: 10.23970/AHRQEPCCER227. PMID: 32338846.
9. Chou R, Deyo R, Friedly J, et al. *Noninvasive treatments for low back pain: Agency for Healthcare Research and Quality (US), Rockville (MD); 2016*.
10. Skelly AC, Chou R, Dettori JR, et al. *Noninvasive nonpharmacological treatment for chronic pain: a systematic review: Agency for Healthcare Research and Quality (US), Rockville (MD); 2018*.
11. Morton SC, Murad MH, O'Connor E, et al. *Quantitative synthesis—an update: Agency for Healthcare Research and Quality (US), Rockville (MD); 2018*.
12. Huizenga HM, Visser I, Dolan CV. Testing overall and moderator effects in random effects meta-regression. *The British journal of mathematical and statistical psychology*. 2011 Feb;64(Pt 1):1-19. doi: 10.1348/000711010x522687. PMID: 21506942.
13. Guyatt GH, Oxman AD, Kunz R, et al. *GRADE guidelines 6. Rating the quality of evidence—imprecision*. *J Clin Epidemiol*. 2011 Dec;64(12):1283-93. doi: 10.1016/j.jclinepi.2011.01.012. PMID: 21839614.
14. Guyatt GH, Norris SL, Schulman S, et al. *Methodology for the development of antithrombotic therapy and prevention of thrombosis guidelines: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines*. *Chest*. 2012 Feb;141(2 Suppl):53s-70s. doi: 10.1378/chest.11-2288. PMID: 22315256.
15. Gerrity M, Fiordalisi C, Pillay J, et al. *AHRQ methods for effective health care. Roadmap for Narratively Describing Effects of Interventions in Systematic Reviews*. Rockville (MD): Agency for Healthcare Research and Quality (US); 2020.
16. Atkins D, Chang SM, Gartlehner G, et al. Assessing applicability when comparing medical interventions: AHRQ and the Effective Health Care Program. *J Clin Epidemiol*. 2011 Nov;64(11):1198-207. doi: 10.1016/j.jclinepi.2010.11.021. PMID: 21463926.

Appendix C. Included Studies List


1. Bestard JA, Toth CC. An open-label comparison of nabilone and gabapentin as adjuvant therapy or monotherapy in the management of neuropathic pain in patients with peripheral neuropathy. *Pain Pract.* 2011 Jul-Aug;11(4):353-68. doi: 10.1111/j.1533-2500.2010.00427.x. PMID: 21087411.
2. Blake DR, Robson P, Ho M, et al. Preliminary assessment of the efficacy, tolerability and safety of a cannabis-based medicine (Sativex) in the treatment of pain caused by rheumatoid arthritis. *Rheumatology (Oxford).* 2006 Jan;45(1):50-2. PMID: 16282192.
3. Campbell G, Hall WD, Peacock A, et al. Effect of cannabis use in people with chronic non-cancer pain prescribed opioids: findings from a 4-year prospective cohort study. *Lancet Public Health.* 2018 Jul;3(7):e341-e50. doi: 10.1016/s2468-2667(18)30110-5. PMID: 29976328.
4. Chaves C, Bittencourt PCT, Pelegrini A. Ingestion of a THC-Rich Cannabis Oil in People with Fibromyalgia: A Randomized, Double-Blind, Placebo-Controlled Clinical Trial. *Pain Med.* 2020;21(10):2212-8. doi: 10.1093/pm/pnaa303. PMID: 33118602.
5. de Vries M, van Rijckevorsel DCM, Vissers KCP, et al. Tetrahydrocannabinol Does Not Reduce Pain in Patients With Chronic Abdominal Pain in a Phase 2 Placebo-controlled Study. *Clin Gastroenterol Hepatol.* 2017 Jul;15(7):1079-86.e4. doi: 10.1016/j.cgh.2016.09.147. PMID: 27720917.
6. Eibach L, Scheffel S, Cardebring M, et al. Cannabidiol for HIV-Associated Neuropathic Pain: A Randomized, Blinded, Controlled Clinical Trial. *Clin Pharmacol Ther.* 2020 Aug 08;109(4):1055-62. doi: 10.1002/cpt.2016. PMID: 32770831.
7. Frank B, Serpell MG, Hughes J, et al. Comparison of analgesic effects and patient tolerability of nabilone and dihydrocodeine for chronic neuropathic pain: randomised, crossover, double blind study. *BMJ.* 2008 Jan 26;336(7637):199-201. doi: 10.1136/bmj.39429.619653.80. PMID: 18182416.
8. Gruber SA, Smith RT, Dahlgren MK, et al. No pain, all gain? Interim analyses from a longitudinal, observational study examining the impact of medical cannabis treatment on chronic pain and related symptoms. *Exp Clin Psychopharmacol.* 2021;29(2):147-56. doi: 10.1037/pha0000435. PMID: 33764103.
9. Kliuk-Ben Bassat O, Schechter M, Ashtamker N, et al. Medical cannabis for pain management in patients undergoing chronic hemodialysis: randomized, double-blind, cross-over, feasibility study. *Clinical Kidney Journal.* 2023;16(4):701-10. doi: 10.1093/ckj/sfac275. PMID: 37007688.
10. Langford RM, Mares J, Novotna A, et al. A double-blind, randomized, placebo-controlled, parallel-group study of THC/CBD oromucosal spray in combination with the existing treatment regimen, in the relief of central neuropathic pain in patients with multiple sclerosis. *J Neurol.* 2013 Apr;260(4):984-97. doi: 10.1007/s00415-012-6739-4. PMID: 23180178.
11. Lee C, Lin M, Martins KJB, et al. Opioid use in medical cannabis authorization adult patients from 2013 to 2018: Alberta, Canada. *BMC Public Health.* 2021;21(1):843. doi: 10.1186/s12889-021-10867-w. PMID: 33933061.
12. Lynch ME, Cesar-Rittenberg P, Hohmann AG. A double-blind, placebo-controlled, crossover pilot trial with extension using an oral mucosal cannabinoid extract for treatment of chemotherapy-induced neuropathic pain. *J Pain Symptom Manage.* 2014 Jan;47(1):166-73. doi: 10.1016/j.jpainsymman.2013.02.018. PMID: 23742737.
13. Merlin JS, Long D, Becker WC, et al. Marijuana Use Is Not Associated With Changes in Opioid Prescriptions or Pain Severity Among People Living With HIV and Chronic Pain. *J Acquir Immune Defic Syndr.* 2019 06 01;81(2):231-7. doi: 10.1097/QAI.0000000000001998. PMID: 30865181.
14. Nurmikko TJ, Serpell MG, Hoggart B, et al. Sativex successfully treats neuropathic pain characterised by allodynia: a randomised, double-blind, placebo-controlled clinical

- trial. *Pain*. 2007 Dec 15;133(1-3):210-20. PMID: 17997224.
15. Pini LA, Guerzoni S, Cainazzo MM, et al. Nabilone for the treatment of medication overuse headache: results of a preliminary double-blind, active-controlled, randomized trial. *J Headache Pain*. 2012 Nov;13(8):677-84. doi: 10.1007/s10194-012-0490-1. PMID: 23070400.
 16. Rintala DH, Fiess RN, Tan G, et al. Effect of dronabinol on central neuropathic pain after spinal cord injury: a pilot study. *Am J Phys Med Rehabil*. 2010 Oct;89(10):840-8. doi: 10.1097/PHM.0b013e3181f1c4ec. PMID: 20855984.
 17. Rog DJ, Nurmikko TJ, Friede T, et al. Randomized, controlled trial of cannabis-based medicine in central pain in multiple sclerosis. *Neurology*. 2005 Sep 27;65(6):812-9. PMID: 16186518.
 18. Schimrigk S, Marziniak M, Neubauer C, et al. Dronabinol Is a Safe Long-Term Treatment Option for Neuropathic Pain Patients. *Eur Neurol*. 2017;78(5-6):320-9. doi: 10.1159/000481089. PMID: 29073592.
 19. Schubert EA, Johnstone MT, Benson MJ, et al. Medicinal cannabis for Australian patients with chronic refractory pain including arthritis. *Br J Pain*. 2023;17(2):206-17. doi: 10.1177/20494637221147115. PMID: 37057257.
 20. Selvarajah D, Gandhi R, Emery CJ, et al. Randomized placebo-controlled double-blind clinical trial of cannabis-based medicinal product (Sativex) in painful diabetic neuropathy: depression is a major confounding factor. *Diabetes Care*. 2010 Jan;33(1):128-30. doi: 10.2337/dc09-1029. PMID: 19808912.
 21. Serpell M, Ratcliffe S, Hovorka J, et al. A double-blind, randomized, placebo-controlled, parallel group study of THC/CBD spray in peripheral neuropathic pain treatment. *Eur J Pain*. 2014 Aug;18(7):999-1012. doi: 10.1002/j.1532-2149.2013.00445.x. PMID: 24420962.
 22. Skrabek RQ, Galimova L, Ethans K, et al. Nabilone for the treatment of pain in fibromyalgia. *J Pain*. 2008 Feb;9(2):164-73. PMID: 17974490.
 23. Tait J, Erridge S, Holvey C, et al. Clinical outcome data of chronic pain patients treated with cannabis-based oils and dried flower from the UK Medical Cannabis Registry. *Expert Rev Neurother*. 2023;1-11. doi: 10.1080/14737175.2023.2195551. PMID: 37021592.
 24. Toth C, Mawani S, Brady S, et al. An enriched-enrolment, randomized withdrawal, flexible-dose, double-blind, placebo-controlled, parallel assignment efficacy study of nabilone as adjuvant in the treatment of diabetic peripheral neuropathic pain. *Pain*. 2012 Oct;153(10):2073-82. doi: 10.1016/j.pain.2012.06.024. PMID: 22921260.
 25. Turcotte D, Doupe M, Torabi M, et al. Nabilone as an adjunctive to gabapentin for multiple sclerosis-induced neuropathic pain: a randomized controlled trial. *Pain Med*. 2015 Jan;16(1):149-59. doi: 10.1111/pme.12569. PMID: 25288189.
 26. Ueberall MA, Essner U, Silván CV, et al. Comparison of the Effectiveness and Tolerability of Nabiximols (THC:CBD) Oromucosal Spray versus Oral Dronabinol (THC) as Add-on Treatment for Severe Neuropathic Pain in Real-World Clinical Practice: Retrospective Analysis of the German Pain e-Registry. *J Pain Res*. 2022a;15:267-86. doi: 10.2147/JPR.S340968. PMID: 35140513.
 27. Ueberall MA, Vila Silvan C, Essner U, et al. Effectiveness, Safety, and Tolerability of Nabiximols Oromucosal Spray vs Typical Oral Long-Acting Opioid Analgesics in Patients with Severe Neuropathic Back Pain: Analysis of 6-Month Real-World Data from the German Pain e-Registry. *Pain Med*. 2022b;23(4):745-60. doi: 10.1093/pm/pnab263. PMID: 34480564.
 28. Vela J, Dreyer L, Petersen KK, et al. Cannabidiol treatment in hand osteoarthritis and psoriatic arthritis: a randomized, double-blind placebo-controlled trial. *Pain*. 2021 Jun 1;163(6):1206-14. PMID: 34510141.
 29. Vigil JM, Stith SS, Adams IM, et al. Associations between medical cannabis and prescription opioid use in chronic pain patients: A preliminary cohort study. *PLoS ONE*. 2017;12(11):e0187795. doi: 10.1371/journal.pone.0187795. PMID: 29145417.

30. Ware MA, Wang T, Shapiro S, et al. Cannabis for the Management of Pain: Assessment of Safety Study (COMPASS). *J Pain*. 2015 Dec;16(12):1233-42. doi: 10.1016/j.jpain.2015.07.014. PMID: 26385201.
31. Wissel J, Haydn T, Muller J, et al. Low dose treatment with the synthetic cannabinoid Nabilone significantly reduces spasticity-related pain : a double-blind placebo-controlled cross-over trial. *J Neurol*. 2006 Oct;253(10):1337-41. PMID: 16988792.
32. Xu DH, Cullen BD, Tang M, et al. The Effectiveness of Topical Cannabidiol Oil in Symptomatic Relief of Peripheral Neuropathy of the Lower Extremities. *Curr Pharm Biotechnol*. 2020;21(5):390-402. doi: 10.2174/1389201020666191202111534. PMID: 31793418.
33. Zajicek JP, Hobart JC, Slade A, et al. Multiple sclerosis and extract of cannabis: results of the MUSEC trial. *J Neurol Neurosurg Psychiatry*. 2012 Nov;83(11):1125-32. doi: 10.1136/jnnp-2012-302468. PMID: 22791906.
34. Zubcevic K, Petersen M, Bach FW, et al. Oral capsules of tetra-hydro-cannabinol (THC), cannabidiol (CBD) and their combination in peripheral neuropathic pain treatment. *Eur J Pain*. 2022doi: 10.1002/ejp.2072. PMID: 36571471.

Appendix D. Literature Flow Diagram

Figure D-1. Literature flow diagram



Note: Numbers in parentheses indicate all records identified up to early July, 2023. Numbers in brackets indicate number of records identified since the previous surveillance report.

Appendix E. Results

Individual Study Summary Tables

Tables E-1 through E-5 present details and results for primary outcomes, serious adverse events, and withdrawals due to adverse events for each included study. Tables E-1 through E-3 provide information for randomized controlled trials and are organized by their respective ratio of tetrahydrocannabinol to cannabidiol. Table E-4 includes details for studies of other cannabinoids, and Table E-5 presents details of observational studies.

Table E-1. Comparable THC to CBD ratio study primary outcomes

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Blake, 2006 Moderate RCT Inflammatory arthritis- rheumatoid arthritis	A: 2.7 mg THC/2.5 mg CBD/100 mcl oromucosal spray, mean dose 5.4 sprays/day (31) B: Placebo (27) 5 weeks Whole plant extracted	Pain severity (mean [SD NR] 0 to 10 NRS scale): 3.1 vs. 4.1, MD -1.04 ^b (95% CI -1.9 to -0.18)	SAE: 0/31 (0%) vs. 2/27 (7.41%), RR 0.18 (95% CI 0.01 to 3.49) WAE: 0/31 (0%) vs. 3/27 (11.11%), RR 0.13 (95% CI 0.01 to 2.32)	Function (mean [SD NR] 0 to 10 28-Joint Disease Activity Score scale): 5 vs. 5.9, MD -0.76 ^c (95% CI -1.23 to -0.28)
Langford, 2013 Low RCT Neuropathic pain- multiple sclerosis	A: 2.7 mg THC/2.5 mg CBD/100 mcl oromucosal spray, mean dose 8.8 sprays/day (167) B: Placebo (172) 15 weeks Whole plant extracted	Pain response ≥30% (NRS scale): 83/167 (49.75%) vs. 77/172 (44.77%), RR 1.11 (95% CI 0.89 to 1.39) Pain severity (mean [SD] 0 to 10 NRS scale): 4.54 (2.24) vs. 4.73 (2.26), MD -0.19 (SE 0.24) (95% CI -0.67 to 0.29)	WAE: 15/167 (8.98%) vs. 12/172 (6.98%), RR 1.29 (95% CI 0.62 to 2.67)	Pain interference (0 to 10 BPI-SF scale): Treatment difference -0.12, p=0.56 Function (0 to 100 SF-36 Physical Functioning scale): Treatment difference -0.45, p=0.785
Lynch, 2014 High RCT (crossover) Neuropathic pain- chemotherapy induced	A: THC/CBD oromucosal spray (dose NR), mean dose 8 sprays/day (8) B: Placebo (8) 4 weeks Whole plant extracted	Pain severity (mean, 0 to 10 NRS-PI scale): 6 (95% CI 6.98 to 5.02) vs. 6.38 (95% CI 5.67 to 7.09)	SAE: 0/8 (0%) vs. 0/8 (0%), RR 1.00 (95% CI 0.02 to 45.13) WAE: 0/8 (0%) vs. 0/8 (0%), RR 1.00 (95% CI 0.02 to 45.13)	Function (mean [SD] 0 to 100 SF-36 Physical Functioning scale): 35.5 (9.19) vs. 46.5 (8.5), MD -11 (4.43) (95% CI -20.49 to -1.51)

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Nurmikko, 2007 Moderate RCT Neuropathic pain- mixed	A: 2.7 mg THC/2.5 mg CBD/100 mcl oromucosal spray, mean dose 10.9 sprays/day (63) B: Placebo (62) 5 weeks Whole plant extracted	Pain response $\geq 30\%$ (NRS scale): 16/73 (25.4%) vs. 9/62 (14.52%), RR 1.75 (95% CI 0.84 to 3.66) Pain severity (mean [SD NR] 0 to 10 NRS scale): 5.82 vs. 6.68, treatment difference -0.96 (95% CI -1.59 to -0.32)	SAE: 1/63 (1.6%) vs. 0/62 (0%), RR 2.95 (95% CI 0.12 to 71.13) WAE: 11/63 (17.46%) vs. 2/62 (3.23%), RR 5.41 (95% CI 1.25 to 23.43)	Function (0 to 70 Pain Disability Index scale): MD -5.85 (95% CI -9.62 to -2.09)
Rog, 2005 Moderate RCT Neuropathic pain- multiple sclerosis	A: 2.7 mg THC/2.5 mg CBD/100 mcl oromucosal spray, mean dose 9.6 sprays/day (34) B: Placebo (32) 5 weeks Whole plant extracted	Pain severity (mean [95% CI] 0 to 10 NRS scale): 3.85 (3.13 to 4.58) vs. 4.96 (4.19 to 5.72), treatment difference -1.25 (95% CI -2.11 to -0.39)	SAE: 0/34 (0%) vs. 0/32 (0%), RR 0.94 (95% CI 0.02 to 46.16) WAE: 2/34 (5.88%) vs. 0/32 (0%), RR 4.71 (95% CI 0.23 to 94.58)	NR
Selvarajah, 2010 High RCT Neuropathic pain- diabetic neuropathy	A: 2.7 mg THC/2.5 mg CBD/100 mcl oromucosal spray, mean dose 7 sprays/day ^d (15) B: Placebo (14) 12 weeks Whole plant extracted	Pain severity (mean [SD] 0 to 100 NPS scale): 51.6 (21.9) vs. 51.9 (24.1), MD -0.3 (SE 8.54) (95% CI -17.83 to 17.23)	NR	Function (mean [SD] 0 to 100 SF-36 Physical Functioning scale): 30.5 (16.6) vs. 36.5 (27.9), MD 6 (SE 8.5) (95% CI -11.35 to 23.35)
Serpell, 2014 Moderate RCT Neuropathic pain- mixed	A: 2.7 mg THC/2.5 mg CBD/100 mcl oromucosal spray, mean dose 8.9 sprays/day (128) B: Placebo (118) 15 weeks Whole plant extracted	Pain response $\geq 30\%$ (NRS scale): 34/123 (27.64%) vs. 19/117 (16.24%), RR 1.7 (95% CI 1.03 to 2.91) Pain severity (mean [SE NR] 0 to 10 NRS scale): Mean reduction -0.34 (0.23) (95% CI -0.79 to 0.11)	SAE: 10/128 (7.81%) vs. 7/118 (6%), RR 1.32 (95% CI 0.52 to 3.35) WAE: 25/128 (19.53%) vs. 25/118 (21.19%), RR 0.92 (95% CI 0.56 to 1.51)	Pain interference (0 to 10 BPI-SF scale): Treatment difference -0.32 (SE 0.241) (95% CI -0.8 to 0.15)

Abbreviations: BPI-SF = brief pain inventory-short form; CBD = cannabidiol; CI = confidence interval; MD = mean difference; NPS = neuropathic pain scale; NR = not reported; NRS = numeric rating scale; NRS-PI = numeric rating scale for pain intensity; SAE = serious adverse events; SD = standard deviation; SE = standard error; SF-36 = short form-36; THC = tetrahydrocannabinol; RCT = randomized controlled trial; RR = risk ratio; WAE = withdrawal due to due adverse events.

^aOther serious adverse events (i.e., psychosis and cannabis use disorder) not reported in any study.

^bDifference in median differences.

^cDifference in mean differences.

^dMean sprays calculated by systematic review team.

Table E-2. High THC to CBD ratio study primary outcomes

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Chaves, 2020 Low RCT Fibromyalgia	A: 1.2 mg THC/0.02 mg CBD sublingual drops, mean 3.6 drops/day (8) B: Placebo (9) 8 weeks Whole plant extracted	Pain severity (mean [SD] 0 to 10 FIQ scale): 3.75 (2.49) vs. 7.67 (1.84), MD -3.92 (1.05) (95% CI -6.17 to -1.68)	WAE: 0/8 (0%) vs. 0/9 (0%), RR 1.11 (95% CI 0.02 to 50.43)	Function (mean [SD] 0 to 10 FIQ scale): 5.83 (2.02) vs. 4.07 (2.25), MD 1.76 (1.04) (95% CI -0.46 to 3.98)
de Vries, 2017 Moderate RCT Visceral pain- chronic pancreatitis and postsurgical abdominal pain	A: THC oral tablet (Dronabinol), range 15 to 24 mg/day (30) B: Placebo (32) 7 weeks Synthetic	Pain severity (mean [SD] 0 to 10 VAS scale): 2.4 (2.28) vs. 3.5 (2.42), MD -1.1 (SE 0.68) (95% CI -2.46 to 0.26)	WAE: 7/30 (23.33%) vs. 2/32 (6.25%), RR 3.73 (95% CI 0.84 to 16.57)	NR
Frank, 2008 Moderate RCT (crossover) Neuropathic pain	A: THC oral capsule (Nabilone), max dose 2 mg/day (48) B: Dihydrocodeine 30 mg, max dose 240 mg/day (48) 6 weeks Synthetic	Pain severity (mean [SD NR] 0 to 100 VAS scale): Treatment effect 5.7 (95% CI 0.5 to 10.9)	SAE: 0/48 (0%) vs. 0/48 (0%), RR 1.00 (95% CI 0.02 to 49.39) WAE: 2/48 (4%) vs. 6/48 (12.5%), RR 0.33 (95% CI 0.07 to 1.57)	Function (mean [SD NR] 0 to 100 SF-36 Physical Functioning scale): Treatment effect 10.8 (95% CI 2.3 to 19.2)

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Pini, 2012 Low RCT (crossover) Headache- medication overuse headache	A: THC 0.5 mg oral capsule (Nabilone) daily (26) B: Ibuprofen 400 mg/day (26) 8 weeks Synthetic	Pain severity (mean [SD] 0 to 10 VAS scale): 5.55 (2.5) vs. 6.75 (2.4), MD -1.2 (0.68) (95% CI -2.57 to 0.17)	WAE: 1/30 (3.33%) vs. 1/30 (3.33%), RR 1.00 (95% CI 0.07 to 15.26)	NR
Rintala, 2010 High RCT (crossover) Neuropathic pain- spinal cord injury	A: THC 5 mg oral capsule (Dronabinol), max dose 20 mg/day (7) B: Diphenhydramine 25 mg, max dose 75 mg/day (5) 47 weeks Synthetic	Pain severity (mean [SD NR] 0 to 10 BPI scale): 5.8 vs. 5.8	SAE: 1/7 (14.29%) vs. 1/5 (20%), RR 0.71 (95% CI 0.06 to 8.91) WAE: 1/7 (14.29%) vs. 0/5 (0%), RR 2.25 (95% CI 0.11 to 46.13)	NR
Schimrigk, 2017 Low RCT Neuropathic pain- multiple sclerosis	A: THC 2.5 mg oral capsule (Dronabinol), mean dose 13 mg/day (124) B: Placebo (116) 16 weeks Synthetic	Pain severity (mean [SD] 0 to 10 NRS scale): 4.48 (2.04) vs. 4.92 (2.04), MD NR, p=0.676	SAE: 12/124 (9.68%) vs. 7/116 (6.03%), RR 1.53 (95% CI 0.63 to 3.76) WAE: 19/124 (15.32%) vs. 12/116 (10.34%), RR 1.48 (95% CI 0.75 to 2.91)	NR
Skrabek, 2008 Moderate RCT Fibromyalgia	A: THC 0.5 mg oral capsule (Nabilone), endpoint dose 2 mg/day (15) B: Placebo (18) 4 weeks Synthetic	Pain severity (mean [SD NR] 0 to 10 VAS scale): 4.8 vs. 5.6, MD -1.43, p<0.05	SAE: 0/15 (0%) vs. 0/18 (0%), RR 1.19 (95% CI 0.02 to 56.54) WAE: 1/20 (5%) vs. 1/20 (5%), RR 1.00 (95% CI 0.07 to 14.90)	NR
Toth, 2012 Low RCT Neuropathic pain- diabetic neuropathy	A: THC 0.5 mg oral capsule (Nabilone), max dose 4 mg/day (13) B: Placebo (13) 5 weeks Synthetic	Pain response ≥30% (NRS scale): 11/13 (84.62%) vs. 5/13 (38.46%), RR 2.2 (95% CI 1.06 to 4.55) Pain severity (mean [SD] 0 to 10 NRS scale): 3.5 (1.3) vs. 5.4 (1.7), MD -1.9 (0.59) (95% CI -3.13 to -0.68)	NR	Pain interference (mean [SD] 0 to 10 MBPI scale): 2.5 (1.6) vs. 3.6 (0.9), MD -1.1 (0.51) (95% CI -2.15 to -0.05)

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Turcotte, 2015 Moderate RCT Neuropathic pain- multiple sclerosis	A: THC 0.5 mg oral capsule (Nabilone), max dose 2 mg/day (8) B: Placebo (7) 9 weeks Synthetic	Pain severity (mean [SD NR] 0 to 100 VAS scale): 35 vs. 57 ^b	SAE: 0/8 (0%) vs. 0/7 (0%), RR 0.89 (95% CI 0.02 to 39.84) WAE: 1/8 (12.5%) vs. 0/7 (0%), RR 2.67 (95% CI 0.13 to 56.63)	Pain interference (mean [SD NR] 0 to 100 VAS impact scale): 41 vs. 40 ^b
Wissel, 2006 High RCT (crossover) Neuropathic pain- multiple sclerosis	A: THC 0.5 mg oral capsule (Nabilone), endpoint dose 1 mg/day (13) B: Placebo (13) 4 weeks Synthetic	Pain severity (median [SD NR] 11 Point Box Test): 4 vs. 6, p<0.05	WAE: 2/13 (15.38%) vs. 0/13 (0%), RR 5.00 (95% CI 0.26 to 95.02)	NR
Zajicek, 2012 Moderate RCT Neuropathic pain- multiple sclerosis	A: THC 2.5 mg capsule, max dose 25 mg/day (143) B: Placebo (134) 12 weeks Whole plant extracted	Pain severity (mean [SD] 0 to 10 CRS scale): 4.1 (2.9) vs. 4.7 (3.0), MD -0.6 (95% CI -1.3 to 0.1)	SAE: 7/143 (4.9%) vs. 3/134 (2.24%), RR 2.19 (95% CI 0.58 to 8.28) WAE: 30/143 (20.98%) vs. 9/134 (6.72%), RR 3.12 (95% CI 1.54 to 6.33)	NR

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Zubcevic, 2022 Low RCT Peripheral neuropathic pain	A: THC 2.5 mg capsule (dronabinol), max dose 25 mg/day (28) B: CBD 5 mg capsule (unknown if synthetic or plant-derived), max dose 50 mg/day (27) C: CBD/THC capsule, max dose 50 mg CBD (unknown of synthetic or plant-derived)/25 mg THC (dronabinol)/day (30) D: Placebo (30)	Pain response $\geq 30\%$ (NRS scale): 12/28 (42.86%) vs. 9/27 (33.34%) vs. 18/30 (60.00%) vs. 17/30 (56/67%), RR (95% CI) A vs. B: 1.29 (0.65 to 2.55) A vs. C: 0.71 (0.43 to 1.20) A vs. D: 0.76 (0.45 to 1.28) B vs. C: 0.56 (0.30 to 1.02) B vs. D: 0.59 (0.32 to 1.09) C vs. D: 1.06 (0.69 to 1.62) Pain severity change from baseline (mean [95% CI] 0 to 10 NRS scale): -1.4 (-2.2 to -0.7) vs. -0.6 (-1.2 to 0.1) vs. -1.9 (-2.7 to -1.2) vs. -1.9 (-2.7 to -1.0)	SAE: 0/28 (0%) vs. 0/27 (0%) vs. 1/30 (3.3%) vs. 0/30 (0%), RR (95% CI) A vs. B: 0.96 (0.02 to 47.01) A vs. C: 0.36 (0.02 to 8.40) A vs. D: 1.07 (0.02 to 52.14) B vs. C: 0.37 (0.02 to 8.70) B vs. D: 1.11 (0.02 to 53.97) C vs. D: 3.00 (0.13 to 70.83) WAE: 1/28 (3.57%) vs. 2/27 (7.41%) vs. 4/30 (13.33%) vs. 0/30 (0%), RR (95% CI) A vs. B: 0.48 (0.05 to 5.01) A vs. C: 0.27 (0.03 to 2.25) A vs. D: 3.21 (0.14 to 75.62) B vs. C: 0.56 (0.11 to 2.80) B vs. D: 5.54 (0.28 to 110.42) C vs. D: 9.00 (0.51 to 160.18)	Pain interference (mean [SD] 0 to 10 Pain Impact on Daily Activities Scale): MD (95% CI) A vs. D: 0.36 (-1.19 to 1.91) B vs. D: 1.24 (-0.32 to 2.81) C vs. D: 0.89 (-0.64 to 2.42)

Abbreviations: BPI = brief pain inventory; CBD = cannabidiol; CI = confidence interval; CRS = category rating scale; FIQ = fibromyalgia impact questionnaire; MBPI = modified brief pain inventory; MD = mean difference; NR = not reported; NRS = numeric rating scale; RCT = randomized controlled trial; SAE = serious adverse events; SD = standard deviation; SE = standard error; THC = tetrahydrocannabinol; RR = risk ratio; VAS = visual analog scale; WAE = withdrawal due to due adverse events.

^aOther serious adverse events (i.e., psychosis and cannabis use disorder) not reported in any study.

^bEstimated from graph.

Table E-3. Low THC to CBD ratio study primary outcomes

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Bassat, 2023 High RCT (crossover) Chronic pain-mixed	A: 3 drops (2.5 mg THC/15 mg CBD) sublingual oil, max dose 18 drops per day (15 mg THC/90 mg CBD) (7) B: 3 drops (2.5 mg THC/15 mg CBD) ≥97% purified sublingual oil, max dose 18 drops per day (15 mg THC/90 mg CBD) (5) C: Placebo (9) 16 weeks Whole plant extracted	NR	SAE: 6/9 (67%) vs. 4/6 (67%) vs. 3/10 (30%) WAE: 1/4 (25%) vs. 1/5 (20%) vs. 1/6 (16.7%)	NR
Vela, 2021 Moderate RCT Musculoskeletal - hand osteoarthritis and psoriatic arthritis	A: CBD oral tablet (20 to 30 mg/day) (68) B: Placebo (61) 12 weeks Synthetic CBD	Pain response ≥30% (VAS 0 to 100 scale): 27/68 (39.7%) vs. 24/61 (39.3%), RR 1.01 (95% CI 0.66 to 1.55) Pain severity (0 to 100 VAS scale): mean NR, MD 0.23 (95% CI -9.41 to 9.9)	SAE: 2/58 (3.4%) vs. 2/61 (3.3%), RR 1.05 (95% CI 0.15 to 7.22) WAE: 0/70 (0%) vs. 2/66 (3%), RR 0.19 (95% CI 0.01 to 3.86)	Function/disability (0 to 3 HAQ-DI scale): mean NR, MD 0.03 (95% CI -0.11 to 0.18)
Xu, 2020 High RCT (crossover) Neuropathic pain- mixed	A: CBD cream (250 mg/3 oz) up to 4 times daily (15) B: Placebo (14) 4 weeks Whole plant extracted	Pain severity (mean [SD] 0 to 10 NPS scale): 3.33 (2.02) vs. 5.55 (2.81), MD -2.22 (95% CI -4.07 to -0.37)	SAE: 0/15 (0%) vs. 0/14 (0%)	NR

Abbreviations: CBD = cannabidiol; CI = confidence interval; HAQ-DI = Health Assessment Questionnaire Disability Index; MD = mean difference; NPS = neuropathic pain scale; NR = not reported; RCT = randomized controlled trial; SAE = serious adverse event; SD = standard deviation; THC = tetrahydrocannabinol.

^aOther serious adverse events (i.e., psychosis and cannabis use disorder) not reported in any study.

Table E-4. Other cannabinoids study primary outcomes

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Eibach, 2020 Moderate RCT (crossover) Neuropathic pain- HIV associated	A: CBDV oral solution (50 mg/mL) 400 mg/day (16) B: Placebo (16) 4 weeks Whole plant extracted	Pain response $\geq 30\%$ (NRS scale): 6/16 (37.5%) vs. 13/16 (81.25%), RR NR Pain severity (mean [SD] 0 to 10 NRS scale): 2.74 (1.47) vs. 3.67 (2.62), MD -0.62 (95% CI -0.27 to 1.51)	SAE: 1/16 (6.25%) vs. 0/16 (0%), RR 3.00 (95% CI 0.13 to 68.57) WAE: 1/16 (6.25%) vs. 0/16 (0%), RR 3.00 (95% CI 0.13 to 68.57)	Pain interference (0 to 10 BPI-SF scale): MD -0.35 (95% CI -1.36 to 0.43)

Abbreviations: BPI-SF = Brief Pain Inventory – Short Form; CBDV = cannabidivarin; CI = confidence interval; MD = mean difference; NR = not reported; RCT = randomized controlled trial; RR = risk ratio; SAE = serious adverse event; SD = standard deviation; WAE = study withdrawals due to adverse events.

^aOther serious adverse events (i.e., psychosis and cannabis use disorder) not reported in any study.

Table E-5. Observational study primary outcomes

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Bestard, 2011 Moderate Prospective cohort Neuropathic pain- mixed	A: THC oral capsule (Nabilone), mean dose 3.05 mg/day (49) B: Gabapentin, mean dose 2,295.5 mg/day (52) C: Gabapentin + THC capsule, mean dose NR + 3.02 mg/day (55) 6 months Synthetic	Pain intensity (mean [SD] 0 to 100 VAS scale): 28.0 (10.5) vs. 33.8 (11.6) vs. 33.1 (20.2), MD -5.8 (95% CI -10.18 to -1.42) for A vs. B, -5.1 (95% CI -11.48 to 1.28) for A vs. C	SAE: 0/49 (0%) vs. 0/52 (0%) vs. 0/55 (0%) WAE: 5/49 (10%) vs. 12/52 (23%) vs. 5/55 (9%), RR 0.44 (95% CI 0.17 to 1.16 for A vs. B, RR 1.12 (95% CI 0.34 to 3.65) for A vs. C, RR 2.54 (95% CI 0.96 to 6.71) for B vs. C	Pain interference (mean [SD] 0 to 10 BPI scale): 4.5 (2.3) vs. 4.6 (2.2) vs. 4.5 (2.2), MD -0.1 (95% CI -0.99 to 0.79) for A vs. B, MD 0.00 (95% CI -0.88 to 0.88) for A vs. C Function (mean [SD] 0 to 100 SF-36 scale ^a): 48.3 (27.2) vs. 46.5 (25.1) vs. 43.7 (26.4), MD 1.80 (95% CI -8.53 to 12.13) for A vs. B, MD 4.60 (95% CI - 5.83 to 15.03) for A vs. C
Campbell, 2018 Moderate	A: Self-reported frequent cannabis use of ≥20 days/mo B: No cannabis use Overall N Baseline: 1,514 4-year followup: 1,217 Groups unclear 4 years Unclear THC concentration; patient- driven choice	A vs. B (reference) Pain intensity (Adjusted mean [SE]; BPI, 0-10 scale): 5.2 (0.14) vs. 4.9 (0.03); Beta: 0.37 (95% CI, -0.23 to 1.10), p=0.20	NR	A vs. B Pain Interference (Adjusted mean [SE]; BPI pain interference, 0 to 10 scale): 5.2 (0.19) vs. 5.4 (0.04); Beta: -0.63 (95% CI, -1.46 to 0.19), p=0.13
Gruber, 2021 High Prospective cohort Mixed (primarily musculoskeletal)	A: THC/CBD: Medicinal cannabis program, mean dose THC 13.3 mg/day, CBD 28.9 mg/day (37) B: Usual care, dose NA (9) 12 weeks Mixed cannabis products	Pain intensity (mean [SD] 0 to 100 VAS scale): 34.07 (22.36) vs. 48.78 (30.42); MD -14.71 (95% CI, -32.71 to 3.29)	NR	A vs. B Function (mean [SD], 0 to 10 PDI scale): 18.13 (12.26) vs. 19.22 (12.73); MD -1.09 (95% CI -10.33 to 8.16) SF-36 Function (mean [SD], 0 to 100 scale ^a): 70.00 (22.87) vs. 69.44 (26.98); MD 0.56 (95% CI -17.17 to 18.29)

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Lee, 2021 ^b Moderate Matched cohort NR	A: Chronic opioid users authorized to use medical cannabis in Canada (5,373) B: Controls who did not receive authorization for medical cannabis in Canada (5,373) 20 months Unknown THC concentration; patient-driven choice	NR	NR	NR
Merlin, 2019 ^b High Prospective cohort Chronic non-cancer pain (HIV)	A: Daily or weekly use of marijuana (55) B: Monthly or 1-2 times a month use of marijuana (65) C: No use (313) 52 weeks Unknown THC concentration; patient-driven choice	NR	NR	NR

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Schubert, 2023 High Prospective cohort Chronic refractory pain, including arthritis	A: CBD only (174) B: CBD dominant (two fold ratio increase in main cannabinoid) (37) C: Balanced (113) D: THC dominant (THC only) (27)	A vs. B vs. C vs. D Pain intensity Score first, mean (SD): 5.9 (2.2) vs. 6.2 (1.8) vs. 6.1 (2.1) vs. 5.6 (2.1) Score final, mean (SD): 5.6 (2.2) vs. 6.1 (2) vs. 5.7 (2) vs. 5.7 (2.2) Improved, n (%): 37 (21.3) vs. 8 (21.6) vs. 27 (23.9) vs. 5 (13.5) Pain impact Raw score first, mean (SD): 31.8 (9.7) vs. 34.9 (10.2) vs. 33.9 (9.5) vs. 30.6 (10.5) Raw score final, mean (SD): 30.9 (9.4) vs. 34.1 (9.8) vs. 32.3 (9.2) vs. 31.4 (9.1) Improved, n (%): 72 (41.4) vs. 14 (37.8) vs. 47 (41.6) vs. 13 (35.1)	NR	A vs. B. vs. C vs. D Pain interference T score first, mean (SD): 64.2 (7.8) vs. 66.6 (7.6) vs. 65.4 (7.4) vs. 63.3 (8.1) T score final, mean (SD): 63.4 (7.4) vs. 65.5 (6.7) vs. 63.7 (7.6) vs. 63.4 (8.1) Improved, n (%): 64 (36.8) vs. 14 (37.8) vs. 49 (43.4) vs. 13 (35.1) Physical function T score first, mean (SD): 37.3 (7.5) vs. 35.7 (9) vs. 35.5 (7.1) vs. 38.5 (8.6) T score final, mean (SD): 37.4 (6.7) vs. 35.1 (7.3) vs. 35.8 (6) vs. 37.3 (8.3) Improved, n (%): 57 (32.8) vs. 11 (29.7) vs. 40 (35.4) vs. 12 (32.4)
Tait, 2023 Moderate Prospective cohort Chronic non-cancer pain	A: A. Cannabis-based sublingual/oral medium-chain triglyceride-based oil containing CBD and THC (348) B. Inhaled dried cannabis flower containing trace CBD and THC (36) C. A + B (377) Dried flower concentration: 20% THC, 0% (trace) CBD Cannabis-based oils ranged from all THC, all CBD, and ratios ranging from 1:1 to 1:20	A vs. B vs. C Pain, VAS score (scale 0-10; median, IQR) 1 month: 7.00 (5.00-8.00) vs. 5.50 (3.00-6.75) vs. 6.00 (5.00- 7.75) 3 months: 6.00 (4.00-7.00) vs. 5.00 (3.00-7.00) vs. 6.00 (4.00- 7.00) 6 months: 6.00 (3.00-7.25) vs. 3.00 (1.50-5.00) vs. 5.00 (3.00- 7.00)	NR	A vs. B. vs. C BPI-Interference score (scale 0-10; median, IQR) 1 month: 5.86 (3.64-7.36) vs. 5.14 (2.21-5.93) vs. 5.79 (3.57-7.43) 3 months: 5.29 (3.43-6.82) vs. 4.14 (2.43-6.29) vs. 5.00 (2.86-6.93) 6 months: 4.71 (3.14-6.43) vs. 2.14 (1.50-3.64) vs. 4.50 (2.64-6.64) Likelihood of improvement, adjusted OR A vs. B: 1.61 (95% CI 0.21 to 12.18); A vs. C: 1.90 (95% CI 0.68 to 5.29)

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events ^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Ueberall, 2022a Moderate Retrospective cohort Peripheral neuropathic pain	A: Nabiximols as an add-on treatment; 16.6 (SD 6.5) mg THC/15.4 (SD 4.1) mg CBD/day (337) B: Dronabinol as an add-on treatment; 17.2 (SD 7.6) mg THC/day (337) 24 weeks Whole plant extracted vs. synthetic	A vs. B Pain intensity index (VAS 0-100 scale) mean relative change (improvement) rates at week 24 83.4% vs. 75.9%, p<0.001 Pain intensity index (VAS 0-100 scale, converted to 0-10) mean difference: 3.50 (95% CI 1.6 to 5.4)	NR	A vs. B Pain-related disabilities (VAS 0-100 scale) mean relative change (improvement) rates at week 24 76.0% vs. 68.3%, p<0.001
Ueberall, 2022b Moderate Retrospective cohort Peripheral neuropathic back pain- mixed	A: 2.7 mg THC/2.5 mg CBD/100 mcl oromucosal spray, mean dose 16.7 mg THC/15.5 mg CBD/day (655) B: Long-acting opioid, MME 69.4 mg/day 24 weeks Whole plant extracted and long-acting opioid	Pain intensity index (mean relative change from baseline at week 24, 0 to 100 VAS scale): -72.3% (SD 30.5) vs. -49.2% (SD 39.9) Pain intensity index (VAS 0-100 scale, converted to 0-10) mean difference: 9.90 (95% CI 8.05 to 11.75)	WAE: 7.9% vs. 29.3%, RR 0.27 (95% CI 0.20 to 0.36)	Pain-related disabilities (mean relative change [improvement] rates at week 24, 0 to 100 VAS scale): -66.1 (28.7) vs. -42.9 (34.5), p<0.001

Author, Year Risk of Bias Study Design Pain Condition	Comparison (n) Followup Duration Derivative	Primary Pain Outcomes (Response, Severity)	Serious Adverse Events and Withdrawals Due to Adverse Events^a	Other Primary Outcomes (Function/Disability, Pain Interference)
Vigil, 2017 ^b High Preliminary historical cohort Mixed musculoskeletal pain	A: THC/CBD: Participation in New Mexico Medical Cannabis Program (37) B: Not participating in medical marijuana program and not using cannabis (29) 21 months Unknown THC concentration	NR	NR	NR
Ware, 2015 High Prospective cohort Chronic non-cancer pain	A: THC 12.5 +/- 1.5% herbal cannabis, median dose 2.5 g/day (215) B: Usual care (216) 13 months Whole plant non- extracted	NR	SAE: 28/215 (13%) vs. 42/216 (19.4%), RR 0.67 (95% CI 0.43 to 1.04) WAE: 10/215 (4.65%) vs. NR (assumed 0)	NR

Abbreviations: BPI = brief pain inventory; CBD = cannabidiol; CI = confidence interval; IQR = interquartile range; MD = mean difference; NA = not applicable; NR = not reported; OR = odds ratio; PDI = Pain Disability Index; SAE = serious adverse events; SD = standard deviation; SF-36= short form-36; THC = tetrahydrocannabinol; VAS = visual analog scale; WAE = withdrawal due to due adverse events.

^aHigher scores indicate better outcomes.


^bOnly included outcome reported was opioid-use.

Meta-Analysis Results

Comparable THC to CBD Ratio Studies


Pooled results and the forest plot for the sensitivity analysis conducted for improvement in pain severity are available upon request by emailing ahmedaz@ohsu.edu.

Figure E-1. Change in pain severity with comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)




Abbreviations: CI = confidence interval; IA = inflammatory arthritis; NPP = neuropathic pain; NR = not reported; PL = profile likelihood; SD = standard deviation.

Figure E-2. Proportion of patients with pain response ($\geq 30\%$ improvement) with comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)




Abbreviations: CI = confidence interval; NPP = neuropathic pain; PL = profile likelihood.

Figure E-3. Overall function: comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)




Abbreviations: CI = confidence interval; IA = inflammatory arthritis; NPP = neuropathic pain; NR = not reported; PL = profile likelihood; SD = standard deviation.

Figure E-4. Any adverse events for comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)




Abbreviations: CI = confidence interval; NPP = neuropathic pain; PL = profile-likelihood.

Figure E-5. Serious adverse events for comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)




Abbreviations: CI = confidence interval; IA = inflammatory arthritis; NPP = neuropathic pain; PL = profile-likelihood.

Figure E-6. Withdrawal due to adverse events for comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)




Abbreviations: CI = confidence interval; IA = inflammatory arthritis; NPP = neuropathic pain; PL = profile-likelihood.

Figure E-7. Dizziness for comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)




Abbreviations: CI = confidence interval; IA = inflammatory arthritis; NPP = neuropathic pain; PL = profile likelihood.

Figure E-8. Nausea for comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)



Abbreviations: CI = confidence interval; IA = inflammatory arthritis; NPP = neuropathic pain; PL = profile likelihood.


Figure E-9. Sedation for comparable THC to CBD ratio versus placebo (short-term, 4 weeks to 6 months followup)



Abbreviations: CI = confidence interval; IA = inflammatory arthritis; NPP = neuropathic pain; PL = profile likelihood.

High THC to CBD Ratio Studies


Figure E-10. Change in pain severity with high THC ratio versus placebo (short-term, 4 weeks to 6 months followup)



Abbreviations: CBD = cannabidiol; CI = confidence interval; EP = end-point; FM = fibromyalgia; NA = not applicable; NPP = neuropathic pain; PD = plant-derived; PL = profile likelihood; SD = standard deviation; TD = total dose; T/C = THC/CBD; THC = tetrahydrocannabinol; VP = visceral pain.

^aNamisol[®] is a purified, plant-based product, but grouped with synthetic dronabinol because they are chemically identical.

Figure E-11. Stratified results on pain severity of RCTs using dronabinol and nabilone (short term, 4 weeks to 6 months followup)



Abbreviations: CBD = cannabidiol; CI = confidence interval; EP = end point; FM = fibromyalgia; NPP = neuropathic pain; NR = not reported; PL = profile likelihood; RCT = randomized controlled trials; SD = standard deviation; TD = total dose; THC = tetrahydrocannabinol; VP = visceral pain.

^aNamisol[®] is a purified, plant-based product, but grouped with synthetic dronabinol because they are chemically identical.


Table E-6. Interaction effect of RCTs assessing synthetic cannabinoids: nabilone versus dronabinol

Group Difference	Coefficient	Standard Error	t-Test	p-Value	95% Confidence Interval
Result	-1.29	0.510	-2.53	0.053	-2.60 to 0.022

Table E-7. Interaction effect of RCTs: synthetic versus plant-extracted interventions


Group Difference	Coefficient	Standard Error	t-Test	p-Value	95% Confidence Interval
Result	-0.986	0.85	-1.16	0.272	-2.87 to 0.90

Figure E-12. Overall function for high THC versus placebo (short-term, 1-6 months followup)



Abbreviations: CBD = cannabidiol; CI = confidence interval; NPP = neuropathic pain; NR = not reported; PL = profile likelihood; SD = standard deviation; TD = total dose; THC = tetrahydrocannabinol.


Figure E-13. Withdrawal due to adverse events for high THC versus placebo (short-term, 4 weeks to 6 months followup)



Abbreviations: CBD = cannabidiol; CI = confidence interval; EP = end point; FM = fibromyalgia; NA = not applicable; NPP = neuropathic pain; PL = profile likelihood; TD = total dose; THC = tetrahydrocannabinol; VP = visceral pain.


^aNamisol® is a purified, plant-based product, but grouped with synthetic dronabinol because they are chemically identical.

Figure E-14. Any adverse event for high THC versus placebo (short-term, 4 weeks to 6 months followup)



Abbreviations: CI = confidence interval; NPP = neuropathic pain.


Figure E-15. Dizziness for high THC versus placebo (short-term, 4 weeks to 6 months followup)



Abbreviations: CBD = cannabidiol; CI = confidence interval; NA = not applicable; NPP = neuropathic pain; PL = profile likelihood; THC = tetrahydrocannabinol; VP = visceral pain.


^aNamisol[®] is a purified, plant-based product, but grouped with synthetic dronabinol because they are chemically identical.

Figure E-16. Sedation for high THC versus placebo (short-term, 4 weeks to 6 months followup)



Abbreviations: CI = confidence interval; EP = end point; FM = fibromyalgia; NA = not applicable; NPP = neuropathic pain; PL = profile likelihood; VP = visceral pain.
^aNamisol[®] is a purified, plant-based product, but grouped with synthetic dronabinol because they are chemically identical.

Figure E-17. Nausea for high THC versus placebo (short-term, 4 weeks to 6 months followup)



Abbreviations: CI = confidence interval; NPP = neuropathic pain; VP = visceral pain.
^aNamisol[®] is a purified, plant-based product, but grouped with synthetic dronabinol because they are chemically identical.

Table E-8. Meta-analysis results and sensitivity analysis using the Bartlett's Correction

THC to CBD Ratio	Outcome	N; k Studies	Point Estimate	PL 95% CI	BC 95% CI	I-Squared
Comparable	Pain severity	N=702; k=7	MD -0.54	-0.95 to -0.19	-1.03 to -0.11	39%
Comparable	Pain response ($\geq 30\%$ improvement)	N=733; k=4	RR 1.18	0.93 to 1.71	0.67 to 2.43	36%
Comparable	Function	N=616; k=6	MD -0.42	-0.73 to -0.16	-0.80 to -0.10	32%
Comparable	Adverse events	N=405; k=2	RR 1.19	1.02 to 1.44	0.74 to 2.03	0%
Comparable	SAEs	N=427; k=3	RR 1.18	0.26 to 3.43	0.02 to 35.25	0%
Comparable	WAEs	N=834; k=5	RR 1.14	0.65 to 3.02	0.31 to 6.16	51%
Comparable	Dizziness	N=866; k=6	RR 3.57	2.42 to 5.60	2.15 to 6.62	0%
Comparable	Nausea	N=866; k=6	RR 1.79	1.19 to 2.77	1.06 to 3.32	0%
Comparable	Sedation	N=866; k=6	RR 5.04	2.10 to 11.89	1.41 to 17.29	0%
High	Pain severity	N=742; k=9	MD -1.12	-1.97 to -0.48	-2.08 to -0.40	65%
High (synthetic)	Pain severity	N=448; k=7	MD -0.95	-1.81 to -0.25	-1.95 to -0.13	60%
High (synthetic - dronabinol)	Pain severity	N=348; k=3	MD -0.35	-1.08 to 0.44	-2.21 to 1.54	45%
High (synthetic - nabilone)	Pain severity	N=100; k=4	MD -1.59	-2.49 to -0.82	-2.21 to -0.39	0%
High (plant-derived)	Pain severity	N=294; k=2	MD -1.97	-5.91 to 1.21	-11.33 to 6.53	85%
High	Function	N=unclear; k=3	MD -0.18	-1.25 to 0.77	-2.23 to 1.78	51%
High	WAEs	N=692; k=6	RR 2.21	1.27 to 4.14	0.96 to 5.58	0%
High (synthetic)	WAEs	N=415; k=5	RR 1.75	0.95 to 4.11	0.50 to 8.88	0%
High (synthetic - dronabinol)	WAEs	N=360; k=3	RR 1.77	0.90 to 5.44	0.25 to 24.91	0%
High (synthetic - nabilone)	WAEs	N=55; k=2	RR 1.54	0.14 to 17.71	0.01 to 280.12	0%
High	Any adverse event	N=266; k=2	RR 1.20	0.96 to 1.48	0.42 to 3.36	0%
High	Dizziness	N=637; k=4	RR 3.57	1.30 to 8.32	0.90 to 11.47	78%
High (synthetic)	Dizziness	N=360; k=3	RR 2.52	1.20 to 4.82	0.42 to 12.00	41%
High	Sedation	N=386; k=4	RR 1.60	1.01 to 42.95	0.74 to 5.09	0%
High (synthetic - dronabinol)	Sedation	N=353; k=	RR 1.46	0.88 to 2.42	0.59 to 3.66	0%
High	Nausea	N=353; k=3	RR 2.22	0.90 to 5.05	0.40 to 11.80	0%

Abbreviations: BC = Bartlett's correction; CBD = cannabidiol; CI = confidence interval; MD = mean difference; PL = profile likelihood; RR = risk ratio; SAEs = serious adverse events; THC = tetrahydrocannabinol; WAEs = study withdrawals due to adverse events.

Appendix F. Evidence Tables

Shown in associated Excel files at <https://effectivehealthcare.ahrq.gov/products/plant-based-chronic-pain-treatment/living-review>.

Appendix G. Risk of Bias Assessment

Shown in associated Excel files at <https://effectivehealthcare.ahrq.gov/products/plant-based-chronic-pain-treatment/living-review>.

Appendix H. Details on Strength of Evidence

Table H-1. KQ1 and 2: Cannabinoids to treat chronic pain – comparable THC to CBD ratio

Comparison	Outcome	Number of Studies (N) and Total Participants	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	SOE Grade
Comparable THC to CBD Ratio vs. Placebo	Pain response (≥30% improvement from baseline)	4 RCTs (N=733) ¹⁻⁴	Moderate	Direct	Consistent	Imprecise	Unknown	Potential small effect, not statistically significant, with THC:CBD 38% vs. 31%, RR 1.18 (0.93 to 1.71); I ² =36%	Low
Comparable THC to CBD Ratio vs. Placebo	Pain severity (change)	7 RCTs (N=878) ¹⁻⁷	Moderate	Direct	Consistent	Precise	Unknown	Small benefit with THC:CBD 0 to 10 scale, MD -0.54 (-0.95 to -0.19; I ² =40%) Subgroup analysis removing high risk of bias studies: Moderate benefit MD -0.64 (-1.15 to -0.24)	Moderate
Comparable THC to CBD Ratio vs. Placebo	Function or Disability	6 RCTs (N=616) ^{1-5,7}	Moderate	Direct	Consistent	Precise	Unknown	Small benefit with THC:CBD, MD -0.42, 95% CI -0.73 to -0.16, I ² =32% (scale 0 to 10)	Moderate
Comparable THC to CBD Ratio vs. Placebo	WAEs	5 RCTs (N=834) ^{1,2,4,5,7}	Moderate	Direct	Consistent	Imprecise	Unknown	Failed to demonstrate or exclude a detrimental effect 13% vs. 10%, RR 1.14 (0.65 to 3.02); I ² =51%	Low
Comparable THC to CBD Ratio vs. Placebo	SAEs	3 RCTs (N=429) ^{2,4,5}	Moderate	Direct	Consistent	Imprecise	Unknown	No effect 5.0% vs. 4.3%, RR 1.18 (0.28 to 3.43; I ² =0%)	Low

Comparison	Outcome	Number of Studies (N) and Total Participants	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	SOE Grade
Comparable THC to CBD Ratio vs. Placebo	Dizziness	6 RCTs (N=866) ^{1,2,4-7}	Moderate	Direct	Consistent	Imprecise	Unknown	Large effect with THC:CBD 30% vs. 8%, RR 3.57 (2.42 to 5.60; I ² =0%)	Low
Comparable THC to CBD Ratio vs. Placebo	Nausea	6 RCTs (N=866) ^{1,2,4-7}	Moderate	Direct	Consistent	Imprecise	Unknown	Moderate effect with THC:CBD 14% vs. 7.5% RR 1.79 (1.19 to 2.77; I ² =0%)	Low
Comparable THC to CBD Ratio vs. Placebo	Sedation	6 RCTs (N=866) ^{1,2,4-7}	Moderate	Direct	Consistent	Imprecise	Unknown	Large effect with THC:CBD RR 5.04 (2.10 to 11.89; I ² =0%)	Low

Abbreviations: BPI-SF = Brief Pain Inventory (Short Form); CBD = cannabidiol; CI = confidence interval; KQ = Key Question; MD = mean difference; RCT = randomized controlled trial; RR = risk ratio; SAE = serious adverse event; SOE = strength of evidence; THC = tetrahydrocannabinol; WAE = withdrawal due to adverse event

Table H-2. KQ1 and 2: Cannabinoids to treat chronic pain – high THC to CBD ratio, synthetic THC

Comparison	Outcome	Number of Studies and Total Participants (N)	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	Strength of Evidence Grade
Synthetic THC vs. Placebo	Pain response (≥30% improvement from baseline)	2 RCTs (N=84) ^{8,9}	Low	Direct	Very serious inconsistency	Imprecise	Unknown	Unable to assess, due to inconsistency from two trials (one trial of nabilone, 85% vs. 38%, RR 2.20 [1.06 to 4.55] and one trial of dronabinol, 43% vs. 57%, RR 0.76 [0.45 to 1.28])	Insufficient (previously low)
Synthetic THC vs. Placebo	Pain severity	7 RCTs (N=448) ⁸⁻¹⁴	Moderate	Direct	Consistent	Imprecise	Unknown	Small effect with synthetic THC 0 to 10 scale, MD -0.95 (-1.81 to -0.25; I ² =60%)	Low
Synthetic THC vs. Placebo	Function/disability	3 RCTs (N=unclear) ^{8,9,13} 1 RCT (N=13) not included in meta-analysis ¹⁴	Moderate	Direct	Consistent	Imprecise	Unknown	No effect (scale 0 to 10) MD: -0.18, -1.25 to 0.77, I ² =51%)	Low
Synthetic THC vs. Placebo	WAEs	5 RCTs (N=415) ⁹⁻¹³	Moderate	Direct	Consistent	Imprecise	Unknown	Potential moderate effect, not statistically significant 14% vs. 7%, RR 1.75 (0.95 to 4.11; I ² =0%)	Low
Synthetic THC vs. Placebo	SAEs	1 RCT (N=240) ¹¹	Low	Direct	Unknown	Imprecise	Unknown	Failed to demonstrate or exclude a detrimental effect 10% vs. 6%, RR 1.60 (0.65 to 3.93)	Insufficient

Comparison	Outcome	Number of Studies and Total Participants (N)	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	Strength of Evidence Grade
Synthetic THC vs. Placebo	Dizziness	3 RCTs (N=360) ⁹⁻¹¹	Low	Direct	Consistent	Imprecise	Unknown	Large effect with dronabinol 29% vs. 11%, RR 2.52 (1.20 to 4.82; I ² =41%)	Moderate
Synthetic THC vs. Placebo	Nausea	3 RCTs (N=302) ⁹⁻¹¹	Low	Direct	Consistent	Imprecise	Unknown	Potential large effect with dronabinol, not statistically significant 11% vs. 5%, RR 2.22 (0.90 to 5.05; I ² =0%)	Low
Synthetic THC vs. Placebo	Sedation	4 RCTs (N=386) ⁹⁻¹²	Moderate	Direct	Consistent	Imprecise	Unknown	Moderate effect with dronabinol 19% vs. 12%, RR 1.60 (1.01 to 2.95; I ² =7.7%)	Low

Abbreviations: CBD = cannabidiol; CI = confidence interval; KQ = Key Question; MD = mean difference; RCT = randomized controlled trial; RR = risk ratio; SAE = serious adverse event; SOE = strength of evidence; THC = tetrahydrocannabinol; WAE = withdrawal due to adverse event.

Table H-3. KQ1 and 2: Cannabinoids to treat chronic pain – high THC to CBD ratio, extracted from whole plant

Comparison	Outcome	Number of Studies and Total Participants (N)	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	Strength of Evidence Grade
Extracted High THC vs. Placebo	Pain severity	2 RCTs (N=297) ^{15,16}	Moderate	Direct	Inconsistent	Imprecise	Unknown	Failed to demonstrate or exclude a detrimental effect MD -1.97 (-5.91 to 1.21; I ² =85%)	Insufficient
	Function/disability	1 RCT (N=18) ¹⁶	High	Direct	Unknown	Imprecise	Unknown	Failed to demonstrate or exclude a detrimental effect MD 1.75 (-0.46 to 3.98)	Insufficient
	WAEs	1 RCT (N=277) ¹⁵	Moderate	Direct	Unknown	Imprecise	Unknown	Large increased risk 13.9% vs. 5.7%, RR 3.12 (1.54 to 6.33)	Low
	SAEs	1 RCT (N=277) ¹⁵	Moderate	Direct	Unknown	Imprecise	Unknown	Failed to demonstrate or exclude a detrimental effect 4.9% vs. 2.2%, RR 2.19 (0.58 to 8.28)	Insufficient
	Dizziness	1 RCT (N=277) ¹⁵	Moderate	Direct	Unknown	Imprecise	Unknown	Large effect 62.2% vs. 7.5%, RR 8.34 (4.53 to 15.34)	Low

Abbreviations: CBD = cannabidiol; CI = confidence interval; KQ = Key Question; MD = mean difference; RCT = randomized controlled trial; RR = risk ratio; SAE = serious adverse event; SOE = strength of evidence; THC = tetrahydrocannabinol; WAE = withdrawal due to adverse event.

Table H-4. KQ1 and 2: Cannabinoids to treat chronic pain – high THC to CBD ratio, combined synthetic and whole-plant extracted studies

Comparison	Outcome	Number of Studies and Total Participants (N)	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	Strength of Evidence Grade
Combined High THC Ratio Studies (Synthetic and Whole-plant extracted)	Pain severity	9 RCTs (N=742) ⁸⁻¹⁶	Moderate	Direct	Consistent	Precise	Unknown	Moderate effect MD -1.12 (-1.97 to -0.48; I ² =65%)	Moderate

Abbreviations: CBD = cannabidiol; CI = confidence interval; KQ = Key Question; MD = mean difference; RCT = randomized controlled trial; SOE = strength of evidence; THC = tetrahydrocannabinol.

Table H-5. KQ1 and 2: Cannabinoids to treat chronic pain – whole plant cannabis

Comparison	Outcome	Number of Studies and Total Participants (N)	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	Strength of Evidence Grade
Whole plant cannabis (standardized to 12% THC) vs. Usual Care	Pain Severity change	1 (N=431, 302 contribute to pain outcome) ¹⁷	High	Direct	Unknown	Imprecise	Unknown	Moderate effect 0 to 10 scale, Adjusted MD at 12 months: -1.10 (-1.56 to -0.72)	Insufficient
	WAE	1 (N=431) ¹⁷	High	Direct	Unknown	Imprecise	Unknown	Large effect with cannabis 4.7% vs. 0%, RR 21.10 (1.24 to 357.80)	Insufficient
	SAE	1 (N=431) ¹⁷	High	Direct	Unknown	Imprecise	Unknown	No effect 13% vs. 19%, OR 0.64 (0.38 to 1.04)	Insufficient
	Dizziness	1 (N=431) ¹⁷	High	Direct	Unknown	Imprecise	Unknown	Failed to demonstrate or exclude a detrimental effect 12.6% vs. 9.7%, RR 1.29 (0.75 to 2.21)	Insufficient
	Nausea	1 (N=431) ¹⁷	High	Direct	Unknown	Imprecise	Unknown	Moderate effect 16.7% vs. 9.7%, RR 1.72 (1.04 to 2.85)	Insufficient
	Sedation	1 (N=431) ¹⁷	High	Direct	Unknown	Imprecise	Unknown	Large effect 13.5% vs. 4.63%, RR 2.91 (1.46 to 5.83)	Insufficient
	Cognitive Disorder	1 (N=431) ¹⁷	High	Direct	Unknown	Imprecise	Unknown	Large effect 13.9% vs. 5.7%, RR 3.12 (1.54 to 6.33)	Insufficient

Abbreviations: CI = confidence interval; KQ = Key Question; MD = mean difference; OR = odds ratio; RCT = randomized controlled trial; RR = risk ratio; SAE = serious adverse event; SOE = strength of evidence; THC = tetrahydrocannabinol; WAE = withdrawal due to adverse event.

Table H-6. KQ1: Cannabinoids to treat chronic pain – low THC to CBD ratio

Comparison	Outcome	Number of Studies (N) and Total Participants	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	Strength of Evidence Grade
Topical, Plant-Extracted CBD vs. Placebo	Pain severity (change)	1 RCT (N=29) ¹⁸	High	Direct	Unknown	Imprecise	Unknown	Small effect with CBD cream MD -0.75, P=0.009 by ANCOVA (0 to 10 scale)	Insufficient
Oral Synthetic CBD vs. Placebo	Pain response (≥30% improvement)	1 RCT (N=136) ¹⁹	Moderate	Direct	Unknown	Imprecise	Unknown	No effect with oral synthetic CBD RR 1.01 (0.66 to 1.55)	Insufficient
Oral CBD or THC/CBD (Unknown If Synthetic or Plant-extracted vs. Placebo)^a	Pain severity (change)	1 RCT (N=87) ⁹	Low	Unclear	Unknown	Imprecise	Unknown	Potential increase in pain for CBD (MD 1.14 [0.11 to 2.19]) and no difference but imprecise for THC/CBD (MD -0.12 [-1.13 to 0.89])	Insufficient
	Pain response (≥30% improvement)	1 RCT (N=87) ⁹	Low	Unclear	Unknown	Imprecise	Unknown	Imprecise estimates for CBD (RR 0.59 [0.32 to 1.09]) and THC/CBD (RR 1.06 [0.69 to 1.62])	Insufficient
	Function/disability	1 RCT (N=87) ⁹	Low	Unclear	Unknown	Imprecise	Unknown	Imprecise estimates for CBD (MD 1.24 [-0.32 to 2.81]) and THC CBD (MD 0.89 [-0.64 to 2.42])	Insufficient

Abbreviations: ANCOVA = analysis of covariance; CBD = cannabidiol; CI = confidence interval; KQ = Key Question; MD = mean difference; RCT = randomized controlled trial; RR = relative risk; SOE = strength of evidence; THC = tetrahydrocannabinol.

^aStudy did not report whether CBD was synthetic or plant-extracted, and did not provide any details about the product composition.

Table H-7. KQ1 and 2: Cannabinoids to treat chronic pain – low THC to CBD ratio

Comparison	Outcome	Number of Studies (N) and Total Participants	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	Strength of Evidence Grade
CBDV vs. Placebo	Pain Response (≥30% improvement from baseline)	1 RCT (N=31) ²⁰	Moderate	Direct	Unknown	Imprecise	Unknown	Large effect, favors placebo 38% vs. 81%, RR 0.46 (95% CI 0.24 to 0.91)	Insufficient
CBDV vs. Placebo	Pain severity (change)	1 RCT (N=31) ²⁰	Moderate	Direct	Unknown	Imprecise	Unknown	Failed to demonstrate or exclude a detrimental effect MD 0.62 (-0.05 to 1.32)	Insufficient

Abbreviations: CBDV = cannabidivarin; CI = confidence interval; KQ = Key Question; MD = mean difference; RCT = randomized controlled trial; RR = risk ratio; SOE = strength of evidence.

Table H-8. KQ1 and 2: Observational studies of cannabinoids to treat chronic pain – unknown THC to CBD ratio (patient-choice)

Comparison	Outcome	Number of Studies (N) and Total Participants	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	SOE Grade
Unknown THC to CBD Ratio vs. Usual Care	Pain response ($\geq 30\%$ improvement from baseline)	No studies	NA	NA	NA	NA	NA	NA	No evidence
Unknown THC to CBD Ratio vs. Usual Care	Pain severity (change) Short-term (3 months)	2 cohort studies: short- to intermediate-term (N=202) ^{21,22}	High	Direct	Inconsistent	Imprecise	Unknown	VAS (0-100): 41.5 vs. 43.6 at 3 months ²¹ 34.1 vs. 48.8; mean difference -14.71 (95% CI, -32.71 to 3.29) ²²	Insufficient
Unknown THC to CBD Ratio vs. Usual Care	Long-term (12 months)	1 cohort (N=1,514) ²³	High	Direct	Unknown	Precise	Unknown	Adjusted mean; BPI, 0-10 scale) 5.2 vs. 4.9; Beta: 0.37 (95% CI -0.23 to 1.10), p=0.20 ²³	Insufficient
Unknown THC to CBD Ratio vs. Usual Care	Function or Disability (SF-36 Physical Function)	2 cohorts = short to medium-term (N=202) ^{21,22}	High	Direct	Consistent	Imprecise	Unknown	SF-36 Physical Functioning (mean, 0 to 100 scale) 46.5 vs. 43.7 at 6 months ²¹ 70.0 vs. 69.4; MD 0.56 (95% CI -17.2 to 18.3) at 3 months ²²	Insufficient
Unknown THC to CBD Ratio vs. Usual Care (Nabilone + Gabapentin vs. Gabapentin Alone)	WAEs	1 cohort study, short- and intermediate-term (N=156) ²¹	Moderate	Direct	Unknown	Imprecise	Unknown	6 months: 23% (12/52) vs. 9% (5/55), RR 2.54 (95% CI 0.95 to 6.71)	Insufficient

Comparison	Outcome	Number of Studies (N) and Total Participants	Study Limitations	Directness	Consistency	Precision	Publication Bias	Main Findings Effect Size (95% CI)	SOE Grade
Unknown THC to CBD Ratio vs. Usual Care (Nabilone + Gabapentin vs. Gabapentin Alone)	SAEs	1 cohort study, short- and intermediate-term (N=156) ²¹	Moderate	Direct	Unknown	Imprecise	Unknown	None in any group	Insufficient
Unknown THC to CBD Ratio vs. Usual Care (Nabilone + gabapentin vs. Gabapentin Alone)	Dizziness	1 cohort study, short- and intermediate-term (N=156) ²¹	Moderate	Direct	Unknown	Imprecise	Unknown	3 months: 33% (17/52) vs. 29% (16/55), RR 1.12 (95% CI 0.64 to 1.98) 6 months: 39% (20/52) vs. 33% (18/55), RR 1.17 (95% CI 0.70 to 0.91)	Insufficient
Unknown THC to CBD Ratio vs. Usual Care (Nabilone + gabapentin vs. Gabapentin Alone)	Nausea	No studies	NA	NA	NA	NA	NA	NA	No evidence
Unknown THC to CBD Ratio vs. Usual Care (Nabilone + Gabapentin vs. Gabapentin Alone)	Sedation	1 cohort study, short- and intermediate-term (N=156) ²¹	Moderate	Direct	Unknown	Imprecise	Unknown	3 months: 54% (28/52) vs. 33% (18/55) RR 1.65 (95% CI 1.04 to 2.59) 6 months: 60% (31/52) vs. 36% (20/55) RR 1.64 (95% CI 1.08 to 2.48)	Insufficient

Abbreviations: BPI-SF = Brief Pain Inventory (Short Form); CBD = cannabidiol; CI = confidence interval; KQ = Key Question; MD = mean difference; NA = not applicable; RCT = randomized controlled trial; RR = risk ratio; SAE = serious adverse event; SOE = strength of evidence; THC = tetrahydrocannabinol; WAE = withdrawal due to adverse event.

Appendix H References

1. Langford RM, Mares J, Novotna A, et al. A double-blind, randomized, placebo-controlled, parallel-group study of THC/CBD oromucosal spray in combination with the existing treatment regimen, in the relief of central neuropathic pain in patients with multiple sclerosis. *J Neurol*. 2013 Apr;260(4):984-97. doi: 10.1007/s00415-012-6739-4. PMID: 23180178.
2. Nurmikko TJ, Serpell MG, Hoggart B, et al. Sativex successfully treats neuropathic pain characterised by allodynia: a randomised, double-blind, placebo-controlled clinical trial. *Pain*. 2007 Dec 15;133(1-3):210-20. PMID: 17997224.
3. Selvarajah D, Gandhi R, Emery CJ, et al. Randomized placebo-controlled double-blind clinical trial of cannabis-based medicinal product (Sativex) in painful diabetic neuropathy: depression is a major confounding factor. *Diabetes Care*. 2010 Jan;33(1):128-30. doi: 10.2337/dc09-1029. PMID: 19808912.
4. Serpell M, Ratcliffe S, Hovorka J, et al. A double-blind, randomized, placebo-controlled, parallel group study of THC/CBD spray in peripheral neuropathic pain treatment. *Eur J Pain*. 2014 Aug;18(7):999-1012. doi: 10.1002/j.1532-2149.2013.00445.x. PMID: 24420962.
5. Blake DR, Robson P, Ho M, et al. Preliminary assessment of the efficacy, tolerability and safety of a cannabis-based medicine (Sativex) in the treatment of pain caused by rheumatoid arthritis. *Rheumatology (Oxford)*. 2006 Jan;45(1):50-2. PMID: 16282192.
6. Lynch ME, Cesar-Rittenberg P, Hohmann AG. A double-blind, placebo-controlled, crossover pilot trial with extension using an oral mucosal cannabinoid extract for treatment of chemotherapy-induced neuropathic pain. *J Pain Symptom Manage*. 2014 Jan;47(1):166-73. doi: 10.1016/j.jpainsymman.2013.02.018. PMID: 23742737.
7. Rog DJ, Nurmikko TJ, Friede T, et al. Randomized, controlled trial of cannabis-based medicine in central pain in multiple sclerosis. *Neurology*. 2005 Sep 27;65(6):812-9. PMID: 16186518.
8. Toth C, Mawani S, Brady S, et al. An enriched-enrolment, randomized withdrawal, flexible-dose, double-blind, placebo-controlled, parallel assignment efficacy study of nabilone as adjuvant in the treatment of diabetic peripheral neuropathic pain. *Pain*. 2012 Oct;153(10):2073-82. doi: 10.1016/j.pain.2012.06.024. PMID: 22921260.
9. Zubcevic K, Petersen M, Bach FW, et al. Oral capsules of tetra-hydro-cannabinol (THC), cannabidiol (CBD) and their combination in peripheral neuropathic pain treatment. *Eur J Pain*. 2022doi: 10.1002/ejp.2072. PMID: 36571471.
10. de Vries M, van Rijckevorsel DCM, Vissers KCP, et al. Tetrahydrocannabinol Does Not Reduce Pain in Patients With Chronic Abdominal Pain in a Phase 2 Placebo-controlled Study. *Clin Gastroenterol Hepatol*. 2017 Jul;15(7):1079-86.e4. doi: 10.1016/j.cgh.2016.09.147. PMID: 27720917.
11. Schimrigk S, Marziniak M, Neubauer C, et al. Dronabinol Is a Safe Long-Term Treatment Option for Neuropathic Pain Patients. *Eur Neurol*. 2017;78(5-6):320-9. doi: 10.1159/000481089. PMID: 29073592.
12. Skrabek RQ, Galimova L, Ethans K, et al. Nabilone for the treatment of pain in fibromyalgia. *J Pain*. 2008 Feb;9(2):164-73. PMID: 17974490.
13. Turcotte D, Doupe M, Torabi M, et al. Nabilone as an adjunctive to gabapentin for multiple sclerosis-induced neuropathic pain: a randomized controlled trial. *Pain Med*. 2015 Jan;16(1):149-59. doi: 10.1111/pme.12569. PMID: 25288189.

14. Wissel J, Haydn T, Muller J, et al. Low dose treatment with the synthetic cannabinoid Nabilone significantly reduces spasticity-related pain : a double-blind placebo-controlled cross-over trial. *J Neurol*. 2006 Oct;253(10):1337-41. PMID: 16988792.
15. Zajicek JP, Hobart JC, Slade A, et al. Multiple sclerosis and extract of cannabis: results of the MUSEC trial. *J Neurol Neurosurg Psychiatry*. 2012 Nov;83(11):1125-32. doi: 10.1136/jnnp-2012-302468. PMID: 22791906.
16. Chaves C, Bittencourt PCT, Pelegrini A. Ingestion of a THC-Rich Cannabis Oil in People with Fibromyalgia: A Randomized, Double-Blind, Placebo-Controlled Clinical Trial. *Pain Med*. 2020;21(10):2212-8. doi: 10.1093/pm/pnaa303. PMID: 33118602.
17. Ware MA, Wang T, Shapiro S, et al. Cannabis for the Management of Pain: Assessment of Safety Study (COMPASS). *J Pain*. 2015 Dec;16(12):1233-42. doi: 10.1016/j.jpain.2015.07.014. PMID: 26385201.
18. Xu DH, Cullen BD, Tang M, et al. The Effectiveness of Topical Cannabidiol Oil in Symptomatic Relief of Peripheral Neuropathy of the Lower Extremities. *Curr Pharm Biotechnol*. 2020;21(5):390-402. doi: 10.2174/1389201020666191202111534. PMID: 31793418.
19. Vela J, Dreyer L, Petersen KK, et al. Cannabidiol treatment in hand osteoarthritis and psoriatic arthritis: a randomized, double-blind placebo-controlled trial. *Pain*. 2021 Jun 1;163(6):1206-14. PMID: 34510141.
20. Eibach L, Scheffel S, Cardebring M, et al. Cannabidivarin for HIV-Associated Neuropathic Pain: A Randomized, Blinded, Controlled Clinical Trial. *Clin Pharmacol Ther*. 2020 Aug 08;109(4):1055-62. doi: 10.1002/cpt.2016. PMID: 32770831.
21. Bestard JA, Toth CC. An open-label comparison of nabilone and gabapentin as adjuvant therapy or monotherapy in the management of neuropathic pain in patients with peripheral neuropathy. *Pain Pract*. 2011 Jul-Aug;11(4):353-68. doi: 10.1111/j.1533-2500.2010.00427.x. PMID: 21087411.
22. Gruber SA, Smith RT, Dahlgren MK, et al. No pain, all gain? Interim analyses from a longitudinal, observational study examining the impact of medical cannabis treatment on chronic pain and related symptoms. *Exp Clin Psychopharmacol*. 2021;29(2):147-56. doi: 10.1037/pha0000435. PMID: 33764103.
23. Campbell G, Hall WD, Peacock A, et al. Effect of cannabis use in people with chronic non-cancer pain prescribed opioids: findings from a 4-year prospective cohort study. *Lancet Public Health*. 2018 Jul;3(7):e341-e50. doi: 10.1016/s2468-2667(18)30110-5. PMID: 29976328.

Appendix I. Excluded Studies List

1. Vaporized Cannabis for chronic pain associated with Sickle Cell Disease. Cannabinoid-based therapy and approaches to quantify pain in Sickle Cell disease. 2013. **Exclusion reason:** Ineligible publication type
2. Cannabis-opioid interaction in the treatment of Fibromyalgia pain “an open label proof of concept study with randomization between treatment groups: Cannabis, Oxycodone or Cannabis/Oxycodone combination. 2019. **Exclusion reason:** Ineligible study design
3. Proof of concept trial of Cannabis derivatives in neuropathic pain. Proof of Concept Trial of Cannabis Derivatives in Neuropathic Pain. 2022. **Exclusion reason:** Ineligible publication type
4. Topical CBD for musculoskeletal pain. Immediate effect of topical CBD for Musculoskeletal pain. 2022. **Exclusion reason:** Ineligible population
5. Cannabinoids for the Reduction of Inflammation and Sickle Cell Related Pain. Dronabinol for the Reduction of Chronic Pain and Inflammation in People With Sickle Cell Disease. 2022. **Exclusion reason:** Ineligible publication type
6. A Phase III study to investigate the effect of EMD-RX5 on symptoms of psychological distress in adults with chronic pain. A multi-site, parallel-arm, randomised, double blind, placebo-controlled study to investigate the effect of EMD-RX5 on symptoms of psychological distress in adults with chronic pain. 2022. **Exclusion reason:** Ineligible publication type
7. Comparison of VER-01 to Opioids in Patients With Chronic Non-specific Low Back Pain. Multicentre, Randomized, Open-label Study to Prove an Additional Benefit of the Full-spectrum Cannabis Extract VER-01 Over Opioids in the Treatment of Patients With Chronic Non-specific Low Back Pain. 2022. **Exclusion reason:** Ineligible publication type
8. Effects of Cannabidiol (CBD) on Resting-state Electroencephalography (EEG) and Neuropathic Pain Severity in People With Spinal Cord Injury (SCI). 2022. **Exclusion reason:** Ineligible publication type
9. Diabetic Neuropathic Pain Relief, 6 Weeks Dosage Sublingual Water-soluble CBD/PEA. A Randomized, Double-Blind, Placebo-Controlled Trial Using Cannabidiol and Palmitoylethanolamide for the Treatment of Painful Diabetic Peripheral Neuropathy of the Feet. 2023. **Exclusion reason:** Ineligible publication type
10. A Double-Blind, Randomised Placebo-Controlled Feasibility Trial Assessing Oral Cannabis for The Relief of Fibromyalgia Symptoms. 2023. **Exclusion reason:** Ineligible publication type
11. A Randomised, Double-Blind, Placebo-Controlled, Parallel-Arm Study to Evaluate the Efficacy, Safety and Tolerability of 150 mg Sublingual Cannabidiol in Adults with Mild-to-Moderate Pain. 2023. **Exclusion reason:** Ineligible publication type
12. Abelev S, Warne LN, Benson M, et al. Medicinal Cannabis for the treatment of chronic refractory pain: an investigation of the adverse event profile and health-related quality of life impact of an oral formulation. *Med Cannabis Cannabinoids*. 2022;5(1):20-31. doi: 10.1159/000521492. PMID: 35950052. **Exclusion reason:** Ineligible comparator
13. Abo Ziad R, Grynbaum MB, Peleg R, et al. The Attitudes and Beliefs of Family Physicians Regarding the Use of Medical Cannabis, Knowledge of Side Effects, and Barriers to Use: A Comparison Between Residents and Specialists. *Am J Ther*. 2020; Publish Ahead of Print doi: 10.1097/MJT.0000000000001236. PMID: 33416237. **Exclusion reason:** Ineligible study design

14. Aboud T, Schuster NM. Pain management in Multiple Sclerosis: a review of available treatment options. *Curr Treat Options Neurol*. 2019 Nov 27;21(12):62. doi: 10.1007/s11940-019-0601-2. PMID: 31773455. **Exclusion reason:** Systematic review used as source document
15. Abrams DI, Couey P, Dixit N, et al. Effect of inhaled Cannabis for pain in adults with Sickle Cell Disease: a randomized clinical trial. *JAMA Netw Open*. 2020 Jul 01;3(7):e2010874. doi: 10.1001/jamanetworkopen.2020.10874. PMID: 32678452. **Exclusion reason:** Inadequate duration
16. Abrams DI, Jay CA, Shade SB, et al. Cannabis in painful HIV-associated sensory neuropathy: a randomized placebo-controlled trial. *Neurology*. 2007 Feb 13;68(7):515-21. doi: 10.1212/01.wnl.0000253187.66183.9c. PMID: 17296917. **Exclusion reason:** Inadequate duration
17. Abuhasira R, Ron A, Sikorin I, et al. Medical Cannabis for older patients—treatment protocol and initial results. *J Clin Med*. 2019 Nov 01;8(11):1819. doi: 10.3390/jcm8111819. PMID: 31683817. **Exclusion reason:** Ineligible population
18. Abuhasira R, Ron A, Sikorin I, et al. Medical Cannabis for older patients—treatment protocol and initial results. *J Clin Med*. 2019;8(11)doi: 10.3390/jcm8111819. PMID: 31683817. **Exclusion reason:** Ineligible population
19. Aebischer JH, Dieckmann NF, Jones KD, et al. Chronic pain clinical and prescriptive practices in the Cannabis era. *Pain Manag Nurs*. 2021 Dec 29;29:29. doi: 10.1016/j.pmn.2021.11.009. PMID: 34973920. **Exclusion reason:** Systematic review used as source document
20. Akgün K, Essner U, Seydel C, et al. Daily practice managing resistant Multiple Sclerosis spasticity with Delta-9-Tetrahydrocannabinol: Cannabidiol oromucosal spray: a systematic review of observational studies. *J Cent Nerv Syst Dis*. 2019;11doi: 10.1177/1179573519831997. PMID: 30886530. **Exclusion reason:** Systematic review used as source document
21. Allan GM, Finley CR, Ton J, et al. Systematic review of systematic reviews for Medical Cannabinoids: pain, nausea and vomiting, spasticity, and harms. *Can Fam Physician*. 2018 Feb;64(2):e78-e94. PMID: 29449262. **Exclusion reason:** Ineligible publication type
22. Almog S, Aharon-Peretz J, Vulfsons S, et al. The pharmacokinetics, efficacy, and safety of a novel selective-dose Cannabis inhaler in patients with chronic pain: a randomized, double-blinded, placebo-controlled trial. *Eur J Pain*. 2020 Sep;24(8):1505-16. doi: 10.1002/ejp.1605. PMID: 32445190. **Exclusion reason:** Inadequate duration
23. Aly E, Masocha W. Targeting the Endocannabinoid system for management of HIV-associated neuropathic pain: a systematic review. *IBRO Neurosci Rep*. 2021 Jun;10:109-18. doi: 10.1016/j.ibneur.2021.01.004. PMID: 34179865. **Exclusion reason:** Systematic review used as source document
24. Amato L, Minozzi S, Mitrova Z, et al. Systematic review of safeness and therapeutic efficacy of Cannabis in patients with Multiple Sclerosis, neuropathic pain, and in oncological patients treated with chemotherapy. *Epidemiol Prev*. 2017;41(5-6)doi: 10.19191/EP17.5-6.AD01.069. PMID: 29119763. **Exclusion reason:** Ineligible publication type
25. AminiLari M, Wang L, Neumark S, et al. Medical Cannabis and Cannabinoids for impaired sleep: a systematic review and meta-analysis of randomized clinical trials. *Sleep*. 2021doi: 10.1093/sleep/zsab234. PMID: 34546363. **Exclusion reason:** Systematic review used as source document
26. Anaya HJM, Ortiz MPT, Valencia DHF, et al. Efficacy of Cannabinoids in Fibromyalgia: a literature review. *Colomb J Anesthesiol*. 2021;49(4)doi: 10.5554/22562087.e980. **Exclusion reason:** Inadequate duration
27. Andreae MH, Carter GM, Shaparin N, et al. Inhaled Cannabis for chronic neuropathic pain: a meta-analysis of individual patient data. *J Pain*. 2015 Dec;16(12):1221-32. doi: 10.1016/j.jpain.2015.07.009. PMID: 26362106. **Exclusion reason:** Inadequate duration

28. Anonymous. National Institute for Health and Care Excellence (UK). 2019 11;11:11. PMID: 35107907. **Exclusion reason:** Systematic review used as source document
29. Arkell TR, Downey LA, Hayley AC, et al. Assessment of Medical Cannabis and Health-Related Quality of Life. *JAMA Netw Open*. 2023;6(5):e2312522. doi: <https://dx.doi.org/10.1001/jamanetworkopen.2023.12522>. **Exclusion reason:** Ineligible study design
30. Arnold JC, McCartney D, Suraev A, et al. The safety and efficacy of low oral doses of cannabidiol: An evaluation of the evidence. *Clinical and translational science*. 2022doi: <https://dx.doi.org/10.1111/cts.13425>. **Exclusion reason:** Systematic review used as source document
31. Aviram J, Atzmony D, Eisenberg E. Long-term effectiveness and safety of Medical Cannabis administered through the metered-dose Syqe Inhaler. *Pain reports*. 2022;7(3):e1011. doi: 10.1097/PR9.0000000000001011. PMID: 35620248. **Exclusion reason:** Ineligible comparator
32. Aviram J, Lewitus GM, Pud D, et al. Specific Phytocannabinoid compositions are associated with analgesic response and adverse effects in chronic pain patients treated with Medical Cannabis. *Pharmacol Res*. 2021 Jul;169:105651. doi: 10.1016/j.phrs.2021.105651. PMID: 34000362. **Exclusion reason:** Ineligible comparator
33. Aviram J, Lewitus GM, Vysotski Y, et al. Sex differences in Medical Cannabis-related adverse effects. *Pain*. 2021doi: 10.1097/j.pain.0000000000002463. PMID: 34538843. **Exclusion reason:** Ineligible comparator
34. Aviram J, Pud D, Gershoni T, et al. Medical Cannabis treatment for chronic pain: outcomes and prediction of response. *Eur J Pain*. 2020 Oct 16;16:16. doi: 10.1002/ejp.1675. PMID: 33065768. **Exclusion reason:** Ineligible comparator
35. Aviram J, Samuely-Leichtag G. Efficacy of Cannabis-based medicines for pain management: a systematic review and meta-analysis of randomized controlled trials. *Pain Physician*. 2017 Sep;20(6):E755-E96. PMID: 28934780. **Exclusion reason:** Systematic review used as source document
36. Bajtel A, Kiss T, Toth B, et al. The safety of Dronabinol and Nabilone: a systematic review and meta-analysis of clinical trials. *Pharmaceuticals (Basel)*. 2022 Jan 14;15(1):14. doi: 10.3390/ph15010100. PMID: 35056154. **Exclusion reason:** Systematic review used as source document
37. Bakewell BK, Sherman M, Binsfeld K, et al. The Use of Cannabidiol in Patients With Low Back Pain Caused by Lumbar Spinal Stenosis: An Observational Study. *Cureus*. 2022;14(9):e29196. doi: <https://dx.doi.org/10.7759/cureus.29196>. **Exclusion reason:** Ineligible study design
38. Balestra AM, Chalk K, Denke C, et al. Influence of Cannabinoid Treatment on Trajectories of Patient-Related Outcomes in Chronic Pain: Pain Intensity, Emotional Distress, Tolerability and Physical Disability. *Brain sci*. 2023;13(4)doi: 10.3390/brainsci13040680. **Exclusion reason:** Ineligible comparator
39. Ball S, Vickery J, Hobart J, et al. The Cannabinoid Use in Progressive Inflammatory brain Disease (CUPID) trial: a randomised double-blind placebo-controlled parallel-group multicentre trial and economic evaluation of cannabinoids to slow progression in multiple sclerosis. *Health Technol Assess*. 2015;19(12):1-187. doi: 10.3310/hta19120. PMID: 25676540. **Exclusion reason:** Ineligible outcome
40. Balu A, Mishra D, Marcu J, et al. Medical Cannabis certification is associated with decreased opiate use in patients with chronic pain: a retrospective cohort study in Delaware. *Cureus*. 2021 Dec;13(12):e20240. doi: 10.7759/cureus.20240. PMID: 35004055. **Exclusion reason:** Ineligible comparator

41. Barakji J, Korang SK, Feinberg J, et al. Cannabinoids versus placebo for pain: A systematic review with meta-analysis and Trial Sequential Analysis. *PLoS ONE*. 2023;18(1):e0267420. doi: <https://dx.doi.org/10.1371/journal.pone.0267420>. **Exclusion reason:** Systematic review used as source document
42. Bargnes V, Hart P, Gupta S, et al. Safety and efficacy of medical cannabis in elderly patients: a retrospective review in a neurological outpatient setting. *Neurology*. 2019;92(15). **Exclusion reason:** Ineligible publication type
43. Barnes MP. Sativex: clinical efficacy and tolerability in the treatment of symptoms of Multiple Sclerosis and neuropathic pain. *Expert Opin Pharmacother*. 2006 Apr;7(5):607-15. doi: [10.1517/14656566.7.5.607](https://doi.org/10.1517/14656566.7.5.607). PMID: 16553576. **Exclusion reason:** Ineligible publication type
44. Becker WC, Li Y, Caniglia EC, et al. Cannabis use, pain interference, and prescription opioid receipt among persons with HIV: a target trial emulation study. *AIDS Care*. 2021 Jun 28:1-9. doi: [10.1080/09540121.2021.1944597](https://doi.org/10.1080/09540121.2021.1944597). PMID: 34180721. **Exclusion reason:** Ineligible population
45. Bellnier T, Brown GW, Ortega TR. Preliminary evaluation of the efficacy, safety, and costs associated with the treatment of chronic pain with Medical Cannabis. *Ment Health Clin*. 2018 Apr 26;8(3):110-5. doi: [10.9740/mhc.2018.05.110](https://doi.org/10.9740/mhc.2018.05.110). PMID: 29955555. **Exclusion reason:** Ineligible comparator
46. Benedict G, Sabbagh A, Conermann T. Medical Cannabis used as an alternative treatment for chronic pain demonstrates reduction in chronic opioid use - a prospective study. *Pain Physician*. 2022 Jan;25(1):E113-E9. PMID: 35051158. **Exclusion reason:** Ineligible comparator
47. Bennici A, Mannucci C, Calapai F, et al. Safety of Medical Cannabis in neuropathic chronic pain management. *Molecules (Basel)*. 2021;26(20):16. doi: [10.3390/molecules26206257](https://doi.org/10.3390/molecules26206257). PMID: 34684842. **Exclusion reason:** Systematic review used as source document
48. Berger AA, Keefe J, Winnick A, et al. Cannabis and Cannabidiol (CBD) for the treatment of Fibromyalgia. *Best Pract Res Clin Anaesthesiol*. 2020doi: [10.1016/j.bpa.2020.08.010](https://doi.org/10.1016/j.bpa.2020.08.010). PMID: 33004171. **Exclusion reason:** Ineligible publication type
49. Berman JS, Symonds C, Birch R. Efficacy of two Cannabis based medicinal extracts for relief of central neuropathic pain from brachial plexus avulsion: results of a randomised controlled trial. *Pain*. 2004 Dec;112(3):299-306. doi: [10.1016/j.pain.2004.09.013](https://doi.org/10.1016/j.pain.2004.09.013). PMID: 15561385. **Exclusion reason:** Inadequate duration
50. Bialas P, Fitzcharles M-A, Klose P, et al. Long-term observational studies with Cannabis-based medicines for chronic non-cancer pain: a systematic review and meta-analysis of effectiveness and safety. *Eur J Pain*. 2022doi: [10.1002/ejp.1957](https://doi.org/10.1002/ejp.1957). PMID: 35467781. **Exclusion reason:** Systematic review used as source document
51. Bicket MC, Stone EM, McGinty EE. Use of Cannabis and Other Pain Treatments Among Adults With Chronic Pain in US States With Medical Cannabis Programs. *JAMA Netw Open*. 2023;6(1):e2249797. doi: <https://dx.doi.org/10.1001/jamanetworkopen.2022.49797>. **Exclusion reason:** Ineligible study design
52. Bilbao A, Spanagel R. Medical Cannabinoids: a pharmacology-based systematic review and meta-analysis for all relevant medical indications. *BMC medicine*. 2022;20(1):259. doi: [10.1186/s12916-022-02459-1](https://doi.org/10.1186/s12916-022-02459-1). PMID: 35982439. **Exclusion reason:** Systematic review used as source document
53. Blake A, Wan BA, Malek L, et al. A selective review of Medical Cannabis in cancer pain management. *Ann Palliat Med*. 2017 Dec;6(Suppl 2):S215-S22. doi: [10.21037/apm.2017.08.05](https://doi.org/10.21037/apm.2017.08.05). PMID: 28866904. **Exclusion reason:** Ineligible population
54. Boehnke KF, Clauw DJ. Cannabinoids for Chronic Pain: Translating Systematic Review Findings Into Clinical Action. *Ann Intern Med*. 2022doi: [10.7326/M22-1512](https://doi.org/10.7326/M22-1512). PMID: 35667063. **Exclusion reason:** Background

55. Boehnke KF, Gagnier JJ, Matallana L, et al. Cannabidiol Use for Fibromyalgia: Prevalence of Use and Perceptions of Effectiveness in a Large Online Survey. *J Pain*. 2021doi: 10.1016/j.jpain.2020.12.001. PMID: 33400996. **Exclusion reason:** Ineligible study design
56. Boehnke KF, Gagnier JJ, Matallana L, et al. Substituting Cannabidiol for opioids and pain medications among individuals with Fibromyalgia: a large online survey. *J Pain*. 2021doi: 10.1016/j.jpain.2021.04.011. PMID: 33992787. **Exclusion reason:** Background
57. Boehnke KF, Hauser W, Fitzcharles M-A. Cannabidiol (CBD) in Rheumatic Diseases (Musculoskeletal Pain). *Curr Rheumatol Rep*. 2022doi: 10.1007/s11926-022-01077-3. PMID: 35503198. **Exclusion reason:** Systematic review used as source document
58. Boehnke KF, Scott JR, Litinas E, et al. High-frequency Medical Cannabis use is associated with worse pain among individuals with chronic pain. *J Pain*. 2020 May - Jun;21(5-6):570-81. doi: 10.1016/j.jpain.2019.09.006. PMID: 31560957. **Exclusion reason:** Ineligible comparator
59. Bonomo Y, Norman A, Collins L, et al. Pharmacokinetics, safety, and tolerability of a Medicinal Cannabis formulation in patients with chronic non-cancer pain on long-term high dose opioid analgesia: a pilot study. *Pain Ther*. 2021;18:18. doi: 10.1007/s40122-021-00344-y. PMID: 34921662. **Exclusion reason:** Ineligible comparator
60. Boychuk DG, Goddard G, Mauro G, et al. The effectiveness of Cannabinoids in the management of chronic nonmalignant neuropathic pain: a systematic review. *J Oral Facial Pain Headache*. 2015;29(1):7-14. doi: 10.11607/ofph.1274. PMID: 25635955. **Exclusion reason:** Ineligible publication type
61. Busse JW, MacKillop J. Medical Cannabis and Cannabinoids for chronic pain: summary of a rapid recommendation. *J Mil Veteran Fam Health*. 2021;7:118-22. doi: 10.3138/jmvfh-2021-0056. **Exclusion reason:** Ineligible publication type
62. Busse JW, Wang L, Kamaleldin M, et al. Opioids for chronic noncancer pain: a systematic review and meta-analysis. *Jama*. 2018 Dec 18;320(23):2448-60. doi: 10.1001/jama.2018.18472. PMID: 30561481. **Exclusion reason:** Systematic review used as source document
63. Campbell CM, Mun CJ, Hamilton KR, et al. Within-subject, double-blind, randomized, placebo-controlled evaluation of combining the cannabinoid dronabinol and the opioid hydromorphone in adults with chronic pain. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2023doi: https://dx.doi.org/10.1038/s41386-023-01597-1. **Exclusion reason:** Ineligible outcome
64. Canavan C, Inoue T, McMahon S, et al. The efficacy, adverse events, and withdrawal rates of the pharmacological management of chronic spinal cord injury pain: a systematic review and meta-analysis. *Pain Med*. 2022 Feb 01;23(2):375-95. doi: 10.1093/pm/pnab140. PMID: 33844010. **Exclusion reason:** Systematic review used as source document
65. Carreira DS, Garden S, Huffman A, et al. Cannabinoids in the Orthopedic Setting: A Literature Review. *Orthopedics*. 2022:1-7. doi: 10.3928/01477447-20220225-11. PMID: 35245146. **Exclusion reason:** Systematic review used as source document
66. Chan CJ. Efficacy of plant based Cannabis in reducing pain in patients with chronic pain: a meta analysis. *Diss Abstr Int*. 2020;81(10-B):No Pagination Specified. **Exclusion reason:** Ineligible publication type
67. Christ MM. Pain medicine: Cannabis is effective in neuropathic pain. *Arzneimitteltherapie*. 2019;37(6):242-3. **Exclusion reason:** Not in English
68. Clermont-Gnamien S, Atlani S, Attal N, et al. The therapeutic use of Δ^9 -tetrahydrocannabinol (dronabinol) in refractory neuropathic pain. *Presse Med*. 2002;31(39 I):1840-5. PMID: 12496714. **Exclusion reason:** Not in English

69. Coates MD, Dalessio S, Walter V, et al. Symptoms and extraintestinal manifestations in active Cannabis users with Inflammatory Bowel Disease. *Cannabis Cannabinoid Res.* 2022;7(4):445-50. doi: 10.1089/can.2020.0155. PMID: 33998892. **Exclusion reason:** Ineligible population
70. Cooper ZD, Abrams DI. Considering abuse liability and neurocognitive effects of Cannabis and Cannabis-derived products when assessing Analgesic efficacy: a comprehensive review of randomized-controlled studies. *Am J Drug Alcohol Abuse.* 2019;45(6):580-95. doi: 10.1080/00952990.2019.1669628. PMID: 31687845. **Exclusion reason:** Systematic review used as source document
71. Corey-Bloom J, Wolfson T, Gamst A, et al. Smoked Cannabis for spasticity in Multiple Sclerosis: a randomized, placebo-controlled trial. *Cmaj.* 2012 Jul 10;184(10):1143-50. doi: 10.1503/cmaj.110837. PMID: 22586334. **Exclusion reason:** Inadequate duration
72. Costales B, van Boemmel-Wegmann S, Winterstein A, et al. Clinical conditions and prescription drug utilization among early medical marijuana registrants in Florida. *J Psychoactive Drugs.* 2021:1-10. doi: 10.1080/02791072.2020.1864069. PMID: 33393877. **Exclusion reason:** Ineligible study design
73. Coughlin LN, Ilgen MA, Jannausch M, et al. Progression of Cannabis withdrawal symptoms in people using Medical Cannabis for chronic pain. *Addiction.* 2021doi: 10.1111/add.15370. PMID: 33400332. **Exclusion reason:** Ineligible study design
74. Crestani F. Medical Cannabis for the treatment of Fibromyalgia. *J Clin Rheumatol.* 2018 Aug;24(5):281. doi: 10.1097/RHU.0000000000000823. PMID: 29757806. **Exclusion reason:** Ineligible study design
75. Cumenal M, Selvy M, Kerckhove N, et al. The safety of medications used to treat peripheral neuropathic pain, part 2 (opioids, Cannabinoids and other drugs): review of double-blind, placebo-controlled, randomized clinical trials. *Expert Opin Drug Saf.* 2020doi: 10.1080/14740338.2021.1842871. PMID: 33103931. **Exclusion reason:** Systematic review used as source document
76. Cunetti L, Manzo L, Peyraube R, et al. Chronic pain treatment with Cannabidiol in kidney transplant patients in Uruguay. *Transplant Proc.* 2018 Mar;50(2):461-4. doi: 10.1016/j.transproceed.2017.12.042. PMID: 29579828. **Exclusion reason:** Ineligible comparator
77. Cunningham CO, Starrels JL, Zhang C, et al. Medical marijuana and opioids (MEMO) study: protocol of a longitudinal cohort study to examine if medical cannabis reduces opioid use among adults with chronic pain. *BMJ Open.* 2020;10(12):e043400. doi: 10.1136/bmjopen-2020-043400. PMID: 33376181. **Exclusion reason:** Ineligible study design
78. Curtis SA, Brandow AM, Deveaux M, et al. Daily Cannabis users with Sickle Cell Disease show fewer admissions than others with similar pain complaints. *Cannabis Cannabinoid Res.* 2020;5(3):255-62. doi: 10.1089/can.2019.0036. PMID: 32923662. **Exclusion reason:** Ineligible study design
79. Darnall BD, Humphreys KN. An experimental method for assessing whether marijuana use reduces opioid use in patients with chronic pain. *Addiction.* 2018 Aug;113(8):1552-3. doi: 10.1111/add.14239. PMID: 29882256. **Exclusion reason:** Ineligible study design
80. Datta S, Ramamurthy PC, Anand U, et al. Wonder or evil?: multifaceted health hazards and health benefits of Cannabis Sativa and its Phytochemicals. *Saudi J Biol Sci.* 28(12):7290-313. doi: 10.1016/j.sjbs.2021.08.036. PMID: 34867033. **Exclusion reason:** Ineligible publication type

81. Degenhardt L, Lintzeris N, Campbell G, et al. Experience of adjunctive Cannabis use for chronic non-cancer pain: findings from the pain and opioids in treatment (POINT) study. *Drug Alcohol Depend.* 2015 Feb 01;147:144-50. doi: 10.1016/j.drugalcdep.2014.11.031. PMID: 25533893. **Exclusion reason:** Ineligible study design
82. Denduluri SK, Woolson ST, Indelli PF, et al. Cannabinoid and Opioid Use Among Total Joint Arthroplasty Patients: A 6-Year, Single-Institution Study. *Orthopedics.* 2020 Oct 01;1-6. doi: 10.3928/01477447-20200928-02. PMID: 33002174. **Exclusion reason:** Ineligible outcome
83. Deshpande A, Mailis-Gagnon A, Zoheiry N, et al. Efficacy and adverse effects of medical marijuana for chronic noncancer pain: systematic review of randomized controlled trials. *Can Fam Physician.* 2015 Aug;61(8):e372-81. PMID: 26505059. **Exclusion reason:** Ineligible publication type
84. Dimitrios L, Aris F. Efficacy, tolerability and safety of Cannabinoids for management of pain in adult patients with Multiple Sclerosis: a systematic review and meta-analysis. *Signa Vitae.* 2021;17:S10. doi: 10.22514/sv.2021.157. **Exclusion reason:** Ineligible publication type
85. Domnic G, Narayanan S, Mohana-Kumaran N, et al. Kratom (*Mitragyna speciosa* Korth.) an overlooked medicinal plant in Malaysia. *J Subst Use.* 2022;27(1):1-6. doi: 10.1080/14659891.2021.1885515. **Exclusion reason:** Systematic review used as source document
86. Durán M, Capellà D. Cannabis and Cannabinoids in the treatment of neuropathic pain. *DOLOR.* 2005;20(4):213-6. **Exclusion reason:** Not in English
87. Dykukha I, Malessa R, Essner U, et al. Nabiximols in chronic neuropathic pain: a meta-analysis of randomized placebo-controlled trials. *Pain Med.* 2021 04 20;22(4):861-74. doi: 10.1093/pm/pnab050. PMID: 33561282. **Exclusion reason:** Systematic review used as source document
88. Eadie L, Lo LA, Christiansen A, et al. Duration of neurocognitive impairment with Medical Cannabis use: a scoping review. *Front Psychiatry.* 2021;12doi: 10.3389/fpsy.2021.638962. PMID: 33790818. **Exclusion reason:** Systematic review used as source document
89. Edinoff AN, Fort JM, Singh C, et al. Alternative Options for Complex, Recurrent Pain States Using Cannabinoids, Psilocybin, and Ketamine: A Narrative Review of Clinical Evidence. *Neurology int.* 2022;14(2):423-36. doi: 10.3390/neurolint14020035. PMID: 35645354. **Exclusion reason:** Ineligible publication type
90. Eichorn NL, Shult HT, Kracht KD, et al. Making a joint decision: Cannabis as a potential substitute for opioids in Obstetrics and Gynecology. *Best practice & research. Clinical obstetrics & gynaecology.* 2022;S1521-6934(22):00100-6. doi: 10.1016/j.bpobgyn.2022.07.002. PMID: 35970747. **Exclusion reason:** Ineligible publication type
91. Ellis RJ, Toperoff W, Vaida F, et al. Smoked Medicinal Cannabis for neuropathic pain in HIV: a randomized, crossover clinical trial. *Neuropsychopharmacology.* 2009 Feb;34(3):672-80. doi: 10.1038/npp.2008.120. PMID: 18688212. **Exclusion reason:** Inadequate duration
92. Ergisi M, Erridge S, Harris M, et al. An updated analysis of clinical outcome measures across patients from the UK Medical Cannabis registry. *Cannabis Cannabinoid Res.* 2022 Jan 24;24:24. doi: 10.1089/can.2021.0145. PMID: 35073160. **Exclusion reason:** Ineligible population
93. Fallon MT, Albert Lux E, McQuade R, et al. Sativex oromucosal spray as adjunctive therapy in advanced cancer patients with chronic pain unalleviated by optimized opioid therapy: two double-blind, randomized, placebo-controlled phase 3 studies. *Br J Pain.* 2017 Aug;11(3):119-33. doi: 10.1177/2049463717710042. PMID: 28785408. **Exclusion reason:** Ineligible population

94. Feingold D, Brill S, Goor-Aryeh I, et al. Depression and Anxiety among chronic pain patients receiving prescription opioids and medical marijuana. *J Affect Disord.* 2017 Aug 15;218:1-7. doi: 10.1016/j.jad.2017.04.026. PMID: 28453948. **Exclusion reason:** Ineligible study design
95. Ferrie ML, Rogers AH, Zvolensky MJ, et al. Alcohol and marijuana co-use among adults with chronic low back pain: Associations with substance misuse, mental health, and pain experience. *The American journal on addictions.* 2022;31(6):546-9. doi: 10.1111/ajad.13343. PMID: 36184876. **Exclusion reason:** Ineligible study design
96. Fiani B, Sarhadi KJ, Soula M, et al. Current application of Cannabidiol (CBD) in the management and treatment of neurological disorders. *Neurol Sci.* 2020 Nov;41(11):3085-98. doi: 10.1007/s10072-020-04514-2. PMID: 32556748. **Exclusion reason:** Background
97. Filippini G, Minozzi S, Borrelli F, et al. Cannabis and Cannabinoids for symptomatic treatment for people with Multiple Sclerosis. *The Cochrane database of systematic reviews.* 2022;5:CD013444. doi: 10.1002/14651858.CD013444.pub2. PMID: 35510826. **Exclusion reason:** Systematic review used as source document
98. First L, Douglas W, Habibi B, et al. Cannabis use and low-back pain: a systematic review. *Cannabis Cannabinoid Res.* 2020;5(4):283-9. doi: 10.1089/can.2019.0077. PMID: 33381642. **Exclusion reason:** Systematic review used as source document
99. Fishbain DA, Cutler RB, Rosomoff HL, et al. Validity of self-reported drug use in chronic pain patients. *Clin J Pain.* 1999 Sep;15(3):184-91. doi: 10.1097/00002508-199909000-00005. PMID: 10524471. **Exclusion reason:** Background
100. Fisher E, Moore RA, Fogarty AE, et al. Cannabinoids, Cannabis, and Cannabis-based medicine for pain management: a systematic review of randomised controlled trials. *Pain.* 2021 Jul 1;162(Suppl 1):S45-s66. doi: 10.1097/j.pain.0000000000001929. PMID: 32804836. **Exclusion reason:** Systematic review used as source document
101. Fitzcharles M-A, Rampakakis E, Sampalis J, et al. Use of Medical Cannabis by patients with Fibromyalgia in Canada after Cannabis legalisation: a cross-sectional study. *Clin Exp Rheumatol.* 2021doi: 10.55563/clinexprheumatol/qcyet7. PMID: 33938797. **Exclusion reason:** Ineligible study design
102. Fitzcharles MA, Baerwald C, Ablin J, et al. Efficacy, tolerability and safety of Cannabinoids in chronic pain associated with rheumatic diseases (Fibromyalgia Syndrome, back pain, Osteoarthritis, Rheumatoid Arthritis): a systematic review of randomized controlled trials. *Schmerz.* 2016 Feb;30(1):47-61. doi: 10.1007/s00482-015-0084-3. PMID: 26767993. **Exclusion reason:** Ineligible publication type
103. Fitzcharles MA, Petzke F, Tolle TR, et al. Cannabis-based medicines and Medical Cannabis in the treatment of nociplastic pain. *Drugs.* 81(18):2103-16. doi: 10.1007/s40265-021-01602-1. PMID: 34800285. **Exclusion reason:** Ineligible publication type
104. Fitzcharles MA, Shir Y, Joseph L, et al. The effects of nabilone on insomnia in fibromyalgia: results of a randomized controlled trial. *Arthritis and rheumatism.* 2009;60:1429. doi: <https://doi.org/10.1002/art.26503>. **Exclusion reason:** Ineligible publication type
105. Fitzcharles MA, Ste-Marie PA, Hauser W, et al. Efficacy, tolerability, and safety of Cannabinoid treatments in the rheumatic diseases: a systematic review of randomized controlled trials. *Arthritis care & research.* 2016 May;68(5):681-8. doi: 10.1002/acr.22727. PMID: 26548380. **Exclusion reason:** Ineligible publication type
106. Flachenecker P, Henze T, Zettl UK. Nabiximols (THC/CBD oromucosal spray, Sativex®) in clinical practice--results of a multicenter, non-interventional study (MOVE 2) in patients with Multiple Sclerosis spasticity. *Eur Neurol.* 2014;71(5-6):271-9. doi: 10.1159/000357427. PMID: 24525548. **Exclusion reason:** Ineligible comparator

107. Flachenecker P, Henze T, Zettl UK. Long-term effectiveness and safety of Nabiximols (tetrahydrocannabinol/cannabidiol oromucosal spray) in clinical practice. *Eur Neurol.* 2014;72(1-2):95-102. doi: 10.1159/000360285. PMID: 24943098. **Exclusion reason:** Ineligible comparator
108. Gado F, Mohamed KA, Meini S, et al. Variously substituted 2-oxopyridine derivatives: extending the structure-activity relationships for allosteric modulation of the Cannabinoid CB2 receptor. *Eur J Med Chem.* 2020;211:113116. doi: 10.1016/j.ejmech.2020.113116. PMID: 33360803. **Exclusion reason:** Ineligible study design
109. Gambino A, Cabras M, Panagiotakos E, et al. Evaluating the suitability and potential efficiency of Cannabis Sativa oil for patients with primary Burning Mouth Syndrome: a prospective, open-label, single-arm pilot study. *Pain Med.* 2020doi: 10.1093/pm/pnaa318. PMID: 33123730. **Exclusion reason:** Ineligible comparator
110. Garmon EH, Olson K. Narrative review of Kratom, an emerging psychoactive substance with perianesthetic implications. *Anesthesia and analgesia.* 2022;XXXdoi: 10.1213/ANE.0000000000006177. PMID: 35986675. **Exclusion reason:** Ineligible publication type
111. Gedin F, Blome S, Ponten M, et al. Placebo Response and Media Attention in Randomized Clinical Trials Assessing Cannabis-Based Therapies for Pain: A Systematic Review and Meta-analysis. *JAMA Netw Open.* 2022;5(11):e2243848. doi: <https://dx.doi.org/10.1001/jamanetworkopen.2022.43848>. **Exclusion reason:** Systematic review used as source document
112. Gershoni T, Pud D, Aviram J, et al. Wellness of patients with chronic pain is not only about pain intensity. *Pain practice : the official journal of World Institute of Pain.* 2022;00:1-10. doi: 10.1111/papr.13168. PMID: 36181347. **Exclusion reason:** Ineligible comparator
113. Gilman JM, Schuster RM, Potter KW, et al. Effect of medical marijuana card ownership on pain, Insomnia, and Affective Disorder symptoms in adults: a randomized clinical trial. *JAMA Netw Open.* 2022;5(3):e222106. doi: 10.1001/jamanetworkopen.2022.2106. PMID: 35302633. **Exclusion reason:** Ineligible intervention
114. Giossi R, Carrara F, Padroni M, et al. Systematic review and meta-analysis seem to indicate that Cannabinoids for chronic primary pain treatment have limited benefit. *Pain Ther.* 2022doi: 10.1007/s40122-022-00434-5. PMID: 36129666. **Exclusion reason:** Systematic review used as source document
115. Goedel WC, Macmadu A, Shihpar A, et al. Association of medical cannabis licensure with prescription opioid receipt: A population-based, individual-level retrospective cohort study. *Int J Drug Policy.* 2021;100:103502. doi: 10.1016/j.drugpo.2021.103502. PMID: 34695720. **Exclusion reason:** Ineligible comparator
116. Greis A, Larsen E, Liu C, et al. Perceived efficacy, reduced prescription drug use, and minimal side effects of Cannabis in patients with chronic orthopedic pain. *Cannabis Cannabinoid Res.* 2021;12:12. doi: 10.1089/can.2021.0088. PMID: 34767730. **Exclusion reason:** Ineligible comparator
117. Greis A, Renslo B, Wilson-Poe AR, et al. Medical Cannabis use reduces opioid prescriptions in patients with chronic back pain. *Cureus.* 2022;14(1):e21452. doi: 10.7759/cureus.21452. PMID: 35223236. **Exclusion reason:** Ineligible comparator
118. Grossman S, Tan H, Gadiwalla Y. Cannabis and orofacial pain: a systematic review. *Br J Oral Maxillofac Surg.* 2021doi: 10.1016/j.bjoms.2021.06.005. PMID: 35305839. **Exclusion reason:** Systematic review used as source document
119. Grotenhermen F. Treatment of severe chronic pain with Cannabis preparations. *Arztl Prax Neurol Psychiatr.* 2002(5):28-30. **Exclusion reason:** Not in English

120. Guillooard M, Authier N, Pereira B, et al. Cannabis use assessment and its impact on pain in Rheumatologic diseases: a systematic review and meta-analysis. *Rheumatology (Oxford)*. 2020doi: 10.1093/rheumatology/keaa534. PMID: 33159797. **Exclusion reason:** Systematic review used as source document
121. Gundin JS, Rubio-Valera M, Romero LG, et al. Off-label use of Cannabinoids efficacy and safety. *Eur J Clin Pharm*. 2017;19(3):158-63. **Exclusion reason:** Ineligible study design
122. Gustavsen S, Hb S, K L, et al. Safety and efficacy of low-dose Medical Cannabis oils in Multiple Sclerosis. *Mult Scler Relat Disord*. 2020;48:102708. doi: 10.1016/j.msard.2020.102708. PMID: 33387864. **Exclusion reason:** Ineligible outcome
123. Gutierrez T, Hohmann AG. Cannabinoids for the treatment of neuropathic pain: Are they safe and effective? *Future Neurol*. 2011;6(2):129-33. doi: 10.2217/fnl.11.6. **Exclusion reason:** Ineligible publication type
124. Habib G, Khazin F, Artul S. The effect of Medical Cannabis on pain level and quality of sleep among Rheumatology clinic outpatients. *Pain Res Manag*. 2021;2021:1756588. doi: 10.1155/2021/1756588. PMID: 34531934. **Exclusion reason:** Ineligible comparator
125. Häckel A. Cannabis for chronic back pain?: Pivotal study for whole Cannabis extract started. *MMW Fortschr Med*. 2021;163(14):63. doi: 10.1007/s15006-021-0197-9. **Exclusion reason:** Ineligible publication type
126. Haddad F, Dokmak G, Karaman R. The efficacy of Cannabis on Multiple Sclerosis-related symptoms. *Life (Basel, Switzerland)*. 2022;12(5)doi: 10.3390/life12050682. PMID: 35629350. **Exclusion reason:** Ineligible publication type
127. Haleem R, Wright R. A scoping review on clinical trials of pain reduction with Cannabis administration in adults. *J Clin Med Res*. 2020 Jun;12(6):344-51. doi: 10.14740/jocmr4210. PMID: 32587650. **Exclusion reason:** Ineligible population
128. Hall N, James B, Bhuiyan MAN, et al. Topical cannabidiol is well tolerated in individuals with a history of elite physical performance and chronic lower extremity pain. *J Cannabis Res*. 2023;5(1):11. doi: https://dx.doi.org/10.1186/s42238-023-00179-8. **Exclusion reason:** Ineligible comparator
129. Hameed M, Prasad S, Jain E, et al. Medical Cannabis for Chronic Nonmalignant Pain Management. *Current pain and headache reports*. 2023doi: https://dx.doi.org/10.1007/s11916-023-01101-w. **Exclusion reason:** Ineligible publication type
130. Hansen JS, Hansen RM, Petersen T, et al. The effect of Cannabis-based medicine on neuropathic pain and spasticity in patients with Multiple Sclerosis and spinal cord injury: study protocol of a national multicenter double-blinded, placebo-controlled trial. *Brain sci*. 2021;11(9)doi: 10.3390/brainsci11091212. PMID: 34573231. **Exclusion reason:** Ineligible publication type
131. Hansen JS, Kasch H, Finnerup NB, et al. The effect of medical cannabis on neuropathic pain and spasticity in patients with multiple sclerosis and spinal cord injury: a danish multicentre double-blinded placebo-controlled study. *Med Cannabis Cannabinoids*. 2019;2(2):79. doi: https://doi.org/10.1159/000500623. **Exclusion reason:** Ineligible publication type
132. Haroutounian S, Arendt-Nielsen L, Belton J, et al. International association for the study of pain presidential task force on Cannabis and Cannabinoid Analgesia: research agenda on the use of Cannabinoids, Cannabis, and Cannabis-based medicines for pain management. *Pain*. 2021 Jul 01;162(Suppl 1):S117-S24. doi: 10.1097/j.pain.0000000000002266. PMID: 34138827. **Exclusion reason:** Background
133. Harris M, Erridge S, Ergisi M, et al. UK Medical Cannabis Registry: an analysis of clinical outcomes of Medicinal Cannabis therapy for chronic pain conditions. *Expert Rev Clin Pharmacol*. 2021:1-13. doi: 10.1080/17512433.2022.2017771. PMID: 34937477. **Exclusion reason:** Ineligible comparator

134. Hassan S, Zheng Q, Rizzolo E, et al. Does integrative medicine reduce prescribed opioid use for chronic pain? A systematic literature review. *Pain Med.* 2020 Apr 01;21(4):836-59. doi: 10.1093/pm/pnz291. PMID: 31755962. **Exclusion reason:** Ineligible intervention
135. Haungs A, Elizondo J. Does smoking Cannabis help with chronic neuropathic pain? *Evid Based Pract.* 2018;21(2):E7-E8. doi: 10.1097/01.EBP.0000541985.24333.b1. **Exclusion reason:** Ineligible publication type
136. Hauser W, Fitzcharles M-A. Register-based studies on Cannabis-based medicines and Medical Cannabis need reliable diagnoses and Cannabis treatment details. *Eur J Pain.* 2022;26(1):3-4. doi: 10.1002/ejp.1881. PMID: 34743399. **Exclusion reason:** Ineligible publication type
137. Hauser W, Fitzcharles M-A, Radbruch L, et al. Cannabinoids in pain management and palliative medicine: an overview of systematic reviews and prospective observational studies. *Dtsch Arztebl Int.* 2017 Sep;114(38):627-34. doi: 10.3238/arztebl.2017.0627. PMID: 29017688. **Exclusion reason:** Systematic review used as source document
138. Hauser W, Fitzcharles MA, Radbruch L, et al. Cannabinoids in Pain Management and Palliative Medicine. *Dtsch Arztebl Int.* 2017 Sep 22;114(38):627-34. doi: 10.3238/arztebl.2017.0627. PMID: 29017688. **Exclusion reason:** Ineligible population
139. Hayes C, Martin JH. Lack of efficacy of Cannabidiol for relieving back pain: time to re-set expectations? *Med J Aust.* 2021doi: 10.5694/mja2.51025. PMID: 33846981. **Exclusion reason:** Ineligible publication type
140. Hefner K, Sofuoglu M, Rosenheck R. Concomitant Cannabis abuse/dependence in patients treated with opioids for non-cancer pain. *The American journal on addictions.* 2015 Sep;24(6):538-45. doi: 10.1111/ajad.12260. PMID: 26246069. **Exclusion reason:** Ineligible outcome
141. Heineman JT, Forster GL, Stephens KL, et al. A randomized controlled trial of topical Cannabidiol for the treatment of thumb basal joint arthritis. *J Hand Surg Am.* 2022 May 27doi: 10.1016/j.jhsa.2022.03.002. PMID: 35637038. **Exclusion reason:** Inadequate duration
142. Hendricks O, Andersen TE, Christiansen AA, et al. Efficacy and safety of Cannabidiol followed by an open label add-on of tetrahydrocannabinol for the treatment of chronic pain in patients with Rheumatoid Arthritis or Ankylosing Spondylitis: protocol for a multicentre, randomised, placebo-controlled study. *BMJ Open.* 2019 Jun 04;9(6):e028197. doi: 10.1136/bmjopen-2018-028197. PMID: 31167870. **Exclusion reason:** Ineligible study design
143. Hershkovich O, Hayun Y, Oscar N, et al. The role of cannabis in treatment-resistant fibromyalgia women. *Pain practice : the official journal of World Institute of Pain.* 2022doi: <https://dx.doi.org/10.1111/papr.13179>. **Exclusion reason:** Ineligible study design
144. Hesselink JM, Kopsky DJ. Enhancing acupuncture by low dose Naltrexone. *Acupunct Med.* 2011 Jun;29(2):127-30. doi: 10.1136/aim.2010.003566. PMID: 21415049. **Exclusion reason:** Ineligible publication type
145. Hill KP, Hurley-Welljams-Dorof WM. Low to moderate quality evidence demonstrates the potential benefits and adverse events of Cannabinoids for certain medical indications. *Evid Based Med.* 2016 Feb;21(1):17. doi: 10.1136/ebmed-2015-110264. PMID: 26490847. **Exclusion reason:** Ineligible publication type
146. Hill KP, Palastro MD, Johnson B, et al. Cannabis and pain: a clinical review. *Cannabis Cannabinoid Res.* 2017;2(1):96-104. doi: 10.1089/can.2017.0017. PMID: 28861509. **Exclusion reason:** Systematic review used as source document

147. Hjorthøj C, La Cour P, Nordentoft M, et al. Cannabis-based medicines and Medical Cannabis for patients with neuropathic pain and other pain disorders: nationwide register-based pharmacoepidemiologic comparison with propensity score matched controls. *Eur J Pain*. 2021doi: 10.1002/ejp.1874. PMID: 34624164. **Exclusion reason:** Ineligible outcome
148. Hjorthøj C, La Cour P, Nordentoft M, et al. Cannabis-based medicines and Medical Cannabis for patients with neuropathic pain and other pain disorders: nationwide register-based pharmacoepidemiologic comparison with propensity score matched controls. *Eur J Pain*. 2022;26(2):480-91. doi: 10.1002/ejp.1874. PMID: 34624164. **Exclusion reason:** Ineligible population
149. Hoggart B, Ratcliffe S, Ehler E, et al. A multicentre, open-label, follow-on study to assess the long-term maintenance of effect, tolerance and safety of THC/CBD oromucosal spray in the management of neuropathic pain. *J Neurol*. 2015 Jan;262(1):27-40. doi: 10.1007/s00415-014-7502-9. PMID: 25270679. **Exclusion reason:** Ineligible study design
150. Hojsted J, Ekholm O, Kurita GP, et al. Addictive behaviors related to opioid use for chronic pain: a population-based study. *Pain*. 2013;154(12):2677-83. doi: 10.1016/j.pain.2013.07.046. PMID: 23906554. **Exclusion reason:** Ineligible intervention
151. Holdcroft A, Smith M, Jacklin A, et al. Pain relief with oral Cannabinoids in familial Mediterranean fever. *Anaesthesia*. 1997 May;52(5):483-6. doi: 10.1111/j.1365-2044.1997.139-az0132.x. PMID: 9165969. **Exclusion reason:** Ineligible study design
152. Horsted T, Lichon Hesthaven K, Leutscher PDC. Safety and Effectiveness of Cannabinoids to Danish Patients with Treatment Refractory Chronic Pain - A Retrospective Observational Real-world Study. *Eur J Pain*. 2022doi: <https://dx.doi.org/10.1002/ejp.2054>. **Exclusion reason:** Ineligible study design
153. Huang IC, Alberts NM, Buckley MG, et al. Change in pain status and subsequent opioid and marijuana use among long-term adult survivors of childhood cancer. *JNCI Cancer Spectr*. 2020;4(6):pkaa070. doi: 10.1093/jncics/pkaa070. PMID: 33409451. **Exclusion reason:** Ineligible study design
154. Hughes PR, Nwokocha J. Medical Cannabis or Cannabinoids for chronic pain: BMJ rapid recommendation. *American Family Physician*. 2022;106(2):208-9. PMID: 35977118. **Exclusion reason:** Ineligible publication type
155. Hunter D, Oldfield G, Tich N, et al. Synthetic transdermal cannabidiol for the treatment of knee pain due to osteoarthritis. *Osteoarthritis Cartilage*. 2018;26:S26. doi: 10.1016/j.joca.2018.02.067. **Exclusion reason:** Ineligible publication type
156. Hwang JK, Clarke H. Cannabis and pain: a review. *J Pain Manag*. 2016;9(4):395-413. **Exclusion reason:** Ineligible publication type
157. Iskedjian M, Bereza B, Gordon A, et al. Meta-analysis of Cannabis based treatments for neuropathic and Multiple Sclerosis-related pain. *Curr Med Res Opin*. 2007 Jan;23(1):17-24. doi: 10.1185/030079906x158066. PMID: 17257464. **Exclusion reason:** Ineligible publication type
158. Issa MA, Narang S, Jamison RN, et al. The subjective psychoactive effects of oral dronabinol studied in a randomized, controlled crossover clinical trial for pain. *Clin J Pain*. 2014 Jun;30(6):472-8. doi: 10.1097/AJP.000000000000022. PMID: 24281276. **Exclusion reason:** Inadequate duration
159. Jain N, Moorthy A. Cannabinoids in Rheumatology: friend, foe or a bystander? *Musculoskeletal care*. 2022doi: 10.1002/msc.1636. PMID: 35476898. **Exclusion reason:** Ineligible publication type

160. Jashinski J, Grossman E, Quaye A, et al. Randomised, pragmatic, waitlist controlled trial of Cannabis added to prescription opioid support on opioid dose reduction and pain in adults with chronic non-cancer pain: study protocol. *BMJ Open*. 2022;12(6):e064457. doi: 10.1136/bmjopen-2022-064457. PMID: 35680252. **Exclusion reason:** Ineligible publication type
161. Jawahar R, Oh U, Yang S, et al. A systematic review of pharmacological pain management in Multiple Sclerosis. *Drugs*. 2013 Oct;73(15):1711-22. doi: 10.1007/s40265-013-0125-0. PMID: 24085618. **Exclusion reason:** Systematic review used as source document
162. Jensen TS, Madsen CS, Finnerup NB. Pharmacology and treatment of neuropathic pains. *Curr Opin Neurol*. 2009 Oct;22(5):467-74. doi: 10.1097/WCO.0b013e3283311e13. PMID: 19741531. **Exclusion reason:** Ineligible publication type
163. Johal H, Devji T, Chang Y, et al. Cannabinoids in chronic non-cancer pain: a systematic review and meta-analysis. *Clin Med Insights Arthritis Musculoskelet Disord*. 2020 Feb 19;13:1179544120906461. doi: 10.1177/1179544120906461. PMID: 32127750. **Exclusion reason:** Systematic review used as source document
164. Jugl S, Okpeku A, Costales B, et al. A mapping literature review of Medical Cannabis clinical outcomes and quality of evidence in approved conditions in the USA from 2016 to 2019. *Med Cannabis Cannabinoids*. 2021;4(1):21-42. doi: 10.1159/000515069. PMID: 34676348. **Exclusion reason:** Systematic review used as source document
165. Kafil TS, Nguyen TM, MacDonald JK, et al. Cannabis for the treatment of Ulcerative Colitis. *Cochrane Database Syst Rev*. 2018 Nov 08;11:CD012954. doi: 10.1002/14651858.CD012954.pub2. PMID: 30406638. **Exclusion reason:** Ineligible population
166. Karst M, Salim K, Burstein S, et al. Analgesic effect of the synthetic cannabinoid CT-3 on chronic neuropathic pain: a randomized controlled trial. *Jama*. 2003 Oct 01;290(13):1757-62. doi: 10.1001/jama.290.13.1757. PMID: 14519710. **Exclusion reason:** Inadequate duration
167. Kaskie B, Kang H, Bhagianadh D, et al. Cannabis use among older persons with Arthritis, Cancer and Multiple Sclerosis: are we comparing apples and oranges? *Brain sci*. 2021;11(5)doi: 10.3390/brainsci11050532. PMID: 33922425. **Exclusion reason:** Ineligible study design
168. Kawka M, Erridge S, Holvey C, et al. Clinical outcome data of first cohort of chronic pain patients treated with Cannabis-based sublingual oils in the United Kingdom - analysis from the UK Medical Cannabis Registry. *J Clin Pharmacol*. 2021doi: 10.1002/jcph.1961. PMID: 34473850. **Exclusion reason:** Ineligible comparator
169. Khurshid H, Qureshi IA, Jahan N, et al. A systematic review of Fibromyalgia and recent advancements in treatment: is Medicinal Cannabis a new hope? *Cureus*. 2021;13(8):e17332. doi: 10.7759/cureus.17332. PMID: 34567876. **Exclusion reason:** Systematic review used as source document
170. Kinnucan JAR, Swaminath A, Sangha MS, et al. Real world dispensary data for IBD patients using medical cannabis. *Am J Gastroenterol*. 2021;116(SUPPL):S436. doi: <https://doi.org/10.14309/01.ajg.0000777220.73380.41>. **Exclusion reason:** Ineligible publication type
171. Kliuk-Ben Bassat O, Brill S, Vaisman N, et al. POS-676 exploratory study to assess the safety of sublingual oil based Medical Cannabis BOL-DP-O-04 in dialysis patients who have chronic pain. *Kidney international reports*. 2021;6(4):S295. doi: 10.1016/j.ekir.2021.03.707. **Exclusion reason:** Ineligible publication type

172. Kocot-Kepska M, Zajackowska R, Mika J, et al. Topical treatments and their molecular/cellular mechanisms in patients with peripheral neuropathic pain-narrative review. *Pharmaceutics*. 2021;13(4)doi: 10.3390/pharmaceutics13040450. PMID: 33810493. **Exclusion reason:** Ineligible publication type
173. Korownyk CS, Montgomery L, Young J, et al. PEER simplified chronic pain guideline: management of chronic low back, osteoarthritic, and neuropathic pain in primary care. *Can Fam Physician*. 2022;68(3):179-90. doi: 10.46747/cfp.6803179. PMID: 35292455. **Exclusion reason:** Systematic review used as source document
174. Kruger DJ, Kruger JS. Consumer experiences with Delta-8-THC: medical use, pharmaceutical substitution, and comparisons with Delta-9-THC. *Cannabis Cannabinoid Res*. 2021;19:19. doi: 10.1089/can.2021.0124. PMID: 34797727. **Exclusion reason:** Ineligible study design
175. Kurlyandchik I, Tiralongo E, Schloss J. Safety and efficacy of Medicinal Cannabis in the treatment of Fibromyalgia: a systematic review. *J Altern Complement Med*. 2020;27(3):198-213. doi: 10.1089/acm.2020.0331. PMID: 33337931. **Exclusion reason:** Systematic review used as source document
176. Lake S, Walsh Z, Kerr T, et al. Frequency of Cannabis and illicit opioid use among people who use drugs and report chronic pain: a longitudinal analysis. *PLoS Med*. 2019 Nov 19;16(11):e1002967. doi: 10.1371/journal.pmed.1002967. PMID: 31743343. **Exclusion reason:** Ineligible study design
177. Lamb LC, Di Fiori MM, Feeney JM. Cannabis to reduce opiate use in patients with acute rib fractures. *Journal of the American College of Surgeons*. 2018;227(4):S259. doi: 10.1016/j.jamcollsurg.2018.07.531. **Exclusion reason:** Ineligible publication type
178. Lee G, Grovey B, Furnish T, et al. Medical Cannabis for neuropathic pain. *Curr Pain Headache Rep*. 2018 Feb 01;22(1):8. doi: 10.1007/s11916-018-0658-8. PMID: 29388063. **Exclusion reason:** Systematic review used as source document
179. Liang AL, Gingham EL, Coleman JS. Medical Cannabis for Gynecologic pain conditions: a systematic review. *Obstet Gynecol*. 2022 Feb 01;139(2):287-96. doi: 10.1097/AOG.0000000000004656. PMID: 35104069. **Exclusion reason:** Ineligible intervention
180. Lichtman AH, Lux EA, McQuade R, et al. Results of a double-blind, randomized, placebo-controlled study of Nabiximols Oromucosal Spray as an adjunctive therapy in advanced cancer patients with chronic uncontrolled pain. *J Pain Symptom Manage*. 2017(pagination)doi: 10.1016/j.jpainsymman.2017.09.001. PMID: 28923526. **Exclusion reason:** Ineligible population
181. Lichtman AH, Lux EA, McQuade R, et al. Results of a double-blind, randomized, placebo-controlled study of nabiximols oromucosal spray as an adjunctive therapy in advanced cancer patients with chronic uncontrolled pain. *J Pain Symptom Manage*. 2018 Feb;55(2):179-88.e1. doi: 10.1016/j.jpainsymman.2017.09.001. PMID: 28923526. **Exclusion reason:** Ineligible population
182. Lick D, Mirza N, Trevor R. Are cannabinoids effective for treatment of noncancer musculoskeletal pain? *Evid Based Pract*. 2022;25(7):40-1. doi: 10.1097/EBP.0000000000001631. PMID: 32655704. **Exclusion reason:** Ineligible publication type
183. Ling H-Q, Chen Z-H, He L, et al. Comparative efficacy and safety of 11 drugs as therapies for adults with neuropathic pain after spinal cord injury: a bayesian network analysis based on 20 randomized controlled trials. *Frontiers in neurology*. 2022;13:818522. doi: 10.3389/fneur.2022.818522. PMID: 35386408. **Exclusion reason:** Systematic review used as source document

184. Link K, Knowles LM, Alschuler KN, et al. Characterizing cannabis use in a sample of adults with multiple sclerosis and chronic pain: An observational study. *Mult Scler Relat Disord*. 2023;75doi: 10.1016/j.msard.2023.104742. **Exclusion reason:** Ineligible outcome
185. Longo R, Oudshoorn A, Befus D. Cannabis for chronic pain: a rapid systematic review of randomized control trials. *Pain Manag Nurs*. 2020;22(2):141-9. doi: 10.1016/j.pmn.2020.11.006. PMID: 33353819. **Exclusion reason:** Systematic review used as source document
186. Longworth J, Szafron M, Gruza A, et al. Cannabis and cannabinoid medications for the treatment of chronic orofacial pain: A scoping review: Cannabinoids and Chronic Orofacial Pain. *Dentistry Review*. 2023;3(1)doi: 10.1016/j.dentre.2023.100063. **Exclusion reason:** Systematic review used as source document
187. Lopez-Sendon Moreno JL, Garcia Caldentey J, Trigo Cubillo P, et al. A double-blind, randomized, cross-over, placebo-controlled, pilot trial with Sativex in Huntington's Disease. *J Neurol*. 2016;263(7):1390-400. doi: 10.1007/s00415-016-8145-9. PMID: 27159993. **Exclusion reason:** Ineligible population
188. Lucas P, Boyd S, Milloy MJ, et al. Cannabis significantly reduces the use of prescription opioids and improves quality of life in authorized patients: results of a large prospective study. *Pain Med*. 2020doi: 10.1093/pm/pnaa396. PMID: 33367882. **Exclusion reason:** Ineligible population
189. Luchetti M, Zanarella C, Moretti C, et al. Cannabinoids for the treatment of neuropathic pain. *Acta Anaesthesiol Ital*. 2008;59(2):187-95. **Exclusion reason:** Not in English
190. Lyes M, Yang KH, Castellanos J, et al. Microdosing psilocybin for chronic pain: a case series. *Pain*. 2022doi: 10.1097/j.pain.0000000000002778. PMID: 36066961. **Exclusion reason:** Ineligible study design
191. Lynch ME, Campbell F. Cannabinoids for treatment of chronic non-cancer pain; a systematic review of randomized trials. *Br J Clin Pharmacol*. 2011 Nov;72(5):735-44. doi: 10.1111/j.1365-2125.2011.03970.x. PMID: 21426373. **Exclusion reason:** Ineligible publication type
192. Lynch ME, Ware MA. Cannabinoids for the treatment of chronic non-cancer pain: an updated systematic review of randomized controlled trials. *J Neuroimmune Pharmacol*. 2015 Jun;10(2):293-301. doi: 10.1007/s11481-015-9600-6. PMID: 25796592. **Exclusion reason:** Ineligible publication type
193. Maayah ZH, Takahara S, Ferdaoussi M, et al. The anti-inflammatory and Analgesic effects of formulated full-spectrum Cannabis extract in the treatment of neuropathic pain associated with Multiple Sclerosis. *Inflamm Res*. 2020 Jun;69(6):549-58. doi: 10.1007/s00011-020-01341-1. PMID: 32239248. **Exclusion reason:** Ineligible publication type
194. MacCallum CA, Eadie L, Barr AM, et al. Practical strategies using Medical Cannabis to reduce harms associated with long term opioid use in chronic pain. *Front Pharmacol*. 2021;12doi: 10.3389/fphar.2021.633168. PMID: 33995035. **Exclusion reason:** Ineligible publication type
195. MacKay M, Zurier R, Kamen D, et al. A Phase 2, Double-blind, Randomized, Placebo-Controlled Multicenter Study to Evaluate Efficacy, Safety, and Tolerability of JBT-101/Lenabasum in Systemic Lupus Erythematosus, an Autoimmunity Centers of Excellence Study (ALE09). *Arthritis rheumatol*. 2022;74:735. doi: <https://doi.org/10.1002/art.42355>. **Exclusion reason:** Ineligible publication type
196. Maida V, Ennis M, Irani S, et al. Adjunctive Nabilone in cancer pain and symptom management: a prospective observational study using propensity scoring. *J Support Oncol*. 2008 Mar;6(3):119-24. PMID: 18402303. **Exclusion reason:** Ineligible population

197. Malik Z, Bayman L, Valestin J, et al. Dronabinol increases pain threshold in patients with functional chest pain: a pilot double-blind placebo-controlled trial. *Dis Esophagus*. 2017 Feb 1;30(2):1-8. doi: 10.1111/dote.12455. PMID: 26822791. **Exclusion reason:** Ineligible population
198. Marková J, Essner U, Akmaz B, et al. Sativex(®) as add-on therapy vs. further optimized first-line ANTispastics (SAVANT) in resistant Multiple Sclerosis spasticity: a double-blind, placebo-controlled randomised clinical trial. *Int J Neurosci*. 2019 Feb;129(2):119-28. doi: 10.1080/00207454.2018.1481066. PMID: 29792372. **Exclusion reason:** Ineligible population
199. Martin-Sanchez E, Furukawa TA, Taylor J, et al. Systematic review and meta-analysis of Cannabis treatment for chronic pain. *Pain Med*. 2009 Nov;10(8):1353-68. doi: 10.1111/j.1526-4637.2009.00703.x. PMID: 19732371. **Exclusion reason:** Ineligible publication type
200. Matarazzo AP, Elisei LMS, Carvalho FC, et al. Mucoadhesive nanostructured lipid carriers as a Cannabidiol nasal delivery system for the treatment of neuropathic pain. *Eur J Pharm Sci*. 2021;105698. doi: 10.1016/j.ejps.2020.105698. PMID: 33406408. **Exclusion reason:** Ineligible study design
201. Maurer M, Henn V, Dittrich A, et al. Delta-9-Tetrahydrocannabinol shows antispastic and analgesic effects in a single case double-blind trial. *Eur Arch Psychiatry Clin Neurosci*. 1990;240(1):1-4. doi: 10.1007/bf02190083. PMID: 2175265. **Exclusion reason:** Inadequate duration
202. Maurotti S, Mare R, Pujia R, et al. Hemp seeds in post-arthroplasty rehabilitation: a pilot clinical study and an in vitro investigation. *Nutrients*. 2021;13(12)doi: 10.3390/nu13124330. PMID: 34959882. **Exclusion reason:** Ineligible intervention
203. Mazza M. Medical Cannabis for the treatment of Fibromyalgia syndrome: a retrospective, open-label case series. *J Cannabis Res*. 2021;3(1):4. doi: 10.1186/s42238-021-00060-6. PMID: 33597032. **Exclusion reason:** Ineligible comparator
204. McDonagh MS, Selph SS, Buckley DI, et al. Nonopioid pharmacologic treatments for chronic pain. Agency for Healthcare Research and Quality (US). 2020 PMID: 32338847. **Exclusion reason:** Systematic review used as source document
205. McDonagh MS, Wagner J, Ahmed AY, et al. Living systematic review on Cannabis and other plant-based treatments for chronic pain - quarterly progress report: May 2021. Agency for Healthcare Research and Quality (US). 2020 12;12:12. PMID: 34228409. **Exclusion reason:** Ineligible publication type
206. McGinty EE, Tormohlen KN, Barry CL, et al. Protocol: mixed-methods study of how implementation of US state Medical Cannabis laws affects treatment of chronic non-cancer pain and adverse opioid outcomes. *Implement Sci*. 2021;16(1):2. doi: 10.1186/s13012-020-01071-2. PMID: 33413454. **Exclusion reason:** Ineligible publication type
207. McMichael BJ, Van Horn RL, Viscusi WK. The impact of Cannabis access laws on opioid prescribing. *J Health Econ*. 2020 Jan;69:102273. doi: 10.1016/j.jhealeco.2019.102273. PMID: 31865260. **Exclusion reason:** Ineligible population
208. McParland AL, Bhatia A, Matelski J, et al. Evaluating the impact of cannabinoids on sleep health and pain in patients with chronic neuropathic pain: a systematic review and meta-analysis of randomized controlled trials. *Regional anesthesia and pain medicine*. 2022doi: <https://dx.doi.org/10.1136/rapm-2021-103431>. **Exclusion reason:** Systematic review used as source document
209. Meade E, Hehir S, Rowan N, et al. Mycotherapy: potential of fungal bioactives for the treatment of mental health disorders and morbidities of chronic pain. *J Fungi (Basel)*. 2022;8(3)doi: 10.3390/jof8030290. PMID: 35330292. **Exclusion reason:** Systematic review used as source document
210. Mechtler L, Hart P, Bargnes V, et al. Medical cannabis treatment in patients with trigeminal neuralgia. *Neurology*. 2019;92(15). **Exclusion reason:** Ineligible publication type

211. Meng H, Johnston B, Englesakis M, et al. Selective Cannabinoids for chronic neuropathic pain: a systematic review and meta-analysis. *Anesth Analg*. 2017 Nov;125(5):1638-52. doi: 10.1213/ANE.0000000000002110. PMID: 28537982. **Exclusion reason:** Ineligible publication type
212. Meng H, Page MG, Ajrawat P, et al. Patient-reported outcomes in those consuming Medical Cannabis: a prospective longitudinal observational study in chronic pain patients. *Canadian journal of anaesthesia = Journal canadien d'anesthésie*. 2021doi: 10.1007/s12630-020-01903-1. PMID: 33469735. **Exclusion reason:** Ineligible population
213. Meuth SG, Henze T, Essner U, et al. Tetrahydrocannabinol and cannabidiol oromucosal spray in resistant Multiple Sclerosis spasticity: consistency of response across subgroups from the SAVANT randomized clinical trial. *International Journal of Neuroscience* 2020;130(12):1199-205. doi: 10.1080/00207454.2020.1730832. PMID: 32065006. **Exclusion reason:** Ineligible population
214. Mistry M, Simpson P, Morris E, et al. Cannabidiol for the Management of Endometriosis and Chronic Pelvic Pain. *J Minim Invasive Gynecol*. 2021;25:25. doi: 10.1016/j.jmig.2021.11.017. PMID: 34839061. **Exclusion reason:** Ineligible publication type
215. Mohiuddin M, Blyth FM, Degenhardt L, et al. General risks of harm with Cannabinoids, Cannabis, and Cannabis-based medicine possibly relevant to patients receiving these for pain management: an overview of systematic reviews. *Pain*. 2021 Jul 01;162(Suppl 1):S80-S96. doi: 10.1097/j.pain.0000000000002000. PMID: 32941319. **Exclusion reason:** Background
216. Montero-Oleas N, Arevalo-Rodriguez I, Nunez-Gonzalez S, et al. Therapeutic use of Cannabis and Cannabinoids: an evidence mapping and appraisal of systematic reviews. *BMC Complement Med Ther*. 2020 Jan 15;20(1):12. doi: 10.1186/s12906-019-2803-2. PMID: 32020875. **Exclusion reason:** Systematic review used as source document
217. Moore RA, Fisher E, Finn DP, et al. Cannabinoids, Cannabis, and Cannabis-based medicines for pain management: an overview of systematic reviews. *Pain*. 2021 Jul 1;162(Suppl 1):S67-s79. doi: 10.1097/j.pain.0000000000001941. PMID: 32804833. **Exclusion reason:** Systematic review used as source document
218. Moreno Torres I, Sanchez AJ, Garcia-Merino A. Evaluation of the tolerability and efficacy of Sativex in Multiple Sclerosis. *Expert Rev Neurother*. 2014 Nov;14(11):1243-50. doi: 10.1586/14737175.2014.971758. PMID: 25331416. **Exclusion reason:** Ineligible publication type
219. Moreno-Sanz G, Madiedo A, Hernandez P, et al. Sex-dependent prescription patterns and clinical outcomes associated with the use of two oral Cannabis formulations in the multimodal management of chronic pain patients in Colombia. *Frontiers in pain research (Lausanne, Switzerland)*. 2022;3:854795. doi: 10.3389/fpain.2022.854795. PMID: 35399153. **Exclusion reason:** Ineligible comparator
220. Moreno-Sanz G, Madiedo A, Lynskey M, et al. "Flower Power": Controlled Inhalation of THC-Predominant Cannabis Flos Improves Health-Related Quality of Life and Symptoms of Chronic Pain and Anxiety in Eligible UK Patients. *Biomedicines*. 2022;10(10)doi: 10.3390/biomedicines10102576. **Exclusion reason:** Ineligible study design
221. Mucke M, Phillips T, Radbruch L, et al. Cannabis-based medicines for chronic neuropathic pain in adults. *Cochrane Database Syst Rev*. 2018 Mar 07;3(3):CD012182. doi: 10.1002/14651858.CD012182.pub2. PMID: 29513392. **Exclusion reason:** Systematic review used as source document
222. Muller C, Reggio PH. An Analysis of the Putative CBD Binding Site in the Ionotropic Cannabinoid Receptors. *Front Cell Neurosci*. 2020;14:615811. doi: 10.3389/fncel.2020.615811. PMID: 33362478. **Exclusion reason:** Ineligible study design

223. Murff HJ. Review: weak evidence of benefits of Cannabis for chronic neuropathic pain; moderate to weak evidence of adverse effects. *Ann Intern Med.* 2017 Dec 19;167(12):JC62. doi: 10.7326/ACPJC-2017-167-12-062. PMID: 29255852. **Exclusion reason:** Ineligible publication type
224. Myers B, Geist T, Hart P, et al. Medical cannabis in the treatment of parkinson's disease. *Neurology.* 2019;92(15). **Exclusion reason:** Ineligible publication type
225. Naftali T, Mechoulam R, Gabay G, et al. Cannabidiol treatment does not effect active crohn's disease. *Gastroenterology.* 2013;144(5):S180. **Exclusion reason:** Ineligible publication type
226. Narang S, Gibson D, Wasan AD, et al. Efficacy of Dronabinol as an adjuvant treatment for chronic pain patients on opioid therapy. *J Pain.* 2008 Mar;9(3):254-64. doi: 10.1016/j.jpain.2007.10.018. PMID: 18088560. **Exclusion reason:** Ineligible study design
227. Neilson LM, Swift C, Swart ECS, et al. Impact of marijuana legalization on opioid utilization in patients diagnosed with pain. *J Gen Intern Med.* 2021doi: 10.1007/s11606-020-06530-6. PMID: 33575906. **Exclusion reason:** Background
228. Nguyen T, Li Y, Greene D, et al. Changes in Prescribed Opioid Dosages Among Patients Receiving Medical Cannabis for Chronic Pain, New York State, 2017-2019. *JAMA Netw Open.* 2023;6(1):e2254573. doi: <https://dx.doi.org/10.1001/jamanetworkopen.2022.54573>. **Exclusion reason:** Ineligible comparator
229. Nielsen S, Picco L, Murnion B, et al. Opioid-sparing effect of Cannabinoids for analgesia: an updated systematic review and meta-analysis of preclinical and clinical studies. *Neuropsychopharmacology* : official publication of the American College of Neuropsychopharmacology. 2022doi: 10.1038/s41386-022-01322-4. PMID: 35459926. **Exclusion reason:** Systematic review used as source document
230. Nielsen S, Picco L, Murnion B, et al. Opioid-sparing effect of Cannabinoids for analgesia: an updated systematic review and meta-analysis of preclinical and clinical studies. *Neuropsychopharmacology.* 2022;47(7):1315-30. doi: 10.1038/s41386-022-01322-4. PMID: 35459926. **Exclusion reason:** Systematic review used as source document
231. Noori A, Miroshnychenko A, Shergill Y, et al. Opioid-sparing effects of Medical Cannabis or Cannabinoids for chronic pain: a systematic review and meta-analysis of randomised and observational studies. *BMJ Open.* 2021 07 28;11(7):e047717. doi: 10.1136/bmjopen-2020-047717. PMID: 34321302. **Exclusion reason:** Systematic review used as source document
232. Novotna A, Mares J, Ratcliffe S, et al. A randomized, double-blind, placebo-controlled, parallel-group, enriched-design study of Nabiximols* (Sativex), as add-on therapy, in subjects with refractory spasticity caused by Multiple Sclerosis. *Eur J Neurol.* 2011;18(9):1122-31. doi: 10.1111/j.1468-1331.2010.03328.x. PMID: 21362108. **Exclusion reason:** Ineligible outcome
233. Nowell WB, Gavigan K, L Silverman S. Cannabis for Rheumatic Disease pain: a review of current literature. *Curr Rheumatol Rep.* 2022;24(5):119-31. doi: 10.1007/s11926-022-01065-7. PMID: 35486218. **Exclusion reason:** Systematic review used as source document
234. Nugent SM, Kansagara D. The effects of Cannabis among adults with chronic pain. *Ann Intern Med.* 2018 Apr 03;168(7):525. doi: 10.7326/L17-0732. PMID: 29610910. **Exclusion reason:** Ineligible publication type
235. Nugent SM, Morasco BJ, O'Neil ME, et al. The effects of Cannabis among adults with chronic pain and an overview of general harms: a systematic review. *Ann Intern Med.* 2017 Sep 05;167(5):319-31. doi: 10.7326/M17-0155. PMID: 28806817. **Exclusion reason:** Systematic review used as source document

236. Nunnari P, Ladiana N, Ceccarelli G, et al. Long-term Cannabis-based oil therapy and pain medications prescribing patterns: an Italian observational study. *Eur Rev Med Pharmacol Sci.* 2022;26(4):1224-34. doi: 10.26355/eurrev_202202_28114. PMID: 35253178. **Exclusion reason:** Ineligible comparator
237. Nurmikko TJ, Serpell MG, Hoggart B, et al. A multi-centre, double-blind, randomized, controlled trial of oro-mucosal Cannabis based medicine in the treatment of neuropathic pain characterized by Allodynia. *Neurology.* no: PO6;64(Suppl 1):A374. **Exclusion reason:** Ineligible publication type
238. Nutt DJ, Phillips LD, Barnes MP, et al. A multicriteria decision analysis comparing pharmacotherapy for chronic neuropathic pain, including Cannabinoids and Cannabis-based medical products. *Cannabis Cannabinoid Res.* 2021 Mar 17;17:17. doi: 10.1089/can.2020.0129. PMID: 33998895. **Exclusion reason:** Ineligible study design
239. O'Connell M, Sandgren M, Frantzen L, et al. Medical Cannabis: effects on opioid and benzodiazepine requirements for pain control. *Ann Pharmacother.* 2019 Nov;53(11):1081-6. doi: 10.1177/1060028019854221. PMID: 31129977. **Exclusion reason:** Ineligible comparator
240. Okusanya BO, Asaolu IO, Ehiri JE, et al. Medical Cannabis for the reduction of opioid dosage in the treatment of non-cancer chronic pain: a systematic review. *Syst Rev.* 2020 Jul 28;9(1):167. doi: 10.1186/s13643-020-01425-3. PMID: 32723354. **Exclusion reason:** Systematic review used as source document
241. Okusanya BO, Lott BE, Ehiri J, et al. Medical Cannabis for the treatment of migraine in adults: a review of the evidence. *Frontiers in Neurology.* 2022;13doi: 10.3389/fneur.2022.871187. PMID: 35711271. **Exclusion reason:** Systematic review used as source document
242. Pankratz T. Impact of telehealth versus traditional visits on pain management patients utilizing medical cannabis during the COVID-19 pandemic. Dissertation Abstracts International Section A: Humanities and Social Sciences. 2023;84(8-A):No-Specified. **Exclusion reason:** Ineligible publication type
243. Paulsingh CN, Mohamed MB, Elhaj MS, et al. The efficacy of marijuana use for pain relief in adults with Sickle Cell Disease: a systematic review. *Cureus.* 2022;14(5):e24962. doi: 10.7759/cureus.24962. PMID: 35706744. **Exclusion reason:** Systematic review used as source document
244. Pellesi L, Licata M, Verri P, et al. Pharmacokinetics and tolerability of oral Cannabis preparations in patients with medication overuse headache (MOH)—a pilot study. *Eur J Clin Pharmacol.* 2018;74(11):1427-36. doi: 10.1007/s00228-018-2516-3. PMID: 29980818. **Exclusion reason:** Ineligible study design
245. Perras C. Sativex for the management of Multiple Sclerosis symptoms. *Issues Emerg Health Technol.* 2005 Sep(72):1-4. PMID: 16317825. **Exclusion reason:** Ineligible publication type
246. Petzke F, Tolle T, Fitzcharles MA, et al. Cannabis-based medicines and Medical Cannabis for chronic neuropathic pain. *CNS Drugs.* 2021;21:21. doi: 10.1007/s40263-021-00879-w. PMID: 34802112. **Exclusion reason:** Systematic review used as source document
247. Pham MN, Hudnall MT, Nadler RB. Marijuana, lower urinary tract symptoms, and pain in the urologic patient. *Urology.* 2020 05;139:8-13. doi: 10.1016/j.urology.2020.01.029. PMID: 32027882. **Exclusion reason:** Systematic review used as source document
248. Phillips KT, Pedula KL, Choi NG, et al. Chronic health conditions, acute health events, and healthcare utilization among adults over age 50 in Hawai'i who use Cannabis: a matched cohort study. *Drug Alcohol Depend.* 2022;234:109387. doi: 10.1016/j.drugalcdep.2022.109387. PMID: 35279458. **Exclusion reason:** Ineligible population

249. Phillips TJC, Cherry CL, Cox S, et al. A metanalysis of randomised controlled clinical trials assessing the effectiveness of pharmacological interventions for painful HIV-associated sensory neuropathy. *European journal of pain supplements*. 2010;4(1):141. doi: [https://doi.org/10.1016/S1754-3207\(10\)70503-4](https://doi.org/10.1016/S1754-3207(10)70503-4). **Exclusion reason:** Systematic review used as source document
250. Philpot LM, Ramar P, Jatoi A, et al. Cannabis in Cancer Survivors Who Report High Impact Chronic Pain: Findings from a 1500+ Patient Survey. *American Journal of Hospice and Palliative Medicine*. 2022doi: 10.1177/10499091221143098. **Exclusion reason:** Ineligible study design
251. Pichini S, Pacifici R, Busardo FP, et al. The challenge of clinical application of FM2 Cannabis oil produced in Italy for the treatment of neuropathic pain. *Eur Rev Med Pharmacol Sci*. 2018 Feb;22(4):863-5. doi: 10.26355/eurrev_201802_14363. PMID: 29509231. **Exclusion reason:** Ineligible publication type
252. Pinsger M, Schimetta W, Volc D, et al. Benefits of an add-on treatment with the synthetic cannabinomimetic nabilone on patients with chronic pain--a randomized controlled trial. *Wien Klin Wochenschr*. 2006;118(11-12):327-35. doi: 10.1007/s00508-006-0611-4. PMID: 16855921. **Exclusion reason:** Not in English
253. Pinsger M, Schimetta W, Volc D, et al. Benefits of an add-on treatment with the synthetic cannabinomimetic nabilone on patients with chronic pain - A randomized controlled trial. *Wien Klin Wochenschr*. 2006;118(11-12):327-35. doi: 10.1007/s00508-006-0611-4. PMID: 16855921. **Exclusion reason:** Not in English
254. Podda G, Constantinescu CS. Nabiximols in the treatment of spasticity, pain and urinary symptoms due to Multiple Sclerosis. *Expert Opin Biol Ther*. 2012 Nov;12(11):1517-31. doi: 10.1517/14712598.2012.721765. PMID: 22954177. **Exclusion reason:** Ineligible publication type
255. Poli P, Carnevale S, Scocca A, et al. Promising health benefits of adjuvant Acmeella and Zingiber extracts combined with Coenzyme Q10 Phytosomes, supplementation in chronic pain treated with Medical Cannabis: a prospective and open-label clinical study. *Evidence-based complementary and alternative medicine : eCAM*. 2022;2022:7099161. doi: 10.1155/2022/7099161. PMID: 35733629. **Exclusion reason:** Ineligible comparator
256. Portenoy RK, Ganae-Motan ED, Allende S, et al. Nabiximols for opioid-treated cancer patients with poorly-controlled chronic pain: a randomized, placebo-controlled, graded-dose trial. *J Pain*. 2012 May;13(5):438-49. doi: 10.1016/j.jpain.2012.01.003. PMID: 22483680. **Exclusion reason:** Ineligible population
257. Prevede E, Hupli A, Marrinan S, et al. Exploring the use of Kratom (*Mitragyna Speciosa*) via the YouTube data tool: A novel netnographic analysis. *Emerg Trends Drugs Addict Health*. 2021 2021/01/01/;1:100007. doi: 10.1016/j.etdah.2021.100007. **Exclusion reason:** Background
258. Price RL, Charlot KV, Frieler S, et al. The efficacy of Cannabis in reducing back pain: a systematic review. *Global Spine J*. 2022doi: 10.1177/21925682211065411. PMID: 35128969. **Exclusion reason:** Systematic review used as source document
259. Prieto Gonzalez JM, Vila Silvan C. Safety and tolerability of nabiximols oromucosal spray: a review of real-world experience in observational studies, registries and case reports. *Expert Rev Neurother*. 2021doi: 10.1080/14737175.2021.1904896. PMID: 33749480. **Exclusion reason:** Systematic review used as source document
260. Prieto Gonzalez JM, Vila Silvan C. Safety and tolerability of nabiximols oromucosal spray: a review of more than 15 years' accumulated evidence from clinical trials. *Expert Rev Neurother*. 2021 Jun 07;07:07. doi: 10.1080/14737175.2021.1935879. PMID: 34092180. **Exclusion reason:** Systematic review used as source document

261. Prozialeck WC, Lamar PC, Krupp M, 2nd, et al. Kratom use within the context of the evolving Opioid Crisis and the COVID-19 Pandemic in the United States. *Front Pharmacol.* 2021;12:729220. doi: 10.3389/fphar.2021.729220. PMID: 34512353. **Exclusion reason:** Background
262. Quintero J-M, Pulido G, Giraldo L-F, et al. A systematic review on Cannabinoids for neuropathic pain administered by routes other than oral or inhalation. *Plants (Basel, Switzerland).* 2022;11(10)doi: 10.3390/plants11101357. PMID: 35631782. **Exclusion reason:** Systematic review used as source document
263. Rabgay K, Waranuch N, Chaiyakunapruk N, et al. The effects of Cannabis, Cannabinoids, and their administration routes on pain control efficacy and safety: a systematic review and network meta-analysis. *J Am Pharm Assoc (2003).* 2020 Jan - Feb;60(1):225-34.e6. doi: 10.1016/j.japh.2019.07.015. PMID: 31495691. **Exclusion reason:** Systematic review used as source document
264. Rapin L, Gamaoun R, El Hage C, et al. Cannabidiol use and effectiveness: real-world evidence from a Canadian medical cannabis clinic. *J Cannabis Res.* 2021;3(1)doi: 10.1186/s42238-021-00078-w. PMID: 34162446. **Exclusion reason:** Ineligible outcome
265. Reisdorf S. Analgesia: Cannabis for neuropathic pain. *MMW Fortschr Med.* 2020;162(7):58. doi: 10.1007/s15006-020-0397-8. **Exclusion reason:** Not in English
266. Renslo B, Greis A, Liu CS, et al. Medical Cannabis use reduces opioid prescriptions in patients with Osteoarthritis. *Cureus.* 2022;14(1):e21564. doi: 10.7759/cureus.21564. PMID: 35228923. **Exclusion reason:** Ineligible comparator
267. Richards BL, Whittle SL, Buchbinder R. Neuromodulators for pain management in Rheumatoid Arthritis. *Cochrane Database Syst Rev.* 2012 Jan 18;1:CD008921. doi: 10.1002/14651858.CD008921.pub2. PMID: 22258992. **Exclusion reason:** Ineligible publication type
268. Richet E, Ferret L, Gaboriau L, et al. Use of dronabinol in the treatment of resistant neuropathic pain: Feedback from patients followed in a multidisciplinary pain center. *Annales Pharmaceutiques Francaises.* 2022doi: 10.1016/j.pharma.2022.12.005. **Exclusion reason:** Not in English
269. Riera R, Pacheco RL, Bagattini AM, et al. Efficacy and safety of therapeutic use of Cannabis derivatives and their synthetic analogs: overview of systematic reviews. *Phytother Res.* 2021;29:29. doi: 10.1002/ptr.7263. PMID: 34841610. **Exclusion reason:** Systematic review used as source document
270. Robinson D, Ritter S, Yassin M. Comparing Sublingual and Inhaled Cannabis Therapies for Low Back Pain: An Observational Open-Label Study. *Rambam Maimonides med.* 2022;13(4)doi: <https://dx.doi.org/10.5041/RMMJ.10485>. **Exclusion reason:** Ineligible study design
271. Rogers AH, Bakhshaie J, Buckner JD, et al. Opioid and Cannabis co-use among adults with chronic pain: relations to substance misuse, mental health, and pain experience. *J Addict Med.* 2019 Jul/Aug;13(4):287-94. doi: 10.1097/ADM.0000000000000493. PMID: 30557213. **Exclusion reason:** Ineligible study design
272. Ross J, Slawek DE, Zhang C, et al. First-year trajectories of Medical Cannabis use among adults taking opioids for chronic pain: an observational cohort study. *Pain Med.* 2021doi: 10.1093/pm/pnab257. PMID: 34411246. **Exclusion reason:** Background
273. Rouhollahi E, Macleod BA, Barr AM, et al. Cannabis extract CT-921 has a high efficacy– adverse effect profile in a neuropathic pain model. *Drug Des Devel Ther.* 2020;14:3351-61. doi: 10.2147/DDDT.S247584. PMID: 32884239 **Exclusion reason:** Ineligible population
274. Russo E. Cannabis and Cannabis based medicine extracts: additional results. *J Cannabis Ther.* 2004;3(4):153-61. doi: 10.1300/J175v03n04_03. **Exclusion reason:** Ineligible study design

275. Russo M, Naro A, Leo A, et al. Evaluating Sativex in neuropathic pain management: a clinical and neurophysiological assessment in Multiple Sclerosis. *Pain Med.* 2016 Jun;17(6):1145-54. doi: 10.1093/pm/pnv080. PMID: 26764336. **Exclusion reason:** Ineligible population
276. Safakish R, Ko G, Salimpour V, et al. Medical Cannabis for the management of pain and quality of life in chronic pain patients: a prospective observational study. *Pain Med.* 2020 Nov 01;21(11):3073-86. doi: 10.1093/pm/pnaa163. PMID: 32556203. **Exclusion reason:** Ineligible study design
277. Sagy I, Bar-Lev Schleider L, Abu-Shakra M, et al. Safety and efficacy of Medical Cannabis in Fibromyalgia. *J Clin Med.* 2019 Jun 05;8(6):807. doi: 10.3390/jcm8060807. PMID: 31195754. **Exclusion reason:** Ineligible comparator
278. Sainsbury B, Bloxham J, Pour MH, et al. Efficacy of Cannabis-based medications compared to placebo for the treatment of chronic neuropathic pain: a systematic review with meta-analysis. *J Dent Anesth Pain Med.* 2021;21(6):479-506. doi: 10.17245/jdapm.2021.21.6.479. PMID: 34909469. **Exclusion reason:** Systematic review used as source document
279. Salmasi V, Nelson LM, Hong J, et al. Association of cannabis and/or opioid with quality of life and healthcare utilization in patients with chronic pain. *Frontiers in pain research (Lausanne, Switzerland).* 2022;3:1015605. doi: <https://dx.doi.org/10.3389/fpain.2022.1015605>. **Exclusion reason:** Ineligible study design
280. Sanchez-Florez JC, Seija-Butnaru D, Valero EG, et al. Pain management strategies in Rheumatoid Arthritis: a narrative review. *J Pain Pall Care Pharmacother.* 2021;35(4):291-9. doi: 10.1080/15360288.2021.1973647. PMID: 34623946. **Exclusion reason:** Systematic review used as source document
281. Santos SA, Kontorinis N, Dieterich DT. Management of chronic Hepatitis C virus in patients with HIV. *Curr Treat Options Gastroenterol.* 2005;8(6):433-41. doi: 10.1007/s11938-005-0029-5. PMID: 16313860. **Exclusion reason:** Ineligible population
282. Schenk M. Chronic neuropathic pain: minimal side effects of therapy with Cannabis. *MMW Fortschr Med.* 2020;162(3):72. doi: 10.1007/s15006-020-0171-y. **Exclusion reason:** Not in English
283. Schindler E. Sustained reductions in headache burden after the limited administration of low dose psilocybin in migraine and cluster headache: results from two preliminary studies. *Neuropsychopharmacology.* 2020;45:57-8. doi: 10.1038/s41386-020-00889-0. **Exclusion reason:** Ineligible publication type
284. Schindler EAD, Sewell RA, Gottschalk CH, et al. Exploratory investigation of a patient-informed low-dose psilocybin pulse regimen in the suppression of cluster headache: Results from a randomized, double-blind, placebo-controlled trial. *Headache.* 2022doi: <https://dx.doi.org/10.1111/head.14420>. **Exclusion reason:** Ineligible intervention
285. Schloss J, Lacey J, Sinclair J, et al. A phase 2 randomised clinical trial assessing the tolerability of two different ratios of Medicinal Cannabis in patients with high grade gliomas. *Front Oncol.* 2021;11doi: 10.3389/fonc.2021.649555. PMID: 34094937. **Exclusion reason:** Ineligible population
286. Schrader NHB, Duipmans JC, Renken RJ, et al. The C4EB study-Transvamix (10% THC / 5% CBD) to treat chronic pain in epidermolysis bullosa: A protocol for an explorative randomized, placebo controlled, and double blind intervention crossover study. *PLoS ONE.* 2022;17(12):e0277512. doi: <https://dx.doi.org/10.1371/journal.pone.0277512>. **Exclusion reason:** Ineligible publication type

287. Schubert EA, Alffenaar JC, Johnstone MT, et al. Medicinal cannabis for patients with chronic non-cancer pain: analysis of safety and concomitant medications. *The International journal of pharmacy practice*. 2022doi: <https://dx.doi.org/10.1093/ijpp/riac073>. **Exclusion reason:** Ineligible study design
288. Schulze-Schiappacasse C, Duran J, Bravo-Jeria R, et al. Are Cannabis, Cannabis-Derived Products, and Synthetic Cannabinoids a Therapeutic Tool for Rheumatoid Arthritis? A Friendly Summary of the Body of Evidence. *J Clin Rheumatol*. 2021doi: 10.1097/RHU.0000000000001745. PMID: 33859125. **Exclusion reason:** Background
289. Scuteri D, Guida F, Boccella S, et al. Nabiximols clinical translation to the treatment of pain and agitation in severe dementia (NACTOPAISD): clinical trial protocol. *Biomedicine & pharmacotherapy*. 2022;153doi: 10.1016/j.biopha.2022.113488. PMID: 36076584. **Exclusion reason:** Ineligible study design
290. Scuteri D, Rombolà L, Hamamura K, et al. Is there a rational basis for Cannabinoids research and development in ocular pain therapy? A systematic review of preclinical evidence. *Biomed Pharmacother*. 2022;146doi: 10.1016/j.biopha.2021.112505. PMID: 34891121. **Exclusion reason:** Ineligible population
291. Seehusen DA, Kehoe K. Cannabis for treatment of chronic pain. *American family physician*. 2022;106(2):202-4. PMID: 35977124. **Exclusion reason:** Ineligible publication type
292. Senderovich H, Wagman H, Zhang D, et al. The effectiveness of Cannabis and Cannabis derivatives in treating lower back pain in the aged population: a systematic review. *Gerontology*. 2021;1-13. doi: 10.1159/000518269. PMID: 34515130. **Exclusion reason:** Systematic review used as source document
293. Senderovich H, Wagman H, Zhang D, et al. The effectiveness of Cannabis and Cannabis derivatives in treating lower back pain in the aged population: a systematic review. *Gerontology*. 2022;68(6):612-24. doi: 10.1159/000518269. PMID: 34515130. **Exclusion reason:** Systematic review used as source document
294. Serrano A, Galvez R, Paremés E, et al. Off-label pharmacological treatment for neuropathic pain: A Delphi study by the Spanish Pain Society Neuropathic Pain Task Force. *Pain practice : the official journal of World Institute of Pain*. 2022doi: <https://dx.doi.org/10.1111/papr.13176>. **Exclusion reason:** Background
295. Shah J, Fermo O. Review of systemic and syndromic complications of cannabis use: A review. *Medicine (United States)*. 2022;101(49):E32111. doi: 10.1097/MD.00000000000032111. **Exclusion reason:** Ineligible publication type
296. Sharon H, Weizman L, Espaniol J, et al. Cannabis induces changes in functional brain connectivity that correlate with increased vagal tone and clinical analgesia in chronic neuropathic pain. *Clinical autonomic research*. 2019;29(5):529. doi: <https://doi.org/10.1007/s10286-019-00631-x>. **Exclusion reason:** Ineligible publication type
297. Shebaby W, Saliba J, Faour WH, et al. In vivo and in vitro anti-inflammatory activity evaluation of Lebanese Cannabis Sativa L. ssp. Indica (Lam.). *J Ethnopharmacol*. 2020;113743. doi: 10.1016/j.jep.2020.113743. PMID: 33359187. **Exclusion reason:** Ineligible study design
298. Shehata I, Hashim A, Elsaedy A, et al. Cannabinoids and Their Role in Chronic Pain Treatment: Current Concepts and a Comprehensive Review. *Health Psychology Research*. 2022;10(4)doi: 10.52965/001c.35848. **Exclusion reason:** Systematic review used as source document

299. Shoshan S, Solt I. Medical Cannabis for Gynecologic pain conditions: a systematic review. *Obstetrics and gynecology*. 2022;139(5):937-8. doi: 10.1097/AOG.0000000000004777. PMID: 35104069. **Exclusion reason:** Ineligible publication type
300. Sihota A, Smith BK, Ahmed SA, et al. Consensus-based recommendations for titrating Cannabinoids and tapering opioids for chronic pain control. *Int J Clin Pract*. 2021;75(8):e13871. doi: 10.1111/ijcp.13871. PMID: 33249713. **Exclusion reason:** Ineligible publication type
301. Smaga S, Gharib A. In adults with chronic low back pain, does the use of inhaled Cannabis reduce overall opioid use? *Evid Based Pract*. 2017;20(1):E10-E1. doi: 10.1097/01.EBP.0000541620.26384.bf. **Exclusion reason:** Ineligible publication type
302. Socias ME, Choi J, Lake S, et al. Cannabis use is associated with reduced risk of exposure to Fentanyl among people on opioid agonist therapy during a community-wide overdose crisis. *Drug Alcohol Depend*. 2020;219:108420. doi: 10.1016/j.drugalcdep.2020.108420. PMID: 33342591. **Exclusion reason:** Ineligible population
303. Sotoodeh R, Waldman LE, Vigano A, et al. Predictors of pain reduction among Fibromyalgia patients using Medical Cannabis: a long-term prospective cohort study. *Arthritis care & research*. 2022doi: 10.1002/acr.24985. PMID: 35876631. **Exclusion reason:** Ineligible comparator
304. Stephens KL, Heineman JT, Forster GL, et al. Cannabinoids and Pain for the Plastic Surgeon: What Is the Evidence? *Annals of plastic surgery*. 2022doi: 10.1097/SAP.0000000000003128. PMID: 35502947. **Exclusion reason:** Systematic review used as source document
305. Stockings E, Campbell G, Hall WD, et al. Cannabis and Cannabinoids for the treatment of people with chronic noncancer pain conditions: a systematic review and meta-analysis of controlled and observational studies. *Pain*. 2018 Oct;159(10):1932-54. doi: 10.1097/j.pain.0000000000001293. PMID: 29847469. **Exclusion reason:** Systematic review used as source document
306. Strand NH, Maloney J, Kraus M, et al. Cannabis for the Treatment of Fibromyalgia: A Systematic Review. *Biomedicines*. 2023;11(6)doi: <https://dx.doi.org/10.3390/biomedicines11061621>. **Exclusion reason:** Systematic review used as source document
307. Sturgeon JA, Khan J, Hah JM, et al. Clinical profiles of concurrent cannabis use in chronic pain: a CHOIR study. *Pain Med*. 2020 Nov 01;21(11):3172-9. doi: 10.1093/pm/pnaa060. PMID: 32232476. **Exclusion reason:** Ineligible population
308. Sun N, Cunha N, Amar S, et al. Synthetic cannabinoid for the treatment of severe chronic noncancer pain in children and adolescents. *Can J Pain*. 2022;6(1):225-31. doi: <https://dx.doi.org/10.1080/24740527.2022.2132138>. **Exclusion reason:** Ineligible study design
309. Svendsen KB, Jensen TS, Bach FW. Does the Cannabinoid Dronabinol reduce central pain in Multiple Sclerosis? Randomised double blind placebo controlled crossover trial. *BMJ*. 2004 Jul 31;329(7460):253. doi: 10.1136/bmj.38149.566979.AE. PMID: 15258006. **Exclusion reason:** Inadequate duration
310. Swogger MT, Smith KE, Garcia-Romeu A, et al. Understanding Kratom use: a guide for healthcare providers. *Front Pharmacol*. 2022;13doi: 10.3389/fphar.2022.801855. PMID: 35308216. **Exclusion reason:** Systematic review used as source document
311. Swogger MT, Walsh Z. Kratom use and mental health: a systematic review. *Drug Alcohol Depend*. 2018 Feb 1;183:134-40. doi: 10.1016/j.drugalcdep.2017.10.012. PMID: 29248691. **Exclusion reason:** Systematic review used as source document

312. Syed Y, McKeage K, Scott L. Delta-9-Tetrahydrocannabinol/Cannabidiol (Sativex(R)): a review of its use in patients with moderate to severe spasticity due to Multiple Sclerosis. *Drugs*. 2014;74(5):563-78. doi: 10.1007/s40265-014-0197-5. PMID: 24671907. **Exclusion reason:** Systematic review used as source document
313. Sznitman S, Mabouk C, Said Z, et al. Opioid and healthcare service use in Medical Cannabis patients with chronic pain: a prospective study. *BMJ Support Palliat Care*. 2021;14:14. doi: 10.1136/bmjspcare-2020-002661. PMID: 34521640. **Exclusion reason:** Ineligible comparator
314. Sznitman SR, Vulfsons S, Meiri D, et al. Medical Cannabis and cognitive performance in middle to old adults treated for chronic pain. *Drug Alcohol Rev*. 2020 Sep 22;22:22. doi: 10.1111/dar.13171. PMID: 32964502. **Exclusion reason:** Ineligible study design
315. Sznitman SR, Vulfsons S, Meiri D, et al. Medical Cannabis and cognitive performance in middle to old adults treated for chronic pain. *Drug Alcohol Rev*. 2021;40(2):272-80. doi: 10.1111/dar.13171. PMID: 32964502. **Exclusion reason:** Ineligible study design
316. Takakuwa KM, Sulak D. A Survey on the effect that Medical Cannabis has on prescription opioid medication usage for the treatment of chronic pain at three medical Cannabis practice sites. *Cureus*. 2020;12(12):e11848. doi: 10.7759/cureus.11848. PMID: 33409086. **Exclusion reason:** Ineligible study design
317. Tetr. The study of effectiveness of Zingiber Cassumunar Roxb.(Plai) mixed Cannabis Sativa l. leaf oil for chronic pain in elderly randomized controlled trial. <https://trialssearch.who.int/Trial2.aspx?TrialID=TCTR20211116004>. 2021. **Exclusion reason:** Ineligible publication type
318. Terrie YC. Medical Cannabis for chronic pain. *US Pharm*. 2020;45(3):24-8. **Exclusion reason:** Ineligible publication type
319. Tervo-Clemmens B, Schmitt W, Wheeler G, et al. Cannabis use and sleep quality in daily life: a daily diary study of adults starting Cannabis for health concerns. medRxiv. 2022 2022-01-01 00:00:00doi: 10.1101/2022.01.19.22269565. **Exclusion reason:** Ineligible population
320. Thomas J. Inhaled Cannabis relieves neuropathic pain. *Aust J Pharm*. 2011;92(1091):88. doi: 10.3316/informit.861962231552691. **Exclusion reason:** Ineligible publication type
321. Thomas PA, Carter GT, Bombardier CH. A scoping review on the effect of Cannabis on pain intensity in people with spinal cord injury. *J Spinal Cord Med*. 2021:1-12. doi: 10.1080/10790268.2020.1865709. PMID: 33465022. **Exclusion reason:** Systematic review used as source document
322. Tsai SHL, Lin C-R, Shao S-C, et al. Cannabinoid Use for Pain Reduction in Spinal Cord Injuries: A Meta-Analysis of Randomized Controlled Trials. *Front Pharmacol*. 2022;13:866235. doi: 10.3389/fphar.2022.866235. PMID: 35571093. **Exclusion reason:** Systematic review used as source document
323. Turcotte D, Chateau D, Doupe M, et al. A randomised, double-blinded, placebo-controlled study evaluating the efficacy and safety of nabilone as an adjunctive to gabapentin in managing multiple sclerosis-induced neuropathic pain. *Mult Scler*. 2011;17(10):S475. doi: <https://doi.org/10.1177/1352458511422301>. **Exclusion reason:** Ineligible study design
324. Turcotte D, Doupe M, Torabi M, et al. A randomized, double-blinded, placebo-controlled study evaluating efficacy and tolerability of nabilone as an adjunctive to gabapentin in the management of multiple sclerosis-induced neuropathic pain. *Mult Scler*. 2013;19(11):112. doi: <https://doi.org/10.1177/1352458513502429>. **Exclusion reason:** Ineligible publication type

325. Turcotte DA, Gomori AJ, Esfahani FE, et al. Randomized, double-blinded, placebo-controlled study evaluating the efficacy and safety of nabilone adjunctive to gabapentin in managing multiple sclerosis-induced neuropathic pain. *European journal of pain supplements*. 2011;5(1):240. doi: [https://doi.org/10.1016/S1754-3207\(11\)70830-6](https://doi.org/10.1016/S1754-3207(11)70830-6). **Exclusion reason:** Ineligible publication type
326. Turcotte DA, Namaka MP, Gomori AJ, et al. A randomized, double-blinded, placebo-controlled study evaluating the efficacy and safety of Nabilone as an adjunctive to Gabapentin in managing Multiple Sclerosis-induced neuropathic pain: an interim analysis. *Pain Res Manag*. 2011;15(2):99. **Exclusion reason:** Ineligible publication type
327. Ueberall MA. A review of scientific evidence for THC: CBD oromucosal spray (Nabiximols) in the management of chronic pain. *J Pain Res*. 2020 Feb 14;13:399-410. doi: 10.2147/JPR.S240011. PMID: 32104061. **Exclusion reason:** Systematic review used as source document
328. Ueberall MA, Essner U, Mueller-Schwefe GH. Effectiveness and tolerability of THC:CBD oromucosal spray as add-on measure in patients with severe chronic pain: analysis of 12-week open-label real-world data provided by the German Pain e-Registry. *J Pain Res*. 2019 May 20;12:1577-604. doi: 10.2147/JPR.S192174. PMID: 31190969. **Exclusion reason:** Ineligible comparator
329. Ueberall MA, Horlemann J, Schuermann N, et al. Effectiveness and tolerability of Dronabinol use in patients with chronic pain - a retrospective analysis of 12-week open-label real-world data provided by the German Pain e-Registry. *Pain Med*. 2022 Feb 01;01:01. doi: 10.1093/pm/pnac010. PMID: 35104881. **Exclusion reason:** Ineligible study design
330. Urits I, Adamian L, Fiocchi J, et al. Advances in the understanding and management of chronic pain in Multiple Sclerosis: a comprehensive review. *Curr Pain Headache Rep*. 2019 Jul 25;23(8):59. doi: 10.1007/s11916-019-0800-2. PMID: 31342191. **Exclusion reason:** Systematic review used as source document
331. Urits I, Charipova K, Gress K, et al. Adverse Effects of Recreational and Medical Cannabis. *Psychopharmacol Bull*. 2021;51(1):94-109. PMID: 33897066. **Exclusion reason:** Ineligible study design
332. van Amerongen G, Kanhai K, Baakman AC, et al. Effects on spasticity and neuropathic pain of an oral formulation of DELTA9-tetrahydrocannabinol in patients with progressive Multiple Sclerosis. *Clin Ther*. 2018 Sep;40(9):1467-82. doi: 10.1016/j.clinthera.2017.01.016. PMID: 28189366. **Exclusion reason:** Ineligible population
333. van Dam CJ, van Velzen M, Kramers C, et al. Cannabis-opioid interaction in the treatment of fibromyalgia pain: an open-label, proof of concept study with randomization between treatment groups: cannabis, oxycodone or cannabis/oxycodone combination-the SPIRAL study. *Trials*. 2023;24(1):64. doi: <https://dx.doi.org/10.1186/s13063-023-07078-6>. **Exclusion reason:** Ineligible publication type
334. Vela J, Kjaer Petersen K, Dreyer L, et al. The effect of Cannabidiol on quantitative sensory testing parameters in patients with Hand Osteoarthritis and Psoriatic Arthritis: a randomized double-blind placebo-controlled trial. *Ann Rheum Dis*. 2022;81:821-2. doi: 10.1136/annrheumdis-2022-eular.1101. **Exclusion reason:** Ineligible publication type
335. Vermersch P, Trojano M. Tetrahydrocannabinol: cannabidiol oromucosal spray for Multiple Sclerosis-related resistant spasticity in daily practice. *Eur Neurol*. 2016;76(5-6):216-26. doi: 10.1159/000449413. PMID: 27732980. **Exclusion reason:** Ineligible comparator
336. Vickery AW, Roth S, Ernenwein T, et al. A large Australian longitudinal cohort registry demonstrates sustained safety and efficacy of oral medicinal cannabis for at least two years. *PLoS ONE*. 2022;17(11):e0272241. doi: <https://dx.doi.org/10.1371/journal.pone.0272241>. **Exclusion reason:** Ineligible study design


337. Vicknasingam B, Chooi WT, Rahim AA, et al. Kratom and pain tolerance: a randomized, placebo-controlled, double-blind study. *Yale J Biol Med.* 2020;93(2):229-38. PMID: 32607084. **Exclusion reason:** Ineligible population
338. Vila Silvan C, Vaney C, Dykukha I. Systematic reviews of randomized controlled trials of Cannabinoid products in chronic pain conditions and for symptoms associated with Multiple Sclerosis: what do they tell us? *Expert Rev Clin Pharmacol.* 2022doi: 10.1080/17512433.2022.2088501. PMID: 35679523. **Exclusion reason:** Systematic review used as source document
339. Villanueva MRB, Joshaghani N, Villa N, et al. Efficacy, safety, and regulation of Cannabidiol on chronic pain: a systematic review. *Cureus.* 2022;14(7):e26913. doi: 10.7759/cureus.26913. PMID: 35860716. **Exclusion reason:** Systematic review used as source document
340. Wade DT, Makela P, Robson P, et al. Do Cannabis-based medicinal extracts have general or specific effects on symptoms in Multiple Sclerosis? A double-blind, randomized, placebo-controlled study on 160 patients. *Mult Scler.* 2004;10(4):434-41. doi: 10.1191/1352458504ms1082oa. PMID: 15327042. **Exclusion reason:** Ineligible population
341. Walitt B, Klose P, Fitzcharles MA, et al. Cannabinoids for Fibromyalgia. *Cochrane Database Syst Rev.* 2016 Jul 18;7:CD011694. doi: 10.1002/14651858.CD011694.pub2. PMID: 27428009. **Exclusion reason:** Ineligible publication type
342. Wallace MS, Marcotte TD, Umlauf A, et al. Efficacy of inhaled Cannabis on painful Diabetic Neuropathy. *J Pain.* 2015 Jul;16(7):616-27. doi: 10.1016/j.jpain.2015.03.008. PMID: 25843054. **Exclusion reason:** Inadequate duration
343. Wang C, Erridge S, Holvey C, et al. Assessment of clinical outcomes in patients with fibromyalgia: Analysis from the UK Medical Cannabis Registry. *Brain and behavior.* 2023:e3072. doi: <https://dx.doi.org/10.1002/brb3.3072>. **Exclusion reason:** Ineligible study design
344. Wang L, Hong PJ, May C, et al. Medical cannabis or cannabinoids for chronic non-cancer and cancer related pain: a systematic review and meta-analysis of randomised clinical trials. *BMJ.* 2021;374:n1034. doi: 10.1136/bmj.n1034. PMID: 34497047. **Exclusion reason:** Systematic review used as source document
345. Wang T, Collet JP, Shapiro S, et al. Adverse effects of medical cannabinoids: a systematic review. *CMAJ.* 2008;178(13):1669-78. doi: 10.1503/cmaj.071178. PMID: 18559804. **Exclusion reason:** Ineligible publication type
346. Wang Y, Jean Jacques J, Li Z, et al. Health outcomes among adults initiating Medical Cannabis for chronic pain: a 3-month prospective study incorporating ecological momentary assessment (EMA). *Cannabis.* 2021;4(2):69-83. doi: 10.26828/cannabis/2021.02.006. PMID: 34671723. **Exclusion reason:** Ineligible comparator
347. Ware MA, Fitzcharles MA, Joseph L, et al. The effects of Nabilone on sleep in Fibromyalgia: results of a randomized controlled trial. *Anesth Analg.* 2010 Feb 01;110(2):604-10. doi: 10.1213/ANE.0b013e3181c76f70. PMID: 20007734. **Exclusion reason:** Inadequate duration
348. Ware MA, Wang T, Shapiro S, et al. Smoked Cannabis for chronic neuropathic pain: a randomized controlled trial. *Cmaj.* 2010 Oct 05;182(14):E694-701. doi: 10.1503/cmaj.091414. PMID: 20805210. **Exclusion reason:** Inadequate duration
349. White CM. Pharmacologic and clinical assessment of Kratom. *Am J Health-Syst Pharm.* 2018 Mar 1;75(5):261-7. doi: 10.2146/ajhp161035. PMID: 29255059. **Exclusion reason:** Background
350. White CM. Pharmacologic and clinical assessment of Kratom: an update. *Am J Health-Syst Pharm.* 2019 Nov 13;76(23):1915-25. doi: 10.1093/ajhp/zxz221. PMID: 31626272. **Exclusion reason:** Background

351. Williams AR, Hill KP. Care of the patient using Cannabis. *Ann Intern Med.* 2020;173(9):ITC65-ITC80. doi: 10.7326/AITC202011030. PMID: 33137270. **Exclusion reason:** Ineligible publication type
352. Williams AR, Mauro CM, Feng T, et al. Adult Medical Cannabinoid Use and Changes in Prescription Controlled Substance Use. *Cannabis Cannabinoid Res.* 2022doi: 10.1089/can.2021.0212. PMID: 35486854. **Exclusion reason:** Ineligible comparator
353. Wilsey B, Marcotte T, Deutsch R, et al. Low-dose vaporized Cannabis significantly improves neuropathic pain. *J Pain.* 2013 Feb;14(2):136-48. doi: 10.1016/j.jpain.2012.10.009. PMID: 23237736. **Exclusion reason:** Inadequate duration
354. Wilsey B, Marcotte T, Tsodikov A, et al. A randomized, placebo-controlled, crossover trial of Cannabis cigarettes in neuropathic pain. *J Pain.* 2008 Jun;9(6):506-21. doi: 10.1016/j.jpain.2007.12.010. PMID: 18403272. **Exclusion reason:** Inadequate duration
355. Wong SSC, Chan WS, Cheung CW. Analgesic effects of Cannabinoids for chronic non-cancer pain: a systematic review and meta-analysis with meta-regression. *J Neuroimmune Pharmacol.* 2020 Dec;15(4):801-29. doi: 10.1007/s11481-020-09905-y. PMID: 32172501. **Exclusion reason:** Systematic review used as source document
356. Xantus G, Gyarmathy V. Cannabidiol in acute low back pain-a protocol and rationale of a clinical effectiveness trial. *Med Cannabis Cannabinoids.* 2020;3(2):134. doi: 10.1159/000511664. PMID: 634190900. **Exclusion reason:** Ineligible publication type
357. Yacyshyn B, Hanauer S, Klassen P, et al. Safety, Pharmacokinetics, and efficacy of Olorinab, a peripherally acting, highly selective, full agonist of the Cannabinoid receptor 2, in a Phase 2a study of patients with chronic abdominal pain associated with Crohn's Disease. *Crohns Colitis* 360. 2021;3(1)doi: 10.1093/crocol/otaa089. **Exclusion reason:** Ineligible intervention
358. Yanes JA, McKinnell ZE, Reid MA, et al. Effects of Cannabinoid administration for pain: a meta-analysis and meta-regression. *Exp Clin Psychopharmacol.* 2019 Aug;27(4):370-82. doi: 10.1037/pha0000281. PMID: 31120281. **Exclusion reason:** Ineligible population
359. Yassin M, Oron A, Robinson D. Effect of adding Medical Cannabis to Analgesic treatment in patients with low back pain related to Fibromyalgia: an observational cross-over single centre study. *Clin Exp Rheumatol.* 2019 Jan-Feb;37(Suppl 116):S13-S20. PMID: 30418116. **Exclusion reason:** Ineligible study design
360. Yimam M, O'Neal A, Horm T, et al. Antinociceptive and Anti-Inflammatory Properties of Cannabidiol Alone and in Combination with Standardized Bioflavonoid Composition. *J Med Food.* 2021doi: 10.1089/jmf.2020.0178. PMID: 33570460. **Exclusion reason:** Ineligible population
361. Yu JS, Premkumar A, Liu S, et al. Rates of self-directed perioperative Cannabidiol use in patients undergoing total hip or knee arthroplasty. *Pain manag.* 2021 Jun 09;09:09. doi: 10.2217/pmt-2021-0018. PMID: 34102871. **Exclusion reason:** Ineligible study design
362. Zajicek J, Fox P, Sanders H, et al. Cannabinoids for treatment of spasticity and other symptoms related to multiple sclerosis (CAMS study): multicentre randomised placebo-controlled trial. *Lancet.* 2003 Nov 8;362(9395):1517-26. doi: 10.1016/s0140-6736(03)14738-1. PMID: 14615106. **Exclusion reason:** Ineligible population
363. Zalai D, Chung SA, Hussain N, et al. Does cannabinoid really improve sleep? Testing the sleep effects of nabilone in chronic pain patients: a placebo-controlled, randomized, pilot study. *Psychotherapy and psychosomatics.* 2015;84:81. doi: <https://doi.org/10.1159/000438780>. **Exclusion reason:** Ineligible publication type
364. Zavori L, Xantus G, Matheson C, et al. Cannabidiol in low back pain: scientific rationale for clinical trials in low back pain. *Expert Rev Clin Pharmacol.* 2021doi: 10.1080/17512433.2021.1917379. PMID: 33861675. **Exclusion reason:** Background

365. Zeraatkar D, Cooper MA, Agarwal A, et al. Long-term and serious harms of Medical Cannabis and Cannabinoids for chronic pain: a systematic review of non-randomized studies. medRxiv. 2021 2021-01-01 00:00:00doi: 10.1101/2021.05.27.21257921. PMID: 35926992. **Exclusion reason:** Systematic review used as source document
366. Zloczower O, Brill S, Zeitak Y, et al. Risk and benefit of Cannabis prescription for chronic non-cancer pain. J Addict Dis. 2021 Aug 02:1-11. doi: 10.1080/10550887.2021.1956673. PMID: 34338621. **Exclusion reason:** Ineligible study design

Appendix J. Funnel Plot of High THC Ratio Studies Included in Meta-Analysis for Pain Severity

Figure J-1. Funnel plot of nine trials of pain severity for high THC ratio products versus placebo



Abbreviations: Groupdiff = group difference; SE = standard error; THC = tetrahydrocannabinol.