



Comparative Effectiveness Review
Number 254

Improving Rural Health Through Telehealth-Guided Provider-to-Provider Communication



Comparative Effectiveness Review

Number 254

Improving Rural Health Through Telehealth-Guided Provider-to-Provider Communication

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
5600 Fishers Lane
Rockville, MD 20857
www.ahrq.gov

Contract No. 75Q80120D00006

Prepared by:

Pacific Northwest Evidence-based Practice Center
Portland, OR

Investigators:

Annette Totten, Ph.D.
Dana M. Womack, Ph.D., R.N.
Marian S. McDonagh, Pharm.D.
Cynthia Davis-O'Reilly, B.S.
Jessica C. Griffin, M.S.
Ian Blazina, M.P.H.
Sara Grusing, M.P.H.
Nancy Elder, M.D., M.S.P.H.

**AHRQ Publication No. 22(23)-EHC023
December 2022**

This report is based on research conducted by the Pacific Northwest Evidence-based Practice Center (EPC) under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. 75Q80120D00006). The National Institutes of Health (NIH) Office of Disease Prevention funded the report. The findings and conclusions in this document are those of the authors, who are responsible for its contents; the findings and conclusions do not necessarily represent the views of AHRQ or NIH. Therefore, no statement in this report should be construed as an official position of AHRQ, NIH, or the U.S. Department of Health and Human Services.

None of the investigators have any affiliations or financial involvement that conflicts with the material presented in this report.

The information in this report is intended to help healthcare decision makers—patients and clinicians, health system leaders, and policymakers, among others—make well-informed decisions and thereby improve the quality of healthcare services. This report is not intended to be a substitute for the application of clinical judgment. Anyone who makes decisions concerning the provision of clinical care should consider this report in the same way as any medical reference and in conjunction with all other pertinent information, i.e., in the context of available resources and circumstances presented by individual patients.

This report is made available to the public under the terms of a licensing agreement between the author and the Agency for Healthcare Research and Quality. This report may be used and reprinted without permission except those copyrighted materials that are clearly noted in the report. Further reproduction of those copyrighted materials is prohibited without the express permission of copyright holders.

AHRQ or U.S. Department of Health and Human Services endorsement of any derivative products that may be developed from this report, such as clinical practice guidelines, other quality enhancement tools, or reimbursement or coverage policies, may not be stated or implied.

AHRQ appreciates appropriate acknowledgment and citation of its work. Suggested language for acknowledgment: This work was based on an evidence report, Improving Rural Health Through Telehealth-Guided Provider-to-Provider Communication, by the Evidence-based Practice Center Program at the Agency for Healthcare Research and Quality (AHRQ).

Suggested citation: Totten A, Womack DM, McDonagh MS, Davis-O'Reilly C, Griffin JC, Blazina I, Grusing S, Elder N. Improving Rural Health Through Telehealth-Guided Provider-to-Provider Communication. Comparative Effectiveness Review No. 254. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. 75Q80120D00006.) AHRQ Publication No. 22(23)-EHC023. Rockville, MD: Agency for Healthcare Research and Quality; December 2022. DOI: <https://doi.org/10.23970/AHRQEPCCER254>. Posted final reports are located on the Effective Health Care Program [search page](#).

Preface

The Agency for Healthcare Research and Quality (AHRQ), through its Evidence-based Practice Centers (EPCs), sponsors the development of evidence reports and technology assessments to assist public- and private-sector organizations in their efforts to improve the quality of healthcare in the United States.

The National Institutes of Health (NIH) Office of Disease Prevention requested this report from the EPC Program at AHRQ. AHRQ assigned this report to the following EPC: Pacific Northwest Evidence-based Practice Center (Contract Number: 75Q80120D00006).

The report was presented at the NIH Office of Disease Prevention public meeting Improving Rural Health Through Telehealth-Guided Provider-to-Provider Communication on October 12–13, 2021.

The reports and assessments provide organizations with comprehensive, evidence-based information on common medical conditions and new healthcare technologies and strategies. They also identify research gaps in the selected scientific area, identify methodological and scientific weaknesses, suggest research needs, and move the field forward through an unbiased, evidence-based assessment of the available literature. The EPCs systematically review the relevant scientific literature on topics assigned to them by AHRQ and conduct additional analyses when appropriate prior to developing their reports and assessments.

To bring the broadest range of experts into the development of evidence reports and health technology assessments, AHRQ encourages the EPCs to form partnerships and enter into collaborations with other medical and research organizations. The EPCs work with these partner organizations to ensure that the evidence reports and technology assessments they produce will become building blocks for healthcare quality improvement projects throughout the Nation. The reports undergo peer review and public comment prior to their release as a final report.

AHRQ expects that the EPC evidence reports and technology assessments, when appropriate, will inform individual health plans, providers, and purchasers as well as the healthcare system as a whole by providing important information to help improve healthcare quality.

If you have comments on this evidence report, they may be sent by mail to the Task Order Officer named below at: Agency for Healthcare Research and Quality, 5600 Fishers Lane, Rockville, MD 20857, or by email to epc@ahrq.hhs.gov.

Robert Otto Valdez, Ph.D., M.H.S.A.
Director
Agency for Healthcare Research and Quality

Arlene S. Bierman, M.D., M.S.
Director
Center for Evidence and Practice
Improvement
Agency for Healthcare Research and Quality

Craig A. Umscheid, M.D., M.S.
Director
Evidence-based Practice Center Program
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Lionel L. Bañez, M.D.
Task Order Officer
Center for Evidence and Practice
Improvement
Agency for Healthcare Research and Quality

Acknowledgments

The authors gratefully acknowledge the following individuals for their contributions to this project: AHRQ Task Order Officers Lionel Bañez, M.D., and Meghan Wagner, Pharm.D.; NIH Office of Disease Prevention (ODP): Kate Winseck, M.S.W., Maria Babirye, M.P.H., Melissa Green Parker, Ph.D., Carrie Klabunde, Ph.D., and Keisha Shropshire, M.P.H; ODP Working Group and NIH partners; Associate Editor Jonathan Treadwell, Ph.D.; Technical Expert Panel members; Peer Reviewers; and Paul Swenson, M.D., Chandler Atchison, M.S., Leah Williams, B.S., and Tracy Dana, M.L.S., for their contributions to the preparation of this report.

Technical Expert Panel

In designing the study questions and methodology at the outset of this report, the EPC consulted several technical and content experts. Broad expertise and perspectives were sought. Divergent and conflicted opinions are common and perceived as healthy scientific discourse that results in a thoughtful, relevant systematic review. Therefore, in the end, study questions, design, methodological approaches, and/or conclusions do not necessarily represent the views of individual technical and content experts.

Technical Experts must disclose any financial conflicts of interest greater than \$5,000 and any other relevant business or professional conflicts of interest. Because of their unique clinical or content expertise, individuals with potential conflicts may be retained. The TOO and the EPC work to balance, manage, or mitigate any potential conflicts of interest identified.

The list of Technical Experts who provided input to this report follows:

Jodi G. Daniel, J.D., M.P.H.

Partner

Crowell & Moring LLP

Washington, DC

Martin Doerfler, M.D.*

Senior Vice President, Clinical Strategy and Development

Associate Chief Medical Officer

Northwell Health

Hyde Park, NY

Tim Gallagher, M.P.H.

Healthy Opportunities of the Piedmont

Piedmont Triad Regional Council

Winston-Salem, NC

Kevin Larsen, M.D.

Senior Vice President of Clinical Innovation and Translation

Optum Labs

Washington, DC

Thomas Tsang, M.D., M.P.H.
CEO, Co-Founder
Valera Health
Brooklyn, NY

Marcia Ward, Ph.D.
Director, Center for Health Policy and Research
College of Public Health
University of Iowa
Iowa City, IA

Jon White
Associate Chief of Staff for Research
U.S. Department of Veterans Affairs
Professor of Medicine
University of Utah School of Medicine
Salt Lake City, UT

*Provided input on Draft Report

Peer Reviewers

Prior to publication of the final evidence report, the EPC sought input from independent Peer Reviewers without financial conflicts of interest. However, the conclusions and synthesis of the scientific literature presented in this report do not necessarily represent the views of individual reviewers.

Peer Reviewers must disclose any financial conflicts of interest greater than \$5,000 and any other relevant business or professional conflicts of interest. Because of their unique clinical or content expertise, individuals with potential nonfinancial conflicts may be retained. The TOO and the EPC work to balance, manage, or mitigate any potential nonfinancial conflicts of interest identified.

The list of Peer Reviewers follows:

Shira H. Fischer, M.D., Ph.D.
Physician Policy Researcher
RAND Corporation
Boston, MA

Clare Liddy, M.D., M.Sc.
Professor and Chair, Department of Family Medicine
Research Chair in eConsult
University of Ottawa
Ottawa, ON, Canada

Christopher M. Shea, Ph.D., M.A., M.P.A.
Gillings School of Global Public Health
University of North Carolina-Chapel Hill
Chapel Hill, NC

John D. Whited, M.D., M.H.S.
Durham Veterans Affairs Health Care System
Associate Professor of Medicine
Duke University School of Medicine
Durham, NC

Improving Rural Health Through Telehealth-Guided Provider-to-Provider Communication

Structured Abstract

Objectives. To assess the use, effectiveness, and implementation of telehealth-supported provider-to-provider communication and collaboration for the provision of healthcare services to rural populations and to inform a scientific workshop convened by the National Institutes of Health Office of Disease Prevention on October 12–14, 2021.

Data sources. We conducted a comprehensive literature search of Ovid MEDLINE®, CINAHL®, Embase®, and Cochrane CENTRAL. We searched for articles published from January 1, 2015, to October 12, 2021, to identify data on use of rural provider-to-provider telehealth (Key Question 1) and the same databases for articles published January 1, 2010, to October 12, 2021, for studies of effectiveness and implementation (Key Questions 2 and 3) and to identify methodological weaknesses in the research (Key Question 4). Additional sources were identified through reference lists, stakeholder suggestions, and responses to a Federal Register notice.

Review methods. Our methods followed the Agency for Healthcare Research and Quality *Methods Guide* (available at <https://effectivehealthcare.ahrq.gov/topics/cer-methods-guide/overview>) and the PRISMA reporting guidelines. We used predefined criteria and dual review of abstracts and full-text articles to identify research results on (1) regional or national use, (2) effectiveness, (3) barriers and facilitators to implementation, and (4) methodological weakness in studies of provider-to-provider telehealth for rural populations. We assessed the risk of bias of the effectiveness studies using criteria specific to the different study designs and evaluated strength of evidence (SOE) for studies of similar telehealth interventions with similar outcomes. We categorized barriers and facilitators to implementation using the Consolidated Framework for Implementation Research (CFIR) and summarized methodological weaknesses of studies.

Results. We included 166 studies reported in 179 publications. Studies on the degree of uptake of provider-to-provider telehealth were limited to specific clinical uses (pharmacy, psychiatry, emergency care, and stroke management) in seven studies using national or regional surveys and claims data. They reported variability across States and regions, but increasing uptake over time.

Ninety-seven studies (20 trials and 77 observational studies) evaluated the effectiveness of provider-to-provider telehealth in rural settings, finding that there may be similar rates of transfers and lengths of stay with telehealth for inpatient consultations; similar mortality rates for remote intensive care unit care; similar clinical outcomes and transfer rates for neonates; improvements in medication adherence and treatment response in outpatient care for depression; improvements in some clinical monitoring measures for diabetes with endocrinology or pharmacy outpatient consultations; similar mortality or time to treatment when used to support emergency assessment and management of stroke, heart attack, or chest pain at rural hospitals; and similar rates of appropriate versus inappropriate transfers of critical care and trauma patients with specialist telehealth consultations for rural emergency departments (SOE: low). Studies of telehealth for education and mentoring of rural healthcare providers may result in intended

changes in provider behavior and increases in provider knowledge, confidence, and self-efficacy (SOE: low). Patient outcomes were not frequently reported for telehealth provider education, but two studies reported improvement (SOE: low). Evidence for telehealth interventions for other clinical uses and outcomes was insufficient.

We identified 67 program evaluations and qualitative studies that identified barriers and facilitators to rural provider-to-provider telehealth. Success was linked to well-functioning technology; sufficient resources, including time, staff, leadership, and equipment; and adequate payment or reimbursement. Some considerations may be unique to implementation of provider-to-provider telehealth in rural areas. These include the need for consultants to better understand the rural context; regional initiatives that pool resources among rural organizations that may not be able to support telehealth individually; and programs that can support care for infrequent as well as frequent clinical situations in rural practices.

An assessment of methodological weaknesses found that studies were limited by less rigorous study designs, small sample sizes, and lack of analyses that address risks for bias. A key weakness was that studies did not assess or attempt to adjust for the risk that temporal changes may impact the results in studies that compared outcomes before and after telehealth implementation.

Conclusions. While the evidence base is limited, what is available suggests that telehealth supporting provider-to-provider communications and collaboration may be beneficial. Telehealth studies report better patient outcomes in some clinical scenarios (e.g., outpatient care for depression or diabetes, education/mentoring) where telehealth interventions increase access to expertise and high-quality care. In other applications (e.g., inpatient care, emergency care), telehealth results in patient outcomes that are similar to usual care, which may be interpreted as a benefit when the purpose of telehealth is to make equivalent services available locally to rural residents. Most barriers to implementation are common to practice change efforts. Methodological weaknesses stem from weaker study designs, such as before-after studies, and small numbers of participants. The rapid increase in the use of telehealth in response to the Coronavirus disease 2019 (COVID-19) pandemic is likely to produce more data and offer opportunities for more rigorous studies.

Contents

Executive Summary	ES-1
Introduction	1
Background	1
Purpose and Scope of This Systematic Review	2
Methods	3
Review Approach	3
Key Questions	3
Analytic Framework	4
Study Selection	4
Data Extraction and Risk of Bias Assessment	4
Data Synthesis and Analysis	5
Grading the Strength of the Body of Evidence	5
Results	7
Description of Included Evidence	7
Key Question 1. What is the uptake of different types of provider-to-provider telehealth in rural areas?	7
Key Points	7
Summary of Findings	7
Key Question 2. What is the effectiveness of provider-to-provider telehealth for rural patients?	10
a. How does provider-to-provider telehealth affect outcomes for patients and populations?	10
b. How does provider-to-provider telehealth affect outcomes for healthcare providers?	10
c. How does provider-to-provider telehealth affect outcomes for private and public payers?	10
Inpatient	11
Outpatient Care and Services	15
Emergency Department/Emergency Medical Services	25
Education and Mentoring	30
Key Question 3. What strategies are effective and what are the barriers and facilitators to implementation and sustainability of provider-to-provider telehealth in rural areas?	36
Key Points	36
Summary of Findings	36
Key Question 4. What are the methodological weaknesses of the included studies of provider-to-provider telehealth for rural patients and what improvements in study design (e.g., focus on relevant comparisons and outcomes) might increase the impact of future research?	39
Key Points	39
Summary of Findings	39
Discussion	44
Findings in Relation to the Decisional Dilemmas	44
Key Question 1: Uptake of Provider-to-Provider Telehealth in Rural Areas	44
Key Question 2: Effectiveness of Provider-to-Provider Telehealth in Improving Health Outcomes for Rural Populations and Providers and Payers	45

Key Question 3: Barriers and Facilitators to implementation and Implementation Strategies of Provider-to-Provider Telehealth	49
Key Question 4: Methodological Weaknesses of Studies of Provider-to-Provider Telehealth for Rural Populations	50
Limitations of the Review Process	50
Limitations of the Evidence Base	51
Applicability	52
Implications for Clinical and Policy Decision Making.....	53
Future Research Needs	54
Summary and Conclusion	55
References.....	56
Abbreviations and Acronyms	73

Tables

Table 1. Key Question 1: summary of provider-to-provider telehealth use in rural areas.....	9
Table 2. Key Question 2: summary of inpatient findings	13
Table 3. Key Question 2: summary of outpatient findings	22
Table 4. Key Question 2: summary of emergency care findings.....	27
Table 5. Key Question 2: summary of remote education and mentoring findings	33
Table 6. Publication study designs by Key Question 2 topic.....	40
Table 7. Publication site types by Key Question 2 topic	40
Table 8. Methodological weaknesses reported in included studies	42
Table 9. Strength of evidence for inpatient effectiveness studies	46
Table 10. Strength of evidence for outpatient effectiveness studies.....	47
Table 11. Strength of evidence for emergency care effectiveness studies.....	48
Table 12. Strength of evidence for education and mentoring effectiveness studies	48

Figures

Figure 1. Analytic framework.....	4
Figure 2. CFIR constructs as facilitators and barriers.....	37

Appendixes

Appendix A. Methods
Appendix B. Results
Appendix C. Included Studies List
Appendix D. Evidence Tables
Appendix E. Risk of Bias Assessment
Appendix F. Details on Strength of Evidence
Appendix G. Excluded Studies List
Appendix H. Appendix References

Executive Summary

Main Points

- **Use of Provider-to-Provider Telehealth for Rural Populations:** Limited research from regional and national surveys and claims data suggest that telehealth for provider-to-provider communication is used to different extents across location for specific clinical care uses such as psychiatry, emergency, and stroke care. Use was increasing even before the COVID-19 pandemic and this seems likely to continue, though research evaluating new and increased use has yet to be published.
- **Effectiveness of Provider-to-Provider Telehealth for Rural Populations:**
 - Telehealth to support direct patient care may provide benefits for inpatient care, for neonates in rural hospitals, outpatient management of depression and diabetes, and emergency care of stroke/heart attack/chest pain and trauma.
 - Evidence on other uses, outcomes, or populations was insufficient to support conclusions. No studies reported harms or unexpected negative outcomes for provider-to-provider telehealth.
 - Use of telehealth for provider education and mentoring, including programs like Extension for Community Healthcare Outcomes (ECHO) that use video for instruction and collaboration, may improve patient outcomes, change provider behavior, and increase provider knowledge and confidence in treating specific conditions.
- **Barriers and Facilitators to Implementation of Provider-to-Provider Telehealth for Rural Populations:** Inadequate provider time, technology, and other resources, as well as limited understanding of the rural context and lack of long-term commitments to telehealth are barriers to broader implementation of provider-to-provider telehealth in rural settings. Telehealth implementation was facilitated when there were sufficient resources, access to knowledge, engagement of leadership and the program addressed patient needs.
- **Methodological Weakness of Studies:** Effectiveness and implementation studies frequently employed less rigorous designs, had small sample sizes, and often did not minimize bias through design or analyses.

Background and Purpose

Numerous studies have documented health disparities for people living in rural areas.¹ Rural populations experience higher mortality²⁻⁵ and morbidity from a wide range of conditions including substance/opioid abuse,^{6, 7} chronic illnesses,⁸⁻¹¹ and HIV/human papillomavirus and other infectious diseases.^{12, 13} The purpose of this review is to identify, summarize, and evaluate the research available on whether telehealth supporting provider-to-provider communication and collaboration can contribute to addressing these disparities and improving the health and well-being of rural communities. The review was commissioned by the National Institutes of Health (NIH) Office of Disease Prevention to inform a Pathways to Prevention workshop, *Improving Rural Health Through Telehealth-Guided Provider-to-Provider Communication*, that was held on October 12-14, 2021 (<https://prevention.nih.gov/research-priorities/research-needs-and-gaps/pathways-prevention/improving-rural-health-through-telehealth-guided-provider-provider-communication>).

Methods

The Key Questions guiding the systematic evidence review were developed by an NIH working group and revised through a topic refinement process. We employed methods consistent with those outlined in the Agency for Healthcare Research and Quality Evidence-based Practice Center Program methods guidance (<https://effectivehealthcare.ahrq.gov/topics/ebp-methods-guide/overview>). Our search included articles published from January 1, 2010, through October 12, 2021. Detailed methods, including the search strategies, are included in the full report and appendixes.

Results

We identified 166 studies reported in 179 publications that addressed the use, effectiveness, implementation of telehealth designed to support provider-to-provider interactions. The extent to which these studies are able to answer each of the questions posed is summarized below.

Key Question 1. What is the uptake of different types of provider-to-provider telehealth in rural areas?

- We did not identify any surveys or national datasets that provided overall counts or estimates of provider-to-provider telehealth usage in the United States.
- Seven published reports provide data from surveys and claims data about rural use of telehealth for specific clinical uses. Examples are: reported rates of use of telepsychiatry (29.2% by U.S. mental health providers), telehealth in emergency care (54% of emergency departments in the United States), and telestroke (8.6 per 1,000 stroke cases for rural residents). Reported use of telehealth in the United States is variable across states and regions, but has been increasing over time.

Key Question 2. What is the effectiveness of provider-to-provider telehealth for rural patients?

- a. How does provider-to-provider telehealth affect outcomes for patients and populations?
- b. How does provider-to-provider telehealth affect outcomes for healthcare providers?
- c. How does provider-to-provider telehealth affect outcomes for private and public payers?

We assessed the research on effectiveness first by healthcare setting (i.e., inpatient, outpatient emergency care, education and mentoring), then by clinical topic within settings as telehealth interventions and outcomes differed across settings based on data from 97 studies.

Inpatient Care

- Telehealth consultations in rural hospitals may result in no difference in length of hospital stay (6 studies) or transfers (3 studies; low strength of evidence [SOE]) compared to usual care, including in-person or phone consultations.

- Telehealth supported care for neonates at rural hospitals may result in no difference in clinical outcomes when compared to transfer and care at a hospital with a Level 4 neonatal intensive care unit (3 studies; low SOE). When telehealth is available, neonate transfers may be more appropriate (2 studies; low SOE).
- Evidence suggests remote intensive care units (ICUs) in rural areas result in no difference in mortality rates compared to transferring patients to more distant locations for ICU care (2 studies; low SOE).

Outpatient Care

- Outpatient telehealth consultations with specialists may result in improvements in clinical outcomes compared to care without specialist involvement:
 - For patients with diabetes:
 - Some improvement in medication adherence and treatment response for patients with depression (3 studies; low SOE).
 - Improvements in A1c and self-management but no effect on blood pressure or cholesterol levels in patients with diabetes (4 studies; low SOE).
 - Improvements in A1c, fasting glucose, and blood pressure in patients with both hypertension and diabetes with pharmacy teleconsultations (2 studies; low SOE).
 - For patients with depression:
 - Higher utilization of telehealth and corresponding costs for outpatient consultations for depression are associated with increased access and cost-effectiveness analyses reported overall benefit (2 studies; low SOE).

Emergency Care

- Telehealth consultations supporting emergency assessment and care of stroke, heart attack, or chest pain at rural hospitals:
 - May result in similar rates of mortality when patients are treated locally as opposed to transferred (5 studies; low SOE).
 - May result in similar time to treatment when patients are treated locally as opposed to transferred (8 studies; low SOE).
- Telehealth consultations by specialists for critical care and trauma patients in rural emergency departments may result in no difference in appropriate or inappropriate transfers (5 studies; low SOE).

Education and Mentoring

- Clinical outcomes: ECHO programs (a specific model that uses video for instruction and case reviews) are associated with better or equivalent patient outcomes (2 studies; low SOE):
 - Reduction in A1c in patients of trainees after ECHO compared to before participation (1 study).
 - Hepatitis C viral response and serious adverse events rates at “spoke” site with ECHO participation were similar to those at an academic medical center (1 study).
- ECHO and non-ECHO video training programs:

- May result in desired changes in provider behavior (e.g., increased appropriate prescribing practices, screening, and patient counseling) (8 studies; low SOE).
- May be associated with increased provider confidence, efficacy, and scores on knowledge tests (13 studies; low SOE).

Key Question 3. What strategies are effective and what are the barriers and facilitators to implementation and sustainability of provider-to-provider telehealth in rural areas??

Sixty-seven program evaluation and qualitative studies using a wide range of methodologies provide information on implementation of provider-to-provider telehealth.

- Barriers and facilitators are similar across provider-to-provider telehealth programs implemented in different settings and for different purposes. The majority are related to available resources, and access to knowledge and information.
- Barriers to rural provider-to-provider telehealth may be addressed by the implementation of specific evidence-based strategies. The following were identified and suggested by the authors of included studies:
 - Consulting providers need to understand the characteristics of rural areas and populations and what resources are available. *Strategy*: rural rotations or periodic in person collaboration.
 - Successful implementation and sustainment require a long-term commitment and resources on a scale that may not be feasible for individual rural organizations. *Strategy*: statewide or regional initiatives with government or philanthropic support.
 - Provider-to-provider telehealth systems may be used for frequent events or serve as a resource for rare events in rural healthcare and the technology and support need to be tailored to frequency of use. *Strategy*: customize system to accommodate frequency of use, such as schedule periodic testing of systems used for rare events.

Key Question 4. What are the methodological weaknesses of the included studies of provider-to-provider telehealth for rural patients and what improvements in study design (e.g., focus on relevant comparisons and outcomes) might increase the impact of future research?

When reviewing studies for Key Questions 2 and 3 we abstracted the limitations cited by the authors and combined these with our risk of bias and applicability assessments in order to identify and categorize the methodological weaknesses.

- Studies of provider-to-provider telehealth for rural areas could be improved by addressing methodological weakness.
- Key weakness: it is often difficult to attribute impact to telehealth because--
 - Weaker study designs are common: randomized controlled trials and cohort studies were identified and accounted for 3/5 of the included studies, but more than 40 percent of the studies were repeated measures (pre/post or before/after) with no other comparison group.

- Lack of control for confounders related to patients, providers, facilities, and differences in telehealth implementation across study sites
- The most frequently identified weakness after overall study design is small sample sizes that result in lack of power to detect differences.
- Studies are also hampered by data limitations related to use of retrospective data, and data produced for care delivery and billing purposes, that can be incomplete or coded differently across organizations.

Strengths and Limitations

The research on provider-to-provider telehealth for communication and collaboration in the delivery of rural healthcare includes studies that directly address the questions asked by this review, but they are spread across settings, many different clinical uses, and evaluated different telehealth outcomes. For these reasons, the existing evidence base is unable to support strong conclusions. Overall, research on telehealth in general is often not based on a clear model of how telehealth is expected to affect outcomes; an outcome-oriented model for telehealth could inform better research. While telehealth should increase patient and provider satisfaction and other outcomes, there is no agreement on how to prioritize across clinical outcomes, resource use, costs, access to care and potential harms. It is also often unclear if the goal of telehealth is to provide care that is as good as care provided without telehealth or if the investment in telehealth requires that outcomes be better.

Implications and Conclusions

The limited available evidence suggests that telehealth supporting provider-to-provider communications and collaboration may produce similar or better results for patients, providers, and payers compared with care without telehealth. Barriers to implementation are known and common to practice change efforts. Methodological weaknesses are due to the use of less rigorous study designs that do not sufficiently address differences in the groups compared and include small numbers of participants. The rapid increase in the use of telehealth in response to the COVID-19 pandemic is likely to produce more data and may offer opportunities for more rigorous studies.

References

1. Rural health. *Health Aff (Millwood)*. 2019 Dec;38(12):1964-5. doi: 10.1377/hlthaff.2019.01365. PMID: 31794298.
2. Singh GK, Siahpush M. Widening rural-urban disparities in all-cause mortality and mortality from major causes of death in the USA, 1969-2009. *J Urban Health*. 2014 Apr;91(2):272-92. doi: 10.1007/s11524-013-9847-2. PMID: 24366854.
3. Garcia MC, Rossen LM, Bastian B, et al. Potentially excess deaths from the five leading causes of death in metropolitan and nonmetropolitan counties - United States, 2010-2017. *MMWR Surveill Summ*. 2019 Nov 8;68(10):1-11. doi: 10.15585/mmwr.ss6810a1. PMID: 31697657.

4. Hall JE, Moonesinghe R, Bouye K, et al. Racial/ethnic disparities in mortality: contributions and variations by rurality in the United States, 2012-2015. *Int J Environ Res Public Health*. 2019 Feb 2;16(3)doi: 10.3390/ijerph16030436. PMID: 30717345.
5. Villapiano N, Iwashyna TJ, Davis MM. Worsening Rural-Urban Gap in Hospital Mortality. *J Am Board Fam Med*. 2017 Nov-Dec;30(6):816-23. doi: 10.3122/jabfm.2017.06.170137. PMID: 29180557.
6. Keyes KM, Cerdá M, Brady JE, et al. Understanding the rural-urban differences in nonmedical prescription opioid use and abuse in the United States. *Am J Public Health*. 2014 Feb;104(2):e52-9. doi: 10.2105/ajph.2013.301709. PMID: 24328642.
7. Monnat SM, Rigg KK. Examining rural/urban differences in prescription opioid misuse among US adolescents. *J Rural Health*. 2016 Spring;32(2):204-18. doi: 10.1111/jrh.12141. PMID: 26344571.
8. Boring MA, Hootman JM, Liu Y, et al. Prevalence of arthritis and arthritis-attributable activity limitation by urban-rural county classification - United States, 2015. *MMWR Morb Mortal Wkly Rep*. 2017 May 26;66(20):527-32. doi: 10.15585/mmwr.mm6620a2. PMID: 28542117.
9. Croft JB, Wheaton AG, Liu Y, et al. Urban-rural county and state differences in chronic obstructive pulmonary disease - United States, 2015. *MMWR Morb Mortal Wkly Rep*. 2018 Feb 23;67(7):205-11. doi: 10.15585/mmwr.mm6707a1. PMID: 29470455.
10. Krishna S, Gillespie KN, McBride TM. Diabetes burden and access to preventive care in the rural United States. *J Rural Health*. 2010 Winter;26(1):3-11. doi: 10.1111/j.1748-0361.2009.00259.x. PMID: 20105262.
11. Primm K, Ferdinand AO, Callaghan T, et al. Congestive heart failure-related hospital deaths across the urban-rural continuum in the United States. *Prev Med Rep*. 2019 Dec;16:101007. doi: 10.1016/j.pmedr.2019.101007. PMID: 31799105.
12. Nelson JA, Kinder A, Johnson AS, et al. Differences in selected HIV care continuum outcomes among people residing in rural, urban, and metropolitan areas-28 US jurisdictions. *J Rural Health*. 2018 Dec;34(1):63-70. doi: 10.1111/jrh.12208. PMID: 27620836.
13. Zahnd WE, Rodriguez C, Jenkins WD. Rural-urban differences in human papillomavirus-associated cancer trends and rates. *J Rural Health*. 2019 Mar;35(2):208-15. doi: 10.1111/jrh.12305. PMID: 29808500.

Introduction

Background

Almost one-fifth of the population of the United States lives in rural areas. Health disparities for people living in rural areas are well studied¹ and include higher mortality²⁻⁵ and morbidity from a wide range of conditions such as substance/opioid abuse,^{6,7} chronic illnesses,⁸⁻¹¹ and HIV/AIDS, human papillomavirus, and other infectious diseases.^{12,13} Rural-urban health inequities have been extensively documented and the subject of research for several decades.^{2,14,15} Despite this recognition, inequities persist today and may be amplified by the COVID-19 pandemic¹⁶ which has increased the risk of morbidity and mortality^{17,18} and continues to impact access to health services.¹⁹⁻²¹ Underlying causes are complex and varied as they are related to macro and micro sociologic-demographic forces²² and economic trends.^{23,24} Research on health disparities continues to evolve from early work describing the extent of disparities, to attempts to understand the underlying reasons disparities exist, and finally to the current focus on identifying and implementing interventions to reduce these disparities.²⁵⁻²⁷ The need for interventions to reduce disparities has garnered additional attention due to impacts of the COVID-19 pandemic,^{28,29} race,³⁰ and structural racism³¹ on access to care³² and health outcomes.³³

Telehealth is the use of information and telecommunications technology to provide healthcare across time and/or distance.³⁴ This broad definition includes many modes of delivery (e.g., asynchronous, real-time video, and many others) and is the starting point for this review. Telehealth is a tool with the potential to increase access, improve the quality of care, increase patient satisfaction, positively impact patient outcomes, and reduce the cost of care. Telehealth includes using technology to directly deliver care but it can encompass broader applications of technologies to healthcare functions such as consultation, distance education, and mentoring, monitoring and data collection, and consumer outreach.³⁵ Provider-to-provider telehealth is a more specific, but still broad, telehealth application and is the focus of this review. We have defined provider-to-provider telehealth as any form of interactive support using telecommunications technology provided to healthcare professionals while they are caring for rural patients and populations. This includes using technology to support clinicians through mechanisms including consultations, mentoring, and education with the goal of improving the care they provide to individual patients or patient populations. More detailed definitions and descriptions that are specific to healthcare settings and clinical indications were used to group similar interventions in our synthesis, and are provided in **Appendix A, Appendix Table A-1**.

Application of telehealth had been steadily increasing in many areas of healthcare before the COVID-19 pandemic³⁶ began in 2020. The potential benefits of telehealth are frequently cited,^{37,38} and there is a sizable body of research on telehealth, including systematic reviews and reviews of reviews.³⁹⁻⁴⁴ Yet implementation and spread have been slow.⁴⁵⁻⁴⁷ Nevertheless, telehealth adoption appeared to be accelerating with improvement in technologies⁴⁸ and expansion of coverage by both public and private payers prior to the pandemic. The increase in use has been accompanied by an increase in the research and published literature on telehealth. Use of provider-to-provider telehealth for consultations has been studied across a range of clinical indications⁴⁹ including specialty care,⁵⁰ acute/emergency care,⁵¹ and intensive care.⁵² Additionally, other forms of provider-to-provider telehealth such as distance learning⁵³ and

Project Extension for Community Healthcare Outcomes^{54, 55} that combines provider education with review of specific cases are increasingly the subject of study.

The COVID-19 pandemic continues to create an environment that requires balancing the need to provide care while minimizing exposure and maximizing resources. Growth in the use of telehealth has been exponential, spurred by this need and supported by temporary changes in payment and regulation,⁵⁶ some of which may become permanent. Documentation of implementation and research on the effectiveness of this recent expansion of telehealth is now becoming available, though with the time lags necessary for data collection and for the measurement of patient-centered outcomes in addition to intermediate, process measures (e.g., numbers of telehealth encounters).

Identifying and synthesizing the available evidence about the use of provider-to-provider telehealth as a means of addressing rural health disparities can support ongoing spread, conversion of telehealth friendly pandemic policies to permanent support, and the identification of potential new areas and approaches for the expansion of telehealth in rural America.⁵⁷

Purpose and Scope of This Systematic Review

The scope of this review is provider-to-provider telehealth interventions designed to address the needs of rural patients and populations. This systematic review is intended to address a number of decisional dilemmas, including areas of uncertainty and lack of accessible evidence to support clinical, organizational and policy decision-making. The review first identified and synthesized the literature regarding the use of telehealth technologies to support provider-to-provider consultation in rural areas. In addition, the report identified known facilitators, barriers and strategies that are effective in promoting the adoption, implementation and sustainability of provider-to-provider telehealth for rural patients and populations. Finally, the project systematically categorized and summarized gaps in the evidence and the strengths and weaknesses of study designs to inform the design and conduct of future research. Narrow definitions of rural and provider-to-provider telehealth were avoided. We accepted study authors' designation of rural. The provider-to-provider telehealth scope is also broadly defined and includes interventions across a range of healthcare settings (inpatient, outpatient, emergency and educational setting) and clinical indications (e.g., stroke, diabetes, mental health).

This review is intended for a broad audience of stakeholders including but not limited to healthcare administrators, clinical leaders and policy makers interested in telehealth as an approach to address rural health disparities and improve the health and wellbeing of rural populations. It was specifically commissioned to inform the Pathways to Prevention workshop: *Improving Rural Health Through Telehealth-Guided Provider-to-Provider Communication*, developed by the National Institutes of Health (NIH) Office of Disease Prevention (ODP) in collaboration with the National Center for Advancing Translational Sciences; National Heart, Lung, and Blood Institute; Federal Office of Rural Health Policy at the Health Resources and Services Administration; and the Office of the Associate Director for Policy and Strategy, Centers for Disease Control and Prevention (hereafter referred to as the NIH/ODP Working Group). The workshop was convened by the NIH on October 12-14, 2021 (<https://prevention.nih.gov/research-priorities/research-needs-and-gaps/pathways-prevention/improving-rural-health-through-telehealth-guided-provider-provider-communication>).

Methods

Review Approach

This systematic review follows the methods suggested in the Agency for Healthcare Research and Quality (AHRQ) *Methods Guide for Effectiveness and Comparative Effectiveness Reviews* (hereafter “AHRQ Methods Guide”).⁵⁸ All methods were determined *a priori*, and a protocol was published on the AHRQ website (<https://effectivehealthcare.ahrq.gov/products/rural-telehealth/protocol>) and included in the PROSPERO systematic reviews registry (registration no. CRD42021233545). Below is a summary of the specific methods used in this review. A detailed description of the methods is provided in **Appendix A**.

Key Questions

The Key Questions for this systematic review are based on questions provided in the statement of work that accompanied the Request for Task Order. The questions were initially developed by the National Institutes of Health/Office of Disease Prevention (NIH/ODP) Working Group, and reviewed, reorganized, and refined by the systematic review project team; they were further revised after additional input from the AHRQ Task Order Officer (TOO) and the NIH/ODP Working Group.

Key Question 1. What is the uptake of different types of provider-to-provider telehealth in rural areas?

Key Question 2. What is the effectiveness of provider-to-provider telehealth for rural patients?

- a. How does provider-to-provider telehealth affect outcomes for patients and populations?
- b. How does provider-to-provider telehealth affect outcomes for healthcare providers?
- c. How does provider-to-provider telehealth affect outcomes for private and public payers?


Key Question 3. What strategies are effective and what are the barriers and facilitators to implementation and sustainability of provider-to-provider telehealth in rural areas?

Key Question 4. What are the methodological weaknesses of the included studies of provider-to-provider telehealth for rural patients and what improvements in study design (e.g., focus on relevant comparisons and outcomes) might increase the impact of future research?

Analytic Framework

The analytic framework (**Figure 1**) shows the target populations, interventions, and outcomes examined; the Key Questions are numbered and indicated in the framework.

Figure 1. Analytic framework



Abbreviation: KQ = Key Question.

Study Selection

We established criteria *a priori* to determine eligibility for inclusion and exclusion of abstracts in accordance with the AHRQ Methods Guide,⁵⁸ based on the Key Questions and population, intervention, comparisons, outcomes, settings, and study designs of interest (PICOS) framework. PICOS elements operationalized in terms of detailed inclusion and exclusion criteria are included in **Appendix Table A-1**.

Data Extraction and Risk of Bias Assessment

After studies were deemed to meet inclusion criteria, data were abstracted, including: study design, year, setting, country, sample size, patient and providers types and characteristics (e.g., age, sex, race, reason for presentation, diagnosis, provider training/background/scope of practice and primary care or specialty type), intervention characteristics (e.g., mode of delivery, duration or frequency, function) and results relevant to each Key Question as outlined in the PICOS

section in **Appendix A**. As Key Question 2 asks about outcomes for three groups: patients and populations, healthcare providers, and payers, outcomes were sorted by these groups. Data abstraction forms were developed after full text review and the data to be included in evidence tables was discussed with the AHRQ TOO and the NIH/ODP Working Group. Information relevant for assessing applicability included the number and diversity of settings or locations as well characteristics of the population, telehealth intervention or implementation strategy, and administering personnel. Sources of funding for all studies were also recorded. All study data was initially abstracted by one team member, then verified for accuracy and completeness by a second team member.

Predefined criteria were used to assess the risk of bias (also referred to as quality or internal validity) for each individual included effectiveness study, using criteria appropriate for the study designs. Controlled trials and observational studies were assessed using *a priori* established criteria consistent with the approach recommended in the chapter “Assessing the Risk of Bias of Individual Studies When Comparing Medical Interventions” in the AHRQ Methods Guide.⁵⁹ Studies were rated as “low risk of bias,” “medium risk of bias,” or “high risk of bias.” We did not exclude studies rated high risk of bias *a priori*, but high risk of bias studies are considered less reliable than low or medium risk of bias studies. For full details about the methods for data extraction and risk of bias assessment, see **Appendix A**.

Data Synthesis and Analysis

We constructed evidence tables identifying the study characteristics (as discussed above), results of interest, and risk of bias ratings for all included studies, and summary tables to highlight the main findings. As the Key Questions varied in nature and scope, our approach to synthesis differed.

For Key Question 2, the question about comparative effectiveness, we applied standard systematic review methods and reviewed and highlighted studies using a hierarchy-of-evidence approach, where the studies with better risk of bias ratings were given more weight in our synthesis for each clinical indication and outcome. Descriptive analysis and interpretation of the results were provided based on the direction and magnitude of effect. Meta-analyses were not performed as they would not producing meaningful results due to limited numbers of studies reporting similar outcomes, and heterogeneity among studies in design, patient population, and interventions.

Our response to Key Question 3 involved identifying and summarizing barriers and facilitators to implementation of provider-to-provider telehealth for rural healthcare. When studies directly compared different strategies, we provide a narrative summary of the studies and their results. When studies only reported on implementation for an individual intervention (not comparative), we abstracted what study authors described as barriers, facilitators, and indicators of implementation success. We applied an existing framework, the Consolidated Framework for Implementation Research (CFIR), to classify barriers and facilitators,⁶⁰ then summarized how frequently they were reported, both across all identified studies and then by setting.

For Key Question 1 (use of telehealth) and Key Question 4 (strengths and weaknesses of included studies) synthesis consists of descriptive narratives and tables, corresponding to the nature of the questions and data.

Grading the Strength of the Body of Evidence

The strength of evidence (SOE) for Key Questions 2a-2c (effectiveness) was determined for each clinical indication and major outcome category. SOE was initially assessed by one researcher and confirmed by a second using the approach described in the AHRQ Methods Guide.⁵⁹ The body of evidence was assessed using the following criteria as they are defined in the AHRQ Methods Guide:

- Study limitations (low, medium, or high level of study limitations)
- Consistency (consistent, inconsistent, or unknown/not applicable)
- Directness (direct or indirect)
- Precision (precise or imprecise)

For definitions of the ratings used to categorize the strength of evidence, see **Appendix Table A-2**.

SOE and the corresponding conclusions are expressed in terms of whether the outcome measured and analyzed in the studies is better, worse, or similar with telehealth compared to typical provider-to-provider interactions without telehealth, often referred to in studies as usual care. However, usual care could have different meanings including in-person interactions or no interactions. For this reason, we have provided detailed descriptions of usual care when they were provided by the study authors.

In studies of telehealth, interpreting results requires consideration of the context and the intended function of telehealth. When outcomes are better or worse, the interpretation is relatively clear. If telehealth consultations are used to provide access to specialized knowledge and patient outcomes are found to be better, telehealth is providing a benefit. If the opposite is true, and a study finds patient outcomes are worse, then telehealth is having a negative impact or causing harm. Drawing a conclusion is less straightforward when patient outcomes are found to be similar with and without telehealth. Telehealth is often used to allow healthcare to be delivered in rural locations rather than transferring the patient or requiring travel. If patient outcomes are similar in rural locations, telehealth may be beneficial, if it reduced travel burdens and this was the intention. However, determining if similar outcomes equals a benefit depends on considering multiple factors, such as resources needed and how perspectives may differ (e.g., what is most important to a patient may not be what is most important to a health system). For this reason, we report when outcomes are similar in results, and then discuss the context to help facilitate conclusions about whether similar outcomes with telehealth can be interpreted as a benefit.

Results

Description of Included Evidence

A total of 6,329 references were reviewed, including 5,973 from electronic database searches and 356 from reference lists and other systematic reviews. After dual review of titles and abstracts, 1,024 articles were selected for full-text review, of which 166 studies in 179 publications were included in this review. Search results and selection of studies are summarized in the literature flow diagram (**Appendix B, Appendix Figure B-1**). Results are arranged by Key Question and then by clinical indication in tables and summarized in accompanying text.

Characteristics of included studies are detailed in **Tables B-1, B-2, and B-3**. A list of included studies can be found in **Appendix C**. Data abstraction of study characteristics and results, quality assessment for all included studies, and details for grading strength of evidence (SOE) are available in **Appendixes D, E, and F**, respectively. A list of excluded studies with reasons for exclusion are included in **Appendix G**. A list of references appearing in the appendix is available in **Appendix H**.

Key Question 1. What is the uptake of different types of provider-to-provider telehealth in rural areas?

Key Points

- We did not identify any surveys or national datasets that provided a comprehensive assessment of the use of provider-to-provider telehealth for rural populations in the United States.
- Seven published reports provide data about regional or national use for specific clinical applications. These include telepsychiatry, telehealth in emergency care, and telehealth use by hospitals for intensive care unit (ICU), stroke and heart attack care.
- While use varies across states and regions, telehealth use remains low, with a minority of rural hospitals reporting telehealth use in the included studies and less use compared to urban areas, though telehealth use has increased over time.

Summary of Findings

To address this Key Question we searched for and included any studies published since 2015 that reported the use of provider-to-provider telehealth in rural areas. We also searched gray literature and asked experts and stakeholders for any known studies or reports (details on the search are included in **Appendix A**). The identified evidence to address Key Question 1 was limited. No national surveys or datasets that provided a comprehensive description or global estimate of provider-to-provider telehealth in rural areas generally were identified through our literature search. Additionally, no supplemental evidence for Key Question 1 was identified following the search process outlined in **Appendix B**.

We did however identify seven studies reporting the uptake of different types of provider-to-provider telehealth for a specific use in rural areas.⁶¹⁻⁶⁷ All studies were regional or national in focus, and all were conducted in the United States. Six studies report the results of surveys,⁶¹⁻⁶⁶ while one study performed a secondary analysis on healthcare claims data.⁶⁷ Telehealth modalities represented across the seven studies include telepsychiatry, emergency telehealth,

remote ICU, telestroke and telecardiology. Risk of bias ratings and grading the SOE were not appropriate for the descriptive nature of this Key Question. The studies that reported on national surveys and trends are summarized in **Table 1** and additional details can be found in **Appendix Tables B-1 and D-1**.

These studies reported relatively low rates of telehealth adoption and most reported lower rates in rural than urban area. All that examined change over time reported increases in rural as well as urban hospital and the results from these surveys underscore variation in use across specialties and regions. Surveys documented that telepsychiatry use increased over time from 15.2 percent of mental health facilities in 2010 to 29.2 percent in 2017. There was wide variability across states, and telehealth was more commonly used by facilities in medically underserved and rural areas.⁶⁴ Rural location, Critical Access Hospital designation, and higher annual total visit volumes were associated with higher likelihood of telepsychiatry use.⁶³

Forty-six percent of rural emergency departments (EDs) responding to a survey reported they were not using telehealth, with the Southern Region of the United States significantly less likely to use telehealth than other regions.⁶⁵ When telehealth was used it was more often in EDs that are rural, that do not have 24-hour per day neurologist availability, and that had an annual volume greater than 20,000 visits.⁶⁶

A study of the use of multiple types of telehealth by hospitals in 2018 found that urban hospitals were twice as likely as rural hospitals to use telehealth for ICU and stroke care.⁶² Similarly, a study of use of telehealth for heart attack and stroke reported that use by rural hospitals increased from 6 percent in 2012 to 16 percent in 2017, though these rates are still lower than those for urban hospitals.⁶¹ In both of these studies, smaller hospitals (fewer beds) were less likely to have adopted telehealth.

The only included study that used claims data, reported that between 2008 and 2015, the proportion of ischemic stroke patients receiving telestroke services increased from 0.4 to 3.8 per 1,000 among fee-for-service Medicare beneficiaries and use of telestroke services increased most rapidly among rural residents. Patients dually eligible for Medicare and Medicaid had a lower proportion of telestroke care use than Medicare-only patients residing in rural and super rural counties as defined by the study authors, but its use was similar among both groups of patients living in urban counties.⁶⁷

Table 1. Key Question 1: summary of provider-to-provider telehealth use in rural areas

Clinical Topic	Modality	Number of Studies	Data Source Sample	Key Findings
Pharmacy	Telepharmacy	1	Data Source Researcher initiated survey and interviews Sample 50 state offices of rural health Followup interviews: 10 states	Use of telepharmacy in rural hospitals varied across states but was not widely adopted ⁶⁸ Telepharmacy models differed according to area, state regulations, hospital ownership, and hospital size and medication order volume ⁶⁸
Mental health	Telepsychiatry	2	Data Sources 2016 National Survey of U.S. Emergency Departments ⁶³ 2010-2017 waves of the National Mental Health Services Survey ⁶⁴ Sample 4,507 of 5,375 (84% response)	20% of all U.S. Emergency Departments reported utilizing telepsychiatry services. The most common applications: admission or discharge decisions (80%) transfer coordination (76%). ⁶³ Use increased over time 2010: 15.2% of mental health facilities (n=1,580) 2017: 29.2% (n=3,385) OR=2.30, 95% CI 1.96 to 2.69, with wide variability among states. ⁶⁴
Emergency Care	Emergency Telehealth	2	Data Sources 2014 National Emergency Department Inventory-New England ⁶⁶ 2016 National Emergency Department Inventory Survey ⁶⁵ Sample 169 of 195 (87% response) ⁶⁶ 977 rural emergency departments ⁶⁵	49% of New England Emergency Departments (12% Rural, 7% Urban) report using telehealth, most commonly for neurology/stroke, pediatrics, psychiatry, and trauma. ⁶⁶ 46% of rural emergency departments in the U.S. did not report using telehealth, with the Southern Region of the U.S. significantly less likely to use telehealth than other regions. ⁶⁵ Telehealth was more often used in emergency departments that are rural, that do not have 24 hour 7 days per week neurologist availability, and with an annual volume of $\geq 20,000$ visits. ⁶⁶
Stroke	Telestroke	1	Data Sources 2008-2015 Medicare fee-for-service administrative claims data ⁶⁷ Sample 1,002,2245 ischemic stroke hospitalizations	2008 to 2015 for fee-for-service Medicare beneficiaries: Increases in ischemic stroke cases receiving telestroke services Overall: 0.4 to 3.8 per 1,000 Rural residents: 0.6 to 8.6 per 1,000 Super rural: 1.0 to 6.1 per 1,000 Urban: 0.3 to 2.3 per 1,000. ⁶⁷

Clinical Topic	Modality	Number of Studies	Data Source Sample	Key Findings
Stroke and ICU	Telestroke and remote ICU	1	Data Source 2018 American Hospital Association Survey ⁶² Sample 781 rural hospitals 2,756 metropolitan and micropolitan*	Metropolitan areas* are more than twice as likely to adopt telehealth for ICU and stroke. eICU 0.48 vs. 0.13 Stroke 0.43 vs. 0.21 Small numbers of beds and lack of health information change capacity explained some of this difference. ⁶²
Stroke and Heart Attack	Telestroke or Telecardiology	1	Data Sources 2012-2017 Health Information Management and System Society Dorenfest Database and Healthcare Cost Report Information System Sample 2,012 rural hospitals 2,096 not rural	Percentage of hospitals with telemedicine, change 2012 to 2017 Rural: 6.31% to 16.45% Urban: 7.30% to 19.42% In rural hospitals government vs. private hospitals, larger hospitals (more beds) and less Medicare patients, and a higher number of ED visits were associated with higher likelihood of telehealth adoption. ⁶¹

Abbreviations: CI = confidence interval; ED = emergency department; ICU = intensive care unit; OR = odds ratio

*Metropolitan areas have at least 1 core urbanized area of at 50,000 people; micropolitan at least 10,000 people but less than 50,000.

Key Question 2. What is the effectiveness of provider-to-provider telehealth for rural patients?

- How does provider-to-provider telehealth affect outcomes for patients and populations?
- How does provider-to-provider telehealth affect outcomes for healthcare providers?
- How does provider-to-provider telehealth affect outcomes for private and public payers?

Key Question 2 presents the available evidence about the effectiveness of provider-to-provider telehealth for rural populations. The following sections summarize the results of the 97 comparative studies in 106 publications we identified using the search and selection process described in the methods section above. Full details of the search and selection process are provided in **Appendix Figure B-1**. We included research that evaluated the effect of telehealth on outcomes for patients, providers, and payers. As these studies include diverse uses of telehealth in different settings for different clinical indications, we have organized results first by setting (i.e., in-patient, out-patient, emergency care and education) and then by clinical indication within settings. This facilitates summarizing evidence that is likely to be comparable. Consideration of findings across settings and broader interpretations are provided in the Discussion section of this report.

Inpatient

Key Points

- Telehealth consultations as part of inpatient care at a rural hospital may result in similar lengths of hospital stay (6 studies; Low SOE) and rates of transfers (3 studies; Low SOE) compared to usual care that involves in person or phone consultations.
- Telehealth supporting care for neonates at rural hospitals may result in similar clinical outcomes when compared to transfer and care at a hospital with a Level 4 neonatal intensive care unit (NICU) (2 studies; Low SOE) and may result in more appropriate transfers (2 studies; Low SOE).
- Remote ICU care in rural hospitals result in no difference in mortality rates compared to transferring patients to more distant locations for ICU care (2 studies; Low SOE) while evidence is insufficient to support conclusions about the impact of remote ICUs in rural area on transfers. (1 study, insufficient evidence).

Summary of Findings

We identified 17 studies reported in 18 articles that evaluated provider-to-provider telehealth for inpatient care of rural populations. Five studies reported in six articles evaluated teleconsultations for care of neonates,⁶⁹⁻⁷⁴ three for infectious disease,⁷⁵⁻⁷⁷ one for spinal fractures,⁷⁸ one for stroke,⁷⁹ one for mental and behavioral disorders,⁸⁰ and two studied remote ICU.^{81, 82} Additionally, five studies evaluated telehealth facilitated rounds and remote consultations for multiple conditions⁸³⁻⁸⁶, include one that focused on pediatric patients.⁸⁴ The majority (13) were studies that compared outcomes before and after the implementation of telehealth,^{69, 72, 75-81, 84-86} two studies reported in three articles were prospective cohort studies,^{70, 71, 74} and three were retrospective cohort studies.^{73, 82, 83} Eleven were conducted in the United States,^{69, 70, 72-74, 76, 77, 82-85} five in Australia^{75, 78, 80, 81, 86} and one in Scotland.⁷⁹ Risk of bias was rated as high for six studies^{69, 72, 75, 77, 78, 85} and medium for eleven studies (**Appendix E**).^{70, 73, 74, 76, 79-84, 86} Additional descriptive information about these studies and key results are included in **Appendix Tables D-2 and D-3**.

The key results are organized by clinical indication in **Table 2** below. Given the small number of studies, we assessed the body of evidence for the same outcome across similar interventions to determine the strength of possible conclusions (SOE details in **Appendix F**).

We considered studies of teleconsultations for infectious disease, stroke, mental and behavioral disorders, spinal fracture and multiple conditions addressed via teleconsultation rounds together as these connected specialists with the inpatient care team to assist in the management of specific patient issues. The evidence was insufficient to support a conclusion about the impact on mortality based on three studies with imprecise and inconsistent results, with one study reporting a decline in mortality after infectious disease teleconsultations were available (odds ratio [OR] 0.3; 95% confidence interval [CI], 0.2 to 0.7)⁷⁶ and two studies reporting no significant differences in mortality after instituting teleconsultation rounds of various types (1.7% vs. 2.3%; 4% vs. 1%).^{83, 85}

Six studies of inpatient consultations provide low SOE that telehealth resulted in similar patient length of stay. The SOE is low as their results were inconsistent with one study reporting an increase in hospitals days with infectious disease teleconsultation versus pre-telehealth phone consultations (OR 1.3; 95% CI, 1.2 to 1.5);⁷⁶ one finding a decrease with remote consultations for spinal fractures versus in-person consultations requiring patient transfer to specialist and back

to local hospital (9.25 vs. 4.2, $p < 0.01$);⁷⁸ and four reported finding no significant difference in length of stay between telehealth and pre-telehealth (mean number of days 2.55 vs. 3.74, $p > 0.05$; median days: 17 [interquartile range [IQR] 9.5 to 31] vs. 29 [IQR 18.75 to 41]; mean (standard deviation [SD]) days: 2.82 (2.49) vs. 2.55 (3.059); $p = 0.136$; mean days: 1.92 vs. 1.94; $p = 0.896$).^{75, 79, 83, 85} Three studies^{76, 80, 85} reported and supported a conclusion that telehealth resulted in similar or fewer patient transfers (Low SOE).

We summarized the results of studies of telehealth consultation and collaborative care of neonates separately as these programs often include mechanisms to transfer imaging and monitoring data as well as video for examination and discussion. One study ($n = 384$ hospitals) reported that mortality declined with telehealth in the hospitals using telehealth and that infant mortality declined state wide.⁷² Two smaller studies ($n = 143$ and 155)^{73, 74} provided consistent but imprecise results that clinical outcomes at the regional center with telehealth were similar to (not different from) those achieved when neonates were transferred to a hospital with level 4 NICU. This was consistent with the findings of a study of neonate care in 9 hospitals in Arkansas before and after telehealth (3 studies; Low SOE). Two studies also reported that with telehealth, transfers were more appropriate. In one study this meant low birth weight babies were not delivered in hospitals without NICUs,⁷² while in the other it meant that few transfers were needed and transferred patients were more stable.⁷⁰ The evidence is insufficient to support a conclusion about the impact of neonatal teleconsultations on length of stay as this outcome was only reported in one study.

Two included studies evaluated remote ICUs. A case control study of U.S. hospitals based on Medicare claims data reported no significant difference in mortality in rural hospitals that had adopted telehealth for ICU compared to control hospitals (OR 1.06 [95% CI 0.99 to 1.13]).⁸² One included study evaluated a remote ICU program in Australia and reported no difference in mortality but a significant reduction in transfers (31.8% to 22.9%, relative risk [RR] 0.88, 95% CI 0.80 to 0.98).⁸¹ As the two studies that consistently reported no difference in mortality with remote ICU were conducted in different countries and used observational designs susceptible bias, we rated the SOE as low (Low SOE). Only one of these studies reported on transfers and we consider this evidence insufficient to support a conclusion.

None of these studies of inpatient provider-to-provider telehealth reported payer outcomes or costs and only one study of teleconsultation for spinal fracture⁷⁸ reported an improvement in provider's knowledge, skills and confidence.

Table 2. Key Question 2: summary of inpatient findings

Clinical Topic N Studies	Provider Specialty	Consultant Specialty	Patient Outcomes: Mortality*	Patient Outcomes: Hospital Use*	Patient Outcomes: Other Clinical*	Provider Outcomes/ Payer Outcomes*
Multiple conditions 4	Physicians, nurse practitioners, nurses	Hospitalist, pharmacist	~ Mortality in hospital ^{83, 85}	~ Transfers ⁸⁵ ~ Length of stay ^{83, 85} ~ Readmission ⁸³	Not reported	+ Communication ratings ⁸⁵ + Hospital Revenue ⁸⁴ + Professional Billing Revenue ⁸⁴ ~ Drug prescribing outcomes ⁸⁶
Infectious Disease 3	Multidisciplinary	Infectious disease specialist physicians and pharmacists	+ Mortality ⁷⁶	~ Transfers ⁷⁶ - Length of stay ^{76†} ~ Length of stay ⁷⁵ ~ 30-day Readmission ⁷⁶	Not reported	+ Provider satisfaction ⁷⁶ + Improved antimicrobial use and infection rate ^{75, 77} ~ Antibiotic use ⁷⁵ + Appropriate prescribing and adherence to guidelines ⁷⁵
Stroke 1	Stroke-unit multidisciplinary team	Stroke physician	Not reported	~ Length of stay ⁷⁹	None reported	+ Cost ⁷⁹
Mental and behavioral disorders 1	Community-based mental health providers	Psychiatrist or senior mental health provider	Not reported	+ Transfers ⁸⁰	Not reported	Not reported
Spinal fracture 1	ED medical officer	Physiotherapist, clinical nurse consultant	Not reported	+ Length of stay ⁷⁸	None reported	+ Knowledge, skills, confidence ⁷⁸

Clinical Topic N Studies	Provider Specialty	Consultant Specialty	Patient Outcomes: Mortality*	Patient Outcomes: Hospital Use*	Patient Outcomes: Other Clinical*	Provider Outcomes/ Payer Outcomes*
Neonates 5	Pediatricians, Nurse practitioners, family medicine, neonatology	Intensivists, Neonatologists	✚ Statewide infant mortality to 1 year and telehealth hospital death before discharge ⁷²	✚ Transfers ^{70, 72} ~ Length of stay ^{73, 74}	~ Enteral feeding ^{73, 74} ~ Ventilation-oxygen ^{73, 74} ~ Proportion of deliveries at community hospitals ⁶⁹ ~ Morbidity ⁷²	None reported
ICU 2	General practitioners and nurses	ICU team	~ Mortality in high dependency unit ⁸¹ ~ Mortality in hospital ⁸¹ ~ Mortality total ⁸¹ ✚ Mortality 90-day ⁸²	✚ Transfers ⁸¹	None reported	None reported

*Symbol meaning: ✚ = Improved Outcome with telehealth; ~ = Similar outcome with telehealth; — = Worse outcome with telehealth

[†]Length of stay was longer with telehealth, and longer stays are usually considered a negative outcome. However mortality decreased which may indicate that more care was appropriate. The authors note that the consultations tended to occur later in the hospital stay in this study and a study in which the consultation happened sooner would provide clearer evidence about the impact on length of stay.

Abbreviations: ICU = intensive care unit; NICU = newborn intensive care unit.

Outpatient Care and Services

Key Points

- Outpatient telehealth consultations with specialists may result in improvements in clinical outcomes compared to care without specialist involvement:
 - For patients with diabetes:
 - Some improvement in medication adherence and treatment response for patients with depression. (3 studies; Low SOE)
 - Improvements in A1c and self-management but no effect on blood pressure or cholesterol levels in patients with diabetes (3 studies; Low SOE)
 - Improvements in A1c, fasting glucose, and blood pressure in patients with hypertension and diabetes with pharmacy teleconsultations. (2 studies; Low SOE)
 - For patients with depression:
 - Higher utilization and corresponding costs for outpatient consultations for depression are associated with increased access from telehealth, but cost-effectiveness analyses report overall benefit. (2 studies; Low SOE)
- Telehealth pharmacy consultations improved guideline adherence and patient outcomes in subgroups of patients with both diabetes and hypertension. (2 studies; Low SOE)

Summary of Findings

Thirty-two studies (in 35 publications) evaluated the use of provider-to-provider telehealth interventions to support outpatient care for rural populations.⁸⁷⁻¹²¹ Four studies assessed diabetes care;^{90, 101, 103, 106} three studies assessed telehealth for depression (in 7 publications);^{93, 95, 97, 109-111, 122} and three studies addressed remote consultations;^{94, 100, 102} two studies each assessed telepharmacy,^{87, 89} rheumatology,^{116, 120} dermatology,^{92, 119} and oncology;^{114, 117} and one study each assessed telehealth for echocardiography,⁸⁸ endoscopy,¹¹⁸ hemodialysis,¹¹³ blood pressure control,¹²¹ fracture,¹⁰⁵ dementia,⁹⁹ hepatitis C,¹¹² attention-deficit/hyperactivity disorder,¹⁰⁷ post-traumatic stress disorder (in two publications),^{96, 108} diabetic retinopathy screening,⁹¹ palliative care,⁹⁸ wound care, and ultrasound in pregnancy.^{104, 115} Sixteen studies (reported in 17 publications) were randomized controlled trials (RCTs),^{88-90, 92, 93, 95-97, 102, 103, 107-110, 116, 119, 121} six were prospective cohort studies,^{91, 94, 98, 99, 111, 120} six were retrospective cohort studies,^{101, 105, 112, 114, 117, 118} four were pre-post study designs (same group measured before and after implementation),^{87, 100, 106, 113} and two were before-after studies (different groups/systems measured before and after implementation).^{104, 115} Eleven of the studies were conducted in the United States;^{87, 89, 92, 93, 95-97, 104, 106, 107, 112, 119, 120} the remainder were conducted in Canada,^{102, 113, 116} Australia,^{91, 98, 105, 117} Korea,^{90, 99, 100} China,¹²¹ Denmark,¹⁰¹ New Zealand,¹¹⁴ Spain,⁹⁴ United Kingdom,^{115, 118} Chile,¹¹¹ Taiwan,¹⁰³ and Sweden.⁸⁸ Risk of bias was rated as high for eight studies,^{88, 100, 101, 104-106, 115, 118} medium for 21 studies,^{87, 89-92, 94, 96-99, 103, 107, 111-114, 116, 117, 119, 120, 123} and low for two studies (**Appendix E**).^{102, 121} Additional details can be found in **Table 3**, and **Appendix Tables D-4 and D-5**.

Depression

Three studies (2 trials^{97, 122} [n=364 and n=395] and 1 prospective cohort¹¹¹ [n=250]) evaluated telehealth-based care for depression in rural settings.^{93, 95, 97, 109-111, 122} The providers in these studies were primary care clinicians, including physicians, psychologists, social workers, midwives, and pharmacists; remote consultations added depression care manager nurses, psychologists, and psychiatrists. All studies compared telehealth-based care with usual practice-based, in-person care without the specialist. One trial reported mixed results with better, but not statistically significantly different, medication adherence at 6 months and 12 months, improved response to treatment at 6 months (but not at 12 months), remission at 12 months (but not at 6 months), 36 item short form survey (SF-36) Mental Component Score at 12 months (but not 6 months), and Quality of Well-Being scores at 6 months (but not at 12 months); treatment satisfaction was superior in the telehealth group at 6 and 12 months.^{96, 122} A subanalysis⁹³ of this trial reported racial differences in response, and found minority group status remaining a significant moderator of the intervention's effect even after adjusting for other factors associated with minority status (adjusted odds ratio [aOR] 6.02; 95% CI 1.48-24.30). Minority participants were more likely to have symptoms improve (i.e., response to treatment) with telehealth but white participants were not. Another subanalysis did not find a difference in depression-free days for either white or nonwhite participants.¹¹⁰ One longer trial reported improved response to treatment (OR 3.26; 95% CI, 1.95 to 5.47) and remission at 18 months (OR 3.15; 95% CI, 1.62 to 6.09),⁹⁷ as well as improvements in SF-36 Mental Component Score (difference 4.75, p=0.002) and Hopkins Symptoms Checklist (difference -0.36, p<0.001), but did not report differences in satisfaction or Quality of Well-Being (difference 0.02, p=0.22). Telehealth-based collaborative care for depression was significantly more expensive than office-based team care when total outpatient costs (difference 391.20, p=0.012),^{95, 109, 110} or depression-related costs were considered in one study (difference 97.42, p=0.013).⁹⁵ However, appropriate utilization increased, especially for depression-related care (difference 1.97, p=0.001),¹⁰⁹ and the incremental cost effectiveness for depression-free days (\$10.78/day) was considered low.

Attention Deficit/Hyperactivity Disorder (ADHD)

One trial compared treatment of ADHD in two telehealth models used to augment primary care.¹⁰⁷ The more intensive intervention included six sessions of pharmacotherapy and caregiver behavior training provided by community therapists with remote supervision by a psychiatrist, while the standard telehealth consultation care group received treatment by their primary care clinicians augmented with a telepsychiatry consultation. Children in both models improved, and although the improvement was greater with the more intensive intervention, the difference was not statistically significant (OR 1.34; 95% CI, 0.55 to 3.24).

Post-Traumatic Stress Disorder (PTSD)

One trial (n=225; reported in two publications) evaluated telehealth-based care for PTSD compared to practice-based care.^{96, 108} Primary care physicians, psychiatric advanced practice nurses, and master's degree-level social workers in the intervention group consulted with remote telephone nurse care managers, clinical pharmacists, psychologists, and psychiatrists. Patients in the telehealth group were more likely to be prescribed a medication for PTSD in the first 6 months (93% vs. 86%) and to be prescribed prazosin hydrochloride at 6 months (27% vs. 15%) and 12 months (32% vs. 11%). Telehealth did not result in differences in adherence (adherent at least 80% of the time; OR 0.91; 95% CI, 0.47 to 1.78) or the proportion of patients with a

psychiatric encounter at 12 months (33% vs. 41%). At 12 months, patients in the telehealth group were more likely to have cognitive processing therapy (OR 18.08; 95% CI, 7.96 to 41.06), and to have received more sessions (OR 7.86; 95% CI, 3.15 to 19.61). They also had greater improvement in Posttraumatic Diagnostic Scale scores (-4.17 [SD=9.82] vs. -1.32 [SD=8.79]; $p=0.04$), Hopkins Symptom Checklist (-0.43 [SD=0.72] vs. -0.23 [SD=0.62]; $p=0.01$), and SF-36 Physical Component Score (-1.02 [SD=8.30] vs. -1.56 [SD=8.30]; $p=0.35$) but did not result in difference in the SF-36 Mental Component Score (2.72 [SD=11.92] vs. 4.05 [SD=10.07]; $p=0.36$). The telehealth group had higher outpatient mental health specialty costs (\$2,964.63 vs. \$2,159.26; $p=0.01$), total outpatient costs (\$8,150.20 vs. \$6,944.13; $p=0.02$), and telephone and non-telephone outpatient PTSD-specific care costs (\$799.61 vs. \$6.94; $p=0.01$ and \$238.00 vs. \$174.96; $p=0.03$). Treatment effectiveness differences were very small (0.008 for Quality of Well-Being and 0.001 for 12 item short form [SF-12] for Veterans), with rather large cost differences (\$2,495/patient) resulting in very high incremental cost-effectiveness ratios (\$185,565 for Quality of Well-Being and \$138,108 for SF-12 for Veterans).

Diabetes

Four studies (two trials^{90, 103} [n=71 and n=95], one retrospective cohort study¹⁰¹ [n=78 telehealth patients compared to national data] and one pre-post study¹⁰⁶ [n=59]) assessed telehealth for diabetes care in rural settings.^{90, 101, 103, 106} Providers in these studies were primary care clinicians,^{103, 106} including nurses.^{90, 101} Consultants involved in the telehealth-based care were nurses, dietitians, and diabetes specialists. Three studies compared telehealth to in-person care; the trials compared telehealth to in-person diabetes education, and the pre-post study compared the time before implementation of the telehealth program, when patients were referred to an endocrinologist in a distant secondary/tertiary hospital, to the period after telehealth implementation.^{90, 103, 106} The three studies all found telehealth was associated with improved management of glycated hemoglobin (A1c; $p=0.002$; -0.5%, $p<0.01$; 0.7 vs. 0.1, $p=0.03$; Effect size -0.49 [95% CI -0.87 to -0.09], $p=0.002$).^{90, 103, 106} Additionally, a pre-post study reported improvements in self-monitoring of blood glucose and in Diabetes Self-Management Education scores with telehealth compared to care provided prior to telehealth implementation (1.65 [95% CI 0.94-2.30]; $p<0.001$).¹⁰⁶ One trial did not report a difference in systolic blood pressure (Change from baseline -1.0 [SD=14.9] vs. -6.7 [SD=17.3]; $p=0.057$),¹⁰³ and the pre-post study did not report a difference in systolic blood pressure (Effect size -0.31 [95% CI -0.74 to 0.11]; $p=0.08$).¹⁰⁶ One study compared telehealth findings to a national diabetes registry (n=78), and reported greater improvements in A1c levels for telehealth participants with Type 1 and Type 2 diabetes.¹⁰¹

An Australian prospective cohort study (n=1,024) also evaluated remote diabetic retinopathy screening compared to usual in-person screening.⁹¹ The use of a remote ophthalmologist to perform diabetic retinopathy screening in cooperation with in-person primary care clinicians was associated with increased rate of referral or reminder for screening, as well as an improved screening rate (100% vs. 41%; RR 2.44; 95% CI, 2.14 to 2.79).

Hepatitis C

One retrospective cohort study (n=80) compared telehealth-based hepatitis C virus treatment at rural sites with treatment at a hepatology clinic.¹¹² Primary care clinicians at the rural sites consulted with remote specialists via videoconference. The study did not report differences in sustained virologic response (55% vs. 43%; $p=0.36$) or in mean weeks of therapy, but did report

a higher proportion of patients completing therapy (78% vs. 53%; $p=0.03$), higher mean number of face-to-face patient visits and face-to-face visits per week, and lower rates of anemia and patient withdrawal due to adverse events (RR 0.42; 95% CI, 0.21 to 0.85).

Rheumatology

Two studies (one trial¹¹⁶ [n=85] and one prospective cohort¹²⁰ [n=85]) evaluated telehealth-based rheumatology in rural settings. Both compared telehealth consultations with in-person rheumatology consultations. The in-person providers in the trial were physical therapists, while those in the cohort study were rheumatology graduate medical education trainees; both studies used rheumatologists to provide the remote consultations. The trial did not report differences in the Rheumatoid Arthritis Disease Activity Index (difference 0.9, 95% CI -2.4 to 0.5), the EQ-5D Questionnaire (difference -0.1, 95% CI -0.1 to 0.4), the modified Health Assessment Questionnaire (difference 0.2, 95% CI -0.5 to 0.1), the Disease Activity in 28 Joints with c-reactive protein (CRP) Measure Score (difference 0.9, 95% CI -3.1 to 1.2), or the 9-item Visit-specific Satisfaction Score (data not reported).¹¹⁶ The prospective cohort study reported benefits of telerheumatology in reducing distances traveled to visit (difference, -384.8 miles/visit; $p<0.01$) and cost of visits (difference, -\$113.80/visit; $p<0.01$).¹²⁰

Oncology

Two retrospective cohort studies (n=110 and n=147) evaluated telehealth-based oncology in rural settings.^{114, 117} One study focused on lung cancer,¹¹⁴ while the other included any cancer diagnosis.¹¹⁷ Both compared telehealth to in-person/usual care. The lung cancer study compared video conference thoracic radiology multidisciplinary meetings with co-located meetings, and did not report any differences in proportion of patients receiving radiotherapy (71% vs. 71%) or in any other management measure (including clinic visits, offering of radiotherapy, receipt of radiotherapy, and completion of radiotherapy).¹¹⁴ The other study of multiple cancers combined reported lower total costs with telehealth compared to hypothetical travel costs (\$442,276 vs. \$762,394; net savings, \$320,118).¹¹⁷

Echocardiology

One small (n=38) trial⁸⁸ compared robot-assisted videoconference echocardiography and teleconsultation to in-person echocardiography and consultation at a secondary/tertiary hospital in patients with symptoms of heart failure in a rural setting. The study compared remote echocardiography performed by a trained sonographer and subsequent teleconsultation among the patient, general practitioner, sonographer, and remote cardiologist with usual care referral to a secondary/tertiary hospital cardiologist. The study reported beneficial reductions in total process time (median 27 vs. 114 days; $p<0.001$) and time from randomization to attaining a specialist consultation (median 12 vs. 86 days; $p<0.001$).

Hemodialysis

One small (n=19) pre-post study compared telehemodialysis to usual care in a rural Canadian First Nation setting.¹¹³ The study compared healthcare utilization before and after introduction of the telehemodialysis program, which provided hospital-based hemodialysis care in the community (by nurses, nutritionists, and general practitioners) with remote supervision by nephrologists in a university dialysis center. The study reported increased amounts of urea

removed (effect size 5.3, $p=0.03$), but did not report differences in other clinical measures, indicating compliance with dialysis best practices.

Fracture

One small ($n=12$) retrospective cohort study compared a telehealth fracture clinic with in-person care in a rural setting.¹⁰⁵ The consultation was a video conference with remote orthopedic specialists in real time. The study reported lower costs compared to transferring patients (savings of \$5941/patient) or sending a specialist to rural locations (savings of \$1779/patient).

Dementia

One prospective cohort study ($n=188$) compared remote dementia care to in-person care in a dementia clinic.⁹⁹ Primary care clinicians and their patients at a rural public health center used a videoconference system to consult with a dementia specialist. The study did not find differences in mean change in Mini-Mental Status Exam (MMSE) scores per year in the study overall (-0.60 vs. -1.03 ; $p=0.29$); however, patients with higher baseline MMSE (15-30) showed significantly less deterioration with telehealth consultations (-0.62 vs. -1.59 ; $p=0.049$).

Wound Care

One study ($n=54$) compared wound care outcomes before and after introduction of a telehealth consultation system.¹¹⁵ Primary care clinicians used the store-and-forward telehealth system to receive consultation from wound specialists in secondary care centers. The study reported decreased waiting times from referral to appointment (median 18 vs. 47 days, p -value not reported), as well as improvements in leg ulcer healing time (median 70 vs. 105 days, p -value not reported), but did not result in differences in cost of care per patient (£665 vs. £651).

Pharmacy

Two studies (one trial⁸⁹ [$n=302$] and one pre-post study⁸⁷ [$n=577$]) assessed telepharmacy in rural settings. The providers were pharmacy technicians⁸⁹ or in-person clinic staff,⁸⁷ and the remote consultants were pharmacists. The trial evaluated pharmacy for a broad range of conditions common to a family medicine clinic, and reported improved guideline adherence with telehealth compared to usual care, though this was not statistically significant (5.0% [SD 2.4%]; 95% CI, -0.5 to 10.4% ; $p=0.07$).⁸⁹ The pre-post study evaluated pharmacy for management of diabetes and hypertension⁸⁷ and reported improvements after telepharmacy was implemented in A1c (-0.2 [SD 1.1]; $p=0.008$), fasting blood glucose (-10.6 [SD 39.4]; $p<0.0001$), and systolic blood pressure (-0.37 [SD 19.4]; $p=0.009$) only in patients with both diabetes and hypertension; telehealth did not result in differences in these outcomes for patients with either diabetes or hypertension alone.⁸⁷

Remote Consultation

Three studies (one trial¹⁰² [$n=113$], one prospective cohort study,⁹⁴ [$n=76$] and one pre-post study¹⁰⁰ [$n=113$]) assessed remote consultations for multiple health conditions. One trial randomly assigned primary care providers to an electronic consultation program or to usual care in which patients traveled to a specialist for an in-person consultation. The eConsult system allowed primary care clinicians to ask patient-specific clinical questions to remote specialists, who are asked to review patient data and respond within 7 days. The study, which aimed to reduce referrals for in person consultations, found that such referrals declined from baseline in

both arms and there was no significant difference in referrals for in person visits for the specialties covered by eConsult (adjusted RR 0.93; 95% CI, 0.85 to 1.03) or overall referrals (adjusted RR 0.94; 95% CI, 0.85 to 1.04) during the followup period.¹⁰² One study of patients with hypertension, hyperlipidemia, or diabetes found improved medication adherence (OR 1.76, 95% CI 1.34 to 2.31) but no difference in healthcare quality of life (OR 1.07, 95% CI 0.90 to 1.27) after implementation of telehealth remote consultations.¹⁰⁰ Another study investigated remote consultations in an integrated care model for older patients (≥ 55 years) recovering from a respiratory or cardiovascular event.⁹⁴ Patients receiving integrated care experienced fewer hospital visits (mean 1.0 [SD=1.1] vs. 2.3 [SD=3.1], p-value not reported), but there were no differences in between the groups in physical function.

Dermatology

One RCT (n=261)¹¹⁹ compared clinical course after store-and-forward teledermatology consultation or conventional consultation. The study did not find significant differences in the proportion of patients with favorable clinical courses (telehealth 36% vs. in-person 38%, p=0.78 at first clinic visit; 82% vs. 83%, p=0.88 at 9 months). A second RCT (n=391)⁹² compared costs between teledermatology referral and the conventional referral process for patients with ambulatory skin conditions. The study did not find significant differences in cost per participant from a U.S. Department of Veteran's Affairs perspective (\$30, 95% CI -\$79 to \$20), and identified very small differences in cost per participant from a societal perspective (-\$82, 95% CI -\$12 to -\$152).

Endoscopy

One study (n=90) compared costs of tele-endoscopy on an island in United Kingdom versus in-person endoscopy on the mainland.¹¹⁸ The study found that when 27 or more patients per year attended tele-endoscopy, the cost was lower with telehealth (£353) than for the mainland clinic (£381). They also estimated that it would take 9.5 years for these savings to pay back the set up costs. The study did not assess the impact of locally available tele-endoscopy on waiting time or missed appointments.

Blood Pressure Control

A cluster-randomized trial (n=1299) compared an integrated mobile health program with usual rural community provider care to control blood pressure in adults with history of stroke in China.¹²¹ Patients in the integrated mobile health program had significantly greater reductions in systolic blood pressure (adjusted mean difference -2.8 mmHg (95% CI -4.8 to -0.9), diastolic blood pressure (adjusted mean difference -2.2 mmHg (95% CI -3.2 to -1.3), and stroke recurrence (RR 0.46 [95% CI 0.32 to 0.66]q).

Palliative Care

One small prospective cohort study (n=21) compared telehealth videoconference assisted palliative care with traditional nurse visit palliative care.⁹⁸ The study did not find significant differences in symptoms or function, but did report that 30 days prior to death, participants receiving telehealth had fewer per capita nursing visits (5.46 vs. 9.32, effect size=0.7), general practitioner visits (0.13 per capita vs. 3.88, effect size=1.34), and hospital admissions (0.02 per capita vs. 0.2, effect size=0.65).

Ultrasound During Pregnancy

One evaluation of a statewide teleprogram (Arkansas, United States) to improve care for patients with high-risk pregnancies tracked whether patients received ultrasounds in the 3 years prior to the program (2001-2003) and the 3 years after its inception (2004-2007). The total number and the percentage of patients receiving comprehensive ultrasounds increased for all patients (9.6% to 11.3%, $p < 0.0001$) and for high-risk patients (16.9% vs. 19.9%, $p < 0.0001$)¹⁰⁴

Table 3. Key Question 2: summary of outpatient findings*

Clinical Topic Number of Studies	Provider Specialty	Consultant Specialty	Patient Outcomes*	Provider Outcomes*	Payer Outcomes*
Telehealth-based collaborative care for depression Studies 3	Primary care clinicians; (physicians, psychologists, social workers, midwives, and nurses; nurse and pharmacist	Depression care managers (nurse, pharmacist, psychologists, psychiatrists); psychiatrists	Patient outcomes M Response (varies by timepoint 6, 12, 18 months) ^{93, 95, 97, 109-111, 122} M Remission (varies by timepoint 6, 12, 18 months) ^{93, 95, 97, 109-111, 122} ~ Depression-free days ^{93, 95, 97, 109-111, 122} ~ Treatment adherence at 3 months, + at 6 months ^{93, 95, 97, 109-111, 122} + Medication adherence 6 & 12 months ^{93, 95, 97, 109-111, 122} ~ Depression symptom score (BDI) at 3 and 6 months ^{93, 95, 97, 109-111, 122} M Quality of Life (varies by timepoint 6, 12, 18 months and scale, Quality of Well-being Scale and SF-36 Mental) ^{93, 95, 97, 109-111, 122}	None reported	~ Utilization: overall, + outpatient only ^{95, 93, 110, 111} ~ Total costs ^{95, 110} - Adjusted total cost ¹⁰⁹ + Incremental cost/depression-free d ¹⁰⁹ - Depression-related primary and mental health costs ^{95, 110}
Telehealth-based collaborative care for PTSD Studies 1	Primary care physicians, psychiatric advanced practice nurses, social workers	Nurse care manager, clinical pharmacist, psychologist, psychiatrist	+ Prescribed any medication for PTSD ⁹⁶ ~ Adherence to medication regimen >80% ⁹⁶ ~ Any psychiatric encounter at 12 months ⁹⁶ + Mean number of CPT sessions at 12 months ⁹⁶ M Quality of Life (varies by scale, Quality of Well-being Scale and SF-36 Mental and Physical scales) ⁹⁶	None reported	- Total outpatient costs ^{96, 108}
Tele-mental health treatment for AD/HD Studies 1	Primary care physicians and caregivers	Child psychiatrists	~ Teacher ratings (VADRS) ¹⁰⁷ ~ Caregiver ratings (VADRS-Role Performance and CIS-P Scale) ¹⁰⁷	None reported	None reported
Telehealth-based diabetes care Studies 4	Primary care clinicians, nurses	Nurse, dietician, and a diabetes specialist, diabetes specialist	+ A1c ^{90, 101, 103, 106} and self-monitoring of blood glucose ¹⁰⁶ + Diabetes Self-Management Education score ^{90, 106} ~ Total cholesterol ^{90, 106} and systolic blood pressure ¹⁰³	None reported	None reported

Clinical Topic Number of Studies	Provider Specialty	Consultant Specialty	Patient Outcomes*	Provider Outcomes*	Payer Outcomes*
Remote diabetic neuropathy screening Studies 1	Primary care clinicians	Ophthalmologist	+ Screening rate ⁹¹	+ Screening referral or reminder ⁹¹	None reported
Telehealth hepatitis C virus consultation Studies 1	Primary care physicians	Hepatologist	~ Sustained virologic response ¹¹² + Completion of therapy ¹¹² ~ Mean weeks of therapy ¹¹² + Mean face-to-face visits ¹¹² + Anemia and withdrawal due to adverse events ¹¹²	None reported	None reported
Tele-rheumatology Studies 2	Rheumatology graduate medical education trainee; physical therapists	Rheumatologist	+ Distance to visit ¹²⁰ ~ Quality of life, disease-related function and visit satisfaction scores ¹¹⁶	None reported	+ Cost of visit ¹²⁰
Teleoncology Studies 2	Local healthcare professionals; Local oncology care team	Oncologists; respiratory physicians, thoracic surgeons, radiation oncologists, medical oncologists and a diagnostic radiologist	~ Receipt of radiotherapy ¹¹⁴	None reported	+ Total costs ¹¹⁷
Tele-echocardiography Studies 1	General practitioners	Cardiologists	+ Total process time ⁸⁸ + Time to specialist consultation ⁸⁸	None reported	None reported
Tele-hemodialysis Studies 1	Nurse, nutritionist, and general practitioners	Nephrologist	+ Amount of urea removed ¹¹³ ~ Other clinical measures ¹¹³	None reported	None reported
Telehealth fracture clinic Studies 1	Registrar (details NR)	Orthopedic surgeon	None reported	None reported	+ Cost compared to transfer or sending a specialist ¹⁰⁵
Telehealth dementia care Studies 1	Local medical staff	Dementia specialists	~ Yearly change in MMSE score among all patients ⁹⁹ + Yearly change in MMSE score among patients with baseline MMSE 15-30 ⁹⁹	None reported	None reported

Clinical Topic Number of Studies	Provider Specialty	Consultant Specialty	Patient Outcomes*	Provider Outcomes*	Payer Outcomes*
Telehealth-based ulcer care Studies 1	Primary care	Secondary care	+ Time from referral to appointment ¹¹⁵ + Leg ulcer healing time ¹¹⁵	None reported	~ Cost per patient ¹¹⁵
Telepharmacy Studies 2	Pharmacy technicians	Pharmacists	~ A1c and systolic blood pressure in patients with either diabetes or hypertension ⁸⁷ + A1c and systolic blood pressure in patients with both diabetes and hypertension ⁸⁷	+ Guideline adherence ⁸⁹	None reported
Remote consultation 3	Family physicians, nurses	Specialists (type NR), multidisciplinary team	+ hospital visits ⁹⁴ + patient function + medication adherence ¹⁰⁰ + health-related quality of life ¹⁰⁰	~ Overall referrals or to eConsult specialties ¹⁰²	None reported
Dermatology 2	Primary care	Dermatology	~ clinical course ¹¹⁹	None reported	M cost per participant ⁹²
Endoscopy 1	Surgeon on anesthetist	ENT	None reported	None reported	~ costs ¹¹⁸
Blood pressure 1	Primary care provider	Physician, specialty not reported	+ blood pressure ¹²¹ + stroke recurrence ¹²¹	None reported	None reported
Palliative care 1	Nurse, general practitioner	Palliative care specialist	~ patient symptoms and function + nurse or general practitioner visits, hospital admissions	None reported	None reported
Ultrasound for pregnancy 1	Community obstetrics providers	Maternal-fetal medicine and radiology	+ increase in comprehensive ultrasound for all and high-risk pregnancies ¹⁰⁴	None reported	None reported

*Symbol meaning: + = Improved Outcome with telehealth; ~ = Similar outcome with telehealth; - = Worse outcome with telehealth, M = Outcomes were not consistent across studies

Abbreviations: CIS-P = Columbia impairment scale-parent version; CRP = c-reactive protein; ENT = ear, nose, throat specialist; ICER = incremental cost-effectiveness ratio; MMSE = mini-mental state examination; NR = not reported; PTSD = post-traumatic stress disorder; QALY = quality adjusted life-years; RN = registered nurse; SF-12 = 12-item short form survey; SF-36 = 36-item short form survey; VADRS = Vanderbilt Attention-Deficit/Hyperactivity Disorder Rating Scale.

Emergency Department/Emergency Medical Services

Key Points

- Telehealth consultations supporting emergency assessment and care of stroke/ST-elevation myocardial infarction (STEMI) and chest pain at a rural hospital:
 - Result in similar rates of mortality when patients are treated locally as opposed to transferred (5 studies; Low SOE)
 - May result in similar time to treatment when patients are treated locally as opposed to transferred (8 studies; Low SOE)
- Telehealth consultations by specialists for rural ED critical care and trauma patients may result in similar appropriate or inappropriate transfers (5 studies; Low SOE)

Summary of Findings

We identified and included 28 studies of provider-to-provider telehealth use in rural areas for the provision of emergency care, either by emergency medical services (EMS) or EDs.^{67, 124-150} Fifteen of these studies evaluated telehealth for the assessment and management of conditions that are time sensitive and require decision about the course and best location for treatment. These include ten studies of stroke,^{67, 127, 136, 139, 140, 142, 143, 145, 148, 149} three studies of STEMI,¹²⁴⁻¹²⁶ and two studies of chest pain/potential myocardial infarction.^{133, 136} Five of the included studies were of the use of teleconsultations for a range of conditions seen in EDs^{132, 138, 140, 144, 150} and five additional studies were of video consultations for critical care,^{128, 129, 131} trauma,¹³⁵ or hand trauma.¹⁴⁶ The three remaining studies also employed video consultations for sepsis or septic shock,¹³⁷ suicidal ideation or attempt¹³⁰ and the evaluation of sexual abuse.¹³⁴

Two of these studies were RCTs^{127, 149}, ten were prospective cohort studies,^{124-126, 132, 133, 136, 137, 142, 143, 145} eight were retrospective cohort studies^{67, 128, 129, 131, 134, 135, 147, 150}, and eight were studies that compared outcomes before and after telehealth initiation.^{130, 138-141, 144, 146, 148} Eighteen of these studies were conducted in the United States^{67, 127-131, 133-138, 141, 144-147, 150} two were conducted in Italy,^{125, 126} two in Australia^{139, 148} and one each in Canada,¹³² Spain,¹⁴² Finland,¹⁴³ Turkey,¹²⁴ Japan,¹⁴⁰ and Germany.¹⁴⁹ Risk of bias was rated as high for one study,¹³⁶ medium for 26 studies,^{67, 124-126, 128-135, 137-150} and low for one trial¹²⁷ (**Appendix E**).

Key results are presented in **Table 4**, and additional study details and results can be found in **Appendix Tables D-6 and D-7**. We grouped studies by similar clinical issues for our qualitative synthesis and SOE assessment.

In studies of telehealth for stroke, STEMI and chest pain combined, five studies reported no significant differences in mortality.^{67, 127, 139, 143, 148} One study of telestroke versus usual care reported no differences in in-hospital mortality for rural or super-rural (areas in the bottom quartile of rural areas as defined by the Centers for Medicare & Medicaid Services Zip code file) patients, (adjusted differences: -0.6, 95% CI, -2.5 to 1.4; 0.4, 95% CI, -2.7 to 3.4), or 30-day all-cause mortality in rural patients (adjusted difference 0.0, 95% CI, -2.6 to 2.7). The study reported a small difference in 30-day all-cause mortality in super-rural patients (adjusted difference -4.5, 95% CI, -8.0 to -1.0).⁶⁷ Similarly, two other studies of telestroke reported small non-significant differences in mortality rates: 13% vs. 10%, $p=0.6$ ¹³⁹ and 10% vs. 7%, $p=0.58$.¹⁴⁸ One study of telehealth consultations between hub and spoke sites versus telephone consultation for stroke patients did not find a difference in 90-day mortality (OR 0.74; 95% CI, 0.35 to 1.55).¹²⁷ The final study compared telestroke with neurologic emergency room care, and did not report a

difference in 3-month mortality (11.5% vs. 10.2%, $p=0.662$).¹⁴³ When combined, these five studies provide a low SOE supporting no difference in mortality with telehealth.

In eight studies time to treatment was either the same or better, with no study reporting that telehealth resulted in treatment delays.^{125-127, 133, 136, 139, 142, 148} No studies reported an increase in harms, such as increases in intracranial bleeds. Only one study reported a payer outcome, finding that mean total expenditures for telestroke patients were higher than with usual care in both rural (\$13,868 vs. \$9,721, $p<0.001$) and super-rural areas (\$14,596 vs. \$10,020, $p<0.001$), likely due to the increase in use of tissue plasminogen activator (tPA).⁶⁷ One study also assessed a provider behavioral outcome, finding that correct thrombolysis decisions were more frequent with telestroke consults than when advice was provided by phone (96% vs. 83%).¹²⁷ While there a number of studies, time to treatment is measured differently and with varying degrees of validity and reliability and as a group these studies provide low SOE that there is either no difference or some improvement in time to treatment with telehealth. telehealth

Three studies of telehealth emergency care consultations for the other clinical indications reported that telehealth reduces unnecessary transfers and increases appropriate transfers,^{131, 132, 141} though two studies did not find differences.^{135, 144} Another key outcome for the use of telehealth was length of stay in the ED, as consultations should help speed disposition decisions. For this outcome even though there are four studies, the evidence is insufficient to support a conclusion as the findings were contradictory with one study reporting shorter length of stay,¹³⁵ one longer length of stay,¹³⁰ and two finding no difference or mixed results.^{138, 147}

Table 4. Key Question 2: summary of emergency care findings

Clinical Topic Number of Studies	Modality	Provider Specialty Consultant Specialty	Patient Outcomes*	Provider Outcomes*	Payer Outcomes*
Stroke 10	TeleED Video- conference	Providers ED clinicians Consultants Neurologist, Stroke expert, ED Physicians, ED Nurses	<p>~ In-hospital, 30-day and 90-day mortality^{67, 127, 139, 143, 148}</p> <p>+ 30-day all-cause mortality, super rural patients⁶⁷</p> <p>+ Discharge to home or rehab¹⁴⁸</p> <p>~ Symptom onset to tPA time^{67, 127, 136, 139, 140, 142, 143, 145, 148}</p> <p>+ tPA within 3 h of symptom onset^{67, 127, 136, 139, 140, 142, 143, 145}</p> <p>~ 90-day modified Rankin Scale^{127, 143}</p> <p>~ post-tPA intracranial hemorrhage^{127, 143, 148}</p> <p>M tPA use^{67, 127, 136, 139, 140, 142, 143, 145}</p> <p>~ Length of stay, overall and rural patients^{67, 127, 136, 139, 140, 142, 143, 145}</p> <p>+ Length of stay, super rural patients⁶⁷</p> <p>+ Appropriate transfer to high-volume center or higher level of care^{140, 149}</p>	+ Correctness of decision-making/accurate triage ^{127, 149}	- Total medical expenditures per event ⁶⁷
Heart attack/ STEMI 2	TeleECG	Providers Paramedics, EMS Consultants Cardiologists	+ On-time and time-to treatment ^{125, 126}	None reported	None reported
Heart attack/ STEMI 1	Text triage	Providers ED clinicians Consultants Cardiologists	<p>+ Time to hospital arrival¹²⁴</p> <p>M Arrival -to-balloon time¹²⁴</p> <p>~ False-positive STEMI¹²⁴</p>	None reported	None reported
Chest pain/ MI 2	TeleED video- conference	Providers ED clinicians Consultants: ED Physicians ED Nurses	<p>M Time from ED arrival to ECG^{133, 136}</p> <p>~ Time from ED arrival to fibrinolytic^{133, 136}</p> <p>+ Likelihood of receiving fibrinolytic when eligible¹³³</p>	None reported	None reported

Clinical Topic Number of Studies	Modality	Provider Specialty Consultant Specialty	Patient Outcomes*	Provider Outcomes*	Payer Outcomes*
Multiple 1	TeleED robot	Providers Nurses, Nurse Practitioners Consultants Pediatric intensivist	+ Transfer ¹³² + Length of stay (robot/transfer vs. no robot/no transfer) ¹³²	None reported	None reported
Multiple 4	TeleED video- conference	Providers Nurse Practitioners, ED clinicians Consultants ED physicians, ED nurses	~ Patient deaths in ER ¹⁴⁴ M Transfer to another facility ^{141,144} + Admit to provider time ¹³⁸ - ED length of stay, nontransferred patients ¹³⁸ + ED length of stay, transferred patients ¹³⁸ - Likelihood of discharge from ED ¹⁴⁴ - Rural hospital admission ¹⁴⁴ - Discharged against medical advice from ED ¹⁴⁴	None reported	~ Total ED patient volume ¹⁴⁴ + Lower ED costs and operating expenses ¹⁵⁰
Critical care 3	TeleED video- conference	Providers ED clinicians Consultants Pediatric critical care physician	+ Transfer avoided ¹³¹ + Transfer to lower level of care ¹³¹ + Parent/guardian satisfaction ¹²⁹ + Accuracy of clinical picture prior to arrival ¹³¹	+ Physician-related ED medication errors ¹²⁸ + Quality of care scores ¹²⁹ + Referring physician satisfaction ¹²⁹	None reported
Trauma 1	TeleED video- conference	Providers ED clinicians Consultants ED Physicians, ED nurses	~ Mortality + Chest tube, intubation ¹³⁵ + ED length of stay ¹³⁵ M Increased diagnostic imaging ¹³⁵ + Time to arrival at final hospital ¹³⁵ ~ Transfer ¹³⁵	None reported	None reported
Hand trauma 1	TeleED video- conference	Providers ED clinicians Consultants Hand surgeon	~ Length of stay at first hospital ¹⁴⁶ ~ Type of transfer, air or ground ¹⁴⁶ + Admitted from ED ¹⁴⁶	None reported	~ Transport cost

Clinical Topic Number of Studies	Modality	Provider Specialty Consultant Specialty	Patient Outcomes*	Provider Outcomes*	Payer Outcomes*
Sepsis/ septic shock 1	TeleED video-conference	Providers ED clinicians Consultants Physician (specialty varied)	None reported	+ Adherence to sepsis treatment bundle ¹³⁷	None reported
Behavioral health; suicidal ideation or attempt 2	TeleED video-conference	Providers ED clinicians Consultants Psychiatric providers, mental health providers	+ ED wait time ¹³⁰ - ED length of stay ¹³⁰ † + Hospital admission ¹⁴⁷ + Involuntary hold placement ¹⁴⁷ ~ 30-day readmission ¹⁴⁷	None reported	None reported
Sexual abuse 1	TeleED video-conference	Providers Rural examiners Consultants Advance practice nurse practitioners	None reported	+ Quality, completeness, accuracy of abuse examination forms ¹³⁴	None reported

*Symbol meaning: + = Improved Outcome with telehealth; ~ = Similar outcome with telehealth; - = Worse outcome with telehealth, M = Outcomes were not consistent across studies

†Difference in ED length of stay exceeds the time for the consult and was not associated with reductions in admissions.

Abbreviations: ECG = electrocardiogram; ED = emergency department; EMS = emergency medical services; MI = myocardial infarction; STEMI = ST-elevated myocardial infarction; TeleED = telehealth emergency department; tPA = tissue plasminogen activator.

Education and Mentoring

Key Points

- Clinical outcomes: Extension for Community Healthcare Outcomes (ECHO) programs are associated with better or similar patient outcomes. (2 studies; Low SOE)
 - Reduction of A1c in patients of trainees after ECHO compared to before participation (1 study)
 - Hepatitis C viral response and serious adverse events rates at “spoke” site with ECHO participation were similar to those at an academic medical center (1 study)
- Provider behavior: ECHO and non-ECHO video training programs result in desired changes in provider behavior (e.g., increased appropriate prescribing practices, screening, and patient counseling). (8 studies; Low SOE)
- Provider knowledge, efficacy, perceptions: ECHO and non-ECHO video training are associated with increased confidence, efficacy, and scores on knowledge tests. (13 studies; Low SOE)

Summary of Findings

Provider-to-provider telehealth for education and mentoring uses communications technology to transmit knowledge, develop capacity and share expertise. Education and mentoring programs usually contain didactic elements and may or may not also include case reviews or supervised or mentored patient care. Telehealth for this purpose may use different technologies, including video, online learning platforms, or simulation devices and complex monitoring. While most uses of telehealth for training are in real time, some asynchronous applications exist. Multiple terms are used to describe use of telehealth technologies for purposes of education and mentoring, including tele-mentoring, distance education, virtual grand rounds, etc.

Twenty-three studies evaluated the use of telehealth for provider-to-provider education and mentoring,¹⁵¹⁻¹⁷³ including three RCTs^{153, 157, 169} and 20 observational studies. Observational study designs included pre-post,^{152, 154, 158-163, 166, 167, 168, 170, 171, 173} before-after,^{155, 165} prospective cohort studies,^{151, 164} and retrospective cohort studies.^{156, 172} All studies were conducted across multiple clinical sites or healthcare organizations. Sixteen studies were performed in the United States,^{151, 152, 154, 156, 158, 159, 161, 162, 164, 165, 167-169, 171-173} four in Australia,^{153, 155, 160, 163} two in Canada,^{166, 170} and one in Vietnam.¹⁵⁷ The duration of the education or mentoring intervention made available to rural care providers ranged from 3 months to 4.5 years, with one study that did not specify the length.¹⁶⁹ All studies evaluated patient clinical or provider outcomes. One study was rated low risk of bias,¹⁶⁶ 11 were rated medium risk of bias,^{151, 153, 156, 157, 164, 165, 168-172} and 11 were rated high risk of bias^{152, 154, 155, 158-163, 167, 173} (**Appendix E**).

Clinical topics addressed through distance education and mentoring included antibiotic therapy,¹⁶⁸ childhood obesity,¹⁶⁵ dermatology,¹⁵⁹ diabetes,¹⁶⁷ liver disease,^{151, 161, 172} mental health,^{152, 153, 155, 156, 158, 162, 163, 166, 169} multiple conditions in primary care,¹⁵⁷ pediatric burns,¹⁶⁰ perioperative care,¹⁶⁴ low vision,¹⁷¹ dementia,¹⁷³ and COVID-19 in long term care.¹⁷⁰ Telehealth modalities used in the studies of remote education and mentoring that we identified include:

- **ECHO Videoconference:** ECHO is a technology-enabled collaborative learning and capacity building model launched in 2003 by the University of New Mexico Health Sciences Center (<https://hsc.unm.edu/echo/>). ECHO is designed to amplify the capacity

to provide best practice care for people in rural and underserved communities. ECHO programs consist of a “hub” site, which is typically housed within an academic medical center, and “spokes” consisting of smaller medical offices, community health centers, nursing homes or other health-related facilities or organizations in rural or underserved areas. ECHO sessions are facilitated by specialist providers and others with clinical subject matter expertise from the “hub” site, and attended by a wide variety of clinicians at “spoke” sites. ECHO sessions are typically comprised of a combination of didactic education sessions and review of clinical cases. The ECHO model, originally designed for improving the care of patients with hepatitis C, had expanded to include many more conditions, and has spread across the United States and the globe through the University of New Mexico ECHO replication program.

- Non-ECHO Videoconference: Videoconference-based education and mentoring that is not reported to be affiliated with or organized by an ECHO replication site and may use a different educational model.
- Online Education Courses: Online education and mentoring includes asynchronous online education courses, or a mix of asynchronous online courses and synchronous interaction with remote mentors via videoconferencing or telephone.
- Short Messaging Service (SMS): Education and mentoring provided through text messaging on a mobile communication device.

Table 5 includes a summary of the clinical and intermediate results from included studies of remote education and mentoring (additional details available in **Appendix Tables D-8 and D-9**). The table is organized by training modality and then by clinical topic. It summarizes the number of articles associated with each by clinical topic, provides information on the type(s) of providers who received remote education and mentoring, and summarizes key clinical and provider outcomes.

Two studies reported only patient outcomes. A prospective cohort study compared sustained viral response and adverse events between patients seen at an ECHO spoke site and those seen at an academic medical center hub site after implementation of a hepatitis C ECHO program. This study did not report differences across the hub and spoke sites (0.70; 95% CI, -9.2 to 10).¹⁵¹ One study reported on patient care, specifically the impact of ECHO programs on the use of direct-viral treatment for hepatitis C, and found that the program was not associated with the odds of treatment among rural vs. urban patients (adjusted OR 1.01; 95% CI 0.99 to 1.02, $p=0.49$)¹⁷²

Three studies reported both patient and provider outcomes, all evaluated videoconference-based delivery of remote education and mentoring, but reported mixed results.^{165, 167, 168} Studies found improved provider self-efficacy and knowledge and improved A1c in diabetic patients (10.2 [SD = 1.4] vs. 8.3 [SD = 0.97], $p < 0.001$),¹⁶⁷ improved prescribing practices but no reduction in mean length of stay or in-hospital mortality in antibiotic therapy,¹⁶⁸ and improved documentation (1; 95% CI 0.9 to 1.2, $p < 0.001$) and counseling behaviors (0.8; 95% CI 0.2 to 1.4, $p=0.01$) but no change in child nutrition or physical activity levels (0.3; 95% CI -0.2 to 0.8) in childhood obesity.¹⁶⁵

Eighteen studies reported a range of provider only outcomes including provider knowledge, efficacy, perception and satisfaction. Education and mentoring modalities utilized in studies reporting only intermediate outcomes include ECHO video-conference (9), Non-ECHO Videoconference (4), online education (4), and SMS (1).^{152-164, 166, 169, 170, 171, 173} Eleven studies reported improvement in provider knowledge, self-efficacy and perception across a range of

clinical topics.^{153-155, 158, 160, 161, 163, 166, 170, 171, 173} A single study of SMS-provided education reported no change in provider knowledge.¹⁵⁷ A single study did not report differences in scores of perioperative skill evaluations among video-conference and in-person participants.¹⁶⁴ Four studies reported improved provider behaviors related to patient screening rates^{152, 161} and prescribing.^{156, 158} A single study of provider capacity to perform dermatology procedures found an increase in the number of providers of biopsies, excisions, shave biopsies, electrocautery, and no change in capacity to provide liquid nitrogen at the spoke site.¹⁵⁹ Four studies reported increased provider satisfaction with remote education and mentoring.^{155, 157, 166, 169} One study¹⁶⁹ reported negative perception of balance between instruction and practice in video-conference-based education and monitoring.

Table 5. Key Question 2: Summary of remote education and mentoring findings

Modality	Clinical Topic Number of Studies	Provider Specialty Consultant/Mentor Specialty	Provider Outcomes*	Patient Outcomes*
ECHO video-conference	Antibiotic therapy 1	Providers Multidisciplinary Consultants Infection disease	+ Antibiotic prescribing ¹⁶⁸	~ In-hospital mortality ¹⁶⁸ ~ Mean length of stay ¹⁶⁸
ECHO video-conference	COVID-19 in long-term care 1	Providers Multidisciplinary Consultants/instructors Long-term care experts	+ Self-efficacy and satisfaction ¹⁷⁰	None
ECHO video-conference	Dementia 1	Providers Multidisciplinary Consultants/instructors Neurology and dementia experts	+ Comfort with assessment and treatment ¹⁷³	None
ECHO video-conference	Diabetes 2	Providers PCPs Community health workers Consultants Multidisciplinary diabetes team; diabetes specialist	+ Self-efficacy in patient coaching/education, and identification of psychosocial treatment barriers ¹⁵⁴	+ A1c ¹⁶⁷
ECHO video-conference	Liver disease 3	Providers Multidisciplinary physicians Consultants Multidisciplinary	+ Hepatitis C Virus awareness, knowledge, abilities and intention to recommend screening for at-risk patients ¹⁶¹	~ Sustained viral response ¹⁵¹ ~ Serious adverse events ¹⁵¹ ~ Access to direct acting antiviral treatment in rural areas ¹⁷¹

Modality	Clinical Topic Number of Studies	Provider Specialty Consultant/Mentor Specialty	Provider Outcomes*	Patient Outcomes*
ECHO video-conference	Mental health 5	Providers Multidisciplinary Consultants Multidisciplinary teams, including addiction and psychiatry specialists	<ul style="list-style-type: none"> + Opioid use disorder diagnosis/ prescribing¹⁵⁶ + Reduction in patients prescribed ≥ 3 psychotropic medications¹⁵⁸ + Provider knowledge and self-efficacy¹⁶⁶ + General development and Autism-specific screening¹⁵² + Pediatric behavioral health knowledge and patient management¹⁵⁸ + Satisfaction with sessions^{158, 166, 169} 	None Reported
Non-ECHO video-conference	Childhood obesity 1	Providers Clinician champions, nurses, medical assistants Consultants/Mentors Healthy Eating Active Living Telehealth Community of Practice faculty and staff	<ul style="list-style-type: none"> + Documentation and counseling¹⁶⁵ ~ Family centered care¹⁶⁵ 	~ Child nutrition and physical activity ¹⁶⁵
Non-ECHO video-conference	Dermatology 1	Providers PCPs, imaging technicians Consultants/mentors Dermatologist	<ul style="list-style-type: none"> + Knowledge of dermatology procedures, ability to provide punch or shave biopsies, excisions, electrocautery¹⁵⁹ ~ Ability to provide liquid nitrogen¹⁵⁹ 	None reported
Non-ECHO video-conference	Mental health 1	Providers Dementia behavior management advisory service Consultants/mentors Mental health services for older people, old age psychiatrist	<ul style="list-style-type: none"> + Confidence in managing behavioral and psychological symptoms of dementia¹⁵⁵ + Satisfaction with educational program¹⁵⁵ 	None reported
Non-ECHO video-conference	Pediatric burns 1	Providers Multidisciplinary Consultants/mentors Burn consultants and nurse	<ul style="list-style-type: none"> + Knowledge of burn prevention, first aid, airway and inhalation injury, and circulation and fluid resuscitation¹⁶⁰ M Chemical and electrical burns, burn wound, pain and itch management¹⁶⁰ 	None reported
Non-ECHO video-conference	Perioperative care 1	Providers Registered nurses Consultants/mentors Trained preceptor nurses	<ul style="list-style-type: none"> ~ Perioperative training scores¹⁶⁴ ~ Rating program as a success¹⁶⁴ 	None reported

Modality	Clinical Topic Number of Studies	Provider Specialty Consultant/Mentor Specialty	Provider Outcomes*	Patient Outcomes*
Online education course	Mental health 3	Providers Multidisciplinary Consultants/mentors Clinical psychologist; mental health clinicians	+ Role adequacy, legitimacy and support ¹⁶² M Role motivation, work satisfaction, and task-specific self-esteem ¹⁶² + Education completion ¹⁵³ + Knowledge, skills, confidence, and utilization of CBT ¹⁵³ + Computer and internet-related skills ¹⁶³ + Knowledge about the roles of mental health service, confidence in responding to mental health problems ¹⁶³	None reported
Online education course	Low vision screening 1	Providers Occupational therapists Consultants/mentors Low vision specialists	+ Knowledge of low vision screening and treatment ¹⁷¹	None reported
Short messaging service	Multiple conditions common in primary care 1	Providers Physician Assistants Consultants/mentors Automated text messages created by specialists	~ Medical knowledge ¹⁵⁷ + Satisfaction with intervention ¹⁵⁷	None reported

*Symbol meaning: **+** = Improved Outcome with telehealth; **~** = Similar outcome with telehealth; **-** = Worse outcome with telehealth; **M** = Outcomes were not consistent across studies

Abbreviations: A1c = glycated hemoglobin; CBT = cognitive behavioral therapy; ECHO = Extension for Community Healthcare Outcomes; PCP = primary care provider.

Key Question 3. What strategies are effective and what are the barriers and facilitators to implementation and sustainability of provider-to-provider telehealth in rural areas?

Key Points

- The majority of barriers and facilitators are similar across provider-to-provider telehealth programs implemented in different settings and for different purposes. These are related to available resources and access to knowledge & information.
- While most barriers and facilitators are likely not unique to rural provider-to-provider telehealth, some may be specific and study authors suggested specific strategies:
 - Consulting providers who are often located in urban areas need to understand the rural context and what resources are available. Strategy: rural rotations or periodic in person collaboration may support success.
 - Successful implementation and sustainment require a long-term commitment and resources on a scale that may not be feasible for individual rural organizations. Strategy: statewide or regional initiatives with government or philanthropic start up support.
 - Provider-to-provider telehealth systems may be used for frequent events or serve as a resource for rare events in rural healthcare and the technology and support need to be tailored to frequency of use. Strategy: schedule periodic testing of systems used for rare events.

Summary of Findings


We identified 67 studies (in 71 publications)^{52, 62, 65, 79, 144, 174-239} that addressed implementation of provider-to-provider telehealth in rural areas. We used these studies to collectively assess the barriers and facilitators that have been encountered in implementing provider-to-provider telehealth for rural populations. As the inclusion criteria for this Key Question were broad, the studies include not only a wide range of topics, but also diverse methods. Many of these studies included for this Key Question are program evaluations that combine data from several sources, such as site visits, observations, surveys and interviews. These evaluations often describe the telehealth program components or experience in more detail than is often included in effectiveness studies. Others were qualitative research studies that analyze interviews, focus groups, or documents and then categorize or catalog specific barriers and facilitators to initial implementation, ongoing operations, longer term sustainment, or spread of the use of telehealth. In some cases, an effectiveness study (included in Key Question 2) collected data and reported on implementation, in which case these studies are included in both Key Questions. Few studies actually compared programs, models or implementation approaches; instead they report on a single experience. Additional details about the included studies can be found in **Appendix Table B-3** and **Appendix Table D-10**.

As outlined in the Methods Section above and described in more detail in **Appendix A**, Methods, we used constructs from a framework called CFIR (Consolidated Framework for Implementation Research) to standardize our description of the facilitators and barriers to implementation and sustainability cited in the included studies. Studies often use different terms

and narratives to describe their experience and using a framework with defined constructs allowed us to group similar concepts and describe patterns.

Figure 2 illustrates how the 219 facilitators and 192 barriers we identified in the included studies map to 19 of 39 possible CFIR constructs. The number next to each construct is the number of times it occurred in our data and the width of the line represents the relative frequency. Short definitions of each construct are included as notes to this figure and the tables in this section. More comprehensive descriptions are available in CFIR articles and tools.^{60, 240} This diagram allows us to represent the important fact that most constructs can be either barriers or facilitators, depending on context and their presence or absence. In our synthesis this is the case for both of the most frequently cited constructs. *Available Resources* is a broad concept including the wide range of investments an organization dedicates to implementing or sustaining an innovation or program. This was cited sixty times, but as represented in the figure, almost equally split between mention as a facilitator and as a barrier. *Access to Knowledge & Information*, was cited 57 times, but most frequently as a facilitator. Additional details regarding facilitators and barriers presented by setting (inpatient, outpatient, EMS/ED, education/mentoring) are found in **Appendix B**.

Figure 2. CFIR constructs as facilitators and barriers*



*Construct definitions: **Access to Knowledge & Information:** Access to digestible information and knowledge about the innovation and how to incorporate it into work tasks; **Formally Appointed Internal Implementation Leaders:** Individuals from within the organization who have been formally appointed with responsibility for implementing an innovation as coordinator, project manager, team leader, or other similar role; **Planning:** Degree to which a scheme or method of behavior and tasks for

implementing an innovation are developed in advance, and the quality of those schemes or methods; **Leadership Engagement:** Commitment, involvement, and accountability of leaders and managers with the implementation of the innovation; **Engaging:** Attracting and involving appropriate individuals in the implementation and use of the innovation through a combined strategy of social marketing, education, role modeling, training, and other similar activities; **Available Resources:** Level of resources organizational dedicated for implementation and on-going operations including physical space and time; **Networks & Communications:** Nature and quality of webs of social networks, and the nature and quality of formal and informal communications within an organization; **Reflecting & Evaluating:** Quantitative and qualitative feedback about the progress and quality of implementation accompanied with regular personal and team debriefing about progress and experience; **Cost:** Costs of the innovation and costs associated with implementing the innovation including investment, supply, and opportunity costs; **External Policy & Incentives:** External strategies to spread innovations including policy and regulations (governmental or other central entity), external mandates, recommendations and guidelines, pay-for-performance, collaboratives, and public or benchmark reporting; **Relative Priority:** Individuals' shared perception of the importance of the implementation within the organization; **Implementation Climate:** Absorptive capacity for change, shared receptivity of involved individuals to an innovation, and the extent to which use of that innovation will be rewarded, supported, and expected within their organization; **Readiness for Implementation:** Tangible and immediate indicators of organizational commitment to its decision to implement an innovation; **Adaptability:** Degree to which an innovation can be adapted, tailored, refined, or reinvented to meet local needs; **Patient Needs & Resources:** Extent to which patient needs, as well as barriers and facilitators to meet those needs, are accurately known and prioritized by the organization; **Compatibility:** Degree of tangible fit between meaning and values attached to the innovation by involved individuals, how those align with individuals' own norms, values, and perceived risks and needs, and how the innovation fits with existing workflows and systems; **Knowledge & Beliefs about the Innovation:** Individuals' attitudes toward and value placed on the innovation, as well as familiarity with facts, truths, and principles related to the innovation; **Complexity:** Perceived difficulty of the innovation, reflected by duration, scope, radicalness, disruptiveness, centrality, and intricacy and number of steps required to implement; **Executing:** Carrying out or accomplishing the implementation according to plan

Abbreviations: CFIR = Consolidated Framework for Implementation Research^{60, 240}

We identified and distilled evaluations and qualitative research studies to help assess the potential of provider-to-provider telehealth to improve the health and well-being of rural populations. These implementation studies and reports on programs provide insight into what is needed to translate what we know about the effectiveness of provider-to-provider telehealth (Key Question 2) to improving actual practice. Reviewing studies and reports of rural provider-to-provider telehealth reveals that there are many facilitators, barriers, and the impacts are likely not specific to rural settings. Telehealth needs to alleviate burden on providers, the technology needs to work, staff resources and reimbursement need to be allocated to provide both start-up infrastructure and ongoing support, and training is needed. Stakeholder engagement and commitment, from patients through to health system leadership and governments seems essential.

There are, however, some issues that do seem to be more specific to implementing provider-to-provider telehealth to serve rural populations. In the United States, lack of broadband internet in rural areas and the need to have offline options remains a barrier to diffusion. Disconnects or lack of understanding of the rural context and environment can reduce the utility of teleconsultations and remote training programs. Solutions that involve periodic in person visits or rotations by the specialists or trainers have been proposed as a way to both assure relevance and develop relationships. While many rural areas face similar changes, most programs are local or in a single health system and networking appears to be minimal. Successful statewide programs have been implemented and sustained, but in some cases uptake or use has been lower than anticipated. Few of the evaluations or studies identified for this review used an implementation framework or reported developing an implementation manual. Use of frameworks could standardize evaluations making comparisons across program evaluations easier and could provide a structure for documenting and disseminating implementation plans that could speed replication and uptake in more sites.

Key Question 4. What are the methodological weaknesses of the included studies of provider-to-provider telehealth for rural patients and what improvements in study design (e.g., focus on relevant comparisons and outcomes) might increase the impact of future research?

Key Points

- Studies of provider-to-provider telehealth for rural areas could be improved by addressing methodological weaknesses.
- A key weakness is that it is often difficult to attribute impact to telehealth because
 - Weaker study designs are common: RCTs and cohort studies were identified, but more than 40 percent of the studies were studies using repeated measures (pre/post or before/after) with no other comparison group.
 - Lack of control for confounders related to patients, providers, facilities, and differences in telehealth implementation across study sites
- The most frequently identified weakness after study design is small sample size and the resulting lack of power to detect differences.
- Studies are also hampered by data limitations related to use of retrospective data, and data produced for care delivery and billing purposes, rather than for research.

Summary of Findings

Conducting effectiveness and implementation research in order to develop an evidence base for telehealth is challenging for many reasons. However, two characteristics are at the root of many of these challenges: First, telehealth includes diverse interventions designed to address different healthcare functions. Even when the scope is narrowed to provider-to-provider telehealth for rural populations the interventions range from remote ICU, which requires significant hardware and other resources, to text message-based remote education using participants' cell phones. Second, telehealth is a means to facilitate communication and deliver health services, but it is not necessarily designed to impact patient health outcomes, provider experience or behavior, or payers independently of the service provided. If the service is inappropriate or care is of low quality, simply using telehealth to deliver it without addressing these issues cannot be expected to improve outcomes. Studies are not always designed to consider if services provided via telehealth vary in quality for some other reason independent of telehealth being the mode of delivery.

The methodological weaknesses of studies of telehealth have been reviewed, cited by other systematic reviews on telehealth,^{44, 49} and study design and methods limitations are often listed by the authors when studies are published, as is the case for many of the studies included in this review. Most of these weaknesses fall into two large categories, 1) the study design employed and 2) how the chosen design was executed. We attempted to identify and describe these two categories of weaknesses in the evidence base for provider-to-provider telehealth for rural populations.

It is generally accepted that an important contributor to methodological weaknesses in the literature evaluating telehealth is choice of study design. While specific designs do not correlate perfectly with weaknesses, and all study designs have some potential to produce high quality results, it is harder to product unambiguous results with less rigorous designs that are more

susceptible to bias. In order to help consider the impact of design on the evidence base for provider-to-provider telehealth, the study designs of the effectiveness studies in Key Question 2 in each setting are listed in **Table 6**. Note that in this analysis the unit is number of publications, rather than studies, as the study design can differ across publications (e.g., one publication may report the main results of an RCT and another publication from the same study may be a pre-post analysis of effects in a subgroup of patients or providers). The table is ordered in the generally accepted hierarchy for lower risk of bias or better internal validity associated with study design. In this review an almost equal number (about 20% or 1/5 each) of the publications were RCTs, prospective cohorts and retrospective cohort studies. The remaining over 2/5s (43%) of the studies used a pre-post or before-after design. These two designs are open to all the biases that are risks with observational designs, but they are also more susceptible to secular trends. This is particularly concerning in a field like telehealth with rapidly evolving technology and in which the environment and organizational context play an important role.

Table 6. Publication study designs by Key Question 2 topic

Study Design	Inpatient	Outpatient	EMS/ED	Education/ Mentoring	Total
RCT	0	15	2	3	20
Prospective Cohort	2	6	10	2	20
Retrospective Cohort	3	6	8	2	19
Pre/Post*	0	4	0	14	18
Before/After[†]	13	3	8	2	26
Total	18	34	28	23	103

* Same patients or providers evaluated pre and post intervention

[†] May not have the same patients or providers in both time points

Abbreviations: ED = emergency department; EMS = emergency medical services; RCT = randomized controlled trial.

Other weaknesses are related to how the study is conducted. One important choice in designing a study is what organization or organizations will be involved. While the number of sites and subjects in a study need to be feasible, studying an intervention in only one setting increases the likelihood that the results could be due to some characteristic or other event at that particular site. When similar results are found across multiple organizations, particularly when the study includes a range of organizations that represent important differences, we are more confident that the result can be attributed to telehealth. **Table 7** shows the distribution of studies in Key Question 2 that were conducted at single sites versus multiple sites, with 17% (18 of 103 studies) reflecting experience at only a single center. The fact that the majority of telehealth studies are including multiple sites is encouraging. Studying a complex intervention at multiple sites enhances the applicability of the findings to other locations.

Table 7. Publication site types by Key Question 2 topic

Number of Sites	Inpatient	Outpatient	EMS/ED	Education/ Mentoring	Total
Multisite	13	27	22	23	85
Single Center	5	7	6	0	18
Total	18	34	28	23	103

Abbreviations: ED = emergency department; EMS = emergency medical services.

The list of potential other weaknesses resulting from how a study is conducted is long. Weaknesses or limitations cited more than once by the authors of included studies are presented in **Table 8** (data abstractions included in **Appendix D**). These are arranged by healthcare setting as a means to potentially identify differences across settings. The table includes the weaknesses cited by the study authors and is not meant to include all possible weaknesses. For example while the inability to blind patients and clinicians in telehealth studies may introduce bias in any setting, it is only included in the row for the setting in **Table 8** if it was explicitly mentioned in included studies.

The most frequently cited weakness across all settings are small sample sizes that result in lack of power to detect statistically significant differences. Other weaknesses include lack of a clearly stated research question and hypothesis, and absence of theory or paradigmatic logic model linking telehealth to outcomes of interest. Additional challenges include historically low utilization of telehealth services, inability to randomize patients to telehealth interventions in many instances, and practical issues that rule out using approaches such as blinding to minimize bias because telehealth use cannot be concealed. Another challenge is that a wide variety of direct and indirect outcomes are utilized in telehealth studies and often measures are not clearly defined or measured. For example, different studies define time-to-treatment outcomes in different ways, and many education and mentoring studies utilized non-validated survey instruments to collect information from participating providers.

One weakness not acknowledged by the publication authors but identified by the review team is the very limited use of evaluation and implementation frameworks or logic models to structure the studies. Frameworks such as CFIR⁶⁰ and Reach, Effectiveness, Adoption, Implementation, and Maintenance (RE-AIM)²⁴¹ use theory-based domains associated with the adoption, implementation and maintenance of clinical and technological interventions to standardize terminology and facilitate comparisons across studies. Analytic frameworks and logic models make explicit how the intervention is expected to impact outcomes and how variables are related. These models usually require acknowledging the potential for harms or unintended consequences, something that was rarely done in the included studies. The fact that few studies in this report have used these frameworks made comparing both implementation and effectiveness across studies difficult and raises concerns that important elements, such as harms, may be missing in the research.

Table 8. Methodological weaknesses reported in included studies

Clinical Topic	Number of Publications	Summary of Identified Weaknesses
Inpatient Effectiveness	18	<p>Limited applicability - Single center studies, small sample sizes</p> <p>Selection bias –may be vulnerable to secular trends; allocation to groups potentially biased, differences in severity of comparison groups</p> <p>Performance bias – e.g., single researcher performed, documented, and analyzed activities (lack of blinding), changes in clinical providers over time (limits consistency of the intervention and therefore internal validity)</p> <p>Detection bias – underpowered to detect outcomes with precision; unblinded assessment of outcomes, outcomes not clearly or inconsistently defined measured, or reported across studies, outcomes not always conceptually linked to telehealth, lag times in outcome measurement or followup not long enough</p> <p>Attrition bias - Incomplete data (electronic or human data capture)</p> <p>Analysis bias – lack of controlling for confounding</p>
Outpatient Effectiveness	34	<p>Limited applicability - Single center studies, small sample sizes</p> <p>Selection bias; randomized by site without appropriate design and analysis for cluster randomization, comparison groups received enhanced usual care, or were a hypothetical reference</p> <p>Attrition bias - High attrition, high refusal to participate in all parts of studies, missing data, between-group differences in followup</p> <p>Performance bias - not all costs identified; unclear if costs (current and future) and outcomes valued appropriately, lack of methods to ensure accurate and reliable data capture from administrative databases</p> <p>Detection bias - Complex intervention without assessment of individual components, followup time inadequate for some outcomes</p> <p>Analysis bias – lack of controlling for confounding by patient characteristics (e.g., race/ethnicity)</p> <p>Other - Unclear methods for enrollment (randomization and allocation concealment), outcome ascertainment, handling of missing data, concerns about fidelity of the intervention to the protocol</p>
Emergency Management Services/ Emergency Department Effectiveness	28	<p>Limited applicability - Single service/center studies, small sample sizes</p> <p>Selection bias –may be vulnerable to secular trends; allocation to groups potentially biased (controls selected based on available data –sites not missing data may not be representative, patient or clinician factors may alter group selection)</p> <p>Performance bias – clinicians knew who received telehealth</p> <p>Retrospective design – analysis was limited to available data;</p> <p>Attrition bias - Missing data, response rate for surveys not reported</p> <p>Analysis bias - lack of control for confounders (e.g., change in practice during study period, severity of illness which may have influenced selection for teleED); comparator not well described</p>

Clinical Topic	Number of Publications	Summary of Identified Weaknesses
Education and Mentoring	23	<p>Limited applicability - Single center studies, small sample sizes</p> <p>Selection bias – analysis may be vulnerable to secular trends (e.g., pre-post designs); allocation to groups potentially biased</p> <p>Detection bias - point-in-time assessment of provider knowledge only (does not assess retention), lack of data on specific provider behavior and comprehensive, consistent and validated outcome measures</p> <p>Analysis bias – does not account for program implementation differences across “spoke” sites, impact of provider technological difficulties with the education and mentoring modality.</p> <p>Other – unclear ‘dosage’ of the intervention; people did not have to attend all sessions.</p>
Barriers & Facilitators	71	<p>Analysis bias – Frequent use of pre post design w/o control for other factors or descriptive studies</p> <p>Selection bias - Few patients or providers willing to be interviewed; differences between groups due to difficulties identifying similar controls across systems; may be vulnerable to secular trends, including perceptions that are likely to change over time</p> <p>Performance bias -Measures of frequency of use maybe misleading if the purpose of the telehealth intervention is to provide capacity to respond to a rare event.</p> <p>Detection bias - Programs are evaluated before long-term results are available; no systematic assessment of the relative impact of different barriers and facilitators; lack of direct measures of patient outcomes; lack of specificity in concepts measured; evaluation is based on impressions of what patients need rather than patient experience, and assessments based in part on a hypothetical scenario</p> <p>Attrition bias - Unknown barriers of provider non-respondents,</p> <p>Analysis bias – does not account for implementation differences across “spoke” sites, few studies use established implementation assessment frameworks that would help facilitate comparisons across sites or across studies</p>

Abbreviation: TeleED = telehealth emergency department.

Discussion

Findings in Relation to the Decisional Dilemmas

While telehealth has the potential to increase access to timely, appropriate, and high quality healthcare for rural populations its promise has yet to be realized. Use of telehealth had been steadily increasing, though many people would say too slowly, over the past three decades. In healthcare, telecommunications technology can be used in a wide variety of ways to expand options for interactions both between patients and clinicians as well as among healthcare providers.

One of the most significant impacts of the COVID-19 pandemic has been the rapid acceleration of telehealth in many forms.^{55, 242, 243} Data on use for direct patient care indicate that the crisis has been a catalyst for changes in technology, regulations, payment/reimbursement, and workflows. While virtual patient visits spiked, then declined somewhat as science and routines adapted to the pandemic, the use of telehealth remains high and is unlikely to return to low, pre pandemic levels.²⁴⁴⁻²⁴⁶ Telehealth use has likely increased as well for different uses among providers, such as remote consultations for inpatient, outpatient and emergency care, virtual care conferences, remote diagnosis and supervision of treatment, and use of technology for education and mentoring of healthcare providers. This review of telehealth for provider-to-provider communications and collaborations was initiated prior to the pandemic. Nevertheless, assembling the available evidence about use (Key Question 1), effectiveness (Key Question 2), implementation (Key Question 3), and methodological weaknesses of research studies (Key Question 4) provides a foundation that, combined with current pandemic experience, can serve as a foundation for future use of telehealth by informing policy and practice initiatives. Which temporary policies should become permanent and where telehealth investments are most likely to help address rural disparities that may have worsened during the pandemic are questions healthcare organizations and policy makers will be grappling with for the coming months and years.

This review underscores that evidence specific to rural telehealth is limited. The number of studies included in the results section of this report is deceptive. If these studies were similar, this would represent a sizable body of evidence; however, the included studies are spread across different settings, clinical indications, and roles in healthcare delivery. Our findings are organized by setting and then by specific clinical indications. We did this because most decisions are made about targeted telehealth programs for a specific use and these distinctions are important for policy and practice. The discussion below strives to provide a high-level overview of the results and an assessment of the context to complement the detail provided earlier in this report.

Key Question 1: Uptake of Provider-to-Provider Telehealth in Rural Areas

How telehealth has been used for provider collaboration in the diagnosis and management of rural patients has not been well documented in the published research literature. Despite requests and searches for unpublished data, we are unable to provide a comprehensive, empirical response to this question. Data on national overall provider-to-provider telehealth use for rural populations was not identified. We did identify studies that included data on specific uses, including six surveys and one analysis of claims data that report on national or regional trends and included

rural location in the analysis or surveyed rural providers. These document that telehealth use is different across specialties, and does appear to be increasing (see **Table 1** for details).

The data sources and designs in the few studies that were identified suggest future approaches to answering this question. Studies were based on national surveys of provider organizations, three survey of emergency departments^{63, 65, 66} one and of mental health services.⁶⁴ These surveys included items about telehealth use and contain information about the responding organizations, including location, which can be used to identify rural providers. One study used claims data to track the extent to which telestroke was used in over 1 million stroke hospitalizations and how this changed over time in rural as well as urban areas.⁶⁷

While these studies provide a very incomplete picture of provider-to-provider telehealth in rural areas, they allow two important conclusions. Telehealth use is increasing and data exists in national surveys and claims that could be used to generate a deeper understanding of recent trends and current use and this understanding could help target future policy initiatives and telehealth program development.

Key Question 2: Effectiveness of Provider-to-Provider Telehealth in Improving Health Outcomes for Rural Populations and Providers and Payers

Appendix Table B-2 provides a description of the characteristics of included studies including geographic distribution, study design, sample size, and mode of telehealth. More details about individual studies are available in the results section above and the evidence tables in **Appendix D**. The assessment of the strength of evidence (SOE) criteria, the overall assessment of the evidence by outcome is in **Appendix F**.

Assessing effectiveness of telehealth is challenging as the modes, functions and outcomes studied and how they are measured are varied. For conclusions and the assessment of the SOE supporting each of these, we attempted to group studies of similar topics that assess similar outcomes. Despite this effort, there are several instances where only one study was located that was not large or rigorous enough to support a conclusion and the overall assessment is “unclear effect” and “insufficient evidence.” Other instances where we have noted “unclear effect” and “insufficient evidence” involve two or more studies that reported conflicting results and are based on small samples (resulting in imprecise estimates). One way SOE is operationalized is the level of confidence that the next study would not change the overall conclusions. Given the lack of multiple, rigorous studies on these topics, for those that we were able to propose a conclusion the SOE is low as even one large, rigorous study could support a different conclusion.

Provider-to-Provider Telehealth for Inpatient Care

Evidence on the impact of inpatient consultations on mortality in studies that involved specialist advice on management for multiple specialties (2 studies) and infection disease (1 study) was insufficient as the results were not consistent across studies. Included studies support a conclusion that telehealth resulted in similar length of hospital stay and the rate of transfers. Two studies of remote intensive care unit (ICU) specifically in rural areas were identified, providing some limited evidence that there is no difference in mortality when patients are treated in rural hospitals using remote ICU rather than being transferred to other hospitals for ICU care (2 studies; Low SOE). Remote monitoring and consultations that allowed neonates to be cared for in rural hospitals results in clinical outcomes similar to those achieved by patients in more

specialized hospitals (Low SOE) and more appropriate transfers (2 studies; Low SOE); however, there were not consistent findings to support conclusions about the impact of teleconsultations on mortality or length of stay (Low SOE, **Table 9** and **Appendix F**).

Table 9. Strength of evidence for inpatient effectiveness studies

Topic	Outcome	Number of Studies (Combined N)	Conclusion	Strength of Evidence (Insufficient, Low, Moderate, High)
Teleconsultation	Mortality	3 (N=1841) ^{76, 83, 85}	Unclear effect	Insufficient
	Length of Stay	6 (N=2867) ^{75, 76, 78, 79, 83, 85}	Similar length of stay with telehealth consultations	Low
	Transfers	3 (N=2032) ^{76, 80, 85}	Similar transfers with telehealth consultations	Low
Remote ICU	Mortality	2(N=525) ⁸¹ (N=83 rural hospitals) ⁸²	Similar mortality rates	Low
	Transfers	1 (N=525) ⁸¹	Unclear effect	Insufficient
Management of Neonates (specialty care)	Mortality	1 (N=317) ⁷⁰	Unclear effect	Insufficient
	Length of Stay	2 (N=298) ^{73, 74}	Unclear effect	Insufficient
	Transfers	2(N=317) ⁷⁰ 1 (N=384 hospitals) ⁷²	Unclear effect	Insufficient
	Clinical Outcomes	2 (N=298) ^{73, 74}	Similar clinical measures with telehealth supported care and a higher level NICU	Low

Abbreviations: ICU = intensive care unit; NICU = neonatal intensive care unit.

Provider-to-Provider Telehealth for Outpatient Care

Studies of outpatient consultants as part of rural care included many different clinical topics. However, we only identified more than one study for depression, diabetes, pharmacy consultations, and remote consultations for multiple conditions. For depression the identified research supports the conclusion that telehealth consultations result in improvement in clinical and patient-centered outcomes including medication adherence, treatment response, and well-being (three studies; Low SOE). These studies of outpatient telehealth also documented that costs increased, though this was expected as telehealth consultations resulted in increased access to ongoing care (e.g., residential treatment, inpatient placement). Specialist support was also shown to produce better outcomes for rural diabetic patients in both laboratory tests (e.g., A1c) and patient behaviors (e.g., glucose monitoring) (4 studies; Low SOE). Similarly, studies in which pharmacists provided consultations resulted in more patients being treated according to established guidelines and improvements in physiologic measures in the more complex patients with both diabetes and hypertension. Two studies of remote consultants for multiple conditions reported mixed effects on outcomes with some better with telehealth (e.g., medication adherence), but not significantly different on others (e.g., function and quality of life) (Insufficient SOE). These are summarized in **Table 10**. For the remaining outpatient topic and outcome combinations only one study was identified and each was small, with most studies including fewer than 200 subjects and the majority less than 100. Details on these studies are included in the Results and Appendixes.

Table 10. Strength of evidence for outpatient effectiveness studies

Topic	Outcome Type	Number of Studies (Combined N)	Conclusion	Strength of Evidence (Insufficient, Low, Moderate, High)
Depression	Clinical	3 (N=974) ^{97, 111, 122}	Improvement in medication adherence, treatment response, satisfaction, and quality of well-being	Low
Depression	Payer	2 (N=724) ^{97, 122}	Higher costs associated with intended higher utilization, acceptable incremental cost effectiveness	Low
Diabetes	Clinical	4 (N=303) ^{90, 101, 103, 106}	Improved A1c, self-monitoring of blood glucose, and Diabetes Self-Management Education score	Low
Pharmacy	Clinical	2 (N=879) ^{87, 89}	Improved guideline adherence; improved A1c, fasting blood glucose, and systolic blood pressure in patients with both diabetes and hypertension (similar improvement in patients with only one of the conditions)	Low
Remote consultations	Clinical	2 (N=189) ^{94, 100}	Unclear effect	Insufficient

Provider-to-Provider Telehealth for Emergency Care

We combined our assessment of programs designed to assess ST-elevated myocardial infarction (STEMI) and chest pain with telestroke as all of these focus on facilitating remote expert review of patients and available data (e.g., imaging, electrocardiogram) with the goal of assuring timely assessment and supporting decisions on how or where patients should be treated. The available evidence supports the safety of these programs, finding that mortality rates were the same for patients treated locally with telehealth compared to those treated at more distant specialized hospitals (5 studies; Low SOE, **Table 11**), though these studies used different mortality metrics (in hospital, 30-day and 90-day). Eight studies included measures of time to treatment for these conditions in which rapid assessment and intervention are key. These studies all found that telehealth and local treatment results in either faster treatment or no difference in the time to treatment, again suggesting that patients can be effectively treated at local hospitals with telehealth support (8 studies; Low SOE).

The other studies of telehealth in emergency care involve providing access to expert advice on whether patients can be treated locally or should be transferred to a higher level hospital. Assessing this type of consultation is challenging because the goal is to both avoid unnecessary transfers and increase transfers when appropriate for the patient. Across these studies there were either improvements in or similar rates of transfer with and without telehealth. (Low SOE). Another metric used in these studies is length of stay in the emergency department (ED), the assumption being it is better to quickly either discharge or admit a patient. Despite identifying multiple studies that reported this outcome, we were unable come to a conclusion and the body of evidence was rated insufficient as the results of each study differed, with studies reporting increases, decreases, or similar length of stay for different subgroups of patients.

Table 11. Strength of evidence for emergency care effectiveness studies

Topic	Outcome Type	Number of Studies (Combined N)	Conclusion	Strength of Evidence (Insufficient, Low, Moderate, High)
EMS/ED: Telestroke/STEMI/Chest Pain	Mortality	5 (N=2,312) ^{67, 127, 139, 143, 148}	Similar mortality with telehealth supported care	Low
	Time to treatment	8 (N=3,725) ^{125-127, 133, 136, 139, 142, 148}	Similar time to treatment. No evidence of increased time to treatment	Low
EMS/ED: Consultation	Transfers	5 (N= 147,910) ^{131, 132, 135, 141, 144}	Similar appropriate patient transfers. No evidence of increased inappropriate patient transfers	Low
	ED length of stay	4 (N=9,094) ^{130, 135, 138, 147}	Unclear effect	Insufficient

Abbreviations: EMS = emergency medical services; ED = emergency department; STEMI = ST-elevation myocardial infarction

Provider-to-Provider Telehealth for Education/Mentoring

Provider-to-provider telehealth for education and mentoring of rural healthcare providers includes Extension for Community Healthcare Outcomes (ECHO) programs as well as training conducted via video or SMS (text messages). In **Table 12** the results are summarized by outcome. When patient outcomes are included in research on education programs, the results either show improvement or that similar results can be achieved in the rural locations as at the more distance hubs with onsite expertise (2 studies; Low SOE, **Table 12**). More frequently studied are provider outcomes. While the studies measured different behaviors specific to particular conditions and programs (e.g., appropriate antibiotic, opioid, and psychotropic prescribing, increasing screening, or initiation of counseling), all studies documented improvement in the desired behavior (8 studies; Low SOE). Additionally, most studies of education and mentoring evaluated provider knowledge, confidence, self-efficacy and satisfaction and report that these measures improved or were positive after participation in the programs (13 studies; Low SOE). All the reports are positive, and while this may be the reality, it is also possible that programs are unlikely to publish negative assessments by participants.

Table 12. Strength of evidence for education and mentoring effectiveness studies

Topic	Outcome Type	Number of Studies	Conclusion	Strength of Evidence (Insufficient, Low, Moderate, High)
Education/Mentoring	Clinical response	2 ^{151, 167}	Patient outcomes improve or are equivalent across spoke and hub sites	Low
	Provider behavior	8 ^{152-154, 156, 158, 159, 165, 168}	Provider behaviors improve	Low
	Provider knowledge, efficacy, perception	13 ^{153, 155, 157-161, 163, 164, 166, 170, 172, 173}	Provider knowledge, self-efficacy and perception improve	Low

Key Question 3: Barriers and Facilitators to Implementation and Implementation Strategies of Provider-to-Provider Telehealth

The original intention of this review was to focus on studies that compared implementation strategies. However, only a very few comparative studies were identified, and these compared models of telehealth for a specific use (remote ICU) and the experiences of different networks implementing telestroke rather than explicit implementations strategies. In order to summarize experience, we expanded the scope to include program evaluations that reported barriers and/or facilitators or the impact of the program and studies that used qualitative research methodology to identify and categorize factors that affected implementation. We continued to restrict our focus to provider-to-provider telehealth for rural populations and excluded publications that simply described telehealth programs without assessing barriers or facilitators to implementation in some way.

This expansion led to identifying and including 67 studies in 71 publications. The barriers, facilitators, and impacts are reported separately for each setting (i.e., inpatient, outpatient, emergency, and education) in detail in the appendix to this report.

Looking across settings, most of the barriers and facilitators are similar to what would be expected for any practice change initiative or new program. These studies confirm that the change needs to fit with the organizational mission, address a specific problem or need, have backing from organizational leadership, a local champion and support from key stakeholders at all levels. Financing and costs are important and all new programs, not just telehealth, struggle with identifying the resources needed for initial investments (e.g., hardware, software, space). Another group of factors highlighted frequently is the ‘fit’ of the intervention on many levels, including the overall mission of the organization, the needs of specific patients and providers, and telehealth’s ability to be integrated easily into the workflows and processes of specific hospitals, practices, or individual providers. Though only occasionally mentioned directly, factors cited as supporting implementation included the need for an understanding of implementation, knowledge of implementation science, explicit planning for implementation, and creating an implementation toolkit or guide that could be used to sustain the program or support its replication.

Telecommunications technology is essential to telehealth, so its role in implementation is specific to telehealth. Aspects of technology were cited as both barriers and facilitators for provider-to-provider telehealth. Technology that does not work reliably, is too expensive, or requires infrastructure that is not available, such as broadband internet, cell phone coverage, or 24/7 tech support not surprisingly slowed or prevented successful implementation. On the other hand, the ability to use existing technology, technology that functioned well, and technology familiar to users all contributed to the success of telehealth programs.

While less frequent, the studies did identify barriers and facilitators and suggest some strategies that are more specific to provider-to-provider telehealth in rural environments. Rural environments differ from the urban environments where consultants or hubs are usually located. Understanding the local culture and policy context, what healthcare and social resources are available and values and preference of patients and providers helps assure telehealth consultations and training are appropriate and relevant. This is likely the motivation for programs to suggest or include some in-person interactions including in-person visits or rotations of the consultants to the rural sites and periods in which rural providers shadow consultants or observe practice at the hub sites.

Another challenge for rural programs is identifying the inputs needed to start-up and then sustain new telehealth programs given the lack of volume, scale, and economic resources in many rural areas. Several of the rural health programs had state government or philanthropic support to start up and depended on changes in reimbursement policy to make programs financially sustainable. Creating regional or statewide networks also allowed programs to share experiences and resources and to develop approaches for rare events or infrequent needs that would not be feasible if attempted individually.

Key Question 4: Methodological Weaknesses of Studies of Provider-to-Provider Telehealth for Rural Populations

This review included an assessment of methodological weaknesses as a Key Question. For this reason, we abstracted the limitations noted by the researchers/authors and looked for trends across studies. The specific weaknesses cited in each study are included in the evidence tables in **Appendix D**. Then these are summarized in the Results chapter and by setting in **Appendix Table B-4** and are grouped by the approximate type of bias or limitation they are likely to cause.

While the majority of studies included more than one site, which supports both applicability and helps assure that site-specific environment is not the sole cause of any impact, many were still small to medium sized studies in a single location. Also, this continues to be a field with few randomized trials. Some elements of classic randomized controlled trials (RCTs), such as blinding participant and providers, are not possible. However, other designs that take advantage of the possibility or need for phased starts across sites (often referred to as step-wedge, delayed start, or wait list designs) could be used to increase the rigor of the evidence base. As pragmatic trial methods evolve and gain acceptance, trials may become more feasible. One of the major sources of methodological weaknesses is the dependence on pre-post (same subjects) or before-after (same organization, different subjects) designs. These rarely include a comparison group or documentation of historical changes in policy or practice, making it difficult to attribute changes in outcomes to telehealth and rule out other explanations.

In addition to the overarching study design, most studies did not provide information on important characteristics of their samples or control for any differences in analyses. Studies were presented with the underlying assumption that telehealth would affect outcomes, but particularly the program evaluations and qualitative studies of implementation rarely make their assumptions or underlying theory or model explicit or use a framework that would facilitate comparing implementation across studies. This is linked to the ongoing challenge of determining what the most appropriate outcomes are for telehealth interventions. In some cases, increasing access is the primary goal and process measure may be most important; in others concerns about safety of objectives related to health disparities require including patient clinical outcomes.

Limitations of the Review Process

There are limitations to this review that are the product of our process and the decisions, tools, and methods available for systematic reviews. Searching for rural provider-to-provider telehealth studies poses several challenges that required assessing whether and how to use specific indexing and key word terms. While telehealth is increasingly indexed, it is a broad term that overlaps with others. Additionally, while the Mesh term “Remote Consultation” exists in Medline this does not capture all possible models of provider-to-provider telehealth. Furthermore, studies do not always include ‘rural’ in titles or abstracts or are not always indexed

as including either rural or underserved populations. Given these challenges we worked with an expert reference librarian with extensive experience with systematic reviews and tested combinations of index terms and key words. Our search strategies are included in **Appendix A**. Despite this approach and our supplemental efforts which included checking references of included studies and other systematic reviews, suggestions from stakeholders, and responses to requests for data, it is possible that relevant studies were missed.

Given the variety of study designs, interventions, outcomes and the lack of detail on comparators in many studies, we were unable to use quantitative synthesis (i.e., meta-analyses). This diversity makes even qualitative synthesis of the effectiveness studies (Key Question 2) challenging. We made decisions to combine studies with similar, but not exactly the same outcome, and to combine some studies that were used for different clinical indications in our SOE assessment. By expanding our scope for the implementation studies to include studies that were not comparative and by including evaluation studies as well as qualitative studies, we were able to provide a broader picture of the state of the science but at the cost of not being able to apply standard risk of bias or strength of evidence assessments. Summarizing and synthesizing in this way requires significant subjective judgements and it is possible that others could interpret the same evidence differently.

Limitations of the Evidence Base

We were able to identify 97 studies (in 106 articles) published in the last 10 years that evaluated the effectiveness or implementation of provider-to-provider telehealth for rural populations. For decades, telehealth has been advanced as an important means of addressing disparities in healthcare access and health outcomes in rural areas. The existence of studies that directly answer the Key Questions is a strength. However, important limitations to this evidence base need to be considered as they impact the utility of this research for practice and policy decisions. Most of the apparent limitations are related to the relative weakness of study designs used in this field, the rigor with which the studies were executed, and the completeness of documentation.

Methodological weaknesses are the focus of a separate Key Question in this review (see Key Question 4 in Results and above) so they are summarized here briefly. Most of the included studies were smaller and are not randomized trials, though the majority involved more than one site. Many studies of telehealth compared outcomes before and after the implementation of telehealth or compare cohorts of patients, clinicians, or organizations with and without telehealth and do not include comparison groups or efforts to isolate the effect of telehealth from historical trends or changes over time. We assessed the risk of bias of all the included effectiveness studies (Key Question 2); very few studies were rated as low risk of bias, with most rated as medium, and several as high risk of bias (**Appendix E**). The criteria that are not met in these studies are those related to selection bias (e.g., whether inclusion based on a random sample or all that meet inclusion criteria and whether analyses account for important potential confounding).

Other weaknesses include the lack of a clear conceptual model and agreement on what outcomes are most important. There have been efforts to develop frameworks for telehealth research and quality measures.^{247, 248} These measures have not consistently been used in the literature published to date. Additionally, the lack of a clear conceptual model and related explicit statements about the intention of telehealth in most studies make interpreting the impact on outcomes challenging. In many studies it is implied, but not explicitly stated that the goal is to achieve patient outcomes that are similar, not necessarily better, than those achieved without

telehealth. But the studies do not provide the context needed to determine if similar outcomes are truly an overall benefit. Additionally most studies, do not acknowledge potential harms or that telehealth could be inferior to face-to-face care in some situations or for some people. Negative outcomes are not defined and measured, making it difficult to determine when in person care may be superior or required. The COVID-19 pandemic resulted in changes in existing telehealth quality measures, in consideration of approaches to measure the quality of telehealth,²⁴⁹ and more explicit statements about telehealth intentions (e.g., provide equivalent care while reducing infection risk). However, these newer measures and frameworks have not yet been integrated into studies that have been completed.

In addition to weaknesses in study design and execution, the lack of key information in published research reports is also a weakness of this evidence base. Both the telehealth interventions and the comparison group or time period are often not well described. This is true in both effectiveness and implementation studies although implementation studies tend to provide more detail. Within each use of telehealth there are a wide range of models. For example, a remote ICU system could use hard wired monitoring in ICU patient rooms or a portable cart or 'robot;' consults for emergency care could be available at the push of a call button or may require paging the appropriate services; and outpatient consultations may or may not involve the patient and providers in real time. These details are often not provided. While more details were often provided in implementation studies (Key Question 3), even these did not always describe the specific telehealth intervention. Combined with the lack of consistent, clearly defined outcomes, lack of information about the specifics of telehealth interventions make it difficult to synthesize results across studies, even qualitatively. Furthermore, identification of facilitators and barriers to implementation was often not the primary purpose of the studies included in Key Question 3. Many studies mentioned facilitators and barriers in the context of pragmatic lessons learned during telehealth deployment. The result is a collection of findings that may not provide a clear, definitive path for decision makers.

Applicability

The scope of this review was defined to include a subset of telehealth (i.e., provider-to-provider communication and collaboration) used to benefit a specific population (i.e., patients in rural areas). As other uses of telehealth and studies that did not explicitly focus on rural populations were excluded, the identified evidence does not differ significantly from the populations, interventions, comparators, outcomes, and settings (PICOS) reflected in the Key Questions. However, there are finer grained applicability issues that should be considered when interpreting and applying this evidence.

One consideration that is rarely acknowledged in summarizing studies related to rural health is that there is diversity across rural settings. The barriers to healthcare access and underlying causes of health disparities as well as the resources that can be marshalled may differ from one rural area to another. Geography, weather, history, natural resources, national policies, and U.S. state and local policies all create an environmental context that could affect whether a telehealth intervention that was successful in one rural area can be translated to another. It may not be reasonable to expect a program that worked for remote island communities to work for dispersed populations in the mountain west of the United States or the expansive Navajo lands.

Additionally, it may be difficult to translate some models that could use different technology or require different infrastructure across or even within clinical applications or care settings, and be confident of obtaining similar results. For example, remote ICUs can be set up using very

different hardware and organizational structures. Another example is that provider-to-provider telehealth for wound care or oral health in long-term care may be more difficult to implement as it requires specialized cameras that may be too expensive or require technical support and infrastructure that are not available.

Perhaps the biggest concern related to applicability is when the included studies were conducted. The time period covered by this review includes 2020 which was characterized by a sudden acceleration in the adoption of telehealth services, concurrent with rapid policy changes regarding reimbursement for healthcare services in response to the COVID-19 pandemic. However, although we included studies published in 2020 they are based on data collected under different policies, before provider and patient familiarity with telehealth services changed significantly. Studies of telehealth in this new context are ongoing, but were not published at the time of this review.

Implications for Clinical and Policy Decision Making

The COVID-19 pandemic has put a spotlight on healthcare systems, revealing numerous cracks in access to and quality of care and subgroups of the population who continue to fall through these cracks. Many observed deficiencies in the U.S. system affect rural populations and healthcare providers and organizations that serve rural areas. While the causes of worse healthcare outcomes in rural America are multi-factorial, telehealth has been and is now even more likely to be viewed as at least part of the solution.

The results of this systematic review do not contradict the idea that provider-to-provider telehealth can contribute to better health outcomes for rural populations. It is reassuring that we did not find any evidence that provider-to-provider telehealth resulted in harms or worse outcomes. However, this must be tempered by the fact that many studies do not explicitly define and then measure potential harms or unintended negative consequences of telehealth. Many of the results that found no significant difference between care that is facilitated by telehealth consultations or remote education can be interpreted that outcomes are “as good as” those resulting from in-person collaboration (although most of these studies were not designed to test equivalence). Other studies were evaluations of programs aiming for better outcomes. This was often the case for before/after or pre/post studies of telehealth programs that were designed to augment existing services, provide services that were not available without telehealth, or provide costly services in a less costly way. The studies may report better outcomes, but the evaluations are not methodologically rigorous, and provide imprecise estimates of effect, which contributed to rating as low SOE.

At the same time, the identified evidence does not provide a clear road map that can be easily followed. Telehealth can be used to address a wide range of problems, from lack of access or expertise to travel burden and the cost implications of patient transfers. Given this breadth, interventions can include all possible settings and a large number of clinical indications and applications. There is not yet enough evidence to conclude where, across all the available problems and options, investments in provider-to-provider telehealth are likely to make the biggest difference. For example, comparing remote ICU care to ECHO programs to decide which should be implemented is difficult.

Within a clinical use there are examples of targeted needs assessments for telehealth. For example, some of the programs that establish statewide telestroke programs have combined information on health status and resources by geographic region in order to identify where hubs and spokes could be placed to best serve the state.¹⁷⁴ Regional assessments of need like this

could extend beyond a single clinical indication or a single form of telehealth to provide more comprehensive guidance and ultimately more efficient targeting of telehealth.

Future Research Needs

A common, but very unsatisfying and impractical response to reviewing a body of evidence dominated by non-randomized studies is to conclude that more RCTs are needed. Identifying where resources should be invested in trials is an important task in any field. Additionally, randomized studies present both ethical and logistical challenges, particularly when underserved or disadvantaged populations are studied. However, addressing the methodological issues identified in the response to Key Question 4 and considering different types of research could also advance the field and promote the health and well-being of rural communities and the people and organizations that serve them, without having to fund expensive RCTs. Major improvements would include:

- Outcome measurement at multiple time points, as effects may become evident initially or over longer periods of time, and because short-term outcomes may differ from longer-term outcomes (e.g., provider retention of knowledge acquired through remote education and mentoring)
- Agreement on common outcomes and goals for similar provider-to provider telehealth interventions. For example, agreeing on: whether remotes ICUs should reduce ICU or all inpatient mortality and length of stay; if ED consultations should reduce the overall rate of transfers or specific types of transfers; what measures of patient access (e.g., time to appointment, care or diagnosis) or patient burden (e.g., travel cost, lost work time, anxiety) can be measured consistently across interventions; and what harms should be measured.
- Use of contemporary comparison groups and the inclusion of multiple sites so that the impact of telehealth could be separated from historical change or the potentially unique characteristics of specific sites or providers/consultants.
- More complete descriptions of both the content of telehealth and comparator interventions and the intention of the research articles would help inform assessments of fit and help clarify when telehealth replaced in-person services or added additional services.

In addition to these specific changes to trials or observational studies, different types of studies or studies of different topics may be useful. Given the need to understand both effectiveness and implementation, hybrid studies seem warranted. Hybrid designs fall on the continuum between purely effectiveness studies to purely implementation studies and include differing degrees of focus on the context and needs for implementation in addition to assessing effectiveness.^{250, 251} Hybrid designs are one way to include more explicit consideration of the impact of the internal environment and local policies. Additionally, studies are needed that consider not just healthcare delivery, but the larger policy context in which care was delivered, including payment, licensing, regulations, and competition across providers.

Perhaps the most pressing future research need is to take advantage of the natural experiment resulting from the COVID-19 pandemic. Rapid telehealth adoption for patient-provider interactions has likely spilled over into provider-to-provider communications and collaborations. Larger studies should now be possible, involving more sites, and changes in reimbursement could lead to better coding or use of currently underused codes for activities like remote

consultations and in turn to more informative claims data that will allow tracking of telehealth consultations and associated patient outcomes. Other changes related to the pandemic have likely also created opportunities for robust studies of provider-to-provider telehealth.

Summary and Conclusion

This review identified and included 166 studies of provider-to-provider telehealth in rural areas. Evidence on where and how provider-to-provider telehealth is being used in the rural United States is sparse and generally limited to a small number of analyses of specific specialties. While these studies all report increasing use of telehealth, they do not provide a complete picture. Studies of effectiveness of provider-to-provider telehealth are dominated by findings of similar or improved outcomes when telehealth is implemented. However, it is not possible to make a universal conclusion across all the different clinical uses, various settings, and outcomes considered in the included research we examined.

Limiting the scope to provider-to-provider collaboration and use in rural areas makes the studies we identified directly applicable; nevertheless, there were too few studies to support clear conclusions with high SOE. We did find sufficient evidence to suggest that telehealth is likely to produce outcomes that are as good as or better in the following situations: inpatient consultations, the care of neonates, outpatient depression and diabetes care, emergency care for stroke/heart attack/chest pain, and emergency consultations for multiple conditions. Also, education and mentoring programs produce improvements in patient clinical outcomes, provider behavior, and provider knowledge and self-efficacy. The strength of evidence for all of these conclusions is low due to inconsistencies in results and the lack of precise estimates of effect given the small number of studies and small sample sizes.

When we searched for and analyzed studies of implementation, we found few comparative studies and our assessment across studies identified barriers and facilitators that are commonly linked to practice change efforts and are not unique to telehealth or rural implementation. Exceptions were the need to understand the rural context and the possible challenges presented by lack of scale. Similarly, assessing methodological weaknesses of the identified and included studies identified the need for larger, more rigorous observational studies, quasi-experimental studies and more trials (including adaptive trials), but also suggested that more studies should combine effectiveness and implementation research, or at minimum provide more detail on the telehealth interventions, comparators, and the context.

The existing evidence base for provider-to-provider telehealth is insufficient to support conclusions in which we can have confidence. However, it is reassuring that the studies we were able to identify do not report harms or negative consequences and do suggest that it is likely that application of telehealth can improve patient outcomes such as access to and quality of care, provider outcomes such as knowledge and self-efficacy, and payer outcomes such as reduced costs or maintenance of payments to rural providers.

References

1. Rural health. *Health Aff (Millwood)*. 2019 Dec;38(12):1964-5. doi: 10.1377/hlthaff.2019.01365. PMID: 31794298.
2. Singh GK, Siahpush M. Widening rural-urban disparities in all-cause mortality and mortality from major causes of death in the USA, 1969-2009. *J Urban Health*. 2014 Apr;91(2):272-92. doi: 10.1007/s11524-013-9847-2. PMID: 24366854.
3. Garcia MC, Rossen LM, Bastian B, et al. Potentially excess deaths from the five leading causes of death in metropolitan and nonmetropolitan counties - United States, 2010-2017. *MMWR Surveill Summ*. 2019 Nov 8;68(10):1-11. doi: 10.15585/mmwr.ss6810a1. PMID: 31697657.
4. Hall JE, Moonesinghe R, Bouye K, et al. Racial/ethnic disparities in mortality: contributions and variations by rurality in the United States, 2012-2015. *Int J Environ Res Public Health*. 2019 Feb 2;16(3)doi: 10.3390/ijerph16030436. PMID: 30717345.
5. Villapiano N, Iwashyna TJ, Davis MM. Worsening Rural-Urban Gap in Hospital Mortality. *J Am Board Fam Med*. 2017 Nov-Dec;30(6):816-23. doi: 10.3122/jabfm.2017.06.170137. PMID: 29180557.
6. Keyes KM, Cerdá M, Brady JE, et al. Understanding the rural-urban differences in nonmedical prescription opioid use and abuse in the United States. *Am J Public Health*. 2014 Feb;104(2):e52-9. doi: 10.2105/ajph.2013.301709. PMID: 24328642.
7. Monnat SM, Rigg KK. Examining rural/urban differences in prescription opioid misuse among US adolescents. *J Rural Health*. 2016 Spring;32(2):204-18. doi: 10.1111/jrh.12141. PMID: 26344571.
8. Boring MA, Hootman JM, Liu Y, et al. Prevalence of arthritis and arthritis-attributable activity limitation by urban-rural county classification - United States, 2015. *MMWR Morb Mortal Wkly Rep*. 2017 May 26;66(20):527-32. doi: 10.15585/mmwr.mm6620a2. PMID: 28542117.
9. Croft JB, Wheaton AG, Liu Y, et al. Urban-rural county and state differences in chronic obstructive pulmonary disease - United States, 2015. *MMWR Morb Mortal Wkly Rep*. 2018 Feb 23;67(7):205-11. doi: 10.15585/mmwr.mm6707a1. PMID: 29470455.
10. Krishna S, Gillespie KN, McBride TM. Diabetes burden and access to preventive care in the rural United States. *J Rural Health*. 2010 Winter;26(1):3-11. doi: 10.1111/j.1748-0361.2009.00259.x. PMID: 20105262.
11. Primm K, Ferdinand AO, Callaghan T, et al. Congestive heart failure-related hospital deaths across the urban-rural continuum in the United States. *Prev Med Rep*. 2019 Dec;16:101007. doi: 10.1016/j.pmedr.2019.101007. PMID: 31799105.
12. Nelson JA, Kinder A, Johnson AS, et al. Differences in selected HIV care continuum outcomes among people residing in rural, urban, and metropolitan areas-28 US jurisdictions. *J Rural Health*. 2018 Dec;34(1):63-70. doi: 10.1111/jrh.12208. PMID: 27620836.
13. Zahnd WE, Rodriguez C, Jenkins WD. Rural-urban differences in human papillomavirus-associated cancer trends and rates. *J Rural Health*. 2019 Mar;35(2):208-15. doi: 10.1111/jrh.12305. PMID: 29808500.
14. Galambos CM. Health care disparities among rural populations: a neglected frontier. *Health Soc Work*. 2005 Aug;30(3):179-81. doi: 10.1093/hsw/30.3.179. PMID: 16190293.
15. Garcia MC, Faul M, Massetti G, et al. Reducing potentially excess deaths from the five leading causes of death in the rural United States. *MMWR Surveill Summ*. 2017 Jan 13;66(2):1-7. doi: 10.15585/mmwr.ss6602a1. PMID: 28081057.
16. Souch JM, Cossman JS. A commentary on rural-urban disparities in COVID-19 testing rates per 100,000 and risk factors. *J Rural Health*. 2020 Apr 13doi: 10.1111/jrh.12450. PMID: 32282964.

17. Centers for Disease Control and Prevention. Why Rural Communities May Be at Higher Risk During the COVID-19 Pandemic. 2020. <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/other-at-risk-populations/rural-communities.html#why-higher-risk>. Accessed September 5, 2020.
18. Murphy T, Marema T. New Covid-19 Cases are on the Rise Again in Rural Counties. 2020. <https://dailyyonder.com/new-covid-19-cases-are-on-the-rise-again-in-rural-counties/2020/09/01/>. Accessed September 5, 2020.
19. Wright B, Fraher E, Holder MG, et al. Will community health centers survive COVID-19? *J Rural Health*. 2020 May 16;16:16. doi: 10.1111/jrh.12473. PMID: 32415879.
20. The Larry A Green Center. Quick COVID-19 Survey. 2020. <https://www.green-center.org/covid-survey>. Accessed September 4, 2020.
21. The Chartis Center for Rural Health. The Rural Health Safety Net Under Pressure: Rural Hospital Vulnerability. 2020. https://www.ivantageindex.com/wpcontent/uploads/2020/02/CCRH_Vulnerability-Research_Final-02.14.20.pdf.
22. Probst JC, Moore CG, Glover SH, et al. Person and place: the compounding effects of race/ethnicity and rurality on health. *Am J Public Health*. 2004 Oct;94(10):1695-703. doi: 10.2105/ajph.94.10.1695. PMID: 15451735.
23. Austin B, Glaeser E, Summers L. Saving the heartland: place-based policies in 21st century America. . BPEA Conference Draft. Spring 2018.
24. Porter E. The Hard Truths of Trying to "Save" the Rural Economy. 2018. <https://www.nytimes.com/interactive/2018/12/14/opinion/rural-america-trump-decline.html>. Accessed September 9, 2020.
25. Carnethon MR, Kershaw KN, Kandula NR. Disparities research, disparities researchers, and health equity. *JAMA*. 2019 Dec 16doi: 10.1001/jama.2019.19329. PMID: 31841578.
26. Jones NL, Breen N, Das R, et al. Cross-cutting themes to advance the science of minority health and health disparities. *Am J Public Health*. 2019 Jan;109(S1):S21-s4. doi: 10.2105/ajph.2019.304950. PMID: 30699031.
27. Srinivasan S, Williams SD. Transitioning from health disparities to a health equity research agenda: the time is now. *Public Health Rep*. 2014 Jan-Feb;129 Suppl 2(Suppl 2):71-6. doi: 10.1177/00333549141291s213. PMID: 24385668.
28. Sirotich E, Hausmann JS. Removing barriers and disparities in health: lessons from the COVID-19 pandemic. *Nat Rev Rheumatol*. 2021 Mar;17(3):125-6. doi: 10.1038/s41584-020-00524-8. PMID: 33028959.
29. Khan SS, Krefman AE, McCabe ME, et al. Association between county-level risk groups and COVID-19 outcomes in the United States: a socioecological study. *BMC Public Health*. 2022 Jan 13;22(1):81. doi: 10.1186/s12889-021-12469-y. PMID: 35027022.
30. Sood L, Sood V. Being African American and Rural: A Double Jeopardy From COVID-19. *J Rural Health*. 2021 Jan;37(1):217-21. doi: 10.1111/jrh.12459. PMID: 32362036.
31. Siegel M, Critchfield-Jain I, Boykin M, et al. Actual Racial/Ethnic Disparities in COVID-19 Mortality for the Non-Hispanic Black Compared to Non-Hispanic White Population in 353 US Counties and Their Association with Structural Racism. *J Racial Ethn Health Disparities*. 2021 Aug 30:1-29. doi: 10.1007/s40615-021-01109-1. PMID: 34462902.
32. Shippee TP, Akosionu O, Ng W, et al. COVID-19 Pandemic: Exacerbating Racial/Ethnic Disparities in Long-Term Services and Supports. *J Aging Soc Policy*. 2020 Jul-Oct;32(4-5):323-33. doi: 10.1080/08959420.2020.1772004. PMID: 32476614.

33. Louis-Jean J, Cenat K, Njoku CV, et al. Coronavirus (COVID-19) and Racial Disparities: a Perspective Analysis. *Journal of racial and ethnic health disparities*. 2020;7(6):1039-45. doi: 10.1007/s40615-020-00879-4. PMID: 33025419.
34. Tuckson RV, Edmunds M, Hodgkins ML. Telehealth. *N Engl J Med*. 2017 Oct 19;377(16):1585-92. doi: 10.1056/NEJMSr1503323. PMID: 29045204.
35. Telemedicine, telehealth, and health information technology. An ATA Issue Paper. American Telemedicine Association. May 2006.
36. Rae M, Cox C, Claxton G. Coverage and utilization of telemedicine services by enrollees in large employer plans. 2020. <https://www.healthsystemtracker.org/brief/coverage-and-utilization-of-telemedicine-services-by-enrollees-in-large-employer-plans/>. Accessed September 5, 2020.
37. Castro D, Miller B, Nager A. Unlocking the Potential of Physician-to-Patient Telehealth Services. 2014. <https://itif.org/publications/2014/05/12/unlocking-potential-physician-patient-telehealth-services>. Accessed September 5, 2020.
38. Lustig T. The role of telehealth in an evolving health care environment - workshop summary. Washington DC: 2012. doi: 10.17226/13466. PMID: 24901186.
39. Bashshur RL, Shannon GW, Smith BR, et al. The empirical foundations of telemedicine interventions for chronic disease management. *Telemed J E Health*. 2014 Sep;20(9):769-800. doi: 10.1089/tmj.2014.9981. PMID: 24968105.
40. Ekland AG, Bowes A, Flottorp S. Effectiveness of telemedicine: a systematic review of reviews. *Int J Med Inf*. 2010 Nov;79(11):736-71. doi: 10.1016/j.ijmedinf.2010.08.006. PMID: 20884286.
41. Hersh WR, Wallace JA, Patterson PK, et al. Telemedicine for the Medicare Population-evidence reports/technology assessments, No. 24. Rockville, MD: July 2001. <http://www.ncbi.nlm.nih.gov/books/NBK33341/>.
42. Hersh WR, Hickman D, Severance S, et al. Telemedicine for the Medicare population: update - evidence reports/technology assessments, No. 131. Rockville, MD.: Feb. 2006. <https://www.ncbi.nlm.nih.gov/books/NBK37953/>.
43. Hersh WR, Wallace JA, Patterson PK, et al. Telemedicine for the medicare population: pediatric, obstetric, and clinician-indirect home interventions. *Evid Rep Technol Assess (Summ)*. 2001 Aug(24 Suppl):1-32. PMID: 11569328.
44. Totten AM, Womack DM, Eden KB, et al. Telehealth: mapping the evidence for patient outcomes from systematic reviews - technical brief No. 26 Pacific Northwest Evidence-based Practice Center. Rockville, MD: June 2016. https://effectivehealthcare.ahrq.gov/sites/default/files/pdf/telehealth_technical-brief.pdf.
45. Scott Kruse C, Karem P, Shifflett K, et al. Evaluating barriers to adopting telemedicine worldwide: a systematic review. *J Telemed Telecare*. 2018 Jan;24(1):4-12. doi: 10.1177/1357633x16674087. PMID: 29320966.
46. Adler-Milstein J, Kvedar J, Bates DW. Telehealth among US hospitals: several factors, including state reimbursement and licensure policies, influence adoption. *Health Aff (Millwood)*. 2014 Feb;33(2):207-15. doi: 10.1377/hlthaff.2013.1054. PMID: 24493762.
47. Broderick A, Lindeman D. Scaling Telehealth Programs: Lessons from Early Adopters. New York, NY: Commonwealth Fund. 2013. <http://www.commonwealthfund.org/Publications/Case-Studies/2013/Jan/Telehealth-Synthesis.aspx>. Accessed July 8, 2016.
48. Beck M. How Telemedicine Is Transforming Health Care. *The Wall Street Journal*; 2016. <https://www.wsj.com/articles/how-telemedicine-is-transforming-health-care-1466993402>.

49. Totten AM, Hansen RN, Wagner J, et al. Telehealth for acute and chronic care consultations. AHRQ Comparative Effectiveness Review No. 216. (Prepared by Pacific Northwest Evidence-based Practice Center under Contract No. 290-2015-00009-I.) AHRQ Publication No. 19-EHC012-EF. Rockville, MD: Agency for Healthcare Research and Quality. 2019 04 PMID: 31577401.
50. Liddy C, Moroz I, Mihan A, et al. A systematic review of asynchronous, provider-to-provider, electronic consultation services to improve access to specialty care available worldwide. *Telemed J E Health*. 2019 03;25(3):184-98. doi: 10.1089/tmj.2018.0005. PMID: 29927711.
51. du Toit M, Malau-Aduli B, Vangaveti V, et al. Use of telehealth in the management of non-critical emergencies in rural or remote emergency departments: a systematic review. *J Telemed Telecare*. 2019 Jan;25(1):3-16. doi: 10.1177/1357633X17734239. PMID: 28980853.
52. Ramnath VR, Khazeni N. Centralized monitoring and virtual consultant models of tele-ICU care: a side-by-side review. *Telemed J E Health*. 2014 Oct;20(10):962-71. doi: 10.1089/tmj.2014.0024. PMID: 25225795.
53. Berndt A, Murray CM, Kennedy K, et al. Effectiveness of distance learning strategies for continuing professional development (CPD) for rural allied health practitioners: a systematic review. *BMC Med Educ*. 2017 Jul 12;17(117):1-13. doi: 10.1186/s12909-017-0949-5. PMID: 28701199.
54. McBain RK, Sousa JL, Rose AJ, et al. Impact of Project ECHO models of medical tele-education: a systematic review. *J Gen Intern Med*. 2019 12;34(12):2842-57. doi: 10.1007/s11606-019-05291-1. PMID: 31485970.
55. Steeves-Reece AL, Elder NC, Graham TA, et al. Rapid deployment of a statewide COVID-19 ECHO program for frontline clinicians: early results and lessons learned. *J Rural Health*. 2020 May 12;37(1):227-30. doi: 10.1111/jrh.12462. PMID: 32396224.
56. Centers for Medicare & Medicaid Services. Coronavirus waivers & flexibilities. 2020. <https://www.cms.gov/about-cms/emergency-preparedness-response-operations/current-emergencies/coronavirus-waivers>. Accessed September 8, 2020.
57. Hirko KA, Kerver JM, Ford S, et al. Telehealth in response to the COVID-19 pandemic: implications for rural health disparities. *J Am Med Inform Assoc*. 2020 Jun 26;27:1816-18. doi: 10.1093/jamia/ocaa156. PMID: 32589735.
58. AHRQ Methods for Effective Health Care. Methods Guide for Effectiveness and Comparative Effectiveness Reviews. Rockville (MD): Agency for Healthcare Research and Quality (US); 2008.
59. Berkman ND, Lohr KN, Ansari MT, et al. Grading the strength of a body of evidence when assessing health care interventions: an EPC update. *J Clin Epidemiol*. 2015 Nov;68(11):1312-24. doi: 10.1016/j.jclinepi.2014.11.023. PMID: 25721570.
60. Damschroder LJ, Aron DC, Keith RE, et al. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci*. 2009 Aug 7;4:50. doi: 10.1186/1748-5908-4-50. PMID: 19664226.
61. Alishahi Tabriz A, Turner K, Williams D, et al. Association of financial factors and telemedicine adoption for heart attack and stroke care among rural and urban hospitals: a longitudinal study. *Telemed J E Health*. 2021 Sep 24;24:24. doi: 10.1089/tmj.2021.0341. PMID: 34559014.
62. Chen J, Amaize A, Barath D. Evaluating telehealth adoption and related barriers among hospitals located in rural and urban areas. *J Rural Health*. 2020 Nov 12;37(1):227-30. doi: 10.1111/jrh.12534. PMID: 33180363.
63. Freeman RE, Boggs KM, Zachrisson KS, et al. National study of telepsychiatry use in U.S. emergency departments. *Psychiatr Serv*. 2020 Jun 01;71(6):540-6. doi: 10.1176/appi.ps.201900237. PMID: 32019430.

64. Spivak S, Spivak A, Cullen B, et al. Telepsychiatry use in U.S. mental health facilities, 2010-2017. *Psychiatr Serv.* 2020 02 01;71(2):121-7. doi: 10.1176/appi.ps.201900261. PMID: 31615370.
65. Zachrison KS, Boggs KM, Hayden EM, et al. Understanding barriers to telemedicine implementation in rural emergency departments. *Ann Emerg Med.* 2020 03;75(3):392-9. doi: 10.1016/j.annemergmed.2019.06.026. PMID: 31474481.
66. Zachrison KS, Hayden EM, Schwamm LH, et al. Characterizing New England emergency departments by telemedicine use. *West J Emerg Med.* 2017 Oct;18(6):1055-60. doi: 10.5811/westjem.2017.8.34880. PMID: 29085537.
67. Zhang D, Wang G, Zhu W, et al. Expansion of telestroke services improves quality of care provided in super rural areas. *Health Aff (Millwood).* 2018 12;37(12):2005-13. doi: 10.1377/hlthaff.2018.05089. PMID: 30633675.
68. Casey MM, Sorensen TD, Elias W, et al. Current practices and state regulations regarding telepharmacy in rural hospitals. *Am J Health-Syst Pharm.* 2010 Jul 01;67(13):1085-92. doi: 10.2146/ajhp090531. PMID: 20554595.
69. Hall RW, Hall-Barrow J, Garcia-Rill E. Neonatal regionalization through telemedicine using a community-based research and education core facility. *Ethn Dis.* 2010;20(1 Suppl 1):S1-136-40. PMID: 20521402.
70. Haynes SC, Dharmar M, Hill BC, et al. The impact of telemedicine on transfer rates of newborns at rural community hospitals. *Acad Pediatr.* 2020 Jul;20(5):636-41. doi: 10.1016/j.acap.2020.02.013. PMID: 32081766.
71. Haynes SC, Hoffman KR, Patel S, et al. The use of telemedicine for stabilization of neonates transferred from rural community hospitals. *Telemed J E Health.* 2021 Mar 01;01:01. doi: 10.1089/tmj.2020.0503. PMID: 33691080.
72. Kim EW, Teague-Ross TJ, Greenfield WW, et al. Telemedicine collaboration improves perinatal regionalization and lowers statewide infant mortality. *J Perinatol.* 2013 Sep;33(9):725-30. doi: 10.1038/jp.2013.37. PMID: 23579490.
73. Makkar A, McCoy M, Hallford G, et al. A hybrid form of telemedicine: a unique way to extend intensive care service to neonates in medically underserved areas. *Telemed J E Health.* 2018 09;24(9):717-21. doi: 10.1089/tmj.2017.0155. PMID: 29298407.
74. Makkar A, McCoy M, Hallford G, et al. Evaluation of neonatal services provided in a level II NICU utilizing hybrid telemedicine: a prospective study. *Telemed J E Health.* 2020 Feb;26(2):176-83. doi: 10.1089/tmj.2018.0262. PMID: 30835166.
75. Avent ML, Walker D, Yarwood T, et al. Implementation of a novel antimicrobial stewardship strategy for rural facilities utilising telehealth. *Int J Antimicrob Agents.* 2021 Jun;57(6):106346. doi: 10.1016/j.ijantimicag.2021.106346. PMID: 33882332.
76. Tande AJ, Berbari EF, Ramar P, et al. Association of a remotely offered infectious diseases eConsult service with improved clinical outcomes. *Open Forum Infect Dis.* 2020 Jan:1-4. doi: 10.1093/ofid/ofaa003. PMID: 31988969.
77. Yam P, Fales D, Jemison J, et al. Implementation of an antimicrobial stewardship program in a rural hospital. *Am J Health-Syst Pharm.* 2012 Jul 01;69(13):1142-8. doi: 10.2146/ajhp110512. PMID: 22722593.
78. Gallagher R, Giles M, Morison J, et al. Telehealth-based model of care redesign to facilitate local fitting and management of patients with a spinal fracture requiring a thoracic lumbar sacral orthosis in rural hospitals in New South Wales. *Aust J Rural Health.* 2018 Jun;26(3):181-7. doi: 10.1111/ajr.12407. PMID: 29573012.
79. Barber M, Frieslick J, Maclean A, et al. The Western Isles Stroke Telerehabilitation (Specialist Medical Consultation) Service—implementation and evaluation. *Eur Res Telemed.* 2015;4(1):19-24.

80. Buckley D, Weisser S. Videoconferencing could reduce the number of mental health patients transferred from outlying facilities to a regional mental health unit. *Aust N Z J Public Health*. 2012 Oct;36(5):478-82. doi: 10.1111/j.1753-6405.2012.00915.x. PMID: 23025371.
81. Panlaqui OM, Broadfield E, Champion R, et al. Outcomes of telemedicine intervention in a regional intensive care unit: a before and after study. *Anaesth Intensive Care*. 2017 09;45(5):605-10. PMID: 28911290.
82. Kahn JM, Le TQ, Barnato AE, et al. ICU Telemedicine and Critical Care Mortality: A National Effectiveness Study. *Med Care*. 2016 Mar;54(3):319-25. doi: 10.1097/MLR.0000000000000485. PMID: 26765148.
83. Boltz M, Cuellar NG, Cole C, et al. Comparing an on-site nurse practitioner with telemedicine physician support hospitalist programme with a traditional physician hospitalist programme. *J Telemed Telecare*. 2019 May;25(4):213-20. doi: 10.1177/1357633X18758744. PMID: 29498301.
84. Dharmar M, Sadorra CK, Leigh P, et al. The financial impact of a pediatric telemedicine program: a children's hospital's perspective. *Telemed J E Health*. 2013 Jul;19(7):502-8. doi: 10.1089/tmj.2012.0266. PMID: 23837516.
85. Kuperman EF, Linson EL, Klefstad K, et al. The virtual hospitalist: a single-site implementation bringing hospitalist coverage to critical access hospitals. *J Hosp Med*. 2018 11 01;13(11):759-63. doi: 10.12788/jhm.3061. PMID: 30255859.
86. Poulson LK, Nissen L, Coombes I. Pharmaceutical review using telemedicine--a before and after feasibility study. *J Telemed Telecare*. 2010;16(2):95-9. doi: 10.1258/jtt.2009.090716. PMID: 20139138.
87. Anderson EJ, Axon DR, Taylor AM, et al. Impact evaluation of a four-year academic-community partnership in provision of medication management and tertiary prevention services for rural patients with diabetes and/or hypertension. *Prev Med Rep*. 2020;17:101038. doi: 10.1016/j.pmedr.2019.101038. PMID: 31956473.
88. Boman K, Olofsson M, Berggren P, et al. Robot-assisted remote echocardiographic examination and teleconsultation: a randomized comparison of time to diagnosis with standard of care referral approach. *JACC Cardiovasc Imaging*. 2014 Aug;7(8):799-803. doi: 10.1016/j.jcmg.2014.05.006. PMID: 25124011.
89. Carter BL, Levy B, Gryzlak B, et al. Cluster-randomized trial to evaluate a centralized clinical pharmacy service in private family medicine offices. *Circ Cardiovasc Qual Outcomes*. 2018 06;11(6):e004188. doi: 10.1161/CIRCOUTCOMES.117.004188. PMID: 29884657.
90. Cho JH, Kwon HS, Kim HS, et al. Effects on diabetes management of a health-care provider mediated, remote coaching system via a PDA-type glucometer and the internet. *J Telemed Telecare*. 2011;17(7):365-70. doi: 10.1258/jtt.2011.100913. PMID: 21933896.
91. Crossland L, Askew D, Ware R, et al. Diabetic retinopathy screening and monitoring of early stage disease in Australian general practice: tackling preventable blindness within a chronic care model. *J Diabetes Res*. 2016;2016:8405395. doi: 10.1155/2016/8405395. PMID: 26798655.
92. Datta SK, Warshaw EM, Edison KE, et al. Cost and Utility Analysis of a Store-and-Forward Teledermatology Referral System: A Randomized Clinical Trial. *JAMA Dermatol*. 2015 Dec 1;151(12):1323-9. doi: 10.1001/jamadermatol.2015.2362. PMID: 26375589.
93. Davis TD, Deen T, Bryant-Bedell K, et al. Does minority racial-ethnic status moderate outcomes of collaborative care for depression? *Psychiatr Serv*. 2011 Nov;62(11):1282-8. doi: 10.1176/ps.62.11.pss6211_1282. PMID: 22211206.
94. De Batlle J, Massip M, Vargiu E, et al. Implementing mobile health-enabled integrated care for complex chronic patients: intervention effectiveness and cost-effectiveness study. *JMIR Mhealth Uhealth*. 2021 01 14;9(1):e22135. doi: 10.2196/22135. PMID: 33443486.

95. Fortney JC, Maciejewski ML, Tripathi SP, et al. A budget impact analysis of telemedicine-based collaborative care for depression. *Med Care*. 2011 Sep;49(9):872-80. doi: 10.1097/MLR.0b013e31821d2b35. PMID: 21623240.
96. Fortney JC, Pyne JM, Kimbrell TA, et al. Telemedicine-based collaborative care for posttraumatic stress disorder: a randomized clinical trial. *JAMA Psychiatry*. 2015 Jan;72(1):58-67. doi: 10.1001/jamapsychiatry.2014.1575. PMID: 25409287.
97. Fortney JC, Pyne JM, Mouden SB, et al. Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. *Focus*. 2017 Jul;15(3):361-72. doi: 10.1176/appi.focus.15306. PMID: 32015700.
98. Jiang B, Bills M, Poon P. Integrated telehealth-assisted home-based specialist palliative care in rural Australia: a feasibility study. *J Telemed Telecare*. 2020 Oct 20;1357633X20966466. doi: 10.1177/1357633X20966466. PMID: 33079611.
99. Kim H, Jhoo JH, Jang JW. The effect of telemedicine on cognitive decline in patients with dementia. *J Telemed Telecare*. 2017 Jan;23(1):149-54. doi: 10.1177/1357633X15615049. PMID: 26541171.
100. Kwak MY, Hwang EJ, Lee TH. Effects of the Physician-Primary-Healthcare Nurse Telemedicine Model (P-NTM) on medication adherence and Health-Related Quality of Life (HRQoL) of patients with chronic disease at remote rural areas. *Int J Environ Res Public Health*. 2021 03 03;18(5):03. doi: 10.3390/ijerph18052502. PMID: 33802513.
101. Levin K, Madsen JR, Petersen I, et al. Telemedicine diabetes consultations are cost-effective, and effects on essential diabetes treatment parameters are similar to conventional treatment: 7-year results from the Svendborg Telemedicine Diabetes Project. *J Diabetes Sci Technol*. 2013 May 1;7(3):587-95. doi: 10.1177/193229681300700302. PMID: 23759390.
102. Liddy C, Moroz I, Keely E, et al. Understanding the impact of a multispecialty electronic consultation service on family physician referral rates to specialists: a randomized controlled trial using health administrative data. *Trials*. 2019 Jun 10;20(1):348. doi: 10.1186/s13063-019-3393-5. PMID: 31182123.
103. Liou JK, Soon MS, Chen CH, et al. Shared care combined with telecare improves glycemic control of diabetic patients in a rural underserved community. *Telemed J E Health*. 2014 Feb;20(2):175-8. doi: 10.1089/tmj.2013.0037. PMID: 24320193.
104. Long MC, Angtuaco T, Lowery C. Ultrasound in telemedicine: its impact in high-risk obstetric health care delivery. *Ultrasound Q*. 2014 Sep;30(3):167-72. doi: 10.1097/RUQ.0000000000000073. PMID: 25148484.
105. McGill AF, North JB. Teleconference fracture clinics: a trial for rural hospitals. *ANZ J Surg*. 2012 Jan-Feb;82(1-2):2-3. doi: 10.1111/j.1445-2197.2011.05952.x. PMID: 22507484.
106. McLendon SF, Wood FG, Stanley N. Enhancing diabetes care through care coordination, telemedicine, and education: evaluation of a rural pilot program. *Public Health Nurs*. 2019 05;36(3):310-20. doi: 10.1111/phn.12601. PMID: 30868661.
107. Myers K, Vander Stoep A, Zhou C, et al. Effectiveness of a telehealth service delivery model for treating attention-deficit/hyperactivity disorder: a community-based randomized controlled trial. *J Am Acad Child Adolesc Psychiatry*. 2015 Apr;54(4):263-74. doi: 10.1016/j.jaac.2015.01.009. PMID: 25791143.

108. Painter JT, Fortney JC, Austen MA, et al. Cost-effectiveness of telemedicine-based collaborative care for posttraumatic stress disorder. *Psychiatr Serv.* 2017 Nov 01;68(11):1157-63. doi: 10.1176/appi.ps.201600485. PMID: 28669290.
109. Pyne JM, Fortney JC, Mouden S, et al. Cost-effectiveness of on-site versus off-site collaborative care for depression in rural FQHCs. *Psychiatr Serv.* 2015 May 01;66(5):491-9. doi: 10.1176/appi.ps.201400186. PMID: 25686811.
110. Pyne JM, Fortney JC, Tripathi SP, et al. Cost-effectiveness analysis of a rural telemedicine collaborative care intervention for depression. *Arch Gen Psychiatry.* 2010 Aug;67(8):812-21. doi: 10.1001/archgenpsychiatry.2010.82. PMID: 20679589.
111. Rojas G, Guajardo V, Martinez P, et al. A remote collaborative care program for patients with depression living in rural areas: open-label trial. *J Med Internet Res.* 2018 04 30;20(4):e158. doi: 10.2196/jmir.8803. PMID: 29712627.
112. Rossaro L, Torruellas C, Dhaliwal S, et al. Clinical outcomes of hepatitis C treated with pegylated interferon and ribavirin via telemedicine consultation in Northern California. *Dig Dis Sci.* 2013 Dec;58(12):3620-5. doi: 10.1007/s10620-013-2810-y. PMID: 24154637.
113. Sicotte C, Moqadem K, Vasilevsky M, et al. Use of telemedicine for haemodialysis in very remote areas: the Canadian First Nations. *J Telemed Telecare.* 2011;17(3):146-9. doi: 10.1258/jtt.2010.100614. PMID: 21303935.
114. Stevens G, Loh J, Kolbe J, et al. Comparison of recommendations for radiotherapy from two contemporaneous thoracic multidisciplinary meeting formats: co-located and video conference. *Intern Med J.* 2012 Nov;42(11):1213-8. doi: 10.1111/j.1445-5994.2012.02817.x. PMID: 22530597.
115. Summerhayes C, McGee JA, Cooper RJ, et al. Introducing leg ulcer telemedicine into rural general practice. *Wounds UK.* 2012;8(2):28-36.
116. Taylor-Gjevre R, Nair B, Bath B, et al. Addressing rural and remote access disparities for patients with inflammatory arthritis through video-conferencing and innovative inter-professional care models. *Musculoskelet* 2018 03;16(1):90-5. doi: 10.1002/msc.1215. PMID: 28967235.
117. Thaker DA, Monypenny R, Olver I, et al. Cost savings from a telemedicine model of care in northern Queensland, Australia. *Med J Aust.* 2013 Sep 16;199(6):414-7. PMID: 24033216.
118. Van der Pol M, McKenzie L. Costs and benefits of tele-endoscopy clinics in a remote location. *J Telemed Telecare.* 2010;16(2):89-94. doi: 10.1258/jtt.2009.090609. PMID: 20139140.
119. Whited JD, Warshaw EM, Kapur K, et al. Clinical course outcomes for store and forward teledermatology versus conventional consultation: a randomized trial. *J Telemed Telecare.* 2013 Jun;19(4):197-204. doi: 10.1177/1357633X13487116. PMID: 23666440.
120. Wood PR, Caplan L. Outcomes, satisfaction, and costs of a rheumatology telemedicine program: a longitudinal evaluation. *J Clin Rheumatol.* 2019 Jan;25(1):41-4. doi: 10.1097/RHU.0000000000000778. PMID: 30461466.
121. Yan LL, Gong E, Gu W, et al. Effectiveness of a primary care-based integrated mobile health intervention for stroke management in rural China (SINEMA): A cluster-randomized controlled trial. *PLoS Med.* 2021 04;18(4):e1003582. doi: 10.1371/journal.pmed.1003582. PMID: 33909607.
122. Fortney JC, Pyne JM, Edlund MJ, et al. A randomized trial of telemedicine-based collaborative care for depression. *J Gen Intern Med.* 2007 Aug;22(8):1086-93. doi: 10.1007/s11606-007-0201-9. PMID: 17492326.

123. Fortney JC, Pyne JM, Mouden SB, et al. Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. *Am J Psychiatry*. 2013 Apr;170(4):414-25. doi: 10.1176/appi.ajp.2012.12050696. PMID: 23429924.
124. Astarcioğlu MA, Sen T, Kilit C, et al. Time-to-reperfusion in STEMI undergoing interhospital transfer using smartphone and WhatsApp messenger. *Am J Emerg Med*. 2015 Oct;33(10):1382-4. doi: 10.1016/j.ajem.2015.07.029. PMID: 26299691.
125. Brunetti ND, Dell'Anno A, Martone A, et al. Prehospital ECG transmission results in shorter door-to-wire time for STEMI patients in a remote mountainous region. *Am J Emerg Med*. 2020 02;38(2):252-7. doi: 10.1016/j.ajem.2019.04.046. PMID: 31079977.
126. Brunetti ND, Di Pietro G, Aquilino A, et al. Pre-hospital electrocardiogram triage with tele-cardiology support is associated with shorter time-to-balloon and higher rates of timely reperfusion even in rural areas: data from the Bari- Barletta/Andria/Trani public emergency medical service 118 registry on primary angioplasty in ST-elevation myocardial infarction. *Europ Heart J Acute Cardiovasc Care*. 2014 Sep;3(3):204-13. doi: 10.1177/2048872614527009. PMID: 24604713.
127. Demaerschalk BM, Raman R, Ernstrom K, et al. Efficacy of telemedicine for stroke: pooled analysis of the Stroke Team Remote Evaluation using a Digital Observation Camera (STRoKE DOC) and STRoKE DOC Arizona telestroke trials. *Telemed J E Health*. 2012 Apr;18(3):230-7. doi: 10.1089/tmj.2011.0116. PMID: 22400970.
128. Dharmar M, Kuppermann N, Romano PS, et al. Telemedicine consultations and medication errors in rural emergency departments. *Pediatrics*. 2013 Dec;132(6):1090-7. doi: 10.1542/peds.2013-1374. PMID: 24276844.
129. Dharmar M, Romano PS, Kuppermann N, et al. Impact of critical care telemedicine consultations on children in rural emergency departments. *Crit Care Med*. 2013 Oct;41(10):2388-95. doi: 10.1097/CCM.0b013e31828e9824. PMID: 23921273.
130. Fairchild RM, Ferng-Kuo SF, Laws S, et al. Telehealth decreases rural emergency department wait times for behavioral health patients in a group of critical access hospitals. *Telemed J E Health*. 2019 12;25(12):1154-64. doi: 10.1089/tmj.2018.0227. PMID: 30735100.
131. Harvey JB, Yeager BE, Cramer C, et al. The impact of telemedicine on pediatric critical care triage. *Pediatr Crit Care Med*. 2017 Nov;18(11):e555-e60. doi: 10.1097/PCC.0000000000001330. PMID: 28922271.
132. Holt T, Sari N, Hansen G, et al. Remote presence robotic technology reduces need for pediatric interfacility transportation from an isolated northern community. *Telemed J E Health*. 2018 11;24(11):927-33. doi: 10.1089/tmj.2017.0211. PMID: 29394155.
133. Miller AC, Ward MM, Ullrich F, et al. Emergency department telemedicine consults are associated with faster time-to-electrocardiogram and time-to-fibrinolysis for myocardial infarction patients. *Telemed J E Health*. 2020 Feb 28;26(12):1440-8. doi: 10.1089/tmj.2019.0273. PMID: 32109200.
134. Miyamoto S, Dharmar M, Boyle C, et al. Impact of telemedicine on the quality of forensic sexual abuse examinations in rural communities. *Child Abuse Negl*. 2014 Sep;38(9):1533-9. doi: 10.1016/j.chiabu.2014.04.015. PMID: 24841062.
135. Mohr NM, Vakkalanka JP, Harland KK, et al. Telemedicine use decreases rural emergency department length of stay for transferred North Dakota trauma patients. *Telemed J E Health*. 2018 03;24(3):194-202. doi: 10.1089/tmj.2017.0083. PMID: 28731843.

136. Mohr NM, Young T, Harland KK, et al. Telemedicine is associated with faster diagnostic imaging in stroke patients: a cohort study. *Telemed J E Health*. 2019 02;25(2):93-100. doi: 10.1089/tmj.2018.0013. PMID: 29958087.
137. Mohr NM, Campbell KD, Swanson MB, et al. Provider-to-provider telemedicine improves adherence to sepsis bundle care in community emergency departments. *J Telemed Telecare*. 2020 Jan 05;1-9. doi: 10.1177/1357633X19896667. PMID: 31903840.
138. Mohr NM, Young T, Harland KK, et al. Emergency department telemedicine shortens rural time-to-provider and emergency department transfer times. *Telemed J E Health*. 2018 08;24(8):582-93. doi: 10.1089/tmj.2017.0262. PMID: 29293413.
139. Nagao KJ, Koschel A, Haines HM, et al. Rural Victorian telestroke project. *Intern Med J*. 2012 Oct;42(10):1088-95. doi: 10.1111/j.1445-5994.2011.02603.x. PMID: 21981424.
140. Nagayoshi Y, Oshima S, Ogawa H. Clinical impact of telemedicine network system at rural hospitals without on-site cardiac surgery backup. *Telemed J E Health*. 2016 11;22(11):960-4. PMID: 27148833.
141. Natafagi N, Mohr NM, Wittrock A, et al. The association between telemedicine and emergency department (ED) disposition: a stepped wedge design of an ED-based telemedicine program in critical access hospitals. *J Rural Health*. 2020 Jun;36(3):360-70. doi: 10.1111/jrh.12370. PMID: 31013552.
142. Pedragosa A, Alvarez-Sabin J, Rubiera M, et al. Impact of telemedicine on acute management of stroke patients undergoing endovascular procedures. *Cerebrovasc Dis*. 2012;34(5-6):436-42. doi: 10.1159/000345088. PMID: 23207552.
143. Sairanen T, Soinila S, Nikkanen M, et al. Two years of Finnish Telestroke: thrombolysis at spokes equal to that at the hub. *Neurology*. 2011 Mar 29;76(13):1145-52. doi: 10.1212/WNL.0b013e318212a8d4. PMID: 21368283.
144. Sterling SA, Seals SR, Jones AE, et al. The impact of the TelEmergency program on rural emergency care: an implementation study. *J Telemed Telecare*. 2017 Jul;23(6):588-94. doi: 10.1177/1357633X16657499. PMID: 27470505.
145. Swanson MB, Miller AC, Ward MM, et al. Emergency department telemedicine consults decrease time to interpret computed tomography of the head in a multi-network cohort. *J Telemed Telecare*. 2019 Nov 04;1-10. doi: 10.1177/1357633X19877746. PMID: 31684801.
146. Tripod M, Tait M, Bracey J, et al. The use of telemedicine decreases unnecessary hand trauma transfers. *Hand*. 2020 May;15(3):422-7. doi: 10.1177/1558944718810877. PMID: 30417694.
147. Vakkalanka JP, Harland KK, Wittrock A, et al. Telemedicine is associated with rapid transfer and fewer involuntary holds among patients presenting with suicidal ideation in rural hospitals: a propensity matched cohort study. *J Epidemiol Community Health*. 2019 11;73(11):1033-9. doi: 10.1136/jech-2019-212623. PMID: 31492762.
148. Bladin CF, Molocziej N, Ermel S, et al. Victorian Stroke Telemedicine Project: implementation of a new model of translational stroke care for Australia. *Intern Med J*. 2015 Sep;45(9):951-6. doi: 10.1111/imj.12822. PMID: 26011155.
149. Helwig SA, Ragoschke-Schumm A, Schwindling L, et al. Prehospital Stroke Management Optimized by Use of Clinical Scoring vs Mobile Stroke Unit for Triage of Patients With Stroke: A Randomized Clinical Trial. *JAMA Neurol*. 2019 Sep 3;76(12):1484-92. doi: 10.1001/jamaneurol.2019.2829. PMID: 31479116.
150. Williams D, Simpson AN, King K, et al. Do Hospitals Providing Telehealth in Emergency Departments Have Lower Emergency Department Costs? *Telemed J E Health*. 2020 Nov 13;13:13. doi: 10.1089/tmj.2020.0349. PMID: 33185503.

151. Arora S, Thornton K, Murata G, et al. Outcomes of treatment for hepatitis C virus infection by primary care providers. *N Engl J Med*. 2011 Jun 09;364(23):2199-207. doi: 10.1056/NEJMoa1009370. PMID: 21631316.
152. Bellesheim KR, Kizzee RL, Curran A, et al. ECHO autism: integrating maintenance of certification with extension for community healthcare outcomes improves developmental screening. *J Dev Behav Pediatr*. 2020 Aug;41(6):420-7. doi: 10.1097/DBP.0000000000000796. PMID: 32735419.
153. Bennett-Levy J, Hawkins R, Perry H, et al. Online cognitive behavioural therapy training for therapists: outcomes, acceptability, and impact of support. *Aust Psychol*. 2012;47(3):174-82. doi: 10.1111/j.1742-9544.2012.00089.x.
154. Bouchonville MF, Hager BW, Kirk JB, et al. Endo ECHO improves primary care provider and community health worker self-efficacy in complex diabetes management in medically underserved communities. *Endocr Pract*. 2018 Jan;24(1):40-6. doi: 10.4158/EP-2017-0079. PMID: 29368967.
155. Doyle C, Jackson D, Loi S, et al. Videoconferencing and telementoring about dementia care: evaluation of a pilot model for sharing scarce old age psychiatry resources. *Int Psychogeriatr*. 2016 09;28(9):1567-74. doi: 10.1017/S1041610216000740. PMID: 27189501.
156. Gadowski A, Anderson J, Chung YK, et al. Full agonist opioid prescribing by primary care clinicians after buprenorphine training. *Subst Abuse*. 2020 Mar 09;1-7. doi: 10.1080/08897077.2020.1736709. PMID: 32150525.
157. Gill CJ, Le Ngoc B, Halim N, et al. The mCME project: a randomized controlled trial of an SMS-based continuing medical education intervention for improving medical knowledge among Vietnamese community based physicians' assistants. *PLoS One* 2016;11(11):e0166293. doi: 10.1371/journal.pone.0166293. PMID: 27861516.
158. Hostutler CA, Valleru J, Maciejewski HM, et al. Improving pediatrician's behavioral health competencies through the Project ECHO teleconsultation model. *Clin Pediatr (Phila)*. 2020 Jun 07;59(12):1049-57. doi: 10.1177/0009922820927018. PMID: 32506939.
159. McFarland LV, Raugi GJ, Taylor LL, et al. Implementation of an education and skills programme in a teler dermatology project for rural veterans. *J Telemed Telecare*. 2012 Mar;18(2):66-71. doi: 10.1258/jtt.2011.110518. PMID: 22198956.
160. McWilliams T, Hendricks J, Twigg D, et al. Burns education for non-burn specialist clinicians in Western Australia. *Burns*. 2015 Mar;41(2):301-7. doi: 10.1016/j.burns.2014.06.015. PMID: 25239848.
161. Momin B, Mera J, Essex W, et al. Implementation of liver cancer education among health care providers and community coalitions in the Cherokee nation. *Prev Chronic Dis*. 2019;16(E112):1-7. doi: 10.5888/pcd16.180671. PMID: 31441770.
162. Puskar KR, Heeyoung L, Mitchell AM, et al. Interprofessional collaborative education for substance use screening: rural areas and challenges. *Online J Rural Nurs Health Care*. 2016;16(1):76-96. doi: 10.14574/ojrnhc.v16i1.385.
163. Robinson T, Hills D, Kelly B. The evaluation of an online orientation to rural mental health practice in Australia. *J Psychiatr Ment Health Nurs*. 2011;18(7):629-36. doi: 10.1111/j.1365-2850.2011.01712.x. PMID: 21848598.
164. Seibert PS, Reddy T, Whitmore T, et al. The use of telemedicine to train perioperative nurses in rural settings. *J Telemed Telecare*. 2013 Sep;19(6):311-4. doi: 10.1177/1357633X13501777. PMID: 24163294.
165. Shaikh U, Nettiksimmons J, Joseph JG, et al. Collaborative practice improvement for childhood obesity in rural clinics: the Healthy Eating Active Living Telehealth Community of Practice (HEALTH COP). *Am J Med Qual*. 2014 Nov-Dec;29(6):467-75. doi: 10.1177/1062860613506252. PMID: 24170936.

166. Sockalingam S, Arena A, Serhal E, et al. Building provincial mental health capacity in primary care: an evaluation of a Project ECHO mental health program. *Acad Psychiatry*. 2018 Aug;42(4):451-7. doi: 10.1007/s40596-017-0735-z. PMID: 28593537.
167. Watts SA, Roush L, Julius M, et al. Improved glycemic control in veterans with poorly controlled diabetes mellitus using a Specialty Care Access Network-Extension for Community Healthcare Outcomes model at primary care clinics. *J Telemed Telecare*. 2016 Jun;22(4):221-4. doi: 10.1177/1357633X15598052. PMID: 26253745.
168. Wilson BM, Banks RE, Crnich CJ, et al. Changes in antibiotic use following implementation of a telehealth stewardship pilot program. *Infect Control Hosp Epidemiol*. 2019 07;40(7):810-4. doi: 10.1017/ice.2019.128. PMID: 31172905.
169. Zittleman L, Curcija K, Sutter C, et al. Building capacity for medication assisted treatment in rural primary care practices: the IT MATTTRs practice team training. *J Prim Care Community Health*. 2020 Jan-Dec;11:2150132720953723. doi: 10.1177/2150132720953723. PMID: 32909491.
170. Lingum NR, Sokoloff LG, Meyer RM, et al. Building long-term care staff capacity during COVID-19 through just-in-time learning: evaluation of a modified ECHO model. *J Am Med Dir Assoc*. 2021 Feb;22(2):238-44.e1. doi: 10.1016/j.jamda.2020.10.039. PMID: 33238143.
171. Nipp CM, Vogtle LK, Warren M. Clinical application of low vision rehabilitation strategies after completion of a computer-based training module. *Occup Ther Health Care*. 2014 Jul;28(3):296-305. doi: 10.3109/07380577.2014.908335. PMID: 24971896.
172. Tran L, Feldman R, Riley T, 3rd, et al. Association of the extension for community healthcare outcomes project with use of direct-acting antiviral treatment among US adults with hepatitis C. *JAMA Netw Open*. 2021 Jul 01;4(7):e2115523. doi: 10.1001/jamanetworkopen.2021.15523. PMID: 34213557.
173. Lindauer A, Wild K, Natanson A, et al. Dementia 360 ECHO: using technology to facilitate diagnosis and treatment. *Gerontol Geriatr Educ*. 2020 Oct 20;1-7. doi: 10.1080/02701960.2020.1835658. PMID: 33078687.
174. Adcock AK, Choi J, Alvi M, et al. Expanding acute stroke care in rural America: a model for statewide success. *Telemed J E Health*. 2020 Jul;26(7):865-71. doi: 10.1089/tmj.2019.0087. PMID: 31596679.
175. Al Kasab S, Adams RJ, Debenham E, et al. Medical university of South Carolina telestroke: a telemedicine facilitated network for stroke treatment in South Carolina-a progress report. *Telemed J E Health*. 2017 08;23(8):674-7. doi: 10.1089/tmj.2016.0229. PMID: 28170316.
176. Alschuler KN, Stobbe GA, Hertz DP, et al. Impact of multiple sclerosis Project ECHO (extension for community healthcare outcomes) on provider confidence and clinical practice. *Int J MS Care*. 2019 Jul-Aug;21(4):143-50. doi: 10.7224/1537-2073.2018-014. PMID: 31474806.
177. Bagot KL, Cadilhac DA, Kim J, et al. Transitioning from a single-site pilot project to a state-wide regional telehealth service: the experience from the Victorian Stroke Telemedicine programme. *J Telemed Telecare*. 2017 Dec;23(10):850-5. doi: 10.1177/1357633X17734004. PMID: 29081268.
178. Banbury A, Smith AC, Mehrotra A, et al. A comparison study between metropolitan and rural hospital-based telehealth activity to inform adoption and expansion. *J Telemed Telecare*. 2021 Mar 26;1357633X21998201. doi: 10.1177/1357633X21998201. PMID: 33765879.
179. Becevic M, Smith E, Golzy M, et al. Melanoma extension for community healthcare outcomes: a feasibility study of melanoma screening implementation in primary care settings. *Cureus*. 2021 May 29;13(5):e15322. doi: 10.7759/cureus.15322. PMID: 34221770.

180. Brunet N, Moore DT, Lendvai Wischik D, et al. Increasing buprenorphine access for veterans with opioid use disorder in rural clinics using telemedicine. *Subst Abus.* 2020 Feb 20;1-8. doi: 10.1080/08897077.2020.1728466. PMID: 32078492.
181. Cadilhac DA, Vu M, Bladin C. Experience with scaling up the Victorian Stroke Telemedicine programme. *J Telemed Telecare.* 2014 Oct;20(7):413-8. doi: 10.1177/1357633X14552389. PMID: 25400003.
182. Conn DK, Madan R, Lam J, et al. Program evaluation of a telepsychiatry service for older adults connecting a university-affiliated geriatric center to a rural psychogeriatric outreach service in Northwest Ontario, Canada. *Int Psychogeriatr.* 2013 Nov;25(11):1795-800. doi: 10.1017/S104161021300118X. PMID: 23870297.
183. Cronin T. Implementing a stroke program using telemedicine. *J Emerg Nurs.* 2013 Nov;39(6):613-8. doi: 10.1016/j.jen.2012.04.012. PMID: 22748668.
184. Evans K, Lerch S, Boyce TW, et al. An innovative approach to enhancing access to medical screening for miners using a mobile clinic with telemedicine capability. *J Health Care Poor Underserved.* 2016;27(4A):62-72. PMID: 27818414.
185. Habashi P, Bouchard S, Nguyen GC. Transforming access to specialist care for inflammatory bowel disease: the PACE telemedicine program. *J Can Assoc Gastroenterol.* 2019 Dec;2(4):186-94. doi: 10.1093/jcag/gwy046. PMID: 31616860.
186. Haozous E, Doorenbos AZ, Demiris G, et al. Role of telehealth/videoconferencing in managing cancer pain in rural American Indian communities. *Psycho-Oncol.* 2012 Feb;21(2):219-23. doi: 10.1002/pon.1887. PMID: 22271543.
187. Haque SN, DeStefano S, Banger A, et al. Factors influencing telehealth implementation and use in frontier critical access hospitals: qualitative study. *JMIR Form Res.* 2021 May 05;5(5):e24118. doi: 10.2196/24118. PMID: 33949958.
188. Hensel JM, Yang R, Rai M, et al. Optimizing electronic consultation between primary care providers and psychiatrists: mixed-methods study. *J Med Internet Res.* 2018 04 06;20(4):e124. doi: 10.2196/jmir.8943. PMID: 29625949.
189. Hilt RJ, Barclay RP, Bush J, et al. A statewide child telepsychiatry consult system yields desired health system changes and savings. *Telemed J E Health.* 2015 Jul;21(7):533-7. doi: 10.1089/tmj.2014.0161. PMID: 25799043.
190. Hofmeyer J, Leider JP, Satorius J, et al. Implementation of telemedicine consultation to assess unplanned transfers in rural long-term care facilities, 2012-2015: a pilot study. *J Am Med Dir Assoc.* 2016 11 01;17(11):1006-10. doi: 10.1016/j.jamda.2016.06.014. PMID: 27477614.
191. Howland M, Tennant M, Bowen DJ, et al. Psychiatrist and psychologist experiences with telehealth and remote collaborative care in primary care: a qualitative study. *J Rural Health.* 2020 Oct 06;06:06. doi: 10.1111/jrh.12523. PMID: 33022079.
192. Jewer J, Parsons MH, Dunne C, et al. Evaluation of a mobile telesimulation unit to train rural and remote practitioners on high-acuity low-occurrence procedures: pilot randomized controlled trial. *J Med Internet Res.* 2019 08 06;21(8):e14587. doi: 10.2196/14587. PMID: 31389340.
193. Jhaveri D, Larkins S, Kelly J, et al. Remote chemotherapy supervision model for rural cancer care: perspectives of health professionals. *Eur J Cancer Care (Engl).* 2016 Jan;25(1):93-8. doi: 10.1111/ecc.12309. PMID: 25871852.
194. Johansson AM, Lindberg I, Soderberg S. The views of health-care personnel about video consultation prior to implementation in primary health care in rural areas. *Prim Health Care Res Dev.* 2014 Apr;15(2):170-9. doi: 10.1017/S1463423613000030. PMID: 23402617.

195. Katzman JG, Gygi K, Swift R, et al. How hands-on pain skills intensive trainings complement ECHO pain and opioid management programs: a program evaluation with the Indian Health Service. *Pain Med.* 2020 Jun 18;21(9):1769-78. doi: 10.1093/pm/pnaa151. PMID: 32556294.
196. Khalil H, Cullen M, Chambers H, et al. Implementation of a successful electronic wound documentation system in rural Victoria, Australia: a subject of collaboration and community engagement. *Int Wound J.* 2014 Jun;11(3):314-8. doi: 10.1111/iwj.12041. PMID: 23418740.
197. Kulcsar M, Gilchrist S, George MG. Improving stroke outcomes in rural areas through telestroke programs: an examination of barriers, facilitators, and state policies. *Telemed J E Health.* 2014 Jan;20(1):3-10. doi: 10.1089/tmj.2013.0048. PMID: 24286197.
198. Leshner AP, Fakhry SM, DuBose-Morris R, et al. Development and evolution of a statewide outpatient consultation service: leveraging telemedicine to improve access to specialty care. *Popul Health Manag.* 2020 02;23(1):20-8. doi: 10.1089/pop.2018.0212. PMID: 31161963.
199. Lim M, Liberali SAC, Calache H, et al. Specialist networks influence clinician willingness to treat individuals with special needs. *JDR Clin Trans Res.* 2021 Jun 21;23800844211020250. doi: 10.1177/23800844211020250. PMID: 34148391.
200. Lucas JAM, Day K, Honey MLL. Clinician's perceptions of telehealth for emergency care on the west coast of New Zealand: findings of a descriptive study. *Emergency Nurse New Zealand.* 2016:6-10.
201. Luscombe GM, Hawthorn J, Wu A, et al. 'Empowering clinicians in smaller sites': A qualitative study of clinician's experiences with a rural Virtual Paediatric Feeding Clinic. *Aust J Rural Health.* 2021 Sep 07;07:07. doi: 10.1111/ajr.12781. PMID: 34490941.
202. MacKinney AC, Ward MM, Ullrich F, et al. The business case for tele-emergency. *Telemed J E Health.* 2015 Dec;21(12):1005-11. doi: 10.1089/tmj.2014.0241. PMID: 26226603.
203. Mahmoud H, Vogt EL, Dahdouh R, et al. Using continuous quality improvement to design and implement a telepsychiatry program in rural Illinois. *Psychiatr Serv.* 2020 Apr 15;71(8):860-3. doi: 10.1176/appi.ps.201900231. PMID: 32290807.
204. Marsh-Feiley G, Eadie L, Wilson P. Paramedic and physician perspectives on the potential use of remotely supported prehospital ultrasound. *Rural Remote Health.* 2018 09;18(3):4574. doi: 10.22605/RRH4574. PMID: 30207737.
205. May S, Jonas K, Fehler GV, et al. Challenges in current nursing home care in rural Germany and how they can be reduced by telehealth - an exploratory qualitative pre-post study. *BMC Health Serv Res.* 2021 Sep 06;21(1):925. doi: 10.1186/s12913-021-06950-y. PMID: 34488746.
206. Moffatt JJ, Eley DS. Barriers to the up-take of telemedicine in Australia--a view from providers. *Rural Remote Health.* 2011;11(2):1581. PMID: 21385004.
207. Morrisette S, Pearlman RL, Kovar M, et al. Attitudes and perceived barriers toward store-and-forward teledermatology among primary care providers of the rural Mississippi. *Arch Dermatol Res.* 2021 Feb 25;25:25. doi: 10.1007/s00403-021-02208-z. PMID: 33630147.
208. Mundt AP, Irarrazaval M, Martinez P, et al. Telepsychiatry consultation for primary care treatment of children and adolescents receiving child protective services in Chile: mixed methods feasibility study. *JMIR Public Health Surveill.* 2021 Jul 22;7(7):e25836. doi: 10.2196/25836. PMID: 34292164.
209. Narva AS, Romancito G, Faber T, et al. Managing CKD by telemedicine: the Zuni telenephrology clinic. *Adv Chronic Kidney Dis.* 2017 Jan;24(1):6-11. doi: 10.1053/j.ackd.2016.11.019. PMID: 28224944.
210. Ness TE, Annese MF, Martinez-Paz N, et al. Using an innovative telehealth model to support community providers who deliver perinatal HIV care. *AIDS Educ Prev.* 2017 12;29(6):516-26. doi: 10.1521/aeap.2017.29.6.516. PMID: 29283274.

211. Newell MC, Strauss CE, Freier T, et al. Design and initial results of the Minneapolis Heart Institute TeleHeart program. *Circ Cardiovasc Qual Outcomes*. 2017;10:1-4. doi: 10.1161/CIRCOUTCOMES.117.003904. PMID: 29021334.
212. Nqala MO, Rout CC, Aldous CM. Remote clinical support by telephone for rural district hospital medical officers in the Eastern Cape. *S Afr Fam Pract*. 2015;57(5):286-90. doi: 10.1080/20786190.2015.1055671.
213. Olenik K, Lehr B. Counteracting brain drain of health professionals from rural areas via teleconsultation: analysis of the barriers and success factors of teleconsultation. *J Public Health*. 2013;21(4):357-64. doi: 10.1007/s10389-013-0565-8.
214. Pandit T, Ray RA, Sabesan S. Managing emergencies in rural North Queensland: the feasibility of teletraining. *Int J Telemed Appl*. 2018;2018doi: 10.1155/2018/8421346. PMID: 29849606.
215. Parks J, Hunter A, Taylor A, et al. Design, development and implementation of the virtual, coordination, access, referral and escalation service in western New South Wales. *Aust J Rural Health*. 2021 Sep 29;29:29. doi: 10.1111/ajr.12800. PMID: 34586702.
216. Peracca SB, Jackson GL, Lamkin RP, et al. Implementing teledermatology for rural veterans: an evaluation using the RE-AIM framework. *Telemed J E Health*. 2020 Apr 27;1-9. doi: 10.1089/tmj.2020.0013. PMID: 32343924.
217. Pimentel CB, Gately M, Barczy SR, et al. GRECC connect: geriatrics telehealth to empower health care providers and improve management of older veterans in rural communities. *Fed Pract*. 2019 Oct;36(10):464-70. PMID: 31768097.
218. Pindyck T, Kalishman S, Flatow-Trujillo L, et al. Treating hepatitis C in American Indians/Alaskan Natives: a survey of Project ECHO(R) (Extension for Community Healthcare Outcomes) utilization by Indian Health Service providers. *SAGE Open Med*. 2015;3:2050312115612805. doi: 10.1177/2050312115612805. PMID: 26770809.
219. Powell KR, Alexander GL. Consequences of rapid telehealth expansion in nursing homes: promise and pitfalls. *Appl Clin Inform*. 2021 Aug;12(4):933-43. doi: 10.1055/s-0041-1735974. PMID: 34614517.
220. Ray KN, Felmet KA, Hamilton MF, et al. Clinician attitudes toward adoption of pediatric emergency telemedicine in rural hospitals. *Pediatr Emerg Care*. 2017 Apr;33(4):250-7. doi: 10.1097/PEC.0000000000000583. PMID: 26785087.
221. Ritter LA, Robinette TR, Cofano J. Evaluation of a statewide telemedicine program. *CJHP*. 2010;8(1):1-9. doi: 10.32398/cjhp.v8i1.2025.
222. Rogove HJ, McArthur D, Demaerschalk BM, et al. Barriers to telemedicine: survey of current users in acute care units. *Telemed J E Health*. 2012 Jan-Feb;18(1):48-53. doi: 10.1089/tmj.2011.0071. PMID: 22082107.
223. Sabesan S, Senko C, Schmidt A, et al. Enhancing chemotherapy capabilities in rural hospitals: implementation of a telechemotherapy model (QReCS) in North Queensland, Australia. *J Oncol Pract*. 2018 07;14(7):e429-e37. doi: 10.1200/JOP.18.00110. PMID: 29996068.
224. Shea CM, Gertner AK, Green SL. Barriers and perceived usefulness of an ECHO intervention for office-based buprenorphine treatment for opioid use disorder in North Carolina: a qualitative study. *Subst Abus*. 2019 Dec 06:1-11. doi: 10.1080/08897077.2019.1694617. PMID: 31809679.
225. Simpson AN, Harvey JB, DiLembo SM, et al. Population health indicators associated with a statewide telestroke program. *Telemed J E Health*. 2020 Feb 11;26(9):1126-33. doi: 10.1089/tmj.2019.0204. PMID: 32045330.
226. Singh R, Mathiassen L, Switzer JA, et al. Assimilation of web-based urgent stroke evaluation: a qualitative study of two networks. *JMIR Med Inform*. 2014 Apr 15;2(1):e6. doi: 10.2196/medinform.3028. PMID: 25601232.

227. Thies KM, Anderson D, Beals-Reid C. Project ECHO chronic pain: a qualitative analysis of recommendations by expert faculty. *Pain Med.* 2019 07 01;20(7):1450-2. doi: 10.1093/pm/pnz024. PMID: 30840075.
228. Trondsen MV, Bolle SR, Stensland GO, et al. Video-confidence: a qualitative exploration of videoconferencing for psychiatric emergencies. *BMC Health Serv Res.* 2014 Oct 31;14:544. doi: 10.1186/s12913-014-0544-y. PMID: 25359404.
229. Trondsen MV, Tjora A, Broom A, et al. The symbolic affordances of a video-mediated gaze in emergency psychiatry. *Soc Sci Med.* 2018 01;197:87-94. doi: 10.1016/j.socscimed.2017.11.056. PMID: 29222999.
230. Tynan A, Deeth L, McKenzie D. An integrated oral health program for rural residential aged care facilities: a mixed methods comparative study. *BMC Health Serv Res.* 2018 07 03;18(1):515. doi: 10.1186/s12913-018-3321-5. PMID: 29970073.
231. Tynan A, Deeth L, McKenzie D, et al. Integrated approach to oral health in aged care facilities using oral health practitioners and teledentistry in rural Queensland. *Aust J Rural Health.* 2018 Apr 16;26:290-4. doi: 10.1111/ajr.12410. PMID: 29660771.
232. Ward MM, Ullrich F, MacKinney AC, et al. Tele-emergency utilization: in what clinical situations is tele-emergency activated? *J Telemed Telecare.* 2016 Jan;22(1):25-31. doi: 10.1177/1357633X15586319. PMID: 26026189.
233. Weaver MS, Neumann ML, Navaneethan H, et al. Human touch via touchscreen: rural nurses' experiential perspectives on telehealth use in pediatric hospice care. *J Pain Symptom Manage.* 2020 Jun 07;60(5):1027-33. doi: 10.1016/j.jpainsymman.2020.06.003. PMID: 32525081.
234. Weigel P, Bhagianadh D, Merchant KA, et al. Tele-emergency behavioural health in rural and underserved areas. *J Telemed Telecare.* 2019 Nov 14:1-10. doi: 10.1177/1357633X19887027. PMID: 31726903.
235. Weigel PA, Merchant KA, Wittrock A, et al. Paediatric tele-emergency care: a study of two delivery models. *J Telemed Telecare.* 2019 Apr 09:1-9. doi: 10.1177/1357633X19839610. PMID: 30966860.
236. White AH, Crowther SA, Lee SH. Supporting rural midwifery practice using a mobile health (mHealth) intervention: a qualitative descriptive study. *Rural Remote Health.* 2019 08;19(3):5294. doi: 10.22605/RRH5294. PMID: 31446762.
237. Williams KM, Kirsh S, Aron D, et al. Evaluation of the Veterans Health Administration's specialty care transformational initiatives to promote patient-centered delivery of specialty care: a mixed-methods approach. *Telemed J E Health.* 2017 07;23(7):577-89. doi: 10.1089/tmj.2016.0166. PMID: 28177858.
238. Wood T, Freeman S, Banner D, et al. Exploring user perspectives of factors associated with use of teletrauma in rural areas. *Australas Emerg Care.* 2021 May 07;07:07. doi: 10.1016/j.auec.2021.04.001. PMID: 33972192.
239. Zhu X, Merchant KAS, Mohr NM, et al. Real-time learning through telemedicine enhances professional training in rural emergency departments. *Telemed J E Health.* 2020 Jun 17;17:1-7. doi: 10.1089/tmj.2020.0042. PMID: 32552479.
240. CFIR Research Team-Center for Clinical Management Research. Consolidated Framework for Implementation Research. 2021. Accessed July 12, 2021.
241. Glasgow RE, Vogt TM, Boles SM. Evaluating the public health impact of health promotion interventions: the RE-AIM framework. *Am J Public Health.* 1999 Sep;89(9):1322-7. doi: 10.2105/ajph.89.9.1322. PMID: 10474547.
242. Dosaj A, Thiagarajan D, Ter Haar C, et al. Rapid Implementation of Telehealth Services During the COVID-19 Pandemic. *Telemed J E Health.* 2021 Feb;27(2):116-20. doi: 10.1089/tmj.2020.0219. PMID: 32706616.

243. Betancourt JA, Rosenberg MA, Zevallos A, et al. The Impact of COVID-19 on Telemedicine Utilization Across Multiple Service Lines in the United States. *Healthcare (Basel)*. 2020 Oct 1;8(4)doi: 10.3390/healthcare8040380. PMID: 33019667.
244. Patel SY, Mehrotra A, Huskamp HA, et al. Trends in outpatient care delivery and telemedicine during the COVID-19 Pandemic in the US. *JAMA Intern Med*. 2021 Mar 1;181(3):388-91. doi: 10.1001/jamainternmed.2020.5928. PMID: 33196765.
245. Bhaskar S, Nurtazina A, Mittoo S, et al. Editorial: Telemedicine during and beyond COVID-19. *Front Public Health*. 2021;9:662617. doi: 10.3389/fpubh.2021.662617. PMID: 33796502.
246. Weiner JP, Bandeian S, Hatef E, et al. In-person and telehealth ambulatory contacts and costs in a large US insured cohort before and during the COVID-19 pandemic. *JAMA Netw Open*. 2021 Mar 1;4(3):e212618. doi: 10.1001/jamanetworkopen.2021.2618. PMID: 33755167.
247. Edmunds M, Tuckson R, Lewis J, et al. An emergent research and policy framework for telehealth. *EGEMS (Wash DC)*. 2017;5(2):1303. doi: 10.13063/2327-9214.1303. PMID: 28459085.
248. National Quality Forum. Creating a Framework to Support Measure Development for Telehealth Department of Health and Human Services. Contract HHSM-500-2012-00009I, Task Order HHSM-500-T0022. Washington, DC. 2017.
249. Hollander J, Neinstein A. Maturation from adoption-based to quality-based telehealth metrics. *NEJM Catalyst*. 2020;1(5).
250. Curran GM, Bauer M, Mittman B, et al. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care*. 2012 Mar;50(3):217-26. doi: 10.1097/MLR.0b013e3182408812. PMID: 22310560.
251. Landes SJ, McBain SA, Curran GM. An introduction to effectiveness-implementation hybrid designs. *Psychiatry Res*. 2019 Oct;280:112513. doi: 10.1016/j.psychres.2019.112513. PMID: 31434011.

Abbreviations and Acronyms

Abbreviation	Definition
ADHD	Attention deficit hyperactivity disorder
AHRQ	Agency for Healthcare Research and Quality
CDC	Centers for Disease Control and Prevention
CFIR	Consolidated Framework for Implementation Research
CI	Confidence interval
CM	Centralized monitoring
COVID-19	Coronavirus disease 2019
CRP	c-reactive protein
ECG	electrocardiogram
ECHO	Extension for Community Healthcare Outcomes
ED	Emergency departments
EMS	Emergency medical services
FORHP	Federal Office of Rural Health Policy
ICU	Intensive care unit
MI	Myocardial infarction
MMSE	Mini-Mental Status Exam
NCATS	National Center for Advancing Translational Sciences
NHLBI	National Heart, Lung and Blood Institute
NICU	Neonatal intensive care unit
NIH	National Institutes of Health
ODP	Office of Disease Prevention
PICOS	Population, intervention, comparison, outcomes, settings and study designs of interest
PTSD	Post-traumatic stress disorder
RCT	Randomized controlled trial
RE-AIM	Reach, Effectiveness, Adoption, Implementation, and Maintenance
RR	Relative risk
SF-12	12-item short form survey
SF-36	36-item short form survey
SMS	Short messaging service
SOE	Strength of evidence
STEMI	ST-elevated myocardial infarction
TOO	Task Order Officer
tPA	Tissue plasminogen activator
VC	Virtual consult

Appendixes

Contents

Appendix A. Methods	A-1
Table A-1. PICOS and corresponding inclusion and exclusion criteria	A-4
Table A-2. Definitions of the grades of overall strength of evidence	A-9
Appendix B. Results	B-1
Figure B-1. Literature flow diagram	B-1
Table B-1. Characteristics of included studies for Key Question 1	B-2
Table B-2. Characteristics of included studies for Key Question 2	B-3
Table B-3. Characteristics of included studies for Key Question 3	B-4
Table B-4. Distribution of CFIR Constructs	B-6
Table B-5. Facilitators and barriers by topic area	B-7
Table B-6. Findings of inpatient implementation studies	B-8
Table B-7. Inpatient: barriers and facilitators by CFIR constructs	B-10
Table B-8. Findings of outpatient implementation studies	B-13
Table B-9. Outpatient: barriers and facilitators by CFIR constructs	B-17
Table B-10. Findings of telestroke and emergency care implementation studies	B-21
Table B-11. Emergency care: barriers and facilitators by CFIR constructs	B-23
Table B-12. Findings of education and mentoring implementation studies	B-26
Table B-13. Education/mentoring: barriers and facilitators by CFIR constructs	B-27
Appendix C. Included Studies List	C-1
Appendix D. Evidence Tables	D-1
Appendix E. Risk of Bias Assessment	E-1
Appendix F. Details on Strength of Evidence	F-1
Table F-1. Strength of Evidence of Selected Outcomes (Key Question 2)	F-1
Appendix G. Excluded Studies List	G-1
Appendix H. Appendix References	H-1

Appendix A. Methods

Details of Study Selection

Search Strategy

Literature Databases: Ovid MEDLINE®, CINAHL®, EMBASE, and Cochrane CENTRAL were searched to capture both published and gray literature. The search strategies were developed by a librarian with expertise in conducting searches for systematic reviews. Searches were peer reviewed by a second librarian.

Search Strategy: Search strategies are included below. In order to focus the search on rural applications of telehealth, we used three specific indexing terms (Rural Health Services, Rural Population, and Medically Underserved Area) as well as key word searches of titles and abstracts for “rural,” “remote,” and “resource limited.” Search and triage results were reviewed with the Technical Expert Panel (TEP) and NIH/ODP Working Group.

Publication Date Range: Searches identified studies published January 1, 2010 to October 12, 2021 for Key Questions 2, 3, and 4 and January 1, 2015 to October 12, 2021 for Key Question 1. This captured studies of systems that rely on more current technology. We included information on the dates the studies were conducted and the technologies used, as well the dates of publication.

Initial searches were conducted through September 2020. These searches were updated during the public comment period of the draft report to capture any new publications. Literature identified during the update search was assessed by following the same process of dual review as all other studies considered for inclusion in the report. New literature was identified for inclusion in the report, and was incorporated into the final submission of the report.

Supplemental Evidence and Data for Systematic review (SEADS): The AHRQ Evidence-based Practice Center (EPC) Scientific Resource Center notified stakeholders about the opportunity to submit information via the SEADS portal. There was also an announcement published in the Federal Register.

Gray Literature: Sources for gray (unpublished) literature included reports produced by government agencies, health care provider organizations, or others. With the help of AHRQ, we contacted the federal government community of practice on telehealth, and other appropriate organizations including, but not limited to the American Telemedicine Association, The Society for Education and the Advancement of Research in Connected Health (SEARCH) and AcademyHealth to make initial inquiries, and we also followed up on any suggestions made by TEP members.

Hand Searching: Reference lists of included articles, selected excluded articles (e.g., narrative reviews), and systematic reviews were reviewed for includable literature.

Contacting Authors: In the event that information regarding methods or results appeared to be omitted from the published results of a study, or if we were aware of unpublished data, we attempted to contact the authors to obtain additional information.

Search Strategies

Database: Ovid MEDLINE(R) ALL

1 Rural Health Services/

2 Rural Population/

3 (rural or "resource limited" or (remote adj5 (population* or community or communities or area*))).ti,ab,kf.

4 Medically Underserved Area/

5 exp Community Health Services/ or Community Health Centers/ or United States Indian Health Service/

6 "Referral and Consultation"/

7 exp Health Services Accessibility/

8 "Delivery of Health Care"/

9 2 or 3

10 or/4-8

11 9 and 10

12 1 or 11

13 exp Telemedicine/

14 Mobile Applications/

15 exp Telecommunications/

16 (telemedicine or telemedical or telehealth or teleconsult* or "e-consult*" or "econsult*" or ((remote or virtual) adj3 (consult* or education or round* or mentor*))).ti,ab,kf.

17 or/13-16

18 9 and 17

19 12 and 17

20 "provider to provider".ti,ab,kf.

21 ("Project ECHO" or "Extension for Community Healthcare Outcomes").ti,ab,kf.

22 or/18-21

23 "journal of telemedicine & telecare".jn.

24 "telemedicine journal & e-health".jn.

25 9 and (23 or 24)

26 "journal of rural health".jn.

27 17 and 26

28 22 or 25 or 27

29 limit 28 to english language

30 limit 29 to yr="2010 - 2021"

Database: EBM Reviews - Cochrane Central Register of Controlled Trials

1 Rural Health Services/

2 Rural Population/

3 (rural or "resource limited" or (remote adj5 (population* or community or communities or area*))).ti,ab.

4 Medically Underserved Area/

5 exp Community Health Services/

6 "Referral and Consultation"/

7 exp Health Services Accessibility/

8 "Delivery of Health Care"/

9 2 or 3

10 or/4-8

11 9 and 10

12 1 or 11
 13 exp Telemedicine/
 14 Mobile Applications/
 15 exp Telecommunications/
 16 (telemedicine or telemedical or telehealth or teleconsult* or "e-consult*" or "econsult*" or ((remote or virtual) adj3 (consult* or education or round* or mentor*))).ti,ab.
 17 or/13-16
 18 9 and 17
 19 12 and 17
 20 "provider to provider".ti,ab.
 21 ("Project ECHO" or "Extension for Community Healthcare Outcomes").ti,ab.
 22 or/18-21
 23 conference abstract.pt.
 24 "journal: conference abstract".pt.
 25 "journal: conference review".pt.
 26 "http://.www.who.int/trialsearch*".so.
 27 "https://clinicaltrials.gov*".so.
 28 23 or 24 or 25 or 26 or 27
 29 22 not 28

Database: EBM Reviews - Cochrane Database of Systematic Reviews

1 (telemedicine or telemedical or telehealth or teleconsult* or "e-consult*" or "econsult*" or ((remote or virtual) adj3 (consult* or education or round* or mentor*))).ti,ab.

Database: EBSCOhost CINAHL Plus

S1 (MH "Rural Health Personnel") OR (MH "Rural Health Centers") OR (MH "Hospitals, Rural") OR (MH "Rural Population") OR (MH "Rural Health Services") OR (MH "Rural Health Nursing") OR (MH "Rural Areas") OR (MH "Rural Health") OR (MH "Frontier Nursing Service")

S2 (MH "Telecommunications+")

S3 S1 and S2

S4 Limiters - Published Date: 20100101-20201231; English Language; Exclude MEDLINE records

Database: Elsevier Embase

('rural population'/exp OR 'rural population' OR 'rural area'/exp OR 'rural area' OR 'medically underserved'/exp OR 'medically underserved') AND ('telehealth'/exp OR telehealth OR 'telemedicine'/exp OR telemedicine OR 'teleconsultation'/exp OR teleconsultation) AND [embase]/lim NOT ([embase]/lim AND [medline]/lim) AND [2010-2020]/py

Inclusion and Exclusion Criteria

The criteria for inclusion and exclusion of individual studies are based on the Key Questions and PICOS described above. Additional details on the scope of this project are provided below and the inclusion and exclusion criteria are outlined in **Table A-1**.

Study Designs: We included comparative studies of any design including comparative trials and observational studies. We included observational cohort studies, including pre-post designs (i.e., the same participants compared across time points) as well as before-after designs (i.e., one group of participants before an intervention/system change compared to a different group after the change). We excluded descriptive studies with no outcomes data or studies that included only data from one point in time (post only). We also excluded modeling studies or studies that used synthetic data. We accessed existing systematic reviews, and reviewed reference lists to identify studies. We also excluded commentaries, letters, and articles that described telehealth systems or implementations but did not assess impact.

Outcomes: The protocol specified that we would include outcomes only for the following conditions: substance abuse/alcohol, HIV/HPV/other infectious diseases, suicide, heart disease, cancer, unintentional injury, chronic lower respiratory disease, and stroke. However, due to a limited body of evidence in some of these conditions, and additional evidence in other conditions, it was decided to include outcomes of interest for any health condition.

Non-English-Language Studies: We restricted to English-language articles, but reviewed English-language abstracts of non-English language articles to identify studies that would otherwise meet inclusion criteria, to assess for the likelihood of language bias.

Table A-1. PICOS and corresponding inclusion and exclusion criteria

PICOS	Include	Exclude
Population	Individual patients and partners of any age and populations with health care needs in geographically rural areas regardless of where the providers are located Rural is broadly defined; any commonly used or endorsed definition is acceptable Providers (clinicians broadly defined or health care organizations) of health care to patients and populations in rural areas Payers (public or private; insurers or self-pay) for health care provided to patients and populations in rural areas	Urban patients or populations Mixed patients and populations that are not separated and predominately urban Interactions between a formal provider and informal/family care partners/givers
Interventions	Provider-to-provider telehealth broadly defined as any form of interactive support using telecommunications technology provided to health care professionals while they are caring for rural patients and populations. This includes: Remote consultations across space (e.g., video) and time (e.g., store and forward) that support diagnosis, treatment, or management of patients Video, audio, or digital only consultations Remote mentoring Remote rounds or group education and case review (e.g., Project ECHOs) Remote continuing education	Use of telehealth for patient encounters involving one clinician (virtual visits) Remote patient monitoring (transmitting data from patient to a single provider) Referrals for services that involve no interaction among providers
Comparators	No service or support Care provided without telehealth In-person activities Time period prior to provider-to-provider telehealth implementation Excluded types of telehealth	KQ2, KQ4: No comparators (e.g., descriptive and cross-sectional studies) KQ1: None

PICOS	Include	Exclude
Outcomes	<p>KQ2, KQ4: Clinical outcomes such as patient-reported outcomes, mortality, morbidity, function, illness recovery, infection rates, or viral load for the identified conditions* Economic outcomes such as return on investment, cost, volume of visits, and resource use Intermediate outcomes such as: Patient satisfaction, behavior, and decisions such as completion of treatment, or satisfaction with less travel to access healthcare Provider satisfaction, behavior, and decisions such as choice of treatment or antibiotic stewardship Time to diagnosis, time to treatment, length of stay (if applicable), 30 days hospitalization Appropriate utilization of health care services and avoided, preventable hospitalizations/readmissions/ED visits/test, treatments, procedures, etc.</p> <p>KQ1: Indicators and measures of uptake of telehealth (e.g., rates of use, timing to implementation) and characteristics of users KQ3: categories and descriptors of barriers and facilitators of telehealth; categories and descriptors of strategies of use of telehealth</p>	<p>KQ2, KQ4: Results of models, simulations, or projections without actual outcomes data Results of cross-sectional studies and surveys that include no comparison (e.g., descriptive statistics) Results from surveys of attitudes or opinions about hypothetical scenarios (i.e., not actual experience) Diagnostic concordance or accuracy or other measures of agreement between in-person and telehealth consultations</p>
Settings	<p>Outpatient (primary care and specialty care) Inpatient (e.g., remote ICU, consultations for hospitalized patients) Prehospital and emergency care (e.g., Telestroke, EMS, ED, urgent care) Post-acute and long-term care (e.g., home care and nursing homes) Studies of health care services delivered outside of health care settings (e.g., social services, churches, schools, prisons)</p> <p>Civilian, Veterans Health Administration, or Military (except battlefield)</p> <p>United States and other countries with similar or more advanced health care systems and resources</p>	<p>Mass casualty and war/battlefield events</p> <p>Countries with significantly different healthcare systems and fewer resources (e.g., low-income countries)</p>
Study types and designs	<p>KQ2, KQ4: Comparative studies including trial and observational studies, including prospective and retrospective cohort studies and before-after studies (i.e., natural experiments)</p> <p>KQ1, KQ3: Comparative or descriptive studies</p>	<p>All: Nonsystematic reviews, commentaries, or letters KQ2, KQ4: Descriptive studies, feasibility assessments</p>
Study years	<p>Published in 2010 or later. Included systematic reviews may include studies prior to 2010 if such studies are relevant to current technology</p> <p>For KQ1: published in 2015 or later</p>	<p>Published prior to 2010 For KQ1: Published prior to 2015</p>

Abbreviations: ED = emergency department; EMS = emergency medical services; ICU = intensive care unit; KQ = Key Question; PICOS = population, intervention, comparator, outcomes, study designs; Project ECHO = Extension for Community Healthcare Outcomes

*Originally noted that only studies which examined outcomes for the following conditions would be included: substance abuse/alcohol, HIV/HPV/other infectious diseases, suicide, heart disease, cancer, unintentional injury, chronic lower respiratory disease, and stroke. However, after review of available body of literature we included any health condition.

Process for Selecting Studies: Pre-established criteria as presented in Table A-1 was used to determine eligibility for inclusion and exclusion of abstracts in accordance with the AHRQ *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.¹ After de-duplication,

we imported all references to DistillerSR for managing abstract and full-text review. To ensure accuracy, all excluded abstracts were dual reviewed. Full text was retrieved for all citations deemed appropriate for inclusion by at least one of the reviewers. All potentially relevant full-text articles were independently reviewed for eligibility by two team members. Any disagreements were resolved by consensus. A flow diagram of study screening and inclusion is below in Appendix B, and a record of studies included in the review and those excluded at the full-text level with reasons for exclusion can be found in Appendix C and G, respectively.

Data Extraction

After studies were deemed to meet inclusion criteria, we abstracted study design, year, setting, country, sample size, patient and providers types and characteristics (e.g., age, sex, race, reason for presentation, diagnosis, and provider specialty), intervention characteristics (e.g., mode of delivery, duration or frequency, function) and results relevant to each Key Question as outlined in the PICOS section in **Table A-1**. As Key Question 2 asks about outcomes for three groups: patients and providers, healthcare providers, and payers, outcomes were sorted by who was impacted. Information relevant for assessing applicability included the number of patients randomized/eligible for inclusion in an observational study relative to the number of patients enrolled or the number and diversity of settings or locations as well characteristics of the population, telehealth intervention or implementation strategy, and administering personnel. Sources of funding for all studies were also recorded. All study data was extracted into Excel and verified for accuracy and completeness by a second team member.

Risk of Bias Assessment of Individual Studies

Predefined criteria were used to assess the risk of bias (also referred to as quality or internal validity) for each individual included study, using criteria appropriate for the study designs. Risk of bias assessment was managed using DistillerSR. Controlled trials and observational studies were assessed using a priori established criteria consistent with the approach recommended in the chapter, *Assessing the Risk of Bias of Individual Studies When Comparing Medical Interventions* in the *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.¹ For controlled trials we assessed adequacy of randomization and allocation concealment, eligibility criteria, baseline differences between groups, intention-to-treat analyses, attrition and adherence levels, blinding methods, reliable and consistently implemented outcome measures, and prespecified and reported outcomes. For observational cohort studies, we assessed eligibility criteria, participant selection, baseline differences between groups, reliable and consistently implemented outcome measures, blinding of outcome assessors or data analysts, amount and handling of missing data, loss-to-follow up and attrition, and prespecified and reported outcomes. We adapted criteria and details for evaluating cost effectiveness studies from Consensus Health Economic Criteria-CHEC List.² We assessed whether competing alternatives were clearly described, appropriateness of study design, whether important and relevant costs for each alternative were identified, appropriateness of costs measured and valued, identified relevant outcomes for each alternative, outcomes measured and valued appropriately, whether an incremental analysis of costs and outcomes of alternatives was performed, whether all future costs and outcomes were discounted, and all important variables whose values are uncertain were subjected to sensitivity analysis. Individual studies were rated as “low risk of bias,” “medium risk of bias,” or “high risk of bias.” Risk of bias ratings can be seen in **Appendix E**.

Studies rated “low risk of bias” are considered to have the least risk of bias, and their results are generally considered valid. “Low risk of bias” studies include clear descriptions of the population, setting, interventions, and comparison groups; a valid method for allocation of patients to treatment; low dropout rates and clear reporting of dropouts; appropriate means for preventing bias; and appropriate measurement of outcomes.

Studies rated “medium risk of bias” are susceptible to some bias, though not enough to invalidate the results. These studies may not meet all the criteria for a rating of low risk of bias, but no flaw is likely to cause major bias. The study may be missing information, making it difficult to assess limitations and potential problems. The “medium risk of bias” category is broad, and studies with this rating will vary in their strengths and weaknesses. The results of some medium risk of bias studies are likely to be valid, while others may be only possibly valid.

Studies rated “high risk of bias” have significant flaws that imply biases of various types that may invalidate the results. They have a serious or “fatal” flaw in design, analysis, or reporting; large amounts of missing information; discrepancies in reporting; or serious problems in the delivery of the intervention. In general, observational studies that do not perform adjustment for potential confounders will be assessed as “high risk of bias.” This is because it is likely the results of these studies are at least as likely to reflect flaws in the study design as the true difference between the compared interventions. We did not exclude studies rated high risk of bias a priori, but high risk of bias studies were considered to be less reliable than low or medium risk of bias studies when synthesizing the evidence, particularly if discrepancies between studies were present.

Two team members independently reviewed included studies, and resolved disagreements by consensus.

Data Synthesis and Analysis

We constructed evidence tables identifying the study characteristics (as discussed above), results of interest, and risk of bias ratings for all included studies, and summary tables to highlight the main findings. As the key questions varied in nature and scope, our approach to synthesis differed.

For the comparative effectiveness question (KQ2), we applied standard systematic review methods. We reviewed and highlighted studies using a hierarchy-of-evidence approach, where the studies with better risk of bias ratings were given more weight in our synthesis for each clinical indication and outcome. Qualitative data were summarized in tables and ranges. Descriptive analysis and interpretation of the results were provided based on the direction and magnitude of effect. Meta-analyses would not produce meaningful results and were not performed due to limited numbers of studies reporting similar outcomes, and heterogeneity among studies in design, patient population, and interventions.

Our response to KQ3 involved identifying and summarizing barriers and facilitators to implementation of provider-to-provider telehealth for rural healthcare. When studies directly compared different strategies, we provide a narrative summary of the studies and their results. When studies only reported on implementation for an individual study (not comparative) we abstracted what was study authors described as barriers, facilitators, and indicators of implementation success. Facilitators and barriers were consistently described in narratives reporting results or discussing implications. These lacked common vocabulary and definitions. To address this and facilitate synthesis, we first abstracted what was reported in each study. Noting conceptual similarities, but different nomenclature across articles, we sought to group

facilitators and barriers into meaningful and informative categories. We selected and used the Consolidated Framework for Implementation Research (CFIR)³ to organize this information. CFIR is composed of 5 categories and 39 defined constructs, designed to aggregate facilitators and barriers into common groupings. Two members of the study team independently matched abstracted descriptions to a CFIR construct using the CFIR code book template available at <https://cfirguide.org/>. Independent coding was followed by collaborative review and arrival at consensus among three members of the study team. Descriptive analysis of study findings and CFIR themes were presented in figure and tables.

For KQ1 (use of telehealth) and KQ4 (strengths and weaknesses of included studies), synthesis consists of descriptive narrative and tables, corresponding to the nature of the questions and data.

Grading the Strength of the Body of Evidence

Key Question 1 (the uptake of telehealth in rural areas) is descriptive and a formal strength of evidence (SOE) assessment was not conducted. We prioritized reports of U.S. national or regional studies over local reports or data from other countries. We summarized the strengths and limitations of the data collection and analyses of the included reports for Key Question 1, with a focus on elements such as the extent the sample represents the population of interest and the completeness and reliability of the data.

The strength of evidence (SOE) for Key Question 2 was assessed using the approach described in the *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.¹ Outcomes were prioritized for SOE assessment based on input from the ODP working group. SOE was initially assessed by one researcher for each outcome. To ensure reliability and validity of the evaluation, the body of evidence was assessed for the following criteria as they are defined in the Methods Guide:

- Study limitations (low, medium, or high level of study limitations)
 - Rated as the degree to which studies for a given outcome are likely to reduce bias with study design and study conduct, based on risk of bias assessments.
- Consistency (consistent, inconsistent, or unknown/not applicable)
 - Rated by degree to which studies find similar magnitude of effect (i.e., range sizes are similar) or same direction (i.e., effect sizes have the same sign) or where there was only one study of a given design, we assessed consistency as “unknown” and downgraded the SOE.
- Directness (direct or indirect)
 - Rated by degree to which evidence assesses a) comparison of interest, b) in the population of interest, and measures the specific outcome of interest.
- Precision (precise or imprecise)
 - Degree of certainty surrounding an effect estimate as it relates to a specific outcome. This may be based on sufficiency of sample size and number of events, and if these are adequate, the interpretation of the confidence interval.
- Publication bias (suspected or undetected)
 - Whether selective publishing of research findings based on favorable direction or magnitude of effects was identified using funnel plots or statistical methods, however, we did not have enough studies to conduct this assessment, so the majority of SOE assessments rated this domain as “unknown.”

The bodies of evidence were assigned an overall SOE grade of high, moderate, low, or insufficient according to a four-level scale by evaluating and weighing the combined results of the above domains (**Table A-2**). Because studies were anticipated to be heterogeneous in the interventions, clinical targets, and outcomes, we did not anticipate that meta-analysis would be possible. As such, the conclusion of findings being similar were based on individual studies not finding statistically significant differences, with consistency across multiple studies in this finding, and that the point estimates were not subjectively viewed as being large.

Table A-2. Definitions of the grades of overall strength of evidence⁴

Grade	Definition
High	We are very confident that the estimate of effect lies close to the true effect for this outcome. The body of evidence has few or no deficiencies. We believe that the findings are stable (i.e., another study would not change the conclusions).
Moderate	We are moderately confident that the estimate of effect lies close to the true effect for this outcome. The body of evidence has some deficiencies. We believe that the findings are likely to be stable, but some doubt remains.
Low	We have limited confidence that the estimate of effect lies close to the true effect for this outcome. The body of evidence has major or numerous deficiencies (or both). We believe that additional evidence is needed before concluding either that the findings are stable or that the estimate of effect is close to the true effect.
Insufficient	We have no evidence, we are unable to estimate an effect, or we have no confidence in the estimate of effect for this outcome. No evidence is available, or the body of evidence has unacceptable deficiencies, precluding reaching a conclusion.

Following the Agency for Healthcare and Quality (AHRQ) Methods Guidance for bodies of evidence consisting of observational studies, the strength rating starts at moderate for harms outcomes, and low for benefit outcomes. Although this allows this evidence to be upgraded under specific circumstances, including if all or most of the studies are low risk of bias and they report consistent, precise estimates. However, this was not the case in this review so no upgrades were made. In cases where there were both RCTs and observational studies were included for a given intervention-outcome pair, we followed the additional guidance on how to weight RCTs over observational studies, how to assess consistency across the two bodies of evidence, and how to come to a final rating.

Given our broad inclusion criteria, the evidence for Key Question 3 consists of studies that used qualitative methods (e.g., interviews, case studies, focus groups) as well as program evaluations that combined qualitative and quantitative information. Very few of the studies were comparative and they are heterogeneous in terms of topics and methods. We considered the GRADE-CERQual⁵ approach to grading qualitative research synthesis to rate our finding for Key Question 3, but the studies included mixed methods, and evaluation studies that were often descriptive or did not adhere to standard qualitative methods. As a result, we were not able to apply this relatively new approach rigorously and reliably.

Assessing Applicability

Applicability was considered according to the approach described in the *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.¹ We used the PICOS framework to consider the applicability of the evidence base for each Key Question, for example, examining the characteristics of the patient populations (e.g., clinical condition) and study setting (e.g., inpatient or outpatient). Variability in the studies may limit the ability to generalize the results to other populations and settings.

Peer Review and Public Commentary

Peer reviewers were invited to provide written comments on the draft report based on their clinical, content, or methodological expertise. The EPC considers all peer review comments on the draft report in preparation of the final report. Peer reviewers do not participate in writing or editing of the final report or other products. The final report does not necessarily represent the views of individual reviewers.


Potential Peer Reviewers must disclose any financial conflicts of interest greater than \$5,000 and any other relevant business or professional conflicts of interest. Invited Peer Reviewers may not have any financial conflict of interest greater than \$5,000. Peer reviewers who disclose potential business or professional conflicts of interest may submit comments on draft reports through the public comment mechanism.

A draft report revised to address comments from peer reviewers was posted for four weeks for public comment. The EPC completed a disposition of comments document containing a summary of peer reviewer comments and author responses to the original draft report as well as point-by-point responses to all public comments on the revised draft report.

Appendix B. Results

Results of Literature Searches

Figure B-1. Literature flow diagram



Note: Five papers are included for more than one Key Question

A total of 6,329 references were identified from electronic database searches. After dual review of abstracts, 1,024 articles were evaluated for inclusion. Search results and selection of studies are summarized in the literature flow diagram above (**Figure B-1**). A total of 166 studies (in 179 publications) were included for at least one key question.⁶⁻¹⁸⁴ Seven studies were included for Key Question 1, 97 studies in 106 publications were included for Key Question 2, and 67 studies in 71 publications were included for Key Question 3. A list of included studies appears in **Appendix C** and excluded studies with reason for exclusion in **Appendix G**.

Description of Included Studies

Key Question 1

The systematic review protocol and a request for unpublished information was posted by AHRQ on the Federal Register Supplemental Evidence and Data (SEADs) webpage. Additionally, we sent emails requesting information to individual federal agencies as well as non governmental organizations involved in telehealth and experts familiar with telehealth practices and policy. Specific program offices contacted included FedTel, the U.S. Federal government working group on Telehealth, the Telehealth Focused Rural Health Research Center Program of the Health Resources and Services Administration (HRSA), and the COVID-19 Telehealth Impact Study organized by the COVID-19 Healthcare Coalition Telehealth Impact Study Work Group with leadership from Mayo Clinic and the MITRE Corporation.

We also explored the possibility of identifying trends through claims data. Although the Centers for Medicare & Medicaid Services has approved Common Procedure Terminology (CPT) codes 99446-99449, 99451 and 99452 for interprofessional electronic assessment and management and referral services provided by a consultative physician, or other qualified health care professional (QHP) and by a patient's treating/requesting physician/QHP (in the case of 99452), there is anecdotal evidence that use of these billing codes is very low (personal communications).¹⁸⁵ It is very likely that informal interprofessional consultations are occurring in a non-compensated manner, but such interactions would not be included in billing records, and literature describing the frequency of informal interprofessional consultations is not currently available.

We did not receive any additional unpublished evidence on provider-to-provider telehealth in the rural U.S. usable for this report. While use of telehealth for patient and provider interactions has been documented, particularly the increase as part of the response to the COVID-19 pandemic,^{186, 187} trends in provider-to-provider telehealth have not yet been documented to the same extent. Details can be found in **Appendix Table D-1**.

Table B-1. Characteristics of included studies for Key Question 1

Characteristic	Categories	Number of Articles - 6	Percentage of Articles	References
Location	Nationwide	6	86%	8, 30, 50, 149, 180, 182
	Regional- New England	1	14%	181
Adoption Category	Use	7	100%	8, 30, 50, 149, 180-182
	Spread	2	29%	149, 182
	Mental Health	2	29%	50, 149
	Emergency Care	2	29%	180, 181
	Stroke	3	43%	8, 30, 182

Key Questions 2 and 4

Study details can be found in **Appendix Tables D-2, D-3, D-4, D-5, D-6, D-7, D-8, D-9, E-1, E-2, E-3, and E-4**.

Table B-2. Characteristics of included studies for Key Question 2

Characteristic	Categories	Number of Articles - 106	Percentage of Articles	References
Geographic Location	United States	63	59%	10, 11, 18, 21, 23, 29, 36, 38-41, 44-49, 51, 55, 58-60, 66, 72, 75, 78, 84, 92, 93, 96, 98, 100, 101, 103-107, 110, 114, 117, 120, 130-132, 139, 142, 143, 150, 153, 154, 158, 159, 164, 167, 173, 175, 176, 178, 182, 184, 35, 172
	Australia	15	14%	13, 19, 20, 27, 34, 42, 52, 70, 97, 99, 111, 122, 128, 136, 156
	Canada	5	5%	65, 82, 85, 145, 148, 155
	United Kingdom	3	3%	81, 152, 165
	Korea	3	3%	31, 76, 79
	Italy	2	2%	25, 26
	Countries with a single study*	14	13%	12, 16, 22, 37, 53, 61, 81, 86, 112, 124, 138, 141, 151, 179
Study Design	RCT	23	22%	19, 22, 29, 31, 35, 36, 38, 45, 47-49, 53, 61, 82, 86, 110, 131, 132, 155, 172, 179, 184
	Observational-before/after	25	24%	13, 16, 20, 27, 41, 42, 44, 52, 55, 75, 78, 87, 105, 111, 112, 114, 120, 122, 128, 143, 150, 152, 154, 159, 178
	Observational-pre/post	18	17%	10, 18, 23, 66, 79, 84, 85, 96, 98, 99, 107, 117, 130, 136, 145, 148, 167, 175
	Observational-prospective cohort	21	20%	11, 12, 25, 26, 34, 37, 59, 60, 65, 70, 76, 93, 100, 103, 106, 124, 138, 141, 142, 153, 176
	Observational-retrospective cohort	19	18%	21, 39, 40, 51, 58, 72, 81, 92, 97, 101, 104, 139, 151, 156, 158, 164, 165, 173, 182
Risk of Bias	Low	5	5%	38, 82, 120, 132, 148
	Medium	75	71%	10-12, 16, 19-21, 25-27, 29, 31, 34-37, 39-41, 44-49, 51, 53, 58-61, 65, 70, 72, 76, 85, 86, 92, 93, 100, 101, 103-105, 110-112, 114, 117, 122, 124, 128, 131, 138, 139, 141-143, 145, 150, 151, 153-156, 158, 159, 164, 172, 173, 175, 176, 179, 182, 184
	High	26	25%	13, 18, 22, 23, 42, 52, 55, 66, 75, 78, 79, 81, 84, 87, 96-99, 106, 107, 130, 136, 152, 165, 167, 178
Sample Size	Under 100	30	28%	18, 19, 22, 23, 25, 31, 37, 52, 66, 70, 81, 84, 86, 96-99, 111, 112, 117, 136, 139, 142, 145, 152, 155, 165, 167, 176, 184
	100-500	48	45%	11, 12, 16, 20, 26, 29, 35, 36, 38-40, 42, 44-49, 58, 59, 61, 65, 72, 76, 78, 79, 82, 85, 92, 93, 101, 110, 120, 124, 130-132, 138, 143, 148, 151, 154, 156, 159, 164, 172, 173, 178
	501-1000	6	6%	10, 53, 75, 103, 122, 153
	1001-10,000	11	10%	21, 27, 34, 51, 100, 104-107, 141, 179
	10,000+	7	7%	41, 53, 55, 87, 114, 150, 158
	Not reported/unclear	2	2%	13, 175, 182
Mode of Telehealth	Video	79	75%	11, 13, 16, 18, 20-23, 27, 31, 34-36, 38-42, 44, 47, 51, 52, 55, 58-61, 65, 66, 70, 72, 75, 76, 78, 79, 81, 84-87, 92, 93, 96-101, 103-107, 111, 114, 120, 122, 124, 128, 132, 139, 141-143, 148, 150, 153, 155, 156, 158, 159, 165, 167, 175, 176, 178, 182, 184
	Data store and forward	4	4%	37, 82, 172, 179
	Electronic chart/record review	3	3%	29, 45, 152
	Mixed modalities	10	12%	19, 46, 48, 49, 110, 112, 131, 138, 145, 154

Characteristic	Categories	Number of Articles - 106	Percentage of Articles	References
	Data streaming	1	1%	25
	Telephone	2	2%	10, 25
	Whats App	1	1%	12
	SMS Based	1	1%	53
	Online Module	3	3%	117, 130, 136
	NR/Unclear	2	2%	164, 173
Clinical category	Inpatient	18	17%	13, 21, 27, 41, 52, 55, 59, 60, 72, 75, 78, 92, 93, 111, 122, 128, 154, 178
	Outpatient	37	35%	10, 22, 29, 31, 34-37, 45-49, 70, 76, 79, 81, 82, 86, 87, 97, 98, 110, 120, 131, 132, 138, 139, 145, 151, 152, 155, 156, 165, 172, 176, 179
	EMS/ED	28	26%	12, 20, 25, 26, 38-40, 44, 58, 61, 65, 100, 101, 103-106, 111, 112, 114, 124, 141, 150, 153, 159, 164, 173, 182
	Education/mentoring	23	22%	11, 19, 23, 42, 51, 53, 66, 84, 85, 96, 99, 107, 117, 128, 130, 136, 142, 143, 148, 158, 167, 175, 184
Outcome categories	Patient	71	67%	10-13, 16, 20-22, 25-27, 29, 31, 34, 36-38, 40, 44, 46, 47, 49, 52, 55, 58-61, 65, 70, 72, 75, 76, 78, 79, 81, 82, 86, 87, 92, 93, 98, 100, 104-106, 110-112, 114, 122, 124, 128, 139, 141, 143, 145, 151-155, 159, 164, 167, 172, 175, 176, 178, 179, 182
	Provider	32	30%	18, 19, 23, 38-42, 51-53, 58, 66, 78, 84, 85, 96, 99, 101, 103, 107, 117, 130, 136, 142, 143, 148, 150, 158, 165, 175, 184
	Payer	13	12%	35, 37, 45, 70, 97, 120, 131, 132, 152, 156, 173, 182, 16

* China, Denmark, Scotland, Finland, New Zealand, Spain, Germany, Chile, Turkey, Japan, Sweden, Taiwan, Vietnam

Key Question 3

Additional study details can be found in **Appendix Table D-10**.

Table B-3. Characteristics of included studies for Key Question 3

Characteristic	Categories	Number of Articles (71 total)	Percentage of Articles	References
Geographic Location	United States	35	49%	6, 7, 9, 17, 24, 30, 33, 56, 57, 63, 64, 67, 73, 77, 80, 90, 91, 108, 115, 126, 127, 129, 134, 135, 144, 146, 147, 150, 157, 166, 169, 170, 174, 180, 183, 185
	Sweden	1	1%	71
	Norway	2	3%	160, 161
	Germany	2	3%	95, 119
	Australia	18	25%	14, 15, 28, 43, 69, 74, 83, 89, 102, 113, 116, 121, 123, 125, 140, 162, 163, 168
	Canada	5	7%	32, 54, 62, 68, 177
	New Zealand	1	1%	88
	Scotland	3	4%	16, 94, 171
	South Africa	1	1%	118
	Multiple	2	2%	133, 137
	Countries with a single study*	1	7%	109
Method	Program statistics	1	1%	91

Characteristic	Categories	Number of Articles (71 total)	Percentage of Articles	References
	Program records	6	8%	54, 63, 74, 126, 162, 163
	Program review	3	4%	113, 116, 123
	Program reporting	1	1%	125
	Program manager observations	1	1%	9
	Patient records	3	4%	54, 68, 140
	Registries	3	4%	6, 7, 146
	Hospital records	1	1%	150
	Administrative data	4	6%	6, 7, 146, 174
	Financial data	1	1%	90
	EHR data	1	1%	125
	EMR data	1	1%	166
	Pre-questionnaire	1	1%	9
	Survey	20	28%	17, 30, 32, 43, 56, 73, 80, 91, 108, 109, 115, 118, 125, 127, 134, 135, 137, 150, 174, 180
	Pilot tests	1	1%	64
	Comparison of two models	2	3%	169, 170
	Interview/Focus groups	28	39%	14-16, 24, 28, 67, 69, 71, 73, 80, 83, 88-90, 94, 95, 102, 119, 121, 129, 135, 144, 160-163, 166, 168, 171, 177, 183
	Exit interviews focused on case presentation	1	1%	9
	Chart review	5	7%	32, 62, 91, 162, 163
	Case study	4	6%	14, 28, 33, 147
	Case reports	2	3%	14, 28
	Case review	3	4%	24, 115, 157
	Site visits	4	6%	57, 90, 135, 166
	Patient and staff evaluations	1	1%	32
	Review of state statutes and regulations	1	1%	77
	Document analysis	2	3%	133, 135
Clinical category	Inpatient	8	11%	16, 118, 133
	Outpatient	30	42%	24, 32, 43, 54, 62-64, 69, 71, 74, 80, 91, 102, 113, 116, 119, 125, 126, 135, 140, 162, 163, 168, 171
	Telestroke and Emergency Care	20	28%	6, 7, 14, 28, 33, 77, 88, 90, 94, 134, 146, 147, 150, 160, 161, 166, 169, 170, 180
	Education/mentoring	13	18%	9, 56, 68, 73, 115, 121, 127, 144, 157, 174, 183
Outcome categories	Facilitators	55	77%	6, 7, 9, 16, 24, 32, 33, 43, 54, 56, 62-64, 68, 69, 71, 73, 74, 77, 80, 88, 90, 91, 94, 102, 113, 115, 116, 118, 119, 121, 125-127, 133-135, 140, 144, 146, 147, 150, 157, 160-163, 166, 168-171, 174, 180, 183
	Barriers	51	72%	6, 7, 9, 16, 24, 32, 33, 54, 56, 62-64, 68, 69, 71, 73, 74, 77, 80, 88, 90, 91, 94, 102, 113, 115, 116, 118, 119, 121, 125-127, 133-135, 140, 144, 146, 147, 150, 157, 162, 163, 166, 168, 170, 171, 174, 180, 183

Table B-4 repeats the number of times a construct was mentioned and adds the number of publications and the number of settings out of the four possible settings (inpatient, outpatient,

EMS/ED, or Education/Mentoring) in which these studies were conducted. This examination demonstrates that the constructs are relevant in all or most of the settings.

We also summarized facilitators and barriers by health care setting (inpatient, outpatient, emergency, and education/mentoring) in two ways. First, **Table B-5** reports the number of barriers and facilitators by setting. Included studies of provider-to-provider telehealth for EMS/ED and education/mentoring reported more facilitators the barriers. This was reversed infor inpatient studies and the number of reports were about equal for outpatient care.

Next, we created tables by setting and clinical indication similar to how the results are organized for Key Question 2. These tables provide the number of studies we identified for each clinical indication with a brief description of the telehealth interventions; basic information about the studies for each topic, including the method, size and location; implementation facilitators and barriers identified in the study as well as the impact cited as an indicator of successful implementation or motivation for sustainment. Not all studies sought to identify all three so the number of facilitators, barriers, and indicators of impact varies by topic. Finally, the studies we identified that compared strategies or interventions we described these in the narrative text for each setting.

Table B-4. Distribution of CFIR constructs

Barrier or Facilitator	CFIR Constructs	# Settings	# Mentions	# Publications
Facilitators	Leadership Engagement	4	13	10
	Implementation Climate	4	13	11
	Patient Needs & Resources	4	32	25
	Planning	4	11	9
	Compatibility	4	33	23
	External Policy & Incentives	4	18	12
	Adaptability	4	9	9
	Knowledge & Beliefs about the Intervention	4	16	11
	Available Resources	4	60	40
	Reflecting & Evaluating	4	12	11
	Access to Knowledge & Information	4	57	36
	Networks & Communications	4	37	30
	Engaging	4	23	18
Barriers	Cost	3	15	13
	Readiness for Implementation	3	17	12
	Formally Appointed Internal Implementation Leaders	3	7	7
	Executing	3	19	12
	Relative Priority	3	8	8
	Complexity	2	11	8

Table B-5. Facilitators and barriers by topic area

Topic	Facilitator or Barrier	# Mentions	# Publications
Inpatient	Barrier	45	9
	Facilitator	24	8
Outpatient	Barrier	95	26
	Facilitator	90	25
ED/EMS	Barrier	28	7
	Facilitator	65	19
Education/Mentoring	Barrier	24	7
	Facilitator	40	13

Inpatient

We identified eight assessments of implementation of provider-to-provider telehealth in rural areas that addressed inpatient care including intensive care, use of anesthesia, stroke rehabilitation, teletrauma, multidisciplinary specialty consultation, and telerobotics (**Table B-6, Appendix Table D-10**). One study compared facilitators and barriers in ICU programs using centralized monitoring (CM) versus virtual consult (VC) models,¹³³ one study used surveys to evaluate phone support consultation in South Africa,¹¹⁸ and one study used focus groups in Scotland to describe user experiences with video team consultations.¹⁶

One study of remote ICUs directly compared the facilitators and barriers for CM, which uses a hub with intensivists and hardwired data transfer and VC that uses portable equipment to connect local providers to relevant specialists. This study analyzed documents collected as part of a systematic review of effectiveness of remote ICU programs that use CM or VC.¹³³ The structural differences in the models drove the differences in barriers and facilitators such as lower cost and faster start-up for VC compared to the CM, but VC required more effort to integrate into workflows (**Table B-6**). Based on surveys of rural physicians who were trained in person and then offered telephone support when they needed to use anesthesia, Ngala et al. identified that an important barrier was remote consultants lacked understanding of the rural environment (**Table B-6**).¹¹⁸ The study of stroke rehabilitation video team consultations reported that lack of technological issues supported implementation and that the consultation increased the patient representative's confidence in the care provided locally (**Table B-6**).¹⁶

Table B-6. Findings of inpatient implementation studies

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Remote ICU 1 CM compared to VC	Document analysis N=91 documents Varied ¹³³	<ul style="list-style-type: none"> • Lower cost, faster start-up (VC) • Evidence supporting efficacy (CM) • No legal issues (both models) • Provided clinical information support (CM) 	<ul style="list-style-type: none"> • Higher fixed costs (CM) • Longer start up time (CM) • More reactive, requiring initiation or scheduling (VC) • Lack of evidence of clinical efficacy (VC) 	Not reported
Anesthesia 1 Phone support following in person training	Survey N=17 rural physicians South Africa ¹¹⁸	<ul style="list-style-type: none"> • In person training prior to implementation • Perception that good advice was available 	<ul style="list-style-type: none"> • Inadequate training • Lack of consultant understanding of environment • No occasion to provide service/no need 	Not reported
Stroke rehabilitation 1 Specialist participation via video in remote team meeting	Focus Group N=12 people; different roles in program Scotland ¹⁶	<ul style="list-style-type: none"> • No staff resistance; clear rationale for use • Technology training and IT involvement • Reliable equipment 	<ul style="list-style-type: none"> • Delay of not having specialist immediately accessible 	<ul style="list-style-type: none"> • Improved decision making • Increased confidence in care (patient representative)
Teletrauma 1	Interview N=14 stakeholders Canada ¹⁷⁷	<ul style="list-style-type: none"> • Flexibility of the technology to receive clinical input • Interprofessional relationships • Seamless integration of technology 	<ul style="list-style-type: none"> • Complicated nature of use • Familiarity with the technology • Workflow changes 	<ul style="list-style-type: none"> • Increased personal and professional support for rural clinicians

Topic Number of Studies	Method N*			
Intervention	Location	Facilitators	Barriers	Impact
Robotic Telemedicine 1	Survey N=38 health care institutions United States, Canada, Ireland ¹³⁷	Not reported	<ul style="list-style-type: none"> • Equipment Cost • Executive administration and leadership hesitancy regarding adoption of robotic telemedicine • Lack of effective leadership • Lack of exposure to robotic telemedicine • Lack of understanding of robotic telemedicine • Patients, Physicians and Nurse hesitancy regarding adoption of robotic telemedicine • Physician lack of incentives to use robotic telemedicine • Potential impact on quality of care • Regulatory barriers (out-of-state licensing, malpractice liability, credentialing, government and nongovernment reimbursement, DEA licensing) • Robotic telemedicine seen as a local threat • Technology issues (usability, reliability, internet connectivity, remote data access, technical support, documentation and billing) 	Not reported
Multidisciplinary Specialty Consultation 3	Interview N=63 hospitals Australia ¹⁵ Secondary analysis national survey data N= 4,608 hospitals U.S. 50 States ³⁰ Method= N=8 hospitals U.S. Montana, Nevada, North Dakota ⁵⁷	<ul style="list-style-type: none"> • Benefits of telehealth well communicated • Investment in services to support telehealth delivery • Specific service agreements • Practitioner Champion • Technical assistance • Training on work flow and infrastructure 	<ul style="list-style-type: none"> • Billing system • Clinician resistance • Confusion regarding policy for out-of-catchment • Cost of using existing computers to provide telehealth • Credentialing process and agreeing to be credentialed • Fragmentation of information technology between tertiary and primary care • Software difficulties • Health Information Exchange capabilities • Cost • Practitioner reluctance • Securing properly credentialed practitioners 	Not reported

*N is used here to represent the unit of analysis, which may be number of individual participants or may be number of health care sites or systems.

Abbreviations: CM = centralized monitoring; ICU = intensive care unit; VC = virtual consult.

Table B-7 provides the barriers and facilitators from studies of provider-to-provider telehealth for inpatient studies standardized by CFIR constructs. While there are fewer studies of inpatient care, the barriers and facilitators are not repeated in multiple studies. The most frequently repeated are *complexity* cited seven times as a barrier and *available resources* cited as a facilitator six times.

Table B-7. Inpatient: barriers and facilitators by CFIR constructs

Type	Facilitator or Barrier Name	Facilitator or Barrier Number of Mentions	Reference Number(s)
Barrier	Access to Knowledge & Information [*]	2	15, 118
	Adaptability [†]	1	118
	Available Resources [‡]	1	118
	Compatibility	3	15, 57, 177
	Complexity [§]	7	15, 30, 133, 177
	Cost	4	15, 57, 133, 137
	Engaging	2	57, 137
	Executing	2	137, 177
	External Policy & Incentives	1	137
	Knowledge & Beliefs about the Innovation [¶]	3	118, 133, 137
	Implementation Climate	2	137
	Leadership Engagement	2	137
	Patient Needs & Resources [#]	3	16, 129, 137
	Networks & Communications ^{**}	1	118
	Planning	2	15
	Reflecting & Evaluating ^{††}	1	118
Facilitator	Access to Knowledge & Information [*]	2	16, 118
	Adaptability	1	177
	Available Resources [‡]	6	15, 16, 57, 133
	Cost	1	133
	Executing	1	177
	External Policy & Incentives	1	15
	Formally Appointed Internal Implementation Leaders	1	57
	Knowledge & Beliefs about the Innovation [¶]	1	16
	Patient Needs & Resources [#]	1	118
	Networks & Communications ^{**}	3	15, 118, 177
	Planning	1	57
	Readiness for Implementation ^{††}	1	133

^{*} Access to digestible information and knowledge about the innovation and how to incorporate it into work tasks.

[†] Degree to which an innovation can be adapted, tailored, refined, or reinvented to meet local needs.

[‡] Level of resources organizational dedicated for implementation and on-going operations including physical space and time.

[§] Perceived difficulty of the innovation, reflected by duration, scope, radicalness, disruptiveness, centrality, and intricacy and number of steps required to implement.

^{||} Costs of the innovation and costs associated with implementing the innovation including investment, supply, and opportunity costs.

[¶] Individuals' attitudes toward and value placed on the innovation, as well as familiarity with facts, truths, and principles related to the innovation.

[#] Extent to which patient needs, as well as barriers and facilitators to meet those needs, are accurately known and prioritized by the organization.

^{**} Nature and quality of webs of social networks, and the nature and quality of formal and informal communications within an organization.

^{††} Quantitative and qualitative feedback about the progress and quality of implementation accompanied with regular personal and team debriefing about progress and experience.

^{‡‡} Tangible and immediate indicators of organizational commitment to its decision to implement an innovation.

Outpatient

We identified 30 studies of implementation of provider-to-provider telehealth for rural outpatient care. These studies all assessed consultations in which one provider, often a specialist, contributed to the diagnosis or management of a patient by another provider, often a primary care physician, nurse or someone lacking specialist certification or extensive experience with the condition or treatment. The barriers and facilitators are grouped and organized by clinical indication in **Table B-8**, and additional details can be found in **Appendix Table D-10**.

Five studies of multi-specialty programs included two statewide programs^{80, 135} and three programs serving a small group of clinics or a single health system.^{71, 102, 119} Psychiatric consultations were the subject of five studies of services that provided expert advice on a range of mental health issues including: medication therapy for opioid use disorder in a group of community clinics that are part of the Veteran Health Administration;²⁴ advice on medications and treatment for children in a state Medicaid program,⁶³ and programs to help diagnose adults and identify and arrange appropriate services.^{32, 62, 91} Five studies were of programs that provided consultations for different aspects of care related to long term services and supports including assessment of whether nursing home residents should be transferred to hospitals,⁶⁴ oral health screening and teledentistry,^{162, 163} wound care,⁷⁴ outpatient geriatric assessment and management,¹²⁶ and pediatric hospice care.¹⁶⁸ The remaining studies each evaluated consultations related to evaluating or managing patients with chronic conditions including cancer,^{69, 140} gastroenterology,⁵⁴ dermatology,¹²⁵ cardiology,¹¹⁶ nephrology,¹¹³ occupational screening of miners,⁴³ and support for midwives managing pre-eclampsia.¹⁷¹

Studies of multispecialty programs included assessments of how one program evolved from a pilot test to a statewide program over years with a mixture of sources of funding.⁸⁰ An evaluation of a 10-year, multi-site initiative to increase access to care in rural areas in California through telehealth reported that organizational barriers contributed to a lack of networking across programs and lower uptake than expected of telehealth services.¹³⁵ Both of these studies demonstrate how implementation and spread requires sustained efforts and commitments from multiple stakeholders and suggests that statewide or regional efforts can be effective. Questions among rural clinicians about whether telehealth was truly patient-centered were a barrier for teleconsultations in rural Sweden as some providers felt it may be easier to send patients to the hospital directly rather than delay hospitalization for a consult.⁷¹ Concerns cited in United States studies were echoed in studies in other countries. A program in Germany cited concerns about time, financing and changes to established workflows as barriers that could be addressed if systems were more usable and training provided.¹¹⁹ A program in Australia illustrated time concerns by documenting that teledermatology consultations take twice as long as in-person assessments and payment does not include this extra time. This program addressed this and other barriers by adding a telehealth coordinator who reduced the need for clinician time and by assuring technical support was available.¹⁰²

Telehealth is often proposed as one solution to the shortage of mental and behavioral health providers and programs in rural areas. Some of the telehealth programs address specific treatments, such as the use of Buprenorphine for opioid use disorder in VA clinics in one state,²⁴ while others are more general. The evaluation telehealth supported Buprenorphine was one of the few that used an implementation science framework to assess their experience and then translate this experience into an implementation tool kit that could be used by others to replicate the program. Another telehealth consult program provided medication review and treatment recommendations for children in a state Medicaid program.⁶³ These programs had to overcome specific barriers including legal concerns related to prescribing and the need for consultants to understand resource availability in other locations. Another program used a continuous quality improvement approach to identify and make workflow adjustments to assure success.⁹¹ Psychiatric teleconsultation services in Ontario, Canada, one for adults⁶² and one focused on geriatric psychiatry³² identified fundamental gaps in organization and culture as barriers, such a lack of integration of the telehealth consultation with telephone and in person visits with the patient⁶² and a concern among providers that telehealth would allow the government to justify the lack of support for increasing local, in-person services for patients.³²

The two articles on provider-to-provider telehealth for cancer care were both reports about the same program in Queensland, Australia. This program allowed chemotherapy to be administered in rural hospitals by local physicians and nurses supported by remote oncologists and chemotherapy nurses.^{69, 140} Starting with a pilot to demonstrate safety, the program expanded to six sites after addressing barriers including lack of role clarity and technology restrictions. Changes included assuring the iCamera could zoom sufficiently to allow checks on chemotherapy bags and provide good visuals during physical exams; structuring the program to provide professional development opportunities for rural nurses; and financial incentives for physicians to participate.

Long term care residents often have limited access to health care services for many reasons including resident's/patient's difficulties traveling and the fact that specialty services are rarely available onsite in nursing homes and other residential care and home-based long-term care. In this context, telehealth consultations and programs may offer services that would not otherwise be available. For example, a multisite program was established by a health system to provide acute assessment and care planning support in order to reduce patient transfers to hospitals.⁶⁴ The program grew from 5 to 34 sites in 4 years by building on the health system's experience with telehealth for other uses and working to change the culture from one that had defaulted to hospitalization to one that accepted treating residents in place. An oral health program provides another example in which a new service was made available. Residents who were not receiving dentistry services were screened by a technician who used a live intra oral camera to transmit images to a remote dentist who could assess what could be done on site and what required travel to a dentist.^{162, 163} This program was able to increase compliance with guidelines and regulations while increasing staff confidence in their ability to manage oral health. Other applications included a geriatric consult service in the Veterans Health Administration that was able to increase assessments by setting up both synchronous and asynchronous consultations.¹²⁶ Implementation of a teleconsult program to support wound care by home and community providers revealed structural barriers to implementation including the need for staff computer literacy and the lack of use of standardized terminology by the home care nurses and consultants.⁷⁴ Adding telehealth consultations to a pediatric hospice program underscored trade-offs and challenges. The program demonstrated the ability to provide multidisciplinary, timely

help to supplement in person care, but found that video consults were limited in their ability to assess family distress and that the consults risk prioritizing expert views over family needs.¹⁶⁸

The remaining outpatient studies included one report each about telehealth consultations for different chronic, or not immediately acute conditions, including a regional cardiology program,¹¹⁶ chronic kidney disease consultations for an Indian Health Service clinic,¹¹³ a local gastroenterology program focused on a single condition,⁵⁴ and a large dermatology program with 15 hubs in the VA.¹²⁵ All these programs were designed to increase access and timeliness of care and all faced hurdles related to lack of staff support, space, and connectivity/bandwidth. Two less common approaches included adding telehealth to a mobile clinic that provides screening for coal miners⁴³ and creating a phone app to supplement support to midwives managing pre-eclampsia in a rural area.¹⁷¹ Both of these programs had to overcome unique technical challenges, but faced common barriers related to limited connectivity.

Table B-8. Findings of outpatient implementation studies

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Multi-specialty 5 Programs that make consultations from a range of specialists available over video, phone or electronic records	Stakeholder (patients and provider) surveys and interviews N=Not reported Statewide hub/spoke program in South Carolina ⁸⁰ Evaluation with surveys, site visits, documentation review, interviews N=10 organizations in 22 counties California ¹³⁵ Focus groups N=5 primary health-care centers; 19 health care personnel Sweden ⁷¹ Interviews N=18; Physicians, administrators, medical students Germany ¹¹⁹ Interviews N=10 expert providers of telehealth Australia ¹⁰²	<ul style="list-style-type: none"> • Private foundation support for start up • Ongoing support from state funds and billing • Purposeful stages of implementation including training and evaluation • Rural providers perceived benefits were worth their investment • Usability of system/ training • Financing plan Coordinator and tech support 	<ul style="list-style-type: none"> • Limits to pro bono consults specialists can provide, need for financial support for specialist time or other concerns about financing • Low networking across programs • Organization barriers and lower utilization than expected • Some specialties and exams not a good fit • Aversion among staff to new ways of working including technology • Time demands Perception it could be easier to transport patient 	<ul style="list-style-type: none"> • Grew from 11 in 2012 to over 1300 consults in 2017 • Travel/cost savings for patients

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Psych/Mental Health 7 Managing opioid use disorder, mental health care planning, for adults and medications and treatment for recommendatio ns for children	Interviews and case review N=3 Clinics; 19 interviews VA in Maine ²⁴ Continuous Quality Improvement Surveys, Chart reviews and program statistics N=1 health system Illinois ⁹¹ Program records N=1 state Medicaid program Washington provision to Wyoming state Medicaid ⁶³ Chart reviews Interviews N=10 Ontario, Canada ⁶² Chart review, patient and staff evaluations, survey to referring MDs, focus groups with community agencies GeroPsych service N=6 communities Ontario, Canada ³² Focus group N=10 Psychiatrists and 4 psychologists across 3 states U.S. Washington, Michigan, Arkansas ⁶⁷ Questionnaire N=8 primary care providers, 4 psychologists Chile ¹⁰⁹	<ul style="list-style-type: none"> • Mission to address issue • Prior teleprescribing experience • Maintenance of local control over patient • Development of tool kit • Ongoing evaluation and adjustment • Peer relationships among clinicians • Detailed recommendations and education • Teamwork • Communication • Provider willingness to collaborate • Punctuality • Staff openness 	<ul style="list-style-type: none"> • Legal concerns • Conflicting interests • Coordination with other programs • Need for team integration • E consults limited to quick or simple inquiries • Need for understanding of local resources • Limited awareness of consults • Discomfort managing mental health in primary care • Perception that telehealth provides justification for not increasing in person access • Ability to track clinical outcomes and cost (2234) • Clinic investment • Communication • Credentialing • Electronic record process • Mobilizing resources • Scheduling support • Staff Turnover • Training • Access to computer and suitable room • Incomplete information on patient • Technical difficulties 	<ul style="list-style-type: none"> • Positive evaluations of ability to see and hear and usefulness by patients • Physician intension to continue to use • Increased ability to include family or an interpreter • Reduction in outlier pediatric psychiatric medications
Cancer 2 Chemotherapy administration at remote sites	Interviews N=19 ⁶⁹ Patient records N=62 ¹⁴⁰ Australia	<ul style="list-style-type: none"> • Opportunity for professional development for rural nurses • Good communication • Implementation management team that developed plan and documentation • Funding support and financial incentives for physicians to participate 	<ul style="list-style-type: none"> • IT did not have capacity to zoom in to check chemotherapy bags and physical exam • Lack of good electronic documentation (fixed after pilot) • Lack of role clarity • Turnover in management and nurses 	<ul style="list-style-type: none"> • Better continuity of care • Spread from pilot to 6 sites • Similar safety to in person care • Project transitioned from special funding to normal financing

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Long-Term Care 3 Acute illness; hospital transfer decisions	Pilot Tests N=1 health system, up to 14 sites Avera Health, Several States ⁶⁴ Interview N=21 administrators and clinicians across 16 facilities U.S. Nationwide ¹²⁹ Interview N=8 Clinicians Germany ⁹⁵	<ul style="list-style-type: none"> • Meetings with leadership • Creation of an implementation plan with stakeholder review • Establishing billing procedures • Leveraging decision support tools • Health system experience with telehealth • Communication • Creating one time slot for visits creates focus, information getting lost • Transmitting patient information 1 day ahead 	<ul style="list-style-type: none"> • Need for bandwidth • Space constraints requiring mobile equipment • Culture change to accept treating in place • Building relationships • Cannot see or hear as well • Clinician reluctance • Difficulty working with cognitively impaired patients • Increased burden on staff • Lack of training • Technology challenges, including internet and connectivity • Workflow changes 	<ul style="list-style-type: none"> • Increase from 5 to 34 sites in 4 years
Oral Health in Long-Term Care residencies 2	Chart review and program records, interviews and focus group N=250 charts, 9 facilities Australian ^{162, 163}	<ul style="list-style-type: none"> • Feedback on compliance • Documentation of cost savings 	<ul style="list-style-type: none"> • Not enough staff time for program management • Oral health staff lack of experience with dementia 	<ul style="list-style-type: none"> • Minimized disruption to residents • Increase compliance • Increased staff confidence in managing oral health
Geriatrics 1	Program records N=12 hubs U.S. Veterans Health Administration ¹²⁶	<ul style="list-style-type: none"> • Prior relationships between hubs and rural clinics 	<ul style="list-style-type: none"> • Multiple contacts needed to establish new relationships • Need to develop case finding approaches to identify patients who could benefit from consultation. 	<ul style="list-style-type: none"> • Increase from 4 to 12 hubs in 4 years
Wound Care 1	Program records N=4 home and community health providers Australia ⁷⁴	<ul style="list-style-type: none"> • Train the trainer model • Staff commitment 	<ul style="list-style-type: none"> • Need for computers and staff computer training/literacy • Need for web access • Need for standardized terminology 	Not reported

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Pediatrics 2	Interview N=15 hospice nurses 1 Midwestern U.S. state ¹⁶⁸ Interview N=1 hub, 7 community health centers Australia ⁸⁹	<ul style="list-style-type: none"> • Ability to involve multiple disciplines and family members • Timely goals of care discussions • Networking and collaboration 	<ul style="list-style-type: none"> • Difficult to assess and address distress • Family perceptions could be usurped by experts • Enough rooms • Funding • Technical issues, being able to tell who was speaking and being able to hear 	Not reported
Chronic Conditions 1 Gastroenterology, care for inflammatory bowel disease	Patient and program records N=99 patients Ontario, Canada ⁵⁴	<ul style="list-style-type: none"> • Multidisciplinary team 	<ul style="list-style-type: none"> • Limited availability of remote telehealth sites • Lack of awareness of program 	<ul style="list-style-type: none"> • Travel cost savings • Reduction in wait times
Dermatology 2 Remote assessment and diagnosis	EHR data, program reporting, online survey N=15 hubs U.S. Veterans Administration ¹²⁵ Survey N=34 Primary care providers U.S. Mississippi ¹⁰⁸	<ul style="list-style-type: none"> • Strong stakeholder support • Use of residents to reduce burden on dermatologists • Communications with primary care 	<ul style="list-style-type: none"> • Understaffing at rural spokes • Lack of space and equipment • Restrictions on funding • Bureaucracy • Concern regarding possibility of misdiagnosis due to poor image quality • Concerns regarding possible loss of patient confidentiality • Insurance coverage • Investment in time needed to master technology • Misconceptions about tele dermatology • Time required to submit consult and response time 	Not reported
Cardiology 1 Case review and remote exam	Program review N=5 sites Minnesota, Wisconsin ¹¹⁶	<ul style="list-style-type: none"> • Using local nurses to prepare patient and chart 	<ul style="list-style-type: none"> • Difference in reimbursement rules for locations • Need for patient and scheduler education 	<ul style="list-style-type: none"> • Able to expand from 1 to 5 sites, one site dropped out

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Nephrology 1 Care review and remote patient/provider appointment	Program review N=1 site Zuni Pueblo, Indian Health Services ¹¹³	<ul style="list-style-type: none"> • Nurse care manager and ancillary staff are key • EHR access • Access to specialist between scheduled clinics • Periodic in-person visits to build rapport and trust 	<ul style="list-style-type: none"> • Does not address need patients with acute needs • Technical issues • Communication components (eye contact, emotional support) 	Not reported
Screening 1 Mobile clinic: mining related exposure and general health	Surveys N=278 (62%) of 4511 mobile clinic with telehealth for miners New Mexico ⁴³	<ul style="list-style-type: none"> • Understanding of occupation risks • Specialist mentor non-specialists • Financially sustainable; most patients have insurance 	<ul style="list-style-type: none"> • Not reported 	<ul style="list-style-type: none"> • Expansion from New Mexico to Wyoming • High percentage return to clinic
Midwifery 1 Using phone app to managing pre-eclampsia	Focus groups, N=18 midwives Scotland ¹⁷¹	<ul style="list-style-type: none"> • Access to up to data information on a relative rare event 	<ul style="list-style-type: none"> • Concerns about using technology • Need for system that works without an internet connection 	Not reported

*N is used here to represent the unit of analysis, which may be number of individual participants or may be number of health care sites or systems.

Abbreviations: EHR = electronic health record; IT = information technology; MD = medical doctor; U.S. = United States; VA = U.S. Department of Veteran's Affairs.

Table B-9 provides a summary of the barriers and facilitators identified in studies of outpatient provider-to-provider telehealth for rural populations by the standardized constructs. As almost half of the studies included for Key Question 3 involved outpatient care, the counts of facilitators and barrier are higher. However, unlike inpatient care some constructs were identified much more frequently than others. *Available resources* was the most frequent barrier, cited 23 times. But others also mapped to higher numbers of cited barriers including *compatibility* (14) and *access to knowledge and information* (9). *Available resources* (15) and *access to knowledge and information* (15) were also cited as common facilitators (11 times), but other important facilitators included *networks & communications*(16) and *patient needs & resource*(9)s. In the case of this last category, a facilitator for use of telehealth was often patients' needs for expertise and services that were not otherwise available without telehealth.

Table B-9. Outpatient: barriers and facilitators by CFIR constructs

Type	Facilitator or Barrier Name	Facilitator or Barrier Number of Mentions	Reference Number(s)
Barrier	Access to Knowledge & Information*	9	62, 67, 69, 74, 102, 129, 162
	Adaptability†	2	62, 67
	Available Resources‡	23	24, 32, 49, 64, 67, 69, 71, 74, 89, 91, 102, 109, 113, 125, 140, 162, 171

Type	Facilitator or Barrier Name	Facilitator or Barrier Number of Mentions	Reference Number(s)
	Compatibility [§]	14	64, 67, 69, 74, 80, 102, 108, 113, 119, 129, 135
	Complexity	3	67, 74, 171
	Cost [¶]	3	80, 116, 119
	Engaging [#]	5	24, 54, 71, 116, 135
	Executing ^{**}	10	32, 89, 108, 109, 119, 129, 140
	External Policy & Incentives ^{††}	5	89, 102, 108, 125, 135
	Implementation Climate ^{‡‡}	1	62
	Knowledge & Beliefs about the Innovation ^{§§}	4	32, 119, 135
	Leadership Engagement	1	108
	Networks & Communications ^{¶¶}	6	24, 67, 113, 125, 126, 129
	Patient Needs & Resources	8	54, 62, 71, 129, 168
	Planning ^{##}	1	135
	Readiness for Implementation ^{***}	3	24, 67, 119
	Reflecting & Evaluating ^{†††}	2	63, 135
	Relative Priority ^{‡‡‡}	3	24, 32, 102
Facilitator	Access to Knowledge & Information [*]	15	24, 32, 43, 62, 64, 69, 71, 74, 91, 125, 126, 135
	Adaptability [†]	1	125
	Available Resources [‡]	11	32, 62, 64, 113, 116, 119, 125, 140, 162
	Compatibility	1	109
	Complexity	1	119
	Cost [¶]	4	43, 91, 119, 163
	Engaging [#]	7	54, 62, 80, 119, 140, 168
	External Policy & Incentives ^{††}	3	80, 140
	Executing	2	95, 109
	Formally Appointed Internal Implementation Leaders ^{§§§}	3	54, 113, 135
	Knowledge & Beliefs about the Innovation ^{§§}	1	32
	Leadership Engagement	4	24, 125, 140
	Patient Needs & Resources	9	32, 43, 108, 113, 119, 126, 168, 171
	Networks & Communications ^{¶¶}	16	32, 62, 67, 69, 80, 89, 95, 109, 113, 125, 126, 129, 135, 140, 162
	Planning ^{##}	4	64, 119, 135, 140
	Readiness for Implementation ^{***}	5	24, 64, 80
	Reflecting & Evaluating ^{†††}	4	64, 91, 140, 163
	Relative Priority ^{‡‡‡}	1	74

- * Access to digestible information and knowledge about the innovation and how to incorporate it into work tasks.
- † Degree to which an innovation can be adapted, tailored, refined, or reinvented to meet local needs.
- ‡ Level of resources organizational dedicated for implementation and on-going operations including physical space and time.
- § Degree of tangible fit between meaning and values attached to the innovation by involved individuals, how those align with individuals' own norms, values, and perceived risks and needs, and how the innovation fits with existing workflows and systems.
- ‡ Perceived difficulty of the innovation, reflected by duration, scope, radicalness, disruptiveness, centrality, and intricacy and number of steps required to implement.
- ¶ Costs of the innovation and costs associated with implementing the innovation including investment, supply, and opportunity costs.
- # Attracting and involving appropriate individuals in the implementation and use of the innovation through a combined strategy of social marketing, education, role modeling, training, and other similar activities.
- ** Carrying out or accomplishing the implementation according to plan.
- †† External strategies to spread innovations including policy and regulations (governmental or other central entity), external mandates, recommendations and guidelines, pay-for-performance, collaboratives, and public or benchmark reporting.
- ‡‡ Absorptive capacity for change, shared receptivity of involved individuals to an innovation, and the extent to which use of that innovation will be rewarded, supported, and expected within their organization.
- §§ Individuals' attitudes toward and value placed on the innovation, as well as familiarity with facts, truths, and principles related to the innovation.
- ‖ Extent to which patient needs, as well as barriers and facilitators to meet those needs, are accurately known and prioritized by the organization.
- ¶¶ Nature and quality of webs of social networks, and the nature and quality of formal and informal communications within an organization.
- ### Degree to which a scheme or method of behavior and tasks for implementing an innovation are developed in advance, and the quality of those schemes or methods.
- *** Tangible and immediate indicators of organizational commitment to its decision to implement an innovation.
- ††† Quantitative and qualitative feedback about the progress and quality of implementation accompanied with regular personal and team debriefing about progress and experience.
- ‡‡‡ Individuals' shared perception of the importance of the implementation within the organization.
- §§§ Individuals from within the organization who have been formally appointed with responsibility for implementing an innovation as coordinator, project manager, team leader, or other similar role.
- ||| Commitment, involvement, and accountability of leaders and managers with the implementation of the innovation.

Telestroke and Emergency Care

One of the most commonly studied applications of provider-to-provider telehealth in rural areas is the diagnosis and management of stroke due to the higher prevalence of stroke and stroke risk factors in rural areas and treatments that require accurate diagnosis and timely administration.^{188, 189} The telestroke programs described in this section bridge ED and inpatient care as they include consultations as part of initial assessment and triage as well treatment decisions and care delivery. We did not identify studies of the implementation of EMS telestroke programs in which consultations focus on prehospital triage and decisions made in the field about where the patient should be transported.

Table B-10 provides an overview of seven studies (reported in eight articles)^{6, 7, 14, 28, 33, 77, 146} of facilitators, barriers and impact related to the implementation of telestroke programs (Additional details in **Appendix Table D-10**). All of the projects studied were hub-spoke models in which one or more hubs where specialists were located were connected with rural hospitals. Evaluations of these programs included studies of statewide programs in West Virginia⁶ and South Carolina,^{7, 146} case studies of networks around a single hub,³³ a comparison of the early implementation of a network in South Carolina to one in Georgia¹⁴⁷ and an assessment of a regional program in Australia.^{14, 28} One evaluation reviewed state laws and regulations in the United States.⁷⁷

The telestroke implementation studies were of successful programs and the evaluations focused on the factors that supported this success. A case study comparing the early (1991) implementation in two networks reported that the networks had not integrated the technology into their care delivery processes and identified enablers which continue to be called out in other,

more recent studies including: resource needs, the key role of performance monitoring and continuous improvement; the importance of a champion and dedicated coordinator at spokes, stakeholder involvement, and tangible goals such as stroke center certification¹⁴⁷ A frequently cited approach included stepped or phased implementation that started with preliminary needs and workflow assessments to inform pilot tests; diversity of engagement and funding, including private and government support; the need for staff to support the program, and training; and the importance of ongoing evaluation and program improvement. Barriers were less frequently cited but included lack of sufficient IT support, lack of integration of records and patient data, and the prohibition on fees or additional reimbursement for telehealth infrastructure in some states. Reports on telestroke implementation also focused on the impact on care and organizational outcomes, citing fewer transfers and certification as a stroke center as motivation to continue to sustain and improve the programs.

We identified studies of implementation of provider-to-provider consultations for general emergency care, pediatric emergencies and psychiatric emergencies in addition to telestroke. Five studies were of models in which a remote specialist or emergency physician advises a generalist physician, nurse practitioner or nurse who staff a rural emergency room.^{88, 90, 150, 166, 180} Follow-up with rural EDs that did and did not use telehealth based on a United States national survey, found that 67% of nonusers had considered implementing telehealth but reported that the single most important reason the ED is not using telehealth was cost (37% of respondents), followed by technologic concerns and the assessment that telehealth is not needed to meet patient's needs (11% each). Despite these barriers, six percent reported they had started to use telehealth since the original survey.¹⁸⁰ Costs cited in this and other studies included the cost of technology but also the cost of the subscription services that provide access to the consultants. Additional barriers included rural providers lack of understanding of telehealth, lack of perception that telehealth will address a clear problem and perceptions that the motivation is to save money, particularly on personnel. For this particular use the identified facilitators were general satisfaction and having a telehealth coordinator who could handle scheduling and technology.

Two studies assessed implementation of telehealth specifically for pediatric emergencies.^{134, 170} One is an example of the few studies that compared different models; one model that provided only pediatric specialty consultations and one in which pediatrics was one of several specialties provided as part of a system wide consult service.¹⁷⁰ This study found that both models were considered successful, but produced very different results as they served different populations. The specialist only model was used for more critical cases while in the other, pediatric consults were used less, but used for both high and low risk cases. As a result, perceptions and measures of the systems differed. These studies identified specific factors that were not emphasized in other studies, such as the need to test technology that is not in frequent use and the importance of building rapport and assuring telehealth fits in the culture of the practice.

Two studies in three articles assessed telehealth psychiatric consultations for patients presenting in EDs. One study evaluated two different network approaches to emergency psychiatric telehealth: a regional network, with assessments and consults available as part of a system that provides multispecialty consults for MI, stroke and other acute illnesses as well as behavior health, by pressing a button compared to a smaller, local system in which a behavioral health specialist was paged when needed.¹⁶⁹ The assessment focused on patient characteristics and confirmed that both models increase access to inpatient care, but the evaluation did not

explore detailed implementation differences in the systems. A study in northern Norway^{160, 161} evaluated a system that made consultation available 24/7 by telephone and video; the study reported that using well established technology and having a safety net system supported the implementation and use of telehealth consults.

We identified one study that specifically addressed implementation of telehealth in prehospital care by EMS. A small study in Scotland⁹⁴ reported several major barriers to the use of remotely guided ultrasound in prehospital care. These included a lack of evidence and lack of documented need for remote guided ultrasound as well as different perceptions of EMS personnel and consulting physicians about skills and priorities.

Table B-10. Findings of telestroke and emergency care implementation studies

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Telestroke 7 Hub:Spoke	<p>Registries and administrative Data N=2 statewide studies West Virginia⁶ South Carolina^{7, 146}</p> <p>Case study N=1 program South Carolina³³</p> <p>Case studies N=2 networks in South Carolina/ Georgia¹⁴⁷</p> <p>Case study including reports and interviews N=16 stakeholder reports, 13 funder reports, 10 protocols, 3 collaborative agreements, 93 meeting minutes Australia^{14, 28}</p> <p>Review of state statutes and regulations N=50 states United States⁷⁷</p>	<ul style="list-style-type: none"> • Stepped implementation: pre, pilot, full, sustainability • Engagement • Diverse support and funding (governmental health and non-health, philanthropic) • Site champions • Pre-implementation clinician surveys • State legislation • Shadowing at hubs • Training for all roles with technology and guidelines • Sustainable technology • Local site coordinator to support start-up at spokes • Expansion locations based on data • Feedback on performance; ongoing evaluation and promotion of success • Focus on application of telehealth that is sustainable and improved care • Licensing and allowing practice across state lines • Same reimbursement for in-person or additional technology fee 	<ul style="list-style-type: none"> • Video conferencing, imaging and clinical records not integrated • IT support not available 24/7 • Prohibition of fees or reimbursement in some states related to telehealth infrastructure 	<ul style="list-style-type: none"> • Increase in consult use • Appropriate triage and treatment (tPA) • Fewer transfers • Certification of additional stroke centers • Diffusion of knowledge • Expansion from pilots to regional or multisite programs

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Emergency Care: Non specific 6 Remote consultation Rural: Primary care provider, Nurse practitioner, nurse or generalist Consultant: Specialists or ED physician or nurse.	Survey of rural EDs N=153/177 telehealth users; 375/453 non users U.S. ¹⁸⁰ Surveys and hospital records Pre/Post implementation N=9 hospitals Mississippi ¹⁵⁰ EMR data, interviews, and site visits N=85 administrator at 26 rural hospitals South Dakota Avera Health ¹⁶⁶ Financial data, interviews, site visits N=1 emergency system; 49 rural hospitals, same interviews as Ward above South Dakota Avera Health ⁹⁰ Interviews N=12 New Zealand ⁸⁸ Program evaluation N = 206 patient records Australia ¹²³	<ul style="list-style-type: none"> • Rural provider satisfaction • Having a telehealth coordinator • Communication • Executive Sponsorship 	<ul style="list-style-type: none"> • Cost of technology or subscription • Lack of understanding of telehealth • No relationship or experience with distance provider • Perception it is about saving money, not improving care • Lack of a clear problem telehealth will solve • Technology issues include lack of bandwidth and power cuts 	<ul style="list-style-type: none"> • Increase in rural hospital admissions (sterling) • -Increase in volume • -Decrease in inappropriate admissions • Low rate of utilization, but used when expert needed • Facilitates transfers, documentation, urgent critical care when physicians not in ED • Maybe profitable if hospital is able to increase revenue admissions and save on local physician back up for NP/PA staffing ED • High level of use (67% of NP used system every shift)
Emergency Care: Pediatrics 2 Pediatrics as part of multispecialty consults service; Dedicated pediatric service	Two models (University of California Davis-specialty/hub vs. Advera-general ED including pediatrics) N=30 hospitals; 15 each ¹⁷⁰ Survey based on themes from interviews N=7 hospitals, 48 interviews, surveys 5 hospitals 104 (34%) of 306 clinicians invited University of Pittsburgh ¹³⁴	<ul style="list-style-type: none"> • Weekly test calls to preemptively solve connection issues • Hub providers build rapport by attending remote ED staff meetings and collaborating on quality initiatives • Perception that telehealth is useful 	<ul style="list-style-type: none"> • Initially use not part of culture or care processes • Lack of provider comfort with telehealth • Technology issues • Lack of time; negative impact on workflows 	<ul style="list-style-type: none"> • Unclear when use is appropriate and whether goal is to reduce transfers and/or discharges

Topic Number of Studies Intervention	Method N* Location	Facilitators	Barriers	Impact
Emergency Care: Psychiatrics 2 studies (3 articles)	Compare 2 ED Behavioral Health Models N=19 spoke hospitals; 2 networks) U.S. Midwest ¹⁶⁹ Interviews N=29 Norway ^{160, 161}	<ul style="list-style-type: none"> Well established technology Ability to confirm initial assessment and collaborative solve problems System provides safety net 	<ul style="list-style-type: none"> Not reported 	<ul style="list-style-type: none"> Increase in admissions to inpatient facilities (interpreted as an increase in access) Immediacy of assessment Engagement with patient Access to specialist/MD that is not available locally Reduced uncertainty
EMS: Ultrasound 1	Interviews N=12 Scotland ⁹⁴	<ul style="list-style-type: none"> Willingness to collaborate in training and care 	<ul style="list-style-type: none"> Difference in EMS and MD perceptions of utility and skills needed Unclear evidence of need and benefits Perception it could result in delay in transport 	Not reported

*N is used here to represent the unit of analysis, which may be number of individual participants or may be number of health care sites or systems.

Abbreviations: ED = emergency department; EMS = emergency medical services; EMR = electronic medical record; IT = information technology; NP = nurse practitioner; PA = physician's assistant; tPA = tissue plasminogen activator; U.S. = United States.

These barriers and facilitators are presented according to the CFIR constructs in **Table B-11**. The barriers are distributed across constructs, with most cited one to three times. The most frequently identified category of barrier was *knowledge & beliefs about the innovation* with this cited five times. The facilitators were more concentrated in categories that were also frequent in in- and out-patient studies; *access to knowledge and information* (11), *available resources* (9), and *patient needs & resources* (7). One construct that was more frequent in emergency care was *engaging* (7) which represents including the right people in the implementation, which may represent the need for emergency care to coordinate activities and processes across organizations such as EMS, multiple hospitals and outpatient care.

Table B-11. Emergency care: barriers and facilitators by CFIR constructs

Type	Facilitator or Barrier Name	Facilitator or Barrier Number of Mentions	Reference Number(s)
Barrier	Access to Knowledge & Information*	2	28, 88
	Adaptability [†]	1	170
	Available Resources [‡]	4	88, 134, 170
	Compatibility [§]	4	94, 134, 180
	Cost	1	180
	Executing	2	28

Type	Facilitator or Barrier Name	Facilitator or Barrier Number of Mentions	Reference Number(s)
	External Policy & Incentives [#]	1	77
	Implementation Climate ^{**}	1	88
	Knowledge & Beliefs about the Innovation ^{††}	5	94, 170, 180
	Networks & Communications ^{‡‡}	2	88, 123
	Readiness for Implementation ^{§§}	3	88, 180
	Relative Priority	3	28, 94, 180
Facilitators	Access to Knowledge & Information [*]	11	7, 14, 28, 33, 88, 134, 146, 147
	Available Resources [†]	9	7, 14, 33, 88, 147, 166, 169, 170
	Cost [‡]	2	77, 90
	Engaging ^{¶¶}	7	14, 28, 33, 147
	Executing ^{¶¶}	1	33
	External Policy & Incentives [#]	4	14, 77
	Formally Appointed Internal Implementation Leaders ^{###}	3	14, 33, 147
	Implementation Climate ^{**}	3	28, 94, 147
	Knowledge & Beliefs about the Innovation ^{††}	1	170
	Leadership Engagement ^{***}	4	28, 123, 147
	Patient Needs & Resources ^{†††}	7	6, 7, 88, 150, 160, 161
	Networks & Communications ^{‡‡}	4	6, 94, 170
	Planning ^{†††}	2	28, 147
	Readiness for Implementation ^{§§}	5	7, 14, 33, 147
	Reflecting & Evaluating ^{§§§}	4	14, 146, 147

* Access to digestible information and knowledge about the innovation and how to incorporate it into work tasks.

† Degree to which an innovation can be adapted, tailored, refined, or reinvented to meet local needs.

‡ Level of resources organizational dedicated for implementation and on-going operations including physical space and time.

§ Degree of tangible fit between meaning and values attached to the innovation by involved individuals, how those align with individuals' own norms, values, and perceived risks and needs, and how the innovation fits with existing workflows and systems.

† Costs of the innovation and costs associated with implementing the innovation including investment, supply, and opportunity costs.

¶ Carrying out or accomplishing the implementation according to plan.

External strategies to spread innovations including policy and regulations (governmental or other central entity), external mandates, recommendations and guidelines, pay-for-performance, collaboratives, and public or benchmark reporting.

** Absorptive capacity for change, shared receptivity of involved individuals to an innovation, and the extent to which use of that innovation will be rewarded, supported, and expected within their organization.

†† Individuals' attitudes toward and value placed on the innovation, as well as familiarity with facts, truths, and principles related to the innovation.

‡‡ Nature and quality of webs of social networks, and the nature and quality of formal and informal communications within an organization.

§§ Tangible and immediate indicators of organizational commitment to its decision to implement an innovation.

|| Individuals' shared perception of the importance of the implementation within the organization.

¶¶ Attracting and involving appropriate individuals in the implementation and use of the innovation through a combined strategy of social marketing, education, role modeling, training, and other similar activities.

Individuals from within the organization who have been formally appointed with responsibility for implementing an innovation as coordinator, project manager, team leader, or other similar role.

*** Commitment, involvement, and accountability of leaders and managers with the implementation of the innovation.

††† Extent to which patient needs, as well as barriers and facilitators to meet those needs, are accurately known and prioritized by the organization.

‡‡‡ Degree to which a scheme or method of behavior and tasks for implementing an innovation are developed in advance, and the quality of those schemes or methods.

§§§ Quantitative and qualitative feedback about the progress and quality of implementation accompanied with regular personal and team debriefing about progress and experience.

Education/Mentoring

Table B-12 includes a summary of information from 13 studies or reports on the use of telehealth for training and mentoring health care providers in rural areas (Additional details in **Appendix D-10**).

Nine of these studies are assessments of ECHO programs. ECHO combines didactic training, case presentations, virtual clinics and peer support to increase capacity and quality of care.^{9, 56, 73, 115, 127, 144, 157, 174} The ECHO programs evaluated in these articles address different topics with four focused on pain or opioids,^{56, 73, 144, 157} one each about Hepatitis C,¹²⁷ Multiple Sclerosis,⁹ and HIV in pregnancy,¹¹⁵ and one program that implemented ECHO as part of a larger expansion of specialist care.¹⁷⁴ While the subject matter and scale of these programs differed, facilitators, challenges and impact were similar. A frequently cited barrier was lack of clinician time to attend sessions or issues with scheduling. The evaluations also acknowledged that while ECHO could increase provider knowledge and skills it could not address all policy and practice barriers to practice change. Most ECHO program evaluations report a high level of stakeholder support and that the programs address rural participants needs for peer interaction, current knowledge, and access to experts. Evaluations of the impact of ECHO programs have documented that participants have changed practice, managed patients they would have referred, engaged in consultations with the expert faculty outside of the ECHO program, and become resources for other providers in their communities.

The three additional evaluations all address the rural clinicians' need for training in emergency care. One evaluation documented how training could be incorporated into consultations, reporting how this assured training was useful and relevant.¹⁸³ An experimental study tested whether training medical students using simulations for relatively rare emergency procedures could be managed by a remote expert trainer and documented that this was feasible and produced similar educational outcomes.⁶⁸ An assessment of remote training for emergency care in Australia documented that training could reduce professional isolation, but that sometimes topics were not relevant to rural working conditions.¹²¹

Table B-12. Findings of education and mentoring implementation studies

Topic Number of studies Intervention	Method N Location	Facilitators	Barriers	Impact
ECHO 9 Program with remote training, case reviews/clinics and peer interaction ECHO as part of a comprehensive program including e-consults	<p>Hepatitis C Survey to participants and non N=32 of 72 contacted; 15 facilities; Indian Health Service¹²⁷</p> <p>Multiple Sclerosis Pre questionnaire, program manager observations, exit interviews focused on case presentations N=8 of 24 clinicians, 13 practice sites; Mississippi, Washington State, Alaska, Montana, Idaho⁹</p> <p>HIV in pregnancy Survey and review of cases N=41 of 53 surveys, 11 cases Perinatal HIV; Washington State, Alaska, Montana, Idaho, Oregon, Colorado¹¹⁵</p> <p>Chronic Pain Content analysis N=Random selection of 25 of 67 cases; 406 data units; U.S. Connecticut. Single federally qualified health center¹⁵⁷</p> <p>Buprenorphine Interviews N=20; U.S. North Carolina¹⁴⁴</p> <p>Pain and Opioid Management with 2-3 in person supplemental training Questionnaires and focus groups N=38 participants; 2 workshops; New Mexico Indian Health Service⁷³</p> <p>Cancer Pain Survey of participants and non-participants N=24(46%) who attended education; 32 (34%) who attended case conferences; U.S. New Mexico⁵⁶</p> <p>Multiple Topics Surveys of clinician leads, administrative data N=180 from 87 sites; U.S. VA¹⁷⁴</p> <p>Melanoma Survey N = 10 Primary care providers; U.S. Missouri¹⁷</p>	<ul style="list-style-type: none"> • Preference or need for <ul style="list-style-type: none"> ◦ Collegial discussion with peers ◦ Being well-informed ◦ Access to experts • Willingness to present cases. conferences • High level of stakeholder support • Adaptability to local needs • Compatibility with existing workflows and systems • Feedback on progress • Quality of networks and communications • Leadership engagement • Access to Knowledge & Information • Networks & Communications 	<ul style="list-style-type: none"> • Lack of time to attend • Scheduling/time of day • Sessions seemed too long • Lack of local leadership support • Negative staff attitudes/stigma toward topic (Opioid use disorder) • ECHO cannot address policy issues (e.g., regulations and payment policy) • Assessment and screening skills require in-person add on 	<ul style="list-style-type: none"> • Increased ability to manage and treat, knowledge and confidence in existing treatment or changed care for case patient • Changed practice for other patients • Managed patients who would have been referred • Content consistent with guidelines • Participants engaged in additional consultations with faculty experts • Participants scored higher on competence than non-participants • Participants serve as resource for other providers

Topic Number of studies Intervention	Method N Location	Facilitators	Barriers	Impact
Telehealth Training 3 Training incorporated as part of the consultation ¹⁸³ Simulation training lead by remote expert ⁶⁸ Training needs assessment ¹²¹	Emergency Care Interviews N=18 hospitals Kansas, Minnesota, Nebraska, North Dakota, South Dakota Avera tele-training during consultation ¹⁸³ High-Acuity Low-Occurrence Procedures Assessments N=69 medical students Randomized to telehealth, in person and no training Newfoundland, Canada ⁶⁸ Emergency Care Interviews N=20 rural physician Australia ¹²¹	<ul style="list-style-type: none"> • Topics of formal training are useful • Real-time training can include exactly what is needed • Respect and supportive teaching from consultants • Interactive sessions preferred over didactic • Ability to reduce professional isolation • Broadcast live procedures • Technology that functions well. 	<ul style="list-style-type: none"> • Timing of live sessions • Workload constraints • Limited relevance to current working conditions 	<ul style="list-style-type: none"> • Asynchronous training courses included in program were valued • Training increased confidence and improved performance • Simulation training is feasible from a distance; results in similar education outcomes
Structured Dentist Network 1	Dentistry Interview and focus group N= 2 Dental specialists and 8 General dentists Australia ⁸³	<ul style="list-style-type: none"> • Networks & Communications 	<ul style="list-style-type: none"> • Access to Knowledge & Information • Planning • Available Resources 	<ul style="list-style-type: none"> • Lower proportion of patients treated at the Special Needs Unit, reflecting increased treatment at local facilities

Abbreviations: ECHO = Extension for Community Healthcare Outcomes; HIV = human immunodeficiency virus; U.S. = United States.

While telehealth interventions for education and mentoring differ in format and outcomes from those that are targeted to care of a specific patient, barriers and facilitators were able to be mapped to CFIR constructs as the framework was designed to be applicable across different types of interventions. **Table B-13** summarizes these. The most frequently cited barrier was *compatibility* (8), being when education did not correspond to the need or environment of the trainees. Not surprisingly, the most common facilitator was *access to knowledge and information* (11) as the telehealth education and mentoring programs were more likely to be implemented when they provided accessible information that could be directly incorporated into the trainees tasks and environment.

Table B-13. Education/mentoring: barriers and facilitators by CFIR constructs

Type	Facilitator or Barrier Name	Facilitator or Barrier Number of Mentions	Article Reference IDs
Barrier	Access to Knowledge & Information*	3	73, 83
	Adaptability*	1	121
	Available Resources [‡]	4	56, 73, 83, 127
	Compatibility [§]	8	121, 127, 144, 183
	External Policy & Incentives	2	144
	Implementation Climate [¶]	3	121, 127
	Knowledge & Beliefs about the Innovation [#]	1	144

Type	Facilitator or Barrier Name	Facilitator or Barrier Number of Mentions	Article Reference IDs
	Leadership Engagement**	1	144
	Planning	1	83
Facilitator	Access to Knowledge & Information*	12	17, 121, 127, 144, 174, 183
	Adaptability*	2	56, 174
	Available Resources‡	2	68, 121
	Compatibility§	2	121, 174
	Engaging††	3	157, 174
	Implementation Climate¶	3	73, 144, 174
	Leadership Engagement**	2	174
	Patient Needs & Resources‡‡	5	9, 115, 121, 127
	Networks & Communications§§	6	17, 83, 127, 157, 174, 183
	Reflecting & Evaluating	1	174
	Relative Priority¶¶	1	174

* Access to digestible information and knowledge about the innovation and how to incorporate it into work tasks.

† Degree to which an innovation can be adapted, tailored, refined, or reinvented to meet local needs.

‡ Level of resources organizational dedicated for implementation and on-going operations including physical space and time.

§ Degree of tangible fit between meaning and values attached to the innovation by involved individuals, how those align with individuals' own norms, values, and perceived risks and needs, and how the innovation fits with existing workflows and systems.

† External strategies to spread innovations including policy and regulations (governmental or other central entity), external mandates, recommendations and guidelines, pay-for-performance, collaboratives, and public or benchmark reporting.

¶ Absorptive capacity for change, shared receptivity of involved individuals to an innovation, and the extent to which use of that innovation will be rewarded, supported, and expected within their organization.

Individuals' attitudes toward and value placed on the innovation, as well as familiarity with facts, truths, and principles related to the innovation.

** Commitment, involvement, and accountability of leaders and managers with the implementation of the innovation.

†† Attracting and involving appropriate individuals in the implementation and use of the innovation through a combined strategy of social marketing, education, role modeling, training, and other similar activities.

‡‡ Extent to which patient needs, as well as barriers and facilitators to meet those needs, are accurately known and prioritized by the organization.

§§ Nature and quality of webs of social networks, and the nature and quality of formal and informal communications within an organization.

|| Quantitative and qualitative feedback about the progress and quality of implementation accompanied with regular personal and team debriefing about progress and experience.

¶¶ Individuals' shared perception of the importance of the implementation within the organization.

Appendix C. Included Studies List

1. Adcock AK, Choi J, Alvi M, et al. Expanding acute stroke care in rural America: a model for statewide success. *Telemed J E Health*. 2020 Jul;26(7):865-71. doi: 10.1089/tmj.2019.0087. PMID: 31596679.
2. Al Kasab S, Adams RJ, Debenham E, et al. Medical university of South Carolina telestroke: a telemedicine facilitated network for stroke treatment in South Carolina-a progress report. *Telemed J E Health*. 2017 08;23(8):674-7. doi: 10.1089/tmj.2016.0229. PMID: 28170316.
3. Alishahi Tabriz A, Turner K, Williams D, et al. Association of financial factors and telemedicine adoption for heart attack and stroke care among rural and urban hospitals: a longitudinal study. *Telemed J E Health*. 2021 Sep 24;24:24. doi: 10.1089/tmj.2021.0341. PMID: 34559014.
4. Alschuler KN, Stobbe GA, Hertz DP, et al. Impact of multiple sclerosis Project ECHO (extension for community healthcare outcomes) on provider confidence and clinical practice. *Int J MS Care*. 2019 Jul-Aug;21(4):143-50. doi: 10.7224/1537-2073.2018-014. PMID: 31474806.
5. Anderson EJ, Axon DR, Taylor AM, et al. Impact evaluation of a four-year academic-community partnership in provision of medication management and tertiary prevention services for rural patients with diabetes and/or hypertension. *Prev Med Rep*. 2020;17:101038. doi: 10.1016/j.pmedr.2019.101038. PMID: 31956473.
6. Arora S, Thornton K, Murata G, et al. Outcomes of treatment for hepatitis C virus infection by primary care providers. *N Engl J Med*. 2011 Jun 09;364(23):2199-207. doi: 10.1056/NEJMoa1009370. PMID: 21631316.
7. Astarcioglu MA, Sen T, Kilit C, et al. Time-to-reperfusion in STEMI undergoing interhospital transfer using smartphone and WhatsApp messenger. *Am J Emerg Med*. 2015 Oct;33(10):1382-4. doi: 10.1016/j.ajem.2015.07.029. PMID: 26299691.
8. Avent ML, Walker D, Yarwood T, et al. Implementation of a novel antimicrobial stewardship strategy for rural facilities utilising telehealth. *Int J Antimicrob Agents*. 2021 Jun;57(6):106346. doi: 10.1016/j.ijantimicag.2021.106346. PMID: 33882332.
9. Bagot KL, Cadilhac DA, Kim J, et al. Transitioning from a single-site pilot project to a state-wide regional telehealth service: the experience from the Victorian Stroke Telemedicine programme. *J Telemed Telecare*. 2017 Dec;23(10):850-5. doi: 10.1177/1357633X17734004. PMID: 29081268.
10. Banbury A, Smith AC, Mehrotra A, et al. A comparison study between metropolitan and rural hospital-based telehealth activity to inform adoption and expansion. *J Telemed Telecare*. 2021 Mar 26;1357633X21998201. doi: 10.1177/1357633X21998201. PMID: 33765879.
11. Barber M, Frieslick J, Maclean A, et al. The Western Isles Stroke Telerehabilitation (Specialist Medical Consultation) Service—implementation and evaluation. *Eur Res Telemed*. 2015;4(1):19-24.
12. Becevic M, Smith E, Golzy M, et al. Melanoma extension for community healthcare outcomes: a feasibility study of melanoma screening implementation in primary care settings. *Cureus*. 2021 May 29;13(5):e15322. doi: 10.7759/cureus.15322. PMID: 34221770.
13. Bellesheim KR, Kizzee RL, Curran A, et al. ECHO autism: integrating maintenance of certification with extension for community healthcare outcomes improves developmental screening. *J Dev Behav Pediatr*. 2020 Aug;41(6):420-7. doi: 10.1097/DBP.0000000000000796. PMID: 32735419.
14. Bennett-Levy J, Hawkins R, Perry H, et al. Online cognitive behavioural therapy training for therapists: outcomes, acceptability, and impact of support. *Aust Psychol*. 2012;47(3):174-82. doi: 10.1111/j.1742-9544.2012.00089.x.

15. Bladin CF, Moloczij N, Ermel S, et al. Victorian Stroke Telemedicine Project: implementation of a new model of translational stroke care for Australia. *Intern Med J*. 2015 Sep;45(9):951-6. doi: 10.1111/imj.12822. PMID: 26011155.
16. Boltz M, Cuellar NG, Cole C, et al. Comparing an on-site nurse practitioner with telemedicine physician support hospitalist programme with a traditional physician hospitalist programme. *J Telemed Telecare*. 2019 May;25(4):213-20. doi: 10.1177/1357633X18758744. PMID: 29498301.
17. Boman K, Olofsson M, Berggren P, et al. Robot-assisted remote echocardiographic examination and teleconsultation: a randomized comparison of time to diagnosis with standard of care referral approach. *JACC Cardiovasc Imaging*. 2014 Aug;7(8):799-803. doi: 10.1016/j.jcmg.2014.05.006. PMID: 25124011.
18. Bouchonville MF, Hager BW, Kirk JB, et al. Endo ECHO improves primary care provider and community health worker self-efficacy in complex diabetes management in medically underserved communities. *Endocr Pract*. 2018 Jan;24(1):40-6. doi: 10.4158/EP-2017-0079. PMID: 29368967.
19. Brunet N, Moore DT, Lendvai Wischik D, et al. Increasing buprenorphine access for veterans with opioid use disorder in rural clinics using telemedicine. *Subst Abus*. 2020 Feb 20:1-8. doi: 10.1080/08897077.2020.1728466. PMID: 32078492.
20. Brunetti ND, Dell'Anno A, Martone A, et al. Prehospital ECG transmission results in shorter door-to-wire time for STEMI patients in a remote mountainous region. *Am J Emerg Med*. 2020 02;38(2):252-7. doi: 10.1016/j.ajem.2019.04.046. PMID: 31079977.
21. Brunetti ND, Di Pietro G, Aquilino A, et al. Pre-hospital electrocardiogram triage with tele-cardiology support is associated with shorter time-to-balloon and higher rates of timely reperfusion even in rural areas: data from the Bari-Barletta/Andria/Trani public emergency medical service 118 registry on primary angioplasty in ST-elevation myocardial infarction. *Europ Heart J Acute Cardiovasc Care*. 2014 Sep;3(3):204-13. doi: 10.1177/2048872614527009. PMID: 24604713.
22. Buckley D, Weisser S. Videoconferencing could reduce the number of mental health patients transferred from outlying facilities to a regional mental health unit. *Aust N Z J Public Health*. 2012 Oct;36(5):478-82. doi: 10.1111/j.1753-6405.2012.00915.x. PMID: 23025371.
23. Cadilhac DA, Vu M, Bladin C. Experience with scaling up the Victorian Stroke Telemedicine programme. *J Telemed Telecare*. 2014 Oct;20(7):413-8. doi: 10.1177/1357633X14552389. PMID: 25400003.
24. Carter BL, Levy B, Gryzlak B, et al. Cluster-randomized trial to evaluate a centralized clinical pharmacy service in private family medicine offices. *Circ Cardiovasc Qual Outcomes*. 2018 06;11(6):e004188. doi: 10.1161/CIRCOUTCOMES.117.004188. PMID: 29884657.
25. Chen J, Amaize A, Barath D. Evaluating telehealth adoption and related barriers among hospitals located in rural and urban areas. *J Rural Health*. 2020 Nov 12;35(4):12-22. doi: 10.1111/jrh.12534. PMID: 33180363.
26. Cho JH, Kwon HS, Kim HS, et al. Effects on diabetes management of a health-care provider mediated, remote coaching system via a PDA-type glucometer and the internet. *J Telemed Telecare*. 2011;17(7):365-70. doi: 10.1258/jtt.2011.100913. PMID: 21933896.

27. Conn DK, Madan R, Lam J, et al. Program evaluation of a telepsychiatry service for older adults connecting a university-affiliated geriatric center to a rural psychogeriatric outreach service in Northwest Ontario, Canada. *Int Psychogeriatr*. 2013 Nov;25(11):1795-800. doi: 10.1017/S104161021300118X. PMID: 23870297.
28. Cronin T. Implementing a stroke program using telemedicine. *J Emerg Nurs*. 2013 Nov;39(6):613-8. doi: 10.1016/j.jen.2012.04.012. PMID: 22748668.
29. Crossland L, Askew D, Ware R, et al. Diabetic retinopathy screening and monitoring of early stage disease in Australian general practice: tackling preventable blindness within a chronic care model. *J Diabetes Res*. 2016;2016:8405395. doi: 10.1155/2016/8405395. PMID: 26798655.
30. Datta SK, Warshaw EM, Edison KE, et al. Cost and Utility Analysis of a Store-and-Forward Teledermatology Referral System: A Randomized Clinical Trial. *JAMA Dermatol*. 2015 Dec 1;151(12):1323-9. doi: 10.1001/jamadermatol.2015.2362. PMID: 26375589.
31. Davis TD, Deen T, Bryant-Bedell K, et al. Does minority racial-ethnic status moderate outcomes of collaborative care for depression? *Psychiatr Serv*. 2011 Nov;62(11):1282-8. doi: 10.1176/ps.62.11.pss6211_1282. PMID: 22211206.
32. de Batlle J, Massip M, Vargiu E, et al. Implementing mobile health-enabled integrated care for complex chronic patients: intervention effectiveness and cost-effectiveness study. *JMIR Mhealth Uhealth*. 2021 01 14;9(1):e22135. doi: 10.2196/22135. PMID: 33443486.
33. Demaerschalk BM, Raman R, Ernstrom K, et al. Efficacy of telemedicine for stroke: pooled analysis of the Stroke Team Remote Evaluation using a Digital Observation Camera (STRoKE DOC) and STRoKE DOC Arizona telestroke trials. *Telemed J E Health*. 2012 Apr;18(3):230-7. doi: 10.1089/tmj.2011.0116. PMID: 22400970.
34. Dharmar M, Kuppermann N, Romano PS, et al. Telemedicine consultations and medication errors in rural emergency departments. *Pediatrics*. 2013 Dec;132(6):1090-7. doi: 10.1542/peds.2013-1374. PMID: 24276844.
35. Dharmar M, Romano PS, Kuppermann N, et al. Impact of critical care telemedicine consultations on children in rural emergency departments. *Crit Care Med*. 2013 Oct;41(10):2388-95. doi: 10.1097/CCM.0b013e31828e9824. PMID: 23921273.
36. Dharmar M, Sadorra CK, Leigh P, et al. The financial impact of a pediatric telemedicine program: a children's hospital's perspective. *Telemed J E Health*. 2013 Jul;19(7):502-8. doi: 10.1089/tmj.2012.0266. PMID: 23837516.
37. Doyle C, Jackson D, Loi S, et al. Videoconferencing and telementoring about dementia care: evaluation of a pilot model for sharing scarce old age psychiatry resources. *Int Psychogeriatr*. 2016 09;28(9):1567-74. doi: 10.1017/S1041610216000740. PMID: 27189501.
38. Evans K, Lerch S, Boyce TW, et al. An innovative approach to enhancing access to medical screening for miners using a mobile clinic with telemedicine capability. *J Health Care Poor Underserved*. 2016;27(4A):62-72. PMID: 27818414.
39. Fairchild RM, Ferng-Kuo SF, Laws S, et al. Telehealth decreases rural emergency department wait times for behavioral health patients in a group of critical access hospitals. *Telemed J E Health*. 2019 12;25(12):1154-64. doi: 10.1089/tmj.2018.0227. PMID: 30735100.
40. Fortney JC, Maciejewski ML, Tripathi SP, et al. A budget impact analysis of telemedicine-based collaborative care for depression. *Med Care*. 2011 Sep;49(9):872-80. doi: 10.1097/MLR.0b013e31821d2b35. PMID: 21623240.

41. Fortney JC, Pyne JM, Edlund MJ, et al. A randomized trial of telemedicine-based collaborative care for depression. *J Gen Intern Med.* 2007 Aug;22(8):1086-93. doi: 10.1007/s11606-007-0201-9. PMID: 17492326.
42. Fortney JC, Pyne JM, Kimbrell TA, et al. Telemedicine-based collaborative care for posttraumatic stress disorder: a randomized clinical trial. *JAMA Psychiatry.* 2015 Jan;72(1):58-67. doi: 10.1001/jamapsychiatry.2014.1575. PMID: 25409287.
43. Fortney JC, Pyne JM, Mouden SB, et al. Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. *Am J Psychiatry.* 2013 Apr;170(4):414-25. doi: 10.1176/appi.ajp.2012.12050696. PMID: 23429924.
44. Fortney JC, Pyne JM, Mouden SB, et al. Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. *Focus.* 2017 Jul;15(3):361-72. doi: 10.1176/appi.focus.15306. PMID: 32015700.
45. Freeman RE, Boggs KM, Zachrison KS, et al. National study of telepsychiatry use in U.S. emergency departments. *Psychiatr Serv.* 2020 Jun 01;71(6):540-6. doi: 10.1176/appi.ps.201900237. PMID: 32019430.
46. Gadowski A, Anderson J, Chung YK, et al. Full agonist opioid prescribing by primary care clinicians after buprenorphine training. *Subst Abus.* 2020 Mar 09;1-7. doi: 10.1080/08897077.2020.1736709. PMID: 32150525.
47. Gallagher R, Giles M, Morison J, et al. Telehealth-based model of care redesign to facilitate local fitting and management of patients with a spinal fracture requiring a thoracic lumbar sacral orthosis in rural hospitals in New South Wales. *Aust J Rural Health.* 2018 Jun;26(3):181-7. doi: 10.1111/ajr.12407. PMID: 29573012.
48. Gill CJ, Le Ngoc B, Halim N, et al. The mCME project: a randomized controlled trial of an SMS-based continuing medical education intervention for improving medical knowledge among Vietnamese community based physicians' assistants. *PLoS One* 2016;11(11):e0166293. doi: 10.1371/journal.pone.0166293. PMID: 27861516.
49. Habashi P, Bouchard S, Nguyen GC. Transforming access to specialist care for inflammatory bowel disease: the PACE telemedicine program. *J Can Assoc Gastroenterol.* 2019 Dec;2(4):186-94. doi: 10.1093/jcag/gwy046. PMID: 31616860.
50. Hall RW, Hall-Barrow J, Garcia-Rill E. Neonatal regionalization through telemedicine using a community-based research and education core facility. *Ethn Dis.* 2010;20(1 Suppl 1):S1-136-40. PMID: 20521402.
51. Haozous E, Doorenbos AZ, Demiris G, et al. Role of telehealth/videoconferencing in managing cancer pain in rural American Indian communities. *Psycho-Oncol.* 2012 Feb;21(2):219-23. doi: 10.1002/pon.1887. PMID: 22271543.
52. Haque SN, DeStefano S, Banger A, et al. Factors influencing telehealth implementation and use in frontier critical access hospitals: qualitative study. *JMIR Form Res.* 2021 May 05;5(5):e24118. doi: 10.2196/24118. PMID: 33949958.
53. Harvey JB, Yeager BE, Cramer C, et al. The impact of telemedicine on pediatric critical care triage. *Pediatr Crit Care Med.* 2017 Nov;18(11):e555-e60. doi: 10.1097/PCC.0000000000001330. PMID: 28922271.
54. Haynes SC, Dharmar M, Hill BC, et al. The impact of telemedicine on transfer rates of newborns at rural community hospitals. *Acad Pediatr.* 2020 Jul;20(5):636-41. doi: 10.1016/j.acap.2020.02.013. PMID: 32081766.
55. Haynes SC, Hoffman KR, Patel S, et al. The use of telemedicine for stabilization of neonates transferred from rural community hospitals. *Telemed J E Health.* 2021 Mar 01;01:01. doi: 10.1089/tmj.2020.0503. PMID: 33691080.

56. Helwig SA, Ragoschke-Schumm A, Schwindling L, et al. Prehospital Stroke Management Optimized by Use of Clinical Scoring vs Mobile Stroke Unit for Triage of Patients With Stroke: A Randomized Clinical Trial. *JAMA Neurol.* 2019 Sep 3;76(12):1484-92. doi: 10.1001/jamaneurol.2019.2829. PMID: 31479116.
57. Hensel JM, Yang R, Rai M, et al. Optimizing electronic consultation between primary care providers and psychiatrists: mixed-methods study. *J Med Internet Res.* 2018 04 06;20(4):e124. doi: 10.2196/jmir.8943. PMID: 29625949.
58. Hilt RJ, Barclay RP, Bush J, et al. A statewide child telepsychiatry consult system yields desired health system changes and savings. *Telemed J E Health.* 2015 Jul;21(7):533-7. doi: 10.1089/tmj.2014.0161. PMID: 25799043.
59. Hofmeyer J, Leider JP, Satorius J, et al. Implementation of telemedicine consultation to assess unplanned transfers in rural long-term care facilities, 2012-2015: a pilot study. *J Am Med Dir Assoc.* 2016 11 01;17(11):1006-10. doi: 10.1016/j.jamda.2016.06.014. PMID: 27477614.
60. Holt T, Sari N, Hansen G, et al. Remote presence robotic technology reduces need for pediatric interfacility transportation from an isolated northern community. *Telemed J E Health.* 2018 11;24(11):927-33. doi: 10.1089/tmj.2017.0211. PMID: 29394155.
61. Hostutler CA, Valleru J, Maciejewski HM, et al. Improving pediatrician's behavioral health competencies through the Project ECHO teleconsultation model. *Clin Pediatr (Phila).* 2020 Jun 07;59(12):1049-57. doi: 10.1177/0009922820927018. PMID: 32506939.
62. Howland M, Tennant M, Bowen DJ, et al. Psychiatrist and psychologist experiences with telehealth and remote collaborative care in primary care: a qualitative study. *J Rural Health.* 2020 Oct 06;06:06. doi: 10.1111/jrh.12523. PMID: 33022079.
63. Jewer J, Parsons MH, Dunne C, et al. Evaluation of a mobile telesimulation unit to train rural and remote practitioners on high-acuity low-occurrence procedures: pilot randomized controlled trial. *J Med Internet Res.* 2019 08 06;21(8):e14587. doi: 10.2196/14587. PMID: 31389340.
64. Jhaveri D, Larkins S, Kelly J, et al. Remote chemotherapy supervision model for rural cancer care: perspectives of health professionals. *Eur J Cancer Care (Engl).* 2016 Jan;25(1):93-8. doi: 10.1111/ecc.12309. PMID: 25871852.
65. Jiang B, Bills M, Poon P. Integrated telehealth-assisted home-based specialist palliative care in rural Australia: a feasibility study. *J Telemed Telecare.* 2020 Oct 20;1357633X20966466. doi: 10.1177/1357633X20966466. PMID: 33079611.
66. Johansson AM, Lindberg I, Soderberg S. The views of health-care personnel about video consultation prior to implementation in primary health care in rural areas. *Prim Health Care Res Dev.* 2014 Apr;15(2):170-9. doi: 10.1017/S1463423613000030. PMID: 23402617.
67. Kahn JM, Le TQ, Barnato AE, et al. ICU Telemedicine and Critical Care Mortality: A National Effectiveness Study. *Med Care.* 2016 Mar;54(3):319-25. doi: 10.1097/MLR.0000000000000485. PMID: 26765148.
68. Katzman JG, Gygi K, Swift R, et al. How hands-on pain skills intensive trainings complement ECHO pain and opioid management programs: a program evaluation with the Indian Health Service. *Pain Med.* 2020 Jun 18;21(9):1769-78. doi: 10.1093/pm/pnaa151. PMID: 32556294.
69. Khalil H, Cullen M, Chambers H, et al. Implementation of a successful electronic wound documentation system in rural Victoria, Australia: a subject of collaboration and community engagement. *Int Wound J.* 2014 Jun;11(3):314-8. doi: 10.1111/iwj.12041. PMID: 23418740.

70. Kim EW, Teague-Ross TJ, Greenfield WW, et al. Telemedicine collaboration improves perinatal regionalization and lowers statewide infant mortality. *J Perinatol*. 2013 Sep;33(9):725-30. doi: 10.1038/jp.2013.37. PMID: 23579490.
71. Kim H, Jhoo JH, Jang JW. The effect of telemedicine on cognitive decline in patients with dementia. *J Telemed Telecare*. 2017 Jan;23(1):149-54. doi: 10.1177/1357633X15615049. PMID: 26541171.
72. Kulcsar M, Gilchrist S, George MG. Improving stroke outcomes in rural areas through telestroke programs: an examination of barriers, facilitators, and state policies. *Telemed J E Health*. 2014 Jan;20(1):3-10. doi: 10.1089/tmj.2013.0048. PMID: 24286197.
73. Kuperman EF, Linson EL, Klefstad K, et al. The virtual hospitalist: a single-site implementation bringing hospitalist coverage to critical access hospitals. *J Hosp Med*. 2018 11 01;13(11):759-63. doi: 10.12788/jhm.3061. PMID: 30255859.
74. Kwak MY, Hwang EJ, Lee TH. Effects of the Physician-Primary-Healthcare Nurse Telemedicine Model (P-NTM) on medication adherence and Health-Related Quality of Life (HRQoL) of patients with chronic disease at remote rural areas. *Int J Environ Res Public Health*. 2021 03 03;18(5):03. doi: 10.3390/ijerph18052502. PMID: 33802513.
75. Leshner AP, Fakhry SM, DuBose-Morris R, et al. Development and evolution of a statewide outpatient consultation service: leveraging telemedicine to improve access to specialty care. *Popul Health Manag*. 2020 02;23(1):20-8. doi: 10.1089/pop.2018.0212. PMID: 31161963.
76. Levin K, Madsen JR, Petersen I, et al. Telemedicine diabetes consultations are cost-effective, and effects on essential diabetes treatment parameters are similar to conventional treatment: 7-year results from the Svendborg Telemedicine Diabetes Project. *J Diabetes Sci Technol*. 2013 May 1;7(3):587-95. doi: 10.1177/193229681300700302. PMID: 23759390.
77. Liddy C, Moroz I, Keely E, et al. Understanding the impact of a multispecialty electronic consultation service on family physician referral rates to specialists: a randomized controlled trial using health administrative data. *Trials*. 2019 Jun 10;20(1):348. doi: 10.1186/s13063-019-3393-5. PMID: 31182123.
78. Lim M, Liberali SAC, Calache H, et al. Specialist networks influence clinician willingness to treat individuals with special needs. *JDR Clin Trans Res*. 2021 Jun 21:23800844211020250. doi: 10.1177/23800844211020250. PMID: 34148391.
79. Lindauer A, Wild K, Natanson A, et al. Dementia 360 ECHO: using technology to facilitate diagnosis and treatment. *Gerontol Geriatr Educ*. 2020 Oct 20:1-7. doi: 10.1080/02701960.2020.1835658. PMID: 33078687.
80. Lingum NR, Sokoloff LG, Meyer RM, et al. Building long-term care staff capacity during COVID-19 through just-in-time learning: evaluation of a modified ECHO model. *J Am Med Dir Assoc*. 2021 Feb;22(2):238-44.e1. doi: 10.1016/j.jamda.2020.10.039. PMID: 33238143.
81. Liou JK, Soon MS, Chen CH, et al. Shared care combined with telecare improves glycemic control of diabetic patients in a rural underserved community. *Telemed J E Health*. 2014 Feb;20(2):175-8. doi: 10.1089/tmj.2013.0037. PMID: 24320193.
82. Long MC, Angtuaco T, Lowery C. Ultrasound in telemedicine: its impact in high-risk obstetric health care delivery. *Ultrasound Q*. 2014 Sep;30(3):167-72. doi: 10.1097/RUQ.0000000000000073. PMID: 25148484.
83. Lucas JAM, Day K, Honey MLL. Clinician's perceptions of telehealth for emergency care on the west coast of New Zealand: findings of a descriptive study. *Emergency Nurse New Zealand*. 2016:6-10.

84. Luscombe GM, Hawthorn J, Wu A, et al. 'Empowering clinicians in smaller sites': A qualitative study of clinician's experiences with a rural Virtual Paediatric Feeding Clinic. *Aust J Rural Health*. 2021 Sep 07;07:07. doi: 10.1111/ajr.12781. PMID: 34490941.
85. MacKinney AC, Ward MM, Ullrich F, et al. The business case for tele-emergency. *Telemed J E Health*. 2015 Dec;21(12):1005-11. doi: 10.1089/tmj.2014.0241. PMID: 26226603.
86. Mahmoud H, Vogt EL, Dahdouh R, et al. Using continuous quality improvement to design and implement a telepsychiatry program in rural Illinois. *Psychiatr Serv*. 2020 Apr 15;71(8):860-3. doi: 10.1176/appi.ps.201900231. PMID: 32290807.
87. Makkar A, McCoy M, Hallford G, et al. A hybrid form of telemedicine: a unique way to extend intensive care service to neonates in medically underserved areas. *Telemed J E Health*. 2018 09;24(9):717-21. doi: 10.1089/tmj.2017.0155. PMID: 29298407.
88. Makkar A, McCoy M, Hallford G, et al. Evaluation of neonatal services provided in a level II NICU utilizing hybrid telemedicine: a prospective study. *Telemed J E Health*. 2020 Feb;26(2):176-83. doi: 10.1089/tmj.2018.0262. PMID: 30835166.
89. Marsh-Feiley G, Eadie L, Wilson P. Paramedic and physician perspectives on the potential use of remotely supported prehospital ultrasound. *Rural Remote Health*. 2018 09;18(3):4574. doi: 10.22605/RRH4574. PMID: 30207737.
90. May S, Jonas K, Fehler GV, et al. Challenges in current nursing home care in rural Germany and how they can be reduced by telehealth - an exploratory qualitative pre-post study. *BMC Health Serv Res*. 2021 Sep 06;21(1):925. doi: 10.1186/s12913-021-06950-y. PMID: 34488746.
91. McFarland LV, Raugi GJ, Taylor LL, et al. Implementation of an education and skills programme in a teledermatology project for rural veterans. *J Telemed Telecare*. 2012 Mar;18(2):66-71. doi: 10.1258/jtt.2011.110518. PMID: 22198956.
92. McGill AF, North JB. Teleconference fracture clinics: a trial for rural hospitals. *ANZ J Surg*. 2012 Jan-Feb;82(1-2):2-3. doi: 10.1111/j.1445-2197.2011.05952.x. PMID: 22507484.
93. McLendon SF, Wood FG, Stanley N. Enhancing diabetes care through care coordination, telemedicine, and education: evaluation of a rural pilot program. *Public Health Nurs*. 2019 05;36(3):310-20. doi: 10.1111/phn.12601. PMID: 30868661.
94. McWilliams T, Hendricks J, Twigg D, et al. Burns education for non-burn specialist clinicians in Western Australia. *Burns*. 2015 Mar;41(2):301-7. doi: 10.1016/j.burns.2014.06.015. PMID: 25239848.
95. Miller AC, Ward MM, Ullrich F, et al. Emergency department telemedicine consults are associated with faster time-to-electrocardiogram and time-to-fibrinolysis for myocardial infarction patients. *Telemed J E Health*. 2020 Feb 28;26(12):1440-8. doi: 10.1089/tmj.2019.0273. PMID: 32109200.
96. Miyamoto S, Dharmar M, Boyle C, et al. Impact of telemedicine on the quality of forensic sexual abuse examinations in rural communities. *Child Abuse Negl*. 2014 Sep;38(9):1533-9. doi: 10.1016/j.chiabu.2014.04.015. PMID: 24841062.
97. Moffatt JJ, Eley DS. Barriers to the uptake of telemedicine in Australia--a view from providers. *Rural Remote Health*. 2011;11(2):1581. PMID: 21385004.
98. Mohr NM, Campbell KD, Swanson MB, et al. Provider-to-provider telemedicine improves adherence to sepsis bundle care in community emergency departments. *J Telemed Telecare*. 2020 Jan 05;1-9. doi: 10.1177/1357633X19896667. PMID: 31903840.
99. Mohr NM, Vakkalanka JP, Harland KK, et al. Telemedicine use decreases rural emergency department length of stay for transferred North Dakota trauma patients. *Telemed J E Health*. 2018 03;24(3):194-202. doi: 10.1089/tmj.2017.0083. PMID: 28731843.

100. Mohr NM, Young T, Harland KK, et al. Emergency department telemedicine shortens rural time-to-provider and emergency department transfer times. *Telemed J E Health*. 2018 08;24(8):582-93. doi: 10.1089/tmj.2017.0262. PMID: 29293413.
101. Mohr NM, Young T, Harland KK, et al. Telemedicine is associated with faster diagnostic imaging in stroke patients: a cohort study. *Telemed J E Health*. 2019 02;25(2):93-100. doi: 10.1089/tmj.2018.0013. PMID: 29958087.
102. Momin B, Mera J, Essex W, et al. Implementation of liver cancer education among health care providers and community coalitions in the Cherokee nation. *Prev Chronic Dis*. 2019;16(E112):1-7. doi: 10.5888/pcd16.180671. PMID: 31441770.
103. Morrisette S, Pearlman RL, Kovar M, et al. Attitudes and perceived barriers toward store-and-forward teler dermatology among primary care providers of the rural Mississippi. *Arch Dermatol Res*. 2021 Feb 25;25:25. doi: 10.1007/s00403-021-02208-z. PMID: 33630147.
104. Mundt AP, Irrarrazaval M, Martinez P, et al. Telepsychiatry consultation for primary care treatment of children and adolescents receiving child protective services in Chile: mixed methods feasibility study. *JMIR Public Health Surveill*. 2021 Jul 22;7(7):e25836. doi: 10.2196/25836. PMID: 34292164.
105. Myers K, Vander Stoep A, Zhou C, et al. Effectiveness of a telehealth service delivery model for treating attention-deficit/hyperactivity disorder: a community-based randomized controlled trial. *J Am Acad Child Adolesc Psychiatry*. 2015 Apr;54(4):263-74. doi: 10.1016/j.jaac.2015.01.009. PMID: 25791143.
106. Nagao KJ, Koschel A, Haines HM, et al. Rural Victorian telestroke project. *Intern Med J*. 2012 Oct;42(10):1088-95. doi: 10.1111/j.1445-5994.2011.02603.x. PMID: 21981424.
107. Nagayoshi Y, Oshima S, Ogawa H. Clinical impact of telemedicine network system at rural hospitals without on-site cardiac surgery backup. *Telemed J E Health*. 2016 11;22(11):960-4. PMID: 27148833.
108. Narva AS, Romancito G, Faber T, et al. Managing CKD by telemedicine: the Zuni telenephrology clinic. *Adv Chronic Kidney Dis*. 2017 Jan;24(1):6-11. doi: 10.1053/j.ackd.2016.11.019. PMID: 28224944.
109. Natafagi N, Mohr NM, Wittrock A, et al. The association between telemedicine and emergency department (ED) disposition: a stepped wedge design of an ED-based telemedicine program in critical access hospitals. *J Rural Health*. 2020 Jun;36(3):360-70. doi: 10.1111/jrh.12370. PMID: 31013552.
110. Ness TE, Annese MF, Martinez-Paz N, et al. Using an innovative telehealth model to support community providers who deliver perinatal HIV care. *AIDS Educ Prev*. 2017 12;29(6):516-26. doi: 10.1521/aeap.2017.29.6.516. PMID: 29283274.
111. Newell MC, Strauss CE, Freier T, et al. Design and initial results of the Minneapolis Heart Institute TeleHeart program. *Circ Cardiovasc Qual Outcomes*. 2017;10:1-4. doi: 10.1161/CIRCOUTCOMES.117.003904. PMID: 29021334.
112. Nipp CM, Vogtle LK, Warren M. Clinical application of low vision rehabilitation strategies after completion of a computer-based training module. *Occup Ther Health Care*. 2014 Jul;28(3):296-305. doi: 10.3109/07380577.2014.908335. PMID: 24971896.
113. Nqala MO, Rout CC, Aldous CM. Remote clinical support by telephone for rural district hospital medical officers in the Eastern Cape. *S Afr Fam Pract*. 2015;57(5):286-90. doi: 10.1080/20786190.2015.1055671.

114. Olenik K, Lehr B. Counteracting brain drain of health professionals from rural areas via teleconsultation: analysis of the barriers and success factors of teleconsultation. *J Public Health*. 2013;21(4):357-64. doi: 10.1007/s10389-013-0565-8.
115. Painter JT, Fortney JC, Austen MA, et al. Cost-effectiveness of telemedicine-based collaborative care for posttraumatic stress disorder. *Psychiatr Serv*. 2017 Nov 01;68(11):1157-63. doi: 10.1176/appi.ps.201600485. PMID: 28669290.
116. Pandit T, Ray RA, Sabesan S. Managing emergencies in rural North Queensland: the feasibility of teletraining. *Int J Telemed Appl*. 2018;2018doi: 10.1155/2018/8421346. PMID: 29849606.
117. Panlaqui OM, Broadfield E, Champion R, et al. Outcomes of telemedicine intervention in a regional intensive care unit: a before and after study. *Anaesth Intensive Care*. 2017 09;45(5):605-10. PMID: 28911290.
118. Parks J, Hunter A, Taylor A, et al. Design, development and implementation of the virtual, coordination, access, referral and escalation service in western New South Wales. *Aust J Rural Health*. 2021 Sep 29;29:29. doi: 10.1111/ajr.12800. PMID: 34586702.
119. Pedragosa A, Alvarez-Sabin J, Rubiera M, et al. Impact of telemedicine on acute management of stroke patients undergoing endovascular procedures. *Cerebrovasc Dis*. 2012;34(5-6):436-42. doi: 10.1159/000345088. PMID: 23207552.
120. Peracca SB, Jackson GL, Lamkin RP, et al. Implementing teledermatology for rural veterans: an evaluation using the RE-AIM framework. *Telemed J E Health*. 2020 Apr 27:1-9. doi: 10.1089/tmj.2020.0013. PMID: 32343924.
121. Pimentel CB, Gately M, Barczy SR, et al. GRECC connect: geriatrics telehealth to empower health care providers and improve management of older veterans in rural communities. *Fed Pract*. 2019 Oct;36(10):464-70. PMID: 31768097.
122. Pindyck T, Kalishman S, Flatow-Trujillo L, et al. Treating hepatitis C in American Indians/Alaskan Natives: a survey of Project ECHO(R) (Extension for Community Healthcare Outcomes) utilization by Indian Health Service providers. *SAGE Open Med*. 2015;3:2050312115612805. doi: 10.1177/2050312115612805. PMID: 26770809.
123. Poulson LK, Nissen L, Coombes I. Pharmaceutical review using telemedicine-a before and after feasibility study. *J Telemed Telecare*. 2010;16(2):95-9. doi: 10.1258/jtt.2009.090716. PMID: 20139138.
124. Powell KR, Alexander GL. Consequences of rapid telehealth expansion in nursing homes: promise and pitfalls. *Appl Clin Inform*. 2021 Aug;12(4):933-43. doi: 10.1055/s-0041-1735974. PMID: 34614517.
125. Puskar KR, Heeyoung L, Mitchell AM, et al. Interprofessional collaborative education for substance use screening: rural areas and challenges. *Online J Rural Nurs Health Care*. 2016;16(1):76-96. doi: 10.14574/ojrnhc.v16i1.385.
126. Pyne JM, Fortney JC, Mouden S, et al. Cost-effectiveness of on-site versus off-site collaborative care for depression in rural FQHCs. *Psychiatr Serv*. 2015 May 01;66(5):491-9. doi: 10.1176/appi.ps.201400186. PMID: 25686811.
127. Pyne JM, Fortney JC, Tripathi SP, et al. Cost-effectiveness analysis of a rural telemedicine collaborative care intervention for depression. *Arch Gen Psychiatry*. 2010 Aug;67(8):812-21. doi: 10.1001/archgenpsychiatry.2010.82. PMID: 20679589.
128. Ramnath VR, Khazeni N. Centralized monitoring and virtual consultant models of tele-ICU care: a side-by-side review. *Telemed J E Health*. 2014 Oct;20(10):962-71. doi: 10.1089/tmj.2014.0024. PMID: 25225795.

129. Ray KN, Felmet KA, Hamilton MF, et al. Clinician attitudes toward adoption of pediatric emergency telemedicine in rural hospitals. *Pediatr Emerg Care*. 2017 Apr;33(4):250-7. doi: 10.1097/PEC.0000000000000583. PMID: 26785087.
130. Ritter LA, Robinette TR, Cofano J. Evaluation of a statewide telemedicine program. *CJHP*. 2010;8(1):1-9. doi: 10.32398/cjhp.v8i1.2025.
131. Robinson T, Hills D, Kelly B. The evaluation of an online orientation to rural mental health practice in Australia. *J Psychiatr Ment Health Nurs*. 2011;18(7):629-36. doi: 10.1111/j.1365-2850.2011.01712.x. PMID: 21848598.
132. Rogove HJ, McArthur D, Demaerschalk BM, et al. Barriers to telemedicine: survey of current users in acute care units. *Telemed J E Health*. 2012 Jan-Feb;18(1):48-53. doi: 10.1089/tmj.2011.0071. PMID: 22082107.
133. Rojas G, Guajardo V, Martinez P, et al. A remote collaborative care program for patients with depression living in rural areas: open-label trial. *J Med Internet Res*. 2018 04 30;20(4):e158. doi: 10.2196/jmir.8803. PMID: 29712627.
134. Rossaro L, Torruellas C, Dhaliwal S, et al. Clinical outcomes of hepatitis C treated with pegylated interferon and ribavirin via telemedicine consultation in Northern California. *Dig Dis Sci*. 2013 Dec;58(12):3620-5. doi: 10.1007/s10620-013-2810-y. PMID: 24154637.
135. Sabesan S, Senko C, Schmidt A, et al. Enhancing chemotherapy capabilities in rural hospitals: implementation of a telechemotherapy model (QReCS) in North Queensland, Australia. *J Oncol Pract*. 2018 07;14(7):e429-e37. doi: 10.1200/JOP.18.00110. PMID: 29996068.
136. Sairanen T, Soinila S, Nikkanen M, et al. Two years of Finnish Telestroke: thrombolysis at spokes equal to that at the hub. *Neurology*. 2011 Mar 29;76(13):1145-52. doi: 10.1212/WNL.0b013e318212a8d4. PMID: 21368283.
137. Seibert PS, Reddy T, Whitmore T, et al. The use of telemedicine to train perioperative nurses in rural settings. *J Telemed Telecare*. 2013 Sep;19(6):311-4. doi: 10.1177/1357633X13501777. PMID: 24163294.
138. Shaikh U, Nettiksimmons J, Joseph JG, et al. Collaborative practice improvement for childhood obesity in rural clinics: the Healthy Eating Active Living Telehealth Community of Practice (HEALTH COP). *Am J Med Qual*. 2014 Nov-Dec;29(6):467-75. doi: 10.1177/1062860613506252. PMID: 24170936.
139. Shea CM, Gertner AK, Green SL. Barriers and perceived usefulness of an ECHO intervention for office-based buprenorphine treatment for opioid use disorder in North Carolina: a qualitative study. *Subst Abus*. 2019 Dec 06:1-11. doi: 10.1080/08897077.2019.1694617. PMID: 31809679.
140. Sicotte C, Moqadem K, Vasilevsky M, et al. Use of telemedicine for haemodialysis in very remote areas: the Canadian First Nations. *J Telemed Telecare*. 2011;17(3):146-9. doi: 10.1258/jtt.2010.100614. PMID: 21303935.
141. Simpson AN, Harvey JB, DiLembo SM, et al. Population health indicators associated with a statewide telestroke program. *Telemed J E Health*. 2020 Feb 11;26(9):1126-33. doi: 10.1089/tmj.2019.0204. PMID: 32045330.
142. Singh R, Mathiassen L, Switzer JA, et al. Assimilation of web-based urgent stroke evaluation: a qualitative study of two networks. *JMIR Med Inform*. 2014 Apr 15;2(1):e6. doi: 10.2196/medinform.3028. PMID: 25601232.
143. Sockalingam S, Arena A, Serhal E, et al. Building provincial mental health capacity in primary care: an evaluation of a Project ECHO mental health program. *Acad Psychiatry*. 2018 Aug;42(4):451-7. doi: 10.1007/s40596-017-0735-z. PMID: 28593537.

144. Spivak S, Spivak A, Cullen B, et al. Telepsychiatry use in U.S. mental health facilities, 2010-2017. *Psychiatr Serv*. 2020 02 01;71(2):121-7. doi: 10.1176/appi.ps.201900261. PMID: 31615370.
145. Sterling SA, Seals SR, Jones AE, et al. The impact of the TelEmergency program on rural emergency care: an implementation study. *J Telemed Telecare*. 2017 Jul;23(6):588-94. doi: 10.1177/1357633X16657499. PMID: 27470505.
146. Stevens G, Loh J, Kolbe J, et al. Comparison of recommendations for radiotherapy from two contemporaneous thoracic multidisciplinary meeting formats: co-located and video conference. *Intern Med J*. 2012 Nov;42(11):1213-8. doi: 10.1111/j.1445-5994.2012.02817.x. PMID: 22530597.
147. Summerhayes C, McGee JA, Cooper RJ, et al. Introducing leg ulcer telemedicine into rural general practice. *Wounds UK*. 2012;8(2):28-36.
148. Swanson MB, Miller AC, Ward MM, et al. Emergency department telemedicine consults decrease time to interpret computed tomography of the head in a multi-network cohort. *J Telemed Telecare*. 2019 Nov 04;1-10. doi: 10.1177/1357633X19877746. PMID: 31684801.
149. Tande AJ, Berbari EF, Ramar P, et al. Association of a remotely offered infectious diseases eConsult service with improved clinical outcomes. *Open Forum Infect Dis*. 2020 Jan:1-4. doi: 10.1093/ofid/ofaa003. PMID: 31988969.
150. Taylor-Gjevne R, Nair B, Bath B, et al. Addressing rural and remote access disparities for patients with inflammatory arthritis through video-conferencing and innovative inter-professional care models. *Musculoskelet* 2018 03;16(1):90-5. doi: 10.1002/msc.1215. PMID: 28967235.
151. Thaker DA, Monypenny R, Olver I, et al. Cost savings from a telemedicine model of care in northern Queensland, Australia. *Med J Aust*. 2013 Sep 16;199(6):414-7. PMID: 24033216.
152. Thies KM, Anderson D, Beals-Reid C. Project ECHO chronic pain: a qualitative analysis of recommendations by expert faculty. *Pain Med*. 2019 07 01;20(7):1450-2. doi: 10.1093/pm/pnz024. PMID: 30840075.
153. Tran L, Feldman R, Riley T, 3rd, et al. Association of the extension for community healthcare outcomes project with use of direct-acting antiviral treatment among US adults with hepatitis C. *JAMA Netw Open*. 2021 Jul 01;4(7):e2115523. doi: 10.1001/jamanetworkopen.2021.15523. PMID: 34213557.
154. Tripod M, Tait M, Bracey J, et al. The use of telemedicine decreases unnecessary hand trauma transfers. *Hand*. 2020 May;15(3):422-7. doi: 10.1177/1558944718810877. PMID: 30417694.
155. Trondsen MV, Bolle SR, Stensland GO, et al. Video-confidence: a qualitative exploration of videoconferencing for psychiatric emergencies. *BMC Health Serv Res*. 2014 Oct 31;14:544. doi: 10.1186/s12913-014-0544-y. PMID: 25359404.
156. Trondsen MV, Tjora A, Broom A, et al. The symbolic affordances of a video-mediated gaze in emergency psychiatry. *Soc Sci Med*. 2018 01;197:87-94. doi: 10.1016/j.socscimed.2017.11.056. PMID: 29222999.
157. Tynan A, Deeth L, McKenzie D. An integrated oral health program for rural residential aged care facilities: a mixed methods comparative study. *BMC Health Serv Res*. 2018 07 03;18(1):515. doi: 10.1186/s12913-018-3321-5. PMID: 29970073.
158. Tynan A, Deeth L, McKenzie D, et al. Integrated approach to oral health in aged care facilities using oral health practitioners and teledentistry in rural Queensland. *Aust J Rural Health*. 2018 Apr 16;26:290-4. doi: 10.1111/ajr.12410. PMID: 29660771.

159. Vakkalanka JP, Harland KK, Wittrock A, et al. Telemedicine is associated with rapid transfer and fewer involuntary holds among patients presenting with suicidal ideation in rural hospitals: a propensity matched cohort study. *J Epidemiol Community Health*. 2019 11;73(11):1033-9. doi: 10.1136/jech-2019-212623. PMID: 31492762.
160. van der Pol M, McKenzie L. Costs and benefits of tele-endoscopy clinics in a remote location. *J Telemed Telecare*. 2010;16(2):89-94. doi: 10.1258/jtt.2009.090609. PMID: 20139140.
161. Ward MM, Ullrich F, MacKinney AC, et al. Tele-emergency utilization: in what clinical situations is tele-emergency activated? *J Telemed Telecare*. 2016 Jan;22(1):25-31. doi: 10.1177/1357633X15586319. PMID: 26026189.
162. Watts SA, Roush L, Julius M, et al. Improved glycemic control in veterans with poorly controlled diabetes mellitus using a Specialty Care Access Network-Extension for Community Healthcare Outcomes model at primary care clinics. *J Telemed Telecare*. 2016 Jun;22(4):221-4. doi: 10.1177/1357633X15598052. PMID: 26253745.
163. Weaver MS, Neumann ML, Navaneethan H, et al. Human touch via touchscreen: rural nurses' experiential perspectives on telehealth use in pediatric hospice care. *J Pain Symptom Manage*. 2020 Jun 07;60(5):1027-33. doi: 10.1016/j.jpainsymman.2020.06.003. PMID: 32525081.
164. Weigel P, Bhagianadh D, Merchant KA, et al. Tele-emergency behavioural health in rural and underserved areas. *J Telemed Telecare*. 2019 Nov 14;1-10. doi: 10.1177/1357633X19887027. PMID: 31726903.
165. Weigel PA, Merchant KA, Wittrock A, et al. Paediatric tele-emergency care: a study of two delivery models. *J Telemed Telecare*. 2019 Apr 09;1-9. doi: 10.1177/1357633X19839610. PMID: 30966860.
166. White AH, Crowther SA, Lee SH. Supporting rural midwifery practice using a mobile health (mHealth) intervention: a qualitative descriptive study. *Rural Remote Health*. 2019 08;19(3):5294. doi: 10.22605/RRH5294. PMID: 31446762.
167. Whited JD, Warshaw EM, Kapur K, et al. Clinical course outcomes for store and forward teledermatology versus conventional consultation: a randomized trial. *J Telemed Telecare*. 2013 Jun;19(4):197-204. doi: 10.1177/1357633X13487116. PMID: 23666440.
168. Williams D, Simpson AN, King K, et al. Do Hospitals Providing Telehealth in Emergency Departments Have Lower Emergency Department Costs? *Telemed J E Health*. 2020 Nov 13;13:13. doi: 10.1089/tmj.2020.0349. PMID: 33185503.
169. Williams KM, Kirsh S, Aron D, et al. Evaluation of the Veterans Health Administration's specialty care transformational initiatives to promote patient-centered delivery of specialty care: a mixed-methods approach. *Telemed J E Health*. 2017 07;23(7):577-89. doi: 10.1089/tmj.2016.0166. PMID: 28177858.
170. Wilson BM, Banks RE, Crnich CJ, et al. Changes in antibiotic use following implementation of a telehealth stewardship pilot program. *Infect Control Hosp Epidemiol*. 2019 07;40(7):810-4. doi: 10.1017/ice.2019.128. PMID: 31172905.
171. Wood PR, Caplan L. Outcomes, satisfaction, and costs of a rheumatology telemedicine program: a longitudinal evaluation. *J Clin Rheumatol*. 2019 Jan;25(1):41-4. doi: 10.1097/RHU.0000000000000778. PMID: 30461466.
172. Wood T, Freeman S, Banner D, et al. Exploring user perspectives of factors associated with use of teletrauma in rural areas. *Australas Emerg Care*. 2021 May 07;07:07. doi: 10.1016/j.aucec.2021.04.001. PMID: 33972192.

173. Yam P, Fales D, Jemison J, et al. Implementation of an antimicrobial stewardship program in a rural hospital. *Am J Health-Syst Pharm*. 2012 Jul 01;69(13):1142-8. doi: 10.2146/ajhp110512. PMID: 22722593.
174. Yan LL, Gong E, Gu W, et al. Effectiveness of a primary care-based integrated mobile health intervention for stroke management in rural China (SINEMA): A cluster-randomized controlled trial. *PLoS Med*. 2021 04;18(4):e1003582. doi: 10.1371/journal.pmed.1003582. PMID: 33909607.
175. Zachrison KS, Boggs KM, Hayden EM, et al. Understanding barriers to telemedicine implementation in rural emergency departments. *Ann Emerg Med*. 2020 03;75(3):392-9. doi: 10.1016/j.annemergmed.2019.06.026. PMID: 31474481.
176. Zachrison KS, Hayden EM, Schwamm LH, et al. Characterizing New England emergency departments by telemedicine use. *West J Emerg Med*. 2017 Oct;18(6):1055-60. doi: 10.5811/westjem.2017.8.34880. PMID: 29085537.
177. Zhang D, Wang G, Zhu W, et al. Expansion of telestroke services improves quality of care provided in super rural areas. *Health Aff (Millwood)*. 2018 12;37(12):2005-13. doi: 10.1377/hlthaff.2018.05089. PMID: 30633675.
178. Zhu X, Merchant KAS, Mohr NM, et al. Real-time learning through telemedicine enhances professional training in rural emergency departments. *Telemed J E Health*. 2020 Jun 17;17:1-7. doi: 10.1089/tmj.2020.0042. PMID: 32552479.
179. Zittleman L, Curcija K, Sutter C, et al. Building capacity for medication assisted treatment in rural primary care practices: the IT MATTTRs practice team training. *J Prim Care Community Health*. 2020 Jan-Dec;11:2150132720953723. doi: 10.1177/2150132720953723. PMID: 32909491.

Appendix D. Evidence Tables

See associated Excel file at <https://effectivehealthcare.ahrq.gov/products/rural-telehealth/research>.

Appendix E. Risk of Bias Assessment

See associated Excel file at <https://effectivehealthcare.ahrq.gov/products/rural-telehealth/research>.

Appendix F. Details on Strength of Evidence

Table F-1. Strength of evidence of selected outcomes (Key Question 2)

Topic	Outcome	Number of Studies (Combined N)	Study Limitations (Low, Medium, High)	Directness (Direct, Indirect)	Consistency (Consistent, Inconsistent, Unknown)	Precision (Precise, Imprecise)	Reporting Bias (Not Detected, Possible, Suspected)	Conclusion	Strength of Evidence Grade (Insufficient, Low, Moderate, High)
Inpatient: Teleconsultation	Mortality	3 (1841) ^{21, 78, 154}	Medium	Direct	Inconsistent	Imprecise	Not detected	Unclear effect	Insufficient
	Length of Stay	6 (2,113) ^{13, 16, 21, 52, 78, 154}	Medium	Direct	Inconsistent	Precise	Not detected	No difference in length of stay with telehealth consultations	Low
	Transfers	3 (2208) ^{27, 78, 154}	Medium	Direct	Consistent	Imprecise	Not detected	No difference or reduction in transfers with telehealth consultations	Low
Inpatient: Remote ICU	Mortality	1 (525 patients) ¹²² 1 (83 rural hospitals) ⁷²	Medium	Direct	Consistent	Imprecise	Not detected	No difference in mortality in for rural hospitals	Low
	Transfers	1 (525) ¹²²	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Inpatient: Neonates (specialty care)	Mortality	1 (384 hospitals) ⁷⁵	High	Indirect	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
	Length of Stay	2 (298) ^{92, 93}	Medium	Direct	Consistent	Imprecise	Not detected	Unclear effect	Insufficient
	Transfers	1 (317 patients) ⁵⁹ 1 (384 hospitals) ⁷⁵	Medium	Direct	Consistent	Imprecise	Not detected	More appropriate transfers	Low

Topic	Outcome	Number of Studies (Combined N)	Study Limitations (Low, Medium, High)	Directness (Direct, Indirect)	Consistency Consistent, Inconsistent, Unknown	Precision (Precise, Imprecise)	Reporting Bias (Not Detected, Possible, Suspected)	Conclusion	Strength of Evidence Grade (Insufficient, Low, Moderate, High)
	Clinical Outcomes	2 (298) ^{92, 93} 1 (384 hospitals) ⁷⁵	Medium	Direct	Consistent	Imprecise	Not detected	No difference in clinical measures between telehealth supported care and a higher level NICU	Low
Outpatient: Depression	Clinical outcomes	3 (974) ^{46, 49, 138}	Medium	Direct	Inconsistent	Imprecise	Not detected	Some improvement in medication adherence, treatment response, satisfaction, and quality of well-being	Low
Outpatient: Depression	Payer outcomes	2 (724) ^{46, 49}	Medium	Direct	Inconsistent	Imprecise	Not detected	Higher costs associated with intended higher utilization, acceptable incremental cost effectiveness	Low
Outpatient: Diabetes	Clinical outcomes	4 (303) ^{31, 81, 86, 98}	Medium	Direct	Inconsistent	Imprecise	Not detected	Improved A1c, self-monitoring of blood glucose, and Diabetes Self-Management Education score;	Low
Outpatient: Diabetes Outpatient: screening	Utilization outcomes	1 (1,024) ³⁴	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Pharmacy	Clinical outcomes	2 (879) ^{10, 29}	Medium	Direct	Inconsistent	Imprecise	Not detected	Improved guideline adherence; improved A1c, fasting blood glucose, and systolic blood pressure in patients with both diabetes and hypertension (no difference in patients with only one of the conditions)	Low
Outpatient: Rheumatology	Clinical outcomes	1 (85) ¹⁵⁵	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Rheumatology	Utilization outcomes	1 (85) ¹⁷⁶	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Oncology	Utilization outcomes	1 (110) ¹⁵¹	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient

Topic	Outcome	Number of Studies (Combined N)	Study Limitations (Low, Medium, High)	Directness (Direct, Indirect)	Consistency Consistent, Inconsistent, Unknown	Precision (Precise, Imprecise)	Reporting Bias (Not Detected, Possible, Suspected)	Conclusion	Strength of Evidence Grade (Insufficient, Low, Moderate, High)
Outpatient: Oncology	Payer outcomes	1 (147) ¹⁵⁶	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Remote Consults	Clinical outcomes	2 (189) ^{37, 79}	High	Direct	Inconsistent	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Remote Consults	Utilization outcomes	1 (113) ⁸²	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Echocardiography	Utilization outcomes	1 (38) ²²	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Hemodialysis	Clinical outcomes	1 (19) ¹⁴⁵	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Fracture	Payer outcomes	1 (12) ⁹⁷	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Dementia	Clinical outcomes	1 (188) ⁷⁶	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Hepatitis C	Clinical outcomes	1 (80) ¹³⁹	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: ADHD	Clinical outcomes	1 (223) ¹¹⁰	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: PTSD	Clinical outcomes	1 (225) ⁴⁷	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: PTSD	Utilization outcomes	1 (225) ⁴⁷	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: PTSD	Payer outcomes	1 (225) ¹²⁰	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Wound care	Clinical outcomes	1 (54) ¹⁵²	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Wound care	Utilization outcomes	1 (54) ¹⁵²	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Wound care	Utilization outcomes	1 (54) ¹⁵²	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Dermatology	Patient outcomes	1 (261) ¹⁷²	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Dermatology	Payer outcomes	1 (392) ³⁵	Medium	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: endoscopy	Payer outcomes	1(90) ¹⁶⁵	High	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient

Topic	Outcome	Number of Studies (Combined N)	Study Limitations (Low, Medium, High)	Directness (Direct, Indirect)	Consistency (Consistent, Inconsistent, Unknown)	Precision (Precise, Imprecise)	Reporting Bias (Not Detected, Possible, Suspected)	Conclusion	Strength of Evidence Grade (Insufficient, Low, Moderate, High)
Outpatient: Blood Pressure Control	Patient outcomes	1 (1299) ¹⁷⁹	Low	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Palliative Care	Patient outcomes	1(n=21) ⁷⁰	High	Direct	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
Outpatient: Ultrasound during pregnancy	Patient outcomes	1 (statewide, over 100,000) ⁸⁷	High	Indirect	Unknown	Imprecise	Not detected	Unclear effect	Insufficient
EMS/ED: Telestroke/ STEMI/ Chest Pain	Mortality	5 (2,312) ^{20, 38, 111, 141, 182}	Medium	Direct	Consistent	Precise	Not detected	No difference in mortality with telehealth supported care	Low
	Time to treatment	8 (3,725) ^{20, 25, 26, 38, 100, 106, 111, 124}	Medium	Direct	Inconsistent	Imprecise	Not detected	No difference or improved time to treatment. No evidence of worsened time to treatment.	Low
EMS/ED: Consultation	Transfers	5 (147,910) ^{58, 65, 104, 114, 150}	Medium	Direct	Inconsistent	Precise	Not detected	No difference or improved appropriate patient transfers. No evidence of increased inappropriate patient transfers.	Low
	ED Length of Stay	4 (9,094) ^{44, 104, 105, 164}	Medium	Direct	Inconsistent	Imprecise	Not detected	Unclear effect	Insufficient
Education/ Mentoring	Clinical response	2 ^{11, 167}	Moderate	Direct	Consistent	Imprecise	Not detected	Patient outcomes (A1c) improve or are equivalent across spoke and hub sites (hepatitis C viral response) after their providers receive remote education and mentoring	Low
	Provider behavior	8 ^{18, 19, 23, 51, 66, 96, 143, 175}	High	Direct	Consistent	Imprecise	Possible	Provider behaviors improve (prescribing, documentation & counseling, screening) after participation in remote education and mentoring	Low

Topic	Outcome	Number of Studies (Combined N)	Study Limitations (Low, Medium, High)	Directness (Direct, Indirect)	Consistency Consistent, Inconsistent, Unknown	Precision (Precise, Imprecise)	Reporting Bias (Not Detected, Possible, Suspected)	Conclusion	Strength of Evidence Grade (Insufficient, Low, Moderate, High)
	Provider knowledge, efficacy, perception	13 ^{19, 42, 53, 66, 84, 85, 96, 99, 107, 136, 142, 148, 158}	High	Direct	Consistent	Imprecise	Not detected	Provider knowledge, self-efficacy and perception improve after participation in remote education and mentoring (diabetes, liver disease, mental health, pediatric burn prevention)	Low

Abbreviations: AD/HD = attention deficit/ hyperactivity disorder; ED = emergency department; EMS = emergency medical services; A1c = glycated hemoglobin; ICU = intensive care unit; NICU = neonatal intensive care unit; PTSD = post-traumatic stress disorder; STEMI = ST-elevation acute myocardial infarction.

Appendix G. Excluded Studies List

1. Abadia de Barbara AH, Nicholas Iv TA, Del Real Colomo A, et al. Virtual simulation training using the Storz C-HUB to support distance airway training for the Spanish Medical Corps and NATO partners. *Stud Health Technol Inform.* 2012;182:1-9. PMID: 23138073. Exclusion: Ineligible population.
2. Abara E, Denton R. Collaborative care of urologic cancer patients through telemedicine in northeastern Ontario. *Urology.* 2013;82(3):S312. Exclusion: Ineligible study design.
3. Abd Allah M, Wahed S, Ammar I, et al. Utility of telemedicine in the treatment of patients with chronic HCV infection using DAAs in remote areas with limited resources. *Liver International.* 2021;41(8):1979-80. doi: 10.1111/liv.14982. PMID: 34153175. Exclusion: Ineligible study design.
4. Abshire DA, Graves JM, Amiri S, et al. Scaling-up telehealth during the COVID-19 pandemic: missed opportunities to leverage spatial data to ascertain the rural reach of services. *J Adolesc Health.* 2020;67(6):871-. doi: 10.1016/j.jadohealth.2020.07.017. PMID: 32839080. Exclusion: Not a study.
5. Adams SJ, Burbridge BE, Badea A, et al. Initial experience using a telerobotic ultrasound system for adult abdominal sonography. *Can Assoc Radiol J.* 2017;68(3):308-14. doi: 10.1016/j.carj.2016.08.002. PMID: 28159435. Exclusion: Ineligible population.
6. Adams SJ, Burbridge B, Chatterson L, et al. Telerobotic ultrasound to provide obstetrical ultrasound services remotely during the COVID-19 pandemic. *J Telemed Telecare.* 2020;1357633X20965422. doi: 10.1177/1357633X20965422. PMID: 33076753. Exclusion: Ineligible comparator.
7. Agley J, Adams ZW, Hulvershorn LA. Extension for Community Healthcare Outcomes (ECHO) as a tool for continuing medical education on opioid use disorder and comorbidities. *Addiction.* 2019;114(3):573-4. doi: 10.1111/add.14494. PMID: 30397977. Exclusion: Not a study.
8. Agley J, Delong J, Janota A, et al. Reflections on project ECHO: qualitative findings from five different ECHO programs. *Med Educ Online.* 2021;26(1):1936435. doi: 10.1080/10872981.2021.1936435. PMID: 34076567. Exclusion: Ineligible comparator.
9. Al Achkar M, Bennett IM, Chwastiak L, et al. Telepsychiatric consultation as a training and workforce development strategy for rural primary care. *Ann Fam Med.* 2020;18(5):438-45. doi: 10.1370/afm.2561. PMID: 32928760. Exclusion: Ineligible study design.
10. Al Kasab S, Harvey JB, Debenham E, et al. Door to needle time over telestroke-a comprehensive stroke center experience. *Telemed J E Health.* 2018;24(2):111-5. doi: 10.1089/tmj.2017.0067. PMID: 28753069. Exclusion: Ineligible comparator.
11. Al Quran HA, Khader YS, Ellauzi ZM, et al. Effect of real-time teledermatology on diagnosis, treatment and clinical improvement. *J Telemed Telecare.* 2015;21(2):93-9. doi: 10.1177/1357633X14566572. PMID: 25589469. Exclusion: Ineligible intervention.
12. AlAzab R, Khader Y. Telenephrology application in rural and remote areas of Jordan: benefits and impact on quality of life. *Rural Remote Health.* 2016;16(1):3646. PMID: 26745230. Exclusion: Ineligible intervention.

13. AlDossary S, Martin-Khan MG, Bradford NK, et al. The development of a telemedicine planning framework based on needs assessment. *J Med Syst.* 2017;41(5):74. doi: 10.1007/s10916-017-0709-4. PMID: 28321589. Exclusion: Not a study.
14. Aligeti M. Telemedicine wellness, intervention, triage, and referral project: a unique intervention strategy to prevent school violence. *J Am Acad Child Adolesc Psychiatry.* 2016;55(10):S12. doi: 10.1016/j.jaac.2016.07.480. Exclusion: Ineligible intervention.
15. Alschuler KN, von Geldern G, Ball D, et al. Rapid transfer of knowledge for multiple sclerosis clinical care during COVID-19: ECHO MS. *Mult Scler Relat Disord.* 2020;46:102600. doi: 10.1016/j.msard.2020.102600. PMID: 33296992. Exclusion: Ineligible outcome.
16. Amorim E, Shih MM, Koehler SA, et al. Impact of telemedicine implementation in thrombolytic use for acute ischemic stroke: the University of Pittsburgh Medical Center telestroke network experience. *J Stroke Cerebrovasc Dis.* 2013;22(4):527-31. doi: 10.1016/j.jstrokecerebrovasdis.2013.02.004. PMID: 23489955. Exclusion: Ineligible population.
17. Anders SH, Woods DD, Schweikhart S, et al. The effects of health information technology change over time: a study of tele-ICU functions. *Appl Clin Inform.* 2012;3(2):239-47. doi: 10.4338/ACI-2011-12-RA-0073. PMID: 23646073. Exclusion: Ineligible outcome.
18. Anderson D, Zlateva I, Davis B, et al. Improving pain care with Project ECHO in community health centers. *Pain Med.* 2017;18(10):1882-9. doi: 10.1093/pm/pnx187. PMID: 29044409. Exclusion: Ineligible population.
19. Anderson D, Villagra V, Coman EN, et al. A cost-effectiveness analysis of cardiology eConsults for Medicaid patients. *Am J Manag Care.* 2018;24(1):e9-e16. PMID: 29350511. Exclusion: Ineligible population.
20. Anderson ER, Smith B, Ido M, et al. Remote assessment of stroke using the iPhone 4. *J Stroke Cerebrovasc Dis.* 2013;22(4):340-4. doi: 10.1016/j.jstrokecerebrovasdis.2011.09.013. PMID: 22018507. Exclusion: Ineligible outcome.
21. Anderson J, Singh J. A case study of using telehealth in a rural healthcare facility to expand services and protect the health and safety of patients and staff. *Healthcare (Basel).* 2021;9(6):15. doi: 10.3390/healthcare9060736. PMID: 34203888. Exclusion: Ineligible intervention.
22. Anderson JB, Martin SA, Gadomski A, et al. Project ECHO and primary care buprenorphine treatment for opioid use disorder: implementation and clinical outcomes. *Subst Abus.* 2021:1-9. doi: 10.1080/08897077.2021.1931633. PMID: 34086529. Exclusion: Ineligible intervention.
23. Anderson LS. The development and implementation of eSchoolCare a novel health care support system for school nurses. *ANS Adv Nurs Sci.* 2013;36(4):289-303. doi: 10.1097/ANS.0000000000000003. PMID: 24169109. Exclusion: Not a study.
24. Anderson SL, Bianco J, DeRemer CE. Adapting ambulatory care learning environments in response to the COVID-19 pandemic. *Am J Health-Syst Pharm.* 2021;78(6):467-71. doi: 10.1093/ajhp/zxab016. PMID: 33483738. Exclusion: Not a study.
25. Anderson T, McClintock AS, McCarrick SS, et al. Working alliance, interpersonal problems, and depressive symptoms in tele-interpersonal psychotherapy for HIV-infected rural persons: evidence for indirect effects. *J Clin Psychol.* 2018;74(3):286-303. doi: 10.1002/jclp.22502. PMID: 28586534. Exclusion: Ineligible intervention.

26. Andrade MV, Maia AC, Cardoso CS, et al. Cost-benefit of the telecardiology service in the state of Minas Gerais: Minas Telecardio Project. *Arq Bras Cardiol.* 2011;97(4):307-16. PMID: 21808852. Exclusion: Ineligible comparater.
27. Andrees V, Klein TM, Augustin M, et al. Live interactive teledermatology compared to in-person care - a systematic review. *J Eur Acad Dermatol Venereol.* 2020;34(4):733-45. doi: 10.1111/jdv.16070. PMID: 31715035. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
28. Anonymous. Telemedicine helps rural EDs access critical neurology expertise for stroke patients. *ED Manag.* 2012;24(3):30-3. PMID: 23687736. Exclusion: Not a study.
29. Anonymous. Technologies assisting in remote consultations for the diagnosis of stroke: a review of the clinical evidence. 2013. <https://www.ncbi.nlm.nih.gov/books/NBK195829/> PMID: 24741717. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
30. Anonymous. Connecting rural and remote services. *Qld Nurse.* 2016;35(1):10. PMID: 27356321. Exclusion: Not a study.
31. Anonymous. Project ECHO: a telementoring network model for continuing professional development: erratum. *J Contin Educ Health Prof.* 2018;38(1):78. doi: 10.1097/CEH.000000000000192. PMID: 29517614. Exclusion: Not a study.
32. Antezana L, Scarpa A, Valdespino A, et al. Rural trends in diagnosis and services for autism spectrum disorder. *Front Psychol.* 2017;8:590. doi: 10.3389/fpsyg.2017.00590. PMID: 28473784. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
33. Arba F, Piccardi B, Baldereschi M, et al. Telemedicine for acute ischaemic stroke. *Cochrane Database Syst Rev.* 2016 (2). Exclusion: Not a study.
34. Arbeille P, Zuj K, Saccomandi A, et al. Remote echography & doppler using tele-operated compact motorised probes & portable echograph. Application to 200 isolated patient in rural areas. *Angeiologie.* 2016;68(3):23-34. Exclusion: Ineligible outcome.
35. Archbald-Pannone LR, Harris DA, Alberio K, et al. COVID-19 collaborative model for an academic hospital and long-term care facilities. *J Am Med Dir Assoc.* 2020;21:939-42. doi: 10.1016/j.jamda.2020.05.044. PMID: 32563752. Exclusion: Ineligible population.
36. Arora S, Kalishman S, Thornton K, et al. Expanding access to hepatitis C virus treatment--Extension for Community Healthcare Outcomes (ECHO) Project: disruptive innovation in specialty care. *Hepatology.* 2010;52(3):1124-33. doi: 10.1002/hep.23802. PMID: 20607688. Exclusion: Ineligible population.
37. Arora S, Kalishman S, Dion D, et al. Partnering urban academic medical centers and rural primary care clinicians to provide complex chronic disease care. *Health Aff (Millwood).* 2011;30(6):1176-84. doi: 10.1377/hlthaff.2011.0278. PMID: 21596757. Exclusion: Not a study.
38. Arora S, Thornton K, Murata G, et al. Project ECHO expands access to hepatitis C (HCV) treatment in rural areas and prisons. Treatment is as effective as a university HCV clinic. *J Hepatol.* 2011;54:S161. doi: 10.1016/S0168-8278(11)60404-2. Exclusion: Not a study.
39. Arora S, Kalishman S, Thornton K, et al. Project ECHO (Project Extension for Community Healthcare Outcomes): a national and global model for continuing professional development. *J Contin Educ Health.* 2016;36 Suppl 1:S48-9. doi: 10.1097/CEH.0000000000000097. PMID: 27584072. Exclusion: Not a study.

40. Arora S, Kalishman SG, Thornton KA, et al. Project ECHO: a telementoring network model for continuing professional development. *J Contin Educ Health Prof.* 2017;37(4):239-44. doi: 10.1097/CEH.000000000000172. PMID: 29189491. Exclusion: Not a study.
41. Arora S. Project ECHO: democratising knowledge for the elimination of viral hepatitis. *Lancet Gastroenterol Hepatol.* 2019;4(2):91-3. doi: 10.1016/S2468-1253(18)30390-X. PMID: 30647014. Exclusion: Not a study.
42. Arora S, Byers EL. Leveraging local expertise to improve rural cancer care outcomes using Project ECHO: a response to Levit et al. *JCO Oncol Pract.* 2020;16(7):399-403. doi: 10.1200/OP.20.00260. PMID: 32574127. Exclusion: Not a study.
43. Arora S, Mate KS, Jones JL, et al. Enhancing collaborative learning for quality improvement: evidence from the improving clinical flow project, a breakthrough series collaborative with Project ECHO. *Jt Comm J Qual Patient Saf.* 2020;46(8):448-56. doi: 10.1016/j.jcjq.2020.04.013. PMID: 32507466. Exclusion: Ineligible population.
44. Arora S, Thornton K. Novel models of hepatitis C virus care delivery: telemedicine, Project ECHO, and integrative care. *Clin Liver Dis (Hoboken).* 2020;16(1):5-7. doi: 10.1002/cld.912. PMID: 32714515. Exclusion: Not a study.
45. Asgary R, Staderini N, Mthethwa-Hleta S, et al. Evaluating smartphone strategies for reliability, reproducibility, and quality of VIA for cervical cancer screening in the Shiselweni region of Eswatini: a cohort study. *PLoS Med.* 2020;17(11):e1003378. doi: 10.1371/journal.pmed.1003378. PMID: 33211691. Exclusion: Ineligible outcome.
46. Atiyeh B, Dibo SA, Janom HH. Telemedicine and burns: an overview. *Ann Burns Fire Disasters.* 2014;27(2):87-93. PMID: 26170782. Exclusion: Not a study.
47. Audebert HJ, Schultes K, Tietz V, et al. Long-term effects of specialized stroke care with telemedicine support in community hospitals on behalf of the Telemedical Project for Integrative Stroke Care (TEMPiS). *Stroke.* 2009;40(3):902-8. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
48. Austad K, Juarez M, Shryer H, et al. Obstetric care navigation: results of a quality improvement project to provide accompaniment to women for facility-based maternity care in rural Guatemala. *BMJ Qual Saf.* 2020;29(2):169-78. doi: 10.1136/bmjqs-2019-009524. PMID: 31678958. Exclusion: Ineligible intervention.
49. Avdalovic MV, Marcin JP. When will telemedicine appear in the ICU? *J Intensive Care Med.* 2019;34(4):271-6. doi: 10.1177/0885066618775956. PMID: 29756527. Exclusion: Not a study.
50. Backhaus R, Schlachetzki F, Rackl W, et al. Intracranial hemorrhage: frequency, location, and risk factors identified in a TeleStroke network. *Neuroreport.* 2015;26(2):81-7. doi: 10.1097/WNR.0000000000000304. PMID: 25536117. Exclusion: Ineligible study design.
51. Baidwan NK, Davlyatov G, Mehta T. Telehealth use among community health centers and cardio-metabolic health outcomes. *Healthcare (Basel).* 2020;8(165):1-10. doi: 10.3390/healthcare8020165. PMID: 32532120. Exclusion: Ineligible intervention.
52. Baker LC, Johnson SJ, Macaulay D, et al. Integrated telehealth and care management program for Medicare beneficiaries with chronic disease linked to savings. *Health Aff (Millwood).* 2011;30(9):1689-97. doi: 10.1377/hlthaff.2011.0216. PMID: 21900660. Exclusion: Ineligible intervention.

53. Bakitas M, Lyons KD, Hegel MT, et al. The project ENABLE II randomized controlled trial to improve palliative care for rural patients with advanced cancer: baseline findings, methodological challenges, and solutions. *Palliat Support Care*. 2009;7(1):75-86. doi: 10.1017/s1478951509000108. PMID: 19619377. Exclusion: Ineligible population.
54. Bakitas MA, Elk R, Astin M, et al. Systematic review of palliative care in the rural setting. *Cancer Control*. 2015;22(4):450-64. PMID: 26678972. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
55. Ball S, Wilson B, Ober S, et al. SCAN-ECHO for pain management: implementing a regional telementoring training for primary care providers. *Pain Med*. 2018;19(2):262-8. doi: 10.1093/pm/pnx122. PMID: 28525633. Exclusion: Ineligible population.
56. Banbury A, Roots A, Nancarrow S. Rapid review of applications of e-health and remote monitoring for rural residents. *Aust J Rural Health*. 2014;22(5):211-22. doi: 10.1111/ajr.12127. PMID: 25303412. Exclusion: Ineligible study design.
57. Bansal M, Singh S, Maheshwari P, et al. Value of interactive scanning for improving the outcome of new-learners in transcontinental tele-echocardiography (VISION-in-Tele-Echo) study. *J Am Soc Echocardiogr*. 2015;28(1):75-87. doi: 10.1016/j.echo.2014.09.001. PMID: 25306222. Exclusion: Ineligible population.
58. Barber M, Maclean A, Frieslick J, et al. Western isles stroke telerehabilitation (specialist medical consultation) service. *Age Ageing*. 2013;42:iii7. doi: 10.1093/ageing/aft096. Exclusion: Not a study.
59. Barlinn J, Gerber J, Barlinn K, et al. Acute endovascular treatment delivery to ischemic stroke patients transferred within a telestroke network: a retrospective observational study. *Int J Stroke*. 2017;12(5):502-9. doi: 10.1177/1747493016681018. PMID: 27899742. Exclusion: Ineligible comparater.
60. Barton AJ, Anderson JL. Meeting the challenge of perinatal care in rural communities. *J Perinat Neonatal Nurs*. 2021;35(2):150-9. doi: 10.1097/JPN.0000000000000562. PMID: 33900245. Exclusion: Not a study.
61. Barton C, Morris R, Rothlind J, et al. Video-telemedicine in a memory disorders clinic: evaluation and management of rural elders with cognitive impairment. *Telemed J E Health*. 2011;17(10):789-93. doi: 10.1089/tmj.2011.0083. PMID: 22023458. Exclusion: Ineligible study design.
62. Basel D. Could telemedicine help with the CMS rule change requiring in-person provider evaluations before transfer from nursing home to hospital? *J Am Med Dir Assoc*. 2016;17(2):179-. doi: 10.1016/j.jamda.2015.11.006. PMID: 26706419. Exclusion: Not a study.
63. Bath B, Lovo Grona S, Milosavljevic S, et al. Advancing interprofessional primary health care services in rural settings for people with chronic low back disorders: protocol of a community-based randomized controlled trial. *JMIR Res Protoc*. 2016;5(4):e212. PMID: 27829573. Exclusion: Not a study.
64. Batsis JA, DiMilia PR, Seo LM, et al. Effectiveness of ambulatory telemedicine care in older adults: a systematic review. *J Am Geriatr Soc*. 2019;67(8):1737-49. doi: 10.1111/jgs.15959. PMID: 31066916. Exclusion: Ineligible intervention.

65. Batsis JA, McClure AC, Weintraub AB, et al. Barriers and facilitators in implementing a pilot, pragmatic, telemedicine-delivered healthy lifestyle program for obesity management in a rural, academic obesity clinic. *Implement Sci Commun.* 2020;1:83. doi: 10.1186/s43058-020-00075-9. PMID: 33015640. Exclusion: Ineligible intervention.
66. Bauer AM, Jakupcak M, Hawrilenko M, et al. Outcomes of a health informatics technology-supported behavioral activation training for care managers in a collaborative care program. *Fam Syst Health.* 2021;39(1):89-100. doi: 10.1037/fsh0000523. PMID: 32853001. Exclusion: Ineligible intervention.
67. Becevic M, Mutrux R, Edison K. Show-Me ECHO: complex disease care capacity-building telehealth program. *Stud Health Technol Inform.* 2016;226:233-6. PMID: 27350513. Exclusion: Ineligible outcome.
68. Beck JR, Mutalik P. Sharing cancer care information across VA health care systems. *Fed Pract.* 2019;36(Suppl 3):S29-S32. PMID: 31138981. Exclusion: Ineligible population.
69. Bedard BA, Younge M, Pettit PA, et al. Using telemedicine for tuberculosis care management: a three county inter-municipal approach. *J Med Syst.* 2017;42(1):12. doi: 10.1007/s10916-017-0872-7. PMID: 29181590. Exclusion: Ineligible study design.
70. Bennett KA, Ong T, Verrall AM, et al. Project ECHO-geriatrics: training future primary care providers to meet the needs of older adults. *J Grad Med Educ.* 2018;10(3):311-5. doi: 10.4300/JGME-D-17-01022.1. PMID: 29946389. Exclusion: Ineligible population.
71. Benson GA, Sidebottom A, Hayes J, et al. Impact of ENHANCED (diEtitiaNs Helping pAtieNs CarE for Diabetes) telemedicine randomized controlled trial on diabetes optimal care outcomes in patients with type 2 diabetes. *J Acad Nutr Diet.* 2019;119(4):585-98. doi: 10.1016/j.jand.2018.11.013. PMID: 30711463. Exclusion: Ineligible intervention.
72. Bergrath S, Reich A, Rossaint R, et al. Feasibility of prehospital teleconsultation in acute stroke--a pilot study in clinical routine. *PLoS One.* 2012;7(5):e36796. doi: 10.1371/journal.pone.0036796. PMID: 22629331. Exclusion: Ineligible population.
73. Berndt A, Murray CM, Kennedy K, et al. Effectiveness of distance learning strategies for continuing professional development (CPD) for rural allied health practitioners: a systematic review. *BMC Med Educ.* 2017;17(117):1-13. doi: 10.1186/s12909-017-0949-5. PMID: 28701199. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
74. Berndt J, Leone P, King G. Using teledentistry to provide interceptive orthodontic services to disadvantaged children. *Am J Orthod Dentofacial Orthop.* 2008;134(5):700-6. doi: 10.1016/j.ajodo.2007.12.023. PMID: 18984404. Exclusion: Ineligible intervention.
75. Bernocchi P, Scalvini S, Tridico C, et al. Healthcare continuity from hospital to territory in Lombardy: TELEMACO project. *Am J Manag Care.* 2012;18(3):e101-8. PMID: 22435961. Exclusion: Ineligible population.
76. Beste LA, Glorioso TJ, Ho PM, et al. Telemedicine specialty support promotes hepatitis C treatment by primary care providers in the Department of Veterans Affairs. *Am J Med.* 2017;130(4):432-8.e3. doi: 10.1016/j.amjmed.2016.11.019. PMID: 27998682. Exclusion: Ineligible population.

77. Bettencourt AF, Ferro RA, Williams JL, et al. Pediatric primary care provider comfort with mental health practices: a needs assessment of regions with shortages of treatment access. *Acad Psychiatry*. 2021;45(4):429-34. doi: 10.1007/s40596-021-01434-x. PMID: 33786779. Exclusion: Ineligible outcome.
78. Bhanot N, Dimitriou G, McAninch L, et al. Perspectives of health care providers in an integrated health care delivery network on inpatient electronic consultation (e-consult) use during the COVID-19 pandemic. *J Patient Exp*. 2021;8:23743735211007696. doi: 10.1177/23743735211007696. PMID: 34179421. Exclusion: Ineligible setting.
79. Bian J, Cristaldi KK, Summer AP, et al. Association of a school-based, asthma-focused telehealth program with emergency department visits among children enrolled in South Carolina Medicaid. *JAMA Pediatr*. 2019;173(11):1041-8. doi: 10.1001/jamapediatrics.2019.3073. PMID: 31498379. Exclusion: Ineligible intervention.
80. Bice-Urbach BJ, Kratochwill TR. Teleconsultation: the use of technology to improve evidence-based practices in rural communities. *J Sch Psychol*. 2016;56:27-43. doi: 10.1016/j.jsp.2016.02.001. PMID: 27268568. Exclusion: Ineligible population.
81. Bidargaddi N, Schrader G, Roeger L, et al. Early effects of upgrading to a high bandwidth digital network for telepsychiatry assessments in rural South Australia. *J Telemed Telecare*. 2015;21(3):174-5. doi: 10.1177/1357633X15569955. PMID: 25680391. Exclusion: Not a study.
82. Biemba G, Chiluba B, Yeboah-Antwi K, et al. Impact of mobile health-enhanced supportive supervision and supply chain management on appropriate integrated community case management of malaria, diarrhoea, and pneumonia in children 2-59 months: a cluster randomised trial in Eastern Province, Zambia. *J Glob Health*. 2020;10(1):010425. doi: 10.7189/jogh.10.010425. PMID: 32509293. Exclusion: Ineligible setting.
83. Biglan KM, Grana MJ, Lauver SC, et al. Implementation of a telemedicine program for Parkinson's disease: process and recommendations. *Mov Disord*. 2012;27:S120. doi: 10.1002/mds.25051. Exclusion: Ineligible intervention.
84. Bikinesi L, O'Bryan G, Roscoe C, et al. Implementation and evaluation of a Project ECHO telementoring program for the Namibian HIV workforce. *Hum Resour Health*. 2020;18(1):61. doi: 10.1186/s12960-020-00503-w. PMID: 32873303. Exclusion: Ineligible setting.
85. Blacquiere D, Lindsay MP, Foley N, et al. Canadian stroke best practice recommendations: telestroke best practice guidelines update 2017. *Int J Stroke*. 2017;12(8):886-95. doi: 10.1177/1747493017706239. PMID: 28441928. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
86. Blech B, O'Carroll CB, Zhang N, et al. Telestroke program participation and improvement in door-to-needle times. *Telemed J E Health*. 2020;26(4):406-10. doi: 10.1089/tmj.2018.0336. PMID: 31287782. Exclusion: Ineligible population.
87. Bolitho L. The changing role of rural consultant physicians and paediatricians. *Intern Med J*. 2010;40:5. doi: 10.1111/j.1445-5994.2010.02186.x. Exclusion: Not a study.

88. Bonnell S, Griggs A, Avila G, et al. Community health workers and use of mHealth: improving identification of pregnancy complications and access to care in the Dominican Republic. *Health Promot Pract*. 2018;19(3):331-40. doi: 10.1177/1524839917708795. PMID: 28578606. Exclusion: Ineligible study design.
89. Borrelli L. How neurologists address patients' needs in rural areas when the practice is three hours away. *Neurology Today*. 2019;19(16):37-9. doi: 10.1097/01.nt.0000579952.52164.8e. Exclusion: Not a study.
90. Bosseila M, Dahroug H. Tele dermatology model in remote and rural communities in Egypt. *J Dermatol Nurses Assoc*. 2020;12(2). Exclusion: Ineligible setting.
91. Bouchonville MF, Paul MM, Billings J, et al. Taking telemedicine to the next level in diabetes population management: a review of the endo ECHO model. *Curr Diab Rep*. 2016;16(10):96. doi: 10.1007/s11892-016-0784-9. PMID: 27549110. Exclusion: Not a study.
92. Bowman DL. Case studies demonstrating stroke telemedicine in the rural emergency department. *J Emerg Nurs*. 2017;43(1):70-1. doi: 10.1016/j.jen.2016.10.008. PMID: 28131354. Exclusion: Not a study.
93. Boxer RJ. Telemedicine: remote cancer care improves communication. *Oncology Times*. 2017;39(2):1-15. doi: 10.1097/01.cot.0000512175.80464.3f. Exclusion: Not a study.
94. Boyle J. Project ECHO, the prison peer education project. *Pain Res Manag*. 2014;19(3):e51. Exclusion: Not a study.
95. Boyle TP, Liu J, Dyer KS, et al. Pilot paramedic survey of benefits, risks, and strategies for pediatric prehospital telemedicine. *Pediatr Emerg Care*. 2020;06:06. doi: 10.1097/PEC.0000000000002099. PMID: 33170566. Exclusion: Ineligible study design.
96. Bradford N, Young J, Armfield NR, et al. A pilot study of the effectiveness of home teleconsultations in paediatric palliative care. *J Telemed Telecare*. 2012;18(8):438-42. doi: 10.1258/jtt.2012.GTH103. PMID: 23155114. Exclusion: Ineligible intervention.
97. Bradford NK, Caffery LJ, Smith AC. Telehealth services in rural and remote Australia: a systematic review of models of care and factors influencing success and sustainability. *Rural Remote Health*. 2016;16(4):4268. PMID: 27817199. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
98. Brandon AR, Song L, Deal AM, et al. Using telehealth to train providers of a cancer support intervention. *Telemed J E Health*. 2015;21(10):793-800. doi: 10.1089/tmj.2014.0208. PMID: 26431258. Exclusion: Ineligible outcome.
99. Brennan JA, Kealy JA, Gerardi LH, et al. Telemedicine in the emergency department: a randomized controlled trial. *J Telemed Telecare*. 1999;5(1):18-22. PMID: 10505365. Exclusion: Ineligible intervention.
100. Brimmer DJ, Jones JF, Boneva R, et al. Assessment of ME/CFS (myalgic encephalomyelitis/chronic fatigue syndrome): a case study for health care providers. *MedEdPORTAL*. 2016;12:10527. doi: 10.15766/mep_2374-8265.10527. PMID: 30984868. Exclusion: Ineligible population.
101. Briscoe JG, Baker S, Menis AV, et al. Comparison of emergency department throughput with standard in-person triage vs virtual telehealth triage. *Acad Emerg Med*. 2020;27:S91. doi: 10.1111/acem.13961. Exclusion: Not a study.

102. Brisson AM, Steinmetz P, Oleskevich S, et al. A comparison of telemedicine teaching to in-person teaching for the acquisition of an ultrasound skill - a pilot project. *J Telemed Telecare*. 2015;21(4):235-9. doi: 10.1177/1357633X15575446. PMID: 25766853. Exclusion: Ineligible population.
103. Brokmann JC, Conrad C, Rossaint R, et al. Treatment of Acute Coronary Syndrome by Telemedically Supported Paramedics Compared With Physician-Based Treatment: A Prospective, Interventional, Multicenter Trial. *J Med Internet Res*. 2016;18(12):e314. doi: 10.2196/jmir.6358. PMID: 27908843. Exclusion: Ineligible population.
104. Brokmann JC, Rossaint R, Hirsch F, et al. Analgesia by telemedically supported paramedics compared with physician-administered analgesia: A prospective, interventional, multicentre trial. *Eur J Pain*. 2016;20(7):1176-84. doi: 10.1002/ejp.843. PMID: 26914284. Exclusion: Ineligible population.
105. Browne E, Nawaz S, Mathelier G, et al. Can acute stroke interventions improve health outcomes in resource-limited rural areas? *J Natl Med Assoc*. 2020;112(5):S34. doi: 10.1016/j.jnma.2020.09.082. Exclusion: Ineligible setting.
106. Browne T, McKinney SH, Duck L, et al. An academic-community interprofessional telehealth online training partnership: impact on students and providers. *J Interprof Care*. 2021:1-8. doi: 10.1080/13561820.2021.1967896. PMID: 34514934. Exclusion: Ineligible population.
107. Browne T, McKinney SH, Duck L, et al. Preparing health professions students to serve southern rural communities in the time of COVID-19 and beyond: a model for interprofessional online telehealth education. *South Med J*. 2021;114(10):665-7. doi: 10.14423/SMJ.0000000000001300. PMID: 34599348. Exclusion: Not a study.
108. Burke BL, Jr., Hall RW, Section On Telehealth C. Telemedicine: pediatric applications. *Pediatrics*. 2015;136(1):e293-308. doi: 10.1542/peds.2015-1517. PMID: 26122813. Exclusion: Not a study.
109. Burns CL, Ward EC, Hill AJ, et al. Randomized controlled trial of a multisite speech pathology telepractice service providing swallowing and communication intervention to patients with head and neck cancer: Evaluation of service outcomes. *Head Neck*. 2017;39(5):932-9. doi: 10.1002/hed.24706. PMID: 28225567. Exclusion: Ineligible intervention.
110. Bursey M, Lyon M. Telemedicine critical care in rural hospitals during the COVID-19 pandemic. *Am J Respir Crit Care Med*. 2021;203(9)doi: 10.1164/ajrccm-conference.2021.203.1_MeetingAbstracts.A2585. Exclusion: Ineligible intervention.
111. Bursey SJ, Beal J, Edwards EK, et al. Delivering CME accredited provincial stroke rounds in Ontario: how OREG has made it work from 2008 - present. *Stroke*. 2013;44(12):e201. Exclusion: Ineligible outcome.
112. Burton DC, Stanley D, Ireson CL. Child advocacy outreach: using telehealth to expand child sexual abuse services in rural Kentucky. *J Telemed Telecare*. 2002;8(Suppl 2):10. PMID: 12217116. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
113. Bussieres S, Tanguay A, Hebert D, et al. Unite de Coordination Clinique des Services Prehospitaliers d'Urgence: a clinical telemedicine platform that improves prehospital and community health care for rural citizens. *J Telemed Telecare*. 2017;23(1):188-94. doi: 10.1177/1357633X15627234. PMID: 27072126. Exclusion: Ineligible population.
114. Buvik A, Bergmo TS, Bugge E, et al. Cost-effectiveness of telemedicine in remote orthopedic consultations: randomized controlled trial. *J Med Internet Res*. 2019;21(2):e11330. doi: 10.2196/11330. PMID: 30777845. Exclusion: Ineligible intervention.

115. Byamba K, Syed-Abdul S, Garcia-Romero M, et al. Mobile teledermatology for a prompter and more efficient dermatological care in rural Mongolia. *Br J Dermatol*. 2015;173(1):265-7. doi: 10.1111/bjd.13607. PMID: 25494968. Exclusion: Not a study.
116. Cadilhac DA, Moloczij N, Denisenko S, et al. Establishment of an effective acute stroke telemedicine program for Australia: protocol for the Victorian Stroke Telemedicine project. *Int J Stroke*. 2014;9(2):252-8. doi: 10.1111/ijss.12137. PMID: 24148281. Exclusion: Not a study.
117. Cai Y, Gong W, He H, et al. Mobile texting and lay health supporters to improve schizophrenia care in a resource-poor community in rural China (LEAN Trial): randomized controlled trial extended implementation. *J Med Internet Res*. 2020;22(12):e22631. doi: 10.2196/22631. PMID: 33258788. Exclusion: Ineligible intervention.
118. Cameron MP, Ray R, Sabesan S. Physicians' perceptions of clinical supervision and educational support via videoconference: a systematic review. *J Telemed Telecare*. 2014;20(5):272-81. PMID: 24906649. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
119. Campbell J, Theodoros D, Hartley N, et al. Implementation factors are neglected in research investigating telehealth delivery of allied health services to rural children: a scoping review. *J Telemed Telecare*. 2019:590-606. doi: 10.1177/1357633X19856472. PMID: 31216211. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
120. Campbell N, Schiffer E, Buxbaum A, et al. Taking knowledge for health the extra mile: participatory evaluation of a mobile phone intervention for community health workers in Malawi. *Glob Health Sci Pract*. 2014;2(1):23-34. doi: 10.9745/GHSP-D-13-00141. PMID: 25276560. Exclusion: Ineligible setting.
121. Canfield C, Galvin S. Bedside nurse acceptance of intensive care unit telemedicine presence. *Crit Care Nurse*. 2018;38(6):e1-e4. doi: 10.4037/ccn2018926. PMID: 30504503. Exclusion: Not a study.
122. Capampangan DJ, Wellik KE, Bobrow BJ, et al. Telemedicine versus telephone for remote emergency stroke consultations: a critically appraised topic. *Neurologist*. 2009;15(3):163-6. PMID: 19430275. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
123. Carallo C, Scavelli FB, Cipolla M, et al. Management of Type 2 Diabetes Mellitus through Telemedicine. *PLoS One*. 2015;10(5):e0126858. doi: 10.1371/journal.pone.0126858. PMID: 25974092. Exclusion: Ineligible population.
124. Casas I. Community dermatology in Argentina. *Dermatol Clin*. 2021;39(1):43-55. doi: 10.1016/j.det.2020.08.005. PMID: 33228861. Exclusion: Ineligible intervention.
125. Casey MM, Sorensen TD, Elias W, et al. Current practices and state regulations regarding telepharmacy in rural hospitals. *Am J Health-Syst Pharm*. 2010;67(13):1085-92. doi: 10.2146/ajhp090531. PMID: 20554595. Exclusion: Ineligible outcome.
126. Cashman H, Mayson E, Kliman D, et al. A shared electronic health record in a rural setting maintains hematological cancer services remotely during the COVID-19 pandemic. *Blood*. 2020;136:5. doi: 10.1182/blood-2020-142209. Exclusion: Ineligible intervention.
127. Cashman H, Sushil S, Milliken S, et al. The impact of covid-19 on a rural hematology outreach service, an Australian experience. *HemaSphere*. 2021;5(SUPPL 2):570. doi: 10.1097/HS9.0000000000000566. Exclusion: Ineligible comparator.

128. Chan AW, Kornder J, Elliott H, et al. Improved survival associated with pre-hospital triage strategy in a large regional ST-segment elevation myocardial infarction program. *JACC Cardiovasc Interv.* 2012;5(12):1239-46. doi: 10.1016/j.jcin.2012.07.013. PMID: 23257372. Exclusion: Ineligible population.
129. Chan BA, Larkins SL, Evans R, et al. Do teleoncology models of care enable safe delivery of chemotherapy in rural towns? *Med J Aust.* 2015;203(10):406-6.e6. PMID: 26561905. Exclusion: Ineligible intervention.
130. Chang HL, Shaw MJ, Lai F, et al. U-Health: an example of a high-quality individualized healthcare service. *Per Med.* 2010;7(6):677-87. doi: 10.2217/pme.10.64. PMID: 29788567. Exclusion: Ineligible study design.
131. Chang LW, Kagaayi J, Arem H, et al. Impact of a mHealth intervention for peer health workers on AIDS care in rural Uganda: a mixed methods evaluation of a cluster-randomized trial. *AIDS Behav.* 2011;15(8):1776-84. doi: 10.1007/s10461-011-9995-x. PMID: 21739286. Exclusion: Ineligible setting.
132. Chao GF, Li KY, Zhu Z, et al. Use of telehealth by surgical specialties during the COVID-19 pandemic. *Obstetrical and Gynecological Survey.* 2021;76(7):410-2. doi: 10.1097/01.ogx.0000767224.19385.74. Exclusion: Ineligible intervention.
133. Charlton M, Schlichting J, Chioreso C, et al. Challenges of rural cancer care in the United States. *Oncology (Williston).* 2015;29(9):633-40. PMID: 26384798. Exclusion: Not a study.
134. Chauhan V, Negi PC, Raina S, et al. Smartphone-based tele-electrocardiography support for primary care physicians reduces the pain-to-treatment time in acute coronary syndrome. *J Telemed Telecare.* 2018;24(8):540-6. doi: 10.1177/1357633X17719395. PMID: 28750576. Exclusion: Ineligible setting.
135. Cheak-Zamora N, Farmer JG, Crossman MK, et al. Provider perspectives on the extension for community healthcare outcomes autism: transition to adulthood program. *J Dev Behav Pediatr.* 2021;42(2):91-100. doi: 10.1097/DBP.0000000000000872. PMID: 33044397. Exclusion: Ineligible population.
136. Chen KC, Yen DH, Chen CD, et al. Effect of emergency department in-hospital tele-electrocardiographic triage and interventional cardiologist activation of the infarct team on door-to-balloon times in ST-segment-elevation acute myocardial infarction. *Am J Cardiol.* 2011;107(10):1430-5. doi: 10.1016/j.amjcard.2011.01.015. PMID: 21414598. Exclusion: Ineligible population.
137. Cheng W, Nazareth S, Flexman JP. Statewide hepatitis C model of care for rural and remote regions. *J Gastroenterol Hepatol.* 2015;30 Suppl 2:1-5. doi: 10.1111/jgh.12863. PMID: 25641223. Exclusion: Not a study.
138. Chern CC, Chen YJ, Hsiao B. Decision tree-based classifier in providing telehealth service. *BMC Med Inf Decis Mak.* 2019;19(1):104. doi: 10.1186/s12911-019-0825-9. PMID: 31146749. Exclusion: Ineligible intervention.
139. Chipps J, Brysiewicz P, Mars M. A systematic review of the effectiveness of videoconference-based tele-education for medical and nursing education. *Worldviews Evid Based Nurs.* 2012;9(2):78-87. doi: 10.1111/j.1741-6787.2012.00241.x. PMID: 22409341. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
140. Cho J, Chung HS, Choa M, et al. A pilot study of the Tele-Airway Management System in a hospital emergency department. *J Telemed Telecare.* 2011;17(1):49-53. doi: 10.1258/jtt.2010.100202. PMID: 21097567. Exclusion: Ineligible population.

141. Choi Yoo SJ, Nyman JA, Cheville AL, et al. Cost effectiveness of telecare management for pain and depression in patients with cancer: results from a randomized trial. *Gen Hosp Psychiatry*. 2014;36(6):599-606. doi: 10.1016/j.genhosppsych.2014.07.004. PMID: 25130518. Exclusion: Ineligible population.
142. Chowdhury M, Birns J, Rudd A, et al. Telemedicine versus face-to-face evaluation in the delivery of thrombolysis for acute ischaemic stroke: a single centre experience. *Postgrad Med J*. 2012;88(1037):134-7. doi: 10.1136/postgradmedj-2011-130060. PMID: 22267526. Exclusion: Ineligible population.
143. Chu K, Maine R, Duvenage R. We asked the experts: the role of rural hospitals in achieving equitable surgical access in low-resourced settings. *World J Surg*. 2021;45(10):3016-8. doi: 10.1007/s00268-021-06271-5. PMID: 34338826. Exclusion: Not a study.
144. Chua CY, Stoll C, Maki J, et al. Opportunities for connection: a scoping review of telemedicine interventions in colorectal cancerscreening in rural and nonrural settings. *Cancer Epidemiol Biomark Prev*. 2020;29(6 SUPPL 1)doi: 10.1158/1538-7755.DIS18-A011. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
145. Cira MK, Tesfay R, Zujewski JA, et al. Promoting evidence-based practices for breast cancer care through web-based collaborative learning. *J Cancer Policy*. 2020;25:1-6. doi: 10.1016/j.jcpo.2020.100242. PMID: 32719736. Exclusion: Ineligible population.
146. Clark S, Piercey C. E-learning provides nursing education in remote areas. *Aust Nurs J*. 2012;20(4):49. PMID: 23252119. Exclusion: Not a study.
147. Cohen GM, Irby MB, Boles K, et al. Telemedicine and pediatric obesity treatment: review of the literature and lessons learned. *Clin Obes*. 2012;2:103-11. PMID: 23227115. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
148. Comfere NI, Sokumbi O, Montori VM, et al. Provider-to-provider communication in dermatology and implications of missing clinical information in skin biopsy requisition forms: a systematic review. *Int J Dermatol*. 2014;53(5):549-57. doi: 10.1111/ijd.12330. PMID: 24116717. Exclusion: Ineligible intervention.
149. Costello A, Daoud AK, Piggott C, et al. A novel adaptation to the Extension for Community Healthcare Outcomes (ECHO) model. *Ann Fam Med*. 2021;19(4):371. doi: 10.1370/afm.2691. PMID: 34264851. Exclusion: Not a study.
150. Costello CM, Cumsy HJL, Maly CJ, et al. Improving access to care through the establishment of a local, teledermatology network. *Telemed J E Health*. 2020;26(7):935-40. doi: 10.1089/tmj.2019.0051. PMID: 31613713. Exclusion: Ineligible population.
151. Cota A, Tarchala M, Parent-Harvey C, et al. Review of 5.5 years' experience using e-mail-based telemedicine to deliver orthopedic care to remote communities. *Telemed J E Health*. 2017;23(1):37-40. doi: 10.1089/tmj.2016.0030. PMID: 27336754. Exclusion: Ineligible study design.
152. Cottrell M, Judd P, Comans T, et al. Comparing fly-in fly-out and telehealth models for delivering advanced-practice physiotherapy services in regional Queensland: an audit of outcomes and costs. *J Telemed Telecare*. 2019:1-7. doi: 10.1177/1357633X19858036. PMID: 31280639. Exclusion: Ineligible intervention.

153. Coustasse A, Sarkar R, Abodunde B, et al. Use of tele dermatology to improve dermatological access in rural areas. *Telemed J E Health*. 2019;25(11):1022-32. doi: 10.1089/tmj.2018.0130. PMID: 30741608. Exclusion: Ineligible intervention.
154. Cox P, Uhrh K, Scott J. Project ECHO hepatitis C: building community partnerships to improve quality care. *Commun Nurs Res*. 2012;45:175-. Exclusion: Ineligible comparater.
155. Craig J. A cohort study of early neurological consultation by telemedicine on the care of neurological inpatients. *J Neurol Neurosurg Psychiatry*. 2004;75(7):1031-5. doi: 10.1136/jnnp.2002.001651. PMID: 15201365. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
156. Crump WJ. Telemedicine: has the time really finally arrived? *J Rural Health*. 2020;37(1):156-7. doi: 10.1111/jrh.12435. PMID: 32277729. Exclusion: Not a study.
157. Curioso WH. Building capacity and training for digital health: challenges and opportunities in Latin America. *J Med Internet Res*. 2019;21(12):e16513. doi: 10.2196/16513. PMID: 31850849. Exclusion: Not a study.
158. Curran JA, Murphy AL, Sinclair D, et al. Factors influencing rural and urban emergency clinicians' participation in an online knowledge exchange intervention. *Rural Remote Health*. 2013;13(1):1-12. PMID: 23374031. Exclusion: Ineligible intervention.
159. Cuttriss N, Walker Walker A, Maahs D, et al. Democratizing type 1 diabetes (T1D) knowledge in rural and underserved communities: Project ECHO T1D. *Diabetes Technol Ther*. 2019;21:A118-A9. doi: 10.1089/dia.2019.2525.abstracts. Exclusion: Not a study.
160. Cuttriss N, Bouchonville MF, Maahs DM, et al. Tele-rounds and case-based training: Project ECHO telementoring model applied to complex diabetes care. *Pediatr Clin North Am*. 2020;67:759-72. doi: 10.1016/j.pcl.2020.04.017. PMID: 32650871. Exclusion: Not a study.
161. Damian AJ, Robinson S, Manzoor F, et al. A mixed methods evaluation of the feasibility, acceptability, and impact of a pilot project ECHO for community health workers (CHWs). *Pilot Feasibility Stud*. 2020;6:132. doi: 10.1186/s40814-020-00678-y. PMID: 32963804. Exclusion: Ineligible comparater.
162. Daniel ML, Julia CM, Sebastian M, et al. High impact of a national psoriasis telementoring clinic on medical practices for patients in underserved areas. *Dermatol Ther*. 2021;34(1):e14575. doi: 10.1111/dth.14575. PMID: 33230929. Exclusion: Ineligible study design.
163. Davis A, Hopkins T, Abrahams Y. Maximizing the impact of telepractice through a multifaceted service delivery model at the Shepherd Centre, Australia. *Volta Rev*. 2012;112(3):383-91. Exclusion: Not a study.
164. Davis RM, Pockl J, Bellis K. Improved diabetic eye care utilizing telemedicine: a randomized controlled trial. *Invest Ophthalmol Vis Sci*. 2003. Exclusion: Ineligible intervention.
165. Dawson AZ, Walker RJ, Campbell JA, et al. Telehealth and indigenous populations around the world: a systematic review on current modalities for physical and mental health. *Mhealth*. 2020;6(30):1-17. doi: 10.21037/mhealth.2019.12.03. PMID: 32632368. Exclusion: Ineligible intervention.
166. Day D, Sibson L, Agarwal S, et al. Clinical review of a network delivered stroke telemedicine thrombolysis service in a rural area. *Int J Stroke*. 2012;7:78. doi: 10.1111/j.1747-4930.2012.00961.x. Exclusion: Not a study.

167. Day SC, Day G, Keller M, et al. Personalized implementation of video telehealth for rural veterans (PIVOT-R). *Began with* 2015. 2021;7:24. doi: 10.21037/mhealth.2020.03.02. PMID: 33898593. Exclusion: Ineligible intervention.
168. Dayal P, Chang CH, Benko WS, et al. Appointment completion in pediatric neurology telemedicine clinics serving underserved patients. *Neurol Clin Pract*. 2019;9(4):314-21. doi: 10.1212/CPJ.0000000000000649. PMID: 31583186. Exclusion: Ineligible intervention.
169. de Batlle J, Massip M, Vargiu E, et al. Implementing mobile health-enabled integrated care for complex chronic patients: patients and professionals' acceptability study. *JMIR Mhealth Uhealth*. 2020;8(11):e22136. doi: 10.2196/22136. PMID: 33216004. Exclusion: Ineligible outcome.
170. DeGuzman PB, Vogel DL, Horton B, et al. Examination of a distress screening intervention for rural cancer survivors reveals low uptake of psychosocial referrals. *J Cancer Surviv*. 2021;13:13. doi: 10.1007/s11764-021-01052-4. PMID: 33983534. Exclusion: Ineligible intervention.
171. Deisz R, Rademacher S, Gilger K, et al. Additional telemedicine rounds as a successful performance-improvement strategy for sepsis management: observational multicenter study. *J Med Internet Res*. 2019;21(1):e11161. doi: 10.2196/11161. PMID: 30664476. Exclusion: Ineligible population.
172. Demaerschalk BM. Telestrokeologists: treating stroke patients here, there, and everywhere with telemedicine. *Semin Neurol*. 2010;30(5):477-91. doi: 10.1055/s-0030-1268869. PMID: 21207340. Exclusion: Not a study.
173. Demaerschalk BM, Kiernan TE, Investigators S. Vascular neurology nurse practitioner provision of telemedicine consultations. *Int J Telemed Appl*. 2010 doi: 10.1155/2010/507071. PMID: 20811594. Exclusion: Ineligible study design.
174. DeMaria AN. Virtual cardiology. *Structural Heart*. 2020;4(5):347-8. doi: 10.1080/24748706.2020.1812256. Exclusion: Not a study.
175. Demeestere J, Sewell C, Rudd J, et al. The establishment of a telestroke service using multimodal CT imaging decision assistance: "turning on the fog lights". *J Clin Neurosci*. 2017;37:1-5. doi: 10.1016/j.jocn.2016.10.018. PMID: 27887976. Exclusion: Ineligible setting.
176. Desai S, Williams ML, Smith AC. Teleconsultation from a secondary hospital for paediatric emergencies occurring at rural hospitals in Queensland. *J Telemed Telecare*. 2013;19(7):405-10. doi: 10.1177/1357633X13506528. PMID: 24218355. Exclusion: Ineligible comparator.
177. Deslich S, Coustasse A. Expanding technology in the ICU: the case for the utilization of telemedicine. *Telemed J E Health*. 2014;20(5):485-92. doi: 10.1089/tmj.2013.0102. PMID: 24621447. Exclusion: Not a study.
178. DeVido J, Glezer A, Branagan L, et al. Telepsychiatry for inpatient consultations at a separate campus of an academic medical center. *Telemed J E Health*. 2016;22(7):572-6. doi: 10.1089/tmj.2015.0125. PMID: 26701608. Exclusion: Ineligible intervention.
179. Dharmasaroja PA, Muengtawepong S, Kommarkg U. Implementation of Telemedicine and Stroke Network in thrombolytic administration: comparison between walk-in and referred patients. *Neurocrit Care*. 2010;13(1):62-6. doi: 10.1007/s12028-010-9360-3. PMID: 20411354. Exclusion: Ineligible population.

180. Di Lieto A, De Falco M, Pontillo M, et al. The wireless tele-ultrasonography in prenatal telemedicine. *G Ital di Ostet e Ginecol.* 2011;33(3):158-62. Exclusion: Not a study.
181. Diaz S, Zhao J, Cronin S, et al. Changes in opioid prescribing behaviors among family physicians who participated in a weekly tele-mentoring program. *J Clin Med.* 2019;9(14):1-11. doi: 10.3390/jcm9010014. PMID: 31861584. Exclusion: Ineligible population.
182. Diego E, McAuliffe P, McGuire K, et al. Telemedicine for delivery of specialized breast care in rural areas: a pilot study. *Ann Surg Oncol.* 2014;21:43-4. doi: 10.1245/s10434-014-3672-z. Exclusion: Ineligible intervention.
183. Dietrich M, Walter S, Ragoschke-Schumm A, et al. Is prehospital treatment of acute stroke too expensive? An economic evaluation based on the first trial. *Cerebrovasc Dis.* 2014;38(6):457-63. doi: 10.1159/000371427. PMID: 25531507. Exclusion: Ineligible study design.
184. Dingwall KM, Puszka S, Sweet M, et al. Evaluation of a culturally adapted training course in Indigenous e-mental health. *Australas Psychiatry.* 2015;23(6):630-5. doi: 10.1177/1039856215608282. PMID: 26423096. Exclusion: Ineligible intervention.
185. Diop IB, Leye M, Ba AT, et al. Telemedicine: tele education in Senegal - what are our needs and realisations? *Cardiovasc J Afr.* 2011;22(3):S7. Exclusion: Ineligible setting.
186. Disler R, Glenister K, Wright J. Rural chronic disease research patterns in the United Kingdom, United States, Canada, Australia and New Zealand: a systematic integrative review. *BMC Public Health.* 2020;20(770):1-7. doi: 10.1186/s12889-020-08912-1. PMID: 32448173. Exclusion: Ineligible intervention.
187. Doherty M, Modanloo S, Evans E, et al. Exploring health professionals' experiences with a virtual learning and mentoring program (Project ECHO) on pediatric palliative care in South Asia. *Glob Pediatr Health.* 2021;8:2333794X211043061. doi: 10.1177/2333794X211043061. PMID: 34485624. Exclusion: Ineligible population.
188. Doherty M, Rayala S, Evans E, et al. Using virtual learning to build pediatric palliative care capacity in South Asia: experiences of implementing a teleteaching and mentorship program (Project ECHO). *JCO Glob Oncol.* 2021;7:210-22. doi: 10.1200/GO.20.00481. PMID: 33555911. Exclusion: Ineligible study design.
189. Donnem T, Ervik B, Magnussen K, et al. Bridging the distance: a prospective tele-oncology study in Northern Norway. *Support Care Cancer.* 2012;20(9):2097-103. doi: 10.1007/s00520-011-1319-1. PMID: 22076621. Exclusion: Ineligible comparater.
190. Dowling MJ, Payne C, Larkin P, et al. Does an interactive, teleconference-delivered, palliative care lecture series improve nursing home staff confidence? *J Palliat Med.* 2020;23(2):179-83. doi: 10.1089/jpm.2018.0549. PMID: 31390293. Exclusion: Ineligible population.
191. Du P, Wang X, Kong L, et al. Can telementoring reduce urban-rural disparities in utilization of direct-acting antiviral agents? *Telemed J E Health.* 2020;02:02. doi: 10.1089/tmj.2020.0090. PMID: 32882154. Exclusion: Ineligible comparater.
192. du Toit M, Malau-Aduli B, Vangaveti V, et al. Use of telehealth in the management of non-critical emergencies in rural or remote emergency departments: a systematic review. *J Telemed Telecare.* 2019;25(1):3-16. doi: 10.1177/1357633X17734239. PMID: 28980853. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.

193. Duarte SS, Nguyen TT, Koch C, et al. Remote obstetric anesthesia: leveraging telemedicine to improve fetal and maternal outcomes. *Telemed J E Health*. 2019;26(8):967-72. doi: 10.1089/tmj.2019.0174. PMID: 31710285. Exclusion: Not a study.
194. DuGan JE. "Keeping you in the know": The effect of an online nursing journal club on evidence-based knowledge among rural registered nurses. *Comput Inform Nurs*. 2019;37(4):190-5. doi: 10.1097/CIN.0000000000000507. PMID: 30688671. Exclusion: Ineligible intervention.
195. Duncanson K, Webster EL, Schmidt DD. Impact of a remotely delivered, writing for publication program on publication outcomes of novice researchers. *Rural Remote Health*. 2018;18(2):4468. doi: 10.22605/RRH4468. PMID: 29793344. Exclusion: Ineligible study design.
196. Durbha S, Commiskey P, Crawley R. Attitudes of emergency medical service personnel towards implementation of technology-enabled stroke care in rural areas. *Eur Stroke J*. 2021;6(1 SUPPL):512. doi: 10.1177/23969873211034932. Exclusion: Ineligible study design.
197. Dutta D, Kendall J, Holmes C, et al. Evaluation of a telephone advice system for remote intravenous thrombolysis in ischemic stroke: data from a United kingdom network. *Stroke*. 2015;46(3):867-9. doi: 10.1161/STROKEAHA.114.008190. PMID: 25604248. Exclusion: Ineligible population.
198. Dy P, Morin PC, Weinstock RS. Use of telemedicine to improve glycemic management in a skilled nursing facility: a pilot study. *Telemed J E Health*. 2013;19(8):643-5. doi: 10.1089/tmj.2012.0274. PMID: 23758078. Exclusion: Ineligible population.
199. Eadie L, Regan L, Mort A, et al. Telestroke assessment on the move: prehospital streamlining of patient pathways. *Stroke*. 2015;46(2):e38-40. doi: 10.1161/STROKEAHA.114.007475. PMID: 25550375. Exclusion: Ineligible population.
200. Eadie L, Mulhern J, Regan L, et al. Remotely supported prehospital ultrasound: a feasibility study of real-time image transmission and expert guidance to aid diagnosis in remote and rural communities. *J Telemed Telecare*. 2018;24(9):616-22. doi: 10.1177/1357633X17731444. PMID: 28920524. Exclusion: Ineligible intervention.
201. Easom LR, Alston G, Coleman R. A rural community translation of a dementia caregiving intervention. *Online J Rural Nurs Health Care*. 2013;13(1):66-91. doi: 10.14574/ojrnhc.v13i1.248. Exclusion: Ineligible intervention.
202. Eaton LH, Gordon DB, Wyant S, et al. Development and implementation of a telehealth-enhanced intervention for pain and symptom management. *Contemp Clin Trials*. 2014;38(2):213-20. doi: 10.1016/j.cct.2014.05.005. PMID: 24846620. Exclusion: Ineligible outcome.
203. Eaton LH, Godfrey DS, Langford DJ, et al. Telementoring for improving primary care provider knowledge and competence in managing chronic pain: A randomised controlled trial. *J Telemed Telecare*. 2020;26(1-2):21-7. doi: 10.1177/1357633x18802978. PMID: 30261805. Exclusion: Ineligible population.
204. Egede LE, Knapp RG, Walker RJ, et al. Randomized controlled trial of technology-assisted case management in low-income adults with type 2 diabetes (TACMDM): effect on quality of life and blood pressure. *Diabetes*. 2020;69doi: 10.2337/db20-651-P. Exclusion: Ineligible intervention.

205. Eibl JK, Gauthier G, Pellegrini D, et al. The effectiveness of telemedicine-delivered opioid agonist therapy in a supervised clinical setting. *Drug Alcohol Depend.* 2017;176:133-8. doi: 10.1016/j.drugalcdep.2017.01.048. PMID: 28535455. Exclusion: Ineligible population.
206. Eisenstein EL, Lobach DF, Kawamoto K, et al. A randomized clinical trial of clinical decision support in a rural community health network serving lower income individuals: study design and baseline characteristics. *Stud Health Technol Inform.* 2009;143:220-6. PMID: 19380940. Exclusion: Ineligible outcome.
207. Ellis MJ, Mendez I, Russell K. Preliminary clinical algorithm to optimise remote delivery of paediatric concussion care in Canada's North. *Int J Circumpolar Health.* 2020;79(1):1832390. doi: 10.1080/22423982.2020.1832390. PMID: 33089768. Exclusion: Ineligible population.
208. Englander H, Patten A, Lockard R, et al. Spreading addictions care across Oregon's rural and community hospitals: mixed-methods evaluation of an interprofessional telementoring ECHO program. *J Gen Intern Med.* 2021;36(1):100-7. doi: 10.1007/s11606-020-06175-5. PMID: 32885371. Exclusion: Ineligible population.
209. Ens CD, Hanlon-Dearman A, Millar MC, et al. Using telehealth for assessment of fetal alcohol spectrum disorder: the experience of two Canadian rural and remote communities. *Telemed J E Health.* 2010;16(8):872-7. doi: 10.1089/tmj.2010.0070. PMID: 20925569. Exclusion: Ineligible intervention.
210. Estai M, Bunt S, Kanagasingam Y, et al. Cost savings from a teledentistry model for school dental screening: an Australian health system perspective. *Aust Health Rev.* 2018;42(5):482-90. doi: 10.1071/AH16119. PMID: 28578759. Exclusion: Ineligible intervention.
211. Etingen B, Patrianakos J, Wirth M, et al. TeleWound practice within the Veterans Health Administration: protocol for a mixed methods program evaluation. *JMIR Res Protoc.* 2020;9(7):e20139. doi: 10.2196/20139. PMID: 32706742. Exclusion: Not a study.
212. Evangelista A, Galuppo V, Mendez J, et al. Hand-held cardiac ultrasound screening performed by family doctors with remote expert support interpretation. *Heart.* 2016;102(5):376-82. doi: 10.1136/heartjnl-2015-308421. PMID: 26802099. Exclusion: Ineligible population.
213. Evans NR, Lebus CS, Day DJ, et al. Delivery of out of hours thrombolysis in a semi-rural area: comparing an in-house centre versus telemedicine service. *Cerebrovasc Dis.* 2014;37:460. doi: 10.1159/000363616. Exclusion: Ineligible study design.
214. Faherty LJ, Rose AJ, Chappel A, et al. Assessing and expanding the evidence base for Project ECHO and ECHO-like models: findings of a technical expert panel. *J Gen Intern Med.* 2020;35(3):899-902. doi: 10.1007/s11606-019-05599-y. PMID: 31925737. Exclusion: Ineligible outcome.
215. Fahey JO, Cohen SR, Holme F, et al. Promoting cultural humility during labor and birth: putting theory into action during PRONTO obstetric and neonatal emergency training. *J Perinat Neonatal Nurs.* 2013;27(1):36-42. doi: 10.1097/JPN.0b013e31827e478d. PMID: 23360940. Exclusion: Ineligible setting.
216. Fairchild R, Ferng-Kuo SF, Rahmouni H, et al. An observational study of telemental care delivery and the context for involuntary commitment for mental health patients in a group of rural emergency departments. *Telemed Rep.* 2020;1(1):22-35. doi: 10.1089/tmr.2020.0005. PMID: 33283206. Exclusion: Ineligible study design.

217. Falconer E, Kho D, Docherty JP. Use of technology for care coordination initiatives for patients with mental health issues: a systematic literature review. *Neuropsychiatr Dis Treat*. 2018;14:2337-49. doi: 10.2147/NDT.S172810. PMID: 30254446. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
218. Fang JL. Use of telemedicine to support neonatal resuscitation. *Semin Perinatol*. 2021;151423. doi: 10.1016/j.sempri.2021.151423. PMID: 33958229. Exclusion: Not a study.
219. Fang JL, Umoren RA, Whyte H, et al. Provider perspectives on the acceptability, appropriateness, and feasibility of teleneonatology. *Am J Perinatol*. 2021;28:28. doi: 10.1055/a-1656-6363. PMID: 34583392. Exclusion: Ineligible comparater.
220. Farshidi D, Craft N, Ochoa MT. Mobile tele dermatology: as doctors and patients are increasingly mobile, technology keeps us connected. *SKINmed*. 2011;9(4):231-8. PMID: 21980708. Exclusion: Not a study.
221. Fatehi F, Jahedi F, Tay-Kearney ML, et al. Teleophthalmology for the elderly population: a review of the literature. *Int J Med Inf*. 2020;136:104089. doi: 10.1016/j.ijmedinf.2020.104089. PMID: 32044698. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
222. Felfeli T, Alon R, Merritt R, et al. Toronto tele-retinal screening program for detection of diabetic retinopathy and macular edema. *Can J Ophthalmol*. 2019;54(2):203-11. doi: 10.1016/j.jcjo.2018.07.004. PMID: 30975344. Exclusion: Ineligible population.
223. Ferguson-Pell M, Armstrong E. Patient and clinician reported outcomes and experiences following advanced tele-rehabilitation assessments conducted in rural Alberta. *Arch Phys Med Rehabil*. 2021;102(10):e10. doi: 10.1016/j.apmr.2021.07.419. Exclusion: Ineligible intervention.
224. Ferrer J, Chaumont T, Trujillo L, et al. New tele-diagnostic model using volume sweep imaging for rural areas. *Conf Proc IEEE Eng Med Biol Soc*. 2017;2017:2622-5. doi: 10.1109/EMBC.2017.8037395. PMID: 29060437. Exclusion: Ineligible study design.
225. Finley BA, Palitsky R, Charteris E, et al. Outpatient telecardiology perceptions among rural, suburban, and urban veterans utilizing in-person cardiology versus telecardiology services: a mixed methods analysis. *J Rural Health*. 2021;18:18. doi: 10.1111/jrh.12586. PMID: 34002404. Exclusion: Ineligible intervention.
226. Fisher E, Hasselberg M, Conwell Y, et al. Telementoring primary care clinicians to improve geriatric mental health care. *Popul Health Manag*. 2017;20(5):342-7. doi: 10.1089/pop.2016.0087. PMID: 28106511. Exclusion: Ineligible population.
227. Flodgren G, Rachas A, Farmer AJ, et al. Interactive telemedicine: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev*. 2016 (12). Exclusion: Ineligible intervention.
228. Flores A, Lazaro SA, Molina-Bastos CG, et al. Teledentistry in the diagnosis of oral lesions: a systematic review of the literature. *J Am Med Inform Assoc*. 2020;27(7):1166-72. doi: 10.1093/jamia/ocaa069. PMID: 32568392. Exclusion: Ineligible setting.
229. Fong WC, Ismail M, Lo JW, et al. Telephone and Teleradiology-Guided Thrombolysis Can Achieve Similar Outcome as Thrombolysis by Neurologist On-site. *J Stroke Cerebrovasc Dis*. 2015;24(6):1223-8. doi: 10.1016/j.jstrokecerebrovasdis.2015.01.022. PMID: 25906936. Exclusion: Ineligible population.

230. Fowler RC, Katzman JG, Comerici GD, et al. Mock ECHO: a simulation-based medical education method. *Teach Learn Med.* 2018;30(4):423-32. doi: 10.1080/10401334.2018.1442719. PMID: 29658798. Exclusion: Ineligible comparater.
231. Francis E, Shifler Bowers K, Buchberger G, et al. Reducing alcohol and opioid use among youth in rural counties: an innovative training protocol for primary health care providers and school personnel. *JMIR Res Protoc.* 2020;9(11):e21015. doi: 10.2196/21015. PMID: 33155572. Exclusion: Ineligible outcome.
232. Freeman BA. Technology, legislators, associations, and companies are encouraging tele-audiology. *Hear J.* 2010;63(6):50-1. Exclusion: Not a study.
233. Friesner DL, Scott DM, Rathke AM, et al. Do remote community telepharmacies have higher medication error rates than traditional community pharmacies? Evidence from the North Dakota Telepharmacy Project. *J Am Pharm Assoc* (2003). 2011;51(5):580-90. doi: 10.1331/JAPhA.2011.10115. PMID: 21896455. Exclusion: Ineligible intervention.
234. Fry MW, Saidi S, Musa A, et al. "Even though I am alone, I feel that we are many" - an appreciative inquiry study of asynchronous, provider-to-provider teleconsultations in Turkana, Kenya. *PLoS One.* 2020;15(9):e0238806. doi: 10.1371/journal.pone.0238806. PMID: 32931503. Exclusion: Ineligible setting.
235. Furlan AD, Pajer KA, Gardner W, et al. Project ECHO: building capacity to manage complex conditions in rural, remote and underserved areas. *Can J Rural Med.* 2019;24(4):115-20. doi: 10.4103/CJRM.CJRM_20_18. PMID: 31552868. Exclusion: Not a study.
236. Furlan AD, Zhao J, Voth J, et al. Evaluation of an innovative tele-education intervention in chronic pain management for primary care clinicians practicing in underserved areas. *J Telemed Telecare.* 2019;25(8):484-92. doi: 10.1177/1357633X18782090. PMID: 29991316. Exclusion: Ineligible population.
237. Furness BW, Goldhammer H, Montalvo W, et al. Transforming primary care for lesbian, gay, bisexual, and transgender people: a collaborative quality improvement initiative. *Ann Fam Med.* 2020;18(4):292-302. doi: 10.1370/afm.2542. PMID: 32661029. Exclusion: Ineligible population.
238. Furness L, Kaltner M. Facilitating student education and support using videoconference. *Aust J Rural Health.* 2015;23(4):249-53. doi: 10.1111/ajr.12148. PMID: 26122604. Exclusion: Ineligible study design.
239. Gabriel K, Hillmann S, Kraft P, et al. Two-year experience of implementing a comprehensive telemedical network within a mainly rural area in Germany: the transregional network for stroke intervention with telemedicine (transit-stroke). *Eur Stroke J.* 2018;3(1):282. doi: 10.1177/2396987318770127. Exclusion: Ineligible study design.
240. Gabriel KMA, Jiru-Hillmann S, Kraft P, et al. Two years' experience of implementing a comprehensive telemedical stroke network comprising in mainly rural region: the Transregional Network for Stroke Intervention with Telemedicine (TRANSIT-Stroke). *BMC Neurol.* 2020;20(104):1-13. doi: 10.1186/s12883-020-01676-6. PMID: 32192438. Exclusion: Ineligible comparater.
241. Gali K, Joshi S, Hueneker S, et al. Barriers, access and management of paediatric epilepsy with telehealth. *J Telemed Telecare.* 2020;1357633X20969531. doi: 10.1177/1357633X20969531. PMID: 33183129. Exclusion: Ineligible comparater.

242. Gallerini S, Marsili L, Groccia V, et al. Appropriateness, safety, and effectiveness of "drip and ship" teleconsultation model in Southeastern Tuscany: a feasibility study. *Neurol Sci.* 2020;41:2961-5. doi: 10.1007/s10072-020-04446-x. PMID: 32383049. Exclusion: Ineligible study design.
243. Galop BN, Fink JL. The impact of telemedicine on pharmacists and implications for controlled substances. *Pharm Times.* 2017;83(12). Exclusion: Not a study.
244. Galván P, Velázquez M, Rivas R, et al. Health diagnosis improvement in remote community health centers through telemedicine. *Med Access Point Care.* 2018;2:1-4. doi: 10.1177/239920261775310. Exclusion: Ineligible study design.
245. Galván P, Rivas R, Grossling B, et al. Building a virtual diagnosis network through a telemedicine platform. *Int J Technol Assess Health Care.* 2019;35:82-3. Exclusion: Ineligible outcome.
246. Galván P, Velázquez M, Rivas R, et al. Telemedicine enhances community hospital response capacity. *Int J Technol Assess Health Care.* 2019;35:62. Exclusion: Ineligible outcome.
247. Gamus A, Keren E, Kaufman H, et al. Synchronous video telemedicine in lower extremities ulcers treatment: a real-world data study. *Int J Med Inf.* 2019;124:31-6. doi: 10.1016/j.ijmedinf.2019.01.009. PMID: 30784424. Exclusion: Ineligible population.
248. Gangadharan S, Lillicrap T, Miteff F, et al. Air vs. road decision for endovascular clot retrieval in a rural telestroke network. *Front Neurol.* 2020;11:628. doi: 10.3389/fneur.2020.00628. PMID: 32765396. Exclusion: Ineligible intervention.
249. Garber R, Garcia E, Deeter L. Pictures influence the decision to transfer: outcomes of a telemedicine program serving an 8 state rural population. *J Burn Care Res.* 2018;39:S20. doi: 10.1093/jbcr/iry006.033. PMID: 32044972. Exclusion: Ineligible comparater.
250. Garg D, Majumdar R, Chauhan S, et al. Teleneurorehabilitation among person with Parkinson's disease in India: the initial experience and barriers to implementation. *Ann Indian Acad Neurol.* 2021;24(4):536-41. doi: 10.4103/aian.AIAN_127_21. PMID: 34728947. Exclusion: Ineligible setting.
251. Garrelts JC, Gagnon M, Eisenberg C, et al. Impact of telepharmacy in a multihospital health system. *Am J Health Syst Pharm.* 2010;67(17):1456-62. doi: 10.2146/ajhp090670. PMID: 20720245. Exclusion: Ineligible population.
252. Gatica JL, Bertoló S, Morales E, et al. Store-and-forward teledermatology in Chile: a contribution to primary health care. *Piel.* 2015;30(3):148-54. doi: 10.1016/j.piel.2014.07.007. Exclusion: Not English language.
253. Gensichen J, von Korff M, Peitz M, et al. Case management for depression by health care assistants in small primary care practices: a cluster randomized trial. *Ann Intern Med.* 2009;151(6):369-78. PMID: 19755362. Exclusion: Ineligible population.
254. Gentry MT, Lapid MI, Rummans TA. Geriatric telepsychiatry: systematic review and policy considerations. *Am J Geriatr Psychiatry.* 2019;27(2):109-27. doi: 10.1016/j.jagp.2018.10.009. PMID: 30416025. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.

255. Gephart SM, Newnam K, Weiss A, et al. Feasibility and acceptability of a Neonatal Project ECHO (NeoECHO) as a dissemination and implementation strategy to prevent necrotizing enterocolitis. *Worldviews Evid Based Nurs.* 2021;23:23. doi: 10.1111/wvn.12529. PMID: 34296821. Exclusion: Ineligible outcome.
256. Gersch WD, Delate T, Bergquist KM, et al. Clinical effectiveness of an outpatient multidisciplinary chronic pain management telementoring service. *Clinical Journal of Pain.* 2021;37(10):740-6. doi: 10.1097/AJP.0000000000000967. PMID: 34265787. Exclusion: Ineligible setting.
257. Giachetto G, Casuriaga AL, Santoro A, et al. Extension for Community Healthcare Outcomes Uruguay: a new strategy to promote best primary care practice for autism. *Glob Pediatr Health.* 2019;6:1-7. doi: 10.1177/2333794X19833734. PMID: 31044151. Exclusion: Ineligible population.
258. Gibson N, Arends R, Voss J, et al. Reinforcing telehealth competence through nurse practitioner student clinical experiences. *J Nurs Educ.* 2020;59(7):413-7. doi: 10.3928/01484834-20200617-12. PMID: 32598014. Exclusion: Ineligible study design.
259. Gilman M, Stensland J. Telehealth and Medicare: payment policy, current use, and prospects for growth. *Medicare Medicaid Res Rev.* 2013;3(4)doi: 10.5600/mmrr.003.04.a04. PMID: 24834368. Exclusion: Ineligible study design.
260. Glass LM, Waljee AK, McCurdy H, et al. Specialty care access network-extension of community healthcare outcomes model program for liver disease improves specialty care access. *Dig Dis Sci.* 2017;62(12):3344-9. doi: 10.1007/s10620-017-4789-2. PMID: 29043594. Exclusion: Ineligible comparater.
261. Gleason LJ, Martinchek M, Long M, et al. An innovative model using telementoring to provide geriatrics education for nurses and social workers at skilled nursing facilities. *Geriatr Nurs.* 2019;40(5):517-21. doi: 10.1016/j.gerinurse.2019.03.018. PMID: 30987777. Exclusion: Ineligible population.
262. Gokula M, Gaspar P, King B, et al. Effect of implementation of the tele-health visiting nurse association project on re-hospitalization rate and cost. *J Am Med Dir Assoc.* 2015;16(3):B26-7. doi: 10.1016/j.jamda.2015.01.058. Exclusion: Ineligible intervention.
263. Golberstein E, Kolvenbach S, Carruthers H, et al. Effects of electronic psychiatric consultations on primary care provider perceptions of mental health care: Survey results from a randomized evaluation. *Healthc (Amst).* 2018;6(1):17-22. doi: 10.1016/j.hjdsi.2017.01.002. PMID: 28162990. Exclusion: Ineligible population.
264. Goldstein KM, Gierisch JM, Zullig LL, et al. Telehealth services designed for women: an evidence map. 2017. <https://www.ncbi.nlm.nih.gov/books/NBK532568/> PMID: 30383345. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
265. Goncalves-Bradley DC, J Maria RA, Ricci-Cabello I, et al. Mobile technologies to support healthcare provider to healthcare provider communication and management of care. *Cochrane Database Syst Rev.* 2020 (8)doi: 10.1002/14651858.CD012927.pub2. PMID: 32813281. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
266. Gopalan P, Auster L, Brockman I, et al. Telemedicine psychiatric consultation on an inpatient obstetrical labor and delivery unit. *J Acad Consult Liaison Psychiatry.* 2021;20:20. doi: 10.1016/j.jaclp.2021.04.001. PMID: 33972195. Exclusion: Ineligible study design.

267. Gordon B, Mason B, Smith SLH. Leveraging telehealth for delivery of palliative care to remote communities: a rapid review. *J Palliat Care*. 2021;8258597211001184. doi: 10.1177/08258597211001184. PMID: 33730904. Exclusion: Ineligible intervention.
268. Gordon SE, Dufour AB, Monti SM, et al. Impact of a videoconference educational intervention on physical restraint and antipsychotic use in nursing homes: results from the ECHO-AGE pilot study. *J Am Med Dir Assoc*. 2016;17(6):553-6. doi: 10.1016/j.jamda.2016.03.002. PMID: 27161317. Exclusion: Ineligible population.
269. Grady B, Singleton M. Telepsychiatry "coverage" to a rural inpatient psychiatric unit. *Telemed J E Health*. 2011;17(8):603-8. doi: 10.1089/tmj.2011.0031. PMID: 21939381. Exclusion: Ineligible population.
270. Grady B. Promises and limitations of telepsychiatry in rural adult mental health care. *World Psychiatry*. 2012;11(3):199-201. doi: 10.1002/j.2051-5545.2012.tb00132.x. PMID: 23024682. Exclusion: Not a study.
271. Gray LC, Fatehi F, Martin-Khan M, et al. Telemedicine for specialist geriatric care in small rural hospitals: preliminary data. *J Am Geriatr Soc*. 2016;64(6):1347-51. doi: 10.1111/jgs.14139. PMID: 27321617. Exclusion: Ineligible study design.
272. Graziane JA, Gopalan P, Cahalane J. Telepsychiatry consultation for medical and surgical inpatient units. *Psychosomatics*. 2018;59(1):62-6. doi: 10.1016/j.psym.2017.08.008. PMID: 28918164. Exclusion: Ineligible study design.
273. Griffiths F, Watkins JA, Huxley C, et al. Mobile consulting (mConsulting) and its potential for providing access to quality healthcare for populations living in low-resource settings of low- and middle-income countries. *Digit Health*. 2020;6:1-17. doi: 10.1177/2055207620919594. PMID: 32341793. Exclusion: Ineligible setting.
274. Grona SL, Bath B, Busch A, et al. Use of videoconferencing for physical therapy in people with musculoskeletal conditions: a systematic review. *J Telemed Telecare*. 2018;24(5):341-55. doi: 10.1177/1357633X17700781. PMID: 28403669. Exclusion: Ineligible intervention.
275. Gross IT, Whitfill T, Redmond B, et al. Comparison of two telemedicine delivery modes for neonatal resuscitation support: a simulation-based randomized trial. *Neonatology*. 2020;117:159-66. doi: 10.1159/000504853. PMID: 31905354. Exclusion: Ineligible population.
276. Gross J, Langford DJ, Tauben DJ. Evolution of a telehealth course during the global pandemic: from temporary elective to permanent rural education program. *Pain Med*. 2021;04:04. doi: 10.1093/pm/pnab266. PMID: 34480577. Exclusion: Not a study.
277. Grubbs KM, Fortney JC, Pyne J, et al. A comparison of collaborative care outcomes in two health care systems: VA clinics and federally qualified health centers. *Psychiatr Serv*. 2018;69(4):431-7. doi: 10.1176/appi.ps.201700067. PMID: 29334874. Exclusion: Ineligible intervention.
278. Gu J, Zhu S, Chen T, et al. Evaluation of the Spring Seedling Project-Zhaotong Program: a study of a novel continuing medical education program for rural doctors in China. *Aust J Rural Health*. 2020;28(5):434-42. doi: 10.1111/ajr.12659. PMID: 32985023. Exclusion: Ineligible setting.
279. Gupta S, Dewan S, Kaushal A, et al. eICU reduces mortality in STEMI patients in resource-limited areas. *Glob Heart*. 2014;9(4):425-7. doi: 10.1016/j.gheart.2014.07.006. PMID: 25592796. Exclusion: Ineligible setting.
280. Gurbeta L, Badnjevic A, Maksimovic M, et al. A telehealth system for automated diagnosis of asthma and chronic obstructive pulmonary disease. *J Am Med Inform Assoc*. 2018;25(9):1213-7. doi: 10.1093/jamia/ocy055. PMID: 29788482. Exclusion: Ineligible intervention.

281. Guthrie CP, Sadler T, Wadle C, et al. Implementation of telemedicine in a rural cystic fibrosis clinic. *Pediatr Pulmonol*. 2020;55(SUPPL 2):360-1. doi: 10.1002/ppul.24592. Exclusion: Not a study.
282. Gutierrez J, Kuperman E, Kaboli PJ. Using telehealth as a tool for rural hospitals in the COVID-19 pandemic response. *J Rural Health*. 2020;37(1):161-4. doi: 10.1111/jrh.12443. PMID: 32277777. Exclusion: Not a study.
283. Hah H, Goldin D, Ha S. Title correction: the association between willingness of frontline care providers' to adaptively use telehealth technology and virtual service performance in provider-to-provider communication: quantitative study. *J Med Internet Res*. 2019;21(11):e17123. doi: 10.2196/17123. PMID: 31743890. Exclusion: Not a study.
284. Hakim A, Gaviria-Agudelo C, Edwards K, et al. Pre-Coronavirus disease 2019 telehealth practices among pediatric infectious diseases specialists in the United States. *J Pediatric Infect Dis Soc*. 2021;10(4):485-91. doi: 10.1093/jpids/piaa146. PMID: 33196843. Exclusion: Ineligible setting.
285. Hall MA, Witkop ML, Rushlow D, et al. Mind the gap: nurse practitioner lead hemophilia treatment center providing healthcare to rural populations utilizing collaborative agreements and telemedicine. *Am J Hematol*. 2014;89(6):E81. doi: 10.1002/ajh.23759. Exclusion: Ineligible comparater.
286. Hallberg S, Harrison D. Telemedicine via continuous remote care: a proactive, patient-centered approach to improve clinical outcomes. *JMIR Diabetes*. 2021;31:31. doi: 10.2196/23646. PMID: 34505578. Exclusion: Not a study.
287. Hallikainen J, Castrén M, Saari A. Emergency medical services in rural areas of Finland. *Notf Rett Med*. 2014;17(3):199-201. doi: 10.1007/s10049-013-1789-4. Exclusion: Not a study.
288. Hamilton SJ, Mills B, Birch EM, et al. Smartphones in the secondary prevention of cardiovascular disease: a systematic review. *BMC Cardiovasc Disord*. 2018;18(1):25. doi: 10.1186/s12872-018-0764-x. PMID: 29415680. Exclusion: Ineligible intervention.
289. Hampton L, Brindley P, Kirkpatrick A, et al. Strategies to improve communication in telementoring in acute care coordination: a scoping review. *Can J Surg*. 2020;63(6):E569-E77. doi: 10.1503/cjs.015519. PMID: 33253511. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
290. Hand LJ. The role of telemedicine in rural mental health care around the globe. *Telemed J E Health*. 2021;31:31. doi: 10.1089/tmj.2020.0536. PMID: 34061678. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies. REVIEW REFERENCE LIST
291. Haridy J, Iyngkaran G, Nicoll A, et al. eHealth technologies for screening, diagnosis and management of viral hepatitis: a systematic review. *Clin Gastroenterol Hepatol*. 2020;04:04. doi: 10.1016/j.cgh.2020.09.011. PMID: 32896632. Exclusion: Ineligible intervention.
292. Hashmi NR, Khan SA. Interventional study to improve diabetic guidelines adherence using mobile health (m-Health) technology in Lahore, Pakistan. *BMJ Open*. 2018;8(5):e020094. doi: 10.1136/bmjopen-2017-020094. PMID: 29858411. Exclusion: Ineligible setting.
293. Hassan S, Carlin L, Zhao J, et al. Promoting an interprofessional approach to chronic pain management in primary care using Project ECHO. *J Interprof Care*. 2020;1-4. doi: 10.1080/13561820.2020.1733502. PMID: 32148115. Exclusion: Ineligible study design.

294. Hasselberg MJ, Fisher E, Conwell Y, et al. Implementing project extension for community healthcare outcomes for geriatric mental healthcare in long-term care facilities. *J Am Med Dir Assoc*. 2019;20(12):1651-3. doi: 10.1016/j.jamda.2019.07.012. PMID: 31466936. Exclusion: Ineligible population.
295. Hastie Norris S, Matthews SW, Hogan MT. Development of a teledermatology service in rural Washington state using the Doctor of Nursing practice skill set and implementation science. *J Dermatol Nurses Assoc*. 2021;13(4):214-8. doi: 10.1097/JDN.0000000000000625. Exclusion: Not a study.
296. Haveland S, Islam S. Key considerations in ensuring a safe regional telehealth care model: a systematic review. *Telemed J E Health*. 2021;05:05. doi: 10.1089/tmj.2020.0580. PMID: 33956524. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
297. Haynes SC, Hoffman K, Smith S, et al. Telemedicine consultations may improve stability of neonates prior to transport. *J Investig Med*. 2020;68(1):A201. doi: 10.1136/jim-2019-WMRC.464. Exclusion: Not a study.
298. Haynes SC, Marcin JP, Dayal P, et al. Impact of telemedicine on visit attendance for paediatric patients receiving endocrinology specialty care. *J Telemed Telecare*. 2020;1357633X20972911. doi: 10.1177/1357633X20972911. PMID: 33226895. Exclusion: Ineligible intervention.
299. Heffner DL, Thirumala PD, Pokharna P, et al. Outcomes of Spoke-Retained Telestroke Patients Versus Hub-Treated Patients After Intravenous Thrombolysis: Telestroke Patient Outcomes After Thrombolysis. *Stroke*. 2015;46(11):3161-7. doi: 10.1161/STROKEAHA.115.009980. PMID: 26396027. Exclusion: Ineligible population.
300. Heifetz LJ, Koppel AB, Kaime EM, et al. Addressing rural disparities in cancer care via telehealth. *J Clin Oncol*. 2020;38(15)doi: 10.1200/JCO.2020.38.15_suppl.e19090. Exclusion: Ineligible intervention.
301. Heitkamp T, Schroeder S, Shogren M, et al. Use of evidence-based technology to assist the behavioral health workforce: a case study on technology transfer systems. *J Addict Nurs*. 2021;32(3):197-204. doi: 10.1097/JAN.0000000000000418. PMID: 34473449. Exclusion: Not a study.
302. Henneker C, Lam WS, Panuccio M. An innovative nursing model to support country people receiving oral treatment for cancer. *Asia Pac J Clin Oncol*. 2018;14:117. doi: 10.1111/ajco.13089. Exclusion: Ineligible intervention.
303. Hensel JM, Ellard K, Koltek M, et al. Digital health solutions for indigenous mental well-being. *Curr Psychiatry Rep*. 2019;21(8):68. doi: 10.1007/s11920-019-1056-6. PMID: 31263971. Exclusion: Not a study.
304. Heppner S, Mohr NM, Carter KD, et al. HRSA's evidence-based tele-emergency network grant program: multi-site prospective cohort analysis across six rural emergency department telemedicine networks. *PLoS ONE*. 2021;16(1):e0243211. doi: 10.1371/journal.pone.0243211. PMID: 33434197. Exclusion: Ineligible comparator.
305. Hermes SS, Rauen J, O'Brien S. Perceptions of school-based telehealth in a rural state: moving forward after COVID-19. *Int J Telerehabil*. 2021;13(1):e6370. doi: 10.5195/ijt.2021.6370. PMID: 34345344. Exclusion: Ineligible intervention.
306. Herrington G, Zardins Y, Hamilton A. A pilot trial of emergency telemedicine in regional Western Australia. *J Telemed Telecare*. 2013;19(7):430-3. doi: 10.1177/1357633X13506531. PMID: 24218359. Exclusion: Ineligible comparator.

307. Hersh WR, Wallace JA, Patterson PK, et al. Telemedicine for the Medicare Population-evidence reports/technology assessments, No. 24. Rockville, MD: July 2001. <http://www.ncbi.nlm.nih.gov/books/NBK33341/>. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
308. Hersh WR, Hickman D, Severance S, et al. Telemedicine for the Medicare population: update - evidence reports/technology assessments, No. 131. Rockville, MD.: Feb. 2006. <https://www.ncbi.nlm.nih.gov/books/NBK37953/>. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
309. Hicken BL, Daniel C, Luptak M, et al. Supporting Caregivers of Rural Veterans Electronically (SCORE). *J Rural Health*. 2017;33(3):305-13. doi: 10.1111/jrh.12195. PMID: 27437642. Exclusion: Ineligible intervention.
310. Higgins WJ, Luczynski KC, Carroll RA, et al. Evaluation of a telehealth training package to remotely train staff to conduct a preference assessment. *J Appl Behav Anal*. 2017;50(2):238-51. doi: 10.1002/jaba.370. PMID: 28090644. Exclusion: Ineligible intervention.
311. Hillman DG, Divers MC, McMurry TL, et al. Developing a telestroke state mapping tool to improve stroke outcomes in rural populations. *Stroke*. 2019;50doi: 10.1161/str.50.suppl_1.WP220. Exclusion: Ineligible intervention.
312. Hilty DM, Marks S, Wegeland J, et al. A randomized, controlled trial of disease management modules, including telepsychiatric care, for depression in rural primary care. *Psychiatry*. 2007;4(2):58-65. PMID: 20805900. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
313. Hinkle H. Beyond the disease symptoms: additional impacts of COVID-19 on rural health and health professions education: a reflection of where we have been and opportunities for the future. *Educ Health*. 2021;34(1):34-6. doi: 10.4103/efh.Efh_216_21. PMID: 34213442. Exclusion: Not a study.
314. Hirko KA, Kerver JM, Ford S, et al. Telehealth in response to the COVID-19 pandemic: implications for rural health disparities. *J Am Med Inform Assoc*. 2020;27:1816-18. doi: 10.1093/jamia/ocaa156. PMID: 32589735. Exclusion: Not a study.
315. Hitt WC, Low GM, Lynch CE, et al. Application of a telecolposcopy program in rural settings. *Telemed J E Health*. 2016;22(10):816-20. PMID: 27128600. Exclusion: Ineligible study design.
316. Hofstetter PJ, Kokesh J, Ferguson AS, et al. The impact of telehealth on wait time for ENT specialty care. *Telemed J E Health*. 2010;16(5):551-6. doi: 10.1089/tmj.2009.0142. PMID: 20575722. Exclusion: Ineligible intervention.
317. Hogan C, Muscat E, Hemington-Gorse S. Adherence to the national burns criteria in a tertiary burns centre in wales. *Br J Surg*. 2020;107(SUPPL 3):145. doi: 10.1002/bjs.11642. Exclusion: Not a study.
318. Holguin E, Stippler M, Yonas H. Teleradiology and online consultation: a solution of care for traumatic brain injury in rural New Mexico. *J Neurotrauma*. 2011;28(6):A97-A8. doi: 10.1089/neu.2011.9945. Exclusion: Ineligible study design.
319. Holmes AN, Chansky PB, Simpson CL. Teledermatology consultation can optimize treatment of cutaneous disease by nondermatologists in under-resourced clinics. *Telemed J E Health*. 2019;26(10):1284-90. doi: 10.1089/tmj.2019.0217. PMID: 31800369. Exclusion: Ineligible population.
320. Hougen HY, Lobo JM, Corey T, et al. Optimizing and validating the technical infrastructure of a novel tele-cystoscopy system. *J Telemed Telecare*. 2016;22(7):397-404. doi: 10.1177/1357633X15610040. PMID: 26489430. Exclusion: Ineligible population.

321. Howland C, Wakefield B. Assessing telehealth interventions for physical activity and sedentary behavior self-management in adults with type 2 diabetes mellitus: an integrative review. *Res Nurs Health*. 2021;44(1):92-110. doi: 10.1002/nur.22077. PMID: 33091168. Exclusion: Ineligible population.
322. Hubert GJ, Meretoja A, Audebert HJ, et al. Stroke thrombolysis in a centralized and a decentralized system (Helsinki and Telemedical Project for Integrative Stroke Care Network). *Stroke*. 2016;47(12):2999-3004. PMID: 27834751. Exclusion: Ineligible population.
323. Hughes AJ, Moloney DP, Fraser C, et al. Remote delivery of the satellite virtual fracture clinic - a pilot report of the first 500 cases. *Injury*. 2021;52(4):782-6. doi: 10.1016/j.injury.2020.11.055. PMID: 33257019. Exclusion: Ineligible population.
324. Huijbregts M, Taylor D, Cameron J, et al. Telehealth delivery of most, a stroke self-management program to remote areas in northern Ontario: results of the pilot evaluation. *Physiotherapy*. 2007;93(Suppl 1):S543. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
325. Hunt RC, Struminger BB, Redd JT, et al. Virtual peer-to-peer learning to enhance and accelerate the health system response to COVID-19: The HHS ASPR Project ECHO COVID-19 Clinical Rounds Initiative. *Ann Emerg Med*. 2021;78(2):223-8. doi: 10.1016/j.annemergmed.2021.03.035. PMID: 34325856. Exclusion: Ineligible population.
326. Hunter J, Scott RE. Considering the socio-cultural impact of e-health. *Stud Health Technol Inform*. 2010;161:77-85. PMID: 21191160. Exclusion: Ineligible setting.
327. Iacono T, Dissanayake C, Trembath D, et al. Family and practitioner perspectives on telehealth for services to young children with autism. *Stud Health Technol Inform*. 2016;231:63-73. PMID: 27782017. Exclusion: Ineligible intervention.
328. Iacono T, Stagg K, Pearce N, et al. A scoping review of Australian allied health research in ehealth. *BMC Health Serv Res*. 2016;16(1):543. PMID: 27716325. Exclusion: Ineligible intervention.
329. Ilko SA, Vakkalanka JP, Ahmed A, et al. Central venous access capability and critical care telemedicine decreases inter-hospital transfer among severe sepsis patients: a mixed methods design. *Crit Care Med*. 2019;47(5):659-67. doi: 10.1097/CCM.0000000000003686. PMID: 30730442. Exclusion: Ineligible population.
330. Imus TL, Northcutt T, Hall-Barrow J. Nurses play integral role in telestroke consultations in a rural state. *Stroke*. 2010;41(4):e395. doi: 10.1161/01.str.0000366116.47141.eb. Exclusion: Ineligible study design.
331. Iredale R, Hilgart JS, Hayward JA, et al. Telegenetics: a systematic review of telemedicine in genetics services. *J Cyber Ther Rehabil*. 2012;5(2):20-1. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
332. Irving M, Stewart R, Spallek H, et al. Using teledentistry in clinical practice as an enabler to improve access to clinical care: a qualitative systematic review. *J Telemed Telecare*. 2018;24(3):129-46. doi: 10.1177/1357633X16686776. PMID: 28092220. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
333. Iyer K, Nisenholtz M, Gutierrez D, et al. Disseminating knowledge in intestinal failure: initial report of the Learn Intestinal Failure Tele-ECHO (LIFT-ECHO) Project. *JPEN J Parenter Enteral Nutr*. 2021;03:03. doi: 10.1002/jpen.2078. PMID: 33533531. Exclusion: Ineligible population.
334. Izquierdo R, Lagua CT, Meyer S, et al. Telemedicine intervention effects on waist circumference and body mass index in the IDEATel project. *Diabetes Technol Ther*. 2010;12(3):213-20. doi: 10.1089/dia.2009.0102. PMID: 20151772. Exclusion: Ineligible intervention.

335. Jacob MK, Larson JC, Craighead WE. Establishing a telepsychiatry consultation practice in rural Georgia for primary care physicians: a feasibility report. *Clin Pediatr (Phila)*. 2012;51(11):1041-7. doi: 10.1177/0009922812441671. PMID: 22523276. Exclusion: Ineligible population.
336. Jacups SP, Kinchin I, Edwards L. Participatory action research applied to an ear, nose, and throat specialty service redesign in remote Australia: a mixed-methods study of key stakeholder perspectives. *Int J Environ Res Public Health*. 2020;18(1):29. doi: 10.3390/ijerph18010167. PMID: 33383696. Exclusion: Ineligible study design.
337. Jafari P, Kostas T, Levine S, et al. ECHO-Chicago geriatrics: using telementoring to "geriatricize" the primary care workforce. *Gerontol Geriatr Educ*. 2019;41(3):333-41. doi: 10.1080/02701960.2019.1572005. PMID: 30706769. Exclusion: Ineligible population.
338. Jagarapu J, Kapadia V, Mir I, et al. TeleNICU: extending the reach of level IV care and optimizing the triage of patient transfers. *J Telemed Telecare*. 2021;1357633X211038153. doi: 10.1177/1357633X211038153. PMID: 34524916. Exclusion: Ineligible comparator.
339. Jahun I, Dirlikov E, Odafe S, et al. Ensuring optimal community HIV testing services in Nigeria using an Enhanced Community Case-Finding Package (ECCP), October 2019-March 2020: acceleration to HIV epidemic control. *HIV AIDS (Auckl)*. 2021;13:839-50. doi: 10.2147/HIV.S316480. PMID: 34471388. Exclusion: Ineligible setting.
340. JaKa MM, Dinh JM, Ziegenfuss JY, et al. Patient and care team perspectives of telemedicine in critical access hospitals. *J Hosp Med*. 2020;15(6):345-8. doi: 10.12788/jhm.3412. PMID: 32490808. Exclusion: Ineligible study design.
341. James HM, Papoutsis C, Wherton J, et al. Spread, scale-up, and sustainability of video consulting in health care: systematic review and synthesis guided by the NASSS framework. *J Med Internet Res*. 2021;23(1):e23775. doi: 10.2196/23775. PMID: 33434141. Exclusion: Ineligible intervention.
342. Jang-Jaccard J, Nepal S, Alem L, et al. Barriers for delivering telehealth in rural Australia: a review based on Australian trials and studies. *Telemed J E Health*. 2014;20(5):496-504. doi: 10.1089/tmj.2013.0189. PMID: 24801522. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
343. Jani PD, Forbes L, Choudhury A, et al. Evaluation of diabetic retinal screening and factors for ophthalmology referral in a telemedicine network. *JAMA Ophthalmol*. 2017;135(7):706-14. doi: 10.1001/jamaophthalmol.2017.1150. PMID: 28520833. Exclusion: Ineligible intervention.
344. Jeerakathil T, Shuaib A, Fang S, et al. Thrombolysis using telehealth has comparable results to non-telehealth thrombolysis across northern Alberta: the Alberta Provincial Stroke Strategy (APSS). *Stroke*. 2012;43(2). Exclusion: Ineligible population.
345. Jew OS, Murthy AS, Danley K, et al. Implementation of a pediatric provider-to-provider store-and-forward teledermatology system: effectiveness, feasibility, and acceptability in a pilot study. *Pediatr Dermatol*. 2020;37(6):1106-12. doi: 10.1111/pde.14226. PMID: 32869352. Exclusion: Ineligible study design.
346. Jewett L, Mirian A, Connolly B, et al. Use of geospatial modeling to evaluate the impact of telestroke on access to stroke thrombolysis in Ontario. *J Stroke Cerebrovasc Dis*. 2017;26(7):1400-6. doi: 10.1016/j.jstrokecerebrovasdis.2017.03.023. PMID: 28478980. Exclusion: Ineligible outcome.

347. Jindal A, Nagarkar N, Thakur A, et al. A hybrid form of telemedicine and quality improvement: a unique way to extend intensive care services to neonates in medically underserved areas. *Pediatr Crit Care Med*. 2021;22(SUPPL 1):29. doi: 10.1097/01.pcc.0000738292.92665.69. Exclusion: Not a study.
348. Joerg L, Davoodi N, Goldberg EM. 65 interviews with emergency physicians on telehealth during COVID-19 and its role in caring for older Americans. *Ann Emerg Med*. 2021;78(2):S32. doi: 10.1016/j.annemergmed.2021.07.067. Exclusion: Ineligible intervention.
349. Johansson AM, Lindberg I, Soderberg S. Healthcare personnel's experiences using video consultation in primary healthcare in rural areas. *Prim Health Care Res Dev*. 2017;18(1):73-83. PMID: 27640522. Exclusion: Ineligible outcome.
350. Johansson T, Wild C. Telemedicine in acute stroke management: systematic review. *Int J Technol Assess Health Care*. 2010;26(2):149-55. doi: 10.1017/S0266462310000139. PMID: 20392317. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
351. Johnson D, Mauldin J, Chang E, et al. Maternal-fetal medicine coverage improved perinatal outcomes in a rural area. *Am J Obstet Gynecol*. 2014;210(1):S197. doi: 10.1016/j.ajog.2013.10.420. Exclusion: Ineligible intervention.
352. Johnson KL, Hertz D, Stobbe G, et al. Project Extension for Community Healthcare Outcomes (ECHO) in multiple sclerosis: increasing clinician capacity. *Int J MS Care*. 2017;19(6):283-9. doi: 10.7224/1537-2073.2016-099. PMID: 29270085. Exclusion: Ineligible population.
353. Johnston DL, Murto K, Kurzawa J, et al. Use of Electronic Consultation System to Improve Access to Care in Pediatric Hematology/Oncology. *J Pediatr Hematol Oncol*. 2017;39(7):e367-e9. doi: 10.1097/mp.0000000000000833. PMID: 28437292. Exclusion: Ineligible study design.
354. Johnston K, Smith D, Preston R, et al. "From the technology came the idea": safe implementation and operation of a high quality teleradiology model increasing access to timely breast cancer assessment services for women in rural Australia. *BMC Health Serv Res*. 2020;20(1):1103. doi: 10.1186/s12913-020-05922-y. PMID: 33256724. Exclusion: Ineligible study design.
355. Jones-Bamman C, Niermeyer S, McConnell K, et al. Teaching helping babies breathe via telehealth: a new application in rural Guatemala. *Biomed Hub*. 2019;4(3):1-6. doi: 10.1159/000502934. PMID: 31993430. Exclusion: Ineligible setting.
356. Jones MS, Goley AL, Alexander BE, et al. Inpatient transition to virtual care during COVID-19 pandemic. *Diabetes Technol Ther*. 2020;22(6):444-8. doi: 10.1089/dia.2020.0206. PMID: 32396395. Exclusion: Ineligible population.
357. Jonk YC, Burgess A, Williamson ME, et al. Telehealth use in a rural state: a mixed-methods study using Maine's All-Payer Claims Database. *J Rural Health*. 2020;21:21. doi: 10.1111/jrh.12527. PMID: 33085154. Exclusion: Ineligible intervention.
358. Jordan LA, Sewell C, Rudd J, et al. Rural tele-thrombolysis protocol implementation: nurses leading change. *Int J Stroke*. 2014;9:5-6. doi: 10.1111/ij.12334. Exclusion: Not a study.
359. Joshi S, Gali K, Radecki L, et al. Integrating quality improvement into the ECHO model to improve care for children and youth with epilepsy. *Epilepsia*. 2020;61(9):1999-2009. doi: 10.1111/epi.16625. PMID: 32860215. Exclusion: Ineligible population.

360. Julien HM, Eberly LA, Adusumalli S. Telemedicine and the forgotten America. *Circulation*. 2020;142(4):312-4. doi: 10.1161/CIRCULATIONAHA.120.048535 . PMID: 32525712. Exclusion: Not a study.
361. Jumah NA, Tyler L, Turuba R, et al. On the path to reclaiming Indigenous midwifery: co-creating the Maternal Infant Support Worker pilot program. *Int J Gynaecol Obstet*. 2021;07:07. doi: 10.1002/ijgo.13918. PMID: 34491574. Exclusion: Ineligible comparator.
362. Kageji T, Obata F, Oka H, et al. Drip-and-ship thrombolytic therapy supported by the telestroke system for acute ischemic stroke patients living in medically under-served areas. *Neurol Med Chir (Tokyo)*. 2016;56(12):753-8. PMID: 27333939. Exclusion: Ineligible comparator.
363. Kalichman SC, Katner H, Eaton LA, et al. Comparative effects of telephone versus in-office behavioral counseling to improve HIV treatment outcomes among people living with HIV in a rural setting. *Transl Behav Med*. 2021;11(3):852-62. doi: 10.1093/tbm/ibaa109. PMID: 33200772. Exclusion: Ineligible intervention.
364. Kamdar NV, Huverserian A, Jalilian L, et al. Development, implementation, and evaluation of a telemedicine preoperative evaluation initiative at a major academic medical center. *Anesth Analg*. 2020;131(6):1647-56. doi: 10.1213/ANE.0000000000005208. PMID: 32841990. Exclusion: Ineligible population.
365. Kanagasingam Y, Bhuiyan A, Abramoff MD, et al. Progress on retinal image analysis for age related macular degeneration. *Prog Retin Eye Res*. 2014;38:20-42. doi: 10.1016/j.preteyeres.2013.10.002. PMID: 24211245. Exclusion: Not a study.
366. Kanagasingam Y. Sight-saving science: specialist eye care using satellite technology. *Med J Aust*. 2015;203(4):164. PMID: 26268277. Exclusion: Ineligible study design.
367. Kane B, Groth K, Randall D. Medical team meetings: utilising technology to enhance communication, collaboration and decision-making. *Behav Inf Technol*. 2011;30(4):437-42. doi: 10.1080/0144929X.2011.591576. Exclusion: Not a study.
368. Karp JF, Hamm ME, Abril F, et al. Improving effective mental health consultation for rural older adults living with depression and pain: learning from the experiences of rural primary care physicians. *Prim Care Companion CNS Disord*. 2021;23(2):18. doi: 10.4088/PCC.20m02725. PMID: 34000175. Exclusion: Irritable
369. Katzman J, Tomedi L, Swift R, et al. Extension for Community Healthcare Outcomes (ECHO) telementoring in the military: where we are now, opportunities and challenges. *Mil Med*. 2021;15:15. doi: 10.1093/milmed/usab010. PMID: 33585892. Exclusion: Ineligible population.
370. Katzman JG, Comerchi G, Jr., Boyle JF, et al. Innovative telementoring for pain management: Project ECHO pain. *J Contin Educ Health Prof*. 2014;34(1):68-75. doi: 10.1002/chp.21210. PMID: 24648365. Exclusion: Ineligible population.
371. Katzman JG, Galloway K, Olivas C, et al. Expanding health care access through education: dissemination and implementation of the ECHO model. *Mil Med*. 2016;181(3):227-35. doi: 10.7205/MILMED-D-15-00044. PMID: 26926747. Exclusion: Ineligible population.
372. Katzman JG, Qualls CR, Satterfield WA, et al. Army and Navy ECHO pain telementoring improves clinician opioid prescribing for military patients: an observational cohort study. *J Gen Intern Med*. 2019;34(3):387-95. doi: 10.1007/s11606-018-4710-5. PMID: 30382471. Exclusion: Ineligible population.

373. Katzman JG, Tomedi LE, Thornton K, et al. Innovative COVID-19 programs to rapidly serve New Mexico: Project ECHO. *Public Health Rep.* 2021;136(1):39-46. doi: 10.1177/0033354920969180. PMID: 33216679. Exclusion: Ineligible outcome.
374. Kaufman T, Geraghty EM, Dullet N, et al. Geospatial Information System analysis of healthcare need and telemedicine delivery in California. *Telemed J E Health.* 2017;23(5):430-4. doi: 10.1089/tmj.2016.0144. PMID: 27835073. Exclusion: Ineligible intervention.
375. Kauth MR, Shipherd JC, Lindsay JA, et al. Teleconsultation and training of VHA providers on transgender care: implementation of a multisite hub system. *Telemed J E Health.* 2015;21(12):1012-8. doi: 10.1089/tmj.2015.0010. PMID: 26171641. Exclusion: Ineligible population.
376. Kawasaki S, Francis E, Mills S, et al. Multi-model implementation of evidence-based care in the treatment of opioid use disorder in Pennsylvania. *J Subst Abuse Treat.* 2019;106:58-64. doi: 10.1016/j.jsat.2019.08.016. PMID: 31540612. Exclusion: Ineligible population.
377. Kazley AS, Wilkerson RC, Jauch E, et al. Access to expert stroke care with telemedicine: REACH MUSC. *Front Neurol.* 2012;3(44):1-6. doi: 10.3389/fneur.2012.00044. PMID: 22461780. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
378. Kenyon JI, Poropatich R, Holtel MR. Cell phones in telehealth and otolaryngology. *Otolaryngol Clin North Am.* 2011;44(6):1351-8, ix. doi: 10.1016/j.otc.2011.08.013. PMID: 22032487. Exclusion: Not a study.
379. Keogh L, Croy S, Newton D, et al. General practitioner knowledge and practice in relation to unintended pregnancy in the Grampians region of Victoria, Australia. *Rural Remote Health.* 2019;19(4):5156. doi: 10.22605/RRH5156. PMID: 31675244. Exclusion: Ineligible intervention.
380. Keough GJ, Matherly AF, Cochran A. Expanding burn care knowledge in a rural region through Project Extension for Community Healthcare Outcomes. *JAMA Surg.* 2017;152(4):401-2. doi: 10.1001/jamasurg.2016.5747. PMID: 28241223. Exclusion: Not a study.
381. Khader YS, Jarrah MI, Al-Shudifat AEM, et al. Telecardiology application in Jordan: Its impact on diagnosis and disease management, patients' quality of life, and time- and cost-savings. *Int J Telemed Appl.* 2014;2014doi: 10.1155/2014/819837. PMID: 25400661. Exclusion: Ineligible intervention.
382. Khan FB, Hall RE, Stamplecoski M, et al. Acute stroke care: Telestroke vs. on-site stroke specialist care. Results from the Ontario telestroke experience. *Stroke.* 2011;42(3):e340. doi: 10.1161/STR.0b013e3182074d9b. Exclusion: Not a study.
383. Khan K, Mathur A, Kaur S, et al. Profile of cases discussed in innovative tele-ECHO mentoring program on addiction management. *Asian J Psychiatr.* 2020;52:102060. doi: 10.1016/j.ajp.2020.102060. PMID: 32339918. Exclusion: Ineligible comparator.
384. Khan S, Usmani A. Remote patient monitoring system with a focus on antenatal care for rural population. *BJOG.* 2014;121:149-50. doi: 10.1111/1471-0528.12787. Exclusion: Ineligible setting.
385. Khatri K, Haddad M, Anderson D. Project ECHO: replicating a novel model to enhance access to hepatitis C care in a community health center. *J Health Care Poor Underserved.* 2013;24(2):850-8. doi: 10.1353/hpu.2013.0093. PMID: 23728050. Exclusion: Ineligible population.
386. Khunlertkit A, Carayon P. Contributions of tele-intensive care unit (tele-ICU) technology to quality of care and patient safety. *J Crit Care.* 2013;28(3):315.e1-12. doi: 10.1016/j.jcrc.2012.10.005. PMID: 23159139. Exclusion: Ineligible population.

387. Kim YK, Kim KY, Lee KH, et al. Clinical outcomes on real-time telemetry system in developing emergency medical service system. *Telemed J E Health*. 2011;17(4):247-53. doi: 10.1089/tmj.2010.0152. PMID: 21480786. Exclusion: Ineligible population.
388. Kingue S, Angandji P, Menanga AP, et al. Efficiency of an intervention package for arterial hypertension comprising telemanagement in a Cameroonian rural setting: the TELEMED-CAM study. *Pan Afr Med J*. 2013;15:153. doi: 10.11604/pamj.2013.15.153.2655. PMID: 24396559. Exclusion: Ineligible setting.
389. Kipnes M, Haffner S, Lightner T, et al. Virtual endocrinologist: hospital diabetes management via telemedicine. *Diabetes*. 2016;65:A328. doi: 10.2337/db16-861-1374. Exclusion: Ineligible population.
390. Kirshenbaum E, Rhee E, Makarov D, et al. Current state of teleurology: prevalence, barriers and reimbursement. *J Urol*. 2020;203:e18-e9. doi: 10.1097/JU.0000000000000816.017. Exclusion: Not a study.
391. Kjelle E, Myklebust AM. Telemedicine remote controlled stroke evaluation and treatment, the experience of radiographers, paramedics and junior doctors in a novel rural stroke management team. *BMC Health Serv Res*. 2021;21(1):554. doi: 10.1186/s12913-021-06591-1. PMID: 34090447. Exclusion: Ineligible comparator.
392. Klein Y, Donchik V, Jaffe D, et al. Management of patients with traumatic intracranial injury in hospitals without neurosurgical service. *J Trauma*. 2010;69(3):544-8. doi: 10.1097/TA.0b013e3181c99936. PMID: 20234328. Exclusion: Ineligible population.
393. Kluesner KM, Wibbenmeyer L, Wu H, et al. Video assisted burn evaluation is a reliable tool for improving initial evaluation and treatment of burn injuries in rural areas. *J Burn Care Res*. 2014;35:S185. doi: 10.1097/01.bcr.0000445189.61189.9d. Exclusion: Not a study.
394. Knapp H, Chan K, Anaya HD, et al. Interactive internet-based clinical education: an efficient and cost-savings approach to point-of-care test training. *Telemed J E Health*. 2011;17(5):335-40. doi: 10.1089/tmj.2010.0187. PMID: 21492031. Exclusion: Ineligible population.
395. Kocanda L, Fisher K, Brown LJ, et al. Informing telehealth service delivery for cardiovascular disease management: exploring the perceptions of rural health professionals. *Aust Health Rev*. 2021;45(2):241-6. doi: 10.1071/AH19231. PMID: 33715764. Exclusion: Ineligible study design.
396. Komaromy M, Ceballos V, Zurawski A, et al. Extension for Community Healthcare Outcomes (ECHO): a new model for community health worker training and support. *J Public Health Policy*. 2018;39(2):203-16. doi: 10.1057/s41271-017-0114-8. PMID: 29263437. Exclusion: Not a study.
397. Konjeti VR, Heuman D, Bajaj JS, et al. Telehealth-based evaluation identifies patients who are not candidates for liver transplantation. *Clin Gastroenterol Hepatol*. 2019;17(1):207-9. doi: 10.1016/j.cgh.2018.04.048. PMID: 29723691. Exclusion: Ineligible population.
398. Kopelovich S, Monroe-DeVita M, Hughes M, et al. Adopting technology and a stepped care approach to advance the implementation of cognitive behavioral therapy for psychosis (CBTP) in community mental health settings: introducing the CBTP Project ECHO clinic. *Schizophr Bull*. 2017;43:S209-S10. Exclusion: Not a study.
399. Korczyk D, McPherson B, Hickey A, et al. Telemedicine in heart failure: heart failure service support for south west Queensland. *Eur J Heart Fail, Supplement*. 2010;9:S221. doi: 10.1093/eurjhf/hsq013. Exclusion: Ineligible study design.

400. Kraus F, Hubert GJ, Haberl RL. "Flying interventionist"-a novel concept for stroke care in rural areas. *Eur Stroke J*. 2017;2(1):310. doi: 10.1177/2396987317705242. Exclusion: Ineligible study design.
401. Kroenke K, Theobald D, Norton K, et al. The Indiana Cancer Pain and Depression (INCPAD) trial design of a telecare management intervention for cancer-related symptoms and baseline characteristics of study participants. *Gen Hosp Psychiatry*. 2009;31(3):240-53. PMID: 19410103. Exclusion: Ineligible intervention.
402. Kroenke K, Theobald D, Wu J, et al. The association of depression and pain with health-related quality of life, disability, and health care use in cancer patients. *J Pain Symptom Manage*. 2010;40(3):327-41. doi: 10.1016/j.jpainsymman.2009.12.023. PMID: 20580201. Exclusion: Ineligible intervention.
403. Kruger C, Niemi M. A telemedicine network to support paediatric care in small hospitals in rural Tanzania. *J Telemed Telecare*. 2012;18(1):59-62. doi: 10.1258/jtt.2011.110312. PMID: 21968000. Exclusion: Ineligible setting.
404. Krum H, Forbes A, Yallop J, et al. Telephone support to rural and remote patients with heart failure: the Chronic Heart Failure Assessment by Telephone (CHAT) study. *Cardiovasc Ther*. 2013;31(4):230-7. doi: 10.1111/1755-5922.12009. PMID: 23061492. Exclusion: Ineligible intervention.
405. Kruse CS, Bouffard S, Dougherty M, et al. Telemedicine use in rural Native American communities in the era of the ACA: a systematic literature review. *J Med Syst*. 2016;40(6):145. doi: 10.1007/s10916-016-0503-8. PMID: 27118011. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
406. Kubat B. Providing rural care, remotely: Avera's eCare initiative. *Caring for the Ages*. 2015;16(7):16. Exclusion: Not a study.
407. Kueider S, Eastwood JS. The use of teleneurology consult for stroke care in a rural community hospital. *Stroke*. 2012;43(2). Exclusion: Ineligible study design.
408. Kulcsar Z, Albert DA, Merrihew K, et al. Tele-rheumatology: despite improved access could there be a potential delay in care without a skilled "presenter"? *Arthritis Rheumatol*. 2014;66:S1008-S9. doi: 10.1002/art.38914. Exclusion: Ineligible intervention.
409. Kulcsar Z, Albert D, Ercolano E, et al. Telerheumatology: a technology appropriate for virtually all. *Semin Arthritis Rheum*. 2016;46(3):380-5. doi: 10.1016/j.semarthrit.2016.05.013. PMID: 27395561. Exclusion: Ineligible setting.
410. Kumar M, Agarwal A, Pant P, et al. Feasibility and accuracy of tele-spirometry: a community based approach. *Am J Respir Crit Care Med*. 2019;199(9). Exclusion: Ineligible intervention.
411. Kumar M, Agarwal A, Singh M, et al. Feasibility and accuracy of Tele-spirometry: a community based approach. *World Allergy Organ J*. 2020;13(8)doi: 10.1016/j.waojou.2020.100151. Exclusion: Not a study.
412. Kunze MG, Machalicek W, Wei Q, et al. Coaching via telehealth: caregiver-mediated interventions for young children on the waitlist for an autism diagnosis using single-case design. *J Clin Med*. 2021;10(8):13. doi: 10.3390/jcm10081654. PMID: 33924440. Exclusion: Ineligible intervention.
413. Laloo C, Diskin C, Ho M, et al. Pediatric Project ECHO: implementation of a virtual medical education program to support community management of children with medical complexity. *Hosp Pediatr*. 2020;10(12):1044-52. doi: 10.1542/hpeds.2020-0067. PMID: 33139305. Exclusion: Ineligible population.

414. Lalloo C, Osei-Twum JA, Rapoport A, et al. Pediatric Project ECHO R: a virtual community of practice to improve palliative care knowledge and self-efficacy among interprofessional health care providers. *J Palliat Med.* 2021;24(7):1036-44. doi: 10.1089/jpm.2020.0496. PMID: 33326309. Exclusion: Ineligible population.
415. Lama T, Karmacharya B, Chandler C, et al. Telephone management of severe wasp stings in rural Nepal: a case report. *J Telemed Telecare.* 2011;17(2):105-8. doi: 10.1258/jtt.2010.100606. PMID: 21139015. Exclusion: Ineligible setting.
416. Lambert D, Gale J, Hartley D, et al. Understanding the business case for telemental health in rural communities. *J Behav Health Serv Res.* 2016;43(3):366-79. doi: 10.1007/s11414-015-9490-7. PMID: 26695645. Exclusion: Ineligible intervention.
417. Landow SM, Oh DH, Weinstock MA. Tele dermatology within the Veterans Health Administration, 2002-2014. *Telemed J E Health.* 2015;21(10):769-73. doi: 10.1089/tmj.2014.0225. PMID: 26083585. Exclusion: Ineligible outcome.
418. Langabeer JR, 2nd, Gonzalez M, Alqusairi D, et al. Telehealth-Enabled Emergency Medical Services Program Reduces Ambulance Transport to Urban Emergency Departments. *West J Emerg Med.* 2016;17(6):713-20. doi: 10.5811/westjem.2016.8.30660. PMID: 27833678. Exclusion: Ineligible population.
419. Langabeer JR, 2nd, Champagne-Langabeer T, Alqusairi D, et al. Cost-benefit analysis of telehealth in pre-hospital care. *J Telemed Telecare.* 2017;23(8):747-51. doi: 10.1177/1357633X16680541. PMID: 27913657. Exclusion: Ineligible population.
420. Lapointe L, Lavallee-Bourget MH, Pichard-Jolicoeur A, et al. Impact of telemedicine on diagnosis, clinical management and outcomes in rural trauma patients: a rapid review. *Can J Rural Med.* 2020;25(1):31-40. doi: 10.4103/CJRM.CJRM_8_19. PMID: 31854340. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
421. Laver KE, Adey-Wakeling Z, Crotty M, et al. Telerehabilitation services for stroke. *Cochrane Database Syst Rev.* 2020 (1)doi: 10.1002/14651858.CD010255.pub3. PMID: 32002991. Exclusion: Ineligible intervention.
422. Lazarus G, Kirchner HL, Siswanto BB. Prehospital tele-electrocardiographic triage improves the management of acute coronary syndrome in rural populations: a systematic review and meta-analysis. *J Telemed Telecare.* 2020;1357633X20960627. doi: 10.1177/1357633X20960627. PMID: 32996348. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
423. Lazarus G, Permana AP, Nugroho SW, et al. Telestroke strategies to enhance acute stroke management in rural settings: a systematic review and meta-analysis. *Brain Behav.* 2020:e01787. doi: 10.1002/brb3.1787. PMID: 32812380. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
424. Le LD, Paulk IR, Axon DR, et al. Comprehensive medication review completion in medically underserved areas and populations. *J Health Care Poor Underserved.* 2021;32(3):1301-11. doi: 10.1353/hpu.2021.0133. PMID: 34421032. Exclusion: Ineligible intervention.
425. Lea JP, Masud T, Someren JT, et al. In-hospital hemodialysis services provided by tele-nephrologists using the Nxstage system one s. *J Am Soc Nephrol.* 2018;29:628. Exclusion: Not a study.

426. LeBlanc M, Petrie S, Paskaran S, et al. Patient and provider perspectives on eHealth interventions in Canada and Australia: a scoping review. *Rural Remote Health*. 2020;20(3):5754. doi: 10.22605/RRH5754. PMID: 32949485. Exclusion: Ineligible population.
427. Lee CH, Huang CC, Huang JT, et al. Live-interactive teledermatology program in Taiwan: one-year experience serving a district hospital in rural Taitung County. *J Formos Med Assoc*. 2021;120:422-8. doi: 10.1016/j.jfma.2020.06.007. PMID: 32563581. Exclusion: Ineligible study design.
428. Lee RH, Lyles KW, Pearson M, et al. Osteoporosis screening and treatment among veterans with recent fracture after implementation of an electronic consult service. *Calcif Tissue Int*. 2014;94(6):659-64. doi: 10.1007/s00223-014-9849-4. PMID: 24699797. Exclusion: Ineligible population.
429. Lee RH, Pearson M, Lyles KW, et al. Geographic scope and accessibility of a centralized, electronic consult program for patients with recent fracture. *Rural Remote Health*. 2016;16(1):3440. PMID: 26745338. Exclusion: Ineligible outcome.
430. Lee SJ, Jung TY, Lee TR, et al. Accepting telemedicine in a circulatory medicine ward in major hospitals in South Korea: patients' and health professionals' perception of real-time electrocardiogram monitoring. *BMC Health Serv Res*. 2018;18(1):293. doi: 10.1186/s12913-018-3105-y. PMID: 29678189. Exclusion: Ineligible intervention.
431. Lee SPH, Cho-Lim JJ, Lee BKY, et al. Improvement of post-radiation therapy cancer care for veteran patients living in a rural area. *Int J Radiat Oncol Biol Phys*. 2017;99(2):E532. doi: 10.1016/j.ijrobp.2017.06.1878. Exclusion: Ineligible study design.
432. Lese A, Sraj S. Rural orthopedics: providing orthopedic care in rural communities. *Orthopedics*. 2019;42(4):e350-e5. doi: 10.3928/01477447-20190624-01. PMID: 31323106. Exclusion: Not a study.
433. Levinson CA, Spoor SP, Keshishian AC, et al. Pilot outcomes from a multidisciplinary telehealth versus in-person intensive outpatient program for eating disorders during versus before the Covid-19 pandemic. *Int J Eat Disord*. 2021;54(9):1672-9. doi: 10.1002/eat.23579. PMID: 34245028. Exclusion: Ineligible population.
434. Levisohn P, Wagner J, Falcone T, et al. Improving mental health services for children and youth with epilepsy: practical solutions from project access. *Epilepsy Curr*. 2014;14:152. Exclusion: Not a study.
435. Lewiecki EM, Boyle JF, Arora S, et al. Telementoring: a novel approach to reducing the osteoporosis treatment gap. *Osteoporos Int*. 2017;28(1):407-11. doi: 10.1007/s00198-016-3708-2. PMID: 27439373. Exclusion: Ineligible study design.
436. Lewiecki EM, Rochelle R, Bouchonville MF, 2nd, et al. Leveraging scarce resources with bone health TeleECHO to improve the care of osteoporosis. *J Endocr Soc*. 2017;1(12):1428-34. doi: 10.1210/js.2017-00361. PMID: 29264466. Exclusion: Ineligible population.
437. Lewiecki EM, Rochelle R. Project ECHO: telehealth to expand capacity to deliver best practice medical care. *Rheum Dis Clin North Am*. 2019;45(2):303-14. doi: 10.1016/j.rdc.2019.01.003. PMID: 30952400. Exclusion: Ineligible population.
438. Lewiecki EM, Rothman MS. COVID-19, medical education, and bone health: insights from Project ECHO. *J Clin Densitom*. 2020;23(3):338-9. doi: 10.1016/j.jocd.2020.04.003. PMID: 32448715. Exclusion: Not a study.
439. Lewinski AA, Crowley MJ, Miller C, et al. Applied rapid qualitative analysis to develop a contextually appropriate intervention and increase the likelihood of uptake. *Med Care*. 2021;59(Suppl 3):S242-S51. doi: 10.1097/MLR.0000000000001553. PMID: 33976073. Exclusion: Ineligible intervention.

440. Li C, Ong C, Morris A, et al. Evaluating the appropriateness of antibiotic treatment of tonsillitis during COVID-19 in the North Wale primary healthcare setting. *J Prim Care Community Health*. 2021;1-6. doi: 10.1177/21501327211003687. PMID: 33733905. Exclusion: Ineligible intervention.
441. Li J, Green DL, Santilli S, et al. Televascular consultation is one answer to rural vascular surgery shortage. *J Telemed Telecare*. 2021;1357633X211003456. doi: 10.1177/1357633X211003456. PMID: 33840280. Exclusion: Ineligible intervention.
442. Liddy C, Smyth C, Poulin PA, et al. Improving access to chronic pain services through eConsultation: a cross-sectional study of the Champlain BASE eConsult service. *Pain Med*. 2016;17(6):1049-57. doi: 10.1093/pm/pnw038. PMID: 27040667. Exclusion: Ineligible study design.
443. Liddy C, McKellips F, Armstrong CD, et al. Improving access to specialists in remote communities: a cross-sectional study and cost analysis of the use of eConsult in Nunavut. *Int J Circumpolar Health*. 2017;76(1):1323493. doi: 10.1080/22423982.2017.1323493. PMID: 28570207. Exclusion: Ineligible study design.
444. Liddy C, Moroz I, Mihan A, et al. A systematic review of asynchronous, provider-to-provider, electronic consultation services to improve access to specialty care available worldwide. *Telemed J E Health*. 2019;25(3):184-98. doi: 10.1089/tmj.2018.0005. PMID: 29927711. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
445. Lim D, Oakley AM, Rademaker M. Better, sooner, more convenient: a successful teledermoscopy service. *Australas J Dermatol*. 2012;53(1):22-5. doi: 10.1111/j.1440-0960.2011.00836.x. PMID: 22309326. Exclusion: Ineligible population.
446. Lin CC, Dievler A, Robbins C, et al. Telehealth in health centers: key adoption factors, barriers, and opportunities. *Health Aff (Millwood)*. 2018;37(12):1967-74. doi: 10.1377/hlthaff.2018.05125. PMID: 30633683. Exclusion: Ineligible intervention.
447. Lindauer A, Croff R, Mincks K, et al. "It took the stress out of getting help": the STAR-C-telemedicine mixed methods pilot. *Care Wkly*. 2018;2:7-14. doi: 10.14283/cw.2018.4. PMID: 30393783. Exclusion: Ineligible population.
448. Linderman DJ, Koff PB, Min S, et al. Proactive integrated care improves evidence-based care and outcomes for COPD patients in rural areas. *Am J Respir Crit Care Med*. 2010;181(1). Exclusion: Ineligible comparator.
449. Lippman JM, Smith SN, McMurphy TL, et al. Mobile telestroke during ambulance transport is feasible in a rural EMS setting: the iTREAT study. *Telemed J E Health*. 2016;22(6):507-13. doi: 10.1089/tmj.2015.0155. PMID: 26600433. Exclusion: Ineligible outcome.
450. Lobo R, B DC, Forbes L, et al. Young Deadly Free: impact evaluation of a sexual health youth peer education program in remote Australian communities. *Sex Health*. 2020;17(5):397-404. doi: 10.1071/SH20069. PMID: 33176903. Exclusion: Ineligible intervention.
451. Loi SM. Dementia behavior management advisory service northern territory pilot program: reaching out to rural communities using technology. *Asia Pac Psychiatry*. 2014;6(2):235-6. doi: 10.1111/appy.12132. PMID: 24753367. Exclusion: Not a study.
452. Lomenick AF, Kuhlman SJ, Barnes JL, et al. Economics of using telemedicine to supplement hospice care in rural areas. *J Palliat Med*. 2021;01:01. doi: 10.1089/jpm.2020.0117. PMID: 33794099. Exclusion: Ineligible intervention.

453. Long LA, Pariyo G, Kallander K. Digital technologies for health workforce development in low- and middle-income countries: a scoping review. *Glob Health Sci Pract.* 2018;6(Suppl 1):S41-S8. doi: 10.9745/GHSP-D-18-00167. PMID: 30305338. Exclusion: Ineligible population.
454. Lopez MS, Baker ES, Milbourne AM, et al. Project ECHO: a telementoring program for cervical cancer prevention and treatment in low-resource settings. *J Glob Oncol.* 2017;3(5):658-65. doi: 10.1200/JGO.2016.005504. PMID: 29094102. Exclusion: Not a study.
455. Lopez Segui F, Franch Parella J, Girones Garcia X, et al. A cost-minimization analysis of a medical record-based, store and forward and provider-to-provider telemedicine compared to usual care in Catalonia: more agile and efficient, especially for users. *Int J Environ Res Public Health.* 2020;17:1-9. doi: 10.3390/ijerph17062008. PMID: 32197434. Exclusion: Ineligible population.
456. Love HE, Schlitt J, Soleimanpour S, et al. Twenty years of school-based health care growth and expansion. *Health Aff (Millwood).* 2019;38(5):755-64. doi: 10.1377/hlthaff.2018.05472. PMID: 31059359. Exclusion: Ineligible intervention.
457. Lowery CL, Bronstein JM, Benton TL, et al. Distributing medical expertise: the evolution and impact of telemedicine in arkansas. *Health Aff (Millwood).* 2014;33(2):235-43. doi: 10.1377/hlthaff.2013.1001. PMID: 24493766. Exclusion: Not a study.
458. Lu AD, Gunzburger E, Glorioso TJ, et al. Impact of longitudinal virtual primary care on diabetes quality of care. *J Gen Intern Med.* 2021;22:22. doi: 10.1007/s11606-020-06547-x. PMID: 33483815. Exclusion: Ineligible intervention.
459. Lunney M, Lee R, Tang K, et al. Impact of telehealth interventions on processes and quality of care for patients with ESRD. *Am J Kidney Dis.* 2018;72(4):592-600. doi: 10.1053/j.ajkd.2018.02.353. PMID: 29699884. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
460. Lustig T. The role of telehealth in an evolving health care environment - workshop summary. Washington DC: 2012. doi: 10.17226/13466. PMID: 24901186. Exclusion: Not a study. Wide ranging conference proceedings; useful background info and potentially KQ2 or Use for background
461. Ly BA, Bourgeault IL, Labonte R, et al. Physicians' perceptions on the impact of telemedicine on recruitment and retention in underserved areas: a descriptive study in Senegal. *Hum Resour Health.* 2017;15(67):1-8. doi: 10.1186/s12960-017-0242-z. PMID: 28923056. Exclusion: Ineligible setting.
462. Lyles A. Rural stroke patients are getting more neurology consults, but mortality outcomes have not improved over time. *Neurology Today.* 2020;20(12):17-22. doi: 10.1097/01.nt.0000682196.80970.4b. Exclusion: Not a study.
463. MacDonald ME, Diallo GS. Socio-cultural contextual factors that contribute to the uptake of a mobile health intervention to enhance maternal health care in rural Senegal. *Reprod Health.* 2019;16(1):141. doi: 10.1186/s12978-019-0800-z. PMID: 31511028. Exclusion: Ineligible setting.
464. Machado SM, Wilson EH, Elliott JO, et al. Impact of a telemedicine eICU cart on sepsis management in a community hospital emergency department. *J Telemed Telecare.* 2018;24(3):202-8. doi: 10.1177/1357633X17691862. PMID: 29278979. Exclusion: Ineligible population.

465. Maddox LJ, Albritton J, Morse J, et al. Implementation and outcomes of a telehealth neonatology program in a single healthcare system. *Front Pediatr*. 2021;9:648536. doi: 10.3389/fped.2021.648536. PMID: 33968852. Exclusion: Ineligible setting.
466. Maddry JK, Sessions D, Heard K, et al. Wartime toxicology: evaluation of a military medical toxicology telemedicine consults service to assist physicians serving overseas and in combat (2005-2012). *J Med Toxicol*. 2014;10(3):261-5. doi: 10.1007/s13181-014-0398-z. PMID: 24752493. Exclusion: Ineligible population.
467. Madu PN, Chang AY, Kayembe MK, et al. Teledermatology as a means to provide multispecialty care: a case of global specialty collaboration. *Pediatr Dermatol*. 2017;34(2):e89-e92. doi: 10.1111/pde.13079. PMID: 28297155. Exclusion: Ineligible study design.
468. Mahadevan J, Shukla L, Chand PK, et al. Innovative virtual mentoring using the Extension for Community Healthcare Outcomes model for primary care providers for the management of alcohol use disorders. *Indian J Med Res*. 2020;151(6):609-12. doi: 10.4103/ijmr.IJMR_1851_18. PMID: 32719236. Exclusion: Ineligible intervention.
469. Mahmood F, Cyr J, Keely E, et al. Teledermatology utilization and integration in residency training over the COVID-19 pandemic. *J Cutan Med Surg*. 2021;12034754211045393. doi: 10.1177/12034754211045393. PMID: 34551623. Exclusion: Ineligible setting.
470. Main D, Dziekan K. Distance education: linking traditional classroom rehabilitation counseling students with their colleagues using hybrid learning models. *Rehabil Res Pract*. 2012;26(4):315-20. doi: 10.1891/2168-6653.26.4.6. Exclusion: Ineligible population.
471. Makkar A, Foulks A, McCoy M, et al. Evaluation of neonatal services provided in a level II NICU utilizing hybrid telemedicine. *J Investig Med*. 2018;66(2):521. doi: 10.1136/jim-2017-000697.417. Exclusion: Not a study.
472. Malavera A, Delcourt C, Paul C, et al. Strategies for improving access to endovascular stroke therapy for rural populations: systematic review and on-going clinical trial. *Int J Stroke*. 2020;15(1 SUPPL):645. doi: 10.1177/1747493020963387. Exclusion: Not a study.
473. Malik MT, Zand R, Vezendy SM, et al. Implementation and follow-up experience of an e-consult program in a rural neurology setting. *Neurol Clin Pract*. 2021;11(4):e430-e7. doi: 10.1212/CPJ.0000000000000993. PMID: 34484941. Exclusion: Ineligible study design.
474. Mamadnabiev I, Imfeld S, Gurke L, et al. A telemedicine communication exchange network in Gorno-Badakhshan Autonomic Oblast in collaboration with Swiss Surgical Teams. *J Telemed Telecare*. 2020;1357633X20948989. doi: 10.1177/1357633X20948989. PMID: 32830613. Exclusion: Ineligible setting.
475. Mandall NA, O'Brien KD, Brady J, et al. Teledentistry for screening new patient orthodontic referrals. Part 1: A randomised controlled trial. *Br Dent J*. 2005;199(10):659-62, discussion 3. doi: 10.1038/sj.bdj.4812930. PMID: 16311569. Exclusion: Ineligible population.
476. Manson J, Ghasemi L, Westerdale E, et al. Evaluating a palliative care education programme for domiciliary care workers. *Nurs Older People*. 2020;35(2):1-11. doi: 10.7748/nop.2020.e1235. PMID: 32347075. Exclusion: Ineligible study design.
477. Manson J, Gardiner C, Taylor P, et al. 'Palliative care education in nursing homes: a qualitative evaluation of telementoring. *BMJ Support Palliat Care*. 2021;24:24. doi: 10.1136/bmjspcare-2020-002727. PMID: 33627368. Exclusion: Ineligible population.

478. Manuel P. A prospective, interventional study of the effectiveness of digital wound imaging, remote consultation and podiatry offloading devices on the healing rates of chronic lower extremity wounds in remote regions of Western Australia. *Wound Practice & Research*. 2012;20(2):103-9. Exclusion: Ineligible study design.
479. Marciano S, Haddad L, Plazzotta F, et al. Implementation of the ECHO R telementoring model for the treatment of patients with hepatitis C. *J Med Virol*. 2017;89(4):660-4. doi: 10.1002/jmv.24668. PMID: 27551942. Exclusion: Ineligible population.
480. Marino BCA, Ribeiro ALP, Alkmim MB, et al. Coordinated regional care of myocardial infarction in a rural area in Brazil: Minas Telecardio Project 2. *Eur Heart J Qual Care Clin Outcomes*. 2016;2(3):215-24. doi: 10.1093/ehjqcco/qcw020. PMID: 29474619. Exclusion: Ineligible intervention.
481. Mars M. Telemedicine and advances in urban and rural healthcare delivery in Africa. *Prog Cardiovasc Dis*. 2013;56(3):326-35. doi: 10.1016/j.pcad.2013.10.006. PMID: 24267440. Exclusion: Ineligible setting.
482. Martinez-Fernandez A, Lobos-Medina I, Diaz-Molina CA, et al. TulaSalud: an m-health system for maternal and infant mortality reduction in Guatemala. *J Telemed Telecare*. 2015;21(5):283-91. doi: 10.1177/1357633X15575830. PMID: 25766857. Exclusion: Ineligible setting.
483. Mashaw A. Implementation of a hospital readmissions prevention program in a rural geriatric population. *J Am Geriatr Soc*. 2014;62(10):1998-9. doi: 10.1111/jgs.13053. PMID: 25333544. Exclusion: Ineligible intervention.
484. Mason JD, Dino G, Boyd J, et al. Amplifying expertise in rural West Virginia through Project ECHO: impactful partnerships and community-engagement. *Progress in Community Health Partnerships*. 2021;15(2):235-42. doi: 10.1353/cpr.2021.0025. PMID: 34248067. Exclusion: 7/
485. Massey CN, Appel SJ, Buchanan KL, et al. Improving diabetes care in rural communities: an overview of current initiatives and a call for renewed efforts. *Clin Diabetes*. 2010;28(1):20-7. doi: 10.2337/diaclin.28.1.20. Exclusion: Not a study.
486. Mattos Sda S, Hazin SM, Regis CT, et al. A telemedicine network for remote paediatric cardiology services in north-east Brazil. *Bull World Health Organ*. 2015;93(12):881-7. doi: 10.2471/BLT.14.148874. PMID: 26668441. Exclusion: Ineligible study design.
487. Matylevich OP, Shushkevich AB, Perevoschikov PA, et al. Project ECHO in Belarus: outcomes of 3-year experience. *Int J Gynecol Cancer*. 2021;31(6):929-31. doi: 10.1136/ijgc-2020-002284. PMID: 33443035. Exclusion: Ineligible setting.
488. Maulik PK, Kallakuri S, Devarapalli S, et al. Increasing use of mental health services in remote areas using mobile technology: a pre-post evaluation of the SMART Mental Health project in rural India. *J Glob Health*. 2017;7(1):010408. doi: 10.7189/jogh.07.010408. PMID: 28400954. Exclusion: Ineligible setting.
489. Mauro E, Marciano S, Torres MC, et al. Telemedicine improves access to hepatology consultation with high patient satisfaction. *J Clin Exp Hepatol*. 2020;10(6):555-62. doi: 10.1016/j.jceh.2020.04.017. PMID: 33311892. Exclusion: Ineligible comparater.
490. Maxwell LG, McFarland MS, Baker JW, et al. Evaluation of the impact of a pharmacist-led telehealth clinic on diabetes-related goals of therapy in a veteran population. *Pharmacotherapy*. 2016;36(3):348-56. doi: 10.1002/phar.1719. PMID: 26877253. Exclusion: Ineligible intervention.

491. Mayberry LS, Lyles CR, Oldenburg B, et al. mHealth interventions for disadvantaged and vulnerable people with type 2 diabetes. *Curr Diab Rep*. 2019;19(12):148. doi: 10.1007/s11892-019-1280-9. PMID: 31768662. Exclusion: Ineligible intervention.
492. Mazighi M, Meseguer E, Labreuche J, et al. TRUST-tPA trial: Telemedicine for remote collaboration with urgentists for stroke-tPA treatment. *J Telemed Telecare*. 2017;23(1):174-80. doi: 10.1177/1357633X15615762. PMID: 26656722. Exclusion: Ineligible population.
493. Mazurek MO, Brown R, Curran A, et al. ECHO autism. *Clin Pediatr (Phila)*. 2017;56(3):247-56. doi: 10.1177/0009922816648288. PMID: 27169714. Exclusion: Ineligible population.
494. Mazurek MO, Stobbe G, Loftin R, et al. ECHO autism transition: enhancing healthcare for adolescents and young adults with autism spectrum disorder. *Autism*. 2020;24(3):633-44. doi: 10.1177/1362361319879616. PMID: 31581793. Exclusion: Ineligible population.
495. Mazzuoccolo LD, Esposito MN, Luna PC, et al. WhatsApp: a real-time tool to reduce the knowledge gap and share the best clinical practices in psoriasis. *Telemed J E Health*. 2019;25(4):294-300. doi: 10.1089/tmj.2018.0059. PMID: 29924684. Exclusion: Ineligible population.
496. McBain RK, Sousa JL, Rose AJ, et al. Impact of Project ECHO models of medical tele-education: a systematic review. *J Gen Intern Med*. 2019;34(12):2842-57. doi: 10.1007/s11606-019-05291-1. PMID: 31485970. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
497. McCord CE, Elliott TR, Brossard DF, et al. Community capacity and teleconference counseling in rural Texas. *Prof Psychol Res Pr*. 2011;42(6):521-7. doi: 10.1037/a0025296. Exclusion: Ineligible intervention.
498. McFarland LV, Raugi GJ, Reiber GE. Primary care provider and imaging technician satisfaction with a teledermatology project in rural Veterans Health Administration clinics. *Telemed J E Health*. 2013;19(11):815-25. doi: 10.1089/tmj.2012.0327. PMID: 24053115. Exclusion: Ineligible comparator.
499. McGraw K, Adler J, Andersen SB, et al. Mental health care for service members and their families across the globe. *Mil Med*. 2019;184(Suppl 1):418-25. doi: 10.1093/milmed/usy324. PMID: 30901426. Exclusion: Ineligible intervention.
500. McKellips F, Keely E, Afkham A, et al. Improving access to allied health professionals through the Champlain BASE TM eConsult service: a cross-sectional study in Canada. *Br J Gen Pract*. 2017;67(664):e757-e63. doi: 10.3399/bjgp17X693125. PMID: 28993307. Exclusion: Ineligible study design.
501. McLendon SF. Interactive video telehealth models to improve access to diabetes specialty care and education in the rural setting: a systematic review. *Diabetes Spectr*. 2017;30(2):124-36. doi: 10.2337/ds16-0004. PMID: 28588379. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
502. McLeod ME, Oladeru OT, Hao J, et al. Leveraging telehealth and medical student volunteers to bridge gaps in education access for providers in limited-resource settings. *Acad Med*. 2021;96(3):390-4. doi: 10.1097/ACM.0000000000003865. PMID: 33264112. Exclusion: Ineligible study design.

503. McPherson K, Nahon I. Telehealth and the provision of pelvic health physiotherapy in regional, rural and remote Australia. *Australian & New Zealand Continence Journal*. 2021;27(3):66-70. doi: 10.33235/anzcj.27.3.66-70. Exclusion: Ineligible intervention.
504. McSweeney S, Pritt JA, Swearingen A, et al. Telestroke: overcoming barriers to lifesaving treatment in rural hospitals. *Perspect Health Inf Manag*. 2017:1-18. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
505. McWilliams T, Hendricks J, Twigg D, et al. Telehealth for paediatric burn patients in rural areas: a retrospective audit of activity and cost savings. *Burns*. 2016;42(7):1487-93. doi: 10.1016/j.burns.2016.03.001. PMID: 27575678. Exclusion: Ineligible study design.
506. Medeiros de Bustos E, Berthier E, Chavot D, et al. Evaluation of a French regional telemedicine network dedicated to neurological emergencies: a 14-year study. *Telemed J E Health*. 2018;24(2):155-60. doi: 10.1089/tmj.2017.0035. PMID: 29346039. Exclusion: Ineligible outcome.
507. Mehraban Far P, Rullo J, Kratky V. Use of telemedicine in the management of life-threatening periorbital necrotizing fasciitis in a remote community. *CJEM*. 2020;22(1):118-20. doi: 10.1017/cem.2019.422. Exclusion: Not a study.
508. Meins AR, Doorenbos AZ, Eaton L, et al. TelePain: a community of practice for pain management. *J Pain Relief*. 2015;4(2):177-81. PMID: 25964869. Exclusion: Ineligible study design.
509. Mennenga HA, Johansen L, Foerster B, et al. Using simulation to improve student and faculty knowledge of telehealth and rural characteristics. *Nurs Educ Perspect*. 2016;37(5):287-8. PMID: 27740564. Exclusion: Ineligible intervention.
510. Mera J, Joshi K, Thornton K, et al. Retrospective study demonstrating high rates of sustained virologic response after treatment with direct-acting antivirals among American Indian/Alaskan Natives. *Open Forum Infect Dis*. 2019;6(7):1-8. doi: 10.1093/ofid/ofz128. PMID: 31289725. Exclusion: Ineligible intervention.
511. Meredith J, Onsrud J, Davidson L, et al. Successful use of telemedicine infectious diseases consultation with an antimicrobial stewardship-led *Staphylococcus aureus* bacteremia care bundle. *Open Forum Infect Dis*. 2021;8(6):ofab229. doi: 10.1093/ofid/ofab229. PMID: 34189171. Exclusion: Ineligible population.
512. Meyer BC, Raman R, Rao R, et al. The STRoKE DOC trial technique: 'video clip, drip, and/or ship'. *Int J Stroke*. 2007;2(4):281-7. Exclusion: Not a study.
513. Migliaretti G, Ciaramitaro P, Berchiolla P, et al. Teleconsulting for minor head injury: the Piedmont experience. *J Telemed Telecare*. 2013;19(1):33-5. doi: 10.1177/1357633X12474738. PMID: 23454822. Exclusion: Ineligible population.
514. Miller KL, Steffen MJ, McCoy KD, et al. Correction to: delivering fracture prevention services to rural US veterans through telemedicine: a process evaluation. *Arch Osteoporos*. 2021;16(1):61. doi: 10.1007/s11657-021-00916-7. PMID: 33825995. Exclusion: Not a study.
515. Miller KL, Steffen MJ, McCoy KD, et al. Delivering fracture prevention services to rural US veterans through telemedicine: a process evaluation. *Arch Osteoporos*. 2021;16(1):27. doi: 10.1007/s11657-021-00882-0. PMID: 33566174. Exclusion: Ineligible population.
516. Milsten J, Ponniah K, McCoy M, et al. Utility of telemedicine for screening congenital heart disease in a level II NICU. *J Investig Med*. 2020;68(2):664. doi: 10.1136/jim-2020-SRM.561. Exclusion: Ineligible population.

517. Ming J, Kelty E, Martin K. 'Half the time it's just guessing': youth worker and youth service manager experiences of sexual health training in the Pilbara, remote Western Australia. *Sex Educ.* 2021;21(2):221-35. doi: 10.1080/14681811.2020.1782184. Exclusion: Ineligible study design.
518. Mishra V, Moeller G. Impacting the opioid crisis through project echo. *J Gen Intern Med.* 2020;35(SUPPL 1):S735-S6. doi: 10.1007/s11606-020-05890-3. Exclusion: Not a study.
519. Mitchell L, Schultz K, Clark D, et al. Telemedicine videoconferencing - breaking down the barriers for cystic fibrosis (CF) patients living in rural and remote areas. *J Cyst Fibros.* 2014;13:S105. Exclusion: Ineligible intervention.
520. Mitchell S, Oliver P, Gardiner C, et al. Community end-of-life care during the COVID-19 pandemic: findings of a UK primary care survey. *BJGP Open.* 2021;5(4)doi: 10.3399/BJGPO.2021.0095. PMID: 34117014. Exclusion: Ineligible comparator.
521. Mitchell S, Oliver P, Gardiner C, et al. Community end-of-life care during COVID-19: findings of a UK primary care survey. *BJGP Open.* 2021;11:11. doi: 10.3399/BJGPO.2021.0095. PMID: 34117014. Exclusion: Ineligible study design.
522. Mitruka K, Thornton K, Cusick S, et al. Expanding primary care capacity to treat hepatitis C virus infection through an evidence-based care model--Arizona and Utah, 2012-2014. *MMWR Morb Mortal Wkly Rep.* 2014;63(18):393-8. PMID: 24807237. Exclusion: Ineligible intervention.
523. Mittal D, Chekuri L, Lu L, et al. Demographic, economic, and clinical correlates of depression treatment response in an underserved primary care population. *J Clin Psychiatry.* 2014;75(8):848-54. doi: 10.4088/JCP.13m08954. PMID: 29676557. Exclusion: Ineligible population.
524. Mitton C, Dionne F, Masucci L, et al. Innovations in health service organization and delivery in northern rural and remote regions: a review of the literature. *Int J Circumpolar Health.* 2011;70(5):460-72. PMID: 22030009. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
525. Miyamoto S, Boyle C, Yang NH, et al. Evaluation of telehealth to support pediatric sexual abuse examinations in rural communities. *J Investig Med.* 2013;61(1):116. doi: 10.231/JIM.0b013e31827d3ac9. Exclusion: Ineligible study design.
526. Miyamoto S, Thiede E, Wright EN, et al. The implementation of the Sexual Assault Forensic Examination Telehealth Center: a program evaluation. *J Forensic Nurs.* 2021;17(3):E24-E33. doi: 10.1097/JFN.0000000000000337. PMID: 34132652. Exclusion: Ineligible population.
527. Moeckli J, Stewart KR, Ono S, et al. Mixed-methods study of uptake of the Extension for Community Health Outcomes (ECHO) telemedicine model for rural veterans with HIV. *J Rural Health.* 2017;33(3):323-31. doi: 10.1111/jrh.12200. PMID: 27557039. Exclusion: Ineligible population.
528. Mohan B, Mehra AO, Naik N, et al. Clinical profile and outcome of critical cardiac patients treated with telemedicine using WhatsApp ECG group. *Indian Heart J.* 2018;70:S100. doi: 10.1016/j.ihj.2018.10.299. Exclusion: Ineligible setting.
529. Mohr NM, Harland KK, Okoro UE, et al. TELEmedicine as an intervention for sepsis in emergency departments: a multicenter, comparative effectiveness study (TELEvised Study). *J Comp Eff Res.* 2021;10(2):77-91. doi: 10.2217/ce-2020-0141. PMID: 33470848. Exclusion: Not a study.

530. Mohsen W, Chan P, Whelan M, et al. Hepatitis C treatment for difficult to access populations: can telementoring (as distinct from telemedicine) help? *Intern Med J*. 2019;49(3):351-7. doi: 10.1111/imj.14072. PMID: 30091164. Exclusion: Ineligible population.
531. Molinari G, Molinari M, Di Biase M, et al. Telecardiology and its settings of application: an update. *J Telemed Telecare*. 2018;24(5):373-81. PMID: 28084886. Exclusion: Not a study.
532. Montelongo A, Becker JL, Roman R, et al. The management of COVID-19 cases through telemedicine in Brazil. *PLoS ONE*. 2021;16(7):e0254339. doi: 10.1371/journal.pone.0254339. PMID: 34260644. Exclusion: Ineligible study design.
533. Moore K, Haboubi M, Wise E, et al. Cerebrovascular trained advance practice nurse practitioners can safely and accurately administer alteplase as well as their neurology physician colleagues via telemedicine. *Neurology*. 2021;96(15 SUPPL 1). Exclusion: Ineligible comparater.
534. Morgan T, Brown AT, Onteddu S, et al. Validation of an education gauge for measuring stroke outreach efforts. *Stroke*. 2018;49. Exclusion: Not a study.
535. Morris BB, Rossi B, Fuemmeler B. The role of digital health technology in rural cancer care delivery: a systematic review. *J Rural Health*. 2021;04:04. doi: 10.1111/jrh.12619. PMID: 34480506. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
536. Morrison JL, Cai Q, Davis N, et al. Clinical and economic outcomes of the electronic intensive care unit: results from two community hospitals. *Crit Care Med*. 2010;38(1):2-8. doi: 10.1097/CCM.0b013e3181b78fa8. PMID: 19730249. Exclusion: Ineligible population.
537. Mort A, Eadie L, Regan L, et al. Combining transcranial ultrasound with intelligent communication methods to enhance the remote assessment and management of stroke patients: framework for a technology demonstrator. *Health Inform J*. 2016;22(3):691-701. doi: 10.1177/1460458215580353. PMID: 25975807. Exclusion: Ineligible outcome.
538. Mottet T, Soubeyran P, Godbert Y, et al. 1613P What are the barriers to routine clinical use of teleconsultation in oncology? A retrospective study on patient's and their physician's satisfaction with 603 video teleconsultations. *Ann Oncol*. 2021;32:S1150. doi: 10.1016/j.annonc.2021.08.1606. Exclusion: Not a study.
539. Moya M, Valdez J, Yonas H, et al. The impact of a telehealth web-based solution on neurosurgery triage and consultation. *Telemed J E Health*. 2010;16(9):945-9. doi: 10.1089/tmj.2010.0044. PMID: 21034238. Exclusion: Ineligible comparater.
540. Muhame RM, Dragulescu A, Nadimpalli A, et al. Cardiac point of care ultrasound in resource limited settings to manage children with congenital and acquired heart disease. *Cardiol Young*. 2021;1-7. doi: 10.1017/S1047951121000834. PMID: 33682650. Exclusion: Ineligible population.
541. Muir SD, de Boer K, Nedeljkovic M, et al. Barriers and facilitators of videoconferencing psychotherapy implementation in veteran mental health care environments: a systematic review. *BMC Health Serv Res*. 2020;20(1):999. doi: 10.1186/s12913-020-05858-3. PMID: 33131495. Exclusion: Ineligible intervention.
542. Mullen-Fortino M, Sites FD, Soisson M, et al. Innovative use of tele-ICU in long-term acute care hospitals. *AACN Adv Crit Care*. 2012;23(3):330-6. PMID: 22828067. Exclusion: Not a study.

543. Muller-Barna P, Hubert GJ, Boy S, et al. TeleStroke units serving as a model of care in rural areas: 10-year experience of the TeleMedical project for integrative stroke care. *Stroke*. 2014;45(9):2739-44. doi: 10.1161/STROKEAHA.114.006141. PMID: 25147327. Exclusion: Ineligible outcome.
544. Müller-Barna P, Boy S, Hubert GJ, et al. Convincing quality of acute stroke care in TeleStroke units. *Eur Res Telemed*. 2015;4(2):53-61. doi: 10.1016/j.eurtel.2015.04.001. Exclusion: Ineligible population.
545. Mwapasa G, Pittalis C, Clarke M, et al. Evaluation of a managed surgical consultation network in Malawi. *World J Surg*. 2021;45(2):356-61. doi: 10.1007/s00268-020-05809-3. PMID: 33026475. Exclusion: Ineligible setting.
546. Myers US, Birks A, Grubaugh AL, et al. Flattening the curve by getting ahead of it: how the VA healthcare system is leveraging telehealth to provide continued access to care for rural veterans. *J Rural Health*. 2021;37:194-6. doi: 10.1111/jrh.12449. PMID: 32282955. Exclusion: Not a study.
547. Nabelsi V, Levesque-Chouinard A, Liddy C, et al. Improving the referral process, timeliness, effectiveness, and equity of access to specialist medical services through electronic consultation: pilot study. *JMIR Med Inform*. 2019;7(3):e13354. doi: 10.2196/13354. PMID: 31293239. Exclusion: Ineligible population.
548. Nadig N, Hynden L, Sterba K, et al. Preliminary development of value scorecards as ICU telemedicine evaluation tools. *J Healthc Manag*. 2021;66(2):124-38. doi: 10.1097/JHM-D-19-00188. PMID: 33692317. Exclusion: Ineligible outcome.
549. Nadig NR, Hynden L, McElligott JT, et al. Value scorecards as a tele-ICU evaluation tool. *Am J Respir Crit Care Med*. 2018;197(MeetingAbstracts). Exclusion: Not a study.
550. Nagata JM. Rapid scale-up of telehealth during the COVID-19 pandemic and implications for subspecialty care in rural areas. *J Rural Health*. 2020;1. doi: 10.1111/jrh.12433. PMID: 32246490. Exclusion: Not a study.
551. Nagayoshi Y, Oshima S, Ogawa H. Response to Pollari et al. re: "clinical impact of telemedicine network system at rural hospitals without on-site cardiac surgery backup". *Telemed J E Health*. 2017;23(3):260-1. doi: 10.1089/tmj.2016.29011.yn. PMID: 27525984. Exclusion: Not a study.
552. Nazareth S, Kontorinis N, Muwanwella N, et al. Successful treatment of patients with hepatitis C in rural and remote Western Australia via telehealth. *J Telemed Telecare*. 2013;19(2):101-6. doi: 10.1258/jtt.2012.120612. PMID: 23528788. Exclusion: Ineligible intervention.
553. Nesbitt TS, Dharmar M, Katz-Bell J, et al. Telehealth at UC Davis--a 20-year experience. *Telemed J E Health*. 2013;19(5):357-62. doi: 10.1089/tmj.2012.0284. PMID: 23343257. Exclusion: Ineligible intervention.
554. Ness T, Annese MF, Martinez-Paz N, et al. Using an innovative telehealth model to support providers in geographically dispersed areas who deliver care to HIV-positive pregnant women. *Am J Trop Med Hyg*. 2016;95(5):63. doi: 10.4269/ajtmh.abstract2016. Exclusion: Not a study.
555. Nethan ST, Hariprasad R, Babu R, et al. Project ECHO: a potential best-practice tool for training healthcare providers in oral cancer screening and tobacco cessation. *J Cancer Educ*. 2019;35:965-71. doi: 10.1007/s13187-019-01549-8. PMID: 31124001. Exclusion: Ineligible population.
556. Nettesheim N, Powell D, Vasios W, et al. Telemedical support for military medicine. *Mil Med*. 2018;183(11-12):e462-e70. doi: 10.1093/milmed/usy127. PMID: 30496581. Exclusion: Not a study.

557. Neufeld J, Case R. Walk-in telemental health clinics improve access and efficiency: a 2-year follow-up analysis. *Telemed J E Health*. 2013;19(12):938-41. doi: 10.1089/tmj.2013.0076. PMID: 24050609. Exclusion: Ineligible intervention.
558. Newman P, Dhaliwall S, Polyakova O, et al. Pharmacy distribution, clinical, and management services: a survey of small hospitals in Canada supported by telepharmacy services. *Can J Hosp Pharm*. 2021;74(3):256-68. doi: 10.4212/CJHP.V74I3.3153. Exclusion: Ineligible comparator.
559. Nguyen-Huynh MN, Klingman JG, Avins AL, et al. Novel Telestroke Program Improves Thrombolysis for Acute Stroke Across 21 Hospitals of an Integrated Healthcare System. *Stroke*. 2018;49(1):133-9. doi: 10.1161/STROKEAHA.117.018413. PMID: 29247142. Exclusion: Ineligible population.
560. Nguyen C, Mbuthia J, Dobson CP. Reduction in medical evacuations from Iraq and Syria following introduction of an asynchronous telehealth system. *Mil Med*. 2020;185:e1693-9. doi: 10.1093/milmed/usaa091. PMID: 32592397. Exclusion: Ineligible population.
561. Nguyen KH, Smith AC, Armfield NR, et al. Cost-effectiveness analysis of a mobile ear screening and surveillance service versus an outreach screening, surveillance and surgical service for indigenous children in Australia. *PLoS One*. 2015;10(9):e0138369. doi: 10.1371/journal.pone.0138369. PMID: 26406592. Exclusion: Ineligible intervention.
562. Nhung LH, Dien TM, Lan NP, et al. Use of Project ECHO telementoring model in continuing medical education for pediatricians in Vietnam: preliminary results. *Health Serv Insights*. 2021;14:11786329211036855. doi: 10.1177/11786329211036855. PMID: 34408433. Exclusion: Ineligible population.
563. Ni Cheallaigh C, O'Leary A, Keating S, et al. Telementoring with project ECHO: a pilot study in Europe. *BMJ Innov*. 2017;3(3):144-51. doi: 10.1136/bmjinnov-2016-000141. PMID: 29445515. Exclusion: Ineligible population.
564. Nietert PJ, Tilley BC, Zhao W, et al. Two pharmacy interventions to improve refill persistence for chronic disease medications: a randomized, controlled trial. *Med Care*. 2009;47(1):32-40. doi: 10.1097/MLR.0b013e3181808c17. PMID: 19106728. Exclusion: Ineligible intervention.
565. Niknejad N, Ismail W, Bahari M, et al. Understanding telerehabilitation technology to evaluate stakeholders' adoption of telerehabilitation services: a systematic literature review and directions for further research. *Arch Phys Med Rehabil*. 2021;102(7):1390-403. doi: 10.1016/j.apmr.2020.12.014. PMID: 33484693. Exclusion: Ineligible population.
566. Nisbet D, Robertson A, Mannil B, et al. Quality management of nuchal translucency ultrasound measurement in Australia. *Aust N Z J Obstet Gynaecol*. 2019;59(1):54-8. doi: 10.1111/ajo.12792. PMID: 29469171. Exclusion: Ineligible population.
567. Nyamtema A, Mwakatundu N, Dominico S, et al. Introducing eHealth strategies to enhance maternal and perinatal health care in rural Tanzania. *Matern Health Neonatol Perinatol*. 2017;3:3. doi: 10.1186/s40748-017-0042-4. PMID: 28116114. Exclusion: Ineligible setting.
568. O'Connor T. Videoconferences bring specialist care closer to home. *Kai Tiaki Nursing New Zealand*. 2013;19(6):15-. Exclusion: Not a study.
569. O'Gorman LD, Hogenbirk JC, Warry W. Clinical telemedicine utilization in Ontario over the Ontario telemedicine network. *Telemed J E Health*. 2016;22(6):473-9. doi: 10.1089/tmj.2015.0166. PMID: 26544163. Exclusion: Ineligible intervention.

570. O'Hara R, Jackson S. Integrating telehealth services into a remote allied health service: a pilot study. *Aust J Rural Health*. 2017;25(1):53-7. doi: 10.1111/ajr.12189. PMID: 25823551. Exclusion: Ineligible study design.
571. O'Neil J, van Ierssel J, Sveistrup H. Remote supervision of rehabilitation interventions for survivors of moderate or severe traumatic brain injury: a scoping review. *J Telemed Telecare*. 2020;26(9):520-35. doi: 10.1177/1357633X19845466. PMID: 31148489. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
572. Obaid M, Zhang Q, Adams SJ, et al. Development and assessment of a teleultrasonography system for musculoskeletal imaging. *Eur Radiol Exp*. 2021;5(1):29. doi: 10.1186/s41747-021-00227-z. PMID: 34312733. Exclusion: Ineligible outcome.
573. Ochiai H, Ohta H, Kanemaru K, et al. Implementation of a telestroke system for general physicians without a nearby stroke center to shorten the time to intravenous thrombolysis for acute cerebral infarction. *Acute Med Surg*. 2020;7(1):e551. doi: 10.1002/ams2.551. PMID: 32802346. Exclusion: Ineligible study design.
574. Ofoma UR, Maddox TM, Perera C, et al. Characteristics of U.S. acute care hospitals that have implemented telemedicine critical care. *Crit Care Explor*. 2021;3(7):e0468. doi: 10.1097/CCE.0000000000000468. PMID: 34235456. Exclusion: Ineligible study design.
575. Olayiwola JN, Knox M, Dubé K, et al. Understanding the Potential for Patient Engagement in Electronic Consultation and Referral Systems: Lessons From One Safety Net System. *Health Serv Res*. 2018;53(4):2483-502. doi: 10.1111/1475-6773.12776. PMID: 28940495. Exclusion: Ineligible population.
576. Oliver K, Beskin K, Noonan L, et al. A quality improvement learning collaborative for human papillomavirus vaccination. *Pediatr Qual Saf*. 2021;6(1):e377. doi: 10.1097/pq9.0000000000000377. PMID: 33409429. Exclusion: Ineligible outcome.
577. Oser T, Litchman ML, Kwan B, et al. Cultural adaptation of a diabetes self-management education and support telehealth intervention for rural Colorado. *Diabetes*. 2021;70(SUPPL 1)doi: 10.2337/db21-466-P. Exclusion: Ineligible intervention.
578. Osman MA, Schick-Makaroff K, Thompson S, et al. Barriers and facilitators for implementation of electronic consultations (eConsult) to enhance access to specialist care: a scoping review. *BMJ Glob Health*. 2019;4(5):e001629. doi: 10.1136/bmjgh-2019-001629. PMID: 31565409. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
579. Osofsky HJ, Osofsky JD, Wells JH, et al. Integrated care: meeting mental health needs after the Gulf oil spill. *Psychiatr Serv*. 2014;65(3):280-3. doi: 10.1176/appi.ps.201300470. PMID: 24584523. Exclusion: Not a study.
580. Osotimehin S, Sokan O, Rodriguez De Bittner M, et al. An innovative transitions of care (TOC) model incorporating pharmacy services by leveraging telehealth. *JACCP*. 2020;3(1):304. doi: 10.1002/jac5.1204. Exclusion: Not a study.
581. Ott KK, Schein RM, Straatmann J, et al. Development of a home-based telerehabilitation service delivery protocol for wheelchair seating and mobility within the Veterans Health Administration. *Mil Med*. 2021;27:27. doi: 10.1093/milmed/usab091. PMID: 33647955. Exclusion: Ineligible study design.

582. Otto C, Shemenski R, Scott JM, et al. Evaluation of tele-ultrasound as a tool in remote diagnosis and clinical management at the Amundsen-Scott South Pole Station and the McMurdo Research Station. *Telemed J E Health*. 2013;19(3):186-91. doi: 10.1089/tmj.2012.0111. PMID: 23480714. Exclusion: Ineligible study design.
583. Owen RR, Woodward EN, Drummond KL, et al. Using implementation facilitation to implement primary care mental health integration via clinical video telehealth in rural clinics: protocol for a hybrid type 2 cluster randomized stepped-wedge design. *Implement Sci*. 2019;14(1):33. doi: 10.1186/s13012-019-0875-5. PMID: 30898129. Exclusion: Not a study.
584. Padayachee A, Ranatunga C, Comans TA. Utilising capacity in a rural hospital to support older people requiring hospital care: Kilcoy connect. *Aust J Rural Health*. 2019;27(4):344-50. doi: 10.1111/ajr.12475. PMID: 30693988. Exclusion: Ineligible comparator.
585. Page K, Qeadan F, Qualls C, et al. Project ECHO revisited: propensity score analysis and HCV treatment outcomes. *Hepat Med*. 2019;11:149-52. doi: 10.2147/HMER.S212855. PMID: 31632162. Exclusion: Ineligible population.
586. Paik AM, Granick MS, Scott S. Plastic surgery telehealth consultation expedites emergency department treatment. *J Telemed Telecare*. 2017;23(2):321-7. doi: 10.1177/1357633X16639459. PMID: 27056907. Exclusion: Ineligible population.
587. Palen BN, Mattox EA, He K, et al. Impact of sleep telementorship in primary care: Sleep VA-ECHO (Veterans Affairs-Extension for Community Healthcare Outcomes). *Int J Environ Res Public Health*. 2021;18(18):21. doi: 10.3390/ijerph18189914. PMID: 34574837. Exclusion: Ineligible population.
588. Palmas W, Shea S, Starren J, et al. Medicare payments, healthcare service use, and telemedicine implementation costs in a randomized trial comparing telemedicine case management with usual care in medically underserved participants with diabetes mellitus (IDEATel). *J Am Med Inform Assoc*. 2010;17(2):196-202. doi: 10.1136/jamia.2009.002592. PMID: 20190064. Exclusion: Ineligible population.
589. Palmer NB, Myers KM, Vander Stoep A, et al. Attention-deficit/hyperactivity disorder and telemental health. *Curr Psychiatry Rep*. 2010;12(5):409-17. doi: 10.1007/s11920-010-0132-8. PMID: 20625857. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
590. Pappesh ND, Gebel J. Benefiting the community; the impact of joining a regional stroke network. *Stroke*. 2012;43(2). Exclusion: Not a study.
591. Papoutsis C, Wherton J, Shaw S, et al. Putting the social back into sociotechnical: case studies of co-design in digital health. *J Am Med Inform Assoc*. 2021;28(2):284-93. doi: 10.1093/jamia/ocaa197. PMID: 33043359. Exclusion: Ineligible setting.
592. Park J, Erikson C, Han X, et al. Are state telehealth policies associated with the use of telehealth services among underserved populations? *Health Aff (Millwood)*. 2018;37(12):2060-8. doi: 10.1377/hlthaff.2018.05101. PMID: 30633679. Exclusion: Ineligible intervention.
593. Parnell K, Kuhlenschmidt K, Madni D, et al. Using telemedicine on an acute care surgery service: improving clinic efficiency and access to care. *Surg Endosc*. 2020;13:13. doi: 10.1007/s00464-020-08055-9. PMID: 33048233. Exclusion: Ineligible setting.

594. Parsons EC, Mattox EA, Beste LA, et al. Development of a sleep telementorship program for rural department of Veterans Affairs primary care providers: sleep Veterans Affairs Extension for Community Healthcare Outcomes. *Ann Am Thorac Soc*. 2017;14(2):267-74. doi: 10.1513/AnnalsATS.201605-361BC. PMID: 27977293. Exclusion: Ineligible population.
595. Patel UK, Malik P, DeMasi M, et al. Multidisciplinary approach and outcomes of tele-neurology: a review. *Cureus*. 2019;11(4):e4410. doi: 10.7759/cureus.4410. PMID: 31205830. Exclusion: Not a study.
596. Patil R, Shrivastava R, Juvekar S, et al. Specialist to non-specialist teleconsultations in chronic respiratory disease management: a systematic review. *J Glob Health*. 2021;11:04019. doi: 10.7189/jogh.11.04019. PMID: 34326988. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
597. Paul MM, Saad AD, Billings J, et al. A telementoring intervention leads to improvements in self-reported measures of health care access and quality among patients with complex diabetes. *J Health Care Poor Underserved*. 2020;31(3):1124-33. doi: 10.1353/hpu.2020.0085. PMID: 33416685. Exclusion: Ineligible intervention.
598. Pearson J, Anderholm K, Bettermann M, et al. Obstetrical care in rural Minnesota: family physician perspectives on factors affecting the ability to provide prenatal, labor, and delivery care. *J Rural Health*. 2021;37(2):362-72. doi: 10.1111/jrh.12478. PMID: 32602949. Exclusion: Ineligible study design.
599. Peiris D, Praveen D, Mogulluru K, et al. SMARThealth India: a stepped-wedge, cluster randomised controlled trial of a community health worker managed mobile health intervention for people assessed at high cardiovascular disease risk in rural India. *PLoS One*. 2019;14(3):e0213708. doi: 10.1371/journal.pone.0213708. PMID: 30913216. Exclusion: Ineligible setting.
600. Pellatt RAF, Bolot R, Sweeny AL, et al. Rural and remote intubations in an Australian Air Medical Retrieval Service: a retrospective cohort study. *Air Med J*. 2021;40(4):251-8. doi: 10.1016/j.amj.2021.03.006. PMID: 34172233. Exclusion: Ineligible intervention.
601. Perry TT, Halterman JS, Brown RH, et al. Breath connection: a school-based telemedicine program for rural children with asthma. *J Allergy Clin Immunol*. 2015;135(2 Suppl 1):AB169. Exclusion: Not a study.
602. Pervez MA, Silva G, Masrur S, et al. Remote supervision of IV-tPA for acute ischemic stroke by telemedicine or telephone before transfer to a regional stroke center is feasible and safe. *Stroke*. 2010;41(1):e18-24. doi: 10.1161/strokeaha.109.560169. PMID: 19910552. Exclusion: Ineligible population.
603. Peterson CD, Scott DM, Rathke A, et al. Establishing a central order entry system for delivering telepharmacy services to remote rural hospitals. *J Pharm Tech*. 2010;26(4):179-86. Exclusion: Ineligible intervention.
604. Phillips D, Matheson L, Pain T, et al. Development of an occupational-therapy-led paediatric burn telehealth review clinic. *Rural Remote Health*. 2021;21(3):6223. doi: 10.22605/RRH6223. PMID: 34392690. Exclusion: Ineligible outcome.
605. Pichard-Jolicoeur A, Mbakop-Nguebou M, Dogba J, et al. Literature review of telemedicine for trauma patients in rural areas. *CJEM*. 2016;18:S113. doi: 10.1017/cem.2016.280. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
606. Pippitt K, Junkins S, Baggaley S. Practical barriers to Project ECHO innovations. *Acad Med*. 2017;92(3):277-8. doi: 10.1097/ACM.0000000000001554. PMID: 28221231. Exclusion: Not a study.

607. Piriz Alvarez G. Technology for improving accessibility of end-of-life care: Extension for Community Healthcare Outcomes Project. *Curr Opin Support Palliat Care*. 2018;12(4):466-71. doi: 10.1097/SPC.0000000000000390. PMID: 30300151. Exclusion: Not a study.
608. Pirkey JM, Levey JA, Newberry SM, et al. Videoconferencing expands nursing students' cultural realm. *J Nurs Educ*. 2012;51(10):586-90. doi: 10.3928/01484834-20120823-01. PMID: 22908884. Exclusion: Ineligible population.
609. Pollari F, Giorlandino MS, Fischlein T. Re: "Clinical impact of telemedicine network system at rural hospitals without on-site cardiac surgery backup" by Nagayoshi et al. (*Telemed J E Health*. 2017;23:260-261). *Telemed J E Health*. 2017;23(3):259-60. doi: 10.1089/tmj.2016.0115. PMID: 27529654. Exclusion: Not a study.
610. Pond D, Higgins I, Mate K, et al. Mobile memory clinic: implementing a nurse practitioner-led, collaborative dementia model of care within general practice. *Aust J Prim Health*. 2021;27(1):6-12. doi: 10.1071/PY20118. PMID: 33517974. Exclusion: Ineligible intervention.
611. Ponnamp HB, Akondi BR. Telemedicine health care delivery in india: a boon during covid-19 pandemic. *Asian J Pharm Res Health Care*. 2021;13(3):227-9. doi: 10.18311/ajprhc/2021/28020. Exclusion: Ineligible setting.
612. Porter KJ, Moon KE, LeBaron VT, et al. A novel behavioral intervention for rural appalachian cancer survivors (weSurvive): participatory development and proof-of-concept testing. *JMIR Cancer*. 2021;7(2):e26010. doi: 10.2196/26010. PMID: 33843597. Exclusion: Ineligible population.
613. Potapov A, Olayiwola JN, Radix AE, et al. Electronic consultations as an educational tool to improve the care of transgender patients in primary care. *J Health Care Poor Underserved*. 2021;32(2):680-7. doi: 10.1353/hpu.2021.0097. PMID: 34120969. Exclusion: Ineligible outcome.
614. Prada C, Izquierdo N, Traipe R, et al. Results of a new telemedicine strategy in traumatology and orthopedics. *Telemed J E Health*. 2020;26(5):665-70. doi: 10.1089/tmj.2019.0090. PMID: 31287786. Exclusion: Ineligible study design.
615. Pradhan T, Six-Workman EA, Law KB. An innovative approach to care: integrating mental health services through telemedicine in rural school-based health centers. *Psychiatr Serv*. 2019;70(3):239-42. doi: 10.1176/appi.ps.201800252. PMID: 30554561. Exclusion: Not a study.
616. Pranata R, Tondas AE, Huang I, et al. Potential role of telemedicine in solving ST-segment elevation dilemmas in remote areas during the COVID-19 pandemic. *Am J Emerg Med*. 2021;42:242-3. doi: 10.1016/j.ajem.2020.06.012. PMID: 32561139. Exclusion: Not a study.
617. Preclaro IA, Tianco EA, Gulmatico-Flores Z. Concordance and reliability of teledermatology using mobile phones in the outpatient clinic in a tertiary hospital: a cross-sectional study. *J Dermatol Nurses Assoc*. 2020;12(2). Exclusion: Not a study.
618. Prinja S, Nimesh R, Gupta A, et al. Impact of m-health application used by community health volunteers on improving utilisation of maternal, new-born and child health care services in a rural area of Uttar Pradesh, India. *Trop Med Int Health*. 2017;22(7):895-907. doi: 10.1111/tmi.12895. PMID: 28510997. Exclusion: Ineligible setting.
619. Puckett HM, Bossaller JS, Sheets LR. The impact of project ECHO on physician preparedness to treat opioid use disorder: a systematic review. *Addict Sci Clin Pract*. 2021;16(1):6. doi: 10.1186/s13722-021-00215-z. PMID: 33482906. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.

620. Quinn EM, Corrigan MA, O'Mullane J, et al. Clinical unity and community empowerment: the use of smartphone technology to empower community management of chronic venous ulcers through the support of a tertiary unit. *PLoS One*. 2013;8(11):e78786. doi: 10.1371/journal.pone.0078786. PMID: 24265716. Exclusion: Ineligible intervention.
621. Rabinowitz T, Murphy KM, Amour JL, et al. Benefits of a telepsychiatry consultation service for rural nursing home residents. *Telemed J E Health*. 2010;16:34-40. doi: 10.1089/tmj.2009.0088. PMID: 20070161. Exclusion: Ineligible intervention.
622. Rada G. Telemedicine: are we advancing the science? *Cochrane Database Syst Rev*. 2015 (9). Exclusion: Not a study. Summarizes the challenges of doing SR in telehealth, good for background?
623. Ragan A, Tran M, Moore T. Impact of mental health clinical pharmacist practitioners to improve veteran access in VA clinical resource hubs. *J Am Coll Clin Pharm*. 2021;4(9):1246. doi: 10.1002/jac5.1481. Exclusion: Not a study.
624. Rainwater L, Lichiello S, Duckworth K, et al. Psychosocial oncology care during a pandemic: effects of offering telemental health in an academic medical center. *Psycho-Oncol*. 2021;30(SUPPL 1):37-8. doi: 10.1002/pon.5637. Exclusion: Ineligible intervention.
625. Rajbhandari H, Joshi S, Malakar S, et al. Epilepsy field workers, a smartphone application and telephone telemedicine: safe and effective epilepsy care in rural Nepal. *Seizure*. 2019;64:54-8. doi: 10.1016/j.seizure.2018.12.005. PMID: 30562653. Exclusion: Ineligible setting.
626. Ramsingh D, Van Gorkom C, Holsclaw M, et al. Use of a smartphone-based augmented reality video conference app to remotely guide a point of care ultrasound examination. *Diagnostics (Basel)*. 2019;9(4):24. doi: 10.3390/diagnostics9040159. PMID: 31652998. Exclusion: Ineligible intervention.
627. Rariy C, Truesdale L, Ahn ER, et al. Bridging the gap by providing access to oncology care to rural communities: a hybrid delivery model combining in-person visits with Telehealth. *Am J Clin Oncol*. 2021;39(15 SUPPL)doi: 10.1200/JCO.2021.39.suppl.e18528. Exclusion: Ineligible intervention.
628. Rashid A, Soda H, Kippnich U, et al. Stroke angel: telemedicine prenotification improves door-to-CT and rate of systemic thrombolysis. *Cerebrovasc Dis*. 2015;39:33. Exclusion: Ineligible intervention.
629. Rawle M, Oliver T, Pighills A, et al. Improving education and supervision of Queensland X-ray Operators through video conference technology: a teleradiography pilot project. *J Med Radiat Sci*. 2017;64(4):244-50. doi: 10.1002/jmrs.229. PMID: 28407440. Exclusion: Ineligible outcome.
630. Ray KN, Demirci JR, Bogen DL, et al. Optimizing telehealth strategies for subspecialty care: recommendations from rural pediatricians. *Telemed J E Health*. 2015;21(8):622-9. doi: 10.1089/tmj.2014.0186. PMID: 25919585. Exclusion: Ineligible outcome.
631. Razavi H, Copeland SP, Turner AW. Increasing the impact of teleophthalmology in Australia: analysis of structural and economic drivers in a state service. *Aust J Rural Health*. 2017;25(1):45-52. doi: 10.1111/ajr.12277. PMID: 26781722. Exclusion: Ineligible intervention.
632. Rebello KE, Gosian J, Salow M, et al. The Rural PILL program: a postdischarge telepharmacy intervention for rural veterans. *J Rural Health*. 2017;33(3):332-9. doi: 10.1111/jrh.12212. PMID: 27606938. Exclusion: Ineligible intervention.

633. Recto P, Lesser J, Zapata J, Jr., et al. The development and implementation of a COVID-19 Project ECHO: a program for community health workers serving populations from rural and medically underserved areas in south Texas. *Issues Ment Health Nurs.* 2021;1-5. doi: 10.1080/01612840.2021.1954449. PMID: 34379563. Exclusion: Ineligible study design.
634. Reddy ER, Bartlett PJ, Harnett JD, et al. Telemedicine and fetal ultrasonography in a remote Newfoundland community. *Can Med Assoc J.* 2000;162(2):206-7. PMID: 10674053. Exclusion: Ineligible outcome.
635. Reeves H, Grandjean Lapierre S, Razafindrina K, et al. Evaluating the use of educational videos to support the tuberculosis care cascade in remote Madagascar. *Int J Tuberc Lung Dis.* 2020;24(1):28-35. doi: 10.5588/ijtld.19.0161. PMID: 32005304. Exclusion: Ineligible setting.
636. Resnick HE, Ilagan PR, Kaylor MB, et al. TEAhM-Technologies for enhancing access to health management: a pilot study of community-based telehealth. *Telemed J E Health.* 2012;18(3):166-74. doi: 10.1089/tmj.2011.0122. PMID: 22364270. Exclusion: Ineligible intervention.
637. Reynolds PM, Rhein E, Nuffer M, et al. Educational methods and technological innovations for introductory experiential learning given the contact-related limitations imposed by the SARS-CoV2/COVID-19 pandemic. *Pharmacy (Basel).* 2021;9(1):25. doi: 10.3390/pharmacy9010047. PMID: 33668864. Exclusion: Ineligible population.
638. Rhodes LA, Register S, Asif I, et al. Alabama Screening and Intervention for Glaucoma and eye Health through Telemedicine (AL-SIGHT): study design and methodology. *J Glaucoma.* 2021;30(5):371-9. doi: 10.1097/IJG.0000000000001794. PMID: 33492893. Exclusion: Not a study.
639. Richman J, Hasselberg M. Using technology to support collaborative approaches to psychiatric problems in rural areas. *Psychother Psychosom.* 2019;88:108. doi: 10.1159/000502467. Exclusion: Not a study.
640. Rikin S, Zhang C, Lipsey D, et al. Impact of an opt-in econsult program on primary care demand for specialty visits: stepped-wedge cluster randomized implementation study. *J Gen Intern Med.* 2020;35:832-8. doi: 10.1007/s11606-020-06101-9. PMID: 32779140. Exclusion: Ineligible population.
641. Riley WT, Keberlein P, Sorenson G, et al. Program evaluation of remote heart failure monitoring: healthcare utilization analysis in a rural regional medical center. *Telemed J E Health.* 2015;21(3):157-62. doi: 10.1089/tmj.2014.0093. PMID: 25025239. Exclusion: Ineligible intervention.
642. Rincon F, Vibbert M, Childs V, et al. Implementation of a model of robotic telepresence (RTP) in the neuro-ICU: effect on critical care nursing team satisfaction. *Neurocrit Care.* 2012;17(1):97-101. doi: 10.1007/s12028-012-9712-2. PMID: 22547040. Exclusion: Ineligible population.
643. Ristevski E, Savage L, Taylor K, et al. Barriers and facilitators to implementing a cancer survivorship model of care in a rural area. *Asia Pac J Clin Oncol.* 2019;15(SUPPL 9):147. doi: 10.1111/ajco.13263. Exclusion: Ineligible intervention.
644. Roberts LJ, Lamont EG, Lim I, et al. Telerheumatology: an idea whose time has come. *Intern Med J.* 2012;42(10):1072-8. doi: 10.1111/j.1445-5994.2012.02931.x. PMID: 22931307. Exclusion: Not a study.
645. Robinson R. The evolution of perioperative telehealth in Katherine, Northern Territory, Australia. *J Perioper Nurs.* 2018;31(3):47-50. Exclusion: Ineligible intervention.

646. Rogers H, Madathil KC, Agnisarman S, et al. A systematic review of the implementation challenges of telemedicine systems in ambulances. *Telemed J E Health*. 2017;23(9):707-17. doi: 10.1089/tmj.2016.0248. PMID: 28294704. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
647. Rohatinsky N, Boyd I, Dickson A, et al. Perspectives of health care use and access to care for individuals living with inflammatory bowel disease in rural Canada. *Rural Remote Health*. 2021;21(2):6358. doi: 10.22605/RRH6358. PMID: 33820422. Exclusion: Ineligible intervention.
648. Rojas SM, Carter SP, McGinn MM, et al. A review of telemental health as a modality to deliver suicide-specific interventions for rural populations. *Telemed J E Health*. 2020;26(6):700-9. doi: 10.1089/tmj.2019.0083. PMID: 31502929. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
649. Rosborg S, Lindberg M, Ramukumba M, et al. Exploring mHealth's fit to workflow in homecare - a case study in Sweden. *Stud Health Technol Inform*. 2019;265:54-9. doi: 10.3233/SHTI190137. PMID: 31431577. Exclusion: Ineligible intervention.
650. Roy A, Das B, Garg P. Project ECHO (extension for community health care outcomes), an online tool for residents' education: a pilot study. *Indian J Ophthalmol*. 2020;68(10):2318-9. doi: 10.4103/ijo.IJO_588_20. PMID: 32971712. Exclusion: Ineligible population.
651. Rubin R. Using telemedicine to treat opioid use disorder in rural areas. *JAMA*. 2019;332(11):1029-31. doi: 10.1001/jama.2019.12574. PMID: 31461125. Exclusion: Not a study.
652. Rudd M, Rodgers H, Curless R, et al. Remote specialist assessment for intravenous thrombolysis of acute ischaemic stroke by telephone. *Emerg Med J*. 2012;29(9):704-8. doi: 10.1136/emmermed-2011-200582. PMID: 22034529. Exclusion: Ineligible population.
653. Ruggiano N, Brown EL, Li J, et al. Rural dementia caregivers and technology: what is the evidence? *Res Gerontol Nurs*. 2018;11(4):216-24. doi: 10.3928/19404921-20180628-04. PMID: 30036405. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
654. Ruiz-Perez I, Bastos A, Serrano-Ripoll MJ, et al. Effectiveness of interventions to improve cardiovascular healthcare in rural areas: a systematic literature review of clinical trials. *Prev Med*. 2019;119:132-44. doi: 10.1016/j.ypmed.2018.12.012. PMID: 30597226. Exclusion: Ineligible intervention.
655. Rush KL, Hatt L, Gorman N, et al. Planning telehealth for older adults with atrial fibrillation in rural communities: understanding stakeholder perspectives. *Clin Nurs Res*. 2019;28(2):130-49. doi: 10.1177/1054773818758170. PMID: 29460639. Exclusion: Ineligible outcome.
656. Sabesan S, Brennan S, Thaker D. Telemedicine for rural and indigenous cancer care. *Intern Med J*. 2011;41:14. doi: 10.1111/j.1445-5994.2011.02486.x. Exclusion: Not a study.
657. Sadaka F, Palagiri A, Trottier S, et al. Telemedicine intervention improves ICU outcomes. *Crit Care Res Pract*. 2013;2013:456389. doi: 10.1155/2013/456389. PMID: 23365729. Exclusion: Ineligible population.
658. Sagi MR, Aurobind G, Prabhat Chand, et al. Innovative telementoring for addiction management for remote primary care physicians: a feasibility study. *Indian J Psychiatry*. 2018;60(4):461-6. doi: 10.4103/psychiatry.IndianJPsychiatry_211_18. PMID: 30581211. Exclusion: Ineligible setting.

659. Sajwani A, Qureshi K, Shaikh T, et al. eHealth for remote regions: findings from Central Asia Health Systems Strengthening Project. *Stud Health Technol Inform.* 2015;209:128-34. PMID: 25980715. Exclusion: Ineligible setting.
660. Salami AC, Barden GM, Castillo DL, et al. Establishment of a Regional Virtual Tumor Board Program to Improve the Process of Care for Patients With Hepatocellular Carcinoma. *J Oncol Pract.* 2015;11(1):e66-74. doi: 10.1200/JOP.2014.000679. PMID: 25466708. Exclusion: Ineligible population.
661. Salcedo MP, Gowen R, Rodriguez AM, et al. Addressing high cervical cancer rates in the Rio Grande Valley along the Texas-Mexico border: a community-based initiative focused on education, patient navigation, and medical provider training/telementoring. *Perspect Public Health.* 2021;1757913921994610. doi: 10.1177/1757913921994610. PMID: 34130548. Exclusion: Ineligible comparater.
662. Saliba V, Legido-Quigley H, Hallik R, et al. Telemedicine across borders: a systematic review of factors that hinder or support implementation. *Int J Med Inf.* 2012;81(12):793-809. doi: 10.1016/j.ijmedinf.2012.08.003. PMID: 22975018. Exclusion: Ineligible setting.
663. Sankaranarayanan J, Murante LJ, Moffett LM. A retrospective evaluation of remote pharmacist interventions in a telepharmacy service model using a conceptual framework. *Telemed J E Health.* 2014;20(10):893-901. doi: 10.1089/tmj.2013.0362. PMID: 24611489. Exclusion: Ineligible comparater.
664. Sarkar S, Zepeda JB, Morris P, et al. Northern California hepatitis C elimination project: tele-mentoring led to quick adoption of hepatitis C treatment. *Hepatology.* 2020;72(1 SUPPL):399A. doi: 10.1002/hep.31579. Exclusion: Ineligible study design.
665. Sasaki R, Yamashita T, Omote Y, et al. A new telestroke network system in northern area of Okayama prefecture. *Neurol Clin Neurosci.* 2021;9(2):166-70. doi: 10.1111/ncn3.12475. Exclusion: Ineligible comparater.
666. Saurman E, Lyle D, Kirby S, et al. Use of a mental health emergency care-rural access programme in emergency departments. *J Telemed Telecare.* 2014;20(6):324-9. doi: 10.1177/1357633X14544422. PMID: 25052388. Exclusion: Ineligible outcome.
667. Savage SR, Youngren K, Kravic J, et al. COVID-19 and Project ECHO driving innovations in rural pain care & education. *Pain Med.* 2021;24:24. doi: 10.1093/pm/pnab191. PMID: 34169310. Exclusion: Not a study.
668. Schepens Niemiec SL, Vigen CLP, Martinez J, et al. Long-term follow-up of a lifestyle intervention for late-midlife, rural-dwelling Latinos in primary care. *Am J Occup Ther.* 2021;75(2):7502205020p1-p11. doi: 10.5014/ajot.2021.042861. PMID: 33657344. Exclusion: Ineligible intervention.
669. Schlemmer HP. Radiomics and AI: future of cancer imaging without radiologists? *Cancer Imaging.* 2018;18doi: 10.1186/s40644-018-0160-x. Exclusion: Ineligible intervention.
670. Schlittenhardt M, Smith SC, Ward-Smith P. Tele-continece care: a novel approach for providers. *Urol Nurs.* 2016;36(5):217-23. PMID: 29240333. Exclusion: Ineligible intervention.
671. Schoen JC, Russi CS, Laack TA. Addressing barriers to telemedicine use in rural emergency medicine: leveraging in situ simulation. *Telemed J E Health.* 2021;19:19. doi: 10.1089/tmj.2021.0026. PMID: 33872089. Exclusion: Ineligible intervention.
672. Scholes-Robertson NJ, Gutman TM, Tong A. Clinician perspectives on access to kidney replacement therapy in rural communities. *J Am Soc Nephrol.* 2020;31:379. Exclusion: Ineligible study design.

673. Schroeder C. Pilot study of telemedicine for the initial evaluation of general surgery patients in the clinic and hospitalized settings. *Surg Open Sci.* 2019;1(2):97-9. doi: 10.1016/j.sopen.2019.06.005. PMID: 32754700. Exclusion: Ineligible intervention.
674. Schwinn D. eNovation at UI health care. *Iowa Med.* 2015;105(2):24. PMID: 30074705. Exclusion: Not a study.
675. Scott A, Woodhouse LD, Watson D, et al. The Southeast Telehealth Network: using technology to overcome the barriers to rural public health practice. *J Public Health Manag Pract.* 2011;17(2):164-6. doi: 10.1097/PHH.0b013e3181f5413c. PMID: 21297412. Exclusion: Ineligible population.
676. Scott DM, Friesner DL, Rathke AM, et al. Differences in medication errors between central and remote site telepharmacies. *J Am Pharm Assoc (2003).* 2012;52(5):e97-e104. PMID: 23023864. Exclusion: Ineligible intervention.
677. Scott DM, Friesner DL, Rathke AM, et al. Medication error reporting in rural critical access hospitals in the North Dakota telepharmacy project. *Am J Health-Syst Pharm.* 2014;71(1):58-67. doi: 10.2146/ajhp120533. PMID: 24352183. Exclusion: Ineligible intervention.
678. Scott DM, Friesner DL, Udem T, et al. Perceived sustainability of community telepharmacy in North Dakota. *J Am Pharm Assoc (2003).* 2017;57(3):362-8.e5. doi: 10.1016/j.japh.2017.02.005. PMID: 28400253. Exclusion: Ineligible intervention.
679. Sekhon H, Sekhon K, Launay C, et al. Telemedicine and the rural dementia population: a systematic review. *Maturitas.* 2021;143:105-14. doi: 10.1016/j.maturitas.2020.09.001. PMID: 33308615. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
680. Sewell C, Jordan LA, Rudd J, et al. Implementation of a tele-thrombolysis service for acute stroke patients of the Manning Rural Referral Hospital: a quasi-experimental study. *Int J Stroke.* 2014;9:44. doi: 10.1111/ijss.12298. Exclusion: Not a study.
681. Shafiee Hanjani L, Bell JS, Freeman C. Undertaking medication review by telehealth. *Aust J Gen Pract.* 2020;49(12):826-31. doi: 10.31128/AJGP-06-20-5461. PMID: 33254216. Exclusion: Not a study.
682. Shafiee Hanjani L, Caffery LJ, Freeman CR, et al. A scoping review of the use and impact of telehealth medication reviews. *Res Social Adm Pharm.* 2020;16(8):1140-53. doi: 10.1016/j.sapharm.2019.12.014. PMID: 31874815. Exclusion: Ineligible intervention.
683. Sharma JJ, Gross G, Sharma P. Extending oncology clinical services to rural areas of Texas via teleoncology. *J Oncol Pract.* 2012;8(2):68. doi: 10.1200/JOP.2011.000436. PMID: 29447100. Exclusion: Not a study.
684. Shea CM, Teal R, Haynes-Maslow L, et al. Assessing the feasibility of a virtual tumor board program: a case study. *J Healthc Manag.* 2014;59(3):177-93. PMID: 24988672. Exclusion: Ineligible population.
685. Shea CM, Tabriz AA, Turner K, et al. Telestroke adoption among community hospitals in North Carolina: a cross-sectional study. *J Stroke Cerebrovasc Dis.* 2018;27(9):2411-7. doi: 10.1016/j.jstrokecerebrovasdis.2018.04.032. PMID: 29784607. Exclusion: Ineligible population.
686. Sherwood AR, MacDonald B. A teleneuropsychology consultation service model for children with neurodevelopmental and acquired disorders residing in rural state regions. *Arch Clin Neuropsychol.* 2020;35(8):1196-203. doi: 10.1093/arclin/acia099. PMID: 33124658. Exclusion: Ineligible intervention.

687. Shimasaki S, Bishop E, Guthrie M, et al. Strengthening the health workforce through the ECHO stages of participation: participants' perspectives on key facilitators and barriers. *J Med Educ Curric Dev.* 2019;6:1-8. doi: 10.1177/2382120518820922. PMID: 30729170. Exclusion: Ineligible population.
688. Shimizu S, Nagai E, Torata N, et al. Global remote education of endoscopic surgery using a worldwide super-fast internet: growing demands in Europe. *Surg Endosc.* 2011;25:S15. doi: 10.1007/s00464-011-1602-8. Exclusion: Ineligible population.
689. Shook LM, Farrell CB, Kalinyak KA, et al. Translating sickle cell guidelines into practice for primary care providers with Project ECHO. *Med Educ Online.* 2016;21:33616. doi: 10.3402/meo.v21.33616. PMID: 27887664. Exclusion: Ineligible population.
690. Shore JH, Thurman MT, Fujinami L, et al. A resident, rural telepsychiatry service: training and improving care for rural populations. *Acad Psychiatry.* 2011;35(4):252-5. doi: 10.1176/appi.ap.35.4.252. PMID: 21804046. Exclusion: Ineligible intervention.
691. Shuaib A, Jeerakathil T, Alberta Mobile Stroke Unit I. The mobile stroke unit and management of acute stroke in rural settings. *CMAJ.* 2018;190(28):E855-E8. doi: 10.1503/cmaj.170999. PMID: 30012801. Exclusion: Not a study.
692. Shura RD, Brearly TW, Tupler LA. Telehealth in response to the COVID-19 pandemic in rural veteran and military beneficiaries. *J Rural Health.* 2020:1-5. doi: 10.1111/jrh.12454. PMID: 32402128. Exclusion: Not a study.
693. Simpson SG, Rochford S, Livingstone A, et al. Tele-web psychology in rural South Australia: the logistics of setting up a remote university clinic staffed by clinical psychologists in training. *Aust Psychol.* 2014;49(4):193-9. doi: 10.1111/ap.12049. Exclusion: Ineligible intervention.
694. Singh G, Pal US, Mishra M, et al. Teleassistance and teleconsultation using smartphones and its contribution in clinical progress of oral and maxillofacial surgery. *Natl J Maxillofac Surg.* 2020;11(1):10-3. doi: 10.4103/njms.NJMS_9_18. PMID: 33041570. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
695. Singh H, McGregor JC, Quale CL, et al. Remote specialty therapeutic management enhances patient-centered heart failure care in a rural community setting. *J Card Fail.* 2015;21(8):S61. Exclusion: Ineligible study design.
696. Singh NN, Chan J, Karazsia BT, et al. Tele-health training of teachers to teach a mindfulness-based procedure for self-management of aggressive behavior to students with intellectual and developmental disabilities. *Int J Dev Disabil.* 2017;63(4):195-203. doi: 10.1080/20473869.2016.1277841. Exclusion: Ineligible population.
697. Singh TP, Vangaveti VN, Kennedy RL, et al. Role of telehealth in diabetic foot ulcer management - a systematic review. *Aust J Rural Health.* 2016;24(4):224-9. doi: 10.1111/ajr.12284. PMID: 27098271. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
698. Sinha N, Cornell M, Wheatley B, et al. Looking through a different lens: patient satisfaction with telemedicine in delivering pediatric fracture care. *J Am Acad Orthop Surg Glob Res Rev.* 2019;3(9):e100. doi: 10.5435/JAAOSGlobal-D-19-00100. PMID: 31773080. Exclusion: Ineligible intervention.
699. Slade CP, O'Toole LJ, Jr., Rho E. State primary stroke center policies in the United States: rural health issues. *Telemed J E Health.* 2012;18(3):225-9. doi: 10.1089/tmj.2011.0141. PMID: 22356528. Exclusion: Ineligible outcome.

700. Slivinski A, Jones R, Whitehead H, et al. Improving access to stroke care in the rural setting: the journey to acute stroke ready designation. *J Emerg Nurs.* 2017;43(1):24-32. doi: 10.1016/j.jen.2016.10.006. PMID: 28131346. Exclusion: Ineligible intervention.
701. Smith-Strøm H, Igland J, Østbye T, et al. The Effect of Telemedicine Follow-up Care on Diabetes-Related Foot Ulcers: A Cluster-Randomized Controlled Noninferiority Trial. *Diabetes Care.* 2018;41(1):96-103. doi: 10.2337/dc17-1025. PMID: 29187423. Exclusion: Ineligible population.
702. Smith D, Johnston K, Carlisle K, et al. Client perceptions of the BreastScreen Australia remote radiology assessment model. *BMC Womens Health.* 2021;21(1):30. doi: 10.1186/s12905-020-01163-7. PMID: 33461562. Exclusion: Ineligible outcome.
703. Smith MA, Smith WT, Stanton M. Universal postoperative hip instruction protocol for rehabilitation in rural skilled nursing facilities. *Prof Case Manag.* 2015;20(5):241-7. doi: 10.1097/NCM.000000000000107. PMID: 26241624. Exclusion: Ineligible intervention.
704. Smith NM, Satyshur RD. Pediatric diabetes telemedicine program improves access to care for rural families: role of APRNs. *Pediatr Nurs.* 2016;42(6):294-9. PMID: 29406653. Exclusion: Ineligible intervention.
705. Snell-Rood C, Pollini RA, Willging C. Barriers to integrated medication-assisted treatment for rural patients with co-occurring disorders: the gap in managing addiction. *Psychiatr Serv.* 2021:appips202000312. doi: 10.1176/appi.ps.202000312. PMID: 33530734. Exclusion: Ineligible intervention.
706. Snoswell CL, North JB, Caffery LJ. Economic advantages of telehealth and virtual health practitioners: return on investment analysis. *JMIR Perioper Med.* 2020;3(1):e15688. doi: 10.2196/15688. PMID: 33393922. Exclusion: Ineligible intervention.
707. Sockalingam S, Kirvan A, Pereira C, et al. The role of tele-education in advancing mental health quality of care: a content analysis of Project ECHO recommendations. *Telemed J E Health.* 2021;27(8):939-46. doi: 10.1089/tmj.2021.0122. PMID: 34227890. Exclusion: Ineligible outcome.
708. Sockalingam S, Rajaratnam T, Zhou C, et al. Building mental health capacity: exploring the role of adaptive expertise in the ECHO virtual learning model. *J Contin Educ Health Prof.* 2021;41(2):104-10. doi: 10.1097/CEH.0000000000000349. PMID: 34009840. Exclusion: Ineligible study design.
709. Sood A, Granick MS, Trial C, et al. The role of telemedicine in wound care: a review and analysis of a database of 5,795 patients from a mobile wound-healing center in Languedoc-Roussillon, France. *Plast Reconstr Surg.* 2016;138(3 Suppl):248S-56S. doi: 10.1097/PRS.0000000000002702. PMID: 27556769. Exclusion: Ineligible study design.
710. Sood A, Pollard C, Suer KL, et al. Caring for miners during the coronavirus disease-2019 (COVID-19) pandemic. *J Rural Health.* 2021;37(1):165-8. doi: 10.1111/jrh.12444. PMID: 32277775. Exclusion: Not a study.
711. Southard EP, Neufeld JD, Laws S. Telemental health evaluations enhance access and efficiency in a critical access hospital emergency department. *Telemed J E Health.* 2014;20(7):664-8. doi: 10.1089/tmj.2013.0257. PMID: 24811858. Exclusion: Ineligible intervention.
712. Sperring B. Using telehealth and photography for wound assessment in Western Australia. *Wounds International.* 2013;4(4):26-9. Exclusion: Not a study.

713. Srisuk N, Cameron J, Ski CF, et al. Randomized controlled trial of family-based education for patients with heart failure and their carers. *J Adv Nurs*. 2017;73(4):857-70. doi: 10.1111/jan.13192. PMID: 27779768. Exclusion: Ineligible intervention.
714. Stading JA, Phan L, Walter A, et al. Initial experience of clinical pharmacy services delivered by computer communication via Cisco Jabber video in a US Veterans Administration medical center. *J Pharm Tech*. 2014;30(3):76-80. doi: 10.1177/8755122513505744. Exclusion: Ineligible study design.
715. Stampehl MR, Jennison SH, Call C, et al. Heart failure care management program utilizing telehealth technology yields readmission rates lower than the state average. *J Card Fail*. 2017;23(8):S80. Exclusion: Ineligible comparater.
716. Steeves-Reece AL, Elder NC, Graham TA, et al. Rapid deployment of a statewide COVID-19 ECHO program for frontline clinicians: early results and lessons learned. *J Rural Health*. 2020;37(1):227-30. doi: 10.1111/jrh.12462. PMID: 32396224. Exclusion: Ineligible study design.
717. Steinman M, Morbeck RA, Pires PV, et al. Impact of telemedicine in hospital culture and its consequences on quality of care and safety. *Einstein (Sao Paulo)*. 2015;13(4):580-6. doi: 10.1590/S1679-45082015GS2893. PMID: 26676268. Exclusion: Ineligible population.
718. Steitz BD, Levy MA. Evaluating the scope of clinical electronic messaging to coordinate care in a breast cancer cohort. *Stud Health Technol Inform*. 2019;264:808-12. doi: 10.3233/SHTI190335. PMID: 31438036. Exclusion: Ineligible study design.
719. Stephens D, Leston J, Terrault NA, et al. An evaluation of hepatitis C virus telehealth services serving tribal communities: patterns of usage, evolving needs, and barriers. *J Public Health Manag Pract*. 2019;25 Suppl 5, Tribal Epidemiology Centers:Advancing Public Health in Indian Country for Over 20 Years:S97-S100. doi: 10.1097/PHH.0000000000001061. PMID: 31348196. Exclusion: Ineligible study design.
720. Sterling SA, Jones AE, King MH, et al. The impact of the TelEmergency program on rural emergency care: an implementation study. *Ann Emerg Med*. 2016;68:S95-S. doi: 10.1016/j.annemergmed.2016.08.259. Exclusion: Not a study.
721. Stern A, Mitsakakis N, Paulden M, et al. Pressure ulcer multidisciplinary teams via telemedicine: a pragmatic cluster randomized stepped wedge trial in long term care. *BMC Health Serv Res*. 2014;14:83. doi: 10.1186/1472-6963-14-83. PMID: 24559218. Exclusion: Ineligible population.
722. Stevenson V, Keenan E, Nutman K, et al. Telemedicine a useful tool to link people with complex disabilities to specialist neurological services. *J Neurol Neurosurg Psychiatry*. 2010;81(11):e42. doi: 10.1136/jnnp.2010.226340.100. Exclusion: Not a study.
723. Steventon A, Ariti C, Fisher E, et al. Effect of telehealth on hospital utilisation and mortality in routine clinical practice: a matched control cohort study in an early adopter site. *BMJ Open*. 2016;6(2):e009221. doi: 10.1136/bmjopen-2015-009221. PMID: 26842270. Exclusion: Ineligible intervention.
724. Stewart M, Marshall JN, Ostbye T, et al. Effectiveness of case-based on-line learning of evidence-based practice guidelines. *Fam Med*. 2005;37(2):131-8. PMID: 15690254. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).

725. Stewart RW, Whiteman LN, Strouse JJ, et al. Improving inpatient care for individuals with sickle cell disease using the Project ECHO model. *South Med J*. 2016;109(9):568-9. doi: 10.14423/SMJ.0000000000000509. PMID: 27598364. Exclusion: Not a study.
726. Stickland M, Jourdain T, Wong EY, et al. Using telehealth technology to deliver pulmonary rehabilitation in chronic obstructive pulmonary disease patients. *Can Respir J*. 2011;18(4):216-20. PMID: 22059179. Exclusion: Ineligible intervention.
727. Stigler Granados P, Pacheco GJ, Nunez Patlan E, et al. Assessing the effectiveness of Chagas disease education for healthcare providers in the United States. *BMC Infect Dis*. 2020;20(1):743. doi: 10.1186/s12879-020-05474-w. PMID: 33036559. Exclusion: Ineligible population.
728. Stingley S, Schultz H. Helmsley trust support for telehealth improves access to care in rural and frontier areas. *Health Aff (Millwood)*. 2014;33(2):336-41. doi: 10.1377/hlthaff.2013.1278. PMID: 24493777. Exclusion: Not a study.
729. Streicher JL, Kini SP, Stoff BK. Innovative dermatopathology teaching in a resource-limited environment. *J Am Acad Dermatol*. 2016;74(5):1024-5. doi: 10.1016/j.jaad.2015.11.010. PMID: 27085239. Exclusion: Ineligible setting.
730. Stuckey R, Domingues-Montanari S. Telemedicine is helping the parents of children with neurodevelopmental disorders living in remote and deprived areas. *Paediatr Int Child Health*. 2017;37(3):155-7. doi: 10.1080/20469047.2017.1315914. PMID: 28498062. Exclusion: Not a study.
731. Su GL, Glass L, Tapper EB, et al. Virtual consultations through the Veterans Administration SCAN-ECHO Project improves survival for veterans with liver disease. *Hepatology*. 2018;68(6):2317-24. doi: 10.1002/hep.30074. PMID: 29729194. Exclusion: Ineligible population.
732. Surdu S, Langelier M. Teledentistry: increasing utilisation of oral-health services for children in rural areas. *J Telemed Telecare*. 2020;1357633X20965425. doi: 10.1177/1357633X20965425. PMID: 33070687. Exclusion: Ineligible comparater.
733. Sutherland JE, Sutphin HD, Rawlins F, et al. A comparison of telesonography with standard ultrasound care in a rural Dominican clinic. *J Telemed Telecare*. 2009;15(4):191-5. PMID: 19471031. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
734. Sykora R, Renza M, Ruzicka J, et al. Audiovisual consults by paramedics to reduce hospital transport after low-urgency calls: randomized controlled trial. *Prehospital Disaster Med*. 2020;35(6):656-62. doi: 10.1017/S1049023X2000117X. PMID: 32985403. Exclusion: Ineligible comparater.
735. Tahan V, Almashhrawi A, Mutrux R, et al. Show Me ECHO-Hepatitis C: a telemedicine mentoring program for patients with hepatitis C in underserved and rural areas in Missouri as a model in developing countries. *Turk J Gastroenterol*. 2015;26(6):447-9. doi: 10.5152/tjg.2015.159000. PMID: 26510085. Exclusion: Not a study.
736. Tan E, Gao L, Tran HN, et al. Telestroke for acute ischaemic stroke: a systematic review of economic evaluations and a de novo cost-utility analysis for a middle income country. *J Telemed Telecare*. 2021;1357633X211032407. doi: 10.1177/1357633X211032407. PMID: 34292801. Exclusion: Ineligible population.
737. Tarver WL, Haggstrom DA. The use of cancer-specific patient-centered technologies among underserved populations in the United States: systematic review. *J Med Internet Res*. 2019;21(4):e10256. doi: 10.2196/10256. PMID: 31012855. Exclusion: Ineligible intervention.

738. Taylor G. Mhealth and video conferencing: better care in rural North Carolina. *Behav Healthc*. 2013;33(5):47-8. PMID: 24298707. Exclusion: Not a study.
739. Taylor L, Waller M, Portnoy JM. Telemedicine for allergy services to rural communities. *J Allergy Clin Immunol Pract*. 2019;7(8):2554-9. doi: 10.1016/j.jaip.2019.06.012. PMID: 31238163. Exclusion: Ineligible intervention.
740. Taylor MA, Knochel ML, Proctor SJ, et al. Pediatric trauma telemedicine in a rural state: lessons learned from a 1-year experience. *J Pediatr Surg*. 2021;56(2):385-9. doi: 10.1016/j.jpedsurg.2020.10.020. PMID: 33228973. Exclusion: Ineligible intervention.
741. Tchero H, Tabue Teguo M, Lannuzel A, et al. Telerehabilitation for stroke survivors: systematic review and meta-analysis. *J Med Internet Res*. 2018;20(10):e10867. doi: 10.2196/10867. PMID: 30368437. Exclusion: Ineligible intervention.
742. Terry B, Polan DL, Nambaziira R, et al. Rapid, remote education for point-of-care ultrasound among non-physician emergency care providers in a resource limited setting. *Afr J Emerg Med*. 2019;9(3):140-4. doi: 10.1016/j.afjem.2019.05.004. PMID: 31528532. Exclusion: Ineligible outcome.
743. Tesfalul M, Littman-Quinn R, Antwi C, et al. Evaluating the impact of a mobile oral telemedicine system on medical management and clinical outcomes of patients with complicated oral lesions in Botswana. *Stud Health Technol Inform*. 2013;192:1074. PMID: 23920848. Exclusion: Ineligible setting.
744. Tetuan CE, Guthrie KD, Stoner SC, et al. Impact of community pharmacist-performed post-discharge medication reviews in transitions of care. *J Am Pharm Assoc (2003)*. 2018;58(6):659-66. doi: 10.1016/j.japh.2018.06.017. PMID: 30056133. Exclusion: Ineligible intervention.
745. Thakur A, Pereira C, Hardy J, et al. Virtual education program to support providers caring for people with intellectual and developmental disabilities during the COVID-19 pandemic: rapid development and evaluation study. *JMIR Ment Health*. 2021;8(10):e28933. doi: 10.2196/28933. PMID: 34617917. Exclusion: Ineligible study design.
746. Thies KM, Anderson D, Beals-Reid C. Project ECHO chronic pain: a qualitative analysis of recommendations by expert faculty and the process of knowledge translation. *J Contin Educ Health Prof*. 2019;39(3):194-200. doi: 10.1097/CEH.0000000000000264. PMID: 31385920. Exclusion: Ineligible population.
747. Thies KM, Gonzalez M, Porto A, et al. Project ECHO COVID-19: vulnerable populations and telehealth early in the pandemic. *J Prim Care Community Health*. 2021;12:21501327211019286. doi: 10.1177/21501327211019286. PMID: 34036832. Exclusion: Ineligible outcome.
748. Thijssing L, van der Heijden JP, Chavannes NH, et al. Telepulmonology: effect on quality and efficiency of care. *Respir Med*. 2014;108(2):314-8. doi: 10.1016/j.rmed.2013.10.017. PMID: 24210893. Exclusion: Ineligible population.
749. Thinyane H, Hansen S, Foster G, et al. Using mobile phones for rapid reporting of zoonotic diseases in rural South Africa. *Stud Health Technol Inform*. 2010;161:179-89. PMID: 21191171. Exclusion: Ineligible study design.
750. Thomas J, Tedder A, Levitt JM, et al. Integration of remote pharmacist telehealth service into 3 distant, rural and underserved primary care practices. *J Am Coll Clin Pharm*. 2021;4(9):1245. doi: 10.1002/jac5.1481. Exclusion: Not a study.

751. Thomson MD, Mariani AC, Williams AR, et al. Factors associated with use of and satisfaction with telehealth by adults in rural Virginia during the COVID-19 pandemic. *JAMA Netw Open*. 2021;4(8):e2119530. doi: 10.1001/jamanetworkopen.2021.19530. PMID: 34351404. Exclusion: Ineligible intervention.
752. Timonen O. The teleconsultation in general practice. A randomized, controlled study of a remote consultation experiment using a videoconferencing system. *Int J Circumpolar Health*. 2004;63(3):289-90. PMID: 15526933. Exclusion: Not a study.
753. Ting L, Wilkes M. Telemedicine for patient management on expeditions in remote and austere environments: a systematic review. *Wilderness Environ Med*. 2021;32(1):102-11. doi: 10.1016/j.wem.2020.09.009. PMID: 33423896. Exclusion: Ineligible setting.
754. Tiyyagura G, Asnes AG, Leventhal JM, et al. Impact of Project ECHO on community ED providers' perceptions of child abuse knowledge and access to subspecialists for child abuse and neglect. *Acad Pediatr*. 2019;19(8):985-7. doi: 10.1016/j.acap.2019.06.018. PMID: 31299431. Exclusion: Ineligible population.
755. Tofighi B, Isaacs N, Byrnes-Enoch H, et al. Expanding treatment for opioid use disorder in publicly funded primary care clinics: exploratory evaluation of the NYC health+hospitals buprenorphine ECHO program. *J Subst Abuse Treat*. 2019;106:1-3. doi: 10.1016/j.jsat.2019.08.003. PMID: 31540604. Exclusion: Ineligible population.
756. Tokuda L, Lorenzo L, Theriault A, et al. The utilization of video-conference shared medical appointments in rural diabetes care. *Int J Med Inf*. 2016;93:34-41. doi: 10.1016/j.ijmedinf.2016.05.007. PMID: 27435945. Exclusion: Ineligible intervention.
757. Toresdahl BG, Young WK, Quijano B, et al. A systematic review of telehealth and sport-related concussion: baseline testing, diagnosis, and management. *HSS J*. 2021;17(1):18-24. doi: 10.1177/1556331620975856. PMID: 33967637. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
758. Torres Diaz A, Lock LJ, Molfenter TD, et al. Implementation for Sustained Impact in Teleophthalmology (I-SITE): applying the NIATx Model for tailored implementation of diabetic retinopathy screening in primary care. *Implement Sci Commun*. 2021;2(1):74. doi: 10.1186/s43058-021-00175-0. PMID: 34229748. Exclusion: Ineligible intervention.
759. Toscano M, Marini TJ, Baran TM, et al. 69 Testing a teliagnostic obstetric ultrasound system without a trained sonographer in a low-resource Peruvian clinic. *Am J Obstet Gynecol*. 2021;224(2):S48-S9. doi: 10.1016/j.ajog.2020.12.068. Exclusion: Ineligible study design.
760. Toscano M, Marini TJ, Drennan K, et al. Testing teliagnostic obstetric ultrasound in Peru: a new horizon in expanding access to prenatal ultrasound. *BMC Pregnancy Childbirth*. 2021;21(1):328. doi: 10.1186/s12884-021-03720-w. PMID: 33902496. Exclusion: Ineligible study design.
761. Totten AM, Womack DM, Eden KB, et al. Telehealth: mapping the evidence for patient outcomes from systematic reviews - technical brief No. 26 Pacific Northwest Evidence-based Practice Center. Rockville, MD: June 2016. https://effectivehealthcare.ahrq.gov/sites/default/files/pdf/telehealth_technical-brief.pdf. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.

762. Totten AM, Hansen RN, Wagner J, et al. Telehealth for acute and chronic care consultations. AHRQ Comparative Effectiveness Review No. 216. (Prepared by Pacific Northwest Evidence-based Practice Center under Contract No. 290-2015-00009-I.) AHRQ Publication No. 19-EHC012-EF. Rockville, MD: Agency for Healthcare Research and Quality. 2019 PMID: 31577401. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
763. Touchett H, Chen P, Lindsay J, et al. Supporting telemental health for rural disabled veterans: bringing rehabilitation expertise to rural mental health providers. *Arch Phys Med Rehabil*. 2021;102(10):e47. doi: 10.1016/j.apmr.2021.07.601. Exclusion: Not a study.
764. Traube DE, Cederbaum JA, Taylor A, et al. Telehealth Training and Provider Experience of Delivering Behavioral Health Services. *J Behav Health Serv Res*. 2021;48(1):93-102. doi: 10.1007/s11414-020-09718-0. PMID: 32596804. Exclusion: Ineligible intervention.
765. Tuckson RV, Edmunds M, Hodgkins ML. Telehealth. *N Engl J Med*. 2017;377(16):1585-92. doi: 10.1056/NEJMSr1503323. PMID: 29045204. Exclusion: Not a study. Has some nice telehealth graphics could be used a background info from NEJM
766. Tuot DS, Leeds K, Murphy EJ, et al. Facilitators and barriers to implementing electronic referral and/or consultation systems: a qualitative study of 16 health organizations. *BMC Health Serv Res*. 2015;15:568. doi: 10.1186/s12913-015-1233-1. PMID: 26687507. Exclusion: Ineligible population.
767. Turner N, Debenham E, Sabatino P, et al. Telestroke consultations in an EMS unit to reduce t-PA decision times. *Stroke*. 2019;50(Supplement 1):AWP481. doi: 10.1161/str.50.suppl_1.WP481. Exclusion: Not a study.
768. Uscher-Pines L, Kahn JM. Barriers and facilitators to pediatric emergency telemedicine in the United States. *Telemed J E Health*. 2014;20(11):990-6. doi: 10.1089/tmj.2014.0015. PMID: 25238565. Exclusion: Ineligible population.
769. Usher R, Payne C, Real S, et al. Project ECHO: Enhancing palliative care for primary care occupational therapists and physiotherapists in Ireland. *Health Soc Care Community*. 2021;15:15. doi: 10.1111/hsc.13372. PMID: 33991147. Exclusion: Ineligible setting.
770. van Gelder VA, Scherpier-de Haan ND, van Berkel S, et al. Web-based consultation between general practitioners and nephrologists: a cluster randomized controlled trial. *Fam Pract*. 2017;34(4):430-6. doi: 10.1093/fampra/cmww131. PMID: 28158576. Exclusion: Ineligible population.
771. Van Oeveren L, Donner J, Fantegrossi A, et al. Telemedicine-assisted intubation in rural emergency departments: a national emergency airway registry study. *Telemed J E Health*. 2017;23(4):290-7. doi: 10.1089/tmj.2016.0140. PMID: 27673565. Exclusion: Ineligible study design.
772. Varon ML, Baker E, Estrada EE, et al. Advancing gynecologic oncology in Latin America through Project ECHO. *Int J Gynecol Cancer*. 2020;30(11):1840-1. doi: 10.1136/ijgc-2020-001559. PMID: 32522773. Exclusion: Not a study.
773. Varon ML, Baker E, Byers E, et al. Project ECHO Cancer Initiative: a Tool to Improve Care and Increase Capacity Along the Continuum of Cancer Care. *J Cancer Educ*. 2021;36(Suppl 1):25-38. doi: 10.1007/s13187-021-02031-0. PMID: 34292501. Exclusion: Not a study.
774. Vaskinn A, Wilsøgard I, Holm A, et al. A feasibility study of a telephone-based screening service for mild cognitive impairment and its uptake by elderly people. *J Telemed Telecare*. 2013;19(1):5-10. doi: 10.1177/1357633X12473904. PMID: 23390214. Exclusion: Ineligible intervention.

775. Vaughan EM, Hyman DJ, Naik AD, et al. A Telehealth-supported, Integrated care with CHWs, and MEDication-access (TIME) program for diabetes improves HbA1c: a randomized clinical trial. *J Gen Intern Med*. 2021;36(2):455-63. doi: 10.1007/s11606-020-06017-4. PMID: 32700217. Exclusion: Ineligible population.
776. Vega S, Marciscano I, Holcomb M, et al. Testing a top-down strategy for establishing a sustainable telemedicine program in a developing country: the Arizona Telemedicine Program-US Army-Republic of Panama Initiative. *Telemed J E Health*. 2013;19(10):746-53. doi: 10.1089/tmj.2013.0025. PMID: 23931731. Exclusion: Ineligible population.
777. Venuthurupalli SK, Rolfe A, Fanning J, et al. Chronic Kidney Disease, Queensland (CKD.QLD) Registry: management of CKD With telenephrology. *Kidney Int Rep*. 2018;3(6):1336-43. doi: 10.1016/j.ekir.2018.07.013. PMID: 30450460. Exclusion: Ineligible intervention.
778. Versleijen M, Martin-Khan MG, Whitty JA, et al. A telegeriatric service in a small rural hospital: a case study and cost analysis. *J Telemed Telecare*. 2015;21(8):459-68. doi: 10.1177/1357633X15611327. PMID: 26556059. Exclusion: Ineligible intervention.
779. Vimalananda VG, Gupte G, Seraj SM, et al. Electronic consultations (e-consults) to improve access to specialty care: a systematic review and narrative synthesis. *J Telemed Telecare*. 2015;21(6):323-30. doi: 10.1177/1357633X15582108. PMID: 25995331. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
780. Vinayak S, Sande J, Nisenbaum H, et al. Training midwives to perform basic obstetric point-of-care ultrasound in rural areas using a tablet platform and mobile phone transmission technology-A WFUMB COE project. *Ultrasound Med Biol*. 2017;43(10):2125-32. doi: 10.1016/j.ultrasmedbio.2017.05.024. PMID: 28716434. Exclusion: Ineligible setting.
781. Vinson AH, Iannuzzi D, Bennett A, et al. Facilitator Reflections on Shared Expertise and Adaptive Leadership in ECHO Autism: Center Engagement. *J Contin Educ Health Prof*. 2021;01:01. doi: 10.1097/CEH.0000000000000395. PMID: 34609351. Exclusion: Ineligible comparater.
782. Vo AT, Gustafson DL. Telepharmacy in oncology care: A scoping review. *J Telemed Telecare*. 2020;1357633X20975257. doi: 10.1177/1357633X20975257. PMID: 33377820. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
783. Vyas KS, Hambrick HR, Shakir A, et al. A systematic review of the use of telemedicine in plastic and reconstructive surgery and dermatology. *Ann Plast Surg*. 2017;78(6):736-68. doi: 10.1097/SAP.0000000000001044. PMID: 28328635. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
784. Wachs JP, Kirkpatrick AW, Tisherman SA. Procedural Telementoring in Rural, Underdeveloped, and Austere Settings: Origins, Present Challenges, and Future Perspectives. *Annu Rev Biomed Eng*. 2021;26:26. doi: 10.1146/annurev-bioeng-083120-023315. PMID: 33770455. Exclusion: Not a study.
785. Wade VA, Karnon J, Elshaug AG, et al. A systematic review of economic analyses of telehealth services using real time video communication. *BMC Health Serv Res*. 2010;10:233. doi: 10.1186/1472-6963-10-233. PMID: 20696073. Exclusion: Ineligible intervention.

786. Wadhwani C, Adomako J. Ghana telemedicine - using mHealth to bridge the gaps in health service availability and accessibility. *Trop Med Int Health*. 2015;20:131-2. doi: 10.1111/tmi.12575. Exclusion: Ineligible setting.
787. Wakefield DV, Sanders T, Wilson E, et al. Initial impact and operational responses to the COVID-19 pandemic by American radiation oncology practices. *Int J Radiat Oncol Biol Phys*. 2020;108(2):356-61. doi: 10.1016/j.ijrobp.2020.06.060. PMID: 32890512. Exclusion: Ineligible intervention.
788. Walensky RP, McQuillen DP, Shahbazi S, et al. Where is the ID in COVID-19? *Ann Intern Med*. 2020;1-3. doi: 10.7326/M20-2684. PMID: 32491920. Exclusion: Not a study.
789. Walker AF, Cuttriss N, Haller MJ, et al. Democratizing type 1 diabetes specialty care in the primary care setting to reduce health disparities: project extension for community healthcare outcomes (ECHO) T1D. *BMJ Open Diabetes Res Care*. 2021;9(1):07. doi: 10.1136/bmjdr-2021-002262. PMID: 34244218. Exclusion: Ineligible population.
790. Wallace P, Haines A, Harrison R, et al. Design and performance of a multi-centre randomised controlled trial and economic evaluation of joint tele-consultations. *BMC Fam Pract*. 2002;3(1)doi: 10.1186/1471-2296-3-1. PMID: 11835692. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
791. Walsh C, Siegler EL, Cheston E, et al. Provider-to-provider electronic communication in the era of meaningful use: a review of the evidence. *J Hosp Med*. 2013;8(10):589-97. doi: 10.1002/jhm.2082. PMID: 24101544. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
792. Walters LEM, Scott RE, Mars M. A teledermatology scale-up framework and roadmap for sustainable scaling: evidence-based development. *J Med Internet Res*. 2018;20(6):e224. doi: 10.2196/jmir.9940. PMID: 29925492. Exclusion: Ineligible study design.
793. Wang S, Parsons M, Stone-McLean J, et al. Augmented reality as a telemedicine platform for remote procedural training. *Sensors (Basel)*. 2017;17:1-21. doi: 10.3390/s17102294. PMID: 28994720. Exclusion: Ineligible setting.
794. Ward MM, Jaana M, Natafagi N. Systematic review of telemedicine applications in emergency rooms. *Int J Med Inf*. 2015;84(9):601-16. doi: 10.1016/j.ijmedinf.2015.05.009. PMID: 26072326. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
795. Warshaw E, Greer N, Hillman Y, et al. Tele dermatology for Diagnosis and Management of Skin Conditions: A Systematic Review of the Evidence. Jan 01 2010. <https://www.hsrd.research.va.gov/publications/esp/telederm-2010.pdf> PMID: 21155201. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
796. Watson M. Project ECHO director responds to concerns. *BMJ*. 2019;366:l5157. doi: 10.1136/bmj.l5157. PMID: 31439623. Exclusion: Not a study.
797. Waugh M, Calderone J, Brown Levey S, et al. Using telepsychiatry to enrich existing integrated primary care. *Telemed J E Health*. 2019;25(8):762-8. doi: 10.1089/tmj.2018.0132. PMID: 30394851. Exclusion: Ineligible population.
798. Wearne S. Remote supervision during general practice training. *Educ Prim Care*. 2016;27(4):333-4. doi: 10.1080/14739879.2016.1207103. PMID: 27412381. Exclusion: Not a study.

799. Weaver DT, Murdock TJ. Telemedicine detection of type 1 ROP in a distant neonatal intensive care unit. *J AAPOS*. 2012;16(3):229-33. doi: 10.1016/j.jaapos.2012.01.007. PMID: 22681938. Exclusion: Ineligible intervention.
800. Wechsler LR, Tsao JW, Levine SR, et al. Teleneurology applications: report of the Telemedicine Work Group of the American Academy of Neurology. *Neurology*. 2013;80(7):670-6. doi: 10.1212/WNL.0b013e3182823361. PMID: 23400317. Exclusion: Not a study.
801. Wechsler LR. Advantages and limitations of teleneurology. *JAMA Neurol*. 2015;72(3):349-54. doi: 10.1001/jamaneurol.2014.3844. PMID: 25580942. Exclusion: Not a study.
802. Weinstock RS, Izquierdo R, Goland R, et al. Improved glycemic control in older adults with diabetes with telemedicine: 5-year results from the IDEATel study. *Diabetes*. 2010. Exclusion: Not a study.
803. Weintraub E, Greenblatt AD, Chang J, et al. Outcomes for patients receiving telemedicine-delivered medication-based treatment for Opioid Use Disorder: A retrospective chart review. *Heroin Addict Relat Clin Probl*. 2021;23(2):5-12. PMID: 33551692. Exclusion: Ineligible intervention.
804. Wessels LE, Roper MT, Ignacio RC, et al. Telementorship in Underway Naval Operations: Leveraging Operational Virtual Health for Tactical Combat Casualty Care. *J Spec Oper Med*. 2021;21(3):93-5. PMID: 34529812. Exclusion: Ineligible intervention.
805. West SP, Lagua C, Trief PM, et al. Goal setting using telemedicine in rural underserved older adults with diabetes: experiences from the informatics for diabetes education and telemedicine project. *Telemed J E Health*. 2010;16(4):405-16. doi: 10.1089/tmj.2009.0136. PMID: 20507198. Exclusion: Ineligible intervention.
806. Whetten J, van der Goes DN, Tran H, et al. Cost-effectiveness of Access to Critical Cerebral Emergency Support Services (ACCESS): a neuro-emergent telemedicine consultation program. *J Med Econ*. 2018;21(4):398-405. doi: 10.1080/13696998.2018.1426591. PMID: 29316820. Exclusion: Ineligible study design.
807. Whitacre BE, Wheeler D, Landgraf C. What can the national broadband map tell us about the health care connectivity gap? *J Rural Health*. 2017;33(3):284-9. doi: 10.1111/jrh.12177. PMID: 26934373. Exclusion: Ineligible study design.
808. White C, McIlpatrick S, Dunwoody L, et al. Supporting and improving community health services-a prospective evaluation of ECHO technology in community palliative care nursing teams. *BMJ Support Palliat Care*. 2019;9(2):202-8. doi: 10.1136/bmjspcare-2015-000935. PMID: 26628535. Exclusion: Ineligible population.
809. White H. Practice-based or telemedicine-based collaborative care for depression. *Am J Psychiatry*. 2013;170(8):926. doi: 10.1176/appi.ajp.2013.13040471. PMID: 23903336. Exclusion: Not a study.
810. Whiteman C, Kiefer C, D'Angelo J, et al. The use of technology to reduce radiation exposure in trauma patients transferred to a level I trauma center. *W V Med J*. 2014;110(3):14-8. PMID: 24984400. Exclusion: Ineligible intervention.
811. Wichman CL, Laszewski A, Doering JJ, et al. Feasibility of model adaptations and implementation of a perinatal psychiatric teleconsultation program. *Gen Hosp Psychiatry*. 2019;59:51-7. doi: 10.1016/j.genhosppsych.2019.05.007. PMID: 31163299. Exclusion: Ineligible population.
812. Wilcox ME, Wiener-Kronish JP. Telemedicine in the intensive care unit: effect of a remote intensivist on outcomes. *JAMA Intern Med*. 2014;174(7):1167-9. doi: 10.1001/jamainternmed.2014.289. PMID: 24819239. Exclusion: Not a study.

813. Williams M, Blanco AA, Hogg R, et al. Creating a virtual community of practice: an evaluation of ophthalmology-optometry Project ECHO. *Eye*. 2018;32(12):1910-1. doi: 10.1038/s41433-018-0184-2. PMID: 30068927. Exclusion: Ineligible study design.
814. Willmitch B, Golembeski S, Kim SS, et al. Clinical outcomes after telemedicine intensive care unit implementation. *Crit Care Med*. 2012;40(2):450-4. doi: 10.1097/CCM.0b013e318232d694. PMID: 22020235. Exclusion: Ineligible population.
815. Wilson K, Dennison C, Struminger B, et al. Building a Virtual Global Knowledge Network during COVID-19: The Infection Prevention and Control Global Webinar Series. *Clinical Infectious Diseases*. 2021;06:06. doi: 10.1093/cid/ciab320. PMID: 33956127. Exclusion: Ineligible study design.
816. Winkler MF, Tappenden KA, Spangenburg M, et al. Learn Intestinal Failure Tele-ECHO Project: An innovative online telementoring and case-based learning clinic. *Nutr Clin Pract*. 2021;36(4):785-92. doi: 10.1002/ncp.10743. PMID: 34159643. Exclusion: Not a study.
817. Wood BR, Mann MS, Martinez-Paz N, et al. Project ECHO: telementoring to educate and support prescribing of HIV pre-exposure prophylaxis by community medical providers. *Sex Health*. 2018;15(6):601-5. doi: 10.1071/SH18062. PMID: 30318034. Exclusion: Ineligible population.
818. Wood D. STORC helps deliver healthy babies: the telemedicine program that serves rural women with high-risk pregnancies. *Telemed J E Health*. 2011;17(1):2-4. doi: 10.1089/tmj.2011.9996. PMID: 21294685. Exclusion: Ineligible comparator.
819. Wood T, Freeman S, Banner D, et al. Factors associated with teletrauma utilization in rural areas: a review of the literature. *Rural Remote Health*. 2021;21(1):6354. doi: 10.22605/RRH6354. PMID: 33721502. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
820. Woodall T, Ramage M, LaBruyere JT, et al. Telemedicine services during COVID-19: considerations for medically underserved populations. *J Rural Health*. 2020;02:1-4. doi: 10.1111/jrh.12466. PMID: 32613657. Exclusion: Not a study.
821. Woodley A. Can tele dermatology meet the needs of the remote and rural population? *Br J Nurs*. 2021;30(10):574-9. doi: 10.12968/bjon.2021.30.10.574. PMID: 34037451. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
822. Woollard M, Pitt K, Hayward AJ, et al. Limited benefits of ambulance telemetry in delivering early thrombolysis: a randomised controlled trial. *Emerg Med J*. 2005;22(3):209-15. doi: 10.1136/emj.2003.013482. PMID: 15735276. Exclusion: Paper published prior to 2010 (KQ1) or 2015 (KQ2a).
823. Xu W, Pan Z, Lu S, et al. Regional heterogeneity of application and effect of telemedicine in the primary care centres in rural China. *Int J Environ Res Public Health*. 2020;17(12):24. doi: 10.3390/ijerph17124531. PMID: 32599689. Exclusion: Ineligible study design.
824. Xyrichis A, Iliopoulou K, Mackintosh NJ, et al. Healthcare stakeholders' perceptions and experiences of factors affecting the implementation of critical care telemedicine (CCT): qualitative evidence synthesis. *Cochrane Database Syst Rev*. 2021;2:CD012876. doi: 10.1002/14651858.CD012876.pub2. PMID: 33599282. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.

825. Yang NH, Dharmar M, Yoo BK, et al. Economic evaluation of pediatric telemedicine consultations to rural emergency departments. *Med Decis Making*. 2015;35(6):773-83. doi: 10.1177/0272989X15584916. PMID: 25952744. Exclusion: Ineligible outcome.
826. Yao Q, Zou L, Lan H, et al. Initial experience of a tele-radiotherapy system for training radiation oncologists in rural areas. *J Cancer Educ*. 2020;28:28. doi: 10.1007/s13187-020-01836-9. PMID: 32725417. Exclusion: Ineligible outcome.
827. Yennurajalingam S, Amos CE, Jr., Weru J, et al. Extension for Community Healthcare Outcomes-Palliative Care in Africa program: improving access to quality palliative care. *J Glob Oncol*. 2019;5:1-8. doi: 10.1200/JGO.19.00128. PMID: 31335237. Exclusion: Ineligible setting.
828. Yoo BK, Kim M, Sasaki T, et al. Selected use of telemedicine in intensive care units based on severity of illness improves cost-effectiveness. *Telemed J E Health*. 2018;24(1):21-36. doi: 10.1089/tmj.2017.0069. PMID: 28661790. Exclusion: Ineligible study design.
829. Yoon MS, Martin KB, Marraccini RL, et al. Provider Satisfaction from a New Telepsychiatry Inpatient Consultation Service. *Telemed J E Health*. 2021;20:20. doi: 10.1089/tmj.2020.0317. PMID: 33471599. Exclusion: Ineligible intervention.
830. Young LB, Chan PS, Lu X, et al. Impact of telemedicine intensive care unit coverage on patient outcomes: a systematic review and meta-analysis. *Arch Intern Med*. 2011;171(6):498-506. doi: 10.1001/archinternmed.2011.61. PMID: 21444842. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
831. Young RJ, Taylor J, Friede T, et al. Pro-Active Call Center Treatment Support (PACCTS) to improve glucose control in type 2 diabetes: a randomized controlled trial. *Diabetes care*. 2005;28(2):278-82. PMID: 15677779. Exclusion: Ineligible population.
832. Yurkiewicz IR, Lappan CM, Neely ET, et al. Outcomes from a US military neurology and traumatic brain injury telemedicine program. *Neurology*. 2012;79(12):1237-43. doi: 10.1212/WNL.0b013e31826aac33. PMID: 22955133. Exclusion: Ineligible population.
833. Zaidi SF, Jumma MA, Urrea XN, et al. Telestroke-guided intravenous tissue-type plasminogen activator treatment achieves a similar clinical outcome as thrombolysis at a comprehensive stroke center. *Stroke*. 2011;42(11):3291-3. doi: 10.1161/strokeaha.111.625046. PMID: 21885843. Exclusion: Ineligible population.
834. Zakus D, Moussa M, Ezechiel M, et al. Clinical evaluation of the use of an mhealth intervention on quality of care provided by community health workers in southwest Niger. *J Glob Health*. 2019;9(1):010812. doi: 10.7189/jogh.09.010812. PMID: 31263555. Exclusion: Ineligible setting.
835. Zangara G, Valentino F, Spinelli G, et al. An Albanian open source telemedicine platform. *Telemed J E Health*. 2014;20(7):673-7. doi: 10.1089/tmj.2013.0239. PMID: 24906118. Exclusion: Ineligible setting.
836. Zangbar B, Pandit V, Rhee P, et al. Smartphone surgery: how technology can transform practice. *Telemed J E Health*. 2014;20(6):590-2. doi: 10.1089/tmj.2013.0234. PMID: 24693938. Exclusion: Not a study.
837. Zarchi K, Haugaard VB, Dufour DN, et al. Expert advice provided through telemedicine improves healing of chronic wounds: prospective cluster controlled study. *J Invest Dermatol*. 2015;135(3):895-900. doi: 10.1038/jid.2014.441. PMID: 25290685. Exclusion: Ineligible population.

838. Zhang LL, Guo YJ, Lin YP, et al. Stroke Care in the First Affiliated Hospital of Chengdu Medical College during the COVID-19 Outbreak. *Eur Neurol.* 2020;83(6):630-5. doi: 10.1159/000513097. PMID: 33341815. Exclusion: Ineligible population.
839. Zhao J, Saleemohamed N, Stinson J, et al. Health care providers' experiences and perceptions participating in a chronic pain telementoring education program: A qualitative study. *Can J Pain.* 2020;4(1):111-21. doi: 10.1080/24740527.2020.1749003. PMID: 33987490. Exclusion: Ineligible study design.
840. Zhou C, Crawford A, Serhal E, et al. The impact of Project ECHO on participant and patient outcomes: a systematic review. *Acad Med.* 2016;91(10):1439-61. PMID: 27489018. Exclusion: Systematic review, secondary analysis, or meta-analysis used as a source document only to identify individual studies.
841. Zhuo C, Wei X, Zhang Z, et al. An antibiotic stewardship programme to reduce inappropriate antibiotic prescribing for acute respiratory infections in rural Chinese primary care facilities: study protocol for a clustered randomised controlled trial. *Trials.* 2020;21(1):394. doi: 10.1186/s13063-020-04303-4. PMID: 32398065. Exclusion: Not a study.
842. Ziemba SJ, Bradley NS, Landry LA, et al. Posttraumatic stress disorder treatment for Operation Enduring Freedom/Operation Iraqi Freedom combat veterans through a civilian community-based telemedicine network. *Telemed J E Health.* 2014;20(5):446-50. doi: 10.1089/tmj.2013.0312. PMID: 24617961. Exclusion: Ineligible intervention.
843. Zöllner JP, Noda AH, McCoy J, et al. Attitudes and barriers towards telemedicine in epilepsy care: a survey among neurological practices. *Zeitschrift für Epileptologie.* 2021;34(3):318-23. doi: 10.1007/s10309-021-00417-0. Exclusion: Not English language.
844. Zubkoff L, Lyons KD, Dionne-Odom JN, et al. A cluster randomized controlled trial comparing Virtual Learning Collaborative and Technical Assistance strategies to implement an early palliative care program for patients with advanced cancer and their caregivers: a study protocol. *Implement Sci.* 2021;16(1):25. doi: 10.1186/s13012-021-01086-3. PMID: 33706770. Exclusion: Ineligible study design.
845. Zupan M, Zaletel M, Štular Ž, et al. The national telestroke network 'telekap' improves the rate of mechanical recanalization in slovenia. *Int J Stroke.* 2020;15(1 SUPPL):317. doi: 10.1177/1747493020963387. Exclusion: Not a study.

Appendix H. Appendix References

1. Berkman ND, Lohr KN, Ansari MT, et al. Grading the strength of a body of evidence when assessing health care interventions: an EPC update. *J Clin Epidemiol*. 2015 Nov;68(11):1312-24. doi: 10.1016/j.jclinepi.2014.11.023. PMID: 25721570.
2. Evers S, Goossens M, de Vet H, et al. Criteria list for assessment of methodological quality of economic evaluations: Consensus on Health Economic Criteria. *Int J Technol Assess Health Care*. 2005 Spring;21(2):240-5. PMID: 15921065.
3. Damschroder LJ, Aron DC, Keith RE, et al. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci*. 2009 Aug 7;4:50. doi: 10.1186/1748-5908-4-50. PMID: 19664226.
4. Berkman ND, Lohr KN, Ansari M, et al. AHRQ Methods for Effective Health Care Grading the Strength of a Body of Evidence When Assessing Health Care Interventions for the Effective Health Care Program of the Agency for Healthcare Research and Quality: An Update. *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*. Rockville (MD): Agency for Healthcare Research and Quality (US); 2013.
5. Lewin S, Booth A, Glenton C, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings: introduction to the series. *Implement Sci*. 2018 Jan 25;13(Suppl 1):2. doi: 10.1186/s13012-017-0688-3. PMID: 29384079.
6. Adcock AK, Choi J, Alvi M, et al. Expanding acute stroke care in rural America: a model for statewide success. *Telemed J E Health*. 2020 Jul;26(7):865-71. doi: 10.1089/tmj.2019.0087. PMID: 31596679.
7. Al Kasab S, Adams RJ, Debenham E, et al. Medical university of South Carolina telestroke: a telemedicine facilitated network for stroke treatment in South Carolina-a progress report. *Telemed J E Health*. 2017 08;23(8):674-7. doi: 10.1089/tmj.2016.0229. PMID: 28170316.
8. Alishahi Tabriz A, Turner K, Williams D, et al. Association of financial factors and telemedicine adoption for heart attack and stroke care among rural and urban hospitals: a longitudinal study. *Telemed J E Health*. 2021 Sep 24;24:24. doi: 10.1089/tmj.2021.0341. PMID: 34559014.
9. Alschuler KN, Stobbe GA, Hertz DP, et al. Impact of multiple sclerosis Project ECHO (extension for community healthcare outcomes) on provider confidence and clinical practice. *Int J MS Care*. 2019 Jul-Aug;21(4):143-50. doi: 10.7224/1537-2073.2018-014. PMID: 31474806.
10. Anderson EJ, Axon DR, Taylor AM, et al. Impact evaluation of a four-year academic-community partnership in provision of medication management and tertiary prevention services for rural patients with diabetes and/or hypertension. *Prev Med Rep*. 2020;17:101038. doi: 10.1016/j.pmedr.2019.101038. PMID: 31956473.
11. Arora S, Thornton K, Murata G, et al. Outcomes of treatment for hepatitis C virus infection by primary care providers. *N Engl J Med*. 2011 Jun 09;364(23):2199-207. doi: 10.1056/NEJMoa1009370. PMID: 21631316.
12. Astarcioğlu MA, Sen T, Kilit C, et al. Time-to-reperfusion in STEMI undergoing interhospital transfer using smartphone and WhatsApp messenger. *Am J Emerg Med*. 2015 Oct;33(10):1382-4. doi: 10.1016/j.ajem.2015.07.029. PMID: 26299691.
13. Avent ML, Walker D, Yarwood T, et al. Implementation of a novel antimicrobial stewardship strategy for rural facilities utilising telehealth. *Int J Antimicrob Agents*. 2021 Jun;57(6):106346. doi: 10.1016/j.ijantimicag.2021.106346. PMID: 33882332.
14. Bagot KL, Cadilhac DA, Kim J, et al. Transitioning from a single-site pilot project to a state-wide regional telehealth service: the experience from the Victorian Stroke Telemedicine programme. *J Telemed Telecare*. 2017 Dec;23(10):850-5. doi: 10.1177/1357633X17734004. PMID: 29081268.

15. Banbury A, Smith AC, Mehrotra A, et al. A comparison study between metropolitan and rural hospital-based telehealth activity to inform adoption and expansion. *J Telemed Telecare*. 2021 Mar 26;1357633X21998201. doi: 10.1177/1357633X21998201. PMID: 33765879.
16. Barber M, Frieslick J, Maclean A, et al. The Western Isles Stroke Telerehabilitation (Specialist Medical Consultation) Service—implementation and evaluation. *Eur Res Telemed*. 2015;4(1):19-24.
17. Becevic M, Smith E, Golzy M, et al. Melanoma extension for community healthcare outcomes: a feasibility study of melanoma screening implementation in primary care settings. *Cureus*. 2021 May 29;13(5):e15322. doi: 10.7759/cureus.15322. PMID: 34221770.
18. Bellesheim KR, Kizzee RL, Curran A, et al. ECHO autism: integrating maintenance of certification with extension for community healthcare outcomes improves developmental screening. *J Dev Behav Pediatr*. 2020 Aug;41(6):420-7. doi: 10.1097/DBP.0000000000000796. PMID: 32735419.
19. Bennett-Levy J, Hawkins R, Perry H, et al. Online cognitive behavioural therapy training for therapists: outcomes, acceptability, and impact of support. *Aust Psychol*. 2012;47(3):174-82. doi: 10.1111/j.1742-9544.2012.00089.x.
20. Bladin CF, Molocziej N, Ermel S, et al. Victorian Stroke Telemedicine Project: implementation of a new model of translational stroke care for Australia. *Intern Med J*. 2015 Sep;45(9):951-6. doi: 10.1111/imj.12822. PMID: 26011155.
21. Boltz M, Cuellar NG, Cole C, et al. Comparing an on-site nurse practitioner with telemedicine physician support hospitalist programme with a traditional physician hospitalist programme. *J Telemed Telecare*. 2019 May;25(4):213-20. doi: 10.1177/1357633X18758744. PMID: 29498301.
22. Boman K, Olofsson M, Berggren P, et al. Robot-assisted remote echocardiographic examination and teleconsultation: a randomized comparison of time to diagnosis with standard of care referral approach. *JACC Cardiovasc Imaging*. 2014 Aug;7(8):799-803. doi: 10.1016/j.jcmg.2014.05.006. PMID: 25124011.
23. Bouchonville MF, Hager BW, Kirk JB, et al. Endo ECHO improves primary care provider and community health worker self-efficacy in complex diabetes management in medically underserved communities. *Endocr Pract*. 2018 Jan;24(1):40-6. doi: 10.4158/EP-2017-0079. PMID: 29368967.
24. Brunet N, Moore DT, Lendvai Wischik D, et al. Increasing buprenorphine access for veterans with opioid use disorder in rural clinics using telemedicine. *Subst Abus*. 2020 Feb 20:1-8. doi: 10.1080/08897077.2020.1728466. PMID: 32078492.
25. Brunetti ND, Dell'Anno A, Martone A, et al. Prehospital ECG transmission results in shorter door-to-wire time for STEMI patients in a remote mountainous region. *Am J Emerg Med*. 2020 02;38(2):252-7. doi: 10.1016/j.ajem.2019.04.046. PMID: 31079977.
26. Brunetti ND, Di Pietro G, Aquilino A, et al. Pre-hospital electrocardiogram triage with tele-cardiology support is associated with shorter time-to-balloon and higher rates of timely reperfusion even in rural areas: data from the Bari- Barletta/Andria/Trani public emergency medical service 118 registry on primary angioplasty in ST-elevation myocardial infarction. *Europ Heart J Acute Cardiovasc Care*. 2014 Sep;3(3):204-13. doi: 10.1177/2048872614527009. PMID: 24604713.
27. Buckley D, Weisser S. Videoconferencing could reduce the number of mental health patients transferred from outlying facilities to a regional mental health unit. *Aust N Z J Public Health*. 2012 Oct;36(5):478-82. doi: 10.1111/j.1753-6405.2012.00915.x. PMID: 23025371.

28. Cadilhac DA, Vu M, Bladin C. Experience with scaling up the Victorian Stroke Telemedicine programme. *J Telemed Telecare*. 2014 Oct;20(7):413-8. doi: 10.1177/1357633X14552389. PMID: 25400003.
29. Carter BL, Levy B, Gryzlak B, et al. Cluster-randomized trial to evaluate a centralized clinical pharmacy service in private family medicine offices. *Circ Cardiovasc Qual Outcomes*. 2018 06;11(6):e004188. doi: 10.1161/CIRCOUTCOMES.117.004188. PMID: 29884657.
30. Chen J, Amaize A, Barath D. Evaluating telehealth adoption and related barriers among hospitals located in rural and urban areas. *J Rural Health*. 2020 Nov 12;12:12. doi: 10.1111/jrh.12534. PMID: 33180363.
31. Cho JH, Kwon HS, Kim HS, et al. Effects on diabetes management of a health-care provider mediated, remote coaching system via a PDA-type glucometer and the internet. *J Telemed Telecare*. 2011;17(7):365-70. doi: 10.1258/jtt.2011.100913. PMID: 21933896.
32. Conn DK, Madan R, Lam J, et al. Program evaluation of a telepsychiatry service for older adults connecting a university-affiliated geriatric center to a rural psychogeriatric outreach service in Northwest Ontario, Canada. *Int Psychogeriatr*. 2013 Nov;25(11):1795-800. doi: 10.1017/S104161021300118X. PMID: 23870297.
33. Cronin T. Implementing a stroke program using telemedicine. *J Emerg Nurs*. 2013 Nov;39(6):613-8. doi: 10.1016/j.jen.2012.04.012. PMID: 22748668.
34. Crossland L, Askew D, Ware R, et al. Diabetic retinopathy screening and monitoring of early stage disease in Australian general practice: tackling preventable blindness within a chronic care model. *J Diabetes Res*. 2016;2016:8405395. doi: 10.1155/2016/8405395. PMID: 26798655.
35. Datta SK, Warshaw EM, Edison KE, et al. Cost and Utility Analysis of a Store-and-Forward Teledermatology Referral System: A Randomized Clinical Trial. *JAMA Dermatol*. 2015 Dec 1;151(12):1323-9. doi: 10.1001/jamadermatol.2015.2362. PMID: 26375589.
36. Davis TD, Deen T, Bryant-Bedell K, et al. Does minority racial-ethnic status moderate outcomes of collaborative care for depression? *Psychiatr Serv*. 2011 Nov;62(11):1282-8. doi: 10.1176/ps.62.11.pss6211_1282. PMID: 22211206.
37. De Batlle J, Massip M, Vargiu E, et al. Implementing mobile health-enabled integrated care for complex chronic patients: intervention effectiveness and cost-effectiveness study. *JMIR Mhealth Uhealth*. 2021 01 14;9(1):e22135. doi: 10.2196/22135. PMID: 33443486.
38. Demaerschalk BM, Raman R, Ernstrom K, et al. Efficacy of telemedicine for stroke: pooled analysis of the Stroke Team Remote Evaluation using a Digital Observation Camera (STRoKE DOC) and STRoKE DOC Arizona telestroke trials. *Telemed J E Health*. 2012 Apr;18(3):230-7. doi: 10.1089/tmj.2011.0116. PMID: 22400970.
39. Dharmar M, Kuppermann N, Romano PS, et al. Telemedicine consultations and medication errors in rural emergency departments. *Pediatrics*. 2013 Dec;132(6):1090-7. doi: 10.1542/peds.2013-1374. PMID: 24276844.
40. Dharmar M, Romano PS, Kuppermann N, et al. Impact of critical care telemedicine consultations on children in rural emergency departments. *Crit Care Med*. 2013 Oct;41(10):2388-95. doi: 10.1097/CCM.0b013e31828e9824. PMID: 23921273.
41. Dharmar M, Sadorra CK, Leigh P, et al. The financial impact of a pediatric telemedicine program: a children's hospital's perspective. *Telemed J E Health*. 2013 Jul;19(7):502-8. doi: 10.1089/tmj.2012.0266. PMID: 23837516.

42. Doyle C, Jackson D, Loi S, et al. Videoconferencing and telementoring about dementia care: evaluation of a pilot model for sharing scarce old age psychiatry resources. *Int Psychogeriatr*. 2016 09;28(9):1567-74. doi: 10.1017/S1041610216000740. PMID: 27189501.
43. Evans K, Lerch S, Boyce TW, et al. An innovative approach to enhancing access to medical screening for miners using a mobile clinic with telemedicine capability. *J Health Care Poor Underserved*. 2016;27(4A):62-72. PMID: 27818414.
44. Fairchild RM, Ferng-Kuo SF, Laws S, et al. Telehealth decreases rural emergency department wait times for behavioral health patients in a group of critical access hospitals. *Telemed J E Health*. 2019 12;25(12):1154-64. doi: 10.1089/tmj.2018.0227. PMID: 30735100.
45. Fortney JC, Maciejewski ML, Tripathi SP, et al. A budget impact analysis of telemedicine-based collaborative care for depression. *Med Care*. 2011 Sep;49(9):872-80. doi: 10.1097/MLR.0b013e31821d2b35. PMID: 21623240.
46. Fortney JC, Pyne JM, Edlund MJ, et al. A randomized trial of telemedicine-based collaborative care for depression. *J Gen Intern Med*. 2007 Aug;22(8):1086-93. doi: 10.1007/s11606-007-0201-9. PMID: 17492326.
47. Fortney JC, Pyne JM, Kimbrell TA, et al. Telemedicine-based collaborative care for posttraumatic stress disorder: a randomized clinical trial. *JAMA Psychiatry*. 2015 Jan;72(1):58-67. doi: 10.1001/jamapsychiatry.2014.1575. PMID: 25409287.
48. Fortney JC, Pyne JM, Mouden SB, et al. Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. *Am J Psychiatry*. 2013 Apr;170(4):414-25. doi: 10.1176/appi.ajp.2012.12050696. PMID: 23429924.
49. Fortney JC, Pyne JM, Mouden SB, et al. Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. *Focus*. 2017 Jul;15(3):361-72. doi: 10.1176/appi.focus.15306. PMID: 32015700.
50. Freeman RE, Boggs KM, Zachrison KS, et al. National study of telepsychiatry use in U.S. emergency departments. *Psychiatr Serv*. 2020 Jun 01;71(6):540-6. doi: 10.1176/appi.ps.201900237. PMID: 32019430.
51. Gadomski A, Anderson J, Chung YK, et al. Full agonist opioid prescribing by primary care clinicians after buprenorphine training. *Subst Abus*. 2020 Mar 09:1-7. doi: 10.1080/08897077.2020.1736709. PMID: 32150525.
52. Gallagher R, Giles M, Morison J, et al. Telehealth-based model of care redesign to facilitate local fitting and management of patients with a spinal fracture requiring a thoracic lumbar sacral orthosis in rural hospitals in New South Wales. *Aust J Rural Health*. 2018 Jun;26(3):181-7. doi: 10.1111/ajr.12407. PMID: 29573012.
53. Gill CJ, Le Ngoc B, Halim N, et al. The mCME project: a randomized controlled trial of an SMS-based continuing medical education intervention for improving medical knowledge among Vietnamese community based physicians' assistants. *PLoS One* 2016;11(11):e0166293. doi: 10.1371/journal.pone.0166293. PMID: 27861516.
54. Habashi P, Bouchard S, Nguyen GC. Transforming access to specialist care for inflammatory bowel disease: the PACE telemedicine program. *J Can Assoc Gastroenterol*. 2019 Dec;2(4):186-94. doi: 10.1093/jcag/gwy046. PMID: 31616860.
55. Hall RW, Hall-Barrow J, Garcia-Rill E. Neonatal regionalization through telemedicine using a community-based research and education core facility. *Ethn Dis*. 2010;20(1 Suppl 1):S1-136-40. PMID: 20521402.

56. Haozous E, Doorenbos AZ, Demiris G, et al. Role of telehealth/videoconferencing in managing cancer pain in rural American Indian communities. *Psycho-Oncol.* 2012 Feb;21(2):219-23. doi: 10.1002/pon.1887. PMID: 22271543.
57. Haque SN, DeStefano S, Banger A, et al. Factors influencing telehealth implementation and use in frontier critical access hospitals: qualitative study. *JMIR Form Res.* 2021 May 05;5(5):e24118. doi: 10.2196/24118. PMID: 33949958.
58. Harvey JB, Yeager BE, Cramer C, et al. The impact of telemedicine on pediatric critical care triage. *Pediatr Crit Care Med.* 2017 Nov;18(11):e555-e60. doi: 10.1097/PCC.0000000000001330. PMID: 28922271.
59. Haynes SC, Dharmar M, Hill BC, et al. The impact of telemedicine on transfer rates of newborns at rural community hospitals. *Acad Pediatr.* 2020 Jul;20(5):636-41. doi: 10.1016/j.acap.2020.02.013. PMID: 32081766.
60. Haynes SC, Hoffman KR, Patel S, et al. The use of telemedicine for stabilization of neonates transferred from rural community hospitals. *Telemed J E Health.* 2021 Mar 01;01:01. doi: 10.1089/tmj.2020.0503. PMID: 33691080.
61. Helwig SA, Ragoschke-Schumm A, Schwindling L, et al. Prehospital Stroke Management Optimized by Use of Clinical Scoring vs Mobile Stroke Unit for Triage of Patients With Stroke: A Randomized Clinical Trial. *JAMA Neurol.* 2019 Sep 3;76(12):1484-92. doi: 10.1001/jamaneurol.2019.2829. PMID: 31479116.
62. Hensel JM, Yang R, Rai M, et al. Optimizing electronic consultation between primary care providers and psychiatrists: mixed-methods study. *J Med Internet Res.* 2018 04 06;20(4):e124. doi: 10.2196/jmir.8943. PMID: 29625949.
63. Hilt RJ, Barclay RP, Bush J, et al. A statewide child telepsychiatry consult system yields desired health system changes and savings. *Telemed J E Health.* 2015 Jul;21(7):533-7. doi: 10.1089/tmj.2014.0161. PMID: 25799043.
64. Hofmeyer J, Leider JP, Satorius J, et al. Implementation of telemedicine consultation to assess unplanned transfers in rural long-term care facilities, 2012-2015: a pilot study. *J Am Med Dir Assoc.* 2016 11 01;17(11):1006-10. doi: 10.1016/j.jamda.2016.06.014. PMID: 27477614.
65. Holt T, Sari N, Hansen G, et al. Remote presence robotic technology reduces need for pediatric interfacility transportation from an isolated northern community. *Telemed J E Health.* 2018 11;24(11):927-33. doi: 10.1089/tmj.2017.0211. PMID: 29394155.
66. Hostutler CA, Valleru J, Maciejewski HM, et al. Improving pediatrician's behavioral health competencies through the Project ECHO teleconsultation model. *Clin Pediatr (Phila).* 2020 Jun 07;59(12):1049-57. doi: 10.1177/0009922820927018. PMID: 32506939.
67. Howland M, Tennant M, Bowen DJ, et al. Psychiatrist and psychologist experiences with telehealth and remote collaborative care in primary care: a qualitative study. *J Rural Health.* 2020 Oct 06;06:06. doi: 10.1111/jrh.12523. PMID: 33022079.
68. Jewer J, Parsons MH, Dunne C, et al. Evaluation of a mobile telesimulation unit to train rural and remote practitioners on high-acuity low-occurrence procedures: pilot randomized controlled trial. *J Med Internet Res.* 2019 08 06;21(8):e14587. doi: 10.2196/14587. PMID: 31389340.
69. Jhaveri D, Larkins S, Kelly J, et al. Remote chemotherapy supervision model for rural cancer care: perspectives of health professionals. *Eur J Cancer Care (Engl).* 2016 Jan;25(1):93-8. doi: 10.1111/ecc.12309. PMID: 25871852.
70. Jiang B, Bills M, Poon P. Integrated telehealth-assisted home-based specialist palliative care in rural Australia: a feasibility study. *J Telemed Telecare.* 2020 Oct 20;1357633X20966466. doi: 10.1177/1357633X20966466. PMID: 33079611.

71. Johansson AM, Lindberg I, Soderberg S. The views of health-care personnel about video consultation prior to implementation in primary health care in rural areas. *Prim Health Care Res Dev.* 2014 Apr;15(2):170-9. doi: 10.1017/S1463423613000030. PMID: 23402617.
72. Kahn JM, Le TQ, Barnato AE, et al. ICU Telemedicine and Critical Care Mortality: A National Effectiveness Study. *Med Care.* 2016 Mar;54(3):319-25. doi: 10.1097/MLR.0000000000000485. PMID: 26765148.
73. Katzman JG, Gygi K, Swift R, et al. How hands-on pain skills intensive trainings complement ECHO pain and opioid management programs: a program evaluation with the Indian Health Service. *Pain Med.* 2020 Jun 18;21(9):1769-78. doi: 10.1093/pm/pnaa151. PMID: 32556294.
74. Khalil H, Cullen M, Chambers H, et al. Implementation of a successful electronic wound documentation system in rural Victoria, Australia: a subject of collaboration and community engagement. *Int Wound J.* 2014 Jun;11(3):314-8. doi: 10.1111/iwj.12041. PMID: 23418740.
75. Kim EW, Teague-Ross TJ, Greenfield WW, et al. Telemedicine collaboration improves perinatal regionalization and lowers statewide infant mortality. *J Perinatol.* 2013 Sep;33(9):725-30. doi: 10.1038/jp.2013.37. PMID: 23579490.
76. Kim H, Jhoo JH, Jang JW. The effect of telemedicine on cognitive decline in patients with dementia. *J Telemed Telecare.* 2017 Jan;23(1):149-54. doi: 10.1177/1357633X15615049. PMID: 26541171.
77. Kulcsar M, Gilchrist S, George MG. Improving stroke outcomes in rural areas through telestroke programs: an examination of barriers, facilitators, and state policies. *Telemed J E Health.* 2014 Jan;20(1):3-10. doi: 10.1089/tmj.2013.0048. PMID: 24286197.
78. Kuperman EF, Linson EL, Klefstad K, et al. The virtual hospitalist: a single-site implementation bringing hospitalist coverage to critical access hospitals. *J Hosp Med.* 2018 11 01;13(11):759-63. doi: 10.12788/jhm.3061. PMID: 30255859.
79. Kwak MY, Hwang EJ, Lee TH. Effects of the Physician-Primary-Healthcare Nurse Telemedicine Model (P-NTM) on medication adherence and Health-Related Quality of Life (HRQoL) of patients with chronic disease at remote rural areas. *Int J Environ Res Public Health.* 2021 03 03;18(5):03. doi: 10.3390/ijerph18052502. PMID: 33802513.
80. Leshner AP, Fakhry SM, DuBose-Morris R, et al. Development and evolution of a statewide outpatient consultation service: leveraging telemedicine to improve access to specialty care. *Popul Health Manag.* 2020 02;23(1):20-8. doi: 10.1089/pop.2018.0212. PMID: 31161963.
81. Levin K, Madsen JR, Petersen I, et al. Telemedicine diabetes consultations are cost-effective, and effects on essential diabetes treatment parameters are similar to conventional treatment: 7-year results from the Svendborg Telemedicine Diabetes Project. *J Diabetes Sci Technol.* 2013 May 1;7(3):587-95. doi: 10.1177/193229681300700302. PMID: 23759390.
82. Liddy C, Moroz I, Keely E, et al. Understanding the impact of a multispecialty electronic consultation service on family physician referral rates to specialists: a randomized controlled trial using health administrative data. *Trials.* 2019 Jun 10;20(1):348. doi: 10.1186/s13063-019-3393-5. PMID: 31182123.
83. Lim M, Liberali SAC, Calache H, et al. Specialist networks influence clinician willingness to treat individuals with special needs. *JDR Clin Trans Res.* 2021 Jun 21;23800844211020250. doi: 10.1177/23800844211020250. PMID: 34148391.
84. Lindauer A, Wild K, Natanson A, et al. Dementia 360 ECHO: using technology to facilitate diagnosis and treatment. *Gerontol Geriatr Educ.* 2020 Oct 20;1-7. doi: 10.1080/02701960.2020.1835658. PMID: 33078687.

85. Lingum NR, Sokoloff LG, Meyer RM, et al. Building long-term care staff capacity during COVID-19 through just-in-time learning: evaluation of a modified ECHO model. *J Am Med Dir Assoc*. 2021 Feb;22(2):238-44.e1. doi: 10.1016/j.jamda.2020.10.039. PMID: 33238143.
86. Liou JK, Soon MS, Chen CH, et al. Shared care combined with telecare improves glycemic control of diabetic patients in a rural underserved community. *Telemed J E Health*. 2014 Feb;20(2):175-8. doi: 10.1089/tmj.2013.0037. PMID: 24320193.
87. Long MC, Angtuaco T, Lowery C. Ultrasound in telemedicine: its impact in high-risk obstetric health care delivery. *Ultrasound Q*. 2014 Sep;30(3):167-72. doi: 10.1097/RUQ.0000000000000073. PMID: 25148484.
88. Lucas JAM, Day K, Honey MLL. Clinician's perceptions of telehealth for emergency care on the west coast of New Zealand: findings of a descriptive study. *Emergency Nurse New Zealand*. 2016:6-10.
89. Luscombe GM, Hawthorn J, Wu A, et al. 'Empowering clinicians in smaller sites': A qualitative study of clinician's experiences with a rural Virtual Paediatric Feeding Clinic. *Aust J Rural Health*. 2021 Sep 07;07:07. doi: 10.1111/ajr.12781. PMID: 34490941.
90. MacKinney AC, Ward MM, Ullrich F, et al. The business case for tele-emergency. *Telemed J E Health*. 2015 Dec;21(12):1005-11. doi: 10.1089/tmj.2014.0241. PMID: 26226603.
91. Mahmoud H, Vogt EL, Dahdouh R, et al. Using continuous quality improvement to design and implement a telepsychiatry program in rural Illinois. *Psychiatr Serv*. 2020 Apr 15;71(8):860-3. doi: 10.1176/appi.ps.201900231. PMID: 32290807.
92. Makkar A, McCoy M, Hallford G, et al. A hybrid form of telemedicine: a unique way to extend intensive care service to neonates in medically underserved areas. *Telemed J E Health*. 2018 09;24(9):717-21. doi: 10.1089/tmj.2017.0155. PMID: 29298407.
93. Makkar A, McCoy M, Hallford G, et al. Evaluation of neonatal services provided in a level II NICU utilizing hybrid telemedicine: a prospective study. *Telemed J E Health*. 2020 Feb;26(2):176-83. doi: 10.1089/tmj.2018.0262. PMID: 30835166.
94. Marsh-Feiley G, Eadie L, Wilson P. Paramedic and physician perspectives on the potential use of remotely supported prehospital ultrasound. *Rural Remote Health*. 2018 09;18(3):4574. doi: 10.22605/RRH4574. PMID: 30207737.
95. May S, Jonas K, Fehler GV, et al. Challenges in current nursing home care in rural Germany and how they can be reduced by telehealth - an exploratory qualitative pre-post study. *BMC Health Serv Res*. 2021 Sep 06;21(1):925. doi: 10.1186/s12913-021-06950-y. PMID: 34488746.
96. McFarland LV, Raugi GJ, Taylor LL, et al. Implementation of an education and skills programme in a teledermatology project for rural veterans. *J Telemed Telecare*. 2012 Mar;18(2):66-71. doi: 10.1258/jtt.2011.110518. PMID: 22198956.
97. McGill AF, North JB. Teleconference fracture clinics: a trial for rural hospitals. *ANZ J Surg*. 2012 Jan-Feb;82(1-2):2-3. doi: 10.1111/j.1445-2197.2011.05952.x. PMID: 22507484.
98. McLendon SF, Wood FG, Stanley N. Enhancing diabetes care through care coordination, telemedicine, and education: evaluation of a rural pilot program. *Public Health Nurs*. 2019 05;36(3):310-20. doi: 10.1111/phn.12601. PMID: 30868661.
99. McWilliams T, Hendricks J, Twigg D, et al. Burns education for non-burn specialist clinicians in Western Australia. *Burns*. 2015 Mar;41(2):301-7. doi: 10.1016/j.burns.2014.06.015. PMID: 25239848.
100. Miller AC, Ward MM, Ullrich F, et al. Emergency department telemedicine consults are associated with faster time-to-electrocardiogram and time-to-fibrinolysis for myocardial infarction patients. *Telemed J E Health*. 2020 Feb 28;26(12):1440-8. doi: 10.1089/tmj.2019.0273. PMID: 32109200.

101. Miyamoto S, Dharmar M, Boyle C, et al. Impact of telemedicine on the quality of forensic sexual abuse examinations in rural communities. *Child Abuse Negl.* 2014 Sep;38(9):1533-9. doi: 10.1016/j.chiabu.2014.04.015. PMID: 24841062.
102. Moffatt JJ, Eley DS. Barriers to the up-take of telemedicine in Australia--a view from providers. *Rural Remote Health.* 2011;11(2):1581. PMID: 21385004.
103. Mohr NM, Campbell KD, Swanson MB, et al. Provider-to-provider telemedicine improves adherence to sepsis bundle care in community emergency departments. *J Telemed Telecare.* 2020 Jan 05;1-9. doi: 10.1177/1357633X19896667. PMID: 31903840.
104. Mohr NM, Vakkalanka JP, Harland KK, et al. Telemedicine use decreases rural emergency department length of stay for transferred North Dakota trauma patients. *Telemed J E Health.* 2018 03;24(3):194-202. doi: 10.1089/tmj.2017.0083. PMID: 28731843.
105. Mohr NM, Young T, Harland KK, et al. Emergency department telemedicine shortens rural time-to-provider and emergency department transfer times. *Telemed J E Health.* 2018 08;24(8):582-93. doi: 10.1089/tmj.2017.0262. PMID: 29293413.
106. Mohr NM, Young T, Harland KK, et al. Telemedicine is associated with faster diagnostic imaging in stroke patients: a cohort study. *Telemed J E Health.* 2019 02;25(2):93-100. doi: 10.1089/tmj.2018.0013. PMID: 29958087.
107. Momin B, Mera J, Essex W, et al. Implementation of liver cancer education among health care providers and community coalitions in the Cherokee nation. *Prev Chronic Dis.* 2019;16(E112):1-7. doi: 10.5888/pcd16.180671. PMID: 31441770.
108. Morrisette S, Pearlman RL, Kovar M, et al. Attitudes and perceived barriers toward store-and-forward teledermatology among primary care providers of the rural Mississippi. *Arch Dermatol Res.* 2021 Feb 25;25:25. doi: 10.1007/s00403-021-02208-z. PMID: 33630147.
109. Mundt AP, Irarrazaval M, Martinez P, et al. Telepsychiatry consultation for primary care treatment of children and adolescents receiving child protective services in Chile: mixed methods feasibility study. *JMIR Public Health Surveill.* 2021 Jul 22;7(7):e25836. doi: 10.2196/25836. PMID: 34292164.
110. Myers K, Vander Stoep A, Zhou C, et al. Effectiveness of a telehealth service delivery model for treating attention-deficit/hyperactivity disorder: a community-based randomized controlled trial. *J Am Acad Child Adolesc Psychiatry.* 2015 Apr;54(4):263-74. doi: 10.1016/j.jaac.2015.01.009. PMID: 25791143.
111. Nagao KJ, Koschel A, Haines HM, et al. Rural Victorian telestroke project. *Intern Med J.* 2012 Oct;42(10):1088-95. doi: 10.1111/j.1445-5994.2011.02603.x. PMID: 21981424.
112. Nagayoshi Y, Oshima S, Ogawa H. Clinical impact of telemedicine network system at rural hospitals without on-site cardiac surgery backup. *Telemed J E Health.* 2016 11;22(11):960-4. PMID: 27148833.
113. Narva AS, Romancito G, Faber T, et al. Managing CKD by telemedicine: the Zuni telenephrology clinic. *Adv Chronic Kidney Dis.* 2017 Jan;24(1):6-11. doi: 10.1053/j.ackd.2016.11.019. PMID: 28224944.
114. Natafagi N, Mohr NM, Wittrock A, et al. The association between telemedicine and emergency department (ED) disposition: a stepped wedge design of an ED-based telemedicine program in critical access hospitals. *J Rural Health.* 2020 Jun;36(3):360-70. doi: 10.1111/jrh.12370. PMID: 31013552.
115. Ness TE, Annese MF, Martinez-Paz N, et al. Using an innovative telehealth model to support community providers who deliver perinatal HIV care. *AIDS Educ Prev.* 2017 12;29(6):516-26. doi: 10.1521/aeap.2017.29.6.516. PMID: 29283274.

116. Newell MC, Strauss CE, Freier T, et al. Design and initial results of the Minneapolis Heart Institute TeleHeart program. *Circ Cardiovasc Qual Outcomes*. 2017;10:1-4. doi: 10.1161/CIRCOUTCOMES.117.003904. PMID: 29021334.
117. Nipp CM, Vogtle LK, Warren M. Clinical application of low vision rehabilitation strategies after completion of a computer-based training module. *Occup Ther Health Care*. 2014 Jul;28(3):296-305. doi: 10.3109/07380577.2014.908335. PMID: 24971896.
118. Nqala MO, Rout CC, Aldous CM. Remote clinical support by telephone for rural district hospital medical officers in the Eastern Cape. *S Afr Fam Pract*. 2015;57(5):286-90. doi: 10.1080/20786190.2015.1055671.
119. Olenik K, Lehr B. Counteracting brain drain of health professionals from rural areas via teleconsultation: analysis of the barriers and success factors of teleconsultation. *J Public Health*. 2013;21(4):357-64. doi: 10.1007/s10389-013-0565-8.
120. Painter JT, Fortney JC, Austen MA, et al. Cost-effectiveness of telemedicine-based collaborative care for posttraumatic stress disorder. *Psychiatr Serv*. 2017 Nov 01;68(11):1157-63. doi: 10.1176/appi.ps.201600485. PMID: 28669290.
121. Pandit T, Ray RA, Sabesan S. Managing emergencies in rural North Queensland: the feasibility of telerehabilitation. *Int J Telemed Appl*. 2018;2018doi: 10.1155/2018/8421346. PMID: 29849606.
122. Panloui OM, Broadfield E, Champion R, et al. Outcomes of telemedicine intervention in a regional intensive care unit: a before and after study. *Anaesth Intensive Care*. 2017 09;45(5):605-10. PMID: 28911290.
123. Parks J, Hunter A, Taylor A, et al. Design, development and implementation of the virtual, coordination, access, referral and escalation service in western New South Wales. *Aust J Rural Health*. 2021 Sep 29;29:29. doi: 10.1111/ajr.12800. PMID: 34586702.
124. Pedragosa A, Alvarez-Sabin J, Rubiera M, et al. Impact of telemedicine on acute management of stroke patients undergoing endovascular procedures. *Cerebrovasc Dis*. 2012;34(5-6):436-42. doi: 10.1159/000345088. PMID: 23207552.
125. Peracca SB, Jackson GL, Lamkin RP, et al. Implementing teledermatology for rural veterans: an evaluation using the RE-AIM framework. *Telemed J E Health*. 2020 Apr 27;1-9. doi: 10.1089/tmj.2020.0013. PMID: 32343924.
126. Pimentel CB, Gately M, Barczi SR, et al. GRECC connect: geriatrics telehealth to empower health care providers and improve management of older veterans in rural communities. *Fed Pract*. 2019 Oct;36(10):464-70. PMID: 31768097.
127. Pindyck T, Kalishman S, Flatow-Trujillo L, et al. Treating hepatitis C in American Indians/Alaskan Natives: a survey of Project ECHO(R) (Extension for Community Healthcare Outcomes) utilization by Indian Health Service providers. *SAGE Open Med*. 2015;3:2050312115612805. doi: 10.1177/2050312115612805. PMID: 26770809.
128. Poulson LK, Nissen L, Coombes I. Pharmaceutical review using telemedicine--a before and after feasibility study. *J Telemed Telecare*. 2010;16(2):95-9. doi: 10.1258/jtt.2009.090716. PMID: 20139138.
129. Powell KR, Alexander GL. Consequences of rapid telehealth expansion in nursing homes: promise and pitfalls. *Appl Clin Inform*. 2021 Aug;12(4):933-43. doi: 10.1055/s-0041-1735974. PMID: 34614517.
130. Puskar KR, Heeyoung L, Mitchell AM, et al. Interprofessional collaborative education for substance use screening: rural areas and challenges. *Online J Rural Nurs Health Care*. 2016;16(1):76-96. doi: 10.14574/ojrnhc.v16i1.385.
131. Pyne JM, Fortney JC, Mouden S, et al. Cost-effectiveness of on-site versus off-site collaborative care for depression in rural FQHCs. *Psychiatr Serv*. 2015 May 01;66(5):491-9. doi: 10.1176/appi.ps.201400186. PMID: 25686811.

132. Pyne JM, Fortney JC, Tripathi SP, et al. Cost-effectiveness analysis of a rural telemedicine collaborative care intervention for depression. *Arch Gen Psychiatry*. 2010 Aug;67(8):812-21. doi: 10.1001/archgenpsychiatry.2010.82. PMID: 20679589.
133. Ramnath VR, Khazeni N. Centralized monitoring and virtual consultant models of tele-ICU care: a side-by-side review. *Telemed J E Health*. 2014 Oct;20(10):962-71. doi: 10.1089/tmj.2014.0024. PMID: 25225795.
134. Ray KN, Felmet KA, Hamilton MF, et al. Clinician attitudes toward adoption of pediatric emergency telemedicine in rural hospitals. *Pediatr Emerg Care*. 2017 Apr;33(4):250-7. doi: 10.1097/PEC.0000000000000583. PMID: 26785087.
135. Ritter LA, Robinette TR, Cofano J. Evaluation of a statewide telemedicine program. *CJHP*. 2010;8(1):1-9. doi: 10.32398/cjhp.v8i1.2025.
136. Robinson T, Hills D, Kelly B. The evaluation of an online orientation to rural mental health practice in Australia. *J Psychiatr Ment Health Nurs*. 2011;18(7):629-36. doi: 10.1111/j.1365-2850.2011.01712.x. PMID: 21848598.
137. Rogove HJ, McArthur D, Demaerschalk BM, et al. Barriers to telemedicine: survey of current users in acute care units. *Telemed J E Health*. 2012 Jan-Feb;18(1):48-53. doi: 10.1089/tmj.2011.0071. PMID: 22082107.
138. Rojas G, Guajardo V, Martinez P, et al. A remote collaborative care program for patients with depression living in rural areas: open-label trial. *J Med Internet Res*. 2018 04 30;20(4):e158. doi: 10.2196/jmir.8803. PMID: 29712627.
139. Rossaro L, Torruellas C, Dhaliwal S, et al. Clinical outcomes of hepatitis C treated with pegylated interferon and ribavirin via telemedicine consultation in Northern California. *Dig Dis Sci*. 2013 Dec;58(12):3620-5. doi: 10.1007/s10620-013-2810-y. PMID: 24154637.
140. Sabesan S, Senko C, Schmidt A, et al. Enhancing chemotherapy capabilities in rural hospitals: implementation of a telechemotherapy model (QReCS) in North Queensland, Australia. *J Oncol Pract*. 2018 07;14(7):e429-e37. doi: 10.1200/JOP.18.00110. PMID: 29996068.
141. Sairanen T, Soinila S, Nikkanen M, et al. Two years of Finnish Telestroke: thrombolysis at spokes equal to that at the hub. *Neurology*. 2011 Mar 29;76(13):1145-52. doi: 10.1212/WNL.0b013e318212a8d4. PMID: 21368283.
142. Seibert PS, Reddy T, Whitmore T, et al. The use of telemedicine to train perioperative nurses in rural settings. *J Telemed Telecare*. 2013 Sep;19(6):311-4. doi: 10.1177/1357633X13501777. PMID: 24163294.
143. Shaikh U, Nettiksimmons J, Joseph JG, et al. Collaborative practice improvement for childhood obesity in rural clinics: the Healthy Eating Active Living Telehealth Community of Practice (HEALTH COP). *Am J Med Qual*. 2014 Nov-Dec;29(6):467-75. doi: 10.1177/1062860613506252. PMID: 24170936.
144. Shea CM, Gertner AK, Green SL. Barriers and perceived usefulness of an ECHO intervention for office-based buprenorphine treatment for opioid use disorder in North Carolina: a qualitative study. *Subst Abus*. 2019 Dec 06:1-11. doi: 10.1080/08897077.2019.1694617. PMID: 31809679.
145. Sicotte C, Moqadem K, Vasilevsky M, et al. Use of telemedicine for haemodialysis in very remote areas: the Canadian First Nations. *J Telemed Telecare*. 2011;17(3):146-9. doi: 10.1258/jtt.2010.100614. PMID: 21303935.
146. Simpson AN, Harvey JB, DiLembo SM, et al. Population health indicators associated with a statewide telestroke program. *Telemed J E Health*. 2020 Feb 11;26(9):1126-33. doi: 10.1089/tmj.2019.0204. PMID: 32045330.

147. Singh R, Mathiassen L, Switzer JA, et al. Assimilation of web-based urgent stroke evaluation: a qualitative study of two networks. *JMIR Med Inform.* 2014 Apr 15;2(1):e6. doi: 10.2196/medinform.3028. PMID: 25601232.
148. Sockalingam S, Arena A, Serhal E, et al. Building provincial mental health capacity in primary care: an evaluation of a Project ECHO mental health program. *Acad Psychiatry.* 2018 Aug;42(4):451-7. doi: 10.1007/s40596-017-0735-z. PMID: 28593537.
149. Spivak S, Spivak A, Cullen B, et al. Telepsychiatry use in U.S. mental health facilities, 2010-2017. *Psychiatr Serv.* 2020 02 01;71(2):121-7. doi: 10.1176/appi.ps.201900261. PMID: 31615370.
150. Sterling SA, Seals SR, Jones AE, et al. The impact of the TelEmergency program on rural emergency care: an implementation study. *J Telemed Telecare.* 2017 Jul;23(6):588-94. doi: 10.1177/1357633X16657499. PMID: 27470505.
151. Stevens G, Loh J, Kolbe J, et al. Comparison of recommendations for radiotherapy from two contemporaneous thoracic multidisciplinary meeting formats: co-located and video conference. *Intern Med J.* 2012 Nov;42(11):1213-8. doi: 10.1111/j.1445-5994.2012.02817.x. PMID: 22530597.
152. Summerhayes C, McGee JA, Cooper RJ, et al. Introducing leg ulcer telemedicine into rural general practice. *Wounds UK.* 2012;8(2):28-36.
153. Swanson MB, Miller AC, Ward MM, et al. Emergency department telemedicine consults decrease time to interpret computed tomography of the head in a multi-network cohort. *J Telemed Telecare.* 2019 Nov 04;1-10. doi: 10.1177/1357633X19877746. PMID: 31684801.
154. Tande AJ, Berbari EF, Ramar P, et al. Association of a remotely offered infectious diseases eConsult service with improved clinical outcomes. *Open Forum Infect Dis.* 2020 Jan:1-4. doi: 10.1093/ofid/ofaa003. PMID: 31988969.
155. Taylor-Gjevre R, Nair B, Bath B, et al. Addressing rural and remote access disparities for patients with inflammatory arthritis through video-conferencing and innovative inter-professional care models. *Musculoskelet* 2018 03;16(1):90-5. doi: 10.1002/msc.1215. PMID: 28967235.
156. Thaker DA, Monypenny R, Olver I, et al. Cost savings from a telemedicine model of care in northern Queensland, Australia. *Med J Aust.* 2013 Sep 16;199(6):414-7. PMID: 24033216.
157. Thies KM, Anderson D, Beals-Reid C. Project ECHO chronic pain: a qualitative analysis of recommendations by expert faculty. *Pain Med.* 2019 07 01;20(7):1450-2. doi: 10.1093/pm/pnz024. PMID: 30840075.
158. Tran L, Feldman R, Riley T, 3rd, et al. Association of the extension for community healthcare outcomes project with use of direct-acting antiviral treatment among US adults with hepatitis C. *JAMA Netw Open.* 2021 Jul 01;4(7):e2115523. doi: 10.1001/jamanetworkopen.2021.15523. PMID: 34213557.
159. Tripod M, Tait M, Bracey J, et al. The use of telemedicine decreases unnecessary hand trauma transfers. *Hand.* 2020 May;15(3):422-7. doi: 10.1177/1558944718810877. PMID: 30417694.
160. Trondsen MV, Bolle SR, Stensland GO, et al. Video-confidence: a qualitative exploration of videoconferencing for psychiatric emergencies. *BMC Health Serv Res.* 2014 Oct 31;14:544. doi: 10.1186/s12913-014-0544-y. PMID: 25359404.
161. Trondsen MV, Tjora A, Broom A, et al. The symbolic affordances of a video-mediated gaze in emergency psychiatry. *Soc Sci Med.* 2018 01;197:87-94. doi: 10.1016/j.socscimed.2017.11.056. PMID: 29222999.
162. Tynan A, Deeth L, McKenzie D. An integrated oral health program for rural residential aged care facilities: a mixed methods comparative study. *BMC Health Serv Res.* 2018 07 03;18(1):515. doi: 10.1186/s12913-018-3321-5. PMID: 29970073.

163. Tynan A, Deeth L, McKenzie D, et al. Integrated approach to oral health in aged care facilities using oral health practitioners and teledentistry in rural Queensland. *Aust J Rural Health*. 2018 Apr 16;26:290-4. doi: 10.1111/ajr.12410. PMID: 29660771.
164. Vakkalanka JP, Harland KK, Wittrock A, et al. Telemedicine is associated with rapid transfer and fewer involuntary holds among patients presenting with suicidal ideation in rural hospitals: a propensity matched cohort study. *J Epidemiol Community Health*. 2019 11;73(11):1033-9. doi: 10.1136/jech-2019-212623. PMID: 31492762.
165. Van der Pol M, McKenzie L. Costs and benefits of tele-endoscopy clinics in a remote location. *J Telemed Telecare*. 2010;16(2):89-94. doi: 10.1258/jtt.2009.090609. PMID: 20139140.
166. Ward MM, Ullrich F, MacKinney AC, et al. Tele-emergency utilization: in what clinical situations is tele-emergency activated? *J Telemed Telecare*. 2016 Jan;22(1):25-31. doi: 10.1177/1357633X15586319. PMID: 26026189.
167. Watts SA, Roush L, Julius M, et al. Improved glycemic control in veterans with poorly controlled diabetes mellitus using a Specialty Care Access Network-Extension for Community Healthcare Outcomes model at primary care clinics. *J Telemed Telecare*. 2016 Jun;22(4):221-4. doi: 10.1177/1357633X15598052. PMID: 26253745.
168. Weaver MS, Neumann ML, Navaneethan H, et al. Human touch via touchscreen: rural nurses' experiential perspectives on telehealth use in pediatric hospice care. *J Pain Symptom Manage*. 2020 Jun 07;60(5):1027-33. doi: 10.1016/j.jpainsymman.2020.06.003. PMID: 32525081.
169. Weigel P, Bhagianadh D, Merchant KA, et al. Tele-emergency behavioural health in rural and underserved areas. *J Telemed Telecare*. 2019 Nov 14:1-10. doi: 10.1177/1357633X19887027. PMID: 31726903.
170. Weigel PA, Merchant KA, Wittrock A, et al. Paediatric tele-emergency care: a study of two delivery models. *J Telemed Telecare*. 2019 Apr 09:1-9. doi: 10.1177/1357633X19839610. PMID: 30966860.
171. White AH, Crowther SA, Lee SH. Supporting rural midwifery practice using a mobile health (mHealth) intervention: a qualitative descriptive study. *Rural Remote Health*. 2019 08;19(3):5294. doi: 10.22605/RRH5294. PMID: 31446762.
172. Whited JD, Warshaw EM, Kapur K, et al. Clinical course outcomes for store and forward teledermatology versus conventional consultation: a randomized trial. *J Telemed Telecare*. 2013 Jun;19(4):197-204. doi: 10.1177/1357633X13487116. PMID: 23666440.
173. Williams D, Simpson AN, King K, et al. Do Hospitals Providing Telehealth in Emergency Departments Have Lower Emergency Department Costs? *Telemed J E Health*. 2020 Nov 13;13:13. doi: 10.1089/tmj.2020.0349. PMID: 33185503.
174. Williams KM, Kirsh S, Aron D, et al. Evaluation of the Veterans Health Administration's specialty care transformational initiatives to promote patient-centered delivery of specialty care: a mixed-methods approach. *Telemed J E Health*. 2017 07;23(7):577-89. doi: 10.1089/tmj.2016.0166. PMID: 28177858.
175. Wilson BM, Banks RE, Crnich CJ, et al. Changes in antibiotic use following implementation of a telehealth stewardship pilot program. *Infect Control Hosp Epidemiol*. 2019 07;40(7):810-4. doi: 10.1017/ice.2019.128. PMID: 31172905.
176. Wood PR, Caplan L. Outcomes, satisfaction, and costs of a rheumatology telemedicine program: a longitudinal evaluation. *J Clin Rheumatol*. 2019 Jan;25(1):41-4. doi: 10.1097/RHU.0000000000000778. PMID: 30461466.
177. Wood T, Freeman S, Banner D, et al. Exploring user perspectives of factors associated with use of teletrauma in rural areas. *Australas Emerg Care*. 2021 May 07;07:07. doi: 10.1016/j.aucc.2021.04.001. PMID: 33972192.

178. Yam P, Fales D, Jemison J, et al. Implementation of an antimicrobial stewardship program in a rural hospital. *Am J Health-Syst Pharm*. 2012 Jul 01;69(13):1142-8. doi: 10.2146/ajhp110512. PMID: 22722593.
179. Yan LL, Gong E, Gu W, et al. Effectiveness of a primary care-based integrated mobile health intervention for stroke management in rural China (SINEMA): A cluster-randomized controlled trial. *PLoS Med*. 2021 04;18(4):e1003582. doi: 10.1371/journal.pmed.1003582. PMID: 33909607.
180. Zachrison KS, Boggs KM, Hayden EM, et al. Understanding barriers to telemedicine implementation in rural emergency departments. *Ann Emerg Med*. 2020 03;75(3):392-9. doi: 10.1016/j.annemergmed.2019.06.026. PMID: 31474481.
181. Zachrison KS, Hayden EM, Schwamm LH, et al. Characterizing New England emergency departments by telemedicine use. *West J Emerg Med*. 2017 Oct;18(6):1055-60. doi: 10.5811/westjem.2017.8.34880. PMID: 29085537.
182. Zhang D, Wang G, Zhu W, et al. Expansion of telestroke services improves quality of care provided in super rural areas. *Health Aff (Millwood)*. 2018 12;37(12):2005-13. doi: 10.1377/hlthaff.2018.05089. PMID: 30633675.
183. Zhu X, Merchant KAS, Mohr NM, et al. Real-time learning through telemedicine enhances professional training in rural emergency departments. *Telemed J E Health*. 2020 Jun 17;17:1-7. doi: 10.1089/tmj.2020.0042. PMID: 32552479.
184. Zittleman L, Curcija K, Sutter C, et al. Building capacity for medication assisted treatment in rural primary care practices: the IT MATTTRs practice team training. *J Prim Care Community Health*. 2020 Jan-Dec;11:2150132720953723. doi: 10.1177/2150132720953723. PMID: 32909491.
185. Personal Communication: April 22, 2021 with Trudy Bearden, Senior Consultant/Telehealth Lead, Comagine Health Idaho Director.
186. Demeke HB, Merali S, Marks S, et al. Trends in use of telehealth among health centers during the COVID-19 pandemic - United States, June 26-November 6, 2020. *MMWR Morb Mortal Wkly Rep*. 2021 Feb 19;70(7):240-4. doi: 10.15585/mmwr.mm7007a3. PMID: 33600385.
187. Weiner JP, Bandeian S, Hatef E, et al. In-person and telehealth ambulatory contacts and costs in a large US insured cohort before and during the COVID-19 pandemic. *JAMA Netw Open*. 2021 Mar 1;4(3):e212618. doi: 10.1001/jamanetworkopen.2021.2618. PMID: 33755167.
188. Hammond G, Luke AA, Elson L, et al. Urban-rural inequities in acute stroke care and in-hospital mortality. *Stroke*. 2020 Jul;51(7):2131-8. doi: 10.1161/strokeaha.120.029318. PMID: 32833593.
189. Howard G, Kleindorfer DO, Cushman M, et al. Contributors to the excess stroke mortality in rural areas in the United States. *Stroke*. 2017 Jul;48(7):1773-8. doi: 10.1161/strokeaha.117.017089. PMID: 28626048.