

Systematic Review on Noninvasive Nonpharmacological Treatment for Chronic Pain: Surveillance Report 1

Update Period: August 2019 through September 2021

Background and Purpose

This is the first update for the 2020 report, *Noninvasive Nonpharmacologic Treatments for Chronic Pain*,¹ (available at <https://effectivehealthcare.ahrq.gov/products/noninvasive-nonpharm-pain-update/research>), covering the period August 2019 through September 2021. The 2020 report addressed benefits and harms of noninvasive nonpharmacological therapy for five common chronic pain conditions: low back pain (LBP), neck pain, knee, hip or hand osteoarthritis (OA), fibromyalgia (FM), and tension headache. Given the clinical and public health importance of this topic, it is important to identify new evidence that could impact practice or policy. The purpose of this surveillance report is to identify new evidence published since the 2020 report and to determine how the new evidence impacts findings of the prior report. Subsequent surveillance reports are planned for January 2022 (based on evidence published from October to December 2021) and April 2022 (based on evidence published from January to March 2022).

Scope

The scope and eligibility criteria established at the time of the original report¹ were utilized for this surveillance report; no changes were made. That report included randomized controlled trials (RCTs) reporting outcomes at least 1 month following the completion of treatment and focused on the use of single, active, noninvasive nonpharmacological interventions (including exercise, mind-body practices, psychological therapies, mindfulness practices, manual therapies, physical modalities, acupuncture, and multidisciplinary rehabilitation) for adults with five common chronic pain conditions, LBP (Key Question 1), neck pain (Key Question 2), knee, hip or hand OA (Key Question 3), FM (Key Question 4), and tension headache (Key Question 5), and addressed:

- whether the interventions work overall compared with sham, waitlist control, attention control, no treatment, or usual care;
- whether the interventions work compared with pharmacological alternatives; and
- how outcomes for individual interventions (e.g., acupuncture) compare with a common comparator (exercise for LBP, neck pain, OA and FM; biofeedback for headache).

In addition, Key Question 6 addressed whether estimates of benefits and harms differ by age, sex, presence of comorbidities (e.g., emotional or mood disorders), or degree of nociplasticity/central sensitization. The full protocol for the original report including detailed inclusion criteria using the PICOTS framework

(<https://www.ncbi.nlm.nih.gov/books/NBK556223/table/ch3.tab1/?report=objectonly>) and full

Key Questions (<https://www.ncbi.nlm.nih.gov/books/n/cer227/ch2/#ch2.s2>) are available on the Agency for Healthcare Research and Quality website (<https://effectivehealthcare.ahrq.gov/products/noninvasive-nonpharm-pain-update/protocol>) and on the PROSPERO systematic reviews registry (CRD42019132457).

Methods

Update searches were conducted to identify evidence published from August 2019 through September 2021. Search strategies from the original report were utilized¹ and we searched the same databases as in the original report (Ovid[®] MEDLINE[®], Cochrane Central Register of Controlled Trials, and Cochrane Database of Systematic Reviews). In addition, to capture articles not yet indexed in Medline, we supplemented the original search strategies with a previously developed² optimized (text-word only) search in pre-Medline, to identify studies not yet indexed with Medical Subject Headings (MeSH). As in the original report, searches on electronic databases were supplemented by review of reference lists of relevant articles. Search strategies are available in Appendix A.

As in the original review, one investigator screened citations identified through searches for eligibility for full-text review (Key Questions and inclusion criteria are available in Appendix B). In addition, to increase efficiency of abstract review, we utilized a machine learning classifier in conjunction with a second investigator to assist in conducting dual reviews. The machine learning classifier was previously shown to have 100 percent recall for identifying eligible studies in update searches for this review.² The machine learning classifier screened all citations; the second investigator performed dual review on all studies except for those classified by the machine learning classifier as very low probability. Any citation identified as potentially eligible by either investigator underwent full-text review to determine final eligibility.

We utilized the same methods for data abstraction and quality assessment as the original report. As in the prior review, we assessed the risk of bias of RCTs using the approaches recommended in the *Cochrane Handbook for Systematic Reviews of Interventions* (Chapter 8.5 Risk of Bias Tool)³ and the *AHRQ Methods Guide for Effectiveness and Comparative Effectiveness Research*,⁴ in conjunction with criteria and methods developed by the Cochrane Back and Neck Group.⁵ The decision to update meta-analyses from the original report was based on the number and sample sizes of new studies eligible for meta-analysis (meta-analysis performed if new evidence was large relative to the studies in the original meta-analysis); consistency in findings between the new studies and the original meta-analysis (meta-analysis performed if findings from new evidence appear inconsistent and new studies were appropriate for pooling based on similarity in populations, interventions, and comparisons, in order to determine whether new studies impact conclusions); or if new evidence could impact the strength of evidence (SOE) (meta-analysis performed if the SOE based on the original meta-analysis was low or insufficient and new evidence could increase the strength of evidence due to increased precision, quality, or other factors). The SOE was based on the totality of evidence (evidence in the original report plus new evidence) and determined using the methods described in the original report. We highlighted any changes in the SOE assessments.

A list of studies included in this surveillance report is provided in Appendix C. Evidence tables providing data from included studies are available in Appendix D, and quality assessments for each study are shown in Appendix E. A list of articles excluded at full-text review along with reasons for exclusion is available in Appendix F. Updated and new meta-analyses can be found in Appendix G.


Results

The update search yielded 5,391 total citations and identified 29 new eligible citations (**Figure 1**). Of those, 25 were new RCTs⁶⁻³⁰ (3 good, 19 fair and 7 poor quality) which reported on primary outcomes of interest, and four (3 LBP, 1 FM) were subsequent publications³¹⁻³⁴ for trials already included in the prior report and reported only secondary outcomes. This surveillance report focuses on the primary outcomes and changes in strength of evidence (SOE) based on new evidence. Secondary outcomes are not summarized in this Addendum but can be found in Appendix E.

For **chronic LBP**, a total of 10 new RCTs^{7,14,15,17,18,20,22,27,28,30} (9 fair and 1 poor quality) were included which compared exercise versus usual care (2 RCTs)^{17,27}; psychological therapy versus usual care (2 RCTs)^{7,27} and versus exercise (1 RCT)²⁷; low-level laser therapy (LLLT) versus sham (1 RCT)¹⁴; transcutaneous electrical nerve stimulation versus sham (1 RCT)³⁰; spinal manipulation versus sham (1 RCT)²⁸; massage versus usual care (1 RCT)¹⁵; yoga versus exercise (2 RCTs)^{20,22}; and acupuncture versus usual care (1 RCT).¹⁸ Only one new, fair-quality RCT was included for **chronic neck pain** and compared manual therapy versus sham and versus exercise.⁸ For **knee OA**, a total of 9 new RCTs^{6,12,13,16,19,21,24,26,29} (3 good, 3 fair and 3 poor quality) were included which compared exercise versus usual care or attention control (3 RCTs)^{19,21,26}; massage versus usual care (1 RCT)²⁴; ultrasound versus sham (1 RCT)¹³; LLLT versus sham (1 RCT)⁶; interferential current versus sham (1 RCT)⁶; Tai Chi versus usual care (1 RCT)¹²; Qigong versus exercise (1 RCT)²⁹; and acupuncture versus sham (1 RCT).¹⁶ For **FM**, a total of five new RCTs^{9-11,23,25} (2 fair and 3 poor quality) were included which compared exercise versus usual care (2 RCTs)^{11,23}; spinal manipulation versus sham (1 RCT)¹⁰; mindfulness-based stress reduction versus usual care (1 RCT)²⁵; basic body awareness training versus usual care (1 RCT)⁹; acupuncture versus usual care and versus exercise (1 RCT)¹¹; and multidisciplinary rehabilitation (MDR) versus usual care and versus exercise (1 RCT).²³ One LBP trial,²⁷ one knee OA trial⁶ and two FM trials^{11,23} had more than two arms and provided data for multiple interventions and/or comparators; thus the number of RCTs cited above are not mutually exclusive.

There were no new trials identified for hip or hand OA or chronic tension-type headache or that compared a nonpharmacological therapy with pharmacological therapy. There was no new evidence for the following interventions for LBP (mindfulness-practices, MDR), neck pain (psychological therapies, physical modalities, mindfulness practices, mind-body practices, acupuncture, and MDR), knee OA (psychological therapies, mindfulness practices, MDR) and FM (psychological therapies, physical modalities). No trials in pregnant or breastfeeding women with pre-existing chronic pain or new trials comparing interventions with topical agents, medical cannabis or muscle relaxants were identified (all Key Questions). No data were available to evaluate nociplasticity as a modifier to treatment effectiveness or safety (Key Question 6).

Figure 1. Literature Flow Diagram



^a Cochrane databases include the Cochrane Central Register of Controlled Trials and the Cochrane Database of Systematic Reviews

^b Other sources include prior reports, reference lists of relevant articles, systematic reviews, etc.

^c Publications may be included or excluded for multiple interventions

^d Used as source documents, studies checked for inclusion eligibility

^e These 4 followup publications (to Groessl 2017, Saper 2017 [chronic low back pain] and McCrae 2019 [fibromyalgia] included in the 2020 report) reported only secondary outcomes of interest and are not summarized in this surveillance report but details can be found in Appendix D.

^f The 4 followup publications are not counted in the number of included trials but are counted under total publications.

Evidence Summary

- Twenty-five new RCTs which reported on primary outcomes of interest were identified for this update (10 in chronic LBP, 1 in chronic neck pain, 9 in knee OA, and 5 in FM). Nine of the new trials (2 in chronic LBP, 1 in chronic neck pain, 2 in knee OA, and 4 in FM; noted below) evaluated interventions, comparators, or timepoints not previously evaluated; otherwise, the trials evaluated comparisons with at least some prior evidence.

Chronic Low Back Pain

- **Exercise:** The addition of two new fair-quality trials comparing exercise with usual care did not change the previous report's conclusions of a small improvement in *function* short term (SOE: moderate) nor did it change the conclusion of no effect with the addition of one new fair-quality trial at intermediate term (SOE: low) based on updated meta-analyses. Similarly, data from these trials did not change the prior report's conclusion of a small improvement in *pain* with exercise short term (adding both new trials) or intermediate term (1 new trial) compared with usual care (SOE: low for both timepoints).
- **Psychologic therapies:** The inclusion of one new fair-quality RCT of pain reprocessing therapy (PRT) to our previous meta-analysis resulted in a similar effect size and conclusion of a small improvement in *function* with psychological therapies short term as found in the 2020 report. We downgraded the SOE from moderate to low based on increased inconsistency. The other new trial of relaxation therapy was excluded from the updated meta-analysis of function short term as an outlier (the effect estimate was substantially larger than those in other included trials). Addition of the PRT study to our previous meta-analysis did not change our previous conclusions of small improvements in *function* at intermediate term and long term (SOE: moderate). PRT was associated with small improvements in *pain* (0 to 10 scale) compared with usual care at short term, intermediate term and long-term (SOE: moderate) which is consistent with the conclusion of the 2020 report. Evidence was insufficient for one new small fair-quality trial comparing psychological therapy with exercise.
- **Physical modalities, low-level laser therapy:** The addition of one new, small poor-quality trial of low-level laser therapy versus sham did not change previous report's conclusions of a small improvement in *function* and moderate improvement in *pain* short term (SOE: low).
- **Physical modalities, transcutaneous electrical nerve stimulation:** The prior report did not contain studies of this modality. One new, fair-quality RCT of transcutaneous electrical nerve stimulation (TENS) compared with sham TENS found no difference in function or pain short term (SOE: low).
- **Manual therapies, spinal manipulation:** Effect size based on updated meta-analysis including one new fair-quality RCT of manipulation and sham laser is consistent with the small improvement in *function* observed short term in the prior report. Similarly, the new trial's finding of no difference in *pain* short term was consistent with the 2020 report. Strength of evidence remained low.
- **Manual therapies, massage:** One new, small fair-quality RCT of massage versus usual care found no difference between groups for function or pain short term. Its addition did

not change our previous conclusion of small improvements in *function and pain* short term (SOE: moderate).

- **Mind-body practices, yoga:** Two new fair-quality RCTs found no differences between yoga and exercise for *function or pain* short term which is consistent with findings of no effect from the 2020 report; the SOE remained low
- **Acupuncture:** One new fair-quality RCT compared two different forms of acupuncture with usual care. Standard acupuncture was associated with moderate improvements in *function and pain* while hand and ear acupuncture was associated with large improvements in *function and pain* versus usual care at short-term followup. Findings from this new RCT are consistent with our prior meta-analyses for the comparison with usual care; SOE remained low for function and moderate for pain

Chronic Neck Pain

- **Exercise:** One new, small fair-quality trial found exercise associated with moderate improvement in function and large improvement in pain versus a sham procedure short term; given the small sample size, it did not change our prior conclusions of no effect for *function or pain* (SOE low).
- **Manual therapies (manipulation):** One new, small fair-quality trial found that manipulation was associated with moderate improvement in *function* and large improvement in *pain* short term compared with a sham procedure (suboccipital inhibition). There was no difference in either *function or pain* short term between manual therapy and exercise. The prior report did not include comparisons of manipulation with sham treatment or with exercise. SOE was low for function and pain across both comparisons.

Knee OA

- **Exercise:** Addition of one poor-quality and two fair-quality RCTs comparing exercise with usual care or attention control in patients with knee OA did not change the 2020 report's conclusions or SOE for small improvement in *function and pain* short- and long-term and moderate improvements in both at intermediate term (SOE moderate for both at short term and low at intermediate and long term).
- **Physical modalities, low-level laser therapy:** One new, good-quality RCT compared LLLT versus sham. Its incorporation into a new meta-analysis for *function* revealed a small improvement at short and intermediate term. Its addition to a prior meta-analysis of two trials for *pain* did not change prior findings of no difference at short or intermediate term. With the addition of the new trial, the SOE was upgraded from insufficient to low for all
- **Physical modalities, ultrasound:** One new good-quality RCT comparing ultrasound with sham reported no difference between groups in *function or pain* short-term; this is consistent with the prior report and SOE remained low.
- **Physical modalities, inferential current:** No trials of this modality were identified for the prior report. The new good-quality RCT found no improvement in either *function or*

pain at short or intermediate terms for interferential current compared with sham (SOE: low).

- **Manual therapies, massage:** One new, small poor-quality RCT of massage versus usual care did not change our conclusion of insufficient evidence.
- **Mind-body therapies, Tai Chi and Qigong:** One new, small poor-quality RCT comparing Tai Chi with usual care does not change conclusions of insufficient evidence from the prior report. Evidence for a new comparison of Qigong with exercise from one fair-quality trial was considered insufficient.
- **Acupuncture:** One new, good quality RCT of acupuncture versus sham reported no difference between groups in *function or pain* short term; this is consistent with the prior report and SOE remained low.

Fibromyalgia

- **Exercise:** Two new poor-quality trials comparing exercise with usual care reported no difference between groups in short term function. Given the poor quality of these trials, our conclusions for a small improvement in *function* short term (SOE: low) did not change.
- **Manual therapy, spinal manipulation.** The prior report did not contain evidence for spinal manipulation versus sham manipulation. One new fair-quality RCT found no difference between groups in either *function or pain* at short or intermediate term (SOE: low).
- **Mindfulness practices, mindfulness-based stress reduction:** One new, poor-quality trial reported a small improvement in *function* following mindfulness-based stress reduction compared with usual care. This is consistent with the original report and SOE remained low.
- **Mind-body therapies, basic body awareness therapy:** No trials of this modality were identified for the prior report. Evidence from one new, small fair-quality trial of basic body awareness therapy was considered insufficient.
- **Acupuncture:** The prior report did not contain evidence for acupuncture versus usual care or versus exercise. Evidence from one new poor-quality RCT was considered insufficient.
- **Multidisciplinary rehabilitation:** One new poor-quality RCT compared MDR with usual care and with exercise and found no difference between MDR and either comparator in function short term. The prior report found a small improvement in *function* short term compared with usual care (SOE: low); given the poor quality of the new RCT, the 2020 report's conclusions and SOE remain unchanged. For the comparison of MDR with exercise, we graded the evidence as insufficient; the prior report did not contain evidence for this comparison at short term.

Harms

- Consistent with the 2020 report, adverse events were not consistently reported in new studies and intervention-related serious life-threatening events requiring medical attention were not reported.

Summary of New Evidence

Table 1 (below) provides the conclusions from the 2020 report and the new findings from studies identified in this surveillance report. Table 1 focuses on Key Questions and interventions with new evidence; the full strength of evidence (SOE) table is available in the prior report (<https://www.ncbi.nlm.nih.gov/books/n/cer227/appg/>).

Table 1. Summary of conclusions and assessments informed by new evidence

Condition	Intervention (Comparator)	Outcome, Timing	Prior (2020) Update ^a	2021 Surveillance Report ^a	Change
LBP	Exercise (vs. UC, AC or placebo)	Function, short term	Small effect SOE: Moderate 10 RCTs (N=940) [Excluding an outlier trial]	2 fair-quality RCTs (N=147) ^{17,27} MA updated 12 RCTs (N=1,011)	No change in conclusions
		Function, intermediate term	No effect SOE: Low 5 RCTs (N=616)	1 fair-quality RCT (N=96) ¹⁷ MA updated 6 RCTs (N=712)	No change in conclusions
		Pain, short term	Moderate effect SOE: Low 11 RCTs (N=981)	1 fair-quality RCT (N=111) ¹⁷ No effect	No change in conclusions
		Pain, intermediate term	Small effect SOE: Low 5 RCTs (N=616)	1 fair-quality RCT (N=96) ¹⁷ No effect	No change in conclusions
	Psych vs. UC	Function, short term	Small effect SOE: Moderate 3 RCTs (N=906)	2 fair-quality RCTs (N=146) ^{7,27} MA updated 4 RCTs (N=1,018) ^b	Small effect SOE: Low (downgraded one level)
		Function, intermediate term	Small effect SOE: Moderate 3 RCTs (N=1,026)	1 fair-quality RCT (N=116) ⁷ MA updated 4 RCTs (N=1,142)	No change in conclusions
		Function, long term	Small effect SOE: Moderate 3 RCTs (N=815)	1 fair-quality RCT (N=124) ⁷ MA updated 4 RCTs (N=939)	No change in conclusions
		Pain, short term	Small effect SOE: Moderate 3 RCTs (N=906)	1 fair-quality RCT (N=112) ⁷ Small effect	No change in conclusions
		Pain, intermediate term	Small effect SOE: Moderate 3 RCTs (N=1,026)	1 fair-quality RCT (N=116) ⁷ Small effect	No change in conclusions
		Pain, long term	Small effect SOE: Moderate 3 RCTs (N=815)	1 fair-quality RCT (N=124) ⁷ Small effect	No change in conclusions
		Psych vs. Exercise	Function, short term	No evidence at short term	1 fair-quality RCT (N=34) ²⁷ No effect
	Physical modalities: LLLT vs. sham	Function, short term	Small effect SOE: Low 1 RCT (N=56)	1 poor-quality RCT (N=34) ¹⁴ Large effect	No change in conclusions
		Pain, short term	Moderate effect SOE: Low 1 RCT (N=56)	1 poor-quality RCT (N=34) ¹⁴ Large effect	No change in conclusions

Condition	Intervention (Comparator)	Outcome, Timing	Prior (2020) Update ^a	2021 Surveillance Report ^a	Change	
	Physical modalities: burst TENS vs. continuous TENS vs. sham	Function, short term	No evidence	1 fair-quality RCT (N=73) ³⁰ No effect	New intervention No effect SOE: Low	
		Pain, short term	No evidence	1 fair-quality RCT (N=73) ³⁰ No effect	New intervention No effect SOE: Low	
	Manual Therapy: Spinal manipulation vs. sham, UC, AC	Function, short term	Small effect SOE: Low 3 RCTs (N=704)	1 fair-quality RCT (N=155) ²⁸ MA updated 4 RCTs (N=859)	No change in conclusions	
		Pain, short term	No effect SOE: Low 3 RCTs (N=530)	1 fair-quality RCT (N=155) ²⁸ No effect	No change in conclusions	
	Manual Therapy: Massage vs. sham, UC, AC	Function, short term	Small effect SOE: Moderate 6 RCTs (N=694)	1 fair-quality RCT (N=59) ¹⁵ No effect	No change in conclusions	
		Pain, short term	Small effect SOE: Moderate 5 RCTs (N=644)	1 fair-quality RCT (N=59) ¹⁵ No effect	No change in conclusions	
	Mind-body practices: Yoga vs. Exercise	Function, short term	No effect SOE: Low 4 RCTs (N=559)	2 fair-quality RCTs (N=252) ^{20,22} No effect	No change in conclusions	
		Pain, short term	No effect SOE: Low 5 RCTs (N=575)	2 fair-quality RCTs (N=252) ^{20,22} No effect	No change in conclusions	
	Acupuncture vs. sham, UC, AC	Function, short term	Small effect SOE: Low 4 RCTs (N=2,066)	1 fair-quality RCT (N=152) ¹⁸ Large effect	No change in conclusions	
		Pain, short term	Small effect SOE: Moderate 5 RCTs (N=2,109)	1 fair-quality RCT (N=152) ¹⁸ Moderate effect	No change in conclusions	
	Neck pain	Exercise vs. AC, WL or NT	Function, short term	No effect [excluding outlier] SOE: Low 3 RCTs (N=444)	1 fair-quality RCT (N=43) ⁸ Moderate effect	No change in conclusions
			Pain, short term	No effect [excluding outlier] SOE: Low 3 RCTs (N=444)	1 fair-quality RCT (N=43) ⁸ Large effect	No change in conclusions
		Manual therapy: Manipulation vs. sham	Function, short term	No evidence	1 fair-quality RCT (N=42) ⁸ Moderate effect	New intervention Moderate effect SOE: Low
			Pain, short term	No evidence	1 fair-quality RCT (N=42) ⁸ Large effect	New intervention Large effect SOE: Low
Manual Therapy: Manipulation vs. exercise		Function, short term	No evidence	1 fair-quality RCT (N=43) ⁸ No effect	New intervention No effect SOE: Low	
		Pain, short term	No evidence	1 fair-quality RCT (N=43) ⁸ No effect	New intervention No effect SOE: Low	
Knee OA			Function, short term	Small effect SOE: Moderate	1 poor-quality RCT (N=84) ²⁶	No change in conclusions

Condition	Intervention (Comparator)	Outcome, Timing	Prior (2020) Update ^a	2021 Surveillance Report ^a	Change
	Exercise vs. UC, AC, sham or NT		8 RCTs (N=748)	MA updated 9 RCTs (N=832)	
		Function, intermediate term	Moderate effect SOE: Low 11 RCTs (N=879)	1 fair-quality RCT (N=265) ¹⁹ MA updated 12 RCTs (N=1,144)	No change in conclusions
		Function, long term	Small effect SOE: Low 4 RCTs (N=1,199)	2 fair-quality RCTs (N=342) ^{19,21} MA updated 6 RCTs (N=1,541)	No change in conclusions
		Pain, short term	Small effect SOE: Moderate 8 RCTs (N=748)	1 poor-quality RCT (N=84) ²⁶ MA updated 9 RCTs (N=832)	No change in conclusions
		Pain, intermediate term	Moderate effect SOE: Low 11 RCTs (N=879)	1 fair-quality RCT (N=261) ¹⁹ MA updated 12 RCTs (N=1,140)	No change in conclusions
		Pain, long term	Small effect SOE: Low 4 RCTs (N=1,199)	2 fair-quality RCTs (N=338) ^{19,21} MA updated 6 RCTs (N=1,537)	No change in conclusions
	Physical Modalities: LLLT vs. sham or UC	Function, short term	Insufficient evidence 1 RCT (N=49)	1 good-quality RCT (N=84) ⁶ New MA 2 RCTs (N=133)	Small effect SOE: Low (upgraded one level)
		Function, intermediate term	Insufficient evidence 2 RCTs (N=109)	1 good-quality RCT (N=84) ⁶ New MA 3 RCTs (N=193)	Small effect SOE: Low (upgraded one level)
		Pain, short term	Insufficient evidence 2 RCTs (N=76)	1 good-quality RCT (N=84) ⁶ MA updated 3 RCTs (N=160)	No effect SOE: Low (upgraded one level)
		Pain, intermediate	Insufficient evidence 2 RCTs (N=109)	1 good-quality RCT (N=84) ⁶ MA updated 3 RCTs (N=193)	No effect SOE: Low (upgraded one level)
	Physical Modalities: continuous and pulsed US vs. sham	Function and pain, short term	No effect SOE: Low 3 RCTs (N=249)	1 good-quality RCT (N=75) ¹³ No effect	No change in conclusions
		Physical Modalities: Interferential current vs. sham	Function, short and intermediate term	No evidence	1 good-quality RCT (N=84) ⁶ No effect
	Pain, short and intermediate term		No evidence	1 good-quality RCT (N=84) ⁶ No effect	New intervention No effect SOE: Low
Manual Therapies: Massage vs. UC	Function, short term	Insufficient evidence 1 RCT (N=125)	1 poor-quality RCT (N=60) ²⁴ No effect	No change in conclusions	

Condition	Intervention (Comparator)	Outcome, Timing	Prior (2020) Update ^a	2021 Surveillance Report ^a	Change
		Pain, short term	Insufficient evidence 1 RCT (N=125)	1 poor-quality RCT (N=60) ²⁴ Small effect	No change in conclusions
	Mind-body Therapies: Tai Chi vs. AC or UC	Function and pain, Intermediate term	Insufficient evidence 1 RCT (N=40)	1 poor-quality RCT (N=92) ¹² Moderate effect	No change in conclusion
	Mind-body Therapies: Qigong vs. Exercise	Function and pain, Intermediate term	No evidence vs. exercise	1 fair-quality trial (N=68) ²⁹ No effect	New comparator Insufficient evidence
	Acupuncture vs. UC, NT, or sham	Function, short term	No effect SOE: Low 5 RCTs (N=944)	1 good-quality RCT (N=83) ¹⁶ No effect	No change in conclusions
		Pain, short term	No effect SOE: Low 6 RCTs (N=1065)	1 good-quality RCT (N=83) ¹⁶ No effect	No change in conclusions
FM	Exercise vs. UC, NT, sham or AC	Function, short term	Small Effect SOE: Low 7 RCTs (N=410)	2 poor-quality RCTs (N=135) ^{11,23} No effect	No change in conclusions
	Manual Therapy: Spinal manipulation vs. sham	Function, short and intermediate term	No evidence	1 fair-quality RCT (N=101) ¹⁰ No effect	New intervention No effect SOE: Low
		Pain, short and intermediate term	No evidence	1 fair-quality RCT (N=101) ¹⁰ No effect	New intervention No effect SOE: Low
	Mindfulness Practices: MBSR, MAT vs. AC or WL	Function, intermediate term	Small effect SOE: Low 1 RCT (N=148)	1 poor-quality RCT (N=98) ²⁵ Small effect	No change in conclusion
	Mind-Body Therapies: BBAT vs. UC	Pain, short and intermediate term	No evidence	1 fair-quality RCT (N=39) ⁹ No effect	New intervention Insufficient evidence
	Acupuncture vs. UC	Function, short term	No evidence vs. usual care ^c	1 poor-quality RCT (N=67) ¹¹ No effect	New comparator Insufficient evidence
	Acupuncture vs. Exercise	Function, short term	No evidence vs. exercise	1 poor-quality RCT (N=67) ¹¹ No effect	New comparator Insufficient evidence
	MDR vs. UC or WL	Function, short term	Small effect SOE: Low 3 RCTs (N=381)	1 poor-quality RCT (N=64) ²³ No effect	No change in conclusion
	MDR vs. Exercise	Function, short term	No evidence at short term	1 poor-quality RCT (N=64) ²³ No effect	New timepoint (short term) Insufficient evidence

Abbreviations: AC = attention control; BBAT = basic body awareness training; LLLT = low-level laser therapy; MA = meta-analysis; MAT = meditation awareness training; MBSR = mindfulness-based stress reduction; MDR = multidisciplinary rehabilitation; NT = no treatment; RCT = randomized controlled trial; SOE = strength of evidence; TENS = transcutaneous electrical nerve stimulation; UC = usual care; US = ultrasound; vs. = versus; WL = waitlist.

^a The sample size (N) reported is as analyzed.

^b One new trial, Shariat 2019, was an outlier and conclusions are based on inclusion of the one new fair-quality RCT (Ashar 2021) which increased heterogeneity leading to downgrade from moderate to low SOE.

^c In the previous report, 3 RCTs were included that compared acupuncture with sham acupuncture and found a small improvement in function short term (Moderate SOE).

Evidence Details

Key Question 1: Chronic Low Back Pain

Exercise for Chronic Low Back Pain

Exercise Compared With Usual Care, an Attention Control, or a Placebo Intervention

Two new fair-quality trials of exercise (pedometer-based walking program [5 consecutive days per week for 12 weeks] and mobility/flexibility exercises [18 sessions over 6 weeks]) versus usual care were identified.^{17,27} One moderate sized RCT (n=96)¹⁷ contributed data on function and pain short and intermediate term. The other smaller RCT (n=36)²⁷ provided data for short-term function only.

In the short term, updated pooled estimates across 12 trials (n=1,011) did not change our previous conclusions of a small improvement in *function* (SOE moderate) with exercise versus usual care (pooled standardized mean difference [SMD] -0.36, 95% CI -0.58 to -0.17, I²=47.3% excluding the same outlier trial as the 2020 report). One of the new trials also reported results at intermediate term.¹⁷ The conclusion of no difference in function between groups (6 RCTs, n=712; updated pooled SMD -0.18, 95% CI -0.42 to 0.03 I²=31.3%) is consistent with the 2020 report and SOE remained low. (See Appendix G, Figure G-1 for updated function plot.) The same trial also found no difference in *pain* between exercise and usual care at short term (n=111; difference -0.6 on a 0 to 10 scale, 95% CI -1.26 to 0.06) or intermediate term (n=96; difference -0.8 on a 0 to 10 scale, 95% CI -1.60 to 0.01). The addition of these data did not change the prior report's conclusion of a small improvement in pain with exercise short term (11 RCTs) or intermediate term (5 RCTs) compared with usual care or SOE of low at either time point.

Of the two new trials, only one reported that no intervention-related adverse events occurred.¹⁷

Psychological Therapies for Chronic Low Back Pain

Psychological Therapies Compared With Usual Care or an Attention Control

Psychological therapies were associated with small functional improvements compared with usual care or attention control at short, intermediate, and long term (SOE moderate for all) in the 2020 report. CBT was the most common psychological treatment reported and three trials reported for each time frame. We identified two new fair-quality trials comparing psychological therapies with usual care that reported on *function*. Patients received relaxation therapy for three, 40–45-minute sessions per week for 6 weeks or continued usual care in one small RCT from Iran (N=34).²⁷ Patients in the other, United States-based trial (N=151 randomized)⁷ received a proprietary pain reprocessing therapy (PRT) that aims to promote patients' reconceptualization of primary chronic pain as a brain-generated false alarm or usual care which consisted of a chronic pain workbook or videos describing placebo treatments combined with an open-label subcutaneous saline injection at the site of greatest back pain. The PRT group received individual hourly sessions delivered twice weekly for 8 weeks.

An updated meta-analysis including the two new trials showed a moderate improvement in short-term *function* (5 RCTs, n=1,052, pooled SMD -0.58, 95% CI -1.27 to 0.01, I²=90.7%) and a substantial increase in heterogeneity compared with the original analyses' I² of 0%. The effect estimate from the new trial of relaxation therapy²⁷ was substantially larger (SMD -1.89) than those in other included trials (SMD range from -0.08 to 0.93). Exclusion of this outlier resulted in less heterogeneity and a smaller pooled effect size (4 RCTs, n=1,018, pooled SMD -0.34, 95% CI -0.74 to 0.02, I²=70.5%) which was more consistent with, but larger than, the original report (prior report analysis, 3 RCTs, n=906, pooled SMD -0.24, 95% CI -0.38 to -0.04, I²=0%); see Appendix G, Figure G-2 for updated plot. While the pooled estimate from the updated meta-analysis including the new PRT trial was similar to the prior report estimate, the heterogeneity was substantially higher. This may in part be due to differences in the therapeutic approach in the PRT trial compared with previously included trials. The bulk of the evidence for psychological therapies in the original report was from large trials focused on cognitive behavioral therapy (CBT). While PRT incorporates aspects of CBT, it also includes other psychological interventions used in pain management and emphasizes exposure-based interventions in conjunction with reappraising pain. Only the new trial of PRT⁷ reported on intermediate- and long-term function. Updated meta-analyses to include this RCT resulted in a small increase in effect size for each time frame but did not change conclusions from the original report that psychological therapies were associated with small improvements in *function* at intermediate term (4 RCTs, n=1,142, pooled SMD -0.30, 95% CI -0.60 to -0.11, I²=0%) and long term (4 RCTs, n=939, pooled SMD -0.31, 95% CI -0.47 to -0.18, I²=0%). PRT was associated with small improvements in *pain* (0 to 10 scale) compared with usual care at short term (difference -0.95, 95% CI -1.43 to -0.46), intermediate term (difference -0.91, 95% CI -1.37 to -0.44) and long term (difference -0.89, 95% CI -1.34 to -0.42), which is consistent with the conclusion of the 2020 report. No adverse events were observed for PRT; the other trial did not report on adverse events.

Psychological Therapies Compared With Exercise

One new small, fair-quality trial (n=36)²⁷ compared relaxation training versus mobility/flexibility exercises (modified from the McKenzie method) over 6 weeks (18 sessions total for both) and found no difference between groups in *function* at short-term followup (difference 0.00 on the 0 to 40 Functional Rating Index, 95% CI -1.09 to 1.09). This is the only trial identified to date evaluating short term outcomes for this comparison. Data were considered insufficient to draw conclusions. Harms were not reported.

Physical Modalities for Chronic Low Back Pain

Low-Level Laser Therapy Compared With Sham

One new poor-quality small (n=34) industry funded trial from Iran compared low-level laser therapy (LLLT) with sham laser.¹⁴ It was rated as poor quality primarily due to unclear concealment of treatment allocation, lack of patient and caregiver blinding and differential loss to followup. They reported a large improvement in *function* (difference -5.70 on 0 to 24 Roland Morris Disability Questionnaire [RDQ], 95% CI -8.47 to -2.93) and large improvement in *pain* (difference -4.40 on a 0 to 10 scale, 95% CI -5.31 to -3.49) short term. The 2020 report found that LLLT was associated with a small improvement in *function* and moderate improvement in *pain* short term (SOE low for both) compared with sham laser based on one small (n=56) fair-quality trial.³⁵ Given the small size and poor quality of the new trial, conclusions from the

original report did not change. In the new trial, about 52% of the patients in the true laser group reported a temporary increase in pain following the first treatment session. Other adverse events were all observed in the sham laser group; authors intimate that they were related to concomitant use of naproxen by participants.

Transcutaneous Electrical Nerve Stimulation Compared With Sham

There were no trials of transcutaneous electrical nerve stimulation (TENS) for chronic LBP in the prior report. One new, fair-quality trial (n=73)³⁰ compared two applications of TENS, burst and conventional, versus sham TENS and found no differences between groups over the short term in *function* as measured by the modified Oswestry Disability Index (ODI) (burst vs. sham TENS: difference -2.90 on a 0 to 50 scale, 95% CI -7.97 to 2.17; conventional vs. sham TENS: difference -2.30 on a 0 to 50 scale, 95% CI -7.77 to 3.17) or in *pain* (burst vs. sham TENS: difference -0.80 on a 0 to 10 scale, 95% CI -2.24 to 0.64; conventional vs. sham TENS: difference -1.30 on a 0 to 10 scale, 95% CI -2.74 to 0.14). Results were similar across other measures of pain. SOE was low. No TENS-associated side-effects occurred in any patient.

Manual Therapies for Chronic Low Back Pain

Spinal Manipulation Compared With Sham Manipulation, Usual Care, an Attention Control, or a Placebo Intervention

One new good-quality RCT (N=155)²⁸ compared groups receiving spinal manipulation, mobilization or sham laser treatment and reported on function and pain short-term. Manipulation consisted of rapid thrust applied to the shoulder and pelvis, mobilization consisted of having the patient gently push the shoulder and pelvis into the clinician's hands. The sham laser treatment was done with the patient in same position as the active interventions and the machine appeared operational to patients and clinicians but delivered no energy. All sessions were twice weekly for 3 weeks. There were no differences in mean change scores for *function* (RDQ, 0 to 24 scale) between manipulation and sham treatment (-0.07, 95% CI -1.43 to 1.29) or mobilization and sham treatment (0.9, 95% CI -0.41 to 2.29). Given this, and the fact that studies in the 2020 report appear to have used a combination of manipulation and mobilization techniques, these active treatment groups were combined to update the meta-analysis. Inclusion of the new trial decreased the effect size estimate somewhat for *functional* improvement (4 RCTs, n=859, pooled SMD -0.24, 95% CI -0.61 to 0.09, I²= 61.8%) and there was greater imprecision compared with the 2020 report (3 RCTs, n=704, pooled SMD -0.34, 95% CI -0.75 to -0.02, I²= 44.6%); see Appendix G, Figure G-3 for updated plot. Our prior conclusion of a small improvement in *function* and low SOE did not change.

There were no differences in mean change scores for numerical rating scale (NRS) *pain* (0 to 10 scale) short term for either manipulation versus sham (difference -0.03, 95% CI -0.65 to 0.59) or mobilization versus sham (difference -0.26, 95% CI -0.38 to 0.85). In the 2020 report, there was no difference in pain short term between spinal manipulation and placebo/sham intervention or attention control across three studies. The addition of the new study did not change this conclusion and SOE remained low. Adverse events were not reported.

Massage Compared With Sham Massage, Usual Care, or an Attention Control

One new, fair-quality trial (n=59)¹⁵ compared Shiatsu massage (4 weekly 1 hour sessions) with usual care and found no differences between groups in *function* (difference in change scores -1.60 on the 0 to 24 RDQ, 95% CI -3.27 to 0.07) or *pain* (difference in change scores -0.80 on a

0 to 10 scale, 95% CI -1.92 to 0.32) at short-term followup. The addition of the new trial did not change the prior report's conclusion (small effect for *function and pain* short term) and SOE remained moderate. Regarding harms, in the Shiatsu massage group, 10% of patients reported muscle pain and 3% reported headache; in the usual care group, 3% of patients reported each of the following events: dizziness, herpes zoster and abdominal pain.

Mind-Body Practices for Chronic Low Back Pain

Yoga Compared With Exercise

Two new fair-quality trials (N=252)^{20,22} comparing yoga with combination exercise (strengthening and stretching with or without stabilization) found no differences between groups in *function or pain* at short-term followup, consistent with the findings of the prior report (SOE remained low). Across trials, patients underwent weekly sessions of different duration (35 and 75 minutes) over 6 and 8 weeks. Both trials reported *function* according to the RDQ (0 to 24 scale); the difference in change scores from baseline was -0.06 (95% CI -1.55 to 1.43) in the larger trial (n=182),²⁰ and in the second trial (n=70) the median (interquartile range [IQR]) scores were 6 (3.5 to 12) versus 6 (1 to 11), p=0.26.²² For *pain*, the difference in change scores was 0.04 on a 0 to 10 VAS (95% CI -0.50 to 0.58) in the larger trial²⁰ and in the second trial the median (IQR) scores on the Defense and Veterans Pain Rating Scale (0 to 10 scale) were 4 (2.5 to 5) versus 4 (2 to 5), p=0.70.²²

No serious adverse events occurred in either trial. The larger trial reported significantly fewer minor events in the yoga group compared with the exercise group which included mild and self-limited exacerbating back pain (7% vs. 30%; RR 0.23, 95% CI 0.10 to 0.50), general pain (4% vs. 18%; RR 0.33, 95% CI 0.11 to 1.00), and muscle pain (0% vs. 8%, p=0.001).²⁰ The second trial reported that three patients experienced a slight increase in pain; the authors did not indicate to which group the patients were randomized.²²

Acupuncture for Chronic Low Back Pain

Acupuncture Compared With Sham Acupuncture, Usual Care, an Attention Control or a Placebo Intervention

One new, fair-quality trial (n=43)¹⁸ compared two types of acupuncture, hand-ear acupuncture and standard acupuncture, with usual care for 7 weeks. At short-term followup, hand-ear acupuncture was associated with large improvements in both *function* (difference -7.34 on the 0-24 RDQ, 95% CI -9.50 to -5.18) and *pain* (difference -2.29 on a 0 to 10 scale, 95% CI -3.01 to -1.57) and standard acupuncture was associated with moderate improvements in both *function* (difference -4.89 on the 0-24 RDQ, 95% CI -7.47 to -2.31) and *pain* (difference -1.15 on a 0 to 10 scale, 95% CI -1.86 to -0.44) compared with usual care. The addition of the new trial did not change the prior report's conclusion (small effects short term) and SOE remained low for function and moderate for pain.

One patient (2%) who received hand-ear acupuncture complained of needling pain and two patients (4%) in the standard acupuncture group experienced anxiety. No other complications were reported.

Key Question 2: Chronic Neck Pain

Exercise for chronic neck pain

Exercise Compared With No Treatment, Waitlist, an Attention Control or Sham Intervention

One new small, fair-quality trial (n=43),⁸ compared therapeutic muscle performance exercise (21 total sessions) with sham suboccipital inhibition for 3 weeks in patients with nonspecific chronic neck pain.⁸ Exercise was associated with a moderate improvement in *function* (difference -16.81 on the 0 to 100 Neck Disability Index [NDI], 95% CI -25.00 to -8.62) and a large improvement in *pain* (difference -2.43 on a 0 to 10 scale, 95% CI -2.69 to -2.17) over the short term. Given the small size of the new trial, its addition did not change the prior report's conclusion (no effect for pain and function) and SOE remained low.

Manual Therapies for chronic neck pain

Spinal Manipulation Compared With Sham Manipulation and With Exercise

There were no trials of spinal manipulation for chronic nonspecific neck pain in the prior report. One new small, fair-quality trial (n=42)⁸ compared spinal manipulation versus sham suboccipital inhibition and versus therapeutic exercise (n=45) for 3 weeks. Spinal manipulation was associated with a moderate improvement in *function* (difference -18.67 on the 0 to 100 NDI, 95% CI -26.04 to -11.30) and a large improvement in *pain* (difference -3.05 on a 0 to 10 scale, 95% CI -3.30 to -2.80) over the short term compared with sham manipulation but when compared with exercise therapy, there were no differences between groups for either outcome. The SOE was low. Harms were not reported.

Key Question 3: Osteoarthritis Knee Pain

Exercise for Osteoarthritis Knee Pain

Exercise Compared With Usual Care, No Treatment, Sham, or an Attention Control

Three new trials comparing group muscle performance exercises with usual care or attention control in patients with knee OA were identified.^{19,21,26} Aquatic exercise was compared with usual care in two trials: cycling 45 minutes twice weekly for 12 weeks (24 sessions) was done in one poor-quality trial (N=102 randomized)²⁶ and one-hour sessions of lower limb resistance training three times per week for 4 months (total of 48 sessions) was done in the other fair-quality trial in post-menopausal women (N=87 randomized).²¹ In the third, fair-quality trial (N=377 randomized),¹⁹ 60-minute high-intensity or low-intensity strength training sessions three times per week for 18 months were compared with an attention control consisting of 24, 60-minute group workshops over the 18 months. To update meta-analyses the high and low intensity groups were combined.

Across time frames, the addition of these new trials did not change the 2020 report's conclusions or strength of evidence regarding *function*.

At short term, the new poor-quality RCT reported no difference in *function* between exercise and usual care.²⁶ Its addition to the previous meta-analysis did not change effect estimates substantially or conclusions that exercise was associated with a small improvement in function

compared with usual care (9 RCTs, n=832, pooled SMD -0.29, 95% CI -0.44 to -0.12, $I^2=4.2\%$, SOE remained moderate). Similarly, one new fair-quality trial reporting function at intermediate term found no difference between groups¹⁹; its addition to the prior meta-analysis did not change the 2020 report's conclusions of moderate functional improvement with exercise (12 RCTs, n=1,144, pooled SMD -0.57, 95% CI -1.07 to -0.08, $I^2=92\%$, excluding the same outlier as the 2020 report, SOE remained low). Neither of the two new, fair-quality trials^{19,21} found a difference in function between exercise and usual care or attention control at long term. Their addition to the prior meta-analysis did not alter our 2020 report's conclusions of small functional improvement with exercise or low SOE (6 RCTs, n=1,541, pooled SMD -0.18, 95% CI -0.38 to -0.03, $I^2=0\%$). See Appendix G, Figure G-4 for updated function plot.

For *pain*, across the new trials, no differences between exercise and usual care or attention control were seen and their contributions to updated meta-analyses did not alter the 2020 report's conclusions. At short term, there continued to be small improvement in pain with exercise compared with usual care, attention control or sham intervention (9 RCTs, n=832, pooled difference 0.50, 95% CI -0.84 to -0.16, $I^2=37.5\%$) with the inclusion of the one new poor-quality trial²⁶ (SOE remained moderate). As seen in the 2020 report, moderate improvement in pain (0 to 10 scale) was seen at intermediate term with the addition of one new fair-quality trial (12 RCTs, n=1,140, pooled difference -1.21, 95% CI -1.96 to -0.44, $I^2=92.1\%$, SOE low) and small improvements in pain that persisted long term with the addition of the two new fair-quality trials (6 RCTs pooled difference -0.26, 95% CI -0.43 to 0.01, $I^2=0\%$, SOE remained low).^{19,21} See Appendix G, Figure G-5 for updated pain plot.

Regarding harms, one new poor-quality trial reported one patient in the exercise group with a history of cardiovascular disease hyperventilated and was hospitalized overnight and continued with training after 2 weeks of rest.²⁶ Twenty patients experienced body pain in the trial of strength training, 19 experienced falls, and 10 experienced muscle strain.¹⁹ The third trial did not report on adverse events.²¹

Manual Therapies for Osteoarthritis Knee Pain

Massage Compared With Usual Care

One new, poor-quality trial (n=60)²⁴ compared six sessions (over 3 weeks) of massage therapy with usual care in elderly patients (mean age 77 years) with knee pain due to OA. There was no difference between groups in short-term *function* according to The Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) physical function (difference -1.63 on a 0 to 68 scale, 95% CI -6.72 to 3.46) but massage was associated with a small improvement in short-term *pain* according to WOMAC pain (difference -1.65 on a 0 to 20 scale, 95% CI -2.93 to -0.37). The trial included in the prior report found no difference between massage and usual care for any outcome short term. The addition of the new poor-quality trial did not change the prior report's conclusion; the evidence remained insufficient to draw conclusions. Harms were not reported.

Physical Modalities for Osteoarthritis Knee Pain

Ultrasound Compared With Sham

One new, good-quality trial (n=75)¹³ compared 24 sessions of pulsed ultrasound (US) compared with sham US over 8 weeks and found no difference between groups in *function* according to WOMAC physical function (difference -2.47 on a 0 to 68 scale, 95% CI -8.07 to

3.13) or in *pain* either at rest (difference -0.03 on a 0 to 10 scale, 95% CI -1.13 to 1.07) or with activities of daily living (difference -0.69 on a 0 to 10 scale, 95% CI -1.95 to 0.57) at short-term followup. The addition of the new trial did not change the prior report's conclusion (no effect for function and pain short term) and SOE remained low. Harms were not reported but two (4%) patients in the sham US group withdrew due to severe knee pain.

Low-level Laser Therapy Compared With Sham

One new good-quality trial ($n=84$)⁶ compared LLLT to sham laser reported on pain and function at short and intermediate term. Evidence in the 2020 report for LLLT versus sham laser, which included one fair-quality trial ($n=49$)³⁶ and two poor-quality trials ($n=27$ and $n=60$),^{37,38} was judged to be insufficient for function and pain. There was a small improvement in *function* at short term between LLLT and sham laser in pooled analysis including the new good-quality and the previously included fair-quality trial (2 RCTs, $n=133$, pooled SMD -0.39 , 95% CI -0.80 to 0.00 , $I^2=0\%$) and at intermediate term when the new trial was added to two previously included trials (3 RCTs, $n=193$, pooled SMD -0.54 , 95% CI -1.19 to 0.05 , $I^2=46.5\%$); see Appendix G, Figure G-6 for new plot. Inclusion of the new trial in meta-analysis resulted in no clear difference in *pain* (0 to 10 scale) between LLLT and sham short term (3 RCTs, $n=160$, difference -1.50 , 95% CI -3.18 to 0.16 , $I^2=76.5\%$) or intermediate term (3 RCTs, $n=193$, difference -1.24 , 95% CI -2.22 to 0.12 , $I^2=6.5\%$); see Appendix G, Figure G-7 for updated plot. Based on the addition of the new good-quality trial, the SOE for short- and intermediate-followup for both *function and pain* was upgraded from insufficient to low. No trial reported long-term outcomes. Adverse events were not reported in the new trial.

Interferential Current Compared With Sham

There were no trials of interferential current for OA knee pain in the prior report. One new, good-quality trial ($n=84$)⁶ compared 12, 50-minute sessions of interferential current with sham intervention (sham interferential current and sham photobiomodulation) over 4 weeks and found no differences between groups on any measure of function or pain at short- and intermediate-term followup (SOE: low). *Function* was measured using the WOMAC total score (0 to 96 scale) (short term: difference 0.55 , 95 % CI -24.31 to 7.05 ; intermediate term: difference 1.42 , 95 % CI -6.73 to 9.58) and the Lequesne functional Index (0 to 24 scale) (short term: difference -1.10 , 95 % CI -3.11 to 0.89 ; intermediate term: difference -0.16 , 95 % CI -2.15 to 1.81). *Pain* was measured on the NRS (0 to 10 scale) both at rest (short term: difference -0.87 , 95 % CI -2.01 to 0.26 ; intermediate term: difference -0.32 , 95 % CI -1.34 to 0.70) and during activity (short term: difference -0.42 , 95 % CI -1.65 to 0.80 ; intermediate term: difference 0.49 , 95 % CI -1.63 to 0.64). Harms were not reported.

Mind-Body Therapies for Osteoarthritis Knee Pain

Tai Chi Compared With Attention Control

One new, poor-quality trial ($n=92$)¹² compared Tai Chi (72 sessions total) versus an attention control (health education lectures and discussion) over a treatment period of 6 months. Tai Chi was associated with moderate improvements in both *function* (difference -17.47 on a 0 to 68 WOMAC physical function scale, 95% CI -22.10 to -12.84) and *pain* (difference -1.07 on a 0 to 10 scale, 95% CI -1.74 to -0.40) at intermediate-term followup. The addition of the new trial does not change the prior report's conclusion; the evidence remained insufficient to draw conclusions. Harms were not reported.

Qigong Compared With Exercise

There were no trials of Qigong compared with exercise for OA knee pain in the prior report. One new, fair-quality trial (n=68)²⁹ compared Qigong with combination exercise (strength and aerobic training) in older participants (mean age 70 years); both groups received treatment 4 days a week for 6 months (96 sessions total). At intermediate-term followup, there were no between group differences on measures of *function* (difference -2.20 on the 0 to 68 WOMAC physical function scale, 95% CI -6.24 to 1.84) or *pain* (difference -1.20 on the 0 to 20 WOMAC pain scale, 95% CI -3.04 to 0.64). The strength of evidence was considered insufficient to draw conclusions. Attrition in this study was over 30 percent. The authors also note that adherence to Qigong was high while adherence in the exercise group was very low, possibly due to differences in the comfort level of older patients for performing the exercise types. This may have impacted the findings. Harms were not reported.

Acupuncture for Osteoarthritis Knee Pain

Acupuncture Compared With Usual Care, Waitlist or Sham

One new, good-quality trial (n=83)¹⁶ compared superficial needling acupuncture with sham acupuncture (10 sessions total over 4 weeks) and found no differences between groups in *function* (difference -45.3 on a 0 to 1700 WOMAC physical function scale, 95% CI -183.8 to 93.3) or *pain* (difference -0.22 on a 0 to 10 scale, 95% CI -1.31 to 0.88) at short-term followup. No differences were seen in the proportion of patients with $\geq 30\%$ (59% vs. 56%) or $\geq 50\%$ (39% vs. 36%) improvement on VAS pain. Similar results were seen across other measures of function and pain. These results are consistent with the prior report; the addition of the new trial did not change the prior report's conclusion (no effect for function and pain short term) and SOE remained low.

Adverse events occurred with greater frequency in the true acupuncture (4.4%) versus the sham acupuncture (0.8%) group; RR 5.86 (95% CI 1.87 to 18.39), due primarily to pain during needle insertion or needle adjustment. Pain during needle removal, bleeding, bruising and numbness were also reported by patients randomized to true acupuncture (range of frequencies, 0.2% to 0.7%).

Key Question 4: Fibromyalgia

Exercise for Fibromyalgia

Exercise Compared With Usual Care, Waitlist, an Attention Control or No Treatment

Two new small, poor-quality trials compared exercise therapies with usual care for fibromyalgia and reported only short-term *function*.^{11,23} One trial (n=69)¹¹ evaluated core stability-based physiotherapy and found no difference between groups on the Fibromyalgia Impact Questionnaire (FIQ) physical function item (difference -0.15 on a 0 to 10 scale, 95% CI -1.05 to 0.75). The second trial (n=66)²³ reported that combination exercise (aerobics, balance, postural, stretching and Pilates) was associated with improved function compared with usual care but that the difference did not reach statistical significance per the authors (mean [standard deviation]: 52.8 [17.9] vs. 68.9 [17.5] on the 0-100 FIQ). The addition of the two new poor-quality trials did not change the prior report's conclusion (small effect for function short term) and SOE remained low.

Only one of the trials reported harms. One patient in the exercise group experienced knee pain exacerbation and had to rest during the last three sessions; no other events were reported.¹¹

Manual Therapies for Fibromyalgia

Spinal Manipulation Compared With Sham Manipulation

There were no trials of spinal manipulation for fibromyalgia in the prior report. One new, fair-quality trial (n=101)¹⁰ comparing six sessions (over 6 weeks) of spinal manipulation versus sham manipulation reported no differences between groups in *function* on the 0 to 100 FIQ at short term (adjusted difference 1.2, 95% CI -4.9 to 7.3) and intermediate term (adjusted difference -1.1, 95% CI -7.9 to 5.6) or in *pain* on a 0 to 10 VAS scale at short term (adjusted difference -0.56, 95% CI -2.21 to 1.08) and intermediate term (adjusted difference -0.50, 95% CI -2.48 to 1.47). Similarly, there were no differences between groups in estimated odds ratios (OR) for global improvement in pain at either timepoint (short term: OR 1.44, 95% CI 0.60 to 3.43; intermediate term: OR 1.51, 95% CI 0.65 to 3.51). Attrition in this trial was high (48% at both timepoints). The strength of evidence was considered low.

Mindfulness Practices for Fibromyalgia

Mindfulness-Based Stress Reduction or Meditation Awareness Training Compared With Usual Care, Waitlist or Attention Control

One new, poor-quality trial (n=98)²⁵ found that mindfulness-based stress reduction (MBSR) (8 weekly sessions) was associated with a small improvement in *function* versus usual care over intermediate-term followup (difference -9.77 on the 0-100 FIQ, 95% CI -17.99 to -1.55). The addition of the new trial does not change the prior report's conclusion (small effect for function intermediate term) and SOE remained low.

Eight (16%) patients randomized to MBSR reported adverse events that occurred during and/or after the intervention. These events occurred at a very low intensity and frequency in five patients (no other information provided) and with "significant frequency" in three patients (mild fatigue, intense palpitations, and a variety of moderate adverse effects such as fatigue, tension, headaches, dizziness, somnolence, gain of weight, and loss of sexual desire).

Mind-Body Therapies for Fibromyalgia

Basic Body Awareness Therapy Compared With Usual Care

There were no trials of basic body awareness therapy (BBAT) for fibromyalgia in the prior report. One new small, fair-quality trial (n=39)⁹ found no difference in *pain* (on a 0 to 10 scale) at short term (mean 6.9 vs. 7.4, p>0.05) and intermediate term (mean 7.1 vs. 7.5, p>0.05) between BBAT (10 sessions over 5 weeks) and usual care. Means were estimated from author figures; no information on variability (e.g., standard deviation) was provided precluding estimation of effect size with corresponding confidence intervals. The data were judged to be insufficient to draw conclusions. Harms were not reported.

Acupuncture for Fibromyalgia

Acupuncture Compared With Usual Care and With Exercise

There were no trials of acupuncture compared with usual care (there were 3 trials versus sham acupuncture) or with exercise in the prior report. One new small, poor-quality trial (n=69)¹¹ compared 10 sessions (over 5 weeks) of Traditional Chinese Medicine needle acupuncture versus usual care and versus core stability exercise and found no differences between groups in short-term *function* according to the FIQ physical function item (versus usual care: difference -0.62 on a 0 to 10 scale, 95% CI -1.62 to 0.38; versus exercise: difference -0.47 on a 0 to 10 scale, 95% CI -1.44 to 0.50). The evidence was considered insufficient to draw conclusions. Intermediate- and long-term data and *pain* outcomes were not reported. No harms were reported in either group.

Multidisciplinary Rehabilitation for Fibromyalgia

Multidisciplinary Rehabilitation Compared With Usual Care or Waitlist and with Exercise

One new, poor-quality trial (n=66)²³ compared 12 sessions (over 12 weeks) of multidisciplinary rehabilitation (MDR) (i.e., CBT and occupational therapy) versus usual care and versus combination exercise (aerobics, balance, postural, stretching and Pilates) and reported only short-term *function*. There was no difference between MDR and usual care (difference -0.30 on the 0 to 100 FIQ, 95% CI -8.61 to 8.00) while MDR was associated with less improvement in function compared with exercise, but the difference did not reach statistical significance per the authors (mean [standard deviation]: 68.6 [15.7] vs. 52.8 [17.9]). The addition of the new trial does not change the prior report's conclusion for MDR versus usual care or waitlist (small effect for function short term, SOE remained low). There was no evidence for function at short-term followup for MDR versus exercise in the prior report; the evidence from this new trial was considered insufficient to draw conclusions.

Conclusions

The original report evaluated noninvasive nonpharmacological treatments for five common chronic pain conditions (LBP, neck pain, OA, FM, and tension headache). Across trials in the prior report, exercise, MDR, acupuncture, CBT and mindfulness, and mind-body practices were most consistently associated with durable small to moderate improvements in function and pain for specific chronic pain conditions, although the data were sparse for many interventions. There was very little evidence for chronic tension headache specifically. The majority of trials compared interventions with usual care with very few trials employing pharmacological treatments or exercise as comparators. Harms were poorly reported across interventions. No serious intervention-related adverse events (e.g., leading to death, disability or requiring intensive medical attention) were identified; reported adverse events were generally minor and time-limited.

New evidence for noninvasive nonpharmacological treatments for chronic LBP, chronic neck pain, knee OA and FM for this surveillance report was generally consistent with the prior report and did not alter its conclusions. For chronic LBP, new trials for psychological interventions versus usual care resulted in downgrading of SOE from moderate to low for short-term function. For knee OA, a new trial of low-level laser therapy with sham allowed us to pool data with

previously included trials, upgrading SOE from insufficient to low for small improvements in function but no effect on pain at short or intermediate term. New interventions or comparators for which there was at least low strength of evidence of no effect for function and/or pain were: TENS versus sham TENS in chronic LBP, spinal manipulation versus exercise for chronic neck pain, inferential current versus sham for knee OA, and spinal manipulation versus sham manipulation for FM. Spinal manipulation was associated with improvements in function and pain short term compared with sham laser for chronic neck pain. SOE for all these findings was low. Consistent with the original report, no trials in pregnant or breastfeeding women with pre-existing chronic pain or trials comparing interventions with opioids, topical agents, medical cannabis or muscle relaxants were identified for this update and no new data were available to evaluate nociplasticity as a modifier to treatment effectiveness or safety.

The next surveillance report is scheduled for February 2022.

DRAFT FINAL

References

1. Skelly AC, Chou R, Dettori JR, et al. Noninvasive Nonpharmacological Treatment for Chronic Pain: A Systematic Review Update. Comparative Effectiveness Review No. 227. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. 290-2015-00009-I.) AHRQ Publication No. 20-EHC009. Rockville, MD: Agency for Healthcare Research and Quality; April 2020. <https://effectivehealthcare.ahrq.gov/products/noninvasive-nonpharm-pain-update/research>. PMID: 32338846.
2. Chou R, Dana T, Shetty KD. Testing a Machine Learning Tool for Facilitating Living Systematic Reviews of Chronic Pain Treatments. Methods Research Report. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. 290-2015-00009-I and the Southern California Evidence-based Practice Center-RAND Corporation under Contract No. 290-2015-00010-I.) AHRQ Publication No. 21-EHC004. Rockville, MD: Agency for Healthcare Research and Quality; November 2020. Posted final reports are located on the Effective Health Care Program search page. DOI: 10.23970/AHRQEPCEMETHTESTINGMAC HINELEARNING.
3. Higgins JPT, Green S, eds. Cochrane Handbook for Systematic Reviews of Interventions. Version 5.1.0 [updated March 2011]. The Cochrane Collaboration. Available from <http://handbook.cochrane.org>; 2011.
4. Methods Guide for Effectiveness and Comparative Effectiveness Reviews. AHRQ Publication No. 10(14)-EHC063-EF. Rockville, MD: Agency for Healthcare Research and Quality. January 2014. Chapters available at: www.effectivehealthcare.ahrq.gov.
5. Furlan AD, Malmivaara A, Chou R, et al. 2015 Updated Method Guideline for Systematic Reviews in the Cochrane Back and Neck Group. Spine (Phila Pa 1976). 2015 Nov;40(21):1660-73. doi: 10.1097/BRS.0000000000001061. PMID: 26208232.
6. Alqualo-Costa R, Rampazo EP, Thome GR, et al. Interferential current and photobiomodulation in knee osteoarthritis: A randomized, placebo-controlled, double-blind clinical trial. Clinical Rehabilitation. 2021 Apr 26;2692155211012004. doi: <https://dx.doi.org/10.1177/02692155211012004>. PMID: 33896234.
7. Ashar YK, Gordon A, Schubiner H, et al. Effect of Pain Reprocessing Therapy vs Placebo and Usual Care for Patients With Chronic Back Pain: A Randomized Clinical Trial. JAMA Psychiatry. 2021 Sep 29;doi: 10.1001/jamapsychiatry.2021.2669. PMID: 34586357.
8. Bernal-Utrera C, Gonzalez-Gerez JJ, Anarte-Lazo E, et al. Manual therapy versus therapeutic exercise in non-specific chronic neck pain: a randomized controlled trial. Trials [Electronic Resource]. 2020 Jul 28;21(1):682. doi: <https://dx.doi.org/10.1186/s13063-020-04610-w>. PMID: 32723399.
9. Bravo C, Skjaerven LH, Espart A, et al. Basic Body Awareness Therapy in patients suffering from fibromyalgia: a randomized clinical trial. Physiotherapy theory and practice. 2019;35(10):919-29. PMID: CN-01980930.
10. Coste J, Medkour T, Maigne JY, et al. Osteopathic medicine for fibromyalgia: a sham-controlled randomized clinical trial. Therapeutic Advances in Musculoskeletal Disease. 2021;13:1759720X211009017. doi: <https://dx.doi.org/10.1177/1759720X211009017>. PMID: 33948127.
11. Garrido-Ardila EM, Gonzalez-Lopez-Arza MV, Jimenez-Palomares M, et al. Effectiveness of acupuncture vs. core stability training in balance and functional capacity of women with fibromyalgia: a randomized controlled trial. Clinical Rehabilitation. 2020 May;34(5):630-45. doi: <https://dx.doi.org/10.1177/0269215520911992>. PMID: 32204612.

12. Hu X, Lai Z, Wang L. Effects of Taichi exercise on knee and ankle proprioception among individuals with knee osteoarthritis. *Research in Sports Medicine*. 2020 Apr-Jun;28(2):268-78. doi: <https://dx.doi.org/10.1080/15438627.2019.1663520>. PMID: 31524502.
13. Karakas A, Dilek B, Sahin MA, et al. The effectiveness of pulsed ultrasound treatment on pain, function, synovial sac thickness and femoral cartilage thickness in patients with knee osteoarthritis: a randomized, double-blind clinical, controlled study. *Clinical Rehabilitation*. 2020 Dec;34(12):1474-84. doi: <https://dx.doi.org/10.1177/0269215520942953>. PMID: 32715744.
14. Kholoosy L, Elyaspour D, Akhgari MR, et al. Evaluation of the Therapeutic Effect of Low Level Laser in Controlling Low Back Pain: A Randomized Controlled Trial. *Journal of Lasers in Medical Sciences*. 2020;11(2):120-5. doi: <https://dx.doi.org/10.34172/jlms.2020.21>. PMID: 32273951.
15. Kobayashi D, Shimbo T, Hayashi H, et al. Shiatsu for chronic lower back pain: Randomized controlled study. *Complementary Therapies in Medicine*. 2019 Aug;45:33-7. doi: <https://dx.doi.org/10.1016/j.ctim.2019.05.019>. PMID: 31331579.
16. Lam WC, Au KY, Qin Z, et al. Superficial needling acupuncture versus sham acupuncture for knee osteoarthritis: a randomized controlled trial. *American Journal of Medicine*. 2021 Jun 11;11:11. doi: <https://dx.doi.org/10.1016/j.amjmed.2021.05.002>. PMID: 34126097.
17. Lang AE, Hendrick PA, Clay L, et al. A randomized controlled trial investigating effects of an individualized pedometer driven walking program on chronic low back pain. *BMC Musculoskeletal Disorders*. 2021 Feb 19;22(1):206. doi: <https://dx.doi.org/10.1186/s12891-021-04060-8>. PMID: 33607979.
18. Luo Y, Yang M, Liu T, et al. Effect of hand-ear acupuncture on chronic low-back pain: a randomized controlled trial. *Journal of traditional chinese medicine = chung i tsa chih ying wen pan*. 2019;39(4):587-98. PMID: CN-02144857.
19. Messier SP, Mihalko SL, Beavers DP, et al. Effect of High-Intensity Strength Training on Knee Pain and Knee Joint Compressive Forces Among Adults With Knee Osteoarthritis: The START Randomized Clinical Trial. *JAMA*. 2021 02 16;325(7):646-57. doi: <https://dx.doi.org/10.1001/jama.2021.0411>. PMID: 33591346.
20. Michalsen A, Jeitler M, Kessler CS, et al. Yoga, Eurythmy Therapy and Standard Physiotherapy (YES-Trial) for Patients With Chronic Non-specific Low Back Pain: A Three-Armed Randomized Controlled Trial. *Journal of Pain*. 2021 Apr 20;20:20. doi: <https://dx.doi.org/10.1016/j.jpain.2021.03.154>. PMID: 33892154.
21. Munukka M, Waller B, Hakkinen A, et al. Effects of progressive aquatic resistance training on symptoms and quality of life in women with knee osteoarthritis: A secondary analysis. *Scandinavian Journal of Medicine & Science in Sports*. 2020 Jun;30(6):1064-72. doi: <https://dx.doi.org/10.1111/sms.13630>. PMID: 31999876.
22. Neyaz O, Sumila L, Nanda S, et al. Effectiveness of Hatha Yoga Versus Conventional Therapeutic Exercises for Chronic Nonspecific Low-Back Pain. *Journal of Alternative & Complementary Medicine*. 2019 Sep;25(9):938-45. doi: <https://dx.doi.org/10.1089/acm.2019.0140>. PMID: 31347920.
23. Patru S, Padureanu R, Dumitrescu F, et al. Influence of multidisciplinary therapeutic approach on fibromyalgia patients. *Experimental & Therapeutic Medicine*. 2021 May;21(5):528. doi: <https://dx.doi.org/10.3892/etm.2021.9960>. PMID: 33815601.
24. Pehlivan S, Karadakovan A. Effects of aromatherapy massage on pain, functional state, and quality of life in an elderly individual with knee osteoarthritis. *Japan journal of nursing science*. 2019;16(4):450-8. PMID: CN-02078086 NEW.

25. Perez-Aranda A, Feliu-Soler A, Montero-Marin J, et al. A randomized controlled efficacy trial of mindfulness-based stress reduction compared with an active control group and usual care for fibromyalgia: the EUDAIMON study. *Pain*. 2019 11;160(11):2508-23. doi: <https://dx.doi.org/10.1097/j.pain.0000000000001655>. PMID: 31356450.
26. Rewald S, Lenssen AFT, Emans PJ, et al. Aquatic Cycling Improves Knee Pain and Physical Functioning in Patients With Knee Osteoarthritis: A Randomized Controlled Trial. *Archives of Physical Medicine & Rehabilitation*. 2020 08;101(8):1288-95. doi: <https://dx.doi.org/10.1016/j.apmr.2019.12.023>. PMID: 32169459.
27. Shariat A, Alizadeh R, Moradi V, et al. The impact of modified exercise and relaxation therapy on chronic lower back pain in office workers: a randomized clinical trial. *Journal of Exercise Rehabilitation*. 2019 Oct;15(5):703-8. doi: <https://dx.doi.org/10.12965/jer.1938490.245>. PMID: 31723560.
28. Thomas JS, Clark BC, Russ DW, et al. Effect of Spinal Manipulative and Mobilization Therapies in Young Adults With Mild to Moderate Chronic Low Back Pain: A Randomized Clinical Trial. *JAMA Network Open*. 2020 08 03;3(8):e2012589. doi: <https://dx.doi.org/10.1001/jamanetworkopen.2020.12589>. PMID: 32756930.
29. Xiao CM, Li JJ, Kang Y, et al. Follow-up of a Wuqinxi exercise at home programme to reduce pain and improve function for knee osteoarthritis in older people: a randomised controlled trial. *Age & Ageing*. 2021 02 26;50(2):570-5. doi: <https://dx.doi.org/10.1093/ageing/afaa179>. PMID: 32931545.
30. Yaksi E, Ketenci A, Baslo MB, et al. Does transcutaneous electrical nerve stimulation affect pain, neuropathic pain, and sympathetic skin responses in the treatment of chronic low back pain? A randomized, placebo-controlled study. *The Korean journal of pain*. 2021 Apr 01;34(2):217-28. doi: <https://dx.doi.org/10.3344/kjp.2021.34.2.217>. PMID: 33785674.
31. Groessl EJ, Liu L, Schmalzl L, et al. Secondary Outcomes from a Randomized Controlled Trial of Yoga for Veterans with Chronic Low-Back Pain. *Int J Yoga Therap*. 2020 Jan 1;30(1):69-76. doi: 10.17761/2020-D-19-00036. PMID: 31509451.
32. Joyce C, Roseen EJ, Keysor JJ, et al. Can Yoga or Physical Therapy for Chronic Low Back Pain Improve Depression and Anxiety Among Adults From a Racially Diverse, Low-Income Community? A Secondary Analysis of a Randomized Controlled Trial. *Archives of Physical Medicine & Rehabilitation*. 2021 Jun;102(6):1049-58. doi: <https://dx.doi.org/10.1016/j.apmr.2021.01.072>. PMID: 33556352.
33. McCrae CS, Curtis AF, Miller MB, et al. Effect of cognitive behavioural therapy on sleep and opioid medication use in adults with fibromyalgia and insomnia. *Journal of Sleep Research*. 2020 12;29(6):e13020. doi: <https://dx.doi.org/10.1111/jsr.13020>. PMID: 32126156.
34. Roseen EJ, Gerlovin H, Femia A, et al. Yoga, Physical Therapy, and Back Pain Education for Sleep Quality in Low-Income Racially Diverse Adults with Chronic Low Back Pain: a Secondary Analysis of a Randomized Controlled Trial. *Journal of General Internal Medicine*. 2020 01;35(1):167-76. doi: <https://dx.doi.org/10.1007/s11606-019-05329-4>. PMID: 31667747.
35. Basford JR, Sheffield CG, Harmsen WS. Laser therapy: a randomized, controlled trial of the effects of low-intensity Nd:YAG laser irradiation on musculoskeletal back pain. *Arch Phys Med Rehabil*. 1999 Jun;80(6):647-52. doi: 10.1016/s0003-9993(99)90167-3. PMID: 10378490.
36. Al Rashoud AS, Abboud RJ, Wang W, et al. Efficacy of low-level laser therapy applied at acupuncture points in knee osteoarthritis: a randomised double-blind comparative trial. *Physiotherapy*. 2014 Sep;100(3):242-8. doi: 10.1016/j.physio.2013.09.007. PMID: 24418801.

37. Hegedus B, Viharos L, Gervain M, et al. The effect of low-level laser in knee osteoarthritis: a double-blind, randomized, placebo-controlled trial. *Photomed Laser Surg.* 2009 Aug;27(4):577-84. doi: 10.1089/pho.2008.2297. PMID: 19530911.
38. Tascioglu F, Armagan O, Tabak Y, et al. Low power laser treatment in patients with knee osteoarthritis. *Swiss Med Wkly.* 2004 May 1;134(17-18):254-8. doi: 2004/17/smw-10518. PMID: 15243853.

DRAFT FINAL

Authors

Andrea C. Skelly, Ph.D., M.P.H.
Erika D. Brodt, B.S.
Shelby Kantner, B.A.
Andrea Diulio-Nakamura, Ph.D.
Kim Mauer, M.D.
Kanaka D. Shetty, M.D., M.S.

Acknowledgments

The authors gratefully acknowledge the following individuals for their contributions to this project: research associate and librarian, Tracy Dana, M.L.S. and research associate Christina Bougatsos M.P.H., all from Oregon Health & Science University; Task Order Officer Suchitra Iyer, Ph.D., at the Agency for Healthcare Research and Quality.

Disclaimers

This report is based on research conducted by the Pacific Northwest Evidence-based Practice Center under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. 75Q80120D00006/Task Order 75Q80121F32008). The findings and conclusions in this document are those of the authors, who are responsible for its contents; the findings and conclusions do not necessarily represent the views of AHRQ. Therefore, no statement in this report should be construed as an official position of AHRQ or of the U.S. Department of Health and Human Services.

None of the investigators have any affiliations or financial involvement that conflicts with the material presented in this report.

The information in this report is intended to help healthcare decision makers—patients and clinicians, health system leaders, and policymakers, among others—make well-informed decisions and thereby improve the quality of health care services. This report is not intended to be a substitute for the application of clinical judgment. Anyone who makes decisions concerning the provision of clinical care should consider this report in the same way as any medical reference and in conjunction with all other pertinent information, i.e., in the context of available resources and circumstances presented by individual patients.

This report is made available to the public under the terms of a licensing agreement between the author and the Agency for Healthcare Research and Quality. This report may be used and reprinted without permission except those copyrighted materials that are clearly noted in the report. Further reproduction of those copyrighted materials is prohibited without the express permission of copyright holders.

AHRQ or U.S. Department of Health and Human Services endorsement of any derivative products that may be developed from this report, such as clinical practice guidelines, other quality enhancement tools, or reimbursement or coverage policies, may not be stated or implied.

AHRQ appreciates appropriate acknowledgment and citation of its work. Suggested language for acknowledgment: This work is a quarterly update report of a living systematic evidence report based on, Opioid Treatments for Chronic Pain, by the Evidence-based Practice Center Program at the Agency for Healthcare Research and Quality (AHRQ).

Suggested citation: Skelly AC, Brodt ED, Kantner S, Diulio-Nakamura A, Mauer K, Shetty KD. Systematic Review on Noninvasive Nonpharmacological Treatment for Chronic Pain: Surveillance Report 1. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. 75Q80120D00006.) AHRQ Publication No. xxx. Rockville, MD: Agency for Healthcare Research and Quality; November 2021. DOI:xxx. Posted final reports are located on the Effective Health Care Program [search page](#).

Afterword

The Agency for Healthcare Research and Quality (AHRQ), through its Evidence-based Practice Centers (EPCs), sponsors the development of systematic reviews to assist public- and private-sector organizations in their efforts to improve the quality of healthcare in the United States. These reviews provide comprehensive, science-based information on common, costly medical conditions, and new healthcare technologies and strategies.

Systematic reviews are the building blocks underlying evidence-based practice; they focus attention on the strength and limits of evidence from research studies about the effectiveness and safety of a clinical intervention. In the context of developing recommendations for practice, systematic reviews can help clarify whether assertions about the value of the intervention are based on strong evidence from clinical studies. For more information about AHRQ EPC systematic reviews, see <https://effectivehealthcare.ahrq.gov/about/epc/evidence-synthesis>.

These quarterly progress reports will provide up-to-date information about the evidence base to inform health plans, providers, purchasers, government programs, and the healthcare system as a whole on the state of the science. Transparency and stakeholder input are essential to the Effective Health Care Program. Please visit the website (www.effectivehealthcare.ahrq.gov) to see draft research questions and reports or to join an email list to learn about new program products and opportunities for input.

If you have comments on this report, they may be sent by mail to the Task Order Officer named below at: Agency for Healthcare Research and Quality, 5600 Fishers Lane, Rockville, MD 20857, or by email to epc@ahrq.hhs.gov. They will be considered in the next version of the report.

David Meyers, M.D.
Acting Director
Agency for Healthcare Research and Quality

Arlene S. Bierman, M.D., M.S.
Director
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Craig A. Umscheid, M.D., M.S.
Director
Evidence-based Practice Center Program
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Suchitra Iyer, Ph.D.
Task Order Officer
Evidence-based Practice Center Program
Center for Evidence and Practice Improvement
Agency for Healthcare Research and Quality

Appendix A. Literature Search Strategies

Database: Ovid MEDLINE(R) and Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Daily and Versions(R), All 2020 through September 30, 2021

1 exp Low Back Pain/ or ((back or spine or spinal) adj2 pain).ti,ab.

2 exp Chronic Pain/

3 Neck Pain/ or neck.ti,ab.

4 exp Osteoarthritis/ or osteoarthritis.ti,ab.

5 Headache/ or headache.ti,ab.

6 Fibromyalgia/ or fibromyalgia.ti,ab.

7 exp Exercise Therapy/

8 exp Physical Therapy Modalities/

9 exp Braces/

10 exp Mind-Body Therapies/

11 exp Acupuncture Therapy/

12 exp Rehabilitation/

13 exp Psychotherapy/

14 exp Musculoskeletal Manipulations/

15 (noninvasive or non-invasive or nonpharmacologic* or non-pharmacologic*).ti,ab.

16 (exercise or physical therapy or cognitive or behavioral or feedback or relaxation or acceptance or commitment or traction or ultrasound or stimulation or laser or magnet* or inferential or electromuscular or diathermy or heat or cold or manipulation or manual or craniosacral or mindfulness or meditation or mind-body or yoga or pilates or Qigong or acupuncture or functional restoration or multidisciplin* or interdisciplin*).ti,ab.

17 rh.fs.

18 or/1-6

19 or/7-17

20 18 and 19

21 randomized controlled trial.pt.

22 controlled clinical trial.pt.

23 clinical trials as topic.sh.

24 (random* or trial or placebo).ti,ab.

25 clinical trials as topic.sh.

26 exp animals/ not humans.sh.

27 or/21-25

28 27 not 26

29 20 and 28

30 limit 29 to english language

31 limit 30 to humans

32 31 and (20171\$ or 2018\$).dt,ed,ep.

33 meta-analysis.pt.

34 meta-analysis/ or systematic review/ or meta-analysis as topic/ or "meta analysis (topic)"/ or "systematic review (topic)"/ or exp technology assessment, biomedical/

35 ((systematic* adj3 (review* or overview*)) or (methodologic* adj3 (review* or overview*))).ti,ab.

36 ((quantitative adj3 (review* or overview* or syntheses*)) or (research adj3 (integrati* or overview*))).ti,ab.
 37 ((integrative adj3 (review* or overview*)) or (collaborative adj3 (review* or overview*)) or (pool* adj3 analy*)).ti,ab.
 38 (data syntheses* or data extraction* or data abstraction*).ti,ab.
 39 (handsearch* or hand search*).ti,ab.
 40 (mantel haenszel or peto or der simonian or dersimonian or fixed effect* or latin square*).ti,ab.
 41 (met analy* or metanaly* or technology assessment* or HTA or HTAs or technology overview* or technology appraisal*).ti,ab.
 42 (meta regression* or metaregression*).ti,ab.
 43 (meta-analy* or metaanaly* or systematic review* or biomedical technology assessment* or bio-medical technology assessment*).mp,hw.
 44 (medline or cochrane or pubmed or medlars or embase or cinahl).ti,ab,hw.
 45 (cochrane or (health adj2 technology assessment) or evidence report).jw.
 46 (meta-analysis or systematic review).ti,ab.
 47 (comparative adj3 (efficacy or effectiveness)).ti,ab.
 48 (outcomes research or relative effectiveness).ti,ab.
 49 ((indirect or indirect treatment or mixed-treatment) adj comparison*).ti,ab.
 50 or/33-49
 51 20 and 50
 52 limit 51 to english language
 53 limit 52 to humans
 54 53 and (20171\$ or 2018\$).dt,ed,ep.
 55 32 or 54

Database: EBM Reviews - Cochrane Central Register of Controlled Trials, All 2020 through September 30, 2021

1 exp Low Back Pain/ or ((back or spine or spinal) adj2 pain).ti,ab.
 2 exp Chronic Pain/
 3 Neck Pain/ or neck.ti,ab.
 4 exp Osteoarthritis/ or osteoarthritis.ti,ab.
 5 Headache/ or headache.ti,ab.
 6 Fibromyalgia/ or fibromyalgia.ti,ab.
 7 exp Exercise Therapy/
 8 exp Physical Therapy Modalities/
 9 exp Braces/
 10 exp Mind-Body Therapies/
 11 exp Acupuncture Therapy/
 12 exp Rehabilitation/
 13 exp Psychotherapy/
 14 exp Musculoskeletal Manipulations/
 15 (noninvasive or non-invasive or nonpharmacologic* or non-pharmacologic*).ti,ab.
 16 (exercise or physical therapy or cognitive or behavioral or feedback or relaxation or acceptance or commitment or traction or ultrasound or stimulation or laser or magnet* or inferential or electromuscular or diathermy or heat or cold or manipulation or manual or

craniosacral or mindfulness or meditation or mind-body or yoga or pilates or Qigong or acupuncture or functional restoration or multidisciplin* or interdisciplin*).ti,ab.

17 rh.fs.

18 or/1-6

19 or/7-17

20 18 and 19

21 limit 20 to yr="2017 -Current"

22 limit 21 to medline records

23 21 not 22

24 limit 23 to english language

Database: EBM Reviews - Cochrane Database of Systematic Reviews, All 2020 through September 30, 2021

1 ((back or spine or spinal) adj2 pain).ti.

2 (neck adj2 pain).ti.

3 osteoarthritis.ti.

4 headache.ti.

5 fibromyalgia.ti.

6 (noninvasive or non-invasive or nonpharmacologic* or non-pharmacologic*).ti,ab. (295)

7 (exercise or physical therapy or cognitive or behavioral or feedback or relaxation or acceptance or commitment or traction or ultrasound or stimulation or laser or magnet* or inferential or electromuscular or diathermy or heat or cold or manipulation or manual or craniosacral or mindfulness or meditation or mind-body or yoga or pilates or Qigong or acupuncture or functional restoration or multidisciplin* or interdisciplin*).ti,ab.

9 6 or 7

10 8 and 9

11 limit 10 to new reviews

Appendix B. Key Questions and Inclusion and Exclusion Criteria

Key Questions

Key Question 1: Adults with chronic low back pain

Key Question 2: Adults with chronic neck pain

Key Question 3: Adults with osteoarthritis-related pain

Key Question 4: Adults with fibromyalgia

Key Question 5: Adults with chronic tension headache

Key Questions 1–5 incorporate the following subquestions:

- a. What are the benefits and harms of noninvasive nonpharmacological therapies compared with sham treatment, no treatment, waitlist, attention control, or usual care?
- b. What are the benefits and harms of noninvasive nonpharmacological therapies compared with pharmacological therapy (e.g., opioids, nonsteroidal anti-inflammatory drugs, acetaminophen, antiseizure medications, antidepressants, topical agents, medical cannabis, and muscle relaxants)?
- c. What are the benefits and harms of noninvasive nonpharmacological therapies compared with exercise or, for headache, biofeedback?

The three-part format for Key Questions 1–5 reflects the following research concepts:

- Part “a” answers the question of whether the various interventions work overall compared with sham, waitlist control, attention control, no treatment, or usual care. For this review, usual care was defined as care that might be provided or recommended by a primary care provider.
- Part “b” answers the question of whether the various interventions work compared with pharmacological alternatives.
- Part “c” answers the question of how outcomes for individual interventions (e.g., acupuncture) compare with a common comparator. Exercise is the most frequent comparison in the literature

for many chronic pain conditions, so it provides a common comparator for analysis. It is also recommended in most guidelines for conditions including low back pain, neck pain, fibromyalgia, and osteoarthritis and is widely available. Exercise served as common comparator for these conditions. For chronic headache, biofeedback provided a common comparator for analysis.

Key Question 6: Do estimates of benefits and harms differ by age, sex, presence of comorbidities (e.g., emotional or mood disorders), or degree of nociplasticity/central sensitization?

Table B-1. Inclusion and exclusion criteria

PICOTS	Inclusion	Exclusion
Population All KQs	General Inclusion Criteria <ul style="list-style-type: none"> Adults with the following chronic pain (defined as pain lasting 12 weeks or longer or pain persisting past the time for normal tissue healing) conditions: low back pain, neck pain, osteoarthritis pain, fibromyalgia, or tension headache. Pregnant or breastfeeding women who have a history of chronic pain prior to pregnancy 	General Exclusion Criteria <ul style="list-style-type: none"> Acute pain Children (<18 years), pregnant or breastfeeding women with pregnancy-related back or pelvic pain or who do not have chronic pain prior to pregnancy; Patients with chronic pain related to “active” cancer, infection, inflammatory arthropathy, <90% of study sample has the defined condition of interest or <90% received the treatment(s) of interest Treatment for addiction Pain at the end of life Neuropathic pain
Population KQ1	KQ1: Low back pain <ul style="list-style-type: none"> Adults with chronic, nonradicular low back pain 	KQ1: Low back pain <ul style="list-style-type: none"> Patients with radiculopathy Low back pain associated with severe or progressive neurological deficits Failed back surgery syndrome
Population KQ2	KQ2: Neck pain <ul style="list-style-type: none"> Adults with chronic neck pain 	KQ2: Neck pain <ul style="list-style-type: none"> Patients with radiculopathy or myelopathy Traumatic spinal cord injury Neck pain associated with progressive neurological deficit, loss of strength
Population KQ3	KQ3: Osteoarthritis <ul style="list-style-type: none"> Adults with osteoarthritis-related pain (primary or secondary osteoarthritis) of the hip, knee or hand 	KQ3: Osteoarthritis <ul style="list-style-type: none"> Other types of arthritis (e.g., rheumatoid) Patients with joint replacement
Population KQ4	KQ4: Fibromyalgia <ul style="list-style-type: none"> Adults with fibromyalgia 	KQ4: Fibromyalgia <ul style="list-style-type: none"> Conditions with generalized pain not consistent with fibromyalgia Systemic exertion intolerance disease, (myalgic encephalomyelitis/chronic fatigue syndrome) Somatization disorder (Briquet’s syndrome)

PICOTS	Inclusion	Exclusion
<p>Population KQ5</p>	<p>KQ5: Headache</p> <ul style="list-style-type: none"> • Adults with primary chronic tension headache (International Classification of Headache Disorders, 3rd edition definition). <ul style="list-style-type: none"> ◦ Primary headaches are attributed to the headache condition itself, not headache caused by another disease or medical condition. Tension headaches are the most common. <p>Chronic headache is defined as 15 or more days each month for at least 12 weeks or history of headache more than 180 days a year.</p>	<p>KQ5: Headache</p> <ul style="list-style-type: none"> • Migraine headache • Mixed headache (also known as coexistent tension and migraine headache, chronic daily headache, transformed migraine) • Trigeminal neuralgia • Cluster headache • Secondary headache types as defined in <i>The International Classification of Headache Disorders</i>, 3rd edition (i.e., headaches due to an underlying pathology such as cancer, prior medical procedures, temporomandibular joint disorders, neck pathology, cervicogenic headache, and medication over-use headache) • Traumatic brain injury
<p>Interventions</p>	<p>All KQs:</p> <ul style="list-style-type: none"> • Exercise (exercise as part of physical therapy, supervised exercise, home exercise, group exercise, formal exercise program) • Psychological therapies (cognitive and/or behavioral therapy, biofeedback, relaxation training) • Physical modalities (traction, ultrasound, transcutaneous electrical nerve stimulation, low-level laser therapy, interferential therapy, electro-muscular stimulation diathermy, superficial heat or cold, bracing for knee, back, neck, hand and magnets) • Manual therapies (musculoskeletal manipulation, massage) • Mindfulness practices (meditation, mindfulness-based stress reduction practices) • Mind-body practices (yoga, tai chi, qigong) • Acupuncture • Multidisciplinary/interdisciplinary rehabilitation^a 	<p>All KQs:</p> <ul style="list-style-type: none"> • Invasive nonsurgical treatments (e.g., injections, nerve block, spinal cord stimulators, parenterally-administered medications) • Surgical interventions (including minimally invasive surgical interventions) • Diet interventions or dietary supplementation • Studies evaluating incremental value of adding a noninvasive nonpharmacological intervention to another noninvasive nonpharmacological intervention • Self-management interventions or programs, self-management education programs • Others not listed for inclusion

PICOTS	Inclusion	Exclusion
<p>Comparators</p>	<p>All KQs, subquestion a</p> <ul style="list-style-type: none"> • Sham treatment • Waitlist • Usual care • No treatment • Attention control intended to control for nonspecific effects (e.g., time, attention, expectations); <p>All KQs subquestion b</p> <ul style="list-style-type: none"> • Commonly used nonopioid pharmacological therapy used to treat chronic pain [NSAIDs, acetaminophen, anti-seizure medications, antidepressants (SNRIs, TCAs), muscle relaxants (including benzodiazepines)] • Topical agents (lidocaine, diclofenac, capsaicin) • Medical cannabis (inhaled, oral, topical); phytocannabinoids (plant derived, THC and CBD); FDA approved synthetic cannabinoids [Dronabinol (THC), Nabilone (similar to THC)] • Opioid analgesics <p>KQs 1-4, 6 subquestion c</p> <ul style="list-style-type: none"> • Exercise^b <p>KQ 5, 6 subquestion c</p> <ul style="list-style-type: none"> • Biofeedback^c 	<p>All KQs:</p> <ul style="list-style-type: none"> • Supplements (e.g., glucosamine, chondroitin, d-ribose, herbal or homeopathic treatments) • Invasive nonsurgical treatments (e.g., injections, nerve block, spinal cord stimulators, parenterally-administered medications) • Antidepressants not typically used for chronic pain including SSRIs and MAOIs • Anti-seizure medications not typically used to treat chronic pain including topiramate, lamotrigine, levetiracetam, phenytoin, valproic acid, zonisamide, tiagabine • Surgical interventions (including minimally invasive surgical interventions) • Studies evaluating incremental value of adding a noninvasive nonpharmacological intervention to another noninvasive nonpharmacological intervention • Comparisons within nonpharmacological intervention types (e.g., comparisons of different types of exercise with each other, different types of massage with each other) • Corticosteroids, biologic drugs • Salicylates (oral and topical) • Topical menthol preparations • Others not listed for inclusion
<p>Outcomes</p>	<p>All KQs:</p> <p>Primary efficacy outcomes; we will focus on outcomes from validated measures for</p> <ul style="list-style-type: none"> • Function/disability/pain interference^d • Pain^d <p>Harms and Adverse effects</p> <p>Secondary outcomes</p> <ul style="list-style-type: none"> • Psychological distress (including measures of depression and anxiety) • Quality of life • Opioid use • Sleep quality, sleep disturbance • Health care utilization 	<p>All KQs:</p> <ul style="list-style-type: none"> • Intermediate outcomes (e.g., biomarkers for inflammation) • Other nonclinical outcomes

PICOTS	Inclusion	Exclusion
Timing	Duration of followup: short term (1 to <6 months), intermediate term (≥6 to <12 months) and long term (≥12 months); focus on longer term (>12 month) effects. Trials lasting ≥6 months that include a supervised intervention followed by continued home treatment as part of the intervention will be included even though the only followup occurs directly after the intervention.	<ul style="list-style-type: none"> • Studies with <1 month followup after treatment
Studies	Randomized controlled trials or high quality systematic reviews of randomized controlled trials published in English; cross-over trials with random assignment of initial treatment will be considered.	All KQs: <ul style="list-style-type: none"> • Studies reporting on intermediate outcomes only • Nonrandomized studies • Abstracts, editorials, letters, conference proceedings • Duplicate publications of the same study that do not report on different outcomes • Single site reports from multicenter trials • White papers • Narrative reviews • Articles identified as preliminary reports when results are published in later versions • Indirect comparisons • Studies with fewer than 15 patients per treatment arm • Systematic reviews on treatment of chronic neck pain, fibromyalgia, chronic headache, or osteoarthritis that are of low methodological quality. Those that do not report outcomes or time frames of interest may be excluded. Systematic reviews may be excluded based on currency or relevance (e.g., if there is a substantial new body of evidence reflected in a later review).
Settings	Any nonhospital setting or in self-directed care	<ul style="list-style-type: none"> • Hospital care, hospice care, emergency department care

CBD = cannabidiol; FDA = Food and Drug Administration; KQ = Key Question; MAOI = monoamine oxidase inhibitor; NSAID = nonsteroidal anti-inflammatory drug; SNRI = serotonin and norepinephrine reuptake inhibitor; SSRI = selective serotonin reuptake inhibitor; TCA = tricyclic antidepressant; THC = tetrahydrocannabinol.

^a Multidisciplinary rehabilitation (MDR) (also known as interdisciplinary rehabilitation), is defined as a coordinated program with biopsychosocial treatment components (e.g., exercise therapy and cognitive-behavioral therapy) provided by professionals from at least two different specialties. Functional restoration training is included as part of MDR.

^b Different forms of exercise will not be compared to each other. Exercise will be compared with nonexercise interventions for low back pain, neck pain, fibromyalgia and osteoarthritis

^c Different forms of biofeedback will not be compared to each other. Biofeedback will be compared with the noninvasive interventions for chronic headache

^d The magnitude of effects for pain and function will be classified using the same system as in the AHRQ-funded noninvasive treatment for low back pain review recognizing that small effects using this system may not meet standard thresholds for clinically meaningful effects. A small effect was defined for pain as a mean between-group difference following treatment of 5 to 10 points on a 0- to 100-point visual analog scale (VAS), 0.5 to 1.0 points on a 0- to 10-point numeric rating scale, or equivalent; for function as a mean difference of 5- to 10-point difference on the 0- to 100-point Oswestry Disability Index (ODI) or 1 to 2 points on the 0- to 24-point Roland-Morris Disability Questionnaire (RDQ), or equivalent; and for any outcome as a standardized mean difference (SMD) of 0.2 to 0.5. A moderate effect was defined for pain as a mean difference of 10 to 20 points on a 0- to 100-point VAS, for function as a mean difference of 10 to 20 points on the ODI or 2 to 5 points on the RDQ, and for any outcome as an SMD of 0.5 to 0.8. Large/substantial effects were defined as greater than moderate. We will apply similar methodology to outcomes measures for the other condition. The clinical relevance of effects classified as small might vary for individual patients depending on preferences, baseline symptom severity, harms, cost, and other factors.

Appendix C. Included Studies List

1. Alqualo-Costa R, Rampazo EP, Thome GR, et al. Interferential current and photobiomodulation in knee osteoarthritis: A randomized, placebo-controlled, double-blind clinical trial. *Clinical Rehabilitation*. 2021 Apr 26;2692155211012004. doi: <https://dx.doi.org/10.1177/02692155211012004>. PMID: 33896234.
2. Ashar YK, Gordon A, Schubiner H, et al. Effect of Pain Reprocessing Therapy vs Placebo and Usual Care for Patients With Chronic Back Pain: A Randomized Clinical Trial. *JAMA Psychiatry*. 2021 Sep 29;doi: 10.1001/jamapsychiatry.2021.2669. PMID: 34586357.
3. Bernal-Utrera C, Gonzalez-Gerez JJ, Anarte-Lazo E, et al. Manual therapy versus therapeutic exercise in non-specific chronic neck pain: a randomized controlled trial. *Trials [Electronic Resource]*. 2020 Jul 28;21(1):682. doi: <https://dx.doi.org/10.1186/s13063-020-04610-w>. PMID: 32723399.
4. Bravo C, Skjaerven LH, Espart A, et al. Basic Body Awareness Therapy in patients suffering from fibromyalgia: a randomized clinical trial. *Physiotherapy theory and practice*. 2019;35(10):919-29. PMID: CN-01980930.
5. Coste J, Medkour T, Maigne JY, et al. Osteopathic medicine for fibromyalgia: a sham-controlled randomized clinical trial. *Therapeutic Advances in Musculoskeletal Disease*. 2021;13:1759720X211009017. doi: <https://dx.doi.org/10.1177/1759720X211009017>. PMID: 33948127.
6. Garrido-Ardila EM, Gonzalez-Lopez-Arza MV, Jimenez-Palomares M, et al. Effectiveness of acupuncture vs. core stability training in balance and functional capacity of women with fibromyalgia: a randomized controlled trial. *Clinical Rehabilitation*. 2020 May;34(5):630-45. doi: <https://dx.doi.org/10.1177/0269215520911992>. PMID: 32204612.
7. Groessl EJ, Liu L, Schmalzl L, et al. Secondary Outcomes from a Randomized Controlled Trial of Yoga for Veterans with Chronic Low-Back Pain. *Int J Yoga Therap*. 2020 Jan 1;30(1):69-76. doi: 10.17761/2020-D-19-00036. PMID: 31509451.
8. Hu X, Lai Z, Wang L. Effects of Taichi exercise on knee and ankle proprioception among individuals with knee osteoarthritis. *Research in Sports Medicine*. 2020 Apr-Jun;28(2):268-78. doi: <https://dx.doi.org/10.1080/15438627.2019.1663520>. PMID: 31524502.
9. Joyce C, Roseen EJ, Keysor JJ, et al. Can Yoga or Physical Therapy for Chronic Low Back Pain Improve Depression and Anxiety Among Adults From a Racially Diverse, Low-Income Community? A Secondary Analysis of a Randomized Controlled Trial. *Archives of Physical Medicine & Rehabilitation*. 2021 Jun;102(6):1049-58. doi: <https://dx.doi.org/10.1016/j.apmr.2021.01.072>. PMID: 33556352.
10. Karakas A, Dilek B, Sahin MA, et al. The effectiveness of pulsed ultrasound treatment on pain, function, synovial sac thickness and femoral cartilage thickness in patients with knee osteoarthritis: a randomized, double-blind clinical, controlled study. *Clinical Rehabilitation*. 2020 Dec;34(12):1474-84. doi: <https://dx.doi.org/10.1177/0269215520942953>. PMID: 32715744.
11. Kholoosy L, Elyaspour D, Akhgari MR, et al. Evaluation of the Therapeutic Effect of Low Level Laser in Controlling Low Back Pain: A Randomized Controlled Trial. *Journal of Lasers in Medical Sciences*. 2020;11(2):120-5. doi: <https://dx.doi.org/10.34172/jlms.2020.21>. PMID: 32273951.
12. Kobayashi D, Shimbo T, Hayashi H, et al. Shiatsu for chronic lower back pain: Randomized controlled study. *Complementary Therapies in Medicine*. 2019 Aug;45:33-7. doi: <https://dx.doi.org/10.1016/j.ctim.2019.05.019>. PMID: 31331579.

13. Lam WC, Au KY, Qin Z, et al. Superficial needling acupuncture versus sham acupuncture for knee osteoarthritis: a randomized controlled trial. *American Journal of Medicine*. 2021 Jun 11;11:11. doi: <https://dx.doi.org/10.1016/j.amjmed.2021.05.002>. PMID: 34126097.
14. Lang AE, Hendrick PA, Clay L, et al. A randomized controlled trial investigating effects of an individualized pedometer driven walking program on chronic low back pain. *BMC Musculoskeletal Disorders*. 2021 Feb 19;22(1):206. doi: <https://dx.doi.org/10.1186/s12891-021-04060-8>. PMID: 33607979.
15. Luo Y, Yang M, Liu T, et al. Effect of hand-ear acupuncture on chronic low-back pain: a randomized controlled trial. *Journal of traditional chinese medicine = chung i tsa chih ying wen pan*. 2019;39(4):587-98. PMID: CN-02144857.
16. McCrae CS, Curtis AF, Miller MB, et al. Effect of cognitive behavioural therapy on sleep and opioid medication use in adults with fibromyalgia and insomnia. *Journal of Sleep Research*. 2020 12;29(6):e13020. doi: <https://dx.doi.org/10.1111/jsr.13020>. PMID: 32126156.
17. Messier SP, Mihalko SL, Beavers DP, et al. Effect of High-Intensity Strength Training on Knee Pain and Knee Joint Compressive Forces Among Adults With Knee Osteoarthritis: The START Randomized Clinical Trial. *JAMA*. 2021 02 16;325(7):646-57. doi: <https://dx.doi.org/10.1001/jama.2021.0411>. PMID: 33591346.
18. Michalsen A, Jeitler M, Kessler CS, et al. Yoga, Eurythmy Therapy and Standard Physiotherapy (YES-Trial) for Patients With Chronic Non-specific Low Back Pain: A Three-Armed Randomized Controlled Trial. *Journal of Pain*. 2021 Apr 20;20:20. doi: <https://dx.doi.org/10.1016/j.jpain.2021.03.154>. PMID: 33892154.
19. Munukka M, Waller B, Hakkinen A, et al. Effects of progressive aquatic resistance training on symptoms and quality of life in women with knee osteoarthritis: A secondary analysis. *Scandinavian Journal of Medicine & Science in Sports*. 2020 Jun;30(6):1064-72. doi: <https://dx.doi.org/10.1111/sms.13630>. PMID: 31999876.
20. Neyaz O, Sumila L, Nanda S, et al. Effectiveness of Hatha Yoga Versus Conventional Therapeutic Exercises for Chronic Nonspecific Low-Back Pain. *Journal of Alternative & Complementary Medicine*. 2019 Sep;25(9):938-45. doi: <https://dx.doi.org/10.1089/acm.2019.0140>. PMID: 31347920.
21. Patru S, Padureanu R, Dumitrescu F, et al. Influence of multidisciplinary therapeutic approach on fibromyalgia patients. *Experimental & Therapeutic Medicine*. 2021 May;21(5):528. doi: <https://dx.doi.org/10.3892/etm.2021.9960>. PMID: 33815601.
22. Pehlivan S, Karadakovan A. Effects of aromatherapy massage on pain, functional state, and quality of life in an elderly individual with knee osteoarthritis. *Japan journal of nursing science*. 2019;16(4):450-8. PMID: CN-02078086 NEW.
23. Perez-Aranda A, Feliu-Soler A, Montero-Marin J, et al. A randomized controlled efficacy trial of mindfulness-based stress reduction compared with an active control group and usual care for fibromyalgia: the EUDAIMON study. *Pain*. 2019 11;160(11):2508-23. doi: <https://dx.doi.org/10.1097/j.pain.0000000000001655>. PMID: 31356450.
24. Rewald S, Lenssen AFT, Emans PJ, et al. Aquatic Cycling Improves Knee Pain and Physical Functioning in Patients With Knee Osteoarthritis: A Randomized Controlled Trial. *Archives of Physical Medicine & Rehabilitation*. 2020 08;101(8):1288-95. doi: <https://dx.doi.org/10.1016/j.apmr.2019.12.023>. PMID: 32169459.

25. Roseen EJ, Gerlovin H, Femia A, et al. Yoga, Physical Therapy, and Back Pain Education for Sleep Quality in Low-Income Racially Diverse Adults with Chronic Low Back Pain: a Secondary Analysis of a Randomized Controlled Trial. *Journal of General Internal Medicine*. 2020 01;35(1):167-76. doi: <https://dx.doi.org/10.1007/s11606-019-05329-4>. PMID: 31667747.
26. Shariat A, Alizadeh R, Moradi V, et al. The impact of modified exercise and relaxation therapy on chronic lower back pain in office workers: a randomized clinical trial. *Journal of Exercise Rehabilitation*. 2019 Oct;15(5):703-8. doi: <https://dx.doi.org/10.12965/jer.1938490.245>. PMID: 31723560.
27. Thomas JS, Clark BC, Russ DW, et al. Effect of Spinal Manipulative and Mobilization Therapies in Young Adults With Mild to Moderate Chronic Low Back Pain: A Randomized Clinical Trial. *JAMA Network Open*. 2020 08 03;3(8):e2012589. doi: <https://dx.doi.org/10.1001/jamanetworkopen.2020.12589>. PMID: 32756930.
28. Xiao CM, Li JJ, Kang Y, et al. Follow-up of a Wuqinxi exercise at home programme to reduce pain and improve function for knee osteoarthritis in older people: a randomised controlled trial. *Age & Ageing*. 2021 02 26;50(2):570-5. doi: <https://dx.doi.org/10.1093/ageing/afaa179>. PMID: 32931545.
29. Yaksi E, Ketenci A, Baslo MB, et al. Does transcutaneous electrical nerve stimulation affect pain, neuropathic pain, and sympathetic skin responses in the treatment of chronic low back pain? A randomized, placebo-controlled study. *The Korean journal of pain*. 2021 Apr 01;34(2):217-28. doi: <https://dx.doi.org/10.3344/kjp.2021.34.2.217>. PMID: 33785674.

Appendix D. Evidence Tables

Shown in associated Excel files at <https://effectivehealthcare.ahrq.gov/products/noninvasive-nonpharm-pain-update/research>.

DRAFT FINAL

Appendix E. Quality Assessment

Shown in associated Excel files at <https://effectivehealthcare.ahrq.gov/products/noninvasive-nonpharm-pain-update/research>.

DRAFT FINAL

Appendix F. Excluded Studies List

1. Abdel-Aal NM, Elsayyad MM, Megahed AA. Short-term effect of adding graston technique to exercise program in treatment of patients with cervicogenic headache: a single-blinded, randomized controlled trial. European journal of physical & rehabilitation medicine. 2021 May 05;05:05. doi: <https://dx.doi.org/10.23736/S1973-9087.21.06595-3>. PMID: 33947825. **Exclusion:** Ineligible population
2. Abdelbasset WK, Nambi G, Alsubaie SF, et al. A Randomized Comparative Study between High-Intensity and Low-Level Laser Therapy in the Treatment of Chronic Nonspecific Low Back Pain. Evidence-Based Complementary & Alternative Medicine: eCAM. 2020;2020:1350281. doi: <https://dx.doi.org/10.1155/2020/1350281>. PMID: 33178306. **Exclusion:** Inadequate duration of followup
3. Ahmadnezhad L, Yalfani A, Gholami Borujeni B. Inspiratory Muscle Training in Rehabilitation of Low Back Pain: A Randomized Controlled Trial. Journal of Sport Rehabilitation. 2020 11 01;29(8):1151-8. doi: <https://dx.doi.org/10.1123/jsr.2019-0231>. PMID: 31910393. **Exclusion:** Inadequate duration of followup
4. Akaltun MS, Altindag O, Turan N, et al. Efficacy of high intensity laser therapy in knee osteoarthritis: a double-blind controlled randomized study. Clinical Rheumatology. 2021 May;40(5):1989-95. doi: <https://dx.doi.org/10.1007/s10067-020-05469-7>. PMID: 33074393. **Exclusion:** Ineligible intervention
5. Alayat MSM, Aly THA, Elsayed AEM, et al. Correction to: Efficacy of pulsed Nd:YAG laser in the treatment of patients with knee osteoarthritis: a randomized controlled trial. Lasers in Medical Science. 2020 Oct;35(8):1875. doi: <https://dx.doi.org/10.1007/s10103-020-03088-x>. PMID: 32647935. **Exclusion:** Not a study
6. Alayat MSM, Atya AM, Ali MME, et al. Correction to: Long-term effect of high-intensity laser therapy in the treatment of patients with chronic low back pain: a randomized blinded placebo-controlled trial. Lasers in Medical Science. 2020 Feb;35(1):297. doi: <https://dx.doi.org/10.1007/s10103-019-02926-x>. PMID: 31788745. **Exclusion:** Not a study
7. Albuquerque NF, Lopes BS. Musculoskeletal applications of infrared thermography on back and neck syndromes: a systematic review. European journal of physical & rehabilitation medicine. 2021 Jun;57(3):386-96. doi: <https://dx.doi.org/10.23736/S1973-9087.20.06287-5>. PMID: 33111511. **Exclusion:** Systematic review used as source document
8. Allen KD, Woolson S, Hoenig HM, et al. Stepped Exercise Program for Patients With Knee Osteoarthritis : A Randomized Controlled Trial. Annals of Internal Medicine. 2021 03;174(3):298-307. doi: <https://dx.doi.org/10.7326/M20-4447>. PMID: 33370174. **Exclusion:** Ineligible intervention
9. Alrwaily M, Schneider M, Sowa G, et al. Stabilization exercises combined with neuromuscular electrical stimulation for patients with chronic low back pain: a randomized controlled trial. Brazilian Journal of Physical Therapy. 2019 Nov - Dec;23(6):506-15. doi: <https://dx.doi.org/10.1016/j.bjpt.2018.10.003>. PMID: 30482602. **Exclusion:** Ineligible intervention
10. Alzayed KA, Alsaadi SM. Efficacy of Pulsed Low-Frequency Magnetic Field Therapy on Patients with Chronic Low Back Pain: A Randomized Double-Blind Placebo-Controlled Trial. Asian Spine Journal. 2020 Feb;14(1):33-42. doi: <https://dx.doi.org/10.31616/asj.2019.0043>. PMID: 31575112. **Exclusion:** Ineligible intervention
11. Amaral DDV, Miyamoto GC, Franco KFM, et al. Examination of a Subgroup of Patients With Chronic Low Back Pain Likely to Benefit More From Pilates-Based Exercises Compared to an Educational Booklet. Journal of Orthopaedic & Sports Physical Therapy. 2020 Apr;50(4):189-97. doi: <https://dx.doi.org/10.2519/jospt.2019.8839>.

- PMID: 31443627. **Exclusion:** Ineligible study design
12. Amaral LKB, Souza MB, Campos MGM, et al. Efficacy of conservative therapy in older people with nonspecific low back pain: A systematic review with meta-analysis and GRADE recommendations. Archives of Gerontology & Geriatrics. 2020 Sep - Oct;90:104177. doi: <https://dx.doi.org/10.1016/j.archger.2020.104177>. PMID: 32682168. **Exclusion:** Systematic review used as source document
 13. An J, Ryu HK, Lyu SJ, et al. Effects of Preoperative Telerehabilitation on Muscle Strength, Range of Motion, and Functional Outcomes in Candidates for Total Knee Arthroplasty: A Single-Blind Randomized Controlled Trial. International Journal of Environmental Research & Public Health [Electronic Resource]. 2021 06 04;18(11):04. doi: <https://dx.doi.org/10.3390/ijerph18116071>. PMID: 34199913. **Exclusion:** Ineligible population
 14. Andersen TE, Ravn SL, Armfield N, et al. Trauma-focused cognitive behavioural therapy and exercise for chronic whiplash with comorbid posttraumatic stress disorder: a randomised controlled trial. Pain. 2021 04 01;162(4):1221-32. doi: <https://dx.doi.org/10.1097/j.pain.00000000000002117>. PMID: 33086286. **Exclusion:** Ineligible comparator
 15. Anheyer D, Klose P, Lauche R, et al. Yoga for Treating Headaches: a Systematic Review and Meta-analysis. Journal of General Internal Medicine. 2020 03;35(3):846-54. doi: <https://dx.doi.org/10.1007/s11606-019-05413-9>. PMID: 31667736. **Exclusion:** Systematic review used as source document
 16. Annaswamy TM, Cunniff KJ, Kroll M, et al. Lumbar Bracing for Chronic Low Back Pain: A Randomized Controlled Trial. American Journal of Physical Medicine & Rehabilitation. 2021 08 01;100(8):742-9. doi: <https://dx.doi.org/10.1097/PHM.00000000000001743>. PMID: 33789322. **Exclusion:** Ineligible intervention
 17. Areeudomwong P, Buttagat V. Proprioceptive neuromuscular facilitation training improves pain-related and balance outcomes in working-age patients with chronic low back pain: a randomized controlled trial. Brazilian Journal of Physical Therapy. 2019 Sep - Oct;23(5):428-36. doi: <https://dx.doi.org/10.1016/j.bjpt.2018.10.005>. PMID: 30361077. **Exclusion:** Inadequate duration of followup
 18. Arguisuelas MD, Lison JF, Domenech-Fernandez J, et al. Effects of myofascial release in erector spinae myoelectric activity and lumbar spine kinematics in non-specific chronic low back pain: randomized controlled trial. Clinical biomechanics. 2019;63:27-33. PMID: CN-01707030. **Exclusion:** Inadequate duration of followup
 19. Atalay SG, Durmus A, Gezginaslan O. The Effect of Acupuncture and Physiotherapy on Patients with Knee Osteoarthritis: A Randomized Controlled Study. Pain Physician. 2021 May;24(3):E269-E78. PMID: 33988943. **Exclusion:** Ineligible comparator
 20. Avendano-Coy J, Comino-Suarez N, Grande-Munoz J, et al. Extracorporeal shockwave therapy improves pain and function in subjects with knee osteoarthritis: A systematic review and meta-analysis of randomized clinical trials. International Journal Of Surgery. 2020 Oct;82:64-75. doi: <https://dx.doi.org/10.1016/j.ijsu.2020.07.055>. PMID: 32798759. **Exclusion:** Systematic review used as source document
 21. Barassi G, Supplizi M, Prosperi L, et al. Dual-wavelength high-power laser therapy and neuromuscular manual therapy in chronic neck pain: a randomized clinical trial. Journal of Biological Regulators & Homeostatic Agents. 2021 Mar-Apr;35(2):767-73. doi: <https://dx.doi.org/10.23812/21-37-L>. PMID: 33902272. **Exclusion:** Ineligible study design
 22. Batistella CE, Bidin F, Giacomelli I, et al. Effects of the Russian current in the treatment of low back pain in women: A randomized clinical trial. Journal of Bodywork & Movement Therapies. 2020 Apr;24(2):118-22. doi: <https://dx.doi.org/10.1016/j.jbmt.2019.10.009>. PMID: 32507136. **Exclusion:** Ineligible study design
 23. Bauer CM, Kankaanpaa MJ, Meichtry A, et al. Efficacy of six months neuromuscular

- exercise on lumbar movement variability - A randomized controlled trial. *Journal of Electromyography & Kinesiology*. 2019 Oct;48:84-93. doi: <https://dx.doi.org/10.1016/j.jelekin.2019.06.008>. PMID: 31252284. **Exclusion:** Ineligible outcomes
24. Baumeister H, Paganini S, Sander LB, et al. Effectiveness of a Guided Internet- and Mobile-Based Intervention for Patients with Chronic Back Pain and Depression (WARD-BP): A Multicenter, Pragmatic Randomized Controlled Trial. *Psychotherapy & Psychosomatics*. 2021;90(4):255-68. doi: <https://dx.doi.org/10.1159/000511881>. PMID: 33321501. **Exclusion:** Ineligible intervention
25. Bellomo TR, Schrepf A, Kruger GH, et al. Pressure Pain Tolerance Predicts the Success of Emotional Awareness and Expression Therapy in Patients With Fibromyalgia. *Clin J Pain*. 2020 Jul;36(7):562-6. doi: 10.1097/AJP.0000000000000829. PMID: 32271184. **Exclusion:** Inadequate duration of followup
26. Beltrame R, Ronconi G, Ferrara PE, et al. Capacitive and resistive electric transfer therapy in rehabilitation: a systematic review. *International Journal of Rehabilitation Research*. 2020 Dec;43(4):291-8. doi: <https://dx.doi.org/10.1097/MRR.000000000000000435>. PMID: 32909988. **Exclusion:** Systematic review used as source document
27. Bendrik R, Kallings LV, Broms K, et al. Physical activity on prescription in patients with hip or knee osteoarthritis: A randomized controlled trial. *Clinical Rehabilitation*. 2021 Apr 11;2692155211008807. doi: <https://dx.doi.org/10.1177/02692155211008807>. PMID: 33843297. **Exclusion:** Ineligible population
28. Bernard S, Gentilcore-Saulnier E, Masse-Alarie H, et al. Is adding pelvic floor muscle training to an exercise intervention more effective at improving pain in patients with non-specific low back pain? A systematic review of randomized controlled trials. *Physiotherapy*. 2021 03;110:15-25. doi: <https://dx.doi.org/10.1016/j.physio.2020.02.005>. PMID: 32349867. **Exclusion:** Systematic review used as source document
29. Bidonde J, Busch AJ, Schachter CL, et al. Mixed exercise training for adults with fibromyalgia. *Cochrane Database of Systematic Reviews*. 2019(5) PMID: 00075320-100000000-11748. **Exclusion:** Systematic review used as source document
30. Bokaeian HR, Esfandiarpour F, Zahednejad S, et al. Effects of an Exercise Therapy Targeting Knee Kinetics on Pain, Function, and Gait Kinetics in Patients With Knee Osteoarthritis: A Randomized Clinical Trial. *Adapted Physical Activity Quarterly*. 2021 Mar 30;38(3):377-95. doi: <https://dx.doi.org/10.1123/apaq.2020-0144>. PMID: 33785660. **Exclusion:** Ineligible comparator
31. Bronfort G, Haas M, Evans RL, et al. Non-invasive physical treatments for chronic/recurrent headache. *Cochrane Database of Systematic Reviews*. 2019(8) PMID: 00075320-100000000-01345. **Exclusion:** Systematic review used as source document
32. Burgess DJ, Evans R, Allen KD, et al. Learning to Apply Mindfulness to Pain (LAMP): Design for a Pragmatic Clinical Trial of Two Mindfulness-Based Interventions for Chronic Pain. *Pain Medicine*. 2020 12 12;21(Suppl 2):S29-S36. doi: <https://dx.doi.org/10.1093/pm/pnaa337>. PMID: 33313730. **Exclusion:** Not a study
33. Buttagat V, Muenpan K, Wiriyasakunphan W, et al. A comparative study of Thai massage and muscle energy technique for chronic neck pain: A single-blinded randomized clinical trial. *J Bodyw Mov Ther*. 2021 Jul;27:647-53. doi: 10.1016/j.jbmt.2021.05.007. PMID: 34391301. **Exclusion:** Inadequate duration of followup
34. Calatayud J, Guzman-Gonzalez B, Andersen LL, et al. Effectiveness of a Group-Based Progressive Strength Training in Primary Care to Improve the Recurrence of Low Back Pain Exacerbations and Function: A Randomised Trial. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2020 11 11;17(22):11. doi: <https://dx.doi.org/10.3390/ijerph17228326>.

- PMID: 33187076. **Exclusion:** Ineligible comparator
35. Callaghan MJ, Palmer E, O'Neill T. Management of patellofemoral joint osteoarthritis using biomechanical device therapy: a systematic review with meta-analysis. *Systematic Reviews*. 2021 06 09;10(1):173. doi: <https://dx.doi.org/10.1186/s13643-021-01708-3>. PMID: 34108025. **Exclusion:** Systematic review used as source document
36. Cejudo J, Garcia-Castillo FJ, Luna P, et al. Using a Mindfulness-Based Intervention to Promote Subjective Well-Being, Trait Emotional Intelligence, Mental Health, and Resilience in Women With Fibromyalgia. *Frontiers in Psychology*. 2019;10:2541. doi: <https://dx.doi.org/10.3389/fpsyg.2019.02541>. PMID: 31798502. **Exclusion:** Ineligible outcomes
37. Chao J, Jing Z, Xuehua B, et al. Effect of Systematic Exercise Rehabilitation on Patients With Knee Osteoarthritis: A Randomized Controlled Trial. *Cartilage*. 2020 Feb 10;1947603520903443. doi: <https://dx.doi.org/10.1177/1947603520903443>. PMID: 32037857. **Exclusion:** Inadequate duration of followup
38. Chen H, Wang Y, Liu C, et al. Benefits of a transtheoretical model-based program on exercise adherence in older adults with knee osteoarthritis: A cluster randomized controlled trial. *Journal of Advanced Nursing*. 2020 Jul;76(7):1765-79. doi: <https://dx.doi.org/10.1111/jan.14363>. PMID: 32202313. **Exclusion:** Ineligible comparator
39. Chen H, Zheng X, Huang H, et al. The effects of a home-based exercise intervention on elderly patients with knee osteoarthritis: a quasi-experimental study. *BMC musculoskeletal disorders*. 2019;20(1):160. PMID: CN-01937404. **Exclusion:** Inadequate duration of followup
40. Chen L, Li M, Fan L, et al. Optimized acupuncture treatment (acupuncture and intradermal needling) for cervical spondylosis-related neck pain: a multicenter randomized controlled trial. *Pain*. 2021 03 01;162(3):728-39. doi: <https://dx.doi.org/10.1097/j.pain.0000000000002071>. PMID: 32947547. **Exclusion:** Ineligible intervention
41. Chen PY, Song CY, Yen HY, et al. Impacts of tai chi exercise on functional fitness in community-dwelling older adults with mild degenerative knee osteoarthritis: a randomized controlled clinical trial. *BMC Geriatr*. 2021 Jul 31;21(1):449. doi: 10.1186/s12877-021-02390-9. PMID: 34332537. **Exclusion:** Inadequate duration of followup
42. Chen Z, Wu J, Wang X, et al. The effects of myofascial release technique for patients with low back pain: A systematic review and meta-analysis. *Complementary Therapies in Medicine*. 2021 Jun;59:102737. doi: <https://dx.doi.org/10.1016/j.ctim.2021.102737>. PMID: 33984499. **Exclusion:** Systematic review used as source document
43. Cheng CA, Chiu YW, Wu D, et al. Effectiveness of Tai Chi on fibromyalgia patients: A meta-analysis of randomized controlled trials. *Complementary Therapies in Medicine*. 2019 Oct;46:1-8. doi: <https://dx.doi.org/10.1016/j.ctim.2019.07.007>. PMID: 31519264. **Exclusion:** Systematic review used as source document
44. Cheung DST, Yeung WF, Suen LK, et al. Self-administered acupressure for knee osteoarthritis in middle-aged and older adults: a pilot randomized controlled trial. *Acupuncture in Medicine*. 2020 04;38(2):75-85. doi: <https://dx.doi.org/10.1177/0964528419883269>. PMID: 31718229. **Exclusion:** Inadequate duration of followup
45. Cho SJ. A Self-Efficacy Reinforcement Stretching Exercise Program for Community-Dwelling Older Women With Osteoarthritis: A Pilot Study. *Rehabilitation Nursing Journal*. 2021 Jan-Feb 01;46(1):11-23. doi: <https://dx.doi.org/10.1097/RNJ.0000000000000290>. PMID: 32932424. **Exclusion:** Ineligible study design
46. Coccetta CA, Sale P, Ferrara PE, et al. Effects of capacitive and resistive electric transfer therapy in patients with knee osteoarthritis: a randomized controlled trial. *International journal of rehabilitation research*. 2019;Internationale Zeitschrift fur Rehabilitationsforschung. *Revue internationale de recherches de readaptation*.

- 42(2):106-11. PMID: CN-01937563. **Exclusion:** Ineligible intervention
47. Corp N, Mansell G, Stynes S, et al. Evidence-based treatment recommendations for neck and low back pain across Europe: A systematic review of guidelines. *European Journal of Pain*. 2021 02;25(2):275-95. doi: <https://dx.doi.org/10.1002/ejp.1679>. PMID: 33064878. **Exclusion:** Systematic review used as source document
48. Corvillo I, Armijo F, Alvarez-Badillo A, et al. Efficacy of aquatic therapy for neck pain: a systematic review. *International Journal of Biometeorology*. 2020 Jun;64(6):915-25. doi: <https://dx.doi.org/10.1007/s00484-019-01738-6>. PMID: 31209599. **Exclusion:** Systematic review used as source document
49. Daneau C, Cantin V, Descarreaux M. Effect of Massage on Clinical and Physiological Variables During Muscle Fatigue Task in Participants With Chronic Low Back Pain: a Crossover Study. *Journal of Manipulative and Physiological Therapeutics*. 2019;42(1):55-65. PMID: CN-02001930. **Exclusion:** Ineligible study design
50. de Campos TF, Pocovi NC, Maher CG, et al. An individualised self-management exercise and education program did not prevent recurrence of low back pain but may reduce care seeking: a randomised trial. *Journal of Physiotherapy*. 2020 Jul;66(3):166-73. doi: <https://dx.doi.org/10.1016/j.jphys.2020.06.006>. PMID: 32709590. **Exclusion:** Ineligible population
51. Deer TR, Esposito MF, McRoberts WP, et al. A Systematic Literature Review of Peripheral Nerve Stimulation Therapies for the Treatment of Pain. *Pain Medicine*. 2020 08 01;21(8):1590-603. doi: <https://dx.doi.org/10.1093/pm/pnaa030>. PMID: 32803220. **Exclusion:** Systematic review used as source document
52. Dharmasri CJ, Griesemer I, Arbeeva L, et al. Acceptability of telephone-based pain coping skills training among African Americans with osteoarthritis enrolled in a randomized controlled trial: a mixed methods analysis. *BMC Musculoskeletal Disorders*. 2020 Aug 14;21(1):545. doi: <https://dx.doi.org/10.1186/s12891-020-03578-7>. PMID: 32795282. **Exclusion:** Ineligible study design
53. Didehdar D, Kamali F, Yoosefinejad AK, et al. The effect of spinal manipulation on brain neurometabolites in chronic nonspecific low back pain patients: a randomized clinical trial. *Irish Journal of Medical Science*. 2020 May;189(2):543-50. doi: <https://dx.doi.org/10.1007/s11845-019-02140-2>. PMID: 31773541. **Exclusion:** Ineligible study design
54. Domingues L, Pimentel-Santos FM, Cruz EB, et al. Is a combined programme of manual therapy and exercise more effective than usual care in patients with non-specific chronic neck pain? A randomized controlled trial. *Clinical Rehabilitation*. 2019 Dec;33(12):1908-18. doi: <https://dx.doi.org/10.1177/0269215519876675>. PMID: 31549519. **Exclusion:** Ineligible intervention
55. Dowsey M, Castle D, Knowles S, et al. The effect of mindfulness training prior to total joint arthroplasty on post-operative pain and physical function: A randomised controlled trial. *Complementary Therapies in Medicine*. 2019 Oct;46:195-201. doi: <https://dx.doi.org/10.1016/j.ctim.2019.08.010>. PMID: 31519279. **Exclusion:** Ineligible population
56. Ehsani F, Hedayati R, Bagheri R, et al. The Effects of Stabilization Exercise on the Thickness of Lateral Abdominal Muscles During Standing Tasks in Women With Chronic Low Back Pain: A Randomized Triple-Blinded Clinical Trial Study. *Journal of Sport Rehabilitation*. 2020 Sep 01;29(7):942-51. doi: <https://dx.doi.org/10.1123/jsr.2019-0058>. PMID: 31821992. **Exclusion:** Ineligible comparator
57. Elshawi AM, Hamada HA, Mosaad D, et al. Effect of pulsed electromagnetic field on nonspecific low back pain patients: a randomized controlled trial. *Brazilian journal of physical therapy*. 2019;23(3):244-9. PMID: CN-01628986. **Exclusion:** Inadequate duration of followup
58. Estevez-Lopez F, Maestre-Cascales C, Russell D, et al. Effectiveness of Exercise on Fatigue and Sleep Quality in Fibromyalgia: A Systematic Review and Meta-analysis of Randomized Trials. *Archives of Physical Medicine & Rehabilitation*. 2021 04;102(4):752-61. doi:

- <https://dx.doi.org/10.1016/j.apmr.2020.06.019>. PMID: 32721388. **Exclusion:** Systematic review used as source document
59. Fan Y, Li Z, Zhang H, et al. Valgus knee bracing may have no long-term effect on pain improvement and functional activity in patients with knee osteoarthritis: a meta-analysis of randomized trials. *Journal of Orthopaedic Surgery*. 2020 Sep 01;15(1):373. doi: <https://dx.doi.org/10.1186/s13018-020-01917-x>. PMID: 32873332. **Exclusion:** Systematic review used as source document
60. Fernandez-Carnero J, Sierra-Silvestre E, Beltran-Alacreu H, et al. Neural Tension Technique Improves Immediate Conditioned Pain Modulation in Patients with Chronic Neck Pain: A Randomized Clinical Trial. *Pain Medicine*. 2019 06 01;20(6):1227-35. doi: <https://dx.doi.org/10.1093/pm/pny115>. PMID: 29945245. **Exclusion:** Inadequate duration of followup
61. Ferreira RM, Torres RT, Duarte JA, et al. Non-Pharmacological and Non-Surgical Interventions for Knee Osteoarthritis: A Systematic Review and Meta-Analysis. *Acta Reumatologica Portuguesa*. 2019 07 29;44(3):173-217. PMID: 31356585. **Exclusion:** Systematic review used as source document
62. Ferro Moura Franco K, Lenoir D, Dos Santos Franco YR, et al. Prescription of exercises for the treatment of chronic pain along the continuum of nociplastic pain: A systematic review with meta-analysis. *European Journal of Pain*. 2021 01;25(1):51-70. doi: <https://dx.doi.org/10.1002/ejp.1666>. PMID: 32976664. **Exclusion:** Systematic review used as source document
63. Fertelli TK, Mollaoglu M, Sahin O. Aquatic Exercise Program for Individuals With Osteoarthritis: pain, Stiffness, Physical Function, Self-Efficacy. *Rehabilitation nursing*. 2019 Sep/Oct;44(5):290-9. doi: 10.1097/rnj.000000000000142. PMID: 29613876. **Exclusion:** Inadequate duration of followup
64. Fisher LR, Alvar BA, Maher SF, et al. Short-term Effects of Thoracic Spine Thrust Manipulation, Exercise, and Education in Individuals With Low Back Pain: A Randomized Controlled Trial. *Journal of Orthopaedic & Sports Physical Therapy*. 2020 Jan;50(1):24-32. doi: <https://dx.doi.org/10.2519/jospt.2020.8928>. PMID: 31810405. **Exclusion:** Inadequate duration of followup
65. Fonseca ACS, Faria PC, Alcantara MA, et al. Effects of aquatic physiotherapy or health education program in women with fibromyalgia: a randomized clinical trial. *Physiotherapy Theory & Practice*. 2021 May;37(5):620-32. doi: <https://dx.doi.org/10.1080/09593985.2019.1639229>. PMID: 31305209. **Exclusion:** Inadequate duration of followup
66. Foo CN, Arumugam M, Lekhraj R, et al. Effectiveness of Health-Led Cognitive Behavioral-Based Group Therapy on Pain, Functional Disability and Psychological Outcomes among Knee Osteoarthritis Patients in Malaysia. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2020 08 26;17(17):26. doi: <https://dx.doi.org/10.3390/ijerph17176179>. PMID: 32858791. **Exclusion:** Ineligible intervention
67. Foster NE, Vertosick EA, Lewith G, et al. Identifying patients with chronic pain who respond to acupuncture: results from an individual patient data meta-analysis. *Acupuncture in Medicine*. 2021 04;39(2):83-90. doi: <https://dx.doi.org/10.1177/0964528420920303>. PMID: 32571096. **Exclusion:** Systematic review used as source document
68. Fraenkel L, Buta E, Suter L, et al. Nonsteroidal Anti-inflammatory Drugs vs Cognitive Behavioral Therapy for Arthritis Pain: A Randomized Withdrawal Trial. *JAMA Internal Medicine*. 2020 09 01;180(9):1194-202. doi: <https://dx.doi.org/10.1001/jamainternmed.2020.2821>. PMID: 32702101. **Exclusion:** Ineligible intervention
69. Francescato Torres S, Brandt de Macedo AC, Dias Antunes M, et al. Effects of electroacupuncture frequencies on chronic low back pain in older adults: triple-blind, 12-months protocol for a randomized controlled trial. *Trials* [Electronic Resource]. 2019 Dec 23;20(1):762. doi: <https://dx.doi.org/10.1186/s13063-019-3813-6>. PMID: 31870456. **Exclusion:** Not a study

70. Frutiger M, Borotkanics R. Systematic Review and Meta-Analysis Suggest Strength Training and Workplace Modifications May Reduce Neck Pain in Office Workers. *Pain Practice*. 2021 01;21(1):100-31. doi: <https://dx.doi.org/10.1111/papr.12940>. PMID: 32657531. **Exclusion:** Systematic review used as source document
71. Furukawa Y. Tasuki for neck pain: An individually-randomized, open-label, waiting-list-controlled trial. *Journal of Occupational Health*. 2020 Jan;62(1):e12097. doi: <https://dx.doi.org/10.1002/1348-9585.12097>. PMID: 31705728. **Exclusion:** Ineligible intervention
72. Galvao-Moreira LV, de Castro LO, Moura ECR, et al. Pool-based exercise for amelioration of pain in adults with fibromyalgia syndrome: A systematic review and meta-analysis. *Modern Rheumatology*. 2021 Jul;31(4):904-11. doi: <https://dx.doi.org/10.1080/14397595.2020.1829339>. PMID: 32990113. **Exclusion:** Systematic review used as source document
73. Garijo IH, Del Barrio SJ, Gomez TM, et al. Effectiveness of non-pharmacological conservative therapies in adults with fibromyalgia: A systematic review of high-quality clinical trials. *Journal of Back & Musculoskeletal Rehabilitation*. 2021 Jun 25;25:25. doi: <https://dx.doi.org/10.3233/BMR-200282>. PMID: 34180405. **Exclusion:** Systematic review used as source document
74. Gati T, Czimer E, Cserhati G, et al. A multicentre randomized controlled follow-up study of the effects of the underwater traction therapy in chronic low back pain. *International Journal of Biometeorology*. 2020 Aug;64(8):1393-400. doi: <https://dx.doi.org/10.1007/s00484-020-01919-8>. PMID: 32361959. **Exclusion:** Ineligible intervention
75. Godley E, Smith MA. Efficacy of acupuncture for chronic low back pain: A systematic review. *Complementary Therapies in Clinical Practice*. 2020 May;39:101146. doi: <https://dx.doi.org/10.1016/j.ctcp.2020.101146>. PMID: 32379678. **Exclusion:** Systematic review used as source document
76. Gohir SA, Eek F, Kelly A, et al. Effectiveness of Internet-Based Exercises Aimed at Treating Knee Osteoarthritis: The iBEAT-OA Randomized Clinical Trial. *JAMA Network Open*. 2021 02 01;4(2):e210012. doi: <https://dx.doi.org/10.1001/jamanetworkopen.2021.0012>. PMID: 33620447. **Exclusion:** Inadequate duration of followup
77. Gomez-de-Regil L, Estrella-Castillo DF. Psychotherapy for Physical Pain in Patients with Fibromyalgia: A Systematic Review. *Pain Research & Management*. 2020;2020:3408052. doi: <https://dx.doi.org/10.1155/2020/3408052>. PMID: 32714478. **Exclusion:** Systematic review used as source document
78. Gonzalez-Rueda V, Hidalgo-Garcia C, Rodriguez-Sanz J, et al. Does Upper Cervical Manual Therapy Provide Additional Benefit in Disability and Mobility over a Physiotherapy Primary Care Program for Chronic Cervicalgia? A Randomized Controlled Trial. *International Journal of Environmental Research & Public Health [Electronic Resource]*. 2020 11 11;17(22):11. doi: <https://dx.doi.org/10.3390/ijerph17228334>. PMID: 33187167. **Exclusion:** Ineligible intervention
79. Gonzalez-Rueda V, Lopez-de-Celis C, Bueno-Gracia E, et al. "Short- and mid-term effects of adding upper cervical manual therapy to a conventional physical therapy program in patients with chronic mechanical neck pain. Randomized controlled clinical trial.". *Clinical Rehabilitation*. 2021 Mar;35(3):378-89. doi: <https://dx.doi.org/10.1177/0269215520965054>. PMID: 33076707. **Exclusion:** Ineligible intervention
80. Gopichandran L, Srivastava AK, Vanamail P, et al. Effectiveness of Progressive Muscle Relaxation and Deep Breathing Exercise on Pain, Disability, and Sleep Among Patients With Chronic Tension-Type Headache: A Randomized Control Trial. *Holistic Nursing Practice*. 2021 May 28;28:28. doi: <https://dx.doi.org/10.1097/HNP.00000000000000460>. PMID: 34054116. **Exclusion:** Inadequate duration of followup
81. Gould HM, Atkinson JH, Chircop-Rollick T, et al. A randomized placebo-controlled trial

- of desipramine, cognitive behavioral therapy, and active placebo therapy for low back pain. *Pain*. 2020 06;161(6):1341-9. doi: <https://dx.doi.org/10.1097/j.pain.0000000000001834>. PMID: 32068667. **Exclusion:** Inadequate duration of followup
82. Groessl EJ, Liu L, Richard EL, et al. Cost-effectiveness of Yoga for Chronic Low Back Pain in Veterans. *Medical Care*. 2020 09;58 Suppl 2 9S:S142-S8. doi: <https://dx.doi.org/10.1097/MLR.0000000000001356>. PMID: 32826784. **Exclusion:** Ineligible outcomes
83. Gueugnon M, Fournel I, Soilly AL, et al. Effectiveness, safety, and cost-utility of a knee brace in medial knee osteoarthritis: the ERGONOMIE randomized controlled trial. *Osteoarthritis & Cartilage*. 2021 04;29(4):491-501. doi: <https://dx.doi.org/10.1016/j.joca.2020.11.009>. PMID: 33524515. **Exclusion:** Ineligible population
84. Guimaraes LS, Costa L, Araujo AC, et al. Photobiomodulation therapy is not better than placebo in patients with chronic nonspecific low back pain: a randomised placebo-controlled trial. *Pain*. 2021 06 01;162(6):1612-20. doi: <https://dx.doi.org/10.1097/j.pain.0000000000002189>. PMID: 33449509. **Exclusion:** Ineligible intervention
85. Guinot M, Maindet C, Hodaj H, et al. Effects of Repetitive Transcranial Magnetic Stimulation and Multicomponent Therapy in Patients With Fibromyalgia: A Randomized Controlled Trial. *Arthritis care & research*. 2021 03;73(3):449-58. doi: <https://dx.doi.org/10.1002/acr.24118>. PMID: 31785190. **Exclusion:** Ineligible intervention
86. Gutierrez Espinoza H, Araya-Quintanilla F, Olguin-Huerta C, et al. Effectiveness of manual therapy in patients with thumb carpometacarpal osteoarthritis: a systematic review and meta-analysis. *Physiotherapy Theory & Practice*. 2021 Jun 01:1-10. doi: <https://dx.doi.org/10.1080/09593985.2021.1926026>. PMID: 34074220. **Exclusion:** Systematic review used as source document
87. Hanel J, Owen PJ, Held S, et al. Effects of Exercise Training on Fear-Avoidance in Pain and Pain-Free Populations: Systematic Review and Meta-analysis. *Sports Medicine*. 2020 Dec;50(12):2193-207. doi: <https://dx.doi.org/10.1007/s40279-020-01345-1>. PMID: 32946074. **Exclusion:** Systematic review used as source document
88. Hansen S, Mikkelsen LR, Overgaard S, et al. Effectiveness of supervised resistance training for patients with hip osteoarthritis - a systematic review. *Danish Medical Journal*. 2020 Jun 01;67(6):01. PMID: 32741435. **Exclusion:** Systematic review used as source document
89. Harada K, Takahashi K, Ikuta F, et al. Efficacy of a deep thermal therapy system for osteoarthritis of the knee. *Journal of Nippon Medical School = Nihon Ika Daigaku Zasshi*. 2020 Sep 30;30:30. doi: <https://dx.doi.org/10.1272/jnms.JNMS.202188-505>. PMID: 32999179. **Exclusion:** Ineligible study design
90. Haugmark T, Hagen KB, Provan SA, et al. Effects of a mindfulness-based and acceptance-based group programme followed by physical activity for patients with fibromyalgia: a randomised controlled trial. *BMJ Open*. 2021 Jun 29;11(6):e046943. doi: 10.1136/bmjopen-2020-046943. PMID: 34187823. **Exclusion:** Ineligible intervention
91. Haugmark T, Hagen KB, Smedslund G, et al. Mindfulness- and acceptance-based interventions for patients with fibromyalgia - A systematic review and meta-analyses. *PLoS ONE [Electronic Resource]*. 2019;14(9):e0221897. doi: <https://dx.doi.org/10.1371/journal.pone.0221897>. PMID: 31479478. **Exclusion:** Systematic review used as source document
92. Hayden JA, Wilson MN, Stewart S, et al. Exercise treatment effect modifiers in persistent low back pain: an individual participant data meta-analysis of 3514 participants from 27 randomised controlled trials. *British Journal of Sports Medicine*. 2020 Nov;54(21):1277-8. doi: <https://dx.doi.org/10.1136/bjsports-2019-101205>. PMID: 31780447. **Exclusion:** Systematic review used as source document
93. Hee SW, Mistry D, Friede T, et al. Identification of subgroup effect with an individual participant data meta-analysis of randomised controlled trials of three different types of therapist-delivered care in low back pain. *BMC Musculoskeletal*

- Disorders. 2021 Feb 16;22(1):191. doi: <https://dx.doi.org/10.1186/s12891-021-04028-8>. PMID: 33593341. **Exclusion:** Systematic review used as source document
94. Hernandez D, Dimaro M, Navarro E, et al. Efficacy of core exercises in patients with osteoarthritis of the knee: A randomized controlled clinical trial. *Journal of Bodywork & Movement Therapies*. 2019 Oct;23(4):881-7. doi: <https://dx.doi.org/10.1016/j.jbmt.2019.06.002>. PMID: 31733777. **Exclusion:** Ineligible comparator
95. Hernando-Garijo I, Ceballos-Laita L, Mingo-Gomez MT, et al. Immediate Effects of a Telerehabilitation Program Based on Aerobic Exercise in Women with Fibromyalgia. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2021 02 20;18(4):20. doi: <https://dx.doi.org/10.3390/ijerph18042075>. PMID: 33672691. **Exclusion:** Inadequate duration of followup
96. Howarth A, Riaz M, Perkins-Porras L, et al. Pilot randomised controlled trial of a brief mindfulness-based intervention for those with persistent pain. *Journal of behavioral medicine*. 2019;42(6):999-1014. PMID: CN-01939048. **Exclusion:** Ineligible population
97. Hu L, Wang Y, Liu X, et al. Tai Chi exercise can ameliorate physical and mental health of patients with knee osteoarthritis: systematic review and meta-analysis. *Clinical Rehabilitation*. 2021 Jan;35(1):64-79. doi: <https://dx.doi.org/10.1177/0269215520954343>. PMID: 32954819. **Exclusion:** Systematic review used as source document
98. Huang CC, Wang HH, Chen KC, et al. Effects of a dynamic combined training on impulse response for middle-aged and elderly patients with osteoporosis and knee osteoarthritis: a randomized control trial. *Aging-Clinical & Experimental Research*. 2021 Jan;33(1):115-23. doi: <https://dx.doi.org/10.1007/s40520-020-01508-0>. PMID: 32100224. **Exclusion:** Ineligible study design
99. Huang JF, Zheng XQ, Chen D, et al. Can Acupuncture Improve Chronic Spinal Pain? A Systematic Review and Meta-Analysis. *Global Spine Journal*. 2020 Oct 09;2192568220962440. doi: <https://dx.doi.org/10.1177/2192568220962440>. PMID: 33034233. **Exclusion:** Systematic review used as source document
100. Huang Z, Liu S, Zhou J, et al. Efficacy and Safety of Acupuncture for Chronic Discogenic Sciatica, a Randomized Controlled Sham Acupuncture Trial. *Pain Medicine*. 2019 11 01;20(11):2303-10. doi: <https://dx.doi.org/10.1093/pm/pnz167>. PMID: 31369674. **Exclusion:** Ineligible population
101. Huppe A, Zeuner C, Karstens S, et al. Feasibility and long-term efficacy of a proactive health program in the treatment of chronic back pain: a randomized controlled trial. *BMC Health Services Research*. 2019 Oct 21;19(1):714. doi: <https://dx.doi.org/10.1186/s12913-019-4561-8>. PMID: 31639016. **Exclusion:** Ineligible intervention
102. Iijima H, Eguchi R, Shimoura K, et al. Transcutaneous Electrical Nerve Stimulation Improves Stair Climbing Capacity in People with Knee Osteoarthritis. *Scientific Reports*. 2020 04 29;10(1):7294. doi: <https://dx.doi.org/10.1038/s41598-020-64176-0>. PMID: 32350320. **Exclusion:** Inadequate duration of followup
103. Izquierdo-Alventosa R, Ingles M, Cortes-Amador S, et al. Low-Intensity Physical Exercise Improves Pain Catastrophizing and Other Psychological and Physical Aspects in Women with Fibromyalgia: A Randomized Controlled Trial. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2020 05 21;17(10):21. doi: <https://dx.doi.org/10.3390/ijerph17103634>. PMID: 32455853. **Exclusion:** Inadequate duration of followup
104. Jafarzadeh A, Ehsani F, Yosephi MH, et al. Concurrent postural training and M1 anodal transcranial direct current stimulation improve postural impairment in patients with chronic low back pain. *Journal of Clinical Neuroscience*. 2019 Oct;68:224-34. doi: <https://dx.doi.org/10.1016/j.jocn.2019.07.017>. PMID: 31350080. **Exclusion:** Ineligible study design

105. Jassi FJ, Del Antonio TT, Azevedo BO, et al. Star-Shape Kinesio Taping Is Not Better Than a Minimal Intervention or Sham Kinesio Taping for Pain Intensity and Postural Control in Chronic Low Back Pain: A Randomized Controlled Trial. *Archives of Physical Medicine & Rehabilitation*. 2021 Jul;102(7):1352-60.e3. doi: <https://dx.doi.org/10.1016/j.apmr.2021.03.007>. PMID: 33819489. **Exclusion:** Ineligible intervention
106. Javdaneh N, Molayei F, Kamranifraz N. Effect of adding motor imagery training to neck stabilization exercises on pain, disability and kinesiophobia in patients with chronic neck pain. *Complementary Therapies in Clinical Practice*. 2021 Feb;42:101263. doi: <https://dx.doi.org/10.1016/j.ctcp.2020.101263>. PMID: 33276225. **Exclusion:** Inadequate duration of followup
107. Jiao J, Russell IJ, Wang W, et al. Ba-Duan-Jin alleviates pain and fibromyalgia-related symptoms in patients with fibromyalgia: results of a randomised controlled trial. *Clinical & Experimental Rheumatology*. 2019 Nov-Dec;37(6):953-62. PMID: 30789154. **Exclusion:** Inadequate duration of followup
108. Kamonseki DH, Lopes EP, van der Meer HA, et al. Effectiveness of manual therapy in patients with tension-type headache. A systematic review and meta-analysis. *Disability & Rehabilitation*. 2020 Sep 12:1-10. doi: <https://dx.doi.org/10.1080/09638288.2020.1813817>. PMID: 32924640. **Exclusion:** Systematic review used as source document
109. Kang TW, Lee JH, Park DH, et al. Effects of a finger exercise program on hand function in automobile workers with hand osteoarthritis: a randomized controlled trial. *Hand surgery & rehabilitation*. 2019;38(1):59-66. PMID: CN-02084202 NEW. **Exclusion:** Ineligible study design
110. Kaplun A, Roitman P, Rosenbloom T. Effects of Brief Guided Imagery on Female Patients Diagnosed with Fibromyalgia: An Exploratory Controlled Trial. *Alternative Therapies in Health & Medicine*. 2021 Jun;27(S1):104-13. PMID: 32827404. **Exclusion:** Inadequate duration of followup
111. Karadag S, Tasci S, Dogan N, et al. Application of heat and a home exercise program for pain and function levels in patients with knee osteoarthritis: A randomized controlled trial. *International Journal of Nursing Practice*. 2019 Oct;25(5):e12772. doi: <https://dx.doi.org/10.1111/ijn.12772>. PMID: 31436359. **Exclusion:** Inadequate duration of followup
112. Kim E, Kim YS, Kim YI, et al. Effectiveness and Safety of Polydioxanone Thread-Embedding Acupuncture as an Adjunctive Therapy for Patients with Chronic Nonspecific Neck Pain: a Randomized Controlled Trial. *Journal of alternative and complementary medicine (New York, N.Y.)*. 2019;25(4):417-26. PMID: CN-01659482. **Exclusion:** Ineligible intervention
113. Kim M, Kim J. Effects of Acupressure on Pain, Flexibility, and Substance P in Middle-Age Women with Chronic Neck Pain. *Journal of Alternative & Complementary Medicine*. 2021 Feb;27(2):160-7. doi: <https://dx.doi.org/10.1089/acm.2020.0413>. PMID: 33296258. **Exclusion:** Inadequate duration of followup
114. Kim S, Hsu FC, Groban L, et al. A pilot study of aquatic prehabilitation in adults with knee osteoarthritis undergoing total knee arthroplasty - short term outcome. *BMC Musculoskeletal Disorders*. 2021 Apr 26;22(1):388. doi: <https://dx.doi.org/10.1186/s12891-021-04253-1>. PMID: 33902505. **Exclusion:** Ineligible population
115. Kim SD. Twelve Weeks of Yoga for Chronic Nonspecific Lower Back Pain: A Meta-Analysis. *Pain Management Nursing*. 2020 12;21(6):536-42. doi: <https://dx.doi.org/10.1016/j.pmn.2020.07.002>. PMID: 32830047. **Exclusion:** Systematic review used as source document
116. Kim SH, Park KN, Kwon OY. Classification-Specific Treatment Improves Pain, Disability, Fear-Avoidance Beliefs, and Erector Spinae Muscle Activity During Walking in Patients With Low Back Pain Exhibiting Lumbar Extension-Rotation Pattern: A Randomized Controlled Trial. *Journal of Manipulative & Physiological Therapeutics*. 2020 02;43(2):123-33. doi:

- <https://dx.doi.org/10.1016/j.jmpt.2019.04.004>. PMID: 32312606. **Exclusion:** Ineligible comparator
117. Kim SK, Min A, Jeon C, et al. Clinical outcomes and cost-effectiveness of massage chair therapy versus basic physiotherapy in lower back pain patients: A randomized controlled trial. *Medicine*. 2020 Mar;99(12):e19514. doi: <https://dx.doi.org/10.1097/MD.00000000000019514>. PMID: 32195952. **Exclusion:** Inadequate duration of followup
118. Kim SY, Busch AJ, Overend TJ, et al. Flexibility exercise training for adults with fibromyalgia. *Cochrane Database of Systematic Reviews*. 2019 Sep 02;9:CD013419. doi: <https://dx.doi.org/10.1002/14651858.CD013419>. PMID: 31476271. **Exclusion:** Systematic review used as source document
119. Kim T, Lee J, Oh S, et al. Effectiveness of Simulated Horseback Riding for Patients With Chronic Low Back Pain: A Randomized Controlled Trial. *Journal of Sport Rehabilitation*. 2020 Feb 01;29(2):179-85. doi: <https://dx.doi.org/10.1123/jsr.2018-0252>. PMID: 30676224. **Exclusion:** Ineligible comparator
120. Kolbe L, Eberhardt T, Leinberger B, et al. Effectiveness of Biofeedback for Primary Headache - A Randomized Controlled Study. *Psychotherapie, Psychosomatik, medizinische Psychologie*. 2020;70(7):300-7. PMID: CN-02073107. **Exclusion:** Not English language but possibly relevant
121. Kong JT, Puetz C, Tian L, et al. Effect of Electroacupuncture vs Sham Treatment on Change in Pain Severity Among Adults With Chronic Low Back Pain: A Randomized Clinical Trial. *JAMA Network Open*. 2020 10 01;3(10):e2022787. doi: <https://dx.doi.org/10.1001/jamanetworkopen.2020.22787>. PMID: 33107921. **Exclusion:** Inadequate duration of followup
122. Krauss I, Hein T, Steinhilber B, et al. A 12-week exercise program for patients with hip osteoarthritis has no influence on gait parameters: A secondary analysis of a randomized controlled trial. *Gait & Posture*. 2020 05;78:6-12. doi: <https://dx.doi.org/10.1016/j.gaitpost.2020.03.001>. PMID: 32151918. **Exclusion:** Inadequate duration of followup
123. Krishna D, Deepeshwar S, Devi B. Yoga-Based Relaxation Technique Facilitates Sustained Attention in Patients with Low Back Pain: A Pilot Study. *Advances in Mind-Body Medicine*. 2020 Summer;34(3):11-7. PMID: 32931457. **Exclusion:** Ineligible study design
124. Kurlyandchik I, Tiralongo E, Schloss J. Safety and Efficacy of Medicinal Cannabis in the Treatment of Fibromyalgia: A Systematic Review. *Journal of Alternative & Complementary Medicine*. 2021 Mar;27(3):198-213. doi: <https://dx.doi.org/10.1089/acm.2020.0331>. PMID: 33337931. **Exclusion:** Systematic review used as source document
125. Kurt V, Aras O, Buker N. Comparison of conservative treatment with and without neural mobilization for patients with low back pain: A prospective, randomized clinical trial. *Journal of Back & Musculoskeletal Rehabilitation*. 2020;33(6):969-75. doi: <https://dx.doi.org/10.3233/BMR-181241>. PMID: 32144973. **Exclusion:** Inadequate duration of followup
126. Lascurain-Aguirrebena I, Newham DJ, Casado-Zumeta X, et al. Immediate effects of cervical mobilisations on neck muscle activity during active neck movements in patients with non-specific neck pain. A double blind placebo controlled trial. *Physiotherapy*. 2021 03;110:42-53. doi: <https://dx.doi.org/10.1016/j.physio.2019.07.003>. PMID: 33131786. **Exclusion:** Inadequate duration of followup
127. Lee E, Lee S. Impact of Cervical Sensory Feedback for Forward Head Posture on Headache Severity and Physiological Factors in Patients with Tension-type Headache: A Randomized, Single-Blind, Controlled Trial. *Medical Science Monitor*. 2019 Dec 15;25:9572-84. doi: <https://dx.doi.org/10.12659/MSM.918595>. PMID: 31838486. **Exclusion:** Inadequate duration of followup
128. Lee K, Lewis GN. Short term relief of multisite chronicpain with Bowen Therapy: A double-blind, randomized controlled trial. *Journal of Bodywork & Movement Therapies*. 2020 Oct;24(4):271-9. doi:

- <https://dx.doi.org/10.1016/j.jbmt.2020.06.025>. PMID: 33218522. **Exclusion:** Ineligible study design
129. Lena O, Todri J, Todri A, et al. The Effectiveness of the Mezieres Method in Elite Rhythmic Gymnastics Athletes With Low Back Pain: A Randomized Controlled Trial. *Journal of Sport Rehabilitation*. 2020 Sep 01;29(7):913-9. doi: <https://dx.doi.org/10.1123/jsr.2019-0204>. PMID: 31711041. **Exclusion:** Ineligible population
130. Letafatkar A, Rabiei P, Alamooti G, et al. Effect of therapeutic exercise routine on pain, disability, posture, and health status in dentists with chronic neck pain: a randomized controlled trial. *International Archives of Occupational & Environmental Health*. 2020 04;93(3):281-90. doi: <https://dx.doi.org/10.1007/s00420-019-01480-x>. PMID: 31654125. **Exclusion:** Inadequate duration of followup
131. Li C, Pei Q, Chen Y, et al. The response-time relationship and covariate effects of acupuncture for chronic pain: A systematic review and model-based longitudinal meta-analysis. *European Journal of Pain*. 2020 10;24(9):1653-65. doi: <https://dx.doi.org/10.1002/ejp.1617>. PMID: 32533885. **Exclusion:** Systematic review used as source document
132. Li R, Chen H, Feng J, et al. Effectiveness of Traditional Chinese Exercise for Symptoms of Knee Osteoarthritis: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2020 10 27;17(21):27. doi: <https://dx.doi.org/10.3390/ijerph17217873>. PMID: 33121082. **Exclusion:** Systematic review used as source document
133. Li YX, Yuan SE, Jiang JQ, et al. Systematic review and meta-analysis of effects of acupuncture on pain and function in non-specific low back pain. *Acupuncture in Medicine*. 2020 08;38(4):235-43. doi: <https://dx.doi.org/10.1136/acupmed-2017-011622>. PMID: 32458717. **Exclusion:** Systematic review used as source document
134. Liao FY, Lin CL, Lo SF, et al. Efficacy of Acupoints Dual-Frequency Low-Level Laser Therapy on Knee Osteoarthritis. *Evidence-Based Complementary & Alternative Medicine: eCAM*. 2020;2020:6979105. doi: <https://dx.doi.org/10.1155/2020/6979105>. PMID: 33029170. **Exclusion:** Inadequate duration of followup
135. Lin KY, Tsai YJ, Hsu PY, et al. Effects of Sling Exercise for Neck Pain: A Systematic Review and Meta-Analysis. *Physical Therapy*. 2021 Apr 30;30:30. doi: <https://dx.doi.org/10.1093/ptj/pzab120>. PMID: 33929540. **Exclusion:** Systematic review used as source document
136. Lin YP, Su YH, Chin SF, et al. Light-emitting diode photobiomodulation therapy for non-specific low back pain in working nurses: A single-center, double-blind, prospective, randomized controlled trial. *Medicine*. 2020 Aug 07;99(32):e21611. doi: <https://dx.doi.org/10.1097/MD.00000000000021611>. PMID: 32769919. **Exclusion:** Ineligible intervention
137. Liu J, Yeung A, Xiao T, et al. Chen-Style Tai Chi for Individuals (Aged 50 Years Old or Above) with Chronic Non-Specific Low Back Pain: a Randomized Controlled Trial. *International journal of environmental research and public health*. 2019;16(3) PMID: CN-01707017. **Exclusion:** Inadequate duration of followup
138. Liu M, Tong Y, Chai L, et al. Effects of Auricular Point Acupressure on Pain Relief: A Systematic Review. *Pain Management Nursing*. 2021 Jun;22(3):268-80. doi: <https://dx.doi.org/10.1016/j.pmn.2020.07.007>. PMID: 32950391. **Exclusion:** Systematic review used as source document
139. Lopez-de-Uralde-Villanueva I, Beltran-Alacreu H, Fernandez-Carnero J, et al. Pain management using a multimodal physiotherapy program including a biobehavioral approach for chronic nonspecific neck pain: a randomized controlled trial. *Physiotherapy Theory & Practice*. 2020 Jan;36(1):45-62. doi: <https://dx.doi.org/10.1080/09593985.2018.1480678>. PMID: 29889599. **Exclusion:** Ineligible intervention
140. Low MY, Lacson C, Zhang F, et al. Vocal Music Therapy for Chronic Pain: A Mixed Methods Feasibility Study. *Journal of Alternative & Complementary Medicine*. 2020 Feb;26(2):113-22. doi:

- <https://dx.doi.org/10.1089/acm.2019.0249>. PMID: 31750726. **Exclusion:** Ineligible population
141. Luan L, Bousie J, Pranata A, et al. Stationary cycling exercise for knee osteoarthritis: A systematic review and meta-analysis. *Clinical Rehabilitation*. 2021 Apr;35(4):522-33. doi: <https://dx.doi.org/10.1177/0269215520971795>. PMID: 33167714. **Exclusion:** Systematic review used as source document
142. Ludvigsson ML, Peterson G, Peolsson A. Neck-specific exercise for radiating pain and neurological deficits in chronic whiplash, a 1-year follow-up of a randomised clinical trial. *Scientific Reports*. 2020 04 21;10(1):6758. doi: <https://dx.doi.org/10.1038/s41598-020-62722-4>. PMID: 32317700. **Exclusion:** Ineligible intervention
143. Lv ZT, Shen LL, Zhu B, et al. Effects of intensity of electroacupuncture on chronic pain in patients with knee osteoarthritis: a randomized controlled trial. *Arthritis research & therapy*. 2019;21(1):120. PMID: CN-01940220. **Exclusion:** Inadequate duration of followup
144. Machado-Oliveira L, da Silva Gauto YO, de Santana Neto FJ, et al. Effects of Different Exercise Intensities on Headache: A Systematic Review. *American Journal of Physical Medicine & Rehabilitation*. 2020 05;99(5):390-6. doi: <https://dx.doi.org/10.1097/PHM.0000000000001349>. PMID: 31725018. **Exclusion:** Systematic review used as source document
145. Madadi-Shad M, Jafarnezhadgero AA, Sheikhalizade H, et al. Effect of a corrective exercise program on gait kinetics and muscle activities in older adults with both low back pain and pronated feet: A double-blind, randomized controlled trial. *Gait & Posture*. 2020 02;76:339-45. doi: <https://dx.doi.org/10.1016/j.gaitpost.2019.12.026>. PMID: 31896537. **Exclusion:** Inadequate duration of followup
146. Mahler EAM, Minten MJ, Leseman-Hoogenboom MM, et al. Effectiveness of low-dose radiation therapy on symptoms in patients with knee osteoarthritis: a randomised, double-blinded, sham-controlled trial. *Annals of the rheumatic diseases*. 2019;78(1):83-90. PMID: CN-02000205. **Exclusion:** Ineligible intervention
147. Marconcin P, Yazigi F, Teles J, et al. The effectiveness of a randomised clinical trial of PLE2 NO self-management and exercise programme for knee osteoarthritis to improve self-efficacy. *Musculoskeletal Care*. 2021 Jun 02;02:02. doi: <https://dx.doi.org/10.1002/msc.1573>. PMID: 34077602. **Exclusion:** Inadequate duration of followup
148. Martimbianco ALC, Porfirio GJ, Pacheco RL, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic neck pain. *Cochrane Database of Systematic Reviews*. 2019 12 12;12:CD011927. doi: <https://dx.doi.org/10.1002/14651858.CD011927.pub2>. PMID: 31830313. **Exclusion:** Systematic review used as source document
149. Martinez-Calderon J, Flores-Cortes M, Morales-Asencio JM, et al. Intervention Therapies to Reduce Pain-Related Fear in Fibromyalgia Syndrome: A Systematic Review of Randomized Clinical Trials. *Pain Medicine*. 2021 02 23;22(2):481-98. doi: <https://dx.doi.org/10.1093/pm/pnaa331>. PMID: 32989450. **Exclusion:** Systematic review used as source document
150. Mascarenhas RO, Souza MB, Oliveira MX, et al. Association of Therapies With Reduced Pain and Improved Quality of Life in Patients With Fibromyalgia: A Systematic Review and Meta-analysis. *JAMA Internal Medicine*. 2021 Jan 01;181(1):104-12. doi: <https://dx.doi.org/10.1001/jamainternmed.2020.5651>. PMID: 33104162. **Exclusion:** Systematic review used as source document
151. McCurry SM, Zhu W, Von Korff M, et al. Effect of Telephone Cognitive Behavioral Therapy for Insomnia in Older Adults With Osteoarthritis Pain: A Randomized Clinical Trial. *JAMA Internal Medicine*. 2021 Apr 01;181(4):530-8. doi: <https://dx.doi.org/10.1001/jamainternmed.2020.9049>. PMID: 33616613. **Exclusion:** Ineligible intervention
152. Mehri A, Letafatkar A, Khosrokiani Z. Effects of Corrective Exercises on Posture, Pain, and Muscle Activation of Patients With Chronic Neck Pain Exposed to Anterior-Posterior Perturbation. *Journal of Manipulative & Physiological Therapeutics*. 2020 05;43(4):311-24. doi:

- <https://dx.doi.org/10.1016/j.jmpt.2018.11.032>. PMID: 32723668. **Exclusion:** Inadequate duration of followup
153. Mohamadi M, Rojhani-Shirazi Z, Assadsangabi R, et al. Can the Positional Release Technique Affect Central Sensitization in Patients With Chronic Tension-Type Headache? A Randomized Clinical Trial. *Archives of Physical Medicine & Rehabilitation*. 2020 10;101(10):1696-703. doi: <https://dx.doi.org/10.1016/j.apmr.2020.05.028>. PMID: 32673652. **Exclusion:** Inadequate duration of followup
154. Moseng T, Dagfinrud H, van Bodegom-Vos L, et al. Low adherence to exercise may have influenced the proportion of OMERACT-OARSI responders in an integrated osteoarthritis care model: secondary analyses from a cluster-randomised stepped-wedge trial. *BMC Musculoskeletal Disorders*. 2020 Apr 13;21(1):236. doi: <https://dx.doi.org/10.1186/s12891-020-03235-z>. PMID: 32284049. **Exclusion:** Ineligible intervention
155. Movahedi Najafabadi M, Ghafari S, Nazari F, et al. The effect of acupressure on quality of life among female nurses with chronic back pain. *Applied Nursing Research*. 2020 02;51:151175. doi: <https://dx.doi.org/10.1016/j.apnr.2019.05.020>. PMID: 31831270. **Exclusion:** Ineligible study design
156. Mu J, Furlan AD, Lam WY, et al. Acupuncture for chronic nonspecific low back pain. *Cochrane Database of Systematic Reviews*. 2020 12 11;12:CD013814. doi: <https://dx.doi.org/10.1002/14651858.CD013814>. PMID: 33306198. **Exclusion:** Systematic review used as source document
157. Mukhtar NB, Meeus M, Gursen C, et al. Effectiveness of Hands-Off Therapy in the Management of Primary Headache: A Systematic Review and Meta-Analysis. *Evaluation & the Health Professions*. 2021 Jan 07;163278720983408. doi: <https://dx.doi.org/10.1177/0163278720983408>. PMID: 33406891. **Exclusion:** Systematic review used as source document
158. Munukka M, Waller B, Hakkinen A, et al. Effects of progressive aquatic resistance training on symptoms and quality of life in women with knee osteoarthritis: A secondary analysis. *Scandinavian Journal of Medicine & Science in Sports*. 2020 Jun;30(6):1064-72. doi: <https://dx.doi.org/10.1111/sms.13630>. PMID: 31999876. **Exclusion:** Inadequate duration of followup
159. Murphy SL, Harris RE, Keshavarzi NR, et al. Self-Administered Acupressure for Chronic Low Back Pain: A Randomized Controlled Pilot Trial. *Pain Medicine*. 2019 12 01;20(12):2588-97. doi: <https://dx.doi.org/10.1093/pm/pnz138>. PMID: 31237610. **Exclusion:** Inadequate duration of followup
160. Nadal-Nicolas Y, Rubio-Arias JA, Martinez-Olcina M, et al. Effects of Manual Therapy on Fatigue, Pain, and Psychological Aspects in Women with Fibromyalgia. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2020 06 26;17(12):26. doi: <https://dx.doi.org/10.3390/ijerph17124611>. PMID: 32604939. **Exclusion:** Ineligible study design
161. Nakamaru K, Aizawa J, Kawarada K, et al. Immediate effects of thoracic spine self-mobilization in patients with mechanical neck pain: a randomized controlled trial. *Journal of bodywork and movement therapies*. 2019;23(2):417-24. PMID: CN-02083937 NEW. **Exclusion:** Inadequate duration of followup
162. Nambi G, Abdelbasset WK, Alqahtani BA, et al. Isokinetic back training is more effective than core stabilization training on pain intensity and sports performances in football players with chronic low back pain: A randomized controlled trial. *Medicine*. 2020 May 22;99(21):e20418. doi: <https://dx.doi.org/10.1097/MD.00000000000020418>. PMID: 32481345. **Exclusion:** Ineligible comparator
163. Nambi G, Abdelbasset WK, Alrawaili SM, et al. Virtual reality or isokinetic training; its effect on pain, kinesiophobia and serum stress hormones in chronic low back pain: A randomized controlled trial. *Technology & Health Care*. 2021;29(1):155-66. doi: <https://dx.doi.org/10.3233/THC-202301>. PMID: 32831210. **Exclusion:** Ineligible comparator

164. Nambi G, Abdelbasset WK, Alsubaie SF, et al. Short-Term Psychological and Hormonal Effects of Virtual Reality Training on Chronic Low Back Pain in Soccer Players. *Journal of Sport Rehabilitation*. 2021 Feb 16:1-10. doi: <https://dx.doi.org/10.1123/jsr.2020-0075>. PMID: 33596538. **Exclusion:** Ineligible comparator
165. Namnaqani FI, Mashabi AS, Yaseen KM, et al. The effectiveness of McKenzie method compared to manual therapy for treating chronic low back pain: a systematic review. *Journal of Musculoskeletal Neuronal Interactions*. 2019 12 01;19(4):492-9. PMID: 31789300. **Exclusion:** Systematic review used as source document
166. Nduwimana I, Nindorera F, Thonnard JL, et al. Effectiveness of walking versus mind-body therapies in chronic low back pain: A systematic review and meta-analysis of recent randomized controlled trials. *Medicine*. 2020 Aug 28;99(35):e21969. doi: <https://dx.doi.org/10.1097/MD.00000000000021969>. PMID: 32871946. **Exclusion:** Systematic review used as source document
167. Nelligan RK, Hinman RS, Kasza J, et al. Effects of a Self-directed Web-Based Strengthening Exercise and Physical Activity Program Supported by Automated Text Messages for People With Knee Osteoarthritis: A Randomized Clinical Trial. *JAMA Internal Medicine*. 2021 Jun 01;181(6):776-85. doi: <https://dx.doi.org/10.1001/jamainternmed.2021.0991>. PMID: 33843948. **Exclusion:** Ineligible intervention
168. Nery M, Natour J, Jennings F, et al. Effects of a progressive resistance exercise program in patients with hand osteoarthritis: A randomized, controlled trial with a blinded assessor. *Clinical Rehabilitation*. 2021 Jul 09:2692155211030622. doi: <https://dx.doi.org/10.1177/02692155211030622>. PMID: 34240642. **Exclusion:** Inadequate duration of followup
169. Niederer D, Engel T, Vogt L, et al. Motor Control Stabilisation Exercise for Patients with Non-Specific Low Back Pain: A Prospective Meta-Analysis with Multilevel Meta-Regressions on Intervention Effects. *Journal of Clinical Medicine*. 2020 Sep 22;9(9):22. doi: <https://dx.doi.org/10.3390/jcm9093058>. PMID: 32971921. **Exclusion:** Systematic review used as source document
170. Noori SA, Rasheed A, Aiyer R, et al. Therapeutic Ultrasound for Pain Management in Chronic Low Back Pain and Chronic Neck Pain: A Systematic Review. *Pain Medicine*. 2020 11 07;21(7):1482-93. doi: <https://dx.doi.org/10.1093/pm/pny287>. PMID: 30649460. **Exclusion:** Systematic review used as source document
171. Norouzi E, Hosseini F, Vaezmosavi M, et al. Zumba dancing and aerobic exercise can improve working memory, motor function, and depressive symptoms in female patients with Fibromyalgia. *European Journal of Sport Science EJSS : Official Journal of the European College of Sport Science*. 2020 Aug;20(7):981-91. doi: <https://dx.doi.org/10.1080/17461391.2019.1683610>. PMID: 31630663. **Exclusion:** Inadequate duration of followup
172. Novak S, Guerron G, Zou Z, et al. New Guidelines for Electrical Stimulation Parameters in Adult Patients With Knee Osteoarthritis Based on a Systematic Review of the Current Literature. *American Journal of Physical Medicine & Rehabilitation*. 2020 08;99(8):682-8. doi: <https://dx.doi.org/10.1097/PHM.00000000000001409>. PMID: 32167955. **Exclusion:** Systematic review used as source document
173. O'Keeffe M, O'Sullivan P, Purtill H, et al. Cognitive functional therapy compared with a group-based exercise and education intervention for chronic low back pain: a multicentre randomised controlled trial (RCT). *British Journal of Sports Medicine*. 2020 Jul;54(13):782-9. doi: <https://dx.doi.org/10.1136/bjsports-2019-100780>. PMID: 31630089. **Exclusion:** Ineligible intervention
174. Oh SL, Kim DY, Bae JH, et al. Effects of rural community-based integrated exercise and health education programs on the mobility function of older adults with knee osteoarthritis. *Aging Clinical & Experimental Research*. 2020 Feb 04;04:04. doi: <https://dx.doi.org/10.1007/s40520-020-01474-7>. PMID: 32020485. **Exclusion:** Inadequate duration of followup
175. Ots T, Kandirian A, Szilagyi I, et al. The selection of dermatomes for sham (placebo)

- acupuncture points is relevant for the outcome of acupuncture studies: a systematic review of sham (placebo)-controlled randomized acupuncture trials. *Acupuncture in Medicine*. 2020 08;38(4):211-26. doi: <https://dx.doi.org/10.1177/0964528419889636>. PMID: 32026725. **Exclusion:** Systematic review used as source document
176. Ouellet P, Lafrance S, Pizzi A, et al. Region-specific exercises versus general exercises approaches in the management of spinal and peripheral musculoskeletal disorders: a systematic review with meta-analyses of randomized controlled trials. *Archives of Physical Medicine & Rehabilitation*. 2021 Mar 05;05:05. doi: <https://dx.doi.org/10.1016/j.apmr.2021.01.093>. PMID: 33684362. **Exclusion:** Systematic review used as source document
177. Paganini S, Lin J, Kahlke F, et al. A guided and unguided internet- and mobile-based intervention for chronic pain: health economic evaluation alongside a randomised controlled trial. *BMJ open*. 2019;9(4):e023390. PMID: CN-01941119. **Exclusion:** Ineligible population
178. Papaconstantinou E, Cancelliere C, Verville L, et al. Effectiveness of non-pharmacological interventions on sleep characteristics among adults with musculoskeletal pain and a comorbid sleep problem: a systematic review. *Chiropractic & manual therapies*. 2021 Jul 08;29(1):23. doi: <https://dx.doi.org/10.1186/s12998-021-00381-6>. PMID: 34238325. **Exclusion:** Systematic review used as source document
179. Pardos-Gascon EM, Narambuena L, Leal-Costa C, et al. Differential efficacy between cognitive-behavioral therapy and mindfulness-based therapies for chronic pain: Systematic review. *International Journal of Clinical & Health Psychology*. 2021 Jan-Apr;21(1):100197. doi: <https://dx.doi.org/10.1016/j.ijchp.2020.08.001>. PMID: 33363580. **Exclusion:** Systematic review used as source document
180. Park S, Park S, Min S, et al. A Randomized Controlled Trial Investigating the Effects of Equine Simulator Riding on Low Back Pain, Morphological Changes, and Trunk Musculature in Elderly Women. *Medicina*. 2020 Nov 13;56(11):13. doi: <https://dx.doi.org/10.3390/medicina56110610>. PMID: 33202928. **Exclusion:** Inadequate duration of followup
181. Park SY, Hwang EH, Cho JH, et al. Comparative Effectiveness of Chuna Manipulative Therapy for Non-Acute Lower Back Pain: A Multi-Center, Pragmatic, Randomized Controlled Trial. *Journal of Clinical Medicine*. 2020 Jan 05;9(1):05. doi: <https://dx.doi.org/10.3390/jcm9010144>. PMID: 31948083. **Exclusion:** Ineligible population
182. Patel K, Sutherland H, Henshaw J, et al. Effects of neurofeedback in the management of chronic pain: A systematic review and meta-analysis of clinical trials. *European Journal of Pain*. 2020 09;24(8):1440-57. doi: <https://dx.doi.org/10.1002/ejp.1612>. PMID: 32502283. **Exclusion:** Systematic review used as source document
183. Paulo LR, Lacerda ACR, Martins FLM, et al. Can a Single Trial of a Thoracolumbar Myofascial Release Technique Reduce Pain and Disability in Chronic Low Back Pain? A Randomized Balanced Crossover Study. *Journal of Clinical Medicine*. 2021 May 07;10(9):07. doi: <https://dx.doi.org/10.3390/jcm10092006>. PMID: 34067152. **Exclusion:** Inadequate duration of followup
184. Pedersini P, Valdes K, Cantero-Tellez R, et al. Effects of Neurodynamic Mobilizations on Pain Hypersensitivity in Patients With Hand Osteoarthritis Compared to Robotic Assisted Mobilization: A Randomized Controlled Trial. *Arthritis care & research*. 2021 02;73(2):232-9. doi: <https://dx.doi.org/10.1002/acr.24103>. PMID: 31675184. **Exclusion:** Ineligible comparator
185. Pei JH, Ma T, Nan RL, et al. Mindfulness-Based Cognitive Therapy for Treating Chronic Pain A Systematic Review and Meta-analysis. *Psychology Health & Medicine*. 2021 03;26(3):333-46. doi: <https://dx.doi.org/10.1080/13548506.2020.1849746>. PMID: 33241941. **Exclusion:** Systematic review used as source document
186. Perlini C, Donisi V, Del Piccolo L. From research to clinical practice: a systematic review of the implementation of psychological interventions for chronic headache in adults. *BMC Health Services Research*. 2020 May 25;20(1):459. doi:

- <https://dx.doi.org/10.1186/s12913-020-05172-y>. PMID: 32450871. **Exclusion:** Systematic review used as source document
187. Perlman A, Fogerite SG, Glass O, et al. Efficacy and Safety of Massage for Osteoarthritis of the Knee: a Randomized Clinical Trial. *Journal of general internal medicine*. 2019;34(3):379-86. PMID: CN-02145187. **Exclusion:** Ineligible study design
188. Petrozzi MJ, Leaver A, Ferreira PH, et al. Addition of MoodGYM to physical treatments for chronic low back pain: A randomized controlled trial. *Chiropractic & manual therapies*. 2019;27:54. doi: <https://dx.doi.org/10.1186/s12998-019-0277-4>. PMID: 31673330. **Exclusion:** Ineligible intervention
189. Petterson S, Plancher K, Klyve D, et al. Low-Intensity Continuous Ultrasound for the Symptomatic Treatment of Upper Shoulder and Neck Pain: A Randomized, Double-Blind Placebo-Controlled Clinical Trial. *Journal of pain research*. 2020;13:1277-87. doi: <https://dx.doi.org/10.2147/JPR.S247463>. PMID: 32606899. **Exclusion:** Ineligible study design
190. Phattharasupharerk S, Purepong N, Eksakulkla S, et al. Effects of Qigong practice in office workers with chronic non-specific low back pain: a randomized control trial. *Journal of bodywork and movement therapies*. 2019;23(2):375-81. PMID: CN-02083942 NEW. **Exclusion:** Inadequate duration of followup
191. Pico-Espinosa OJ, Aboagye E, Cote P, et al. Deep tissue massage, strengthening and stretching exercises, and a combination of both compared with advice to stay active for subacute or persistent non-specific neck pain: A cost-effectiveness analysis of the Stockholm Neck trial (STONE). *Musculoskeletal Science & Practice*. 2020 04;46:102109. doi: <https://dx.doi.org/10.1016/j.msksp.2020.102109>. PMID: 31989965. **Exclusion:** Ineligible outcomes
192. Pietrosimone B, Luc-Harkey BA, Harkey MS, et al. Using TENS to Enhance Therapeutic Exercise in Individuals with Knee Osteoarthritis. *Medicine & Science in Sports & Exercise*. 2020 10;52(10):2086-95. doi: <https://dx.doi.org/10.1249/00000000000002353>. PMID: 32251254. **Exclusion:** Ineligible comparator
193. Pitsillides A, Stasinopoulos D, Giannakou K. The effects of cognitive behavioural therapy delivered by physical therapists in knee osteoarthritis pain: A systematic review and meta-analysis of randomized controlled trials. *Journal of Bodywork & Movement Therapies*. 2021 Jan;25:157-64. doi: <https://dx.doi.org/10.1016/j.jbmt.2020.11.002>. PMID: 33714488. **Exclusion:** Systematic review used as source document
194. Polaski AM, Phelps AL, Smith TJ, et al. Integrated Meditation and Exercise Therapy: A Randomized Controlled Pilot of a Combined Nonpharmacological Intervention Focused on Reducing Disability and Pain in Patients with Chronic Low Back Pain. *Pain Medicine*. 2021 02 23;22(2):444-58. doi: <https://dx.doi.org/10.1093/pm/pnaa403>. PMID: 33621332. **Exclusion:** Inadequate duration of followup
195. Prado ERA, Meireles SM, Carvalho ACA, et al. Influence of isostretching on patients with chronic low back pain. A randomized controlled trial. *Physiotherapy Theory & Practice*. 2021 Feb;37(2):287-94. doi: <https://dx.doi.org/10.1080/09593985.2019.1625091>. PMID: 31161855. **Exclusion:** Inadequate duration of followup
196. Pujol J, Ramos-Lopez D, Blanco-Hinojo L, et al. Testing the effects of gentle vibrotactile stimulation on symptom relief in fibromyalgia. *Arthritis Research & Therapy*. 2019 06 14;21(1):148. doi: <https://dx.doi.org/10.1186/s13075-019-1932-9>. PMID: 31200775. **Exclusion:** Inadequate duration of followup
197. Qing W, Shi X, Zhang Q, et al. Effect of Therapeutic Ultrasound for Neck Pain: A Systematic Review and Meta-Analysis. *Archives of Physical Medicine & Rehabilitation*. 2021 Mar 17;17:17. doi: <https://dx.doi.org/10.1016/j.apmr.2021.02.009>. PMID: 33722564. **Exclusion:** Systematic review used as source document
198. Racine M, Jensen MP, Harth M, et al. Operant Learning Versus Energy Conservation Activity Pacing Treatments in a Sample of Patients With Fibromyalgia

- Syndrome: a Pilot Randomized Controlled Trial. *Journal of pain*. 2019;20(4):420-39. PMID: CN-02145231 NEW. **Exclusion:** Ineligible comparator
199. Rae L, Dougherty P, Evertz N. Yoga vs Stretching in Veterans With Chronic Lower Back Pain and the Role of Mindfulness: A Pilot Randomized Controlled Trial. *Journal of Chiropractic Medicine*. 2020 Jun;19(2):101-10. doi: <https://dx.doi.org/10.1016/j.jcm.2019.10.005>. PMID: 33318728. **Exclusion:** Ineligible study design
200. Raghava Neelapala YV, Bhagat M, Shah P. Hip Muscle Strengthening for Knee Osteoarthritis: A Systematic Review of Literature. *Journal of Geriatric Physical Therapy*. 2020 Apr/Jun;43(2):89-98. doi: <https://dx.doi.org/10.1519/JPT.00000000000000214>. PMID: 30407271. **Exclusion:** Systematic review used as source document
201. Rampazo EP, Martignago CCS, de Noronha M, et al. Transcutaneous electrical stimulation in neck pain: a systematic review and meta-analysis. *European Journal of Pain*. 2021 Jul 20;20:20. doi: <https://dx.doi.org/10.1002/ejp.1845>. PMID: 34288255. **Exclusion:** Systematic review used as source document
202. Raposo F, Ramos M, Lucia Cruz A. Effects of exercise on knee osteoarthritis: A systematic review. *Musculoskeletal Care*. 2021 Mar 05;05:05. doi: <https://dx.doi.org/10.1002/msc.1538>. PMID: 33666347. **Exclusion:** Systematic review used as source document
203. Razzaq A, Sajjad AG, Yasin S, et al. Comparison of Cyriax manipulation with traditional physical therapy for the management of cervical discogenic problems. A randomized control trial. *JPMA - Journal of the Pakistan Medical Association*. 2020 Aug;70(8):1329-33. doi: <https://dx.doi.org/10.5455/JPMA.13250>. PMID: 32794481. **Exclusion:** Inadequate duration of followup
204. Rickardsson J, Gentili C, Holmstrom L, et al. Internet-delivered acceptance and commitment therapy as microlearning for chronic pain: A randomized controlled trial with 1-year follow-up. *European Journal of Pain*. 2021 05;25(5):1012-30. doi: <https://dx.doi.org/10.1002/ejp.1723>. PMID: 33460240. **Exclusion:** Ineligible population
205. Rodriguez-Roca B, Urcola-Pardo F, Anguas-Gracia A, et al. Impact of Reducing Sitting Time in Women with Fibromyalgia and Obesity: A Randomized Controlled Trial. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2021 Jun 09;18(12):09. doi: <https://dx.doi.org/10.3390/ijerph18126237>. PMID: 34207661. **Exclusion:** Ineligible intervention
206. Roseen EJ, Gerlovin H, Felson DT, et al. Which Chronic Low Back Pain Patients Respond Favorably to Yoga, Physical Therapy, and a Self-care Book? Responder Analyses from a Randomized Controlled Trial. *Pain Medicine*. 2021 02 04;22(1):165-80. doi: <https://dx.doi.org/10.1093/pm/pnaa153>. PMID: 32662833. **Exclusion:** Inadequate duration of followup
207. Rosenberg NR, Petersen SB, Begtrup LM, et al. Early Occupational Intervention for People with Low Back Pain in Physically Demanding Jobs: 1-year Follow-up Results of the Randomized Controlled GOBACK Trial. *Spine*. 2021 Mar 15;46(6):347-55. doi: <https://dx.doi.org/10.1097/BRS.00000000000003793>. PMID: 33181779. **Exclusion:** Ineligible population
208. Rubira A, Rubira MC, Rubira LA, et al. Comparison of the effects of low-level laser and pulsed and continuous ultrasound on pain and physical disability in chronic non-specific low back pain: a randomized controlled clinical trial. *Advances in Rheumatology*. 2019 12 17;59(1):57. doi: <https://dx.doi.org/10.1186/s42358-019-0099-z>. PMID: 31847915. **Exclusion:** Inadequate duration of followup
209. Salm DC, Belmonte LAO, Emer AA, et al. Aquatic exercise and Far Infrared (FIR) modulates pain and blood cytokines in fibromyalgia patients: A double-blind, randomized, placebo-controlled pilot study. *Journal of Neuroimmunology*. 2019 12 15;337:577077. doi: <https://dx.doi.org/10.1016/j.jneuroim.2019.577077>. PMID: 31655422. **Exclusion:** Ineligible intervention

210. Sander LB, Paganini S, Terhorst Y, et al. Effectiveness of a Guided Web-Based Self-help Intervention to Prevent Depression in Patients With Persistent Back Pain: The PROD-BP Randomized Clinical Trial. *JAMA Psychiatry*. 2020 10 01;77(10):1001-11. doi: <https://dx.doi.org/10.1001/jamapsychiatry.2020.1021>. PMID: 32459348. **Exclusion:** Ineligible intervention
211. Sarig Bahat H, Hadar D, Treleaven J. Predictors for Positive Response to Home Kinematic Training in Chronic Neck Pain. *Journal of Manipulative & Physiological Therapeutics*. 2020 10;43(8):779-90. doi: <https://dx.doi.org/10.1016/j.jmpt.2019.12.008>. PMID: 32829943. **Exclusion:** Ineligible study design
212. Sarmiento CVM, Moon S, Pfeifer T, et al. The therapeutic efficacy of Qigong exercise on the main symptoms of fibromyalgia: A pilot randomized clinical trial. *Integrative Medicine Research*. 2020 Dec;9(4):100416. doi: <https://dx.doi.org/10.1016/j.imr.2020.100416>. PMID: 32455108. **Exclusion:** Ineligible study design
213. Sato T, Shimizu K, Shiko Y, et al. Effects of Nintendo Ring Fit Adventure Exergame on Pain and Psychological Factors in Patients with Chronic Low Back Pain. *Games for Health Journal*. 2021 Jun;10(3):158-64. doi: <https://dx.doi.org/10.1089/g4h.2020.0180>. PMID: 33891508. **Exclusion:** Inadequate duration of followup
214. Sauch Valmana G, Vidal-Alaball J, Poch PR, et al. Effects of a Physical Exercise Program on Patients Affected with Fibromyalgia. *Journal of Primary Care & Community Health*. 2020 Jan-Dec;11:2150132720965071. doi: <https://dx.doi.org/10.1177/2150132720965071>. PMID: 33084477. **Exclusion:** Inadequate duration of followup
215. Schemer L, Schroeder A, Ornbol E, et al. Exposure and cognitive-behavioural therapy for chronic back pain: an RCT on treatment processes. *European journal of pain* (London, England). 2019;23(3):526-38. PMID: CN-01690849. **Exclusion:** Ineligible comparator
216. Schiller J, Karst M, Kellner T, et al. Combination of acupuncture and medical training therapy on tension type headache: Results of a randomised controlled pilot study. *Cephalalgia*. 2021 Jul;41(8):879-93. doi: <https://dx.doi.org/10.1177/0333102421989620>. PMID: 33563049. **Exclusion:** Ineligible population
217. Schlenk EA, Fitzgerald GK, Rogers JC, et al. Promoting Physical Activity in Older Adults With Knee Osteoarthritis and Hypertension: A Randomized Controlled Trial. *Journal of Aging & Physical Activity*. 2020 09 04;29(2):207-18. doi: <https://dx.doi.org/10.1123/japa.2019-0498>. PMID: 32887850. **Exclusion:** Ineligible intervention
218. Schlicker S, Baumeister H, Buntrock C, et al. A Web- and Mobile-Based Intervention for Comorbid, Recurrent Depression in Patients With Chronic Back Pain on Sick Leave (Get.Back): Pilot Randomized Controlled Trial on Feasibility, User Satisfaction, and Effectiveness. *JMIR Mental Health*. 2020 Apr 15;7(4):e16398. doi: <https://dx.doi.org/10.2196/16398>. PMID: 32293577. **Exclusion:** Ineligible intervention
219. Schmid AA, Fruhauf CA, Sharp JL, et al. Yoga for People With Chronic Pain in a Community-Based Setting: A Feasibility and Pilot RCT. *Journal of Evidence-based Integrative Medicine*. 2019 Jan-Dec;24:2515690X19863763. doi: <https://dx.doi.org/10.1177/2515690X19863763>. PMID: 31394910. **Exclusion:** Ineligible population
220. Schmid AA, Van Puymbroeck M, Fruhauf CA, et al. Yoga improves occupational performance, depression, and daily activities for people with chronic pain. *Work*. 2019;63(2):181-9. doi: <https://dx.doi.org/10.3233/WOR-192919>. PMID: 31156199. **Exclusion:** Ineligible population
221. Schulze NB, Salemi MM, de Alencar GG, et al. Efficacy of Manual Therapy on Pain, Impact of Disease, and Quality of Life in the Treatment of Fibromyalgia: A Systematic Review. *Pain Physician*. 2020 09;23(5):461-76. PMID: 32967389. **Exclusion:** Systematic review used as source document
222. Seguin-Fowler R, Graham M, Ward J, et al. Feasibility of a yoga intervention to decrease

- pain in older women: a randomized controlled pilot study. *BMC Geriatrics*. 2020 10 12;20(1):400. doi: <https://dx.doi.org/10.1186/s12877-020-01818-y>. PMID: 33046009. **Exclusion:** Ineligible population
223. Shamsi M, Shahsavari S, Safari A, et al. A randomized clinical trial for the effect of static stretching and strengthening exercise on pelvic tilt angle in LBP patients. *Journal of Bodywork & Movement Therapies*. 2020 Jul;24(3):15-20. doi: <https://dx.doi.org/10.1016/j.jbmt.2020.02.001>. PMID: 32825981. **Exclusion:** Ineligible comparator
224. Sherman KJ, Wellman RD, Hawkes RJ, et al. Tai Chi for Chronic Low Back Pain in Older Adults: A Feasibility Trial. *Journal of Alternative & Complementary Medicine*. 2020 Mar;26(3):176-89. doi: <https://dx.doi.org/10.1089/acm.2019.0438>. PMID: 32013530. **Exclusion:** Ineligible outcomes
225. Shimo K, Hasegawa M, Mizutani S, et al. Effects of a 12-week workplace counseling program on physical activity and low back pain: A pilot randomized controlled study. *Journal of Back & Musculoskeletal Rehabilitation*. 2021 Apr 30;30:30. doi: <https://dx.doi.org/10.3233/BMR-200178>. PMID: 33935064. **Exclusion:** Ineligible intervention
226. Shimoura K, Iijima H, Suzuki Y, et al. Immediate Effects of Transcutaneous Electrical Nerve Stimulation on Pain and Physical Performance in Individuals With Preradiographic Knee Osteoarthritis: a Randomized Controlled Trial. *Archives of physical medicine and rehabilitation*. 2019;100(2):300-6.e1. PMID: CN-02000235 NEW. **Exclusion:** Inadequate duration of followup
227. Simoni G, Bozzolan M, Bonnini S, et al. Effectiveness of standard cervical physiotherapy plus diaphragm manual therapy on pain in patients with chronic neck pain: A randomized controlled trial. *Journal of Bodywork & Movement Therapies*. 2021 Apr;26:481-91. doi: <https://dx.doi.org/10.1016/j.jbmt.2020.12.002>. PMID: 33992285. **Exclusion:** Ineligible intervention
228. Sitthipornvorakul E, Sihawong R, Waongenngarm P, et al. The effects of walking intervention on preventing neck pain in office workers: A randomized controlled trial. *Journal of Occupational Health*. 2020 Jan;62(1):e12106. doi: <https://dx.doi.org/10.1002/1348-9585.12106>. PMID: 31849170. **Exclusion:** Ineligible population
229. Skillgate E, Pico-Espinosa OJ, Cote P, et al. Effectiveness of deep tissue massage therapy, and supervised strengthening and stretching exercises for subacute or persistent disabling neck pain. The Stockholm Neck (STONE) randomized controlled trial. *Musculoskeletal Science & Practice*. 2020 02;45:102070. doi: <https://dx.doi.org/10.1016/j.msksp.2019.102070>. PMID: 31655314. **Exclusion:** Ineligible population
230. Smith J, Faux SG, Gardner T, et al. Reboot Online: A Randomized Controlled Trial Comparing an Online Multidisciplinary Pain Management Program with Usual Care for Chronic Pain. *Pain Medicine*. 2019 12 01;20(12):2385-96. doi: <https://dx.doi.org/10.1093/pm/pnz208>. PMID: 31498393. **Exclusion:** Ineligible population
231. Smith SL, Langen WH. A Systematic Review of Mindfulness Practices for Improving Outcomes in Chronic Low Back Pain. *International Journal of Yoga*. 2020 Sep-Dec;13(3):177-82. doi: https://dx.doi.org/10.4103/ijoy.IJOY_4_20. PMID: 33343146. **Exclusion:** Systematic review used as source document
232. Song HJ, Seo HJ, Kim D. Effectiveness of high-intensity laser therapy in the management of patients with knee osteoarthritis: A systematic review and meta-analysis of randomized controlled trials. *Journal of Back & Musculoskeletal Rehabilitation*. 2020;33(6):875-84. doi: <https://dx.doi.org/10.3233/BMR-191738>. PMID: 32831189. **Exclusion:** Systematic review used as source document
233. Sonmezer E, Ozkoslu MA, Yosmaoglu HB. The effects of clinical pilates exercises on functional disability, pain, quality of life and lumbopelvic stabilization in pregnant women with low back pain: A randomized controlled study. *Journal of Back &*

- Musculoskeletal Rehabilitation. 2021;34(1):69-76. doi: <https://dx.doi.org/10.3233/BMR-191810>. PMID: 32986655. **Exclusion:** Ineligible population
234. Stausholm MB, Naterstad IF, Joensen J, et al. Efficacy of low-level laser therapy on pain and disability in knee osteoarthritis: systematic review and meta-analysis of randomised placebo-controlled trials. *BMJ Open*. 2019 10 28;9(10):e031142. doi: <https://dx.doi.org/10.1136/bmjopen-2019-031142>. PMID: 31662383. **Exclusion:** Systematic review used as source document
235. Suman A, Schaafsma FG, van Dongen JM, et al. Effectiveness and cost-utility of a multifaceted eHealth strategy to improve back pain beliefs of patients with non-specific low back pain: a cluster randomised trial. *BMJ Open*. 2019 12 05;9(12):e030879. doi: <https://dx.doi.org/10.1136/bmjopen-2019-030879>. PMID: 31811006. **Exclusion:** Ineligible population
236. Teychenne M, Lamb KE, Main L, et al. General strength and conditioning versus motor control with manual therapy for improving depressive symptoms in chronic low back pain: A randomised feasibility trial. *PLoS ONE* [Electronic Resource]. 2019;14(8):e0220442. doi: <https://dx.doi.org/10.1371/journal.pone.0220442>. PMID: 31369613. **Exclusion:** Ineligible population
237. Thompson AR, Christopherson Z, Marshall LM, et al. A Pilot Randomized Controlled Trial for Aerobic and Strengthening Exercises on Physical Function and Pain for Hip Osteoarthritis. *Pm & R*. 2020 03;12(3):229-37. doi: <https://dx.doi.org/10.1002/pmrj.12262>. PMID: 31600429. **Exclusion:** Inadequate duration of followup
238. Tonye-Geoffroy L, Mauboussin Carlos S, Tuffet S, et al. Efficacy of a combination of hypnosis and transcutaneous electrical nerve stimulation for chronic non-cancer pain: A randomized controlled trial. *Journal of Advanced Nursing*. 2021 Jun;77(6):2875-86. doi: <https://dx.doi.org/10.1111/jan.14833>. PMID: 33783846. **Exclusion:** Ineligible population
239. Tse M, Li Y, Tang SK, et al. An Exploration of the Effectiveness of a Peer-Led Pain Management Program (PAP) for Nursing Home Residents with Chronic Pain and an Evaluation of Their Experiences: A Pilot Randomized Controlled Trial. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2020 06 08;17(11):08. doi: <https://dx.doi.org/10.3390/ijerph17114090>. PMID: 32521785. **Exclusion:** Ineligible population
240. Tsuboi Y, Oka T, Nakatsuka K, et al. Effectiveness of workplace active rest programme on low back pain in office workers: a stepped-wedge cluster randomised controlled trial. *BMJ Open*. 2021 06 25;11(6):e040101. doi: <https://dx.doi.org/10.1136/bmjopen-2020-040101>. PMID: 34172540. **Exclusion:** Ineligible study design
241. Tu JF, Wang LQ, Shi GX, et al. Effect of acupuncture on knee injury and osteoarthritis outcome score in patients with knee osteoarthritis. *Zhongguo zhen jiu* [Chinese acupuncture & moxibustion]. 2021;41(1):27-30. PMID: CN-02232691 NEW. **Exclusion:** Not English language but possibly relevant
242. Tu JF, Wang LQ, Shi GX, et al. [Effect of acupuncture on knee injury and osteoarthritis outcome score in patients with knee osteoarthritis]. *Zhongguo Zhenjiu*. 2021 Jan 12;41(1):27-30. doi: <https://dx.doi.org/10.13703/j.0255-2930.20191212-0001>. PMID: 33559438. **Exclusion:** Not English language but possibly relevant
243. Turner MN, Hernandez DO, Cade W, et al. The Role of Resistance Training Dosing on Pain and Physical Function in Individuals With Knee Osteoarthritis: A Systematic Review. *Sports & Health*. 2020 Mar/Apr;12(2):200-6. doi: <https://dx.doi.org/10.1177/1941738119887183>. PMID: 31850826. **Exclusion:** Systematic review used as source document
244. Udina-Cortes C, Fernandez-Carnero J, Romano AA, et al. Effects of neuro-adaptive electrostimulation therapy on pain and disability in fibromyalgia: A prospective, randomized, double-blind study. *Medicine*. 2020 Dec 18;99(51):e23785. doi: <https://dx.doi.org/10.1097/MD.0000000000>


- [023785](#). PMID: 33371148. **Exclusion:** Ineligible intervention
245. Uebelacker LA, Van Noppen D, Tremont G, et al. A pilot study assessing acceptability and feasibility of hatha yoga for chronic pain in people receiving opioid agonist therapy for opioid use disorder. *Journal of Substance Abuse Treatment*. 2019 10;105:19-27. doi: <https://dx.doi.org/10.1016/j.jsat.2019.07.015>. PMID: 31443887. **Exclusion:** Ineligible population
246. Vaegter HB, Ussing K, Johansen JV, et al. Improvements in clinical pain and experimental pain sensitivity after cognitive functional therapy in patients with severe persistent low back pain. *The Pain Report*. 2020 Jan-Feb;5(1):e802. doi: <https://dx.doi.org/10.1097/PR9.00000000000000802>. PMID: 32072097. **Exclusion:** Ineligible study design
247. Valera-Calero A, Lluch Girbes E, Gallego-Izquierdo T, et al. Endocrine response after cervical manipulation and mobilization in people with chronic mechanical neck pain: a randomized controlled trial. *European journal of physical and rehabilitation medicine*. 2019;55(6):792-805. PMID: CN-01690485. **Exclusion:** Inadequate duration of followup
248. Van Dyke BP, Newman AK, Morais CA, et al. Heterogeneity of Treatment Effects in a Randomized Trial of Literacy-Adapted Group Cognitive-Behavioral Therapy, Pain Psychoeducation, and Usual Medical Care for Multiply Disadvantaged Patients With Chronic Pain. *Journal of Pain*. 2019 10;20(10):1236-48. doi: <https://dx.doi.org/10.1016/j.jpain.2019.04.006>. PMID: 31022555. **Exclusion:** Ineligible population
249. Veronese N, Smith L, Bolzetta F, et al. Efficacy of conservative treatments for hand osteoarthritis : An umbrella review of interventional studies. *Wiener Klinische Wochenschrift*. 2021 Mar;133(5-6):234-40. doi: <https://dx.doi.org/10.1007/s00508-020-01702-0>. PMID: 32607645. **Exclusion:** Systematic review used as source document
250. Villafaina S, Collado-Mateo D, Dominguez-Munoz FJ, et al. Effects of exergames on heart rate variability of women with fibromyalgia: A randomized controlled trial. *Scientific Reports*. 2020 03 20;10(1):5168. doi: <https://dx.doi.org/10.1038/s41598-020-61617-8>. PMID: 32198423. **Exclusion:** Ineligible outcomes
251. Vining R, Long CR, Minkalis A, et al. Effects of Chiropractic Care on Strength, Balance, and Endurance in Active-Duty U.S. Military Personnel with Low Back Pain: A Randomized Controlled Trial. *Journal of Alternative & Complementary Medicine*. 2020 Jul;26(7):592-601. doi: <https://dx.doi.org/10.1089/acm.2020.0107>. PMID: 32543211. **Exclusion:** Inadequate duration of followup
252. Walsh N, Jones L, Phillips S, et al. Facilitating Activity and Self-management for people with Arthritic knee, hip or lower back pain (FASA): A cluster randomised controlled trial. *Musculoskeletal Science & Practice*. 2020 12;50:102271. doi: <https://dx.doi.org/10.1016/j.msksp.2020.102271>. PMID: 33068901. **Exclusion:** Ineligible intervention
253. Wang F, Zhang X, Tong X, et al. The effects on pain, physical function, and quality of life of quadriceps strengthening exercises combined with Baduanjin qigong in older adults with knee osteoarthritis: a quasi-experimental study. *BMC Musculoskeletal Disorders*. 2021 Mar 29;22(1):313. doi: <https://dx.doi.org/10.1186/s12891-021-04179-8>. PMID: 33781238. **Exclusion:** Ineligible study design
254. Williams ACC, Fisher E, Hearn L, et al. Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane Database of Systematic Reviews*. 2020 08 12;8:CD007407. doi: <https://dx.doi.org/10.1002/14651858.CD007407.pub4>. PMID: 32794606. **Exclusion:** Systematic review used as source document
255. Wippert PM, Drieslein D, Beck H, et al. The Feasibility and Effectiveness of a New Practical Multidisciplinary Treatment for Low-Back Pain: A Randomized Controlled Trial. *Journal of Clinical Medicine*. 2019 Dec 31;9(1):31. doi: <https://dx.doi.org/10.3390/jcm9010115>. PMID: 31906224. **Exclusion:** Ineligible population
256. Wu YL, Fang SC, Chen SC, et al. Effects of Neurofeedback on Fibromyalgia: A Randomized Controlled Trial. *Pain*

- Management Nursing. 2021 Feb 09;09:09. doi: <https://dx.doi.org/10.1016/j.pmn.2021.01.004>. PMID: 33579615. **Exclusion:** Inadequate duration of followup
257. Xiao C, Zhuang Y, Kang Y. Effects of Wu Qin xi Qigong exercise on physical functioning in elderly people with knee osteoarthritis: A randomized controlled trial. *Geriatrics & gerontology international*. 2020 Oct;20(10):899-903. doi: <https://dx.doi.org/10.1111/ggi.14007>. PMID: 32886828. **Exclusion:** Inadequate duration of followup
258. Xie SH, Wang Q, Wang LQ, et al. Effect of Internet-Based Rehabilitation Programs on Improvement of Pain and Physical Function in Patients with Knee Osteoarthritis: Systematic Review and Meta-analysis of Randomized Controlled Trials. *Journal of Medical Internet Research*. 2021 01 05;23(1):e21542. doi: <https://dx.doi.org/10.2196/21542>. PMID: 33399542. **Exclusion:** Systematic review used as source document
259. Yang Q, Yu S, Wang J, et al. Effects of Baduanjin on patients with chronic nonspecific low back pain: A randomized controlled trial. *Medicine*. 2021 Jan 29;100(4):e24448. doi: <https://dx.doi.org/10.1097/MD.00000000000024448>. PMID: 33530252. **Exclusion:** Not a study
260. Yao C, Li Z, Zhang S, et al. Effects of Wuqinxi in the Patients with Chronic Low Back Pain: A Randomized Controlled Trial. *Evidence-Based Complementary & Alternative Medicine: eCAM*. 2020;2020:1428246. doi: <https://dx.doi.org/10.1155/2020/1428246>. PMID: 32908554. **Exclusion:** Ineligible comparator
261. Ye J, Zheng Q, Zou L, et al. Mindful Exercise (Baduanjin) as an Adjuvant Treatment for Older Adults (60 Years Old and Over) of Knee Osteoarthritis: A Randomized Controlled Trial. *Evidence-Based Complementary & Alternative Medicine: eCAM*. 2020;2020:9869161. doi: <https://dx.doi.org/10.1155/2020/9869161>. PMID: 32617115. **Exclusion:** Inadequate duration of followup
262. Yeh SW, Hong CH, Shih MC, et al. Low-Level Laser Therapy for Fibromyalgia: A Systematic Review and Meta-Analysis. *Pain Physician*. 2019 05;22(3):241-54. PMID: 31151332. **Exclusion:** Systematic review used as source document
263. Yoo SA, Kim CY, Kim HD, et al. Effects of progressive muscle relaxation therapy with home exercise on pain, fatigue, and stress in subjects with fibromyalgia syndrome: A pilot randomized controlled trial. *Journal of Back & Musculoskeletal Rehabilitation*. 2021 Jun 15;15:15. doi: <https://dx.doi.org/10.3233/BMR-191703>. PMID: 34151818. **Exclusion:** Inadequate duration of followup
264. You Y, Liu J, Tang M, et al. Effects of Tai Chi exercise on improving walking function and posture control in elderly patients with knee osteoarthritis: A systematic review and meta-analysis. *Medicine*. 2021 Apr 23;100(16):e25655. doi: <https://dx.doi.org/10.1097/MD.00000000000025655>. PMID: 33879749. **Exclusion:** Systematic review used as source document
265. Zaworski K, Latosiewicz R. The effectiveness of manual therapy and proprioceptive neuromuscular facilitation compared to kinesiotherapy: a four-arm randomized controlled trial. *European journal of physical & rehabilitation medicine*. 2021 Apr;57(2):280-7. doi: <https://dx.doi.org/10.23736/S1973-9087.21.06344-9>. PMID: 33650840. **Exclusion:** Ineligible population
266. Zhang C, Li Y, Zhong Y, et al. Effectiveness of motor control exercise on non-specific chronic low back pain, disability and core muscle morphological characteristics: A meta-analysis of randomized controlled trials. *European journal of physical & rehabilitation medicine*. 2021 May 07;07:07. doi: <https://dx.doi.org/10.23736/S1973-9087.21.06555-2>. PMID: 33960180. **Exclusion:** Systematic review used as source document
267. Zhang L, Yuan H, Zhang L, et al. Effect of acupuncture therapies combined with usual medical care on knee osteoarthritis. *Journal of traditional chinese medicine = chung i tsa chih ying wen pan*. 2019;39(1):103-10. PMID: CN-02133212 NEW. **Exclusion:** Inadequate duration of followup

268. Zhang Q, Fang J, Chen L, et al. Different kinds of acupuncture treatments for knee osteoarthritis: a multicentre, randomized controlled trial. *Trials* [Electronic Resource]. 2020 Mar 14;21(1):264. doi: <https://dx.doi.org/10.1186/s13063-019-4034-8>. PMID: 32171318. **Exclusion:** Not a study
269. Zhao L, Cheng K, Wu F, et al. Effect of Laser Moxibustion for Knee Osteoarthritis: A Multisite, Double-blind Randomized Controlled Trial. *Journal of Rheumatology*. 2021 Jun;48(6):924-32. doi: <https://dx.doi.org/10.3899/jrheum.200217>. PMID: 32611673. **Exclusion:** Ineligible intervention
270. Zheng Z, Gibson S, Helme RD, et al. Effects of Electroacupuncture on Opioid Consumption in Patients with Chronic Musculoskeletal Pain: a Multicenter Randomized Controlled Trial. *Pain medicine* (Malden, Mass.). 2019;20(2):397-410. PMID: CN-01943577. **Exclusion:** Ineligible population
271. Zhu F, Zhang M, Wang D, et al. Yoga compared to non-exercise or physical therapy exercise on pain, disability, and quality of life for patients with chronic low back pain: A systematic review and meta-analysis of randomized controlled trials. *PLoS ONE* [Electronic Resource]. 2020;15(9):e0238544. doi: <https://dx.doi.org/10.1371/journal.pone.0238544>. PMID: 32870936. **Exclusion:** Systematic review used as source document

Appendix G. Updated or New Meta-analyses

Figure G-1. Exercise versus usual care, an attention control, or a placebo intervention for chronic low back pain: effects on function, excluding outlier trial^a




AC = attention control; AE = aerobic exercise (walking); CI = confidence interval; CPGS –BD = Von Korff Chronic Pain Grade Score Back Disability; DP = directional preference; GE= general exercise; MC = motor control; MF = mobility/flexibility; MI = minimal intervention; N = number; NE = no exercise; NM = neuromuscular re-education; ODI = Oswestry Disability Index; PDI = Pain Disability Index; RDQ = Roland-Morris Disability Questionnaire; SD = standard deviation; SMD = standardized mean difference; Spilates = selective Pilates; Strng=Strength training; UC = usual care; WL = waitlist

^a Areeudomwong 2017, included in prior report.

^b New trials: Shariat 2019, Lang 2021

Figure G-2. Psychological therapy versus usual care or an attention control for chronic low back pain: effects on function, excluding outlier trial^a




AC = attention control; CB = cognitive-behavioral therapy; CI = confidence interval; MRDQ = Modified Roland-Morris Disability Questionnaire; N = number; ODI = Oswestry Disability Index; PI = placebo intervention; PRT = pain reprocessing therapy; RDQ = Roland-Morris Disability Questionnaire; RPT = respondent therapy (progressive relaxation); SD = standard deviation; SMD = standardized mean difference; UC = usual care

^a Shariat 2019, new trial.

^b New trial: Ashar 2021


Figure G-3. Spinal manipulation versus sham manipulation, usual care, an attention control, or a placebo intervention for chronic low back pain: effects on function



AC = attention control; CI = confidence interval; N = number; ODI = Oswestry Disability Index; PI = placebo intervention; RDQ = Roland-Morris Disability Questionnaire; SD = standard deviation; SMD = standardized mean difference; SP= sham manipulation; UC = usual care; UK BEAM = UK Back pain exercise and manipulation trial; VF = Von Korff functional disability

^b New trial: Thomas 2020

Figure G-4. Exercise versus usual care, no treatment, sham, or an attention control for osteoarthritis knee pain: effects on function, excluding outlier trial^a




AC = attention control; APC = Arthritis Impact Measurement Scale (AIMS) physical activity component; CI = confidence interval; COM = combination exercise therapy; KADL = Knee Injury and Osteoarthritis Outcome Score (KOOS) ADL subscore; LI = Lequesne Index; LLFDI = Late Life Function and Disability Index Basic Lower Limb Function Score; ME = mobility exercise; MP = muscle performance exercise; NR = neuromuscular reeducation exercise; NT = no treatment; OKS = Oxford Knee Score; SD = standard deviation; SMD = standardized mean difference; UC = usual care; WOMAC = Western Ontario and McMaster's Universities Osteoarthritis Index

^a Dias 2003, included in prior report.

^b New trials: Messier 2021, Munukka 2020 and Rewald 2019.


Figure G-5. Exercise versus usual care, no treatment, sham, or an attention control for osteoarthritis knee pain: effects on pain



AC = attention control; CI = confidence interval; COM = combination exercise therapy; ME = mobility exercise; MP = muscle performance exercise; NR = neuromuscular re-education exercise; NT = no treatment; SD = standard deviation; SMD = standardized mean difference; UC = usual care

^a New trials: Messier 2021, Munukka 2020 and Rewald 2019.


Figure G-6. Low-level laser therapy versus usual care or sham for osteoarthritis knee pain: effects on function [NEW]



CI = confidence interval; SD = standard deviation; UC = usual care
^a New trial: Alqualio-Costa 2021

DRAFT

Figure G-7. Low-level laser therapy versus usual care or sham for osteoarthritis knee pain: effects on pain



CI = confidence interval; SD = standard deviation; UC = usual care

^a New trial: Alqualio-Costa 2021

DRAFT