

Effective Health Care

Interventions for Substance Use Disorders in Adolescents

Results of Topic Selection Process & Next Steps

The nominator, American Academy of Child & Adolescent Psychiatry (AACAP), is interested in a new systematic review on the treatment of substance abuse disorders in children and adolescents to inform an update of their practice parameter.

The topic fulfills the Effective Health Care Program selection criteria. This topic, Substance Use Disorders in Adolescents, will go forward for refinement as a systematic review. The scope of this topic, including populations, interventions, comparators, and outcomes, will be further developed in the systematic review.

Topic Brief

Topic #: 0676

Nomination Date: 06/02/2016

Nominator: American Academy of Child & Adolescent Psychiatry

Date: August 2016, January 2018

Authors:

Kara Winchell Karli Kondo Rose Relevo Mark Helfand Christine Chang (update)

Conflict of Interest: None of the investigators have any affiliations or financial involvement that conflicts with the material presented in this report

Summary of Key Findings:

- Appropriateness and importance: The nomination is both appropriate and important.
- <u>Duplication:</u> Three completed systematic reviews were identified related to KQ 1. All three pertain to interventions for substance use disorders. Two of the systematic reviews examine motivational interviewing as a treatment for substance use disorders, and the third included two trials of disulfiram for the treatment of alcohol dependency in adolescents. Our search did not find any evidence reviews which met inclusion criteria for KQ 2, examining subgroup variation in the treatment of substance use disorders.
 - <u>Duplication, updated search:</u> The updated duplication search identified 14 additional systematic reviews. These reviews do not cover the scope of the nomination, such as types of substance abuse, interventions, and subgroups. In addition seven of the reviews have searches that are not sufficiently recent to likely be useful to the nominator. Depending on the results of the in-process systematic reviews on brief interventions the scope may be further narrowed.

Feasibility:

- Size/scope of review: We estimate that the total size of the relevant literature (2011-present) may be approximately 64 studies across key questions (low confidence). The estimate is not firm because all titles and abstracts were not reviewed for this document. In addition during scoping of the review issues and concerns from other perspectives, including those of health systems may broaden the review scope.
- o ClinicalTrials.gov: We identified 7 relevant trials on ClinicalTrials.gov.
- Cochrane RCT filter results: We identified 14 additional RCTs covering interventions for SUDs, as well as interventions for subgroups (including comorbidities and ethnic minorities).
- Impact: The nomination has high impact potential if a new systematic review provides new evidence where it was previously lacking from the AACAP's 2005 practice parameters.
- <u>Value</u>: The potential for value is high, given that AACAP will use a new AHRQ systematic review to update their clinical practice guideline on a topic that is of concern to their constituency. Given the prevalence and challenges in treating substance abuse disorders in adolescents, a review on this topic may be of value to other stakeholder groups such as health systems.

Table of Contents

Introduction	4
Methods	5
Appropriateness and Importance	5
Desirability of New Review/Duplication	5
Impact of a New Evidence Review	5
Feasibility of New Evidence Review	5
Value	5
Compilation of Findings	5
Results	6
Appropriateness and Importance	6
Desirability of New Review/Duplication	6
Impact of a New Evidence Review	6
Feasibility of a New Evidence Review	6
Value	7
Summary of Findings	7
References	8
Appendix A. Selection Criteria Summary	A-1
Appendix B. Search for Systematic Reviews (Duplication)	B-1
Appendix C. Search Strategy & Results (Feasibility)	C-1

Introduction

There is a rich literature examining optimal treatment programs for adults with various substance use disorders, but the available evidence for adolescents and children is less clear. As adults and adolescents should be treated differently, there is a need for a comprehensive treatment guide for substance use disorders in adolescents. The National Institute on Drug Abuse (NIDA) assert that approximately 15.2% of children and adolescents who begin drinking alcohol by age 14 will eventually develop alcohol dependence, and 25% of those who begin abusing prescription drugs before or at the age of 13 will develop a substance use disorder at some time in their lives. The NIDA report asserts that adolescent substance use needs to be identified as soon as possible and treatment should address larger life needs related to medical, social and psychological well-being, not just drug use. Treatment should also address cooccurring mental health conditions. A support system consisting of family, friends, community, and health providers is an important factor in treating substance use disorders. The NIDA report includes a section entitled "Evidence-based Approaches to Treating Adolescent Substance Use Disorders" that lists 5 behavioral treatments, 5 family-based approaches, various medications, and 4 recovery support services that have been "shown to be effective."

This nomination was received on 06/02/2016. The nominator is interested in using a rigorous systematic review process to inform the update of 2005 American Academy of Child & Adolescent Psychiatry (AACAP) practice parameters² pertaining to the interventions for substance use disorders in adolescents. They have specifically requested that we exclude tobacco use from the scope. They are interested in the range of interventions, including brief interventions.

The questions for this nomination are:

Key Question 1. What are the benefits and harms and comparative benefits and harms of interventions for substance use disorders (generally and by substance specific) in adolescents?

Key Question 2. Do the benefits and harms vary by subpopulation (e.g., age, gender, race/ethnicity, comorbidity)?

To define the inclusion criteria for the key questions we specify the population, interventions, comparators, and outcomes (PICO) of interest. See Table 1.

Table 1. Key Questions and PICOs

Key Questions	1. What are the (a) benefits and harms and (b) comparative benefits and harms of interventions for substance use disorders (generally and by substance specific) in adolescents?	2. Do the benefits and harms vary by subpopulation (e.g., age, gender, race/ethnicity, comorbidity)?
Population	Adolescents with substance use disorder (e.g., alcohol use disorder, cannabis use disorder, opioid use disorder)	Adolescents with substance use disorder (e.g., alcohol use disorder, cannabis use disorder, opioid use disorder)
Interventions	Non-pharmacological and pharmacologic interventions	Non-pharmacological and pharmacologic interventions
Comparators	Combination of treatments, placebo, no treatment	Combination of treatments, placebo, no treatment
Outcomes	Lower rates of substance use, adverse events, quality of life	Lower rates of substance use, adverse events, quality of life

Methods

To assess this topic nomination for priority for a systematic review or other AHRQ EHC report, we used a modified process based on established criteria. Our assessment is hierarchical in nature, with the findings of our assessment determining the need for further evaluation. Details related to our assessment are provided in Appendix A.

- 1. Determine the appropriateness of the nominated topic for inclusion in the EHC program.
- 2. Establish the overall *importance* of a potential topic as representing a health or healthcare issue in the United States.
- 3. Determine the *desirability of new evidence review* by examining whether a new systematic review or other AHRQ product would be duplicative.
- 4. Assess the potential impact a new systematic review or other AHRQ product.
- 5. Assess whether the *current state of the evidence* allows for a systematic review or other AHRQ product (feasibility).
- 6. Determine the potential value of a new systematic review or other AHRQ product.

Appropriateness and Importance

We assessed the nomination for appropriateness and importance (see Appendix A).

Desirability of New Review/Duplication

We searched for high-quality, completed or in-process evidence reviews pertaining to the key questions of the nomination. Table 2 includes the citations for the reviews that were determined to address the key questions. Appendix B includes the list of the sources searched and potentially relevant titles identified by our research librarian.

The search was updated on 1/29/2018. See Appendix B for sources searched.

Impact of a New Evidence Review

The impact of a new evidence review was assessed by analyzing the current standard of care, the existence of potential knowledge gaps, and practice variation. We considered whether it was hypothetically possible for this review to influence the current state of practice through various dissemination pathways (practice recommendation, clinical guidelines, etc.).

Feasibility of New Evidence Review

We conducted a literature search in PubMed and PsycInfo from June 2011 and June 2016.

Because a large number of articles were identified, we reviewed a random sample of 200 titles and abstracts for inclusion and classified identified studies by study design, to assess the size and scope of a potential evidence review. We then calculated the projected total number of included studies based on the proportion of studies included from the random sample. See Table 2, Feasibility Column, Size/Scope of Review Section for the citations of included studies. See Appendix C for the PubMed (and PsycInfo) search strategy and links to the ClinicalTrials.gov search.

Value

We assessed the nomination for value (see Appendix A). We considered whether or not the topic would inform clinical policy in community and/or clinical settings, and if there was a partner organization that would use this evidence review to do disseminate this policy.

Compilation of Findings

We constructed a table outlining the selection criteria as they pertain to this nomination (see Appendix A).

Results

Appropriateness and Importance

This is an appropriate and important topic. In a nationally representative samples of US youth the lifetime prevalence of alcohol use disorders is approximately 8% and illicit drug use disorders is 2-3%.

Desirability of New Review/Duplication

A new evidence review would not be duplicative of an existing product. Three completed systematic reviews were identified related to KQ 1. All three pertain to interventions for substance use disorders. Two of the systematic reviews examine motivational interviewing as a treatment for substance use disorders, and the third included two trials of disulfiram for the treatment of alcohol dependency in adolescents. Our search did not find any evidence reviews which met inclusion criteria for KQ 2, examining subgroup variation in the treatment of substance use disorders. See Table 2, Duplication column for the systematic review citations that were determined to address the key questions.

The updated duplication search identified 14 additional completed and in-process systematic reviews. All but one pertain to KQ 1. Five focused on non-opioid treatment; one on alcohol and cannabis; one on cannabis; two on alcohol; and four on substance abuse treatment broadly. All focused on non-pharmacologic treatments, including CBT, motivational interviewing, types of family therapy, computerized interventions, and brief interventions.

Despite these reviews, a new evidence review would not be duplicative.

- End dates of searches for seven of the reviews were not sufficiently recent, ranging from 2011-2014.
- The remaining reviews do not cover the scope of the nomination including pharmacologic interventions, treatment of opioid abuse, and most subgroups of interest.
 - Three reviews cover single interventions: motivational interviewing, music therapy, and computerized brief interventions.
 - Three reviews are in-process and all focus on brief interventions. These could partly duplicate a part of this nomination.

See Appendix D for the references for the additional systematic reviews.

Impact of a New Evidence Review

A new systematic review may have high impact, as it will inform a practice through a practice parameters and potentially fill gaps in evidence from the previous practice parameter.

Feasibility of a New Evidence Review

A new evidence review examining is feasible. Our PubMed feasibility search identified 1,253 unique articles published between June 2011 and June 2016. See Table 2, Feasibility column for the citations that were determined to address the key questions. We estimate that the total size of the relevant literature (2011-present) may be approximately 64 studies across key questions. We identified 14 additional RCTs covering interventions for SUDs, as well as interventions for subgroups (including comorbidities and ethnic minorities) in the Cochrane RCT filter. We identified 7 relevant trials on ClinicalTrials.gov.

Table 2. Nomination key questions, with the identified evidence reviews and original research

Table 2. Nomination key questions, with the identified evidence reviews and original research		
Key Question	Duplication (Completed or In-	
	Process Evidence Reviews)	Ongoing Research)
KQ 1: Benefits and harms and comparative benefits and harms of interventions for SUD in adolescents	Total number of completed and in-progress systematic reviews: 16	Size/scope of review Relevant Studies Identified: 3 RCT – 2 ^{6,7} Retrospective Cohort - 1 ⁸ Projected Total: 24
		ClinicalTrials.gov Relevant Trials: 7 Recruiting – 4 ⁹⁻¹² Active, not recruiting – 1 ¹³ Completed - 2 ^{14,15}
KQ 2: Variation by Subpopulation	Total number of completed and in-progress systematic reviews: 1	Size/scope of review Relevant Studies Identified: 1 Retrospective Cohort 18 Projected Total: 8 ClinicalTrials.gov Relevant Trials: 3 Recruiting - 1 ¹² Completed - 2 ^{14,15}

Abbreviations: KQ=Key Question; RCT=Randomized Controlled Trial; SUD=Substance Use Disorder

Value

The potential for value is high given the given that AACAP will use a new AHRQ systematic review to update their clinical practice guideline. This organization has previously produced high quality, evidence based guidelines, and are transparent about their methodology.

Summary of Findings

- Appropriateness and importance: The nomination is both appropriate and important.
- <u>Duplication:</u> We identified 17 completed and in-process reviews. However these
 did not cover the breadth of types of substance abuse, interventions and
 subgroups of interest. In addition the searches for seven of the reviews were not
 sufficiently recent to likely be useful to the nominator. Depending on the results of
 the in-process systematic reviews on brief interventions the scope may be further
 narrowed.
- Feasibility:
 - Size/scope of review: We estimate that the total size of the relevant literature (2011-present) may be approximately 64 studies across key questions (low confidence). The estimate is not firm because all titles and abstracts were not reviewed for this document. In addition during scoping of the review issues and concerns from other perspectives, including those of health systems may broaden the review scope.
 - o ClinicalTrials.gov: We identified 7 relevant trials on ClinicalTrials.gov.
 - Cochrane RCT filter results: We identified 14 additional RCTs covering interventions for SUDs, as well as interventions for subgroups (including comorbidities and ethnic minorities).
- <u>Impact:</u> The nomination has high impact potential if a new systematic review provides new evidence where it was previously lacking from the AACAP's 2005 practice parameters.

<u>Value</u>: The potential for value is high, given that AACAP will use a new AHRQ systematic review to update their clinical practice guideline on a topic of concern to their constituency. Given the prevalence of substance abuse disorders in adolescents, a review on this topic may be of value to other stakeholder groups such as health systems.

References

- 1. National Institute on Drug Abuse. Principles of Adolescent Substance Use Disorder Treatment: A Research-Based Guide. In: Abuse NIoD, ed: National Institutes of Health; 2014.
- 2. Bukstein OG, Bernet W, Arnold V, et al. Practice parameter for the assessment and treatment of children and adolescents with substance use disorders. *J Am Acad Child Adolesc Psychiatry*. 2005;44(6):609-621.
- 3. Barnett E, Sussman S, Smith C, Rohrbach LA, Spruijt-Metz D. Motivational Interviewing for adolescent substance use: a review of the literature. *Addict Behav.* 2012;37(12):1325-1334.
- 4. Jensen CD, Cushing CC, Aylward BS, Craig JT, Sorell DM, Steele RG. Effectiveness of motivational interviewing interventions for adolescent substance use behavior change: a meta-analytic review. *J Consult Clin Psychol*. 2011;79(4):433-440.
- 5. Skinner MD, Lahmek P, Pham H, Aubin HJ. Disulfiram efficacy in the treatment of alcohol dependence: a meta-analysis. *PLoS One*. 2014;9(2):e87366.
- 6. Brown PC, Budney AJ, Thostenson JD, Stanger C. Initiation of abstinence in adolescents treated for marijuana use disorders. *J Subst Abuse Treat*. 2013;44(4):384-390.
- 7. Robbins MS, Feaster DJ, Horigian VE, et al. Brief strategic family therapy versus treatment as usual: results of a multisite randomized trial for substance using adolescents. *J Consult Clin Psychol.* 2011;79(6):713-727.
- 8. Becker SJ, Stein GL, Curry JF, Hersh J. Ethnic differences among substance-abusing adolescents in a treatment dissemination project. *J Subst Abuse Treat*. 2012;42(3):328-336.
- 9. Medical University of South Carolina. Integrative Risk Reduction and Treatment for Teen Substance Use Problems and PTSD (RRFT). *ClinicalTrials.gov*. 2015:NCT01751035.
- 10. Dartmouth-Hitchcock Medical Center. Behavioral Treatment of Adolescent Substance Use (SMART). *ClinicalTrials.gov.* 2015;NCT02063984.
- 11. Oregon Research Institute. Family and Adolescent Motivational Incentives for Leveraging Youth (FAMILY). *ClinicalTrials.gov.* 2016;NCT01736995.
- 12. Oregon Research Institute. Family Therapy Via Video Teleconference for Substance-Abusing Rural Adolescents (RAFT). *ClinicalTrials.gov.* 2016;NCT01751217.
- 13. University of Connecticut Health Center. Adaptive Treatment for Adolescent Cannabis Use Disorders. *ClinicalTrials.gov.* 2016;NCT01656707.
- 14. University of New Mexico. Assessing the Fit of Motivational Interviewing by Cultures With Adolescents (AMICA). *ClinicalTrials.gov*. 2015;NCT01948700.
- 15. University of Miami. Culturally Informed Family Based Treatment of Adolescents: A Randomized Trial (CIFFTA). *ClinicalTrials.gov*. 2015;NCT01823250.

Appendix A. Selection Criteria Summary

Selection Criteria	Supporting Data
1. Appropriateness	
1a. Does the nomination represent a health care drug, intervention, device, technology, or health care system/setting available (or soon to be available) in the U.S.?	Yes, this topic represents a health care drug and intervention available in the U.S.
1b. Is the nomination a request for a systematic review?	Yes, this topic is a request for a systematic review.
1c. Is the focus on effectiveness or comparative effectiveness?	The focus of this review is on both effectiveness and comparative effectiveness.
1d. Is the nomination focus supported by a logic model or biologic plausibility? Is it consistent or coherent with what is known about the topic?	Yes, it is biologically plausible. Yes, it is consistent with what is known about the topic.
2. Importance	
2a. Represents a significant disease burden; large proportion of the population	Yes, this topic represents a significant burden. AACAP states nationally representative samples of US youth reveal that the lifetime prevalence of alcohol use disorders is approximately 8% and illicit drug use disorders is 2-3%.
2b. Is of high public interest; affects health care decision making, outcomes, or costs for a large proportion of the US population or for a vulnerable population	Yes, this topic affects heath care decisions for a large, vulnerable population.
2c. Represents important uncertainty for decision makers	Yes, this topic represents important uncertainty for decision makers.
2d. Incorporates issues around both clinical benefits and potential clinical	Yes, this nomination addresses both benefits and potential harms of prevention interventions, pharmacological interventions, and non-pharmacological treatments for substance abuse disorders.
2e. Represents high costs due to common use, high unit costs, or high associated costs to consumers, to patients, to health care systems, or to payers	Yes this mental health diagnosis represents high cost due to the high rate of prevalence, and the increasing costs of treatment options.
Desirability of a New Evidence Review/Duplication	
3. Would not be redundant (i.e., the proposed topic is not already covered by available or soon-to-be available high-quality systematic review by AHRQ or others)	Three systematic reviews ³⁻⁵ were identified related to the key questions of the nomination—all three are complete and all three pertain to interventions for substance use disorders. Two of the systematic reviews ^{3,4} examine motivational interviewing as a treatment for substance use disorders. The third ⁵ examined disulfiram for the treatment of alcohol dependency; it included 2 trials in adolescents, one published in 2008 and the other in 2003. Our search did not find any evidence reviews examining subgroup variation in the treatment of substance use disorders (Key Question 2).

Selection Criteria	Supporting Data
4. Impact of a New Evidence Review	
4a. Is the standard of care unclear (guidelines not available or guidelines	Yes, the standard of care is unclear due to conflicting data and
inconsistent, indicating an information gap that may be addressed by a new	recommendations among authors and experts.
evidence review)?	
4b. Is there practice variation (guideline inconsistent with current practice,	Yes, there is practice variation due to conflicting data and existing
indicating a potential implementation gap and not best addressed by a new	recommendations.
evidence review)?	
5. Primary Research	
5. Effectively utilizes existing research and knowledge by considering:	Size/scope of review: We estimate that the total size of the relevant
- Adequacy (type and volume) of research for conducting a systematic	literature (2011-present) may be approximately 64 studies across key
review	questions (low confidence).
- Newly available evidence (particularly for updates or new technologies)	Official Trials and Mark Lough and Calabara Official Trials and
	ClinicalTrials.gov: We identified 7 relevant trials on ClinicalTrials.gov.
	Coobrana PCT filter regults: We identified 14 additional PCTs severing
	Cochrane RCT filter results: We identified 14 additional RCTs covering
	interventions for SUDs, as well as interventions for subgroups (including comorbidities and ethnic minorities).
6. Value	comorbidities and etimic minorities).
6a. The proposed topic exists within a clinical, consumer, or policy-making	Yes, this topic will inform clinical decision-making on treating substance
context that is amenable to evidence-based change	abuse disorders among children and adolescents across community and
Context that is amenable to evidence based change	clinical settings.
6b. Identified partner who will use the systematic review to influence	Yes, AACAP will use a systematic review to update their 2005 clinical
practice (such as a guideline or recommendation)	practice guidelines on the assessment and treatment of bipolar disorder
practice (cach as a galacinic of recommendation)	among children and adolescents.
Abbreviations, AACAD, American Academy of Child 9 Adolescent Development	

Abbreviations: AACAP= American Academy of Child & Adolescent Psychiatry; AHRQ=Agency for Healthcare and Research Quality; RCT=Randomized Controlled Trial

Appendix B. Search for Systematic Reviews (Duplication), June 2016 and January 2018

Listed below are the sources searched and results of our search for existing guidance. A research librarian conducted the search and selected potentially relevant evidence based on the key question in the nomination and the associated PICOTS. An investigator reviewed each of the links to evidence below for inclusion. The links below do not represent the evidence selected for inclusion (see main topic brief).

Substance Use Disorders in Adolescents	Search for Duplication: June 7, 2016
Source	Evidence
AHRQ and Other Federal Products	
AHRQ: Evidence reports and technology assessments, USPSTF recommendations, and related DEcIDE projects, and Horizon Scan	Technical Brief: Medication-Assisted Treatment Models of Care for Opioid Use Disorder in Primary Care Settings https://www.effectivehealthcare.ahrq.gov/ehc/products/636/2225/opioid-use-disorder-draft-report-160513.pdf https://www.effectivehealthcare.ahrq.gov/ehc/products/636/2225/opioid-use-disorder-draft-appendixes-160513.pdf
	Primary Care Behavioral Interventions to Prevent or Reduce Illicit Drug and Nonmedical Pharmaceutical Use in Children and Adolescents http://www.ncbi.nlm.nih.gov/books/NBK195068/
	Alcohol Misuse: Screening and Behavioral Counseling Interventions in Primary Care, May 2013 http://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/alcohol-misuse-screening-and-behavioral-counseling-interventions-in-primary-care
	Drug Use, Illicit: Primary Care Interventions for Children and Adolescents, March 2014 http://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/drug-use-illicit-primary-care-interventions-for-children-and-adolescents
	Drug Use, Illicit: Screening, January 2008 http://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/drug-use-illicit-screening
	From the Horizon Scanning Update: https://effectivehealthcare.ahrq.gov/ehc/products/393/880/AHRQ-Healthcare-Horizon-Scanning-Status-Update-150930.pdf Interactive text messaging program for prevention of hazardous alcohol use.
VA Products: PBM, and HSR&D (ESP) publications, and VA/DoD EBCPG Program	Nothing with adolescents
Cochrane Database and Others	
Cochrane Systematic Reviews and Protocols http://www.cochranelibrary.com/	Multisystemic Therapy for social, emotional, and behavioral problems in youth aged 10-17 http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD004797.pub4/full
	Primary prevention for alcohol misuse in young people http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD003024.pub2/full [Withdrawn from publication]

Substance Use Disorders in Adolescents	Search for Duplication: June 7, 2016
PubMed Health http://www.ncbi.nlm.nih.gov/pubmedhealth/	It is premature to draw conclusions about the effectiveness of MST compared with other services http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0013029/
	Mentoring adolescents to prevent drug or alcohol use http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0016418/
	Treatments for co-occurring depression and substance use in young people: a systematic review http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0076229/
	A review of 25 long-term adolescent tobacco and other drug use prevention program evaluations http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0020282/
	A systematic review of the effectiveness of brief interventions with substance using adolescents by type of drug http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0020268/
	The effects of interventions to prevent substance use among adolescents: a systematic review http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0052718/
	A systematic review of school-based alcohol and other drug prevention programs facilitated by computers or the Internet http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0058877/
	Effective family-based interventions for adolescents with substance use problems: a systematic review http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0022609/
	Drug prevention with vulnerable young people: a review http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0022417/
	Psychosocial and Developmental Alcohol Misuse Prevention in Schools can be effective http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0014879/
	Interventions delivered to young people in non-school settings for the prevention of drug use http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0013190/
	School-based prevention for illicit drug use http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0011759/
	Prevention of substance use in children/adolescents with mental disorders: a systematic review http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0052509/
	Alcohol-Use Disorders Diagnosis, Assessment and Management of Harmful Drinking and Alcohol Dependence http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0042164/
	Motivational interviewing for adolescent substance use: a review of the literature http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0050602/
	Adolescent substance abuse treatment: a synthesis of controlled evaluations http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0020812/
	Effectiveness of motivational interviewing interventions for adolescent substance use behavior change: a meta-analytic review http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0033193/
	A critical review of adolescent substance abuse group treatments http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0029002/

Substance Use Disorders in Adolescents	Search for Duplication: June 7, 2016	
HTA (CRD database): Health Technology	The effects of Positive Youth Development interventions on substance use, violence and inequalities: systematic review of theories of change, processes and	
Assessments	outcomes	
http://www.crd.york.ac.uk/crdweb/	http://www.crd.york.ac.uk/CRDWeb/ShowRecord.asp?AccessionNumber=32016000690&UserID=0	
	A systematic review of computerised serious educational games about alcohol and other drugs for adolescents	
	http://www.crd.york.ac.uk/CRDWeb/ShowRecord.asp?AccessionNumber=12014000596&UserID=0	
	A systematic review of effective interventions for reducing multiple health risk behaviors in adolescence http://www.ncbi.nlm.nih.gov/pubmed?term=24625172	
	http://www.hcbi.htm.nini.gov/pubmed.remi=24625172	
	Disulfiram efficacy in the treatment of alcohol dependence: a meta-analysis	
	http://www.ncbi.nlm.nih.gov/pubmed?term=24520330	
	Fifther than 1 and	
	Effectiveness of brief interventions as part of the Screening, Brief Intervention and Referral to Treatment (SBIRT) model for reducing the nonmedical use of psychoactive substances: a systematic review	
	http://www.ncbi.nlm.nih.gov/pubmed?term=24887418	
	Pravention des alkoholmissbrauchs von kindern, jugendlichen und jungen erwachsenen [Prevention of alcohol misuse among children, youths and young adults]	
	http://www.crd.york.ac.uk/CRDWeb/ShowRecord.asp?AccessionNumber=32011001381&UserID=0	
	Adolescent outpatient treatment and continuing care: main findings from a randomized clinical trial	
	http://www.crd.york.ac.uk/CRDWeb/ShowRecord.asp?AccessionNumber=22010001299&UserID=0	
	Community-based interventions to reduce substance misuse among vulnerable and disadvantaged children and young people http://www.crd.york.ac.uk/CRDWeb/ShowRecord.asp?AccessionNumber=32011000437&UserID=0	
PROSPERO Database (international prospective	A systematic review of interventions to reduce youth substance use through parent training: applications in the adolescent health arena	
register of systematic reviews and protocols)	http://www.crd.york.ac.uk/prospero/display_record.asp?/ID=CRD42014013069	
http://www.crd.york.ac.uk/prospero/		
	Systematic review of universal school-based resilience interventions targetisting adolescent tobacco, alcohol or illicit drug use: review protocol	
	http://www.crd.york.ac.uk/prospero/display_record.asp?ID=CRD42014004906	
	Modifiable parenting factors for reducing adolescent alcohol use: a systematic review and meta-analysis of longitudinal studies	
	Based on the literature, what parenting factors can be modified in order to delay adolescents' alcohol initiation and prevent their subsequent alcohol misuse?	
	http://www.crd.york.ac.uk/prospero/display_record.asp?ID=CRD42015016723	
CADTH (Canadian Agency for Drugs and	Prevention Strategies for Substance Misuse Harm Reduction A Review of Clinical Evidence	
Technologies in Health) https://www.cadth.ca/	https://www.cadth.ca/prevention-strategies-substance-misuse-harm-reduction-review-clinical-evidence	
intps.//www.cautii.ca/	Team-Based Behavioural Support for Youth Experiencing Mental Health or Addiction Issues: Clinical Effectiveness and Guidelines	
	https://www.cadth.ca/team-based-behavioural-support-youth-experiencing-mental-health-or-addiction-issues-clinical	

Substance Use Disorders in Adolescents	Search for Duplication: June 7, 2016
DoPHER (Database of promoting health effectiveness reviews) http://eppi.joe.ac.uk/webdatabases4/Intro.aspx?ID=9	Adolescent substance abuse: a review of the past 10 years. http://www.ncbi.nlm.nih.gov/pubmed/9519629
nttp://eppi.loe.ac.uk/weboatabases4/intro.aspx?ID=9	Adolescent substance abuse prevention programs: Theories, models, and research in the encouraging 80's. http://www.ncbi.nlm.nih.gov/pubmed/24258623
	Adolescent substance use: reviewing the effectiveness of prevention strategies. http://www.ncbi.nlm.nih.gov/pubmed/15281689
	Brief school-based interventions and behavioural outcomes for substance-using adolescents. http://www.ncbi.nlm.nih.gov/pubmed/26787125
	A comparison of family interventions to address adolescent risky behaviors: a literature review. http://www.ncbi.nlm.nih.gov/pubmed/23166186
	Etiology and prevention of substance use among Asian American youth. http://www.ncbi.nlm.nih.gov/pubmed/11519375
	Family-based preventive intervention: an approach to preventing substance use and antisocial behavior. http://www.ncbi.nlm.nih.gov/pubmed/10439843
	Family-strengthening approaches for the prevention of youth problem behaviors. http://www.ncbi.nlm.nih.gov/pubmed/12971192
	Independent Evaluation of Middle School-Based Drug Prevention Curricula: A Systematic Review. http://www.ncbi.nlm.nih.gov/pubmed/26367105
	Interventions for reducing adolescent alcohol abuse: a meta-analytic review. http://www.ncbi.nlm.nih.gov/pubmed/20048247
	Meta-analysis of motivational interviewing for adolescent health behavior: efficacy beyond substance use. http://www.ncbi.nlm.nih.gov/pubmed/24841861
	A meta-analysis of motivational interviewing interventions for pediatric health behavior change. http://www.ncbi.nlm.nih.gov/pubmed/24547922
	Preventing adolescent substance abuse through drug education. http://www.ncbi.nlm.nih.gov/pubmed/6419117
	Psychosocial approaches to substance abuse prevention: theoretical foundations and empirical findings. http://www.ncbi.nlm.nih.gov/pubmed/2661151
	The role of mass media in preventing adolescent substance abuse. http://www.ncbi.nlm.nih.gov/pubmed/6419119
	SBIRT for adolescent drug and alcohol use: current status and future directions. http://www.ncbi.nlm.nih.gov/pubmed/23352110
	Screening instruments for substance use and brief interventions targeting adolescents in primary care: a literature review. http://www.ncbi.nlm.nih.gov/pubmed/23454877
	Strengthening the role of two key institutions in the prevention of adolescent substance abuse. http://www.ncbi.nlm.nih.gov/pubmed/11755797
	Substance dependency treatment for adolescents: practice and research. http://www.ncbi.nlm.nih.gov/pubmed/11138716
	Systematic review of interventions focusing on Indigenous adolescent mental health and substance use. http://www.ncbi.nlm.nih.gov/pubmed/26425865
	A systematic review of the effectiveness of brief interventions with substance using adolescents by type of drug. http://www.ncbi.nlm.nih.gov/pubmed/15385228
ECRI institute https://www.ecri.org/Pages/default.aspx	None identified.

Substance Use Disorders in Adolescents	Search for Duplication: June 7, 2016
Secondary Sources checked on an as needed basis	
BlueCross BlueShield Foundation Massachusetts	None identified.
http://bluecrossfoundation.org/	
Campbell Collaboration	None identified.
http://www.campbellcollaboration.org/	
CMS Policies https://www.cms.gov/	None identified.
Hayes, inc.	None identified.
http://www.hayesinc.com/hayes/	
IOM (Institute of Medicine)	None identified.
http://iom.nationalacademies.org/	
McMaster Health System Evidence	None identified.
https://www.healthsystemsevidence.org/	
Robert Wood Johnson	None identified.
http://www.rwjf.org/	Mars ideatified
Systematic Reviews (Journal) : protocols and reviews	None identified.
http://systematicreviewsjournal.biomedcentral.com/	
UBC Centre for Health Services and Policy	None identified.
Research	Note definited.
http://chspr.ubc.ca/	
WHO Health Evidence Network	None identified.
http://www.euro.who.int/en/data-and-	
evidence/evidence-informed-policy-making/health-	
evidence-network-hen	
National Heart, Lung, and Blood Institute	None identified.
http://www.nhlbi.nih.gov/	
National Cancer Institute	None identified.
http://www.cancer.gov/	
CINAHL (EBSCO)	None identified.
PsycINFO (Ovid)	None identified.

Sources searched for updated duplication search, January 2018

Substance Use Disorders in Adolescents
Source

Search for Duplication: January 29, 2018

AHRQ: Evidence reports and technology assessments, USPSTF recommendations

Cochrane Systematic Reviews and Protocols

http://www.cochranelibrary.com/

PubMed Health

http://www.ncbi.nlm.nih.gov/pubmedhealth/

PROSPERO Database (international prospective register of systematic reviews and protocols) http://www.crd.york.ac.uk/prospero/

CADTH (Canadian Agency for Drugs and Technologies in Health)

https://www.cadth.ca/

DoPHER (Database of promoting health effectiveness reviews)

http://eppi.ioe.ac.uk/webdatabases4/Intro.aspx?ID=9

Campbell Collaboration

http://www.campbellcollaboration.org/

Systematic Reviews (Journal): protocols and reviews http://systematicreviewsjournal.biomedcentral.com/

Appendix C. Search Strategy & Results (Feasibility)

Topic: Substance Use Disorders in Children and Adolescents

Date: June 9, 2016

Database Searched: PubMed

Concept	Search String
Substance Use Disorder Treatment	(((((((treatment[Title] OR intervention[Title] OR therapy[Title] OR recovery[Title])) AND (dependence[Title] OR addiction[Title] OR habituation[Title] OR withdrawal[Title])) AND (substance[Title] OR drug[Title] OR drugs[Title] OR alcohol[Title] OR amphetamine[Title] OR cocaine[Title] OR inhalant[Title] OR marijuana[Title] OR opioid[Title] OR heroin[Title] OR morphine[Title])))) OR "Substance-Related Disorders/therapy"[Mesh]
Not Editorials, etc.	(((((("Letter"[Publication Type]) OR "News"[Publication Type]) OR "Patient Education Handout"[Publication Type]) OR "Comment"[Publication Type]) OR "Editorial"[Publication Type])) OR "Newspaper Article"[Publication Type]
Limit to Children, last 5 years, Human, English	Filters activated: published in the last 5 years, Humans, English, Child: birth-18 years.
N=2514	
Systematic Review N=166	PubMed subsection "Systematic [sb]"
Randomized Controlled Trials N=1095	Cochrane Sensitive Search Strategy for RCT's "((((((((((((((((((((((((((((((((((((
Other N=1253	

ClinicalTrials.gov

96 studies found for: Substance-Related Disorders | Child | received on or after 06/01/2011 https://clinicaltrials.gov/ct2/results?term=&recr=&rslt=&type=&cond=Substance-Related+Disorders&intr=&titles=&outc=&spons=&lead=&id=&state1=&cntry1=&state2=&cntry2=&state3=&cntry3=&locn=&gndr=&age=0&rcv_s=06%2F01%2F2011&rcv_e=&lup_s=&lup_e=

Appendix D. Additional Systematic Reviews identified in January 2018 search

KQ 1: Completed systematic reviews

- Filges T, Knudsen ASD, Svendsen MM, Kowalski K, Benjaminsen L, Jørgensen AMK. Cognitive-Behavioural Therapies for Young People in Outpatient Treatment for Non-Opioid Drug Use: A Systematic Review. Campbell Systematic Reviews 2015:3. 10.4073/csr.2015.3
- Filges T, Andersen D, Jørgensen, AMK. Functional Family Therapy (FFT) for Young People in Treatment for Non-opioid Drug Use: A Systematic Review. Campbell Systematic Reviews 2015:14 DOI: 10.4073/csr.2015.14
- Filges T, Rasmussen P, Andersen, D. Multidimensional Family Therapy (MDFT) for Young People in Treatment for Non-opioid Drug Abuse: A Systematic Review. Campbell Systematic Reviews 2015:8 10.4073/csr.2015.8
- Smedslund G, Wollcheid S, Fang L, Nilsen W, Steiro A, Larun L. Effects of early, computerized brief interventions on risky alcohol use and risky cannabis use among young people. Campbell Systematic Reviews 2017:6 DOI: 10.4073/csr.2017.6
- European Monitoring Centre for Drugs and Drug Addiction (2014), *Multidimensional family* therapy for adolescent drug users: a systematic review, EMCDDA Papers, Publications Office of the European Union, Luxembourg.
- Hohmann L, Bradt J, Stegemann T, Koelsch S (2017) Effects of music therapy and music-based interventions in the treatment of substance use disorders: A systematic review. PLoS ONE 12(11): e0187363. https://doi.org/10.1371/journal.pone.0187363
- Lindstrøm M, Saidj M, Kowalski K, Filges T, Rasmussen PS, Jørgensen A-MK. Brief Strategic Family Therapy (BSFT) for Young People in Treatment for Non-Opioid Drug Use: A Systematic Review. Campbell Systematic Reviews 2013:07 10.4073/csr.2013.07
- Lindstrøm M, Saidj M, Kowalski K, Filges T, Rasmussen P. S., Jørgensen A.K. Family Behavior Therapy (FBT) for young people in treatment for non-opioid drug use. Campbell Systematic Reviews 2015:9 10.4073/csr.2011.09
- Li, L., Zhu, S., Tse, N., Tse, S., and Wong, P. (2016) Effectiveness of motivational interviewing to reduce illicit drug use in adolescents: a systematic review and meta-analysis. Addiction, 111: 795–805. doi: 10.1111/add.13285.
- Tanner-Smith EE, Lipsey MW. Brief Alcohol Interventions for Adolescents and Young Adults: A
 Systematic Review and Meta-analysis. *J Subst Abuse Treat*. 2015 April; 51: 1–18.
 doi:10.1016/j.jsat.2014.09.001.

KQ 1: In-process systematic reviews

- Final Research Plan: Drug Use in Adolescents and Adults, Including Pregnant Women: Screening.
 U.S. Preventive Services Task Force. October 2016.
 https://www.uspreventiveservicestaskforce.org/Page/Document/final-research-plan/drug-use-in-adolescents-and-adults-including-pregnant-women-screening
- Final Research Plan: Unhealthy Alcohol Use in Adolescents and Adults, Including Pregnant Women: Screening and Behavioral Counseling Interventions. U.S. Preventive Services Task Force. August 2017. https://www.uspreventiveservicestaskforce.org/Page/Document/final-research-plan/unhealthy-alcohol-use-in-adolescents-and-adults-including-pregnant-women-screening-and-behavioral-counseling-interventions
- Halladay J, Petker T, Fein A, Munn C, MacKillop J. Brief interventions for cannabis use in emerging adults. PROSPERO 2018 CRD42018085412 Available from: http://www.crd.york.ac.uk/PROSPERO/display record.php?ID=CRD42018085412

KQ 2, Completed Systematic Review:

• Steinka-Fry, Katarzyna T. et al. Culturally sensitive substance use treatment for racial/ethnic minority youth: A meta-analytic review. Journal of Substance Abuse Treatment, Volume 75, 22 - 37